Short Report:

Comparison of alternative solvents for *in situ* extraction of hydrocarbons from the colonial green alga *Botryococcus braunii* race B (Showa)

Takehiro A. Ozawa-Uyeda[a], Sebastian Overmans[b], Bárbara Bastos de Freitas[b], Edmundo Lozoya-Gloria[a]** and Kyle J. Lauersen[b]*

[a] Genetic Engineering Department, CINVESTAV-IPN Irapuato Unit, C.P. 36824 Irapuato, Gto., Mexico

[b] Bioengineering Program, Biological and Environmental Sciences and Engineering Division, King Abdullah University of Science and Technology (KAUST), Thuwal 23955-6900, Saudi Arabia

Corresponding author: kyle.lauersen@kaust.edu.sa

****Co-corresponding author:** edmundo.lozoya@cinvestav.mx

Highlights

- Effective *in situ* extraction of *B. braunii* hydrocarbons with bromodecane underlay.
- γ-Terpinene is an alternative solvent to hexane for 'milking' algal hydrocarbons.
- Green solvents are best for botryococcene extraction but degrade chlorophyll.
- Higher botryococcene extractability using modified Chu-13 medium with 5% RSW

Abstract

Background:

The colony-forming, green microalga *Botryococcus braunii* secretes large amounts of petroleum-like hydrocarbons, which enables non-destructive *in situ* extraction (milking) of these extracellular hydrocarbons during culture growth continuously without cell lysis. This study compares the suitability of 15 different solvents, including alkanes, halogens, and green solvents, for *in situ* extraction of *B. braunii* race B (Showa strain) hydrocarbons after acclimation to moderate salinity stress.

Findings:

After 24 h of extraction, bio-based terpene 'green' solvents such as ɣ-terpinene showed the highest hydrocarbon extraction capacity, with GC peak areas of botryococcenes around 10-fold greater than with conventional alkane solvents. Brominated alkanes and liquid perfluorocarbons (FCs) formed a lower phase to algal cultures rather than an upper phase as with other solvents, but only bromodecane effectively captured extracellular hydrocarbons as is the case with conventional alkane solvents. However, bromodecane and all green solvents were too toxic for two-phase continuous culture contact hydrocarbon milking, leading to 33–100% chlorophyll content loss in algal cultures.

Conclusions:

To overcome the adverse effects of these promising solvents, future research should focus on their application in short-term extraction period-milking systems to minimize algal-solvent contact and enable continuous extraction.

Keywords:

Botryococcus braunii, microalgae, in situ extraction, hydrocarbons, green solvents

Introduction

The colony-forming green microalga *Botryococcus braunii* race B produces and secretes large amounts of liquid triterpene hydrocarbons, botryococcenes, during its exponential growth phase which are stored in its extracellular matrix (ECM) (Lozoya-Gloria *et al*., 2019). Botryococcenes can be used to produce petroleum-equivalent fuels suitable for combustion engines (Jin *et al*., 2016; Lozoya-Gloria *et al*., 2019). In contrast to most other oleaginous microalgae which accumulate oils within the cell, extracellular *B. braunii* hydrocarbons are products that can be harvested without requiring cell lysis (Suzuki *et al*., 2013; Ennaceri *et al*., 2023). This trait makes it a candidate for non-destructive *in situ* extraction, 'milking', of lipophilic compounds. Milking is a promising approach to improve the cost- and energy-requirements associated with algal biomass downstream processing: harvesting, dewatering and cell disruption (Jackson *et al*., 2017; Kleinert and Griehl 2022; Ennaceri *et al*., 2023).

B. braunii typically exhibits slow growth rates compared to other photosynthetic microbes, which has limited its broader exploitation (Lozoya-Gloria *et al*., 2019). The colony surface of *B. braunii* race B is also covered with an ultrastructure called a retaining wall, which is decorated with a fibrillar sheath composed mainly of arabinose-galactose polysaccharides (Weiss *et al*., 2012). This barrier hinders the entry of many organic solvents, limiting milking efficiencies (Furuhashi *et al*., 2016b). One pretreatment that was found to improve the extraction rate of botryococcenes is cultivation of the alga in brackish medium (Furuhashi *et al*., 2013; 2016a). It was proposed that the increased salinity decreased the production or integrity of the polysaccharides, increasing colony size and flotation of *B. braunii* race B (Furuhashi *et al*., 2016a; 2016b).

Extraction of hydrocarbons from *B. braunii* has mainly been tested with commonly used

short-chain non-polar solvents, such as hexane (C6) and heptane (C7) (Jackson *et al*.,

2017; Ennaceri *et al*., 2023). These petroleum-based solvents showed high

botryococcene extraction efficiencies but low biocompatibility, resulting in rapid loss of cell

viability after long contact times (Kleinert and Griehl, 2021). Dodecane (C12) is

biocompatible with *B. braunii* cultures but has reduced hydrocarbon extractability

compared to hexane (Mehta *et al*., 2019). Alkane solvents have significant drawbacks that

can hinder scaling their application to larger culture volumes: flammability, the

requirement for corrosion-resistant infrastructure, and emulsion formation at the culture-

solvent interface (Lauersen, 2019).

It was recently reported that halogenated perfluorocarbon (FC) liquid solvents can enable

milking of sesqui- and di-terpenoids from engineered algal cultures (Overmans and

Lauersen, 2022), but these have not yet been tested on *B. braunii*. Green solvents are

another class of chemicals that are bio-based in origin and have gained popularity in

process chemistry due to their biodegradability (Kumar *et al*., 2017; De Jesus and Maciel

Filho, 2020). Monoterpenes derived from essential oils such as D-limonene, α-pinene, and

p-cymene are considered 'green' solvents because they are obtained from renewable

feedstocks, like plants and crop wastes (Kumar *et al*., 2017). Lipid extraction of microalgae

and oilseeds using terpene green solvents was shown to have equal or superior quality

and yields than with hexane (De Jesus and Maciel Filho, 2020). In this study, we evaluated

the suitability of halogenated solvents and green solvents as alternatives to conventional

alkane solvents for botryococcene extraction from *B. braunii* race B (Showa strain)

cultures grown in brackish medium derived from medium-sea water mixtures.

Materials and Methods

Reagents

The solvents tested in this work were grouped into straight-chain alkanes, halogenated solvents, and green solvents. Alkanes included n-hexane, n-decane, n-dodecane (≥99%), and n-pentadecane, which were obtained from Acros Organics (Geel, Belgium), VWR International (Fontenay-sous-Bois, France), Sigma-Aldrich (St. Louis, USA) and Alfa Aesar (Heysham, England), respectively. Halogenated solvents included brominated alkanes (bromododecane and bromododecane) and chlorinated alkane (chlorodecane), acquired from Sigma Aldrich (St. Louis, USA). Additionally, two perfluorocarbon liquids, hereafter referred to as FCs, were used in this work: the perfluorinated ether FC-770 and the perfluorinated amine FC-3283, obtained from Fluorochem (Glossop, UK) and Acros Organics (Geel, Belgium), respectively. The green solvents included three monoterpenes (para-cymene (99%), gamma-terpinene, and alpha-pinene) and three alcohols (n-heptanol, nerol, and nopol), all purchased from Sigma-Aldrich (Taufkirchen, Germany). For chlorophyll quantification, ethanol (96% vol) and methanol were purchased from VWR International (Fontenay-sous-Bois, France).

Microalgal cultures and growth conditions

Batch cultures of *Botryococcus braunii* race B, Showa strain (Nonomura, 1988), were grown in 1-L Erlenmeyer flasks in Algem photobioreactors (Algenuity©, UK) at 25 °C under 130 µmol photons $m^{-2} s^{-1}$ with a 12h:12h light:dark cycle and sparged with a 7% CO_2 in air mixture. The gas mix was supplied periodically to cultures at 25 cc min^{-1} according to the pH control level set on the photobioreactor, with the online pH measured continuously using a calibrated pH probe (Broadley James, UK) to maintain a constant

pH of 7.5 in the culture media. *B. braunii* was cultured in biological triplicates (n = 3) in

three different media: modified Chu-13 medium as the standard freshwater medium based

on the recipe in (Furuhashi *et al.*, 2013), modified Chu-13 medium with 37.5 mM NaCl,

and modified Chu-13 medium with 5% Red Sea water (RSW). Brackish medium was

prepared by mixing filtered (0.2 μm) and autoclaved RSW with modified Chu-13 medium.

The concentration of total dissolved solids in RSW was previously determined to be

around 750 mM NaCl (Abdel-Aal *et al.*, 2015). To get a final NaCl concentration of 30–40

mM (i.e., 0.3–0.4% seawater salinity), identical to the concentrations used in Furuhashi *et*

al. (2016a; 2022), Chu-13 medium was adjusted to a final concentration of 5% RSW (37.5

mM NaCl). All cultures were grown in the Algem photobioreactors with a working volume

of 400 mL for 18 days before being recovered for further analyses.

Determination of growth parameters

Final algal biomass was determined gravimetrically by obtaining a pellet of 5 mL algal

culture after adding an equal volume of 96% ethanol into a pre-weighed glass tube, mixed

manually several times, and centrifuged at 3,000 $\times$ g for 1 min to sediment all algal cells.

The addition of ethanol helps to precipitate the floating layer of algal colonies, but the

incubation time should be as short as possible to minimize extraction of undesired

pigments. After discarding the ethanol, the remaining algal pellet was dried in an oven at

110 °C for 24 h and cooled for 30 min before weighing. The dry cell weight (DCW; in mg)

of the microalgae was obtained by subtracting the dry weight of the empty glass tube from

the loaded glass tube.

Total chlorophyll content was obtained by adding 1 mL 96% ethanol to 1 mL culture

samples, mixed by inversion several times, and centrifuged at 11,000 $\times$ g for 1 min to

sediment all suspended algal cells. After discarding the ethanol, the pellet was

resuspended in 1 mL 100% methanol and incubated for 30 min at 45 ºC in the dark to

extract chlorophyll. The cell debris was pelleted by centrifugation at 11,000 $\times$ g for 1 min

while the supernatant was used to measure the absorbance at 666, 653, and 470 nm with

a spectrophotometer (TECAN Infinite M1000 PRO). Total chlorophyll content was

calculated according to Wellburn (1994).

Determination of hydrocarbon content

Algal pellet samples were collected by centrifugation as described above, and DCW was

determined. Hydrocarbons were extracted from the dried algal sample by adding 5 mL of

n-hexane and incubating for 30 min at room temperature with manual shaking at 10 min

intervals. The extraction solvent, which turned yellowish-orange due to the carotenoids

found within the ECM and the botryococcene hydrocarbons (Eroglu and Melis, 2010), was

separated from the algal residue and collected in a separate glass flask. This process was

repeated until the solvent phase became colorless. All n-hexane extracts were combined

and evaporated in an oven at 110 ºC overnight. Hydrocarbon amounts were calculated

gravimetrically and expressed as a percentage of biomass DCW. For positive

identification of botryococcenes in samples, the extracts were resuspended in 1 mL of n-

hexane or n-dodecane, and then transferred to gas chromatography (GC) vials (n = 3)

and stored at -20 °C until further analysis as described below.

Preliminary screening of solvents for two-phase 'milking' of *B. braunii* race B in low salinity culture medium

Two-phase living extractions (i.e., 'milking') of extracellular hydrocarbons were performed with alkanes, halogenated-, and green solvents by gently adding 250 µL of each respective solvent (25% v/v) onto 750 µL liquid cultures of *B. braunii* that were previously grown in brackish medium for 18 days. All two-phase cultures were grown for 24 h in duplicates in 2-mL microcentrifuge tubes on an orbital shaker at 120 rpm with a 12h:12h light:dark cycle at room temperature. The solvent layer was then collected following centrifugation at 4,500 × *g* for 5 min to obtain two separate layers. From each sample, 150 µL of the solvent fraction was transferred into individual GC vials and stored at -20 °C until further analysis, as described below.

Alternative solvents for *in situ* extraction of hydrocarbons produced by *B. braunii* race B

The two most efficient extraction solvents of each group were selected from the initial screening. Two-phase living extractions of extracellular hydrocarbons were performed by gently adding 1 mL of each respective solvent onto 9 mL liquid cultures of *B. braunii* (10% v/v) that were previously grown in a brackish medium for 18 days. The two-phase cultures were cultivated for 24 h in triplicates in 50-mL Erlenmeyer flasks on an orbital shaker at 120 rpm with a 12h:12h light:dark cycle at room temperature. The solvent and algal samples were obtained at two different time points: 2 h and 24 h of cultivation. At each time point, the solvent layer was collected after centrifugation at 4,500 × *g* for 5 min. From each sample, 150 µL of the solvent fraction was transferred into separate GC vials and stored at -20 °C until further analysis.

Gas chromatography

All solvent extracts were analyzed using a gas chromatograph equipped with a mass

spectrometer and a flame ionization detector (GC-MS-FID) with modifications to a

previously described protocol (Thapa *et al.*, 2016). Briefly, the GC-MS-FID analyses were

performed using an Agilent 7890A gas chromatograph connected to a 5975C inert MSD

with a triple-axis detector. The system was equipped with a DB-5MS column (30 m ×

0.25 mm i.d., 0.25 μm film thickness) (Agilent J&W, USA). The temperatures were set to:

injector (260 °C), interface (250 °C), and ion source (200 °C). One μL of liquid solvent

sample was injected in splitless mode with an autosampler (Model G4513A, Agilent).

Column flow was kept constant at 2.5 mL min^{-1} with He as a carrier gas. The initial GC

oven temperature of 220 °C was held for 1 min, then raised to 280 °C at a rate of 5 °C

min^{-1}, followed by 2 °C min^{-1} to 300 °C and held for 5 min. Mass spectra were recorded

after a 3 min solvent delay using a scanning range of 50–750 m/z at 20 scans s^{-1}.

Gas chromatograms were manually reviewed for quality control before botryococcene GC

peak areas were integrated using MassHunter Workstation software version B.08.00

(Agilent Technologies, USA). The NIST library (National Institute of Standards and

Technology, Gaithersburg, MD, USA) was used to identify botryococcenes.

Statistical analysis

204 Differences between the mean values obtained from each assay were assessed through

205 one-way analysis of variance (ANOVA) followed by Tukey's HSD post-hoc test with a

206 confidence level of 95%. All statistical analyses were performed using R software v3.4.3

207 (https://www.r-project.org/) through the RStudio interface.

Results and Discussion

B. braunii growth and hydrocarbon extractability in different media

After 18 days of cultivation, there were no significant differences in specific growth rates of *B. braunii* cultured in modified Chu13 medium (0.077 ± 0.002 day^{-1}; mean $\pm$ SEM) and in 5% RSW (0.064 ± 0.007 day^{-1}), but the algae cultured in 37.5 mM NaCl (0.050 ± 0.005 day^{-1}) showed a significant decrease ($p < 0.01$) in the specific growth rate compared to the other treatments (Fig. 1A). *B. braunii* cultured in 5% RSW had significant decreases in chlorophyll content (1.7 ± 0.19 mg L^{-1}) and appeared less healthy (chlorotic phenotype) compared to algae cultured in non-saline medium (3.0 ± 0.07 mg L^{-1}). *B. braunii* cultured in 37.5 mM NaCl contained less chlorophyll (1.2 ± 0.10 mg L^{-1}) than in 5% RSW treatment (Fig. 1B). The hydrocarbon content in dry biomass of *B. braunii* cultured in the brackish medium was 33 ± 0.5% DCW compared to 40 ± 2.1% in modified Chu13 medium (Fig. 1C). However, the amount of botryococcenes in 5% RSW was at least 2-fold greater than in the algae cultured in non-saline and 37.5 mM NaCl conditions (Fig. 1D). The presence of C_{34} and C_{32} botryococcenes from suspended *B. braunii* cells cultured in 5% RSW was confirmed by the appearance of two distinct peaks (9.7 min and 10.2 min, respectively) with appropriate mass spectra in GC-MS chromatograms (Fig. 1E). These results indicate that culturing *B. braunii* in modified Chu-13 medium with 5% RSW is possible without strongly reducing its growth and encourages the production of extracellular botryococcenes. This is in line with previous reports (Furuhashi *et al.*, 2013; 2016; 2022), where *B. braunii* cultivation in brackish medium reduced the cultures' growth rate but improved hydrocarbon extractability from algal biomass compared to cultivation in non-saline medium.

Figure 1. Analysis of *B. braunii* (Showa strain) cultured for 18 days in three different media. (**A**) Specific growth rate, (**B**) total chlorophyll content, and (**C**) extracellular hydrocarbon content of *B. braunii* race B cells grown in modified Chu-13 medium (Chu13), modified Chu-13 medium with 37.5 mM NaCl (37.5 mM NaCl), and modified Chu-13 medium with 5% Red Sea water (5% RSW). (**D**) Total GC-FID peak area of extracted botryococcenes. Peak areas are the sum of two detected botryococcenes peaks. (**E**) Representative GC-FID chromatograms of botryococcenes extracted with commonly used alkane solvents obtained from wet *B. braunii* algal samples. Error bars represent standard error of the mean of three biological replicates (SEM; n = 3). Statistical significance was determined by one-way ANOVA with Tukey's HSD post-hoc test (p < 0.05). Within each panel, different letters above bars indicate instances where means were significantly different (*p* < 0.05).

Suitability screening of solvents for two-phase cultivation of *B. braunii* race B in

brackish culture medium

A recent study has shown that the Showa strain is one of the most promising for *in situ* extraction because of its high solvent resistance, hydrocarbon extractability, and lipid productivity (Kleinert and Griehl 2021). Our results showed that modified Chu-13 medium with 5% RSW was the best for growing and improving botryococcene accumulation around cultured *B. braunii* Showa (Fig. 1). Because *B. braunii* hydrocarbons are mainly (90–95%) stored in the ECM, most of them can be extracted by soaking the algae in a non-polar organic solvent such as hexane that does not permeate the cells (Furuhashi *et al.*, 2013; Suzuki *et al.*, 2013). Hydrocarbons obtained in this way are typically referred to as extracellular hydrocarbons. After recovering the extracellular hydrocarbons, a small amount of intracellular hydrocarbons can also be extracted with a mixture of chloroform and methanol that can reach inside the cells (Suzuki *et al.*, 2013; Thapa *et al.*, 2016). For the initial solvent screening, cultures grown in 5% RSW were treated with 14 different non-polar solvents at 25% (v/v), whose physicochemical properties are shown in Table 1.

After 2 and 24 h extraction, the solvents exhibited a highly variable performance in extracting botryococcene hydrocarbons from the algal cells (Fig. 2A). This was visually indicated by the co-extraction of extracellular carotenoids that had previously been reported to be associated with the botryococcene fraction from the Showa strain (Eroglu and Melis, 2010). Most of the solvents showed yellowish coloration at different levels in the upper phase (Fig. 2A). Brominated alkanes and FCs formed a lower phase (Fig. 2A). Alkane solvents showed a similar but relatively poor extraction of botryococcenes (Fig. 2B). Halogenated solvents were worse than alkanes except for bromodecane, which was the best of all previously tested solvents (Fig. 2B). However, the amount of

botryococcenes extracted by green solvents was found to be up to ~10-fold and ~3.8-fold greater than with dodecane and bromodecane, respectively (Fig. 2B). The exception was nopol which did not extract botryococcenes (Fig. 2B). This indicates that green solvents, such as ɣ-terpinene and α-pinene, were more efficient at extracting botryococcenes *in situ* compared to conventional alkane solvents and their halogen derivatives (Fig. 2B).

Figure 2. Extraction and toxicity properties of all different solvents. (**A**) *B. braunii* race B cells grown in culture medium with 5% RSW after 2 h and 24 h extraction with 25% (v/v) of solvents. (**B**) GC-FID peak areas of botryococcenes in solvents recovered after 24 h extractions. These are the sum of two compound peaks identified as botryococcenes. (**C**) Total chlorophyll content in *B. braunii* cultured in 5% RSW medium after 24 h extraction with each solvent. Error bars represent the standard error of the mean of two biological replicates (SEM; n = 2). Statistical significance was determined by one-way ANOVA with post-hoc Tukey's HSD test. Letters above the bars indicate significantly different mean concentrations ($p < 0.05$).

Although both α-pinene and ɣ-terpinene yielded the highest accumulation of botryococcenes in this screening (Fig. 2B), we observed a severe degradation of chlorophyll in all green solvents extracts after 24 h of incubation compared to alkanes and halogenated solvents (Fig. 2C). The total chlorophyll content in *B. braunii* after two-phase cultivation with solvents was measured to evaluate solvent biocompatibility. The addition of solvents to *B. braunii* led to differing extents of chlorosis. Of the different two-phase cultivations, algal cultures with alkanes showed a slight decrease in the chlorophyll content compared to cultivations without solvent (Fig. 2C). Algae with brominated alkanes and chlorododecane demonstrated a stronger reduction in chlorophyll content (3.6–4.5 mg L^{-1}) than those with straight-chain alkanes (6.3–7.5 mg L^{-1}) while cultures with FCs had chlorophyll contents (8.3–8.9 mg L^{-1}) similar to the no-solvent treatment (9.3 ± 0.17 mg L^{-1}) (Fig. 2C). Green solvents were the most toxic solvents, particularly the alcohols, leading to a drastic decrease in chlorophyll content to around 0–1 mg L^{-1} (Fig. 2C). Our results indicate that green solvents are too toxic for two-phase cultivations, unlike alkanes and halogenated solvents. A previous study reported similar results after comparing the effects of decane, its alcohol derivative decanol, and the green solvent D-

limonene, where decanol was the most toxic solvent, followed by D-limonene and decane,

(Concha *et al.*, 2018). Here, brominated alkanes and FCs formed a lower phase or

'underlay' with algal cultures, unlike other solvents that formed a distinct layer on top of

the culture (Fig. 2A). Of the heavy solvents, only bromodecane could capture

hydrocarbons from wet algal samples (Fig. 2B). In contrast, previous work reported that

FCs could replace the use of dodecane to extract the sesqui- and diterpenoids produced

by engineered green microalgae (Overmans and Lauersen, 2022).

Selected solvents for *in situ* extraction of hydrocarbons produced by *B. braunii*

race B

We evaluated *in situ* hydrocarbon extraction in larger culture volumes at two incubation

periods, 2 h and 24 h, with six solvents: hexane, dodecane, bromodecane,

bromododecane, p-cymene and ɣ-terpinene. These were used at concentrations of 10%

(v/v), the standard culture:solvent ratio used to capture terpenoids or hydrocarbons from

engineered algae (Overmans and Lauersen, 2022; Yunus *et al.*, 2022). Regardless of

solvent density, we observed the formations of emulsions in cultures with all solvents after

shaking the mixtures at 120 rpm even after 2 h (Fig. 3A). This formation was previously

identified as one technical limitation for isoprenoid milking with hydrophobic solvents on

algal culture (Lauersen, 2019). After 2 h of extraction, the alkane solvents dodecane and

hexane showed a yellow coloration, indicating hydrocarbon extraction, while the aqueous

lower phase contained algae and culture medium. After 24 h of extraction, algal cultures

were only green in dodecane, which also exhibited a more intense yellow color

accumulation than hexane, where cells were chlorotic or dead (Fig. 3A).

Brominated solvents formed a lower phase to the aqueous algal culture where colonies

clumped, but no hydrocarbon extraction occurred from either solvent. Terpene green

solvents were found to extract the hydrocarbons, with aggregation of colonies visible at 2

h and severe chlorophyll degradation at 24 h of extraction (Fig. 3A). After a 2 h contact

time between solvents and the 5% RSW-acclimated *B. braunii* cells, we detected the

highest hydrocarbon content in hexane, followed by p-cymene and ɣ-terpinene (Fig. 3B).

The concentrations of botryococcenes in extractions with dodecane and brominated

alkanes were found to be much higher after 24 h than after 2 h (Fig. 3B). Bromodecane

showed a similar hydrocarbon accumulation compared to the two alkanes after 24 h of

incubation. Even though the extraction of botryococcenes with bromodecane resulted in

a 3.8-fold increase compared to bromododecane (Fig. 3B), both of the tested brominated

alkanes caused a similar decrease in chlorophyll content compared to dodecane from 2.5

$\pm$ 0.05 mg L^{-1} to ~1.8 mg L^{-1} (Fig. 3C). *B. braunii* race B cells cultured with hexane and

p-cymene showed a fast decrease in chlorophyll content (2.42 $\pm$ 0.11 mg L^{-1} and 1.78 $\pm$

0.06 mg L^{-1}, respectively), whereas algal cells with ɣ-terpinene (2.86 $\pm$ 0.04 mg L^{-1})

showed a similar chlorophyll content to cells with dodecane after 2 h of incubation (2.97 $\pm$

0.05 mg L^{-1}) (Fig. 3A and 3C). Longer contact time (24 h) with ɣ-terpinene was detrimental

to algal cells (0.45 $\pm$ 0.04 mg L^{-1}), as could be seen in the hexane treatment (Fig. 3A and

3C). This result indicates that the bromodecane underlay was amenable to accumulating

extracellular hydrocarbons and carotenoids from *B. braunii* cells, but it was more

aggressive as a two-phase culture system than the conventional dodecane overlay.

Figure 3. Extraction and toxicity properties of selected solvents. (**A**) *B. braunii* cells grown in culture medium with 5% RSW after a contact time of 2 h and 24 h with 10% (v/v) of either (1) hexane, (2) dodecane, (3) bromododecane, (4) bromodecane, (5) p-cymene, or (6) γ-terpinene. (**B**) Total GC-FID peak areas of two botryococcenes peaks after 2 h and 24 h of two-phase living *B. braunii* extraction cultures with solvents. (**C**) Total chlorophyll content of *B. braunii* cultured in medium with 5% RSW after 2 h and 24 h extraction time with solvents. Error bars represent standard error of the mean of three biological replicates (SEM; n = 3). Statistical significance was determined by one-way ANOVA with post-hoc Tukey's HSD test. Lower- and upper-case letters above the bars indicate the post hoc results of the 2 h and 24 h data sets, respectively. Means were considered significantly different at level $p < 0.05$.

The highest botryococcene contents were detected in p-cymene and ɣ-terpinene extracts

after 24 h of incubation, both showing GC peak areas of botryococcenes 13- and 15-fold

greater than hexane and dodecane, respectively (Fig. 3B). As far as we know, this is the

first report of using γ-terpinene in botryococcenes extraction. On the other hand, p-

cymene has been used to extract lipids from different biomass sources, including

oleaginous microalgae (De Jesus and Maciel Filho, 2020). Even though we found a similar

hydrocarbon accumulation between p-cymene and ɣ-terpinene when the solvent content

was 10% v/v (Fig. 3B), p-cymene was more aggressive on the cells than ɣ-terpinene (Fig.

3C). It has been previously reported that p-cymene was more toxic against insects than

γ-terpinene, indicating the latter will likely be more appropriate for future botryococcene

milking setups (Gong and Ren, 2020).

The results indicate that ɣ-terpinene could be used as an alternative solvent to replace

hexane for *in situ* extractions of hydrocarbons, but only for short extraction times. Different

bioprocess designs to the *in situ* extraction procedure used in our study could help to

reduce the toxic effects of solvent contact on algal cells. For example, it was recently

shown that *B. braunii* cultures grown in a bubble column bioreactor attached to an

extraction column could achieve a long-term *in situ* extraction of extracellular

hydrocarbons over 30 days by using a daily extraction time of 5 h with hexane (Kleinert

and Griehl 2022). Other *in situ* extraction systems and setups, such as membrane

filtration, have shown to be successful in extracting exopolysaccharides (EPS) from *B.*

braunii CCALA778 by using a microfiltration hollow-fiber membrane without requiring

solvent (García-Cubero *et al.,* 2018). A similar system was used to continuously extract

the heterologous sesquiterpenoid patchoulol with a hydrophobic hollow-fiber membrane

acting as a physical interaction contact matrix between engineered algal cells and the

solvent dodecane (Overmans *et al.*, 2022).

Hexane is a suitable extraction solvent due to its low boiling point (69 °C), low cost, and

high extraction efficiency (Jackson *et al.*, 2017; Ennaceri *et al.*, 2023). Using hexane

facilitates solvent separation from the botryococcenes by distillation, as they have boiling

points well above hexane (> 300 °C; Jackson *et al.*, 2017). Hexane would be superior to

brominated alkanes and the terpene green solvents with boiling points > 150 °C, but only

if the downstream process relies on energy-intensive distillation (Table 1). Other emerging

techniques, including organic solvent nanofiltration and alternative wetting of membranes,

may enable low-energy separations of botryococcenes, similar to those recently reported

for sesquiterpene alcohols (Jackson *et al.*, 2017; Alduraiei *et al.*, 2021; Overmans *et al.*,

2022). Terpene green solvents used in this work (134–136 g/mol; Table 1) have smaller

molecular weights than botryococcenes (e.g., 466.8 g/mol $C_{34}H_{58}$), which would facilitate

their downstream separation by these techniques. Furthermore, the improved extraction

efficiencies and sustainable nature of green solvents make them attractive alternatives to

hexane.

Conclusions

Using brackish cultivation to improve hydrocarbon extractability is a step-change in milking efficiency potential for *B. braunii* cultures, but the solvent choice was limited to the use of straight-chain alkanes. Here, we show that terpene green solvents and heavy solvents like bromodecane could also be used as alternative solvents to hexane for milking botryococcenes from *B. braunii* race B cultured in brackish medium. However, they are less biocompatible than dodecane, leading to 33–100% chlorophyll content loss. Our results warrant follow-up testing of these parameters in short-time scale solvent contactor systems that may improve the *B. braunii* hydrocarbon milking processes.

Abbreviations

FC: Perfluorocarbon; ECM: Extracellular matrix; RSW: Red Sea water; DCW: Dry cell weight; GC-MS-FID: Gas chromatography–mass spectrometry-flame ionization detector.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

All the authors consented to the publication of this work and its submission to Bioresources and Bioprocessing

Availability of data and materials

All data generated or analyzed during this study are included in this article and its supplementary information file.

Competing Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Funding

TAO-U is financially supported by CONAHCYT (Consejo Nacional de Humanidades, Ciencias y Tecnologías) Mexico with a PhD fellowship (CVU 780487) and the KAUST Visiting Student Research Program (VSRP). The research reported in this publication was supported by KAUST baseline funding awarded to KJL.

Authors' contributions

TAO-U performed the experiments, contributed to the research planning, experimental design, data collection and visualization, formal analysis, and wrote the original draft manuscript. SO contributed to the research planning, experimental design, GC-MS analysis, and data visualization, and reviewed the manuscript. BBdF contributed to experimental design, assisted in the operation of Algem photobioreactors, and reviewed

the manuscript. EL-G contributed to funding acquisition and data visualization, and

reviewed the manuscript. The research was mainly conducted in the laboratory of KJL,

who was responsible for experimental design, project scope, funding acquisition and

reviewing the manuscript. All authors read and approved the final manuscript.

Acknowledgments

We would like to thank Chandrasekaran Lakshmipathy of the KAUST Lab Equipment

Maintenance (LEM) team for assistance in upgrading and initializing the GC-FID-MS unit.

References

1. Abdel-Aal, E. A., Farid, M. E., Hassan, F. S., & Mohamed, A. E. (2015). Desalination of Red Sea water using both electrodialysis and reverse osmosis as complementary methods. *Egyptian Journal of Petroleum*, *24*(1), 71-75. https://doi.org/10.1016/j.ejpe.2015.02.007

2. Alduraiei, F., Manchanda, P., Pulido, B., Szekely, G., & Nunes, S. P. (2021). Fluorinated thin-film composite membranes for nonpolar organic solvent nanofiltration. *Separation and Purification Technology*, *279*, 119777. https://doi.org/10.1016/j.seppur.2021.119777

3. Concha, E., Heipieper, H. J., Wick, L. Y., Ciudad, G. A., & Navia, R. (2018). Effects of limonene, n-decane and n-decanol on growth and membrane fatty acid composition of the microalga *Botryococcus braunii*. *AMB Express*, *8*, 189. https://doi.org/10.1186/s13568-018-0718-9

4. De Jesus, S. S., & Maciel Filho, R. (2020). Recent advances in lipid extraction using green solvents. *Renewable and Sustainable Energy Reviews*, *133*, 110289. https://doi.org/10.1016/j.rser.2020.110289

5. Ennaceri, H., Nwoba, E. G., Ogbonna, C. N., Bahri, P. A., & Moheimani, N. R. (2023). Progress of non-destructive hydrocarbon extraction technology of *Botryococcus braunii*. *Algal Research*, *73*, 103156. https://doi.org/10.1016/j.algal.2023.103156

6. Eroglu, E., & Melis, A. (2010). Extracellular terpenoid hydrocarbon extraction and quantitation from the green microalgae *Botryococcus braunii* var. Showa. *Bioresource Technology*, *101*(7), 2359-2366. https://doi.org/10.1016/j.biortech.2009.11.043

7. Furuhashi, K., Saga, K., Okada, S., & Imou, K. (2013). Seawater-cultured *Botryococcus braunii* for efficient hydrocarbon extraction. *PLoS One*, *8*(6), e66483. https://doi.org/10.1371/journal.pone.0066483

8. Furuhashi K, Hasegawa F, Saga K, Okada S, Kaizu Y, & Imou K. (2016a). Effects of culture medium salinity on the hydrocarbon extractability, growth, and morphology of *Botryococcus braunii*. *Biomass and Bioenergy, 91*, 83–90. https://doi.org/10.1016/j.biombioe.2016.05.007

9. Furuhashi, K., Noguchi, T., Okada, S., Hasegawa, F., Kaizu, Y., & Imou, K. (2016b). The surface structure of *Botryococcus braunii* colony prevents the entry of extraction solvents into the colony interior. *Algal Research*, 16, 160-166. https://doi.org/10.1016/j.algal.2016.02.021

10. Furuhashi, K., Magota, A., Liu, Y., Hasegawa, F., Okada, S., Kaizu, Y., & Imou, K. (2022). Thinning of *Botryococcus braunii* Colony Sheath by Pretreatment Enhances Solvent-Based Hydrocarbon Recovery. *Phycology*, *2*(4), 363-373. https://doi.org/10.3390/phycology2040020

11. García-Cubero, R., Wang, W., Martín, J., Bermejo, E., Sijtsma, L., Togtema, A., ... & Kleinegris, D. M. (2018). Milking exopolysaccharides from *Botryococcus braunii* CCALA778 by membrane filtration. *Algal Research*, *34*, 175-181. https://doi.org/10.1016/j.algal.2018.07.018

12. Gong, X., & Ren, Y. (2020). Larvicidal and ovicidal activity of carvacrol, p-cymene, and γ-terpinene from *Origanum vulgare* essential oil against the cotton bollworm, *Helicoverpa armigera* (Hübner). *Environmental Science and Pollution Research*, *27*, 18708-18716. https://doi.org/10.1007/s11356-020-08391-2

13. Jackson, B. A., Bahri, P. A., & Moheimani, N. R. (2017). Repetitive non-destructive
milking of hydrocarbons from *Botryococcus braunii*. *Renewable and Sustainable Energy*
Reviews, *79*, 1229-1240. https://doi.org/10.1016/j.rser.2017.05.130

14. Jin, J., Dupré, C., Yoneda, K., Watanabe, M. M., Legrand, J., & Grizeau, D. (2016).
Characteristics of extracellular hydrocarbon-rich microalga *Botryococcus braunii* for
biofuels production: Recent advances and opportunities. *Process Biochemistry*, *51*(11),
1866-1875. https://doi.org/10.1016/j.procbio.2015.11.026

15. Kleinert, C., & Griehl, C. (2021). Identification of suitable Botryococcus braunii strains for
non-destructive in situ hydrocarbon extraction. *Journal of Applied Phycology*, *33*(2), 785-
798. https://doi.org/10.1007/s10811-020-02342-7

16. Kleinert, C., & Griehl, C. (2022). In situ extraction (milking) of the two promising
Botryococcus braunii strains Showa and Bot22 under optimized extraction time. *Journal*
of Applied Phycology, *34*(1), 269-283. https://doi.org/10.1007/s10811-021-02633-7

17. Kumar, S. J., Kumar, G. V., Dash, A., Scholz, P., & Banerjee, R. (2017). Sustainable
green solvents and techniques for lipid extraction from microalgae: A review. *Algal*
Research, *21*, 138-147. https://doi.org/10.1016/j.algal.2016.11.014

18. Lauersen, K. J. (2019). Eukaryotic microalgae as hosts for light-driven heterologous
isoprenoid production. *Planta*, *249*, 155-180. https://doi.org/10.1007/s00425-018-3048-
x

19. Lozoya-Gloria, E., Morales-de la Cruz, X. and Ozawa-Uyeda, T. A. (2019). "The Colonial
Microalgae *Botryococcus braunii* as Biorefinery". Ch. 7. In: *Microalgae - From Physiology*
to Application. Vítová M. (Ed.) IntechOpen, London, UK. Pp. 123-144.
https://doi.org/10.5772/intechopen.88206

20. Mehta, P., Jackson, B. A., Nwoba, E. G., Vadiveloo, A., Bahri, P. A., Mathur, A. S., &
Moheimani, N. R. (2019). Continuous non-destructive hydrocarbon extraction from
Botryococcus braunii BOT-22. *Algal Research*, *41*, 101537.
https://doi.org/10.1016/j.algal.2019.101537

21. Nonomura, A. M. (1988). *Botryococcus braunii* var. Showa (Chlorophyceae) from
Berkeley, California, United States of America. *Japanese Journal of Phycology*, *36*, 285-
291. https://patents.google.com/patent/USPP6169P/en

22. Overmans, S., Ignacz, G., Beke, A. K., Xu, J., Saikaly, P. E., Szekely, G., & Lauersen,
K. J. (2022). Continuous extraction and concentration of secreted metabolites from
engineered microbes using membrane technology. *Green Chemistry*, *24*(14), 5479-
5489. https://doi.org/10.1039/D2GC00938B

23. Overmans, S., & Lauersen, K. J. (2022). Biocompatible fluorocarbon liquid underlays for
in situ extraction of isoprenoids from microbial cultures. *RSC Advances*. *12*(26), 16632-
16639. https://doi.org/10.1039/D2RA01112C

24. Suzuki, R., Ito, N., Uno, Y., Nishii, I., Kagiwada, S., Okada, S., & Noguchi, T. (2013).
Transformation of lipid bodies related to hydrocarbon accumulation in a green alga,
Botryococcus braunii (Race B). *PLoS One*, *8*(12), e81626.
https://doi.org/10.1371/journal.pone.0081626

25. Thapa, H. R., Naik, M. T., Okada, S., Takada, K., Molnár, I., Xu, Y., & Devarenne, T. P.
(2016). A squalene synthase-like enzyme initiates production of tetraterpenoid
hydrocarbons in *Botryococcus braunii* Race L. *Nature Communications*, *7*, 11198.
https://doi.org/10.1038/ncomms11198

26. Weiss, T. L., Roth, R., Goodson, C., Vitha, S., Black, I., Azadi, P., ... & Goodenough, U.
(2012). Colony organization in the green alga *Botryococcus braunii* (Race B) is specified

by a complex extracellular matrix. *Eukaryotic Cell*, *11*(12), 1424-1440.
https://doi.org/10.1128/ec.00184-12
27. Wellburn, R. W. (1994). The spectral determination of chlorophylls a and b, as well as
total carotenoids, using various solvents with spectrophotometers of different resolution.
Journal of Plant Physiology, *144*(3), 307-313. https://doi.org/10.1016/S0176-
1617(11)81192-2
28. Yunus, I. S., Anfelt, J., Sporre, E., Miao, R., Hudson, E. P., & Jones, P. R. (2022).
Synthetic metabolic pathways for conversion of CO_2 into secreted short-to medium-chain
hydrocarbons using cyanobacteria. *Metabolic engineering*, *72*, 14-23.
https://doi.org/10.1016/j.ymben.2022.01.017
542

 Tables

544 **Table 1**. Physico-chemical properties of the solvents used in this work.

	Solvent	Molecular formula	Molecular weight (g/mol)	Density (g/mL)	Boiling point (ºC)	Water solubility at 20ºC (mg/L)
Alkane solvents	Decane	$C_{10}H_{22}$	142	0.73	174	0.052
	Dodecane	$C_{12}H_{26}$	170	0.75	216	0.0037
	Pentadecane	$C_{15}H_{32}$	212	0.77	271	0.00004
Halogenated solvents	Bromodecane	$C_{10}H_{21}Br$	221	1.07	241	Insoluble
	Bromododecane	$C_{12}H_{25}Br$	249	1.04	276	Insoluble
	Chlorododecane	$C_{12}H_{25}Cl$	205	0.73	< 1000	0.052
	FC-770	C_4F_9NO	399	1.79	95	1.3
	FC-3283	$C_9F_{21}N$	521	1.82	128	< 5.0
Green solvernts	α-Pinene	$C_{10}H_{16}$	136	0.86	156	2.49
	γ-Terpinene	$C_{10}H_{16}$	136	0.85	183	8.72
	p-Cymene	$C_{10}H_{14}$	134	0.86	177	23.4
	Heptanol	$C_7H_{16}O$	116	0.82	176	1670
	Nerol	$C_{10}H_{18}O$	154	0.88	225	1311
	Nopol	$C_{13}H_{18}O$	166	0.97	230–240	333.5
Control	Hexane	C_6H_{14}	86	0.66	69	0.01

545 Hexane is the standard botryococcene extraction solvent.