Skin lactate dynamics and lactate threshold during exercise

Lactate Dynamics in Skin Interstitial Fluid Predicts Lactate Threshold in Exercising Rats

Shohei Dobashi[1,2]*, Daisuke Funabashi[1,3]*, Kazuki Sameshima[1], Noriko Tsuruoka[4], Takashi

Matsui[1,2,3]

[1] Exercise Biochemistry & Sport Neurobiology Division, Institute of Health and Sport Sciences,

University of Tsukuba, Tsukuba, Japan

[2] R&D Center for Lifestyle Innovation, University of Tsukuba, Tsukuba, Japan

[3] Advanced Research Initiative for Human High Performance, University of Tsukuba, Tsukuba,

Japan

[4] Graduate School of Biomedical Engineering, Tohoku University, Sendai, Japan.

*These authors contributed equally to this work.

Correspondence:

Takashi Matsui Ph.D.

Assistant Professor

Exercise Biochemistry & Sport Neurobiology Division,

Institute of Health and Sport Sciences, University of Tsukuba

E-mail: matsui.takashi.ga@u.tsukuba.ac.jp   Tel: +81-29-853-2750

Skin lactate dynamics and lactate threshold during exercise

Abstract

The blood lactate threshold (LT) is a crucial marker of metabolic shifts and stress transitions
during exercise, influencing optimal intensity determination and chronic exercise impact
evaluation across diverse individuals. Although recent studies have explored methods for
estimating blood lactate levels without blood sampling, accurate solutions for LT estimation
remain undeveloped. Here, we tested the analytical capability of an ultra-minimally invasive
technique for LT measurement that harnesses skin interstitial lactate dynamics using a single
microneedle perfusion system. The rats were acclimated to exercise and underwent external
jugular vein cannulation. Following comprehensive recovery, a 1 mm single microneedle was
implanted into the back skin. This allowed for the simultaneous measurement of lactate
concentrations in the blood and skin interstitial fluid during incremental treadmill exercise. Skin
lactate progressively increased in tandem with blood lactate during incremental exercise, albeit
with lingering post-exercise clearance that did not correlate with blood lactate. Remarkably, LT
extrapolated from skin lactate showed significant alignment with blood LT. Our findings reveal
the potential of skin interstitial dynamics to accurately assess LT during exercise using a single
microneedle, integrating the discoveries of exercise physiology into real-world athletic and/or
healthy practices.

Keywords
Exercise, Lactate, Blood, Skin interstitial fluid, Single microneedle, Rat

Graphical abstract

Representative graphical image of exercise rats with inserted an ultra-minimally invasive single
microneedle (*A*). Representative graph of LT assessed by the blood (*B*) and skin fluid lactate (*C*)
dynamics. The findings suggest that LT determination using the perfusate collected by the single
microneedle perfusion system can be an effective alternative to current methods for measuring
blood lactate, which may be a breakthrough in the development of sports science.

Skin lactate dynamics and lactate threshold during exercise

Introduction

In the field of sports science, where the enhancement of athletic performance and promotion of

physical well-being are paramount objectives, the evaluation of exercise intensity and assessment

of chronic exercise impact assumes a pivotal role. Traditionally, blood lactate concentration has

been used as a fundamental physiological marker, with particular emphasis on its dynamic

behavior during periods of rest, exercise, and recovery. In this context, universal indices that

denote aerobic and anaerobic work capacities encompassing the lactate threshold (LT), peak

lactate concentration, and maximal lactate steady state (MLSS) during constant and incremental

load exercises have been developed and used in a diverse population of athletes and exercise

enthusiasts[1,2]. Hence, continuous, real-time assessment of blood lactate concentration during

exercise is important for the development of sports medicine and science. However, continuous

measurements are needed to mandate the insertion of indwelling needles [3,4] or a high-frequency

blood sampling regime[5]. Therefore, despite the acknowledged significance of continuous blood

lactate monitoring during exercise, there is a considerable gap between theoretical recognition

and practical applications.

In recent sports science, there has been a noteworthy surge in the rapid exploration of

alternative biomarkers as replacements for blood lactate concentration, along with the

development of robust quantitative assessment methodologies. Devices such as mouthpieces,

eyewear, and tattoo-seal-types have emerged as non-invasive platforms for the collection of saliva

and sweat lactate[6-8]. Seki et al. confirmed that LT calculated from sweat lactate levels during

exercise was significantly associated with LT derived from blood lactate concentrations[9].

However, the same research group also reported that there were some cases in which sweat LT

failed to align with blood LT, attributable to individual variations in sweat patterns[10]. Moreover,

the evaluation of sweat lactate cannot be performed in scenarios where exercise occurs in cold

environments, during swimming activities, or in low-intensity workouts, where sweat production

is minimal or infrequent.

We then focused our attention on the skin interstitial fluid (ISF). The primary constituents of

ISF are plasma, and low-molecular-weight substances such as blood lactate and blood glucose are

believed to exhibit profiles in skin tissue fluid similar to their blood concentrations, as they can

diffuse across vessel walls[11,12]. Indeed, previous studies have reported correlations between the

dynamics of glucose and lactate concentrations in human skin ISF and the blood[13,14]. More

recently, systems have been developed that allow rapid and continuous monitoring of skin ISF

lactate as an electrical current using enzyme electrodes within wearable sensors[15,16]. Furthermore,

a recent investigation has demonstrated that the changes in ISF lactate during exercise have

recently commenced, and suggested similarities to the blood lactate concentration dynamics[17].

However, it has been reported that there is a time lag between the variations in lactate in the skin
interstitial fluid during exercise and those in the blood lactate concentration[18,19]. This time lag has
been attributed to the extended collection of interstitial fluid due to the spread of wounds around
the indwelling needle during exercise, resulting in collection of the same volume of interstitial
fluid over an extended period[18,19]. Recent wearable devices have relatively minimally invasive
needles, and many of them have been designed with multiple projections. This can potentially
increase user discomfort during prolonged physical activity and may lead to the expansion of
sensor insertion site wounds, thereby introducing time delays in the immediate monitoring of
lactate levels in the skin interstitial fluid.

In contrast, Tsuruoka et al. designed a single ultra-minimally invasive micro needle with an
outer diameter of approximately 0.2 mm and a length of 1-2 mm[20]. The microneedle perfusion
system can collect skin ISF using circulating saline solution. The authors administered a lactate
solution to mice with a micro needle inserted, and lactate concentrations were quantified directly
outside the body using a lactate analysis device. The results indicated a close temporal alignment
between post-lactate administration blood lactate concentrations and skin fluid lactate
concentrations with a significant positive correlation. Additionally, the range of blood lactate
concentrations that exhibited a significant correlation with skin fluid lactate concentrations at this
time exceeded the onset of blood lactate accumulation (OBLA) (4 mM), suggesting that the
microneedle perfusion system of Tsuruoka et al. may offer the potential to predict LT from skin
ISF lactate levels during incremental load exercises[20]. Such a system appears to minimize the
impact of enzyme electrode drawbacks, which are susceptible to interference due to prolonged
wear[21], while significantly reducing the user's burden during LT evaluation. However, it remains
uncertain whether lactate collected in the skin ISF through the microneedle perfusion system
accurately reflects the LT during incremental load exercises.

Here, we provide the first scientific evidence of the analytical capability of an ultra-minimally
invasive technique for LT measurement that harnesses skin interstitial lactate dynamics using a
single microneedle perfusion system. Previous studies have investigated the possibility of sweat
contamination in ISF samples collected from the epidermis[22] as well as the potential negative
effects on exercise performance due to repeated blood sampling during traditional incremental
exercise, leading to increased pain and stress[23]. To address these concerns, we conducted
incremental exercise in rats without sweat production in the epidermis. The results of this study
suggest the potential use of skin ISF lactate as an effective alternative to current methods for
measuring blood lactate levels during exercise, offering groundbreaking, world-first evidence
with the potential to significantly contribute to the advancement of sports medicine and exercise
science research.
142

Results

Lactate concentration increases in the skin fluid and blood during incremental treadmill

running.

Following running habituation, the rats with inserted jugular catheters and single microneedles

were subjected to incremental running on the treadmill (Fig. 1*A-C*). The treadmill running speed

was gradually increased every 3 min until exhaustion (running time:34.4 $\pm$ 1.9 min; peak

speed:40.3 $\pm$ 1.6 m/min). Blood lactate concentrations significantly increased as running speed

increased, reaching 8.2 $\pm$ 0.8 mM at exhaustion (Fig. 2*A*). Following the exercise test, the blood

lactate concentrations rapidly decreased to baseline levels. Skin fluid lactate concentrations also

gradually increased during incremental treadmill running but did not decrease to baseline levels

even 30 min after the exercise test (Fig. 2*B*). We confirmed that any fluid leaked subcutaneously

or into the skin after removing the microneedle (Fig. 1*D*), and the amount of skin fluid collected

did not change among the time points (Fig. 2*C*).

Skin fluid lactate dynamics were significantly associated with blood lactate dynamics during

the exercise test but not afterward.

We then examined the relationship between lactate dynamics in blood and skin fluid (Fig. 3). A

very high correlation was observed between them in individual rats (Fig. 3*A-B*). The pooled

analysis also revealed a moderately significant correlation coefficient (Fig. 3*C*). However, these

significant correlations disappeared in the recovery phase following the exercise test in both

individual and pooled analyses (Fig. 3*D-F*).

Lactate threshold calculated from skin fluid lactate are closely associated with those

calculated from blood.

We attempted to calculate LT derived from skin fluid and examined its validity by performing a

correlation analysis with LT calculated from blood lactate dynamics. First, we performed LT

determination using the *V-slope* method[24], calculating the running speed and lactate levels in the

blood and skin fluid separatory (Fig. 4). The absolute and relative (% peak) running speed and

lactate levels corresponding to each LT calculated from blood and skin fluid were similar in

typical participants (Fig.4*A-D*). Moreover, no significant differences were observed in the

absolute and relative running speeds corresponding to the LT determined from the blood and skin

fluids (Fig. 5*A, C*). Furthermore, we confirmed that skin LT was significantly associated with

blood LT levels (Fig. 5*B, D*).

Discussion

This study tested the analytical capability of an ultra-minimally invasive technique for LT

measurement that harnesses skin interstitial lactate dynamics using a single microneedle perfusion

system (Fig. 1). Our results show that lactate levels in rat skin fluid collected using a single

microneedle perfusion system increased during incremental treadmill running, which was

significantly associated with an increase in blood lactate levels (Figs. 2-3). We also confirmed

that the LT determined from the blood and skin fluids were comparable and closely related (Figs.

4-5). As we discovered these findings in rats without sweat production in the epidermis, our results

strictly proved that skin ISF can be applied for LT prediction during exercise.

Validity of the rat model for incremental exercise to determine lactate threshold.

The incremental exercise gradually increased blood lactate concentration in rats, and the LT

calculated by *V-slope* method revealed 22.8 ± 0.8 m/min (absolute running speed) and 56.4 ±

2.2% (relative to the running speed at exhaustion) (Fig. 5*A, C*). We conducted an exercise test

using an experimental protocol modified from a previous study, and both the absolute and relative

LT were higher than those in the study[25]. Although our study used rats that conducted only running

habituation without any specific training, the relative LT was similar to that of untrained humans

(approx. 50-60% peak speed)[26]. Hence, the incremental exercise model in the present study seems

to be a reasonable experimental model for simulating LT determination using the human

incremental exercise test.

Lactate dynamics in the perfusate from the epidermal tissue are correlated with blood

during exercise.

Notably, skin fluid lactate concentration increases as well as blood lactate during the exercise test

(Fig. 2*A-B*). Moreover, lactate dynamics in skin fluid were significantly associated with those in

blood (Fig. 3*A-C*). These results are in line with those of a previous study that investigated the

relationship between blood and skin fluid lactate in resting mice after intraperitoneal injection of

lactate solution[20]. Since the components of ISF are comparable to the plasma[11,12], our results

indicate that the lactate levels in perfusate reflect the concentration in skin ISF, and the

microneedle perfusion system successfully collected skin ISF even during exercise. Thus, the

microneedle perfusion system might be effective in assessing lactate dynamics even during

exercise. Previous studies have demonstrated that lactate dynamics in the skin ISF during and

after exercise is delayed compared to that in the blood[18,19]. The authors noted that this was because

the wound spread and it took a long time to collect ISF [18,19]. The lactate concentrations in skin

fluid were related to those in blood at time points that were assumed to correspond theoretically,

suggesting that the measurements in this study were accomplished without a time lag. This may
be due to the microperfusion systems that consist of only a single microneedle and its extremely
low invasiveness (Fig. 1*B-D*). Moreover, as the lactate concentrations in perfusates collected in
vitro were directly quantified by the fluorescent staining method[20], we believe that the present
study strictly proved the relationship between blood and skin fluid lactate.

While a high positive correlation of lactate dynamics during exercise was observed between
blood and skin fluid within each individual rat, the correlation coefficient of the pooled analysis
decreased to moderate levels (but it was still statistically significant). The reason for this is
assumed to be that the slope of the regression line between blood and skin fluid lactate levels was
different for each rat. Although it may be possible to estimate the responsiveness of blood lactate
dynamics from skin fluid dynamics, absolute quantification of blood lactate concentration is
currently considered difficult based on skin fluid lactate dynamics.

The lactate thresholds calculated from the blood and skin fluid are closely correlated.
Both the absolute values of LT calculated from the blood and skin perfusate and their relative
values to peak speed were not significantly different (Fig. 5*A,C*). Furthermore, there was an
extremely strong positive correlation between the two (Fig. 5*B, D*). Accordingly, LT can be
estimated from the dynamics of skin ISF during exercise. A previous study pointed out that ISF
collected from the epidermal tissue may be contaminated by sweat[22]; therefore, we used rats that
did not produce sweat from the epidermis as subjects in this study. From these observations, this
study may be the first to directly prove the effectiveness of skin fluid LT as an alternative marker
of blood LT.

Furthermore, the skin fluid LT corresponded to 50-65 % of peak speed as well as blood LT, and
the results of the correlation analysis suggested that the individual differences were clearly
captured in that range (Fig. 5*D*). Exercise at moderate intensity including 50-65% of maximal
intensity, is widely used in training programs for amateur-level athletes and health-enhancing
exercisers[27], and it has been suggested that LT assessed by skin fluid may be applicable in many
sports settings. Future studies should examine the validity of the system used in this study for
highly athletic individuals with LT corresponding to 70-90% of maximal effort[26].

The lactate levels in the skin fluid did not decrease to baseline levels after the exercise test.
Interestingly, the increase in blood lactate concentration rapidly decreased to baseline levels (Fig.
2*A*), but those in the skin fluid decreased slowly and remained significantly higher 30 min after
the completion of the exercise test (Fig. 2*B*). Additionally, the significant relationship of the
lactate dynamics between the blood and skin fluid observed during exercise disappeared during
the recovery phase after exercise (Fig. 3*D-F*). Since Ming et al. confirmed a slow reduction in

skin ISF lactate concentrations after acute exercise in humans, our results support this response[17].
The authors speculated that the reason for this is that changes in components in the ISF
accompanied by a relative reduction in the perfusion rate of ISF to the skin after exercise might
be involved[17]. However, if the flow rate had been decreased, the skin fluid lactate concentrations
would have been diluted in saline circulating in the microneedle perfusion system. Thus, the
relatively high concentrations of skin fluid lactate after the exercise test; nevertheless, almost the
same amount of skin perfusate was collected, suggesting that clearance of lactate in the skin tissue
may have been slower than that in skeletal muscles rather than the reduction in perfusion rate of
ISF.

Growing evidence suggests that lactate induces several physiological adaptations via its
receptor (i.e., G-protein coupled receptor 81) in various tissues[28,29]. In addition, a recent study
suggested that regular exercise rejuvenated aging skin conditions[30]. These physiological
adaptations in skin tissue might be occurred by the lactate accumulation after exercise. While
further investigations should be required for testing the hypothesis, the microneedle perfusion
system used in this study might be a great tool for addressing it.

Perspectives
Collectively, our findings indicate that skin fluid lactate reflects the dynamics of blood lactate
during exercise and may be a novel valuable biomarker to estimate LT with high accuracy. This
suggests that LT could be assessed without the need for blood sampling by the single microneedle
perfusion system. However, since the dynamics following the exercise tests were different
between the blood and skin fluid, it seems difficult to assess the lactate clearance ability after
exercise and lactate clamp[31]. Moreover, we had measured lactate concentrations in perfusate,
collected in vitro, after the completion of exercise test. Thus, the current microneedle perfusion
system could not provide immediately and continuous feedback. In the future, further
miniaturization and development of sensors that enable immediate evaluation are needed for
human applications.

Nonetheless, it is noteworthy that the micro perfusion system is theoretically capable of
evaluating the dynamics of other metabolites and endocrine products (e.g., ketones, testosterone,
corticosterone, growth hormone, and so on) as well as lactate. Hence, the experimental systems
may be able to observe the dynamics of secretion and/or metabolites during exercise and daily
living including sleep, which may shed light on the development of mechanistic investigations in
the field of exercise physiology and biochemistry.

Limitations
This study has at least two limitations. First, the time resolution of the lactate assessment in skin

perfusate was low. The microneedle perfusion system used in this study required a flow rate of

5.0 μl/min in the tubes to ensure a stable ISF amount[20]. Moreover, a minimum of 20 μl of perfusate

was required for the quantification of lactate concentrations, so the collection frequency was less

than that of blood. Thus, insufficient skin perfusate collection, especially at the running speed

close to all-out, to collect enough amount to calculate LT2, the point at acceleration of lactate

accumulation[1,2]. LT2 is an index suggesting that the exercise intensity is physiologically high and

is particularly important in training elite athletes[1,2]. However, we have observed only a few cases

in where the rat could keep running even high-intensity were able to collect sufficient points of

skin perfusate to calculate LT2 (Supplemental Fig. 1). Accordingly, the time resolution should be

improved to enhance accurate measurements of skin fluid lactate concentrations at various

exercise intensities. Another issue is that this study did not perform the analysis of ventilatory

responses. Ventilatory threshold (VT) is also used to assess the physiological exercise intensity in

the training field, and a previous study in where the LT calculated using sweat lactate was

significantly associated with the VT in human subjects[9]. While it is classically known that LT and

VT approximate each other[1,2], inclusion of ventilatory gas analysis would further enhance the

physiological validity of LT determination using the skin fluid.

Conclusion

In this study, we tested the analytical capability of an ultra-minimally invasive ISF collecting

technique using a single microneedle perfusion system on LT measurements in exercising rats.

Our findings reveal the potential of skin interstitial dynamics to accurately assess LT during

exercise using a single microneedle. Currently, our group is challenging to develop a wearable

sensor capable of real-time measurement with the aim of applying this system to humans. The

implementation of this system may lead to the construction of various training programs and

evaluation of training effects through the evaluation of not only LT but also MLSS. We believe

that our findings will provide valuable insights for integrating the discoveries of exercise

physiology into real-world athletic and/or healthy practices.

Methods

Animals and running habituation.

Adult male Wistar rats (250–300 g) (SLC), housed and cared for in an animal facility, were fed a

standard pellet diet (MF; Oriental Yeast) and provided water ad libitum. Room temperature was

maintained between 22 and 24 °C under a 12-h light–12-h dark cycle (lights on, 0700 to 1900 h).

All experimental protocols were conducted in accordance with the guidelines of the University of

Tsukuba Animal Experiment Committee. The rats were habituated to running on a treadmill

(Natsume, Tokyo, Japan) for five sessions over 6 days for 30 min/day. The running speed was

gradually increased from 5 to 25 m/min[32-34].

Surgery to insert jugular catheter.

After habituation to treadmill running, the rats were anesthetized with isoflurane and a silicone

catheter was inserted into the jugular vein and fixed with a silk thread suture (32 mm). The

external distal end of the catheter was fixed at the nape of the neck. Subsequent experiments and

analyses were carried out 2 d after surgery[32-34].

Inserting the ultra-minimally invasive microneedle and incremental treadmill running test.

Two days after the surgery to insert catheter, the ultra-minimally invasive microneedle was

inserted into the back skin of the rats under anesthetization 30 min before start of the exercise test.

The specification of the microneedle was previously described[20]. To place the needle in the

subepidermal tissue, the needle was inserted along the skin surface (Fig. 1B). The inlet syringe

was filled with saline as the perfusate. Perfusion at a flow rate of 5 µl/min was started immediately

after the needle was secured to the rat's back. This flow rate was determined considering the

efficient collecting perfusate (skin ISF mixture saline)[20]. After the completion of inserting

microneedle into the back skin in the rats under anesthesia, the rats were moved on the treadmill

and ends of the tubes were connected to syringes attached to push (ESP-36, Eicom, Kyoto, Japan)

and pull (Nexus 3000, Chemyx Inc., USA) syringe pumps. 25-30 min after waking up, the animals

were further kept rest for 18 min before the running test (Fig. 1A). The initial treadmill velocity

was 5 m/min and was increased 5 m/min every 3 min until the speed at 20 m/min. Thereafter, the

treadmill speed was increased 2.5 m/min every 3 min until the point of exhaustion. Exhaustion

was considered to have occurred when the rat was unable to keep pace with the treadmill and

stayed on the grid positioned at the back of the treadmill for a period of 30 s despite being gently

pushed with sticks or breathed on. This protocol is modified from an incremental treadmill

exercise protocol to determine blood LT for rats[25]. Following the reached exhaustion, the rats were

kept rest for 35 min on the treadmill.

Blood and skin fluid sampling.

Blood samples were collected from the catheter every 3 min; however, it was collected 30 seconds

before each speed increase to determine blood LT during the exercise test. Due to the specification

of the microneedle perfusion system in this study, perfusing at 5 µl/min would result in a time

delay of 5 min compared to a real concentration change of skin ISF. Moreover, as the assay kit

for detecting the lactate concentration needed at least 20 µl of perfusate per sample, we collected

perfusate every 6 min, the collection was conducted 5 min after blood sampling from the catheter

(Fig. 1*C*). Theoretically, the components in the collected perfusate reflects the average values of

3 points of blood samples (5, 8, and 11 min before the skin fluid sampling). Therefore, the

collected perfusate at 35 min after the exhaustion reflects until 30 min after the exercise test. In a

previous study, it is suggested that the flow rate of perfusate could be changed during the recovery

phase after acute exercise[17]. Thus, we measured collected amount of the skin fluid in this study.

We also confirmed that any fluid leaked subcutaneously or into the skin after removing the

microneedle (Fig. 1D).

Lactate measurement and determination of lactate threshold.

Blood lactate levels were measured using an automated lactate analyzer (YSI 2500, YSI Inc.,

Yellow Springs, USA). Skin fluid lactate levels were determined using a commercial fluorescent

stain kit (L-Lactate Assay Kit, Cayman Chemical Company, USA), according to the

manufacturer's instructions. Fluorometric determination was performed using a fluorescence

microplate reader (Varioskan LUX multimode microplate reader, Thermo Fisher Scientific, USA).

LT was determined using the V-slope method[24], calculating the running speed corresponding to

the intersection of two linear regression lines derived separately from the data points of the

running speed and the lactate level.

Statistical analysis.

One-way analysis of variance (ANOVA) was performed for the time course changes in blood and

skin fluid lactate concentrations, and skin fluid sample amount. Pearson's correlation analysis

between the lactate concentrations in blood and skin samples was conducted separately during

and following the exercise test. As mentioned above, the collected skin fluid might be reflected

the 3 points of blood samples due to the methodological specification; therefore, we conducted

the correlation analysis using the average concentration of 3 blood lactate per 1 skin fluid lactate.

We also compared the absolute and relative (% peak) treadmill speed corresponding to the LT

calculated from the blood and skin by paired t-test and examined their relationships between the

LT calculated from blood and skin fluid by Pearson's test. All statistical analysis was performed

using the GraphPad Prism ver. 10.0.3 (MDF, Japan).

Data availability

The data that support the findings of this study are available from the corresponding author upon

reasonable request.

Acknowledgements
This research was supported in part by the Leading-edge Research Project for Sports Medicine
and Science (LRP) by Japan Sports Agency to T.M., Grant-in-Aid for Scientific Research (B) by
JSPS KAKENHI (19H03994 and 22H03478) to T.M., and Fusion Oriented Research for
disruptive Science and Technology (FOREST) by Japan Science and Technology Agency (JST)
to T.M. (JPMJFR205M) and N.T.(JPMJFR215).
401

Author contributions
T.M. and N.T. conceived and designed the study. N.T. prepared a microneedle for the collection
of subepidermal interstitial fluid. D.F., S.D., K.S., and T.M. collected data. D.F. and S.D.
determined the blood and skin lactate concentrations. K.S. measured skin fluid amount. D.F. and
T.M. conducted data analysis and interpreted the data. S.D., T.M., and D.F. drafted the manuscript.
All authors edited and revised the manuscript, and read and approved the final version of this
manuscript.
409

410 **Competing interests**
411 The authors declare no competing interests.
412

References

1) Poole D, Rossiter HB, Brooks GA, Gladden LB. The anaerobic threshold: 50+ years of

controversy. J Physiol. 2021, 599: 737-767.

2) Jaminick NA, Pettitt RW, Granata C., Pyne DB, Bishop DJ. An examination and critique of

current methods to determine exercise intensity. Sports Med. 2020, 50: 1729-1756.

3) Smith EW, Skelton MS, Kremer DE, Pascoe DD, Gladden LB. Lactate distribution in the blood

during progressive exercise. Med Sci Sport Exerc. 1997, 29: 654-660.

4) Goodwin ML., Harris JE, Hernández A, Gladden LB. Blood lactate measurements and analysis

during exercise: a guide for clinicians. J Diabetes Sci Technol. 2007, 1: 558-569.

5) Robergs RA, Chwalbinska-Moneta J, Mitchell JB, Pascoe DD, Houmard J, Costill DL. Blood

lactate threshold differences between arterialized and venous blood. Int J Sports Med. 1990, 11:

446-451.

6) Kim J, Valdés-Ramírez G, Bandodkar AJ, Jia W, Martinez AG, Ramírez J, Mercier P, Wang J.

Non-invasive mouthguard biosensor for continuous salivary monitoring of metabolites. Analyst.

2014, 139: 1632-1636.

7) Jia W, Bandodkar AJ, Valdés-Ramírez G, Windmiller JR, Yang Z, Ramírez J, Chan G, Wang J.

Electrochemical tattoo biosensors for real-time noninvasive lactate monitoring in human

perspiration. Anal Chem. 2013. 85: 6553-6560.

8) He W, Wang C, Wang H, Jian M, Lu W, Liang X, Zhang X, Yang F, Zhang Y. Integrated textile

sensor patch for real-time and multiplex sweat analysis. Sci Adv. 2019, 5: eaax0649.

9) Seki Y, Nakashima D, Shiraishi Y, Ryuzaki T, Ikura H, Miura K, Suzuki M, Watanabe T, Nagura

T, Matsumoto M, Nakamura M, Sato K, Fukuda K, Katsumata Y. A novel device for detecting

anaerobic threshold using sweat lactate during exercise. Sci Rep. 2021, 11: 4929.

10) Maeda Y, Okawara H, Sawada T, Nakashima D, Nagahata J, Fujitsuka H, Ikeda K, Hoshino

S, Kobari Y, Katsumata Y, Nakamura M, Nagura T. Implications of the onset of sweating on the

sweat lactate threshold. Sensors. 2023, 23: 3378.

11) Scallan J, Huxley VH, Korthuis RJ. Capillary fluid exchange: regulation, functions, and
pathology. Chaptor 2 The Interstitium. In Integrated Systems Physiology: from Molecule to
Function to Disease (Ganger DN and Granger JP, eds), Morgan & Vlaypool Life Sciences. 2010.
452
453   12) Miller PR, Taylor RM, Tran BQ, Boyd G, Glaros T, Chavez VH, Krishnakumar R, Sinha A,
454   Poorey K, Williams KP, Branda SS, Baca JT, Polsky R. Extraction and biomolecular analysis of
455   dermal interstial fluid collected with hollow microneedles. Commun Biol, 2018, 1: 173.
456
457   13) Petersen LJ. Interstitial lactate levels in human skin at rest and during an oral glucose load: a
458   microdialysis study. Clin Physiol. 1999, 19: 246-250.
459
460   14) Enderle B, Moser I, Kannan C, Schwab KO, Urban G. Interstitial glucose and lactate levels
461   are inversely correlated with the body mass index: need for in vivo calibration of glucose sensor
462   results with blood values in obese patients. J Diabetes Sci Technol. 2018, 12: 341-348.
463
464   15) Wang Y, Aursi IR, Wang Z, Derry C, Tang XS. Towards a transdermal membrane biosensor
465   for the detection of lactate in body fluids. Sens. Actuators B Chem. 2020, 308: 127645.
466
467   16) Chien MN, Fan SH, Huang CH, Wu CC, Huang JT. Continuous lactate monitoring system
468   based on percutaneous microneedle array. Sensors. 2022, 22:1 468.
469
470   17) Ming DK, Jangam S, Gowers SAN, Wilson R, Freeman DME, Boutelle MG, Cass AEG,
471   O'Hare D, Holmes AH. Real-time continuous measurement of lactate through a minimally
472   invasive microneedle patch: a phase I clinical study. BMJ Innov. 2022, 8:87-94.
473
474   18) de Boer J, Plijter-Groendijk H, Visser KR, Mook GA, Korf J. Continuous monitoring of
475   lactate during exercise in humans using subcutaneous and transcutaneous microdialysis. Eur J
476   Appl Physiol Occup Physiol. 1994, 69: 281-286.
477
478   19) Ellmerer M, Schaupp L, Trajanoski Z, Jobst G, Moser I, Urban G, Skrabal F, Wach P.
479   Continuous measurement of subcutaneous lactate concentration during exercise by combining
480   open-flow microperfusion and thin-film lactate sensors. Biosens Bioelectron. 1998, 13:1007-
481   1013.
482
20) Tsuruoka N, Ishii K, Matsunaga T, Nagatomi R, Haga Y. Lactate and glucose measurement in
subepidermal tissue using minimally invasive microperfusion needle. Biomed Microdevices.

2016, 18: 19.

21) Rathee K, Dhull V, Dhull R, Singh S. Biosensors based on electrochemical lactate detection:

A comprehensive review. Biochem Biophys Rep. 2015, 5: 35-34.

22) Kim J, Sempionatto JR, Imani S, Hartel MC, Barfdokht A, Tang G, Campbell AS, Mercier PP,

Wang J. Simultaneous monitoring of sweat and interstitial fluid using a single wearable biosensor

platform. Adv Sci. 2018, 5: 1800880.

23) Xuan X, Chen C, Molinero-Fernandez A, Ekelund E, Cardinale D, Swarén M, Wedholm L,

Cuartero M, Crespo GA. Fully integrated wearable device for continuous sweat lactate monitoring

in sports. ACS Sens. 2023, 8: 2401-2409.

24) Beaver WL, Wasserman K, Whipp BJ. A new method for detecting anaerobic threshold by

gas exchange. J Appl Physiol. 1986, 60: 2020-2027.

25) Soya H, Mukai A, Deocaris CC, Ohiwa N, Chang H, Nishijima T, Fujikawa T, Togashi K,

Saito T. Threshold-like pattern of neuronal activation in the hypothalamus during treadmill

running: establishment of a minimum running stress (MRS) rat model. Neurosci Res. 2007, 58:

341-348.

26) Messonnier LA, Emhoff CA, Fattor JA, Horning MA, Carlson TJ, Brooks GA. Lactate

kinetics at the lactate threshold in trained and untrained men. J Appl Physiol. 2013, 114: 1593-

1602.

27) Garber GE, Blissmer B, Deschenes MR, Franklin BA, Lamonte MJ, Lee IM, Nieman DC,

Swain DP, American College of Sports Medicine. American College of Sports Medicine position

stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory,

musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing

exercise. Med Sci Sports Exerc. 2011, 43: 1334-1359.

28) Morland C, Andersson KA, Haugen ØP, Hadzic A, Kleppa L, Gille A, Rinholm JE, Palibrk V,

Diget EH, Kennedy LH, Stølen T, Hennestad E, Moldestad O, Cai Y, Puchades M, Offermanns S,

Vervaeke K, Bjørås M, Wisløff U, Storm-Mathisen J, Bergersen LH. Exercise induces cerebral

VEGF and angiogenesis via the lactate receptor HCAR1. Nat Commun. 2017, 8: 15557.

29) Ohno Y, Nakatani M, Ito T, Natsui Y, Ando K, Suda Y, Ohashi K, Yokoyama S, Goto K.
Activation of lactate receptor positively regulates skeletal muscle mass in mice. Physiol Res. 2023,
72: 465-473.
524
525   30) Nishikiori S, Yasuda J, Murata K, Takegaki J, Harada Y, Shirai Y, Fujita S. Resistance training
526   rejuvenates aging skin by reducing circulating inflammatory factors and enhancing dermal
527   extracellular matrices. Sci Rep. 2023, 13: 10214.
528
529   31) Emholf CAW, Messonier LA. Concepts of lactate metabolic clearance rate and lactate clamp
530   for metabolic inquiry: a mini-review. Nutrients. 2023, 15: 3213
531
532   32) Matsui T, Soya S, Okamoto M, Ichitani Y, Kawanaka K, Soya H. Brain glycogen decreases
533   during prolonged exercise. J Physiol. 2011, 589: 3383-3393.
534
535   33) Matsui T, Ishikawa T, Ito H, Okamoto M, Inoue K, Lee MC, Fujikawa T, Ichitani Y, Kawanaka
536   K, Soya H. Brain glycogen supercompensation following exhaustive exercise. J Physiol. 2012,
537   590: 607-616.
538
539   34) Matsui T, Omuro H, Liu YF, Soya M, Shima T, McEwen BS, Soya H. Astrocytic glycogen-
540   derived lactate fuels the brain during exhaustive exercise to maintain endurance capacity. Proc
541   Natl Acad Sci USA. 2017, 114: 6358-6363.

542 **Figures**

Fig. 1. The overview of experimental procedures using a jugular catheter and an ultra-
minimally microneedle perfusion system.

(*A*) The experimental settings of rats with inserted catheter and microneedle perfusion system
during exercise and experimental protocol of incremental running to determine lactate threshold
for rats. (*B*) The specification of microneedle. (*C*) The part of skin with the microneedle inserted
and its fixation. (*D*) The part of skin after removing microneedle. (*C* and *D*) indicate no significant
inflammation and bleeding, suggesting that the harmful effect of inserting the single microneedle
perfusion system may be minimum.

Fig. 2. Lactate concentration increases in the skin fluid and blood during incremental treadmill running. Time course changes in blood (*A*) and skin fluid (*B*) lactate concentrations and skin fluid amount (*C*) pre, during, and post incremental running test. Each plot and shaded regions represent mean values and continuous standard error bands, respectively. *$p < 0.05$ vs. 0 min (immediately before the start of the exercise test).

Fig. 3. Skin fluid lactate dynamics were significantly associated with blood lactate dynamics during the exercise test but not afterward. The correlation coefficient between the blood and skin fluid lactate concentrations in typical case A and B, and pooled data of all cases during (A-C) and following (*D-F*) exercise. Shaded regions represent 95% simultaneous confidence bands. *p < 0.05, n.s., not significant.

Fig. 4. The lactate dynamics in skin fluid can determine lactate threshold as well as blood.
Representative LT profiles calculated by the dynamics from blood and skin fluid in typical cases
A (panel *A, B*) and B (panel *C, D*). LT in each graph was calculated by *V-slop* method (Beaver et
al., 1986).

Fig. 5. Lactate threshold calculated from skin fluid lactate are closely associated with those calculated from blood. The absolute running speed corresponding to the LT calculated from blood and skin fluid (A) and their relationship (B). The running speed (C) and its relationship (D) relative to peak speed during incremental exercise corresponding to LT calculated from blood and skin fluid, respectively. The values of bar graph (*A, C*) represent means and individual plots. Shaded regions represent 95% simultaneous confidence bands. n.s., not significant, *p < 0.05.

704 **Supplemental Fig. 1 Representative case detecting lactate threshold 2 from blood and skin**

705 **fluid lactate dynamics during the incremental exercise test.**