1 **Exercise intensity-dependent lactate elevation in skin interstitial fluid and its long-**
2 **lasting effect**
3

4 Shohei Dobashi[1,2,5], Daisuke Funabashi[1,3,5], Kazuki Sameshima[1], Noriko Tsuruoka[4],
5 Takashi Matsui[1,2,3,5]∗
6
[1] Exercise Biochemistry & Sport Neurobiology Division, Institute of Health and Sport
Sciences, University of Tsukuba, Tsukuba, Japan
[2] R&D Center for Lifestyle Innovation, University of Tsukuba, Tsukuba, Japan
[3] Advanced Research Initiative for Human High Performance, University of Tsukuba,
Tsukuba, Japan
[4] Graduate School of Engineering, Tohoku University, Sendai, Japan.
[5] These authors contributed equally to this work.
14
Correspondence:
Takashi Matsui Ph.D.
Assistant Professor
Exercise Biochemistry & Sport Neurobiology Division,
Institute of Health and Sport Sciences, University of Tsukuba
E-mail: matsui.takashi.ga@u.tsukuba.ac.jp Tel: +81-29-853-2750

Abstract

Regular exercise promotes various anti-ageing adaptations in skin tissue. Although the underlying mechanisms of that might associate to the acute exercise-induced lactate signaling in the skin, it remains uncertain the profile of skin interstitial fluid (ISF) lactate dynamics during and following acute exercise. Here, we investigated whether the skin ISF lactate level increases in association with blood lactate during acute incremental exercise using a single microneedle perfusion system. The rats were acclimated to treadmill running exercise and underwent external jugular vein cannulation. Following comprehensive recovery, a 1 mm single microneedle was implanted into the back skin. Skin ISF lactate progressively increased in tandem with blood lactate during the incremental exercise but did not decrease to baseline levels until 30 minutes following the exercise unlike blood. Moreover, lactate threshold (LT), is a crucial marker of athletic aerobic performance during acute exercise, extrapolated from skin ISF showed significant alignment with blood LT. Our findings reveal that the skin ISF lactate increases associated with blood lactate and the long-lasting elevation state following acute exercise. Moreover, lactate dynamics in skin ISF can predict LT. These findings would be a milestone for elucidating regular exercise-induced physiological adaptations in the skin and evaluating athletic performance.

Key words: Lactic acid, Treadmill running, Microneedle perfusion system, Lactate threshold

Graphical Abstract

Lactate levels in rat skin interstitial fluid (ISF) increased in association with blood lactate
during acute incremental exercise, but the increase in skin ISF lactate lasted 30 min
following acute exercise. Lactate threshold (LT), a crucial marker of aerobic athletic
performance, can be predicted from skin fluid dynamics during acute exercise. These
findings would be a milestone for elucidating regular exercise-induced physiological
adaptations in the skin and evaluating athletic performance.

Introduction

The skin plays a pivotal role, such as safeguarding the body from external environmental threats, regulating body temperature, maintaining hydration, and houses sensory receptors (1); therefore, the maintenance and/or enhancement in skin tissue function are important to achieve a longer healthy lifespan. Regular exercise is believed as real polypill, which has various benefits for physical and psychological health, including skin health (2). For instance, a recent study demonstrated that regular aerobic and resistance exercises rejuvenated aging skin by reducing inflammatory factors and enhancing dermal extracellular matrices (3).

Although the underlying mechanisms of chronic exercise-induced benefits on skin health remains unknown, "lactate" is sought as one of candidates. Lactate is known not only as the end product of glycolysis, but also as an exercise-induced bioactive compound ("exerkine") (2, 4). As the exercise intensity increases, lactate is released into the bloodstream, primarily from the exercising muscles, with a concomitant enhancement in physiological adaptations of the various tissues (5). Moreover, previous studies have reported that acute exercise-induced lactate is taken up via monocarboxylate transporters (MCTs) or binds to lactate receptors (G protein-coupled receptors, GPR81) in skeletal muscle and brain, the repetition of which results in chronic beneficial adaptations (2,6). MCTs and GPR81 are also expressed in skin (7,8) and lactate promotes mitochondrial biogenesis and attenuates inflammation there (9). Hence, the chronic exercise-induced skin benefits may also be due to the accumulation of blood-derived lactate transported to the skin, as well as to other tissues. As the constituents in blood diffuse into skin interstitial fluid (ISF) (10), we tempted to hypothesize that the skin ISF lactate level during acute exercise increases in association with blood lactate.

Recent technologies have developed to collect skin ISF and the validity of ISF lactate has been investigated (10). Among them, a novel minimally invasive microneedle perfusion system by Tsuruoka et al. appears promising, which demonstrates that the lactate dynamics in skin ISF collected using the microneedle system are consistent with those in the blood in mice after intraperitoneal lactate administration (11). Since the lactate dynamics in skin ISF were also associated with the dynamics in blood above the onset of blood lactate accumulation (OBLA) (4 mM) at this time, the microneedle perfusion system may be able to monitor dynamics of skin ISF lactate during acute exercise.

Here, we tested the hypothesis that the skin ISF lactate level increases in association with blood lactate during acute incremental exercise using the minimally

invasive single microneedle perfusion system. If the results are as hypothesized, lactate dynamics in the skin ISF would allow us to predict lactate threshold (LT), which are known as universal indices that predict athletic aerobic performance and used to evaluate the exercise intensity and assessment of chronic exercise impact (12). Thus, we also aimed to examine the validity of LT determination derived from skin ISF lactate dynamics during exercise. To test this, we used rats, devoid of sweat production, because a previous study pointed out that ISF collected from the epidermal tissue may be contaminated by sweat which can interfere with adequate assessment in skin ISF lactate dynamics (13). Then, the rats were cannulated with external jugular vein and the microneedle was implanted into the back skin. This allowed for the simultaneous measurement of lactate concentrations in the blood and skin ISF during the incremental treadmill exercise.

The results of this study demonstrate that skin ISF lactate concentrations increase in association with blood lactate levels during acute incremental exercise. However, the increased skin lactate concentrations were not completely decreased to baseline levels even 30 minutes after the exercise test. We also confirmed that LT calculated from skin fluid lactate is closely related to those calculated from blood. These findings provide new insights into assessing chronic exercise-induced physiological adaptations of the skin and athletic performance using lactate dynamics in skin ISF.

Methods

Animals and running habituation

Adult male Wistar rats (n = 6, 250–300 g) (SLC), housed and cared for in an animal facility, were fed a standard pellet diet (MF; Oriental Yeast) and provided water ad libitum. Room temperature was maintained between 22 and 24 °C under a 12-h light–12-h dark cycle (lights on, 0700 to 1900 h). All experimental protocols were conducted in accordance with the guidelines of the University of Tsukuba Animal Experiment Committee. The rats were habituated to running on a treadmill (Natsume, Tokyo, Japan) for five sessions over 6 days for 30 min/day. The running speed was gradually increased from 5 to 25 m/min (14-16).

Surgery to insert jugular catheter

After habituation to treadmill running, the rats were anesthetized with isoflurane and a silicone catheter was inserted into the jugular vein and fixed with a silk thread suture (32 mm). The external distal end of the catheter was fixed at the nape. Subsequent experiments and analyses were carried out 2 d after surgery (14-16).

Inserting the minimally invasive microneedle and incremental treadmill running test.

Two days after the surgery to insert catheter, the ultra-minimally invasive microneedle was inserted into the back skin of the rats under anesthetization 30 min before start of the exercise test (Figure 1A). This single microneedle system has an outer diameter of approximately 0.2 mm and a length of 1-2 mm can collect ISF through saline perfusion (Figure 1B) (11). To place the needle in the subepidermal tissue, the needle was inserted along the skin surface (Figure 1C). The inlet syringe was filled with saline as the perfusate. Perfusion at a flow rate of 5 µl/min was started immediately after the needle was secured to the rat's back. This flow rate was determined considering the efficient collecting perfusate (skin ISF mixture saline) (11).

After the completion of inserting microneedle into the back skin in the rats under anesthesia, the rats were moved on the treadmill and ends of the tubes were connected to syringes attached to push (ESP-36, Eicom, Kyoto, Japan) and pull (Nexus 3000, Chemyx Inc., USA) syringe pumps. 25-30 min after waking up, the animals were further kept rest for 18 min before the running test (Figure 1A). The initial treadmill velocity was 5 m/min and was increased 5 m/min every 3 min until the speed at 20 m/min. Thereafter, the treadmill speed was increased 2.5 m/min every 3 min until the point of exhaustion. Exhaustion was considered to have occurred when the rat was unable to keep pace with the treadmill and stayed on the grid positioned at the back of the treadmill for a period of 30 s despite being gently pushed with sticks or breathed on. This protocol is modified from an incremental treadmill exercise protocol to determine blood LT for rats (17). Following the reached exhaustion, the rats were kept rest for 35 min on the treadmill.

Blood and skin fluid sampling

Blood samples were collected from the catheter every 3 min; however, it was collected 30 seconds before each speed increase to determine blood LT during the exercise test (Figure 1A). Due to the specification of the microneedle perfusion system in this study, perfusing at 5 µl/min would result in a time delay of 5 min compared to a real concentration change of skin ISF. Moreover, as the assay kit for detecting the lactate concentration needed at least 20 µl of perfusate per sample, we collected perfusate every 6 min, the collection was conducted 5 min after blood sampling from the catheter (Figure 1A). Theoretically, the components in the collected perfusate reflects the average values of 3 points of blood samples (5, 8, and 11 min before the skin fluid sampling). Therefore, the collected perfusate at 35 min after the exhaustion reflects until 30 min after the

exercise test. In a previous study, it is suggested that the flow rate of perfusate could be
changed during the recovery phase after acute exercise (18). Thus, we measured collected
amount of the skin fluid in this study. We also confirmed that any fluid leaked
subcutaneously or into the skin after removing the microneedle (Figure 1*D*).

Lactate measurement and determination of lactate threshold
Immediately after each blood sampling during the exercise test, lactate levels in the whole
blood were measured using an automated lactate analyzer (YSI 2500, YSI Inc., Yellow
Springs, USA). Skin fluid lactate levels were determined using a commercial fluorescent
stain kit (L-Lactate Assay Kit, Cayman Chemical Company, USA), according to the
manufacturer's instructions. Fluorometric determination was performed using a
fluorescence microplate reader (Varioskan LUX multimode microplate reader, Thermo
Fisher Scientific, USA).
171   We evaluated not only lactate dynamics in skin ISF with blood, but also
determined lactate threshold (LT) using the modified *V-slope* method (19) in the previous
study (17), calculating the running speed corresponding to the intersection of two linear
regression lines derived separately from the data points of the running speed and the
lactate level.

Statistical analysis
One-way analysis of variance (ANOVA) was performed for the time course changes in
blood and skin fluid lactate concentrations, and skin fluid sample amount. Pearson's
correlation analysis between the lactate concentrations in blood and skin samples was
conducted separately during and following the exercise test. As mentioned above, the
collected skin fluid might be reflected the 3 points of blood samples due to the
methodological specification; therefore, we conducted the correlation analysis using the
average concentration of 3 blood lactate per 1 skin fluid lactate (11). We also compared
the absolute and relative (% peak) treadmill speed corresponding to the LT calculated
from the blood and skin by paired t-test and examined their relationships between the LT
calculated from blood and skin fluid by Pearson's test. All statistical analysis was
performed using the GraphPad Prism ver. 10.0.3 (MDF, Japan).

Results

Lactate concentration increases in the skin fluid and blood during incremental
treadmill running

Following the treadmill running habituation, the rats with inserted jugular catheters and
single microneedles were subjected to incremental running on the treadmill (Figure 1).
The treadmill running speed was gradually increased every 3 min until exhaustion
(running time: 34.4 ± 1.9 min; peak speed: 40.3 ± 1.6 m/min). Blood lactate
concentrations significantly increased as running speed increased, reaching 8.2 ± 0.8 mM
at exhaustion (Figure 2A). Skin fluid lactate concentrations gradually increased during
incremental treadmill running associated with blood but did not decrease to baseline
levels even 30 min after the exercise test like as blood (Figure 2A, B). The collected
amount of the skin fluid was similar throughout the experiment (Figure 2C).

Skin fluid lactate dynamics are significantly associated with blood lactate dynamics
during the exercise test but not afterward
We then examined the relationship between lactate dynamics in blood and skin fluid
(Figure 3). A very high correlation was observed between them during exercise in
individual rats (Figure 3A, B). The pooled analysis also revealed a moderately significant
correlation coefficient (Figure 3C). However, these significant correlations disappeared
in the recovery phase following the exercise test in both individual and pooled analyses
(Figure 3D-F).

Lactate threshold calculated from skin fluid lactate are closely related with those
calculated from blood
We also attempted to calculate LT derived from skin fluid and examined its relationship
with LT calculated from blood lactate dynamics by performing a correlation analysis
(Figure 4). The absolute and relative (% peak) running speed and lactate levels
corresponding to each LT calculated from blood and skin fluid were similar in typical
participant (Figure 4A, B). Moreover, no significant differences were observed in the
absolute and relative running speeds corresponding to the LT determined from the blood
and skin fluids (Figure 4C, E). Furthermore, we confirmed that skin LT was significantly
associated with blood LT levels (Figure 4D, F).

Discussion
We tested the hypothesis that the skin ISF lactate level increases in association with blood
lactate during acute exercise using the minimally invasive single microneedle perfusion
system, and the results were in line with that (Figure 2). However, the increase in skin
ISF lactate did not decrease to baseline levels even 30 min after the exercise like as blood
lactate (Figure 2). The different linkage of lactate dynamics in blood and skin ISF during

and after the exercise test was also observed in the correlation analysis (Figure 3).

Additionally, we confirmed that the LT determined from the blood and skin fluids were

comparable and closely related (Figure 4). As we discovered these findings in rats without

sweat production in the epidermis to prevent contamination, our study provides the first

evidence that unveils the skin ISF lactate dynamics response to acute exercise.

Lactate dynamics in the perfusate from the epidermal tissue are correlated with

blood during exercise

In this study, skin ISF lactate concentrations progressively increased in tandem with blood

lactate during the incremental exercise (Figure 2*A, B*). As we used the rats and the lactate

concentrations in perfusates collected in vitro were directly quantified by the fluorescent

staining method, our present study strictly proved that skin ISF lactate concentrations

itself increases as well as blood lactate. While the previous studies have pointed out that

acute dynamics of lactate concentrations in ISF have delayed from the dynamics of that

in blood (20), we observed that lactate dynamics in skin ISF during the acute exercise test

could measure without the time delay from that in blood in this study. In this regard, a

previous study demonstrated that acute exercise increased skin blood flow (21). Although

further study should be required to measure skin blood flow, we speculate that the acute

increase in skin blood flow due to exercise may facilitate the diffusion of water-soluble

components from the blood to the skin, thereby allowing continuous lactate measurement

without time delay.

Moreover, the lactate dynamics in skin fluid were significantly associated with

those in blood (Figure 3*A-C*). These results are in line with those of a previous study that

investigated the relationship between blood and skin fluid lactate in resting mice after

intraperitoneal injection of lactate solution (11). While our study could not properly

distinguish whether the skin ISF lactate was derived from the blood or the skin itself, the

significant relationship between lactate dynamics in blood and skin fluids suggests that

the acute exercise-induced increase in skin ISF lactate may have been derived the blood

rather than in the skin. A previous study demonstrated that acute hypoxic exposure

increased skin ISF lactate concentrations in association with the blood in rats (22), which

supports our present findings. However, a high positive correlation of lactate dynamics

during the exercise test was observed between blood and skin fluid within each individual

rat (Figure 3*A-B*), the correlation coefficient of pooled analysis decreased to moderate

levels (but it was still statistically significant) (Figure 3*C*). The reason for this presumably

because the slope of the regression line of lactate dynamics between the blood and the

skin ISF differed among the rats due to individual differences in the diffusion rate of the

components from blood to skin ISF and skin blood flow rate during acute exercise. This
may affect the individual variability in skin adaptation to chronic exercise. Further studies
should be needed to clarify the mechanisms.

The lactate thresholds calculated from the blood and skin fluid are closely correlated
We also observed that the LT calculated from lactate dynamics in skin ISF was
corresponded to 50-65 % of peak speed as well as blood LT, and the results of the
correlation analysis suggested that the individual differences were clearly captured in that
range (Figure 4*F*). Although the microneedle perfusion system does not aim to estimate
absolute blood lactate concentrations from skin ISF lactate concentrations, our results
indicate that the lactate dynamics in skin ISF during acute exercise can predict LT. Our
study used rats that conducted only running habituation without any specific training, but
the relative LT was similar to that of untrained humans (approx. 50-60% peak speed) (23).
Hence, the incremental exercise model in the present study seems to be a reasonable
experimental model for simulating LT determination using the human incremental
exercise test, and our results can be applied for human. Additionally, the exercise-
intensity corresponding to the calculated skin LT is widely used in training programs for
amateur-level athletes and health-enhancing exercisers (24). Steps still need to be taken
toward practical application in humans, but LT assessed by skin fluid might be applicable
in many sports settings in the future.

The lactate levels in the skin fluid did not decrease to baseline levels after the exercise
test
Interestingly, the increase in blood lactate concentrations rapidly decreased to baseline
levels (Figure 2*A*), but those in the skin fluid decreased slowly and remained significantly
higher even 30 min after the completion of the exercise test (Figure 2*C*). Additionally, the
significant relationship of the lactate dynamics between the blood and skin fluid was
observed during the exercise (Figure 3*A-C*) but disappeared during the recovery phase
after the exercise (Figure 3*D-F*). Although it was assumed the possibility that the changes
in collected amount of perfusate due to the reduction in perfusion rate of ISF following
acute exercise affected lactate concentrations in the skin ISF, we observed that almost the
same amount of skin perfusate was collected throughout the experiment (Figure 2*C*). This
result suggests that clearance of lactate in the skin tissue may have been slower than that
in tissues (mainly skeletal muscle) and the blood rather than the reduction in perfusion
rate of ISF. This explanation should be supported by a rapid post-exercise decrease in skin
blood flow (21), which affects the clearance of water-soluble substances from the skin

into the blood (10). In any case, our results show a unique response in skin, where high
concentrations of lactate last for a long-time following acute exercise.
304

New hypotheses for lactate-mediated regular exercise-induced skin physiological adaptation

The present results of skin ISF lactate dynamics in this study can provide new hypotheses.
Growing evidence suggests that chronic exercise improved skin conditions (2,3), and
lactate acts as a signaling molecule that interacts with mitochondrial biogenesis and
turnover and tunes the ability of cells to respond to stressors (9). These findings and the
present results suggest that the beneficial physiological adaptation of skin tissue to
chronic exercise might be due to the accumulated effects of acute exercise-induced long-
lasting lactate elevation in the skin ISF. It has been demonstrated that regular higher-
intensity exercise to increase blood lactate production elicited beneficial adaptations in
the skeletal muscles and brain (4,5). Furthermore, the previous study that reported
improved skin conditions with chronic resistance exercise also employed an intensity of
exercise to increase blood lactate levels (3,25). From the fact that the lactate levels in the
skin ISF increased in tandem with the blood during incremental exercise and they were
closely related, regular higher-intensity exercise can be effective for inducing chronic
exercise-induced skin adaptations as well as other tissue of the body. The microneedle
perfusion system used in this study should be a great tool to test the new hypotheses.
322

Perspectives

The microneedle perfusion system used in this study required a flow rate of 5.0 μl/min in
the tubes to ensure a stable ISF amount (11). Moreover, a minimum of 20 μl of perfusate
was required for the quantification of lactate concentrations by a fluorescent staining
method, so the collection frequency of skin fluid had to be less than that of blood.
Although our present results clearly proved that skin ISF lactate increases in tandem with
blood lactate during the incremental running test, the relatively low time resolution of
determination of skin ISF lactate concentrations using this system might make our
interpretation difficult. In the future, a sensor that can continuously and immediately
measure lactate in the perfusate should be incorporated with the microneedle to
investigate the lactate dynamics in skin ISF during exercise with great precision.
Moreover, the futured microneedle perfusion system should also be required to implement
the technology to prevent sweat contamination for human application. The improved
system will advance our understanding of lactate dynamics in skin during exercise and
the resulting physiological responses. The system can also detect maximal lactate steady

state (MLSS), which are used for assessing aerobic endurance capacity (26), as well as
LT in real time. In this regard, we believe that our present findings to confirm the validity
of LT determination derived from skin ISF dynamics will be a milestone toward
realization of such innovative systems.
342

Conclusion

Our findings reveal that skin ISF lactate concentrations increase in association with blood
lactate levels during acute incremental exercise. However, the lactate levels in skin ISF,
rather than blood, shows long-lasting elevation following acute exercise, suggesting a
potential mechanism of chronic exercise-induced beneficial adaptations of the skin.
Moreover, LT can be predicted from skin fluid dynamics during acute exercise without
blood sampling. Collectively, we provide new insights for assessing chronic exercise-
induced physiological adaptations of the skin and athletic performance using lactate
dynamics in skin ISF.
352

Author contributions

T.M. and N.T. conceived and designed the study. N.T. prepared a microneedle for the
collection of subepidermal interstitial fluid. D.F., S.D., K.S., and T.M. collected data. D.F.
and S.D. determined the blood and skin lactate concentrations. K.S. measured skin fluid
amount. D.F. and T.M. conducted data analysis and interpreted the data. S.D., T.M., and
D.F. drafted the manuscript. All authors edited and revised the manuscript and read and
approved the final version of this manuscript.
360

Acknowledgements

362 This research was supported in part by the Leading-edge Research Project for Sports
363 Medicine and Science (LRP) by Japan Sports Agency to T.M., Grant-in-Aid for Scientific
364 Research (B) by JSPS KAKENHI (19H03994 and 22H03478) to T.M., and Fusion
365 Oriented Research for disruptive Science and Technology (FOREST) by Japan Science
366 and Technology Agency (JST) to T.M. (JPMJFR205M) and N.T.(JPMJFR215).
367

Competing interests

369 The authors declare that no competing interests exist.
370

Data availability

372 The data that support the findings of this study are available from the corresponding
373 author upon reasonable request.

References

1. Sorg H. Tilkorn DJ, Hager S, Hauser J, Mirastschijski U. Skin wound healing: An update on the current knowledge and concepts. *Eur Surg Res*. 2017;58(1-2):81-94.

2. Brooks GA, Osmond AD, Arevalo JA, Duong JJ, Curl CC, Moreno-Santillan DD, Leija RG. Lactate as a myokine and exerkine: drivers and signals of physiology and metabolism. *J Appl Physiol*. 2023;134(3):529-48.

3. Nishikiori S, Yasuda J, Murata K, Takegaki J, Harada Y, Shirai Y, Fujita S. Resistance training rejuvenates aging skin by reducing circulating inflammatory factors and enhancing dermal extracellular matrices. *Sci Rep*. 2023;13(1):10214.

4. Hashimoto T, Tsukamoto H, Ando S, Ogoh S. Effect of exercise on brain health: the potential role of lactate as a myokine. *Metabolites*. 2021;11(12):813.

5. Maclnnis MJ, Gibala MJ. Physiological adaptations to interval training and the role of exercise intensity. *J Physiol*. 2017;595(9):2915-30.

6. Morland C, Andersson KA, Haugen ØP, Hadzic A, Kleppa L, Gille A, Rinholm JE, Palibrk V, Diget EH, Kennedy LH, Stølen T, Hennestad E, Moldestad O, Cai Y, Puchades M, Offermanns S, Vervaeke K, Bjørås M, Wisløff U, Storm-Mathisen J, Bergersen LH. Exercise induces cerebral VEGF and angiogenesis via the lactate receptor HCAR1. *Nat Commun*. 2017;8:15557.

7. Bonen A, Heynen M, Hatta H. Distribution of monocarboxylate transporters MCT1-MCT8 in rat tissues and human skeletal muscle. *Appl Physiol Nutr Metab*. 2006;31(1): 31-9.

8. Qian J, Li X, Yin Z, Dai Y, Zhang H, Li H, Peng C, Chen W. Yogurt alleviates imiquimod-induced psoriasis by activating the lactate/GPR81 signaling axis in mice. *J Agric Food Chem*. 2024;72(2):1055-66.

9. Chirumbolo S, Bertossi D, Magistretti P. Insights on the role of L-lactate as a signaling molecule in skin aging. *Biogerontology*. 2023;24(5):709-26.

10. Friedel M, Thompson IAP, Kasting G, Polsky R, Cunningham D, Soh HT, Heikenfeld J. Opportunities and challenges in the diagnostic utility of dermal interstitial fluid. *Nat Biomed Eng*. 2023;7(12):1541-55.

11. Tsuruoka N, Ishii K, Matsunaga T, Nagatomi R, Haga Y. Lactate and glucose measurement in subepidermal tissue using minimally invasive microperfusion needle. *Biomed Microdevices*. 2016;18(1):19.

12. Poole D, Rossiter HB, Brooks GA, Gladden LB. The anaerobic threshold: 50+ years of controversy. *J Physiol*. 2021;599(3):737-67.

13. Kim J, Sempionatto JR, Imani S, Hartel MC, Barfdokht A, Tang G, Campbell AS,
Mercier PP, Wang J. Simultaneous monitoring of sweat and interstitial fluid using a
single wearable biosensor platform. *Adv Sci.* 2018;5(10):1800880.

14. Matsui T, Soya S, Okamoto M, Ichitani Y, Kawanaka K, Soya H. Brain glycogen
decreases during prolonged exercise. *J Physiol.* 2011; 589 (Pt13):3383-93.

15. Matsui T, Ishikawa T, Ito H, Okamoto M, Inoue K, Lee MC, Fujikawa T, Ichitani Y,
Kawanaka K, Soya H. Brain glycogen supercompensation following exhaustive
exercise. *J Physiol.* 2012;590(3):607-16.

16. Matsui T, Omuro H, Liu YF, Soya M, Shima T, McEwen BS, Soya H. Astrocytic
glycogen-derived lactate fuels the brain during exhaustive exercise to maintain
endurance capacity. *Proc Natl Acad Sci USA*. 2017;114(24):6358-63.

17. Soya H, Mukai A, Deocaris CC, Ohiwa N, Chang H, Nishijima T, Fujikawa T, Togashi
K, Saito T. Threshold-like pattern of neuronal activation in the hypothalamus during
treadmill running: establishment of a minimum running stress (MRS) rat model.
Neurosci Res.2007; 58(4):341-8.

18. Chien MN, Fan SH, Huang CH, Wu CC, Huang JT. Continuous lactate monitoring
system based on percutaneous microneedle array. *Sensors*. 2022;22(4):1468.

19. Beaver WL, Wasserman K, Whipp BJ. A new method for detecting anaerobic
threshold by gas exchange. *J Appl Physiol*. 1986;60(6):2020-7.

20. Dror N, Weidling J, White S, Ortenzio F, Shreim S, Keating MT, Pharm H, Random-
Aizik S, Botvinick E. Clinical evaluation of a novel subcutaneous lactate monitor. *J*
Clin Monit Comput. 2022;36(2):537-43.

21. Stapleton J, Gagnon D, Kenny GP. Short-term exercise training does not improve
whole-body heat loss when rate of metabolic heat production is considered. *Eur J*
Appl Physiol. 2010;109(3):437-46.

22. Tigchelaar F, Groen H, Westgren M, Huinink KD, Cremers T, vanden Berg PP. A new
microdialysis probe for continuous lactate measurement during fetal monitoring:
Proof of concept in an animal model. *Acta Obstet Gynecol Scand*. 2020;99(10):1411-
6.

23. Messonnier LA, Emhoff CA, Fattor JA, Horning MA, Carlson TJ, Brooks GA.
Lactate kinetics at the lactate threshold in trained and untrained men. *J Appl Physiol*.
2013;114(11): 1593-602.

24. Garber GE, Blissmer B, Deschenes MR, Franklin BA, Lamonte MJ, Lee IM, Nieman
DC, Swain DP. American College of Sports Medicine. American College of Sports
Medicine position stand. Quantity and quality of exercise for developing and
maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently

healthy adults: guidance for prescribing exercise. *Med Sci Sports Exerc*.
2011;43(7):1334-59.
25. Juel C, Holten MK, Dela F. Effects of strength training on muscle lactate release and
MCT1 and MCT4 content in healthy and type 2 diabetic humans. *J Physiol*.
2004;556(Pt 1):297-304.
26. Jaminick NA, Pettitt RW, Granata C, Pyne DB, Bishop DJ. An examination and
critique of current methods to determine exercise intensity. *Sports Med*.
2020;50(10):1729-56.

Figure legends

Figure 1 The overview of experimental procedures using a jugular catheter and an
ultra-minimally microneedle perfusion system.

(*A*) The experimental settings of rats with inserted catheter and microneedle perfusion
system during exercise and experimental protocol of acute incremental running to
determine lactate threshold for rats. (*B*) The specification of microneedle. (*C*) The part of
skin with the microneedle inserted and its fixation. (*D*) The part of skin after removing
microneedle. (*C* and *D*) indicate no obvious inflammation and bleeding, suggesting that
the harmful effect of inserting the single microneedle perfusion system may be minimum.

Figure 2 Lactate concentration increases in the skin fluid and blood during incremental treadmill running.

Time course changes in blood (*A*) and skin fluid (*B*) lactate concentrations and skin fluid amount (*C*) pre, during, and post incremental running test. Each plot and shaded regions represent mean values and continuous standard error bands, respectively. *P < 0.05 vs. 0 min (immediately before the start of the exercise test).

Figure 3 Skin fluid lactate dynamics were significantly associated with blood lactate
dynamics during the exercise test but not afterward.
The correlation coefficient between the blood and skin fluid lactate concentrations in
typical case A and B, and pooled data of all cases during (*A-C*) and following (*D-F*) the
exercise. Shaded regions represent 95% simultaneous confidence bands. *P < 0.05, n.s.,
not significant.

Figure 4 Lactate threshold calculated from skin fluid lactate are closely associated with those calculated from blood.

(*A*, *B*) Representative LT profiles calculated by the dynamics from blood and skin fluid. The absolute running speed corresponding to the LT calculated from blood and skin fluid (*C*) and their relationship (*D*). The running speed (*E*) and its relationship (*F*) relative to peak speed during incremental exercise corresponding to LT calculated from blood and skin fluid, respectively. The values of bar graph (*C*, *E*) represent means and individual plots. Shaded regions represent 95% simultaneous confidence bands. n.s., not significant, *$P < 0.05$.