**Diurnal control of iron responsive element (IRE)-containing mRNAs through iron regular**

**proteins IRP1 and IRP2 is mediated by feeding rhythms**

Hima Priyanka Nadimpalli[1,#], Georgia Katsioudi[1,#], Enes S. Arpa[1], Lies Chikhaoui[1], A. Bulak

Arpat[1], Angelica Liechti[1], Gaël Palais[2], Claudia Tessmer[3], Ilse Hofmann[3], Bruno Galy[2], David

Gatfield[1]*

Affiliations:

[1] Center for Integrative Genomics, University of Lausanne, 1015 Lausanne, Switzerland.

[2] German Cancer Research Center (DKFZ), Division of Virus-associated Carcinogenesis, Im

Neuenheimer Feld 280, 69120, Heidelberg, Germany.

[3] German Cancer Research Center (DKFZ), Core Facility Antibodies, Im Neuenheimer Feld 280,

69120, Heidelberg, Germany.

# equal contribution

* Corresponding author

Abstract

A central mechanism for cellular iron homeostasis involves iron regulatory proteins, IRP1 and

IRP2, that interact with iron regulatory elements (IREs) on mRNAs. IRPs act as sensors of

metabolic cues (foremost of iron itself) and can regulate either translation or transcript stability,

depending on IRE location within the mRNA. In liver, IRP2 is widely considered the principle

player mediating IRE regulation. Here, we uncover high-amplitude diurnal rhythms in the

regulation of several prominent IRE-containing mRNAs in liver, compatible with maximal IRP

activity at the onset of the dark phase of the day (a timepoint likely underrepresented in previous

studies on IRP/IRE regulation). We find that IRP2 protein abundance itself oscillates over the day,

yet ribosome profiling in livers from IRP2-deficient mice uncovers that IRP2 is dispensable for

maximal repression of its rhythmically regulated target mRNAs at the time of its highest

abundance, i.e. the light-dark transition. These findings are compatible with temporally controlled

redundancy in IRP activities, with IRP2 exclusively mediating regulation of IRE-containing

transcripts in the light phase and redundancy, conceivably with IRP1, at dark onset. We further

unveil that the identified diurnal IRP/IRE regulation can ensue in the absence of a functional

circadian clock as long as feeding is rhythmic, and that it follows feeding rhythms also in the

presence of a clock, thus identifying feeding-associated signals as the dominant driver of IRP/IRE

rhythmicity. Altogether, our study identifies a novel level of regulatory control and complexity in a

metabolic pathway that had been considered well-understood since a long time.

## Introduction

Mammalian behaviour, physiology and metabolism undergo distinct changes over the course of the day (reviewed in [1]). These rhythmic patterns are orchestrated to large extent via the body's internal timekeeping system, the circadian clock. Molecular circadian oscillators can be found in virtually all cell types of the body and are hierarchically organised. The master clock in the brain's suprachiasmatic nuclei (SCN) is entrained to the environmental light-dark cycle through signals that it receives via projections from the retina. In turn, the SCN entrains (via e.g. hormones, body temperature cycles, feeding-fasting cycles) the peripheral oscillators that are active in the remainder of the organism. Across cell types, the rhythm-generating core clock has a similar molecular makeup that is based on negative transcriptional feedback loops in gene expression (reviewed in [2]). In the main loop, the negative feedback involves the transcriptional activators BMAL1:CLOCK (and/or BMAL1:NPAS2 in some neurons) that drive the rhythmic expression of the PERIOD and CRYPTOCHROME proteins which act as repressors of BMAL1:CLOCK-mediated transcription. Downstream of the core clock mechanism, BMAL1:CLOCK and other rhythmic transcriptional activities drive daily cycles in gene expression at hundreds to thousands of loci known as clock-controlled genes (CCGs). The resulting transcriptome-scale oscillations in mRNA abundance have been extensively studied across a variety of organs and cell types, genetic mouse models, light-dark regimens, metabolic conditions and dietary treatments. Notably, metabolism is a key rhythmically regulated process in organs such as the liver, with many links to human disease [3].

While rhythmic transcription is thought to be the principal driver of oscillating gene expression, various roles for post-transcriptional mechanisms have emerged as well (reviewed in [4, 5]). Conceptually, mRNA accumulation can be rendered rhythmic by time-of-day-dependent modulation of its stability, rather than of its transcription; and proteins can display daily changes in abundance in the absence of rhythmic mRNA, due to time-of-day-dependent control of translation and/or protein stability. To this end, previous studies using ribosome profiling (ribo-seq) in mouse liver around-the-clock have identified transcripts and pathways that are rhythmically regulated specifically at the level of translation, uncovering ribosomal protein and translation factor biosynthesis as the most prominent group of proteins regulated by this mechanism [6, 7].

An intriguing case of rhythmic translation that we noted in a previous study, are three transcripts involved in iron metabolism, *Ferritin light chain 1 (Ftl1), Ferritin heavy chain 1 (Fth1)* and *Aminolevulinic acid synthase 2 (Alas2)* [6]. The observation that Ferritin mRNAs (encoding the main iron storage proteins) are translationally repressed and activated, respectively, in response

to low and high iron concentrations, led to the discovery of the translational control element known
as the IRE (iron-responsive element) more than 40 years ago [8, 9]; *Alas2*, encoding the enzyme
that catalyses the first step in the heme biosynthetic pathway especially in erythrocytes, was
identified as IRE-containing a few years later [10] (please note that *Alas2* is distinct in its regulation
from *Alas1*, which specifies the housekeeping version of the enzyme and is well-known for its
very high transcriptional rhythmicity [11]). Since then, several other IRE-containing transcripts
have been identified and validated, and many more predicted ([12] and see below). IREs act
together with IRE-binding proteins (IRPs; iron-regulatory proteins) in response to physiological
iron availability and other cues ($O_2$, $H_2O_2$, NO) (reviewed in [13, 14]). The liver, which is the major
iron storage site in mammals, has been used as a model organ to study IRE/IRP regulation. From
the analysis of livers of mice deficient for the known IRPs, IRP1 (encoded by the *Aco1* gene) and
IRP2 (encoded by *Ireb2*), it has been concluded that IRP2 is the main actor in hepatic IRE
regulation, whereas IRP1 has a limited role, unable to fully replace IRP2 in the regulation of the
most prominent IRE-containing transcripts (e.g. [15-17]). Importantly, none of these studies
considered a possible time-of-day component in the regulatory mechanism.
In the present work, we first explore in a systematic fashion to what extent IRE-containing
transcripts beyond *Ftl1, Fth1* and *Alas2* are subject to rhythmic regulation. We then analyse
dependency of rhythmic regulation on the two IRPs using ribo-seq in livers collected at defined
timepoints in the light vs. dark phase from *Aco1/Irp1* and *Ireb2/Irp2* knockout mice. Finally, we
analyse previously published datasets to infer to what extent rhythmic IRE-regulation is a
circadian clock- vs. feeding-driven phenomenon. Our findings lead to a re-evaluation of current
models of IRP1 vs. IRP2-dependence of hepatic IRE-regulation, a key mechanism in organismal
iron homeostasis, and they expand our knowledge of the intricate connections between rhythmic
gene expression and metabolism.
95
**Results**
*A subset of iron response element-containing transcripts is rhythmically translated in mouse liver*
In our previous study [6] we used RNA-seq and ribo-seq on mouse liver samples collected around-
the-clock to identify and compare daily gene expression rhythms occurring at the level of mRNA
abundance and translation (Fig. 1A). These analyses had revealed rhythmicity across various
iron homeostatic genes, from cellular iron uptake to intracellular iron transport and storage,
hepatic heme and Fe-S cluster biosynthesis, cellular iron export, and systemic hormonal iron
homeostasis [6]. Of the almost 20 identified rhythmically regulated iron metabolic genes, the
majority showed similar oscillations in both RNA-seq and ribo-seq data, indicating that their

oscillations were driven at the level of transcription (and/or RNA stability). However, three
transcripts formed a notable exception – *Ftl1, Fth1* and *Alas2* were invariable in their mRNA
abundance, yet they showed robust rhythms in ribosome occupancy. All three transcripts contain
an IRE in their 5' untranslated region (5' UTR) [12]. Such 5' IREs act in concert with iron regulatory
proteins, IRP1 and IRP2, which bind to the IREs and inhibit translation under conditions of low
intracellular iron and as a function of other physiological cues (Fig. 1B-C) [12]. IRE sequences
fold into specific hairpin structures (Fig. 1D) that are crucial for IRP binding and have been used
for IRE predictions [18, 19]. In mammals, more than a dozen IRE-containing mRNAs have been
reported, mostly in 5' UTRs. In rare cases, IREs are in the 3' UTR. In *Transferrin receptor (Tfrc)*
mRNA, multiple 3' IREs regulate transcript stability in an IRP-dependent manner, involving
endonucleolytic cleavage (Fig. 1C) [20]. Single 3' IREs have also been identified in *Slc11a2* (also
known as *Dmt1,* for *Divalent metal transporter 1*) [21] and *Profilin 2 (Pfn2)* [22], but their
mechanisms of action appear to be distinct from regulation of *Tfrc*.
With *Ftl1, Fth1* and *Alas2* translation showing robust oscillations [6], we first wished to investigate
whether rhythmic translation was more commonly found across IRE-containing transcripts. To
this end, we used our liver [6] and kidney [23] RNA-seq and ribo-seq datasets to compare time-
resolved gene expression profiles of additional IRE-containing transcripts. As compared to the
robust rhythms in translation seen for *Fth1, Ftl1* and *Alas2* in liver (Fig. 1E, upper), other 5' IRE-
containing mRNAs lacked high-amplitude oscillations in ribosome occupancy. Thus, the
translation of *Ferroportin (Slc40a1)* mRNA showed, at best, low-amplitude oscillations, whereas
*Aconitase 2 (Aco2)* and *Hif2a/Endothelial PAS domain-containing protein 1 (Epas1)* were non-
rhythmic (Fig. 1F, upper), despite harbouring well-characterized 5' IREs [24-26]. We next
compared the time-resolved gene expression profiles from liver to those in kidney (Fig. 1E-F,
lower). *Ftl1* and *Fth1* showed blunted rhythmic profiles in kidney, resembling the low amplitudes
seen for *Slc40a1* (please note that low expression of *Alas2* in kidney precluded drawing strong
conclusions); *Aco2* and *Epas1* were clearly non-rhythmic in kidney at RNA abundance and
translational levels. In summary, we concluded that, first, rhythmic translational regulation in liver
only occurred on a subset of 5' IRE-containing transcripts. Second, rhythmic translational
regulation showed tissue-specificity, with higher amplitudes in liver as compared to kidney.
We considered it a plausible hypothesis that distinct features in IRE sequence or structure could
underlie the observed differences in regulation. We inspected whether the IRE sequences and
secondary structure predictions showed specific characteristics correlating with rhythmic vs. non-
rhythmic translation (Fig. 1G-H). Despite individual differences across IREs with regard to the
length of the paired stem, the positions of additional bulge nucleotides, or the location of the IRE

relative to the 5' end of the mRNA or the translation start codon, we did not identify a particular
feature that would distinguish the rhythmicity-associated IREs (Fig. 1G) from the other IREs (Fig.
1H).
142
*Rhythmic abundance of 3' IRE-containing* Transferrin receptor *mRNA is controlled post-*
*transcriptionally and altered in nonsense-mediated mRNA decay mutant livers*
Next, we analysed the daily expression profiles of *Tfrc* mRNA, which harbours multiple 3' IREs
(Fig. 2A) that regulate transcript stability (Fig. 1C) [20]. *Tfrc* mRNA abundance was robustly
rhythmic in both liver and kidney and ribosome footprints closely matched the mRNA profiles,
indicating that no additional rhythmicity was imparted at the level of translation (Fig. 2B). We then
confirmed that the observed *Tfrc* mRNA rhythms were of post-transcriptional origin, in line with
the 3' IRE mechanism and previous observations [6, 27]. To this end, we leveraged two around-
the-clock mouse liver datasets [28, 29] in which we had carried out RNA-seq on total RNA,
allowing the quantification of both unspliced pre-mRNA (intron-mapping reads) and of mature
mRNA (exon-mapping reads). With pre-mRNA levels serving as a proxy for nascent
RNA/transcription at the *Tfrc* locus, both datasets confirmed that *Tfrc* was transcribed in a non-
rhythmic fashion, yet its mRNA accumulated rhythmically around-the-clock (Fig. 2C-D, data from
control mice). We concluded that *Tfrc* rhythms were generated post-transcriptionally by
modulating transcript stability in a diurnal fashion. This finding is further supported by a recent
preprint [30].
One of the analysed wild-type liver datasets (Fig. 2C) was part of a study that contained a
matching cohort of animals that were deficient in miRNA biogenesis due to the hepatocyte-
specific deletion of *Dicer* [28]. It has previously been proposed that *Tfrc* mRNA stability is
regulated by the interplay between IRPs and miR-7/miR-141 binding at the 3' IRE stem-loops
[31]. However, because *Tfrc* mRNA profiles were unaltered in *Dicer* knockouts (Fig. 2C), we deem
it unlikely that miRNA-mediated mechanisms make strong contributions to hepatic 3' IRE
regulation. The other analysed dataset (Fig. 2D) contained a matching cohort of animals that
expressed a catalytically inactive, mutant version of the endonuclease SMG6; this enzyme
degrades transcripts targeted through the nonsense-mediated decay (NMD) pathway [29].
Interestingly, at all timepoints *Tfrc* mRNA abundances were post-transcriptionally increased in
*Smg6* mutant livers, leading to >2-fold higher mRNA levels; still, mutants showed similar
rhythmicity dynamics and peak-to-trough amplitudes in the range of ca. 3.2-fold (mutants) vs. 2.6-
fold (controls) (Fig. 2D). This finding could indicate that SMG6 is implicated, directly or indirectly,
in the regulation of *Tfrc* mRNA stability, potentially as a part or independently of the Roquin- and

Regnase-1-dependent pathways previously implicated in *Tfrc* mRNA decay [32, 33]. Another possibility is that the *Tfrc* locus produces high levels of unproductively spliced transcript isoforms that are subsequently cleared by NMD; indeed, several *Tfrc* mRNA variants with retained introns are annotated in Ensembl (www.ensembl.org/ - locus ENSMUSG00000022797). To evaluate the impact of NMD on mature TFRC levels, we analysed its protein abundance by Western blot in control and NMD-deficient livers around-the-clock (Fig. 2E, F). TFRC protein abundance in wild-type animals was minimal and maximal at *Zeitgeber* Time (ZT) 12 and 20, respectively, with a 2-fold peak-to-trough amplitude that is consistent with its mRNA rhythmicity (Fig. 2D). In *Smg6* mutant livers, the around-the-clock pattern of TFRC levels was irregular with a 1.8-fold peak-to-trough amplitude, and its average abundance over-the-day was increased 1.4-fold as compared to controls (Fig. 2F). Hence, the effect of NMD loss-of-function on TFRC protein was quantitatively considerably smaller than what could have been expected from the *Tfrc* mRNA increase (Fig. 2D). We thus propose that most likely the main role of NMD in *Tfrc* regulation consists in the clearance of unproductive mRNA isoforms, although some effect on regulating mature, translated *Tfrc* mRNA is plausible as well. Altogether, involvement of NMD factors in *Tfrc* mRNA regulation is not unprecedented [34] and remains intriguing, in particular because of recent work that identified TFRC and iron homeostasis as regulators that act *upstream* of NMD, through a mechanism that relies on the Fe-S cluster-containing ribosome recycling factor ABCE1 [35]. One may thus speculate about a possible feedback mechanism, with (i) TFRC protein controlling cellular iron uptake, (ii) intracellular iron availability controlling NMD activity, and (iii) NMD factors implicated in regulating *Tfrc* mRNA stability.

*Rhythmic translation/mRNA abundance predicts phases of maximal and minimal IRP activity*

The above analyses established that high-amplitude rhythmic regulation was particularly prominent for four IRE-containing transcripts in mouse liver. Of these, the three translationally regulated transcripts *(Ftl1, Fth1, Alas2)* showed similar rhythmic phase, with a peak in translation occurring at approximately ZT4-6 and a trough at ~ZT12 (Fig. 1E). Given that the association with IRPs inhibits the translation of IRE-containing transcripts (Fig. 1B-C), we would expect IRP binding/activity on *Ftl1, Fth1* and *Alas2* mRNAs to be minimal at ZT4-6 and maximal at ZT12. For *Tfrc*, the binding of IRPs to the IREs would increase the stability of the mRNA and (assuming constant rates in *Tfrc* mRNA transcription) maximal IRP activity would be predicted during the rising phase of *Tfrc* mRNA abundance, i.e. around ZT12 as well. Therefore, all four rhythmic gene expression events would be consistent with a mechanism involving rhythmic IRP activity on IREs that is minimal and maximal at ZT4-6 and ZT12, respectively (Fig. 2G).

The observed diurnal regulation could be mediated simply by daily changes in IRP protein
abundance; we had dismissed this mechanism, however, in our previous study in which we had
observed no rhythmicity (or, at most, very modest daily changes) for liver IRP1/ACO1 or
IRP2/IREB2 at the level of mRNA, ribosome footprints or protein [6]. Since publication of our
study, however, it has been reported that the antibody we had used to quantify IRP2/IREB2
(abcam ab1811539) not only detects IRP2/IREB2 (as marketed by the vendor at the time), but
also IRP1/ACO1 [36]. Therefore, our original analysis of around-the-clock IRP2 protein levels was
undoubtedly flawed, representing an overlay of both IRPs rather than IRP2 specifically. Moreover,
it has been reported that in colon cancer cells, the *Ireb2* gene is circadianly transcribed through
BMAL1:CLOCK, engendering *Ireb2* mRNA and IRP2/IREB2 protein oscillations that subsequently
regulate *Tfrc* stability in a rhythmic fashion [37]. The ensemble of these findings prompted us to
revisit the temporal dynamics of IRP expression across our datasets. Thus, our RNA-seq and
ribo-seq data showed modest rhythmicity (<1.5-fold peak-to-trough amplitudes) for *Aco1* and
*Ireb2* in liver but not in kidney (Fig. 2H). For *Ireb2*, the mRNA phase – maximal abundance around
ZT10 – was indeed compatible with the BMAL1:CLOCK regulation that has been reported from
colon cancer cells [37]. However, the shallow peak-trough amplitudes that *Ireb2* displays for both
RNA and pre-mRNA (Fig. 2I, lower) indicated that rhythmic transcriptional activation through
BMAL1:CLOCK would not appear to be an important driver of *Ireb2* mRNA synthesis in liver. For
comparison, in the same dataset *Arntl/Bmal1* itself, and well-established BMAL1:CLOCK targets
such as *Dbp* [38], present very high amplitude mRNA and pre-mRNA oscillations (Fig. 2J).
The main regulation of IRP1 activity and of IRP2 abundance occurs through post-translational
mechanisms (Fig. 1B) [14]. Given the issues with antibody specificity described above, we re-
evaluated IRP1 and IRP2 protein levels around-the-clock using highly specific monoclonal
antibodies that are free of cross-reactivity (see later, Fig. 4G, J). Using the same time series of
pooled liver protein extracts as for TFRC quantification (Fig. 2E), no high-amplitude rhythmicity
was apparent for ACO1/IRP1 (Fig. 2K, L). By contrast, the Western blot for IREB2/IRP2 revealed
high-amplitude rhythmic protein accumulation in the control animals (*Smg6*$^{wt/wt}$), with minimal and
maximal protein abundances around ZT4 and ZT12, respectively (Fig. 2N, O). In the matching
time series from *Smg6*$^{mut/mut}$ animals, the IRP2 rhythm was blunted (Fig. 2N, O). The difference
in IRP2 protein levels quantified from single liver samples (ZT4 and ZT12; control animals)
showed a ca. 10-fold amplitude with some inter-individual variability (Fig. 2M, P). Next, we
analysed the levels of FBXL5, which is the F-box protein that targets IREB2/IRP2 for proteasomal
degradation and is key for the regulation of IRP2 stability and abundance; according to the
established mechanism, FBXL5 acts as an iron sensor that is stabilized by Fe binding under high

iron conditions, yet unfolded and degraded during iron scarcity [39, 40]. In addition, an FBXL5-

independent mechanism of IREB2/IRP2-dependent target regulation has recently been identified

as well [41]. However, our western blot analysis in the control liver time series and in single liver

samples indicated that FBXL5 protein levels did not show the same marked rhythmicity as

displayed by IREB2/IRP2 (Fig. 2Q-S). Thus, we deem it unlikely that the observed changes in

IREB2/IRP2 (Fig. 2N, O) are a direct outcome of FBXL5 availability. Interestingly, $Smg6^{mut/mut}$

animals displayed blunted IRP2 rhythmicity compared to wild-type (Fig. 2N, O). This suggests

that the NMD pathway may somehow contribute to daily changes in IRP2 abundance in liver,

which together with recent report of FBXL5-independent mechanism of IREB2/IRP2-dependent

target regulation [41], highlights the complexity of IRP2 regulation.

*In vivo recording of diurnal gene expression oscillations shows unperturbed circadian*

*mPER2::LUC reporter rhythms in* Aco1 *and* Ireb2 *knockout mice*

We concluded from the above analyses that rhythmicity of the IRE-containing transcripts *Ftl1,*

*Fth1, Alas2* and *Tfrc* was compatible with diurnal regulation through IRPs, with a trough and peak

of IRP activity in the early light phase and at the light-dark transition, respectively. Conceivably,

the mechanism underlying these daily changes could be related to oscillations in IREB2/IRP2

levels that followed a similar rhythm (at least in the control animal time series) with lower IRP2 in

the light phase and higher IRP2 in the dark phase (Fig. 2N, O). In support of this hypothesis,

impact on IRE regulation has been reported mainly upon IRP2 deficiency and not so upon IRP1

loss [15, 42]. Yet, IRP1 has in the past been suggested to be particularly abundant in liver, and

our ribosome footprint data confirmed that ACO1/IRP1 is more highly expressed than

IREB2/IRP2, with a translational output for ACO1/IRP1 >5-fold higher than that of IREB2/IRP2

(Fig. 2H; compare maximal liver footprint Reads Per Kilobase per Million mapped reads of ca.

150 for *Aco1* and of ca. 25 for *Ireb2*). Importantly, previous studies have not addressed the

possibility of temporal control of IRP1-mediated IRE regulation in liver. Hence, depending on the

affinities that IRP1 and IRP2 have for individual IREs, and on additional regulatory mechanisms

that may be in place, either of the IRPs, or both, may contribute to rhythmic regulation, even if

only IREB2/IRP2 appears to undergo daily changes in protein abundance.

We set out to determine experimentally how rhythmic target regulation would be changed in the

absence of IRP1 or IRP2. To this end, we made use of constitutive *Aco1* and *Ireb2* knockout

mouse strains [43]. These knockout mice present with systemic alterations to iron homeostasis

and phenotypic abnormalities, including in liver (reviewed in [44]); moreover, there are

documented links between iron and the circadian timekeeping mechanism itself. For example,

several core clock proteins contain heme [11, 45, 46] and the transcription of *Per1* and *Per2* has
been reported as sensitive to iron accumulation through a mechanism that involves histone
methylation state [47]. Therefore, we first explored whether whole-body *Aco1* or *Ireb2* knockout
engendered phenotypes in entrained liver clock rhythms that could potentially confound the
specific analysis of rhythmic IRP/IRE regulation. We used a method for the real-time recording of
liver gene expression in freely moving mice [48, 49] that relies on luciferase reporters, luciferin
delivery via an osmotic minipump, and highly sensitive bioluminescence photon counting (Fig.
3A). Into the *Aco1* and *Ireb2* knockout lines, we crossed the *mPer2::luc* knock-in allele [50] that
expresses a protein fusion between core clock component PER2 and firefly luciferase, serving as
a precise readout of diurnal oscillations. We carried out real-time recordings of liver rhythms under
conditions that ensured light-entrainment of the master (SCN) clock to an external 24-hour light-
dark cycle by means of a skeleton photoperiod, i.e. two 30 min light pulses applied at times
corresponding to the beginning and to the end of the light phase in a 12h-light-12h-dark (LD12:12)
cycle (Fig. 3B). Our experiments revealed rhythmic bioluminescence rhythms in wild-type
animals, as expected, and also in littermate animals that were knockouts for either of the IRPs
(Fig. 3C-F). For $Aco1^{-/-}$ mice, the phase of entrained mPER2::LUC reporter oscillations was
virtually indistinguishable from controls (Fig. 3D). In the $Ireb2^{-/-}$ animals, the entrained
mPER2::LUC phase was minimally delayed (~0.5 h) as compared to controls (Fig. 3F), yet no
other overt differences were apparent. We nevertheless verified that the liver clock in *Ireb2*
knockouts was fully functional by recording free-running rhythms from liver organ explants (Fig.
3G). Free-running bioluminescence rhythms from *Ireb2* knockouts and controls (Fig. 3H) and their
quantification (Fig. 3I) indicated overall comparable oscillations *ex vivo*, yet with higher variability
in period length between individual explants in the knockouts. Altogether, our data suggested that
diurnal oscillations in liver were comparable between IRP knockouts and their respective controls,
and in particular that entrained clock phases between genotypes showed good correspondence.
300
*Timepoint-specific gene expression datasets from IRP-deficient livers allow analysis of diurnal*
*regulation*
For our further analyses we selected the two timepoints when rhythmic IRP/IRE activity would be
low (ZT5) and high (ZT12), respectively, leading to high and low translation efficiency *(Ftl1, Fth1,*
*Alas2)* and mRNA turnover *(Tfrc)* (Fig. 4A). We reasoned that in the simplest scenario, one of the
IRPs (e.g. IRP2, the most potent IRE-regulator in liver according to the literature) was singly
responsible for rhythmic target regulation. This rhythm would then be detectable in the control
livers when comparing the two timepoints, yet absent in the livers from animals deficient for the

rhythm-relevant IRP. More complex regulatory scenarios are conceivable, too, such as redundant

regulation by the two IRPs, which would lead to no, or partial deregulation of rhythmicity in the

knockouts. We collected 3 livers per timepoint and genotype ($Aco1^{-/-}$ and $Ireb2^{-/-}$; plus their

respective, matched $Aco1^{+/+}$ and $Ireb2^{+/+}$ controls carrying the same genetic background) and

prepared RNA-seq and ribo-seq libraries (Fig. 4B). After sequencing and analysis, quantification

of $Aco1$-mapping ribosome protected fragments (RPFs) in $Aco1^{-/-}$ livers confirmed the knockout

at the level of ACO1/IRP1 translation/protein biosynthesis (Fig. 4C, top panel). Because in the

knockout allele the open reading frame is disrupted yet still produces an mRNA from the locus

[43], the reduction in $Aco1$ in the RNA-seq data was less pronounced (Fig. 4C, middle panel).

$Aco1$ translation efficiency (TE; defined as RPF/RNA ratio) was decreased in the knockouts as

expected (Fig. 4C, lower panel). In livers from $Ireb2$-deficient animals, $Aco1$ RNA and RPFs were

decreased by 10-20% as compared to their littermate controls (Fig. 4D), indicating that

IREB2/IRP2 loss did not lead to compensatory higher IRP1 biosynthesis. Overall analogous

observations were made for $Ireb2$ expression, which showed the expected decrease in the $Ireb2^{-}$

$^{/-}$ animals at RPF and TE level (Fig. 4F, upper and lower panels), but not for the RNA (Fig. 4F,

middle panel). In $Aco1$ knockouts, $Ireb2$ expression was largely unchanged at the RPF level,

indicating that there was no substantial compensatory upregulation (Fig. 4E). We next validated

knockout and possible compensatory effects by Western blot. These analyses confirmed the

expected loss of ACO1/IRP1 in $Aco1^{-/-}$ mice (Fig. 4G, H). In these animals, IREB2/IRP2

abundance showed a tendency towards increased levels that did not reach statistical significance

(Fig. 4G, I). Higher abundance of IREB2/IRP2 at ZT12 vs. ZT5 was evident as well, although

quantitively less pronounced than in the around-the-clock time series analysed before (compare

Fig. 4G, I with Fig. 2M-P). In the $Ireb2$ cohort, knockouts were devoid of IREB2/IRP2 protein as

expected (Fig. 4J, L). In the control animals from this cohort, IREB2/IRP2 increased in abundance

from ZT5 to ZT12 in a quantitatively minor (yet statistically significant) fashion (Fig. 4L). In

summary, we concluded that the datasets obtained from the two timepoints and genotypes would

allow the analysis of interactions between IRP function and time-of-day and, more specifically, to

investigate whether the identified rhythmic regulation could be attributed specifically to one of the

IRPs. Moreover, our experiments validated the specificity of the IRP1 and IRP2 antibodies.

Finally, although we observed a significant difference in IRP2 abundance between light-phase

and dark-phase timepoints across different independent datasets, the fold-changes seen in the

three cohorts (Fig. 2O, P; Fig. 4I, L) were highly variable, pointing towards limited robustness of

IRP2 rhythmicity.

*Previous observations on IRE-IRP regulation were likely biased towards mechanisms occurring*

*during the light phase of the day*

A number of previous studies have quantified how the expression of IRE-containing transcripts is

altered in IRP knockout livers (e.g. [15-17, 43]). These studies focused in particular on Ferritin

protein levels and on *Tfrc* mRNA abundance as readouts in IRP1- or IRP2-deficient mice. A main

outcome from these studies was that the targeted deletion of *Ireb2/Irp2*, but not of *Aco1/Irp1*, is

associated with increased hepatic Ferritin protein and reduced *Tfrc* mRNA levels. These findings

led to the model that IREB2/IRP2 is the main actor in hepatic IRE regulation and that ACO1/IRP1

cannot compensate for its loss. None of these studies, however, used ribo-seq to specifically

quantify translation rates, nor did they consider a possible temporal component in regulation. As

the timing of animal sacrifice is not reported in previous publications on hepatic IRE regulation,

we estimated a likely time window during which liver sampling could have been carried out.

Standard animal housing usually follows an LD12:12 schedule that is approximately in sync with

geophysical time (e.g., the light phase is set from 7 am to 7 pm). We can assume that animals

were sacrificed during "core working hours", i.e. between early morning and late afternoon. This

would correspond to an approximate time window from ZT2 to ZT10 (which translates, as an

example, to 9 am – 5 pm in the authors' current animal facility at the University of Lausanne). In

particular the timepoint for which we predict the strongest IRP activity on its rhythmic targets

(ZT12; see Fig. 4A) is unlikely to have been included in previously published work, given that it

lies at the beginning of the dark phase (7 pm in our facility). Despite decades of research on the

IRP/IRE system, specific regulatory effects and mechanisms occurring at the onset of, or during

the dark phase may thus have gone unnoticed. Dark phase timepoints are, however, particularly

relevant when working with nocturnal species, as this is when mice are most active and ingest

most of their food. Also in the knockout and control animals used in our experiments, most activity

is seen between ZT12-24 as judged by the *in vivo* recordings of locomotor activity (Fig. 3D, F;

lower panels). That particular experiment was conducted using skeleton photoperiod illumination

(i.e., for most of the "subjective day", the animal was nevertheless in the dark). Due to the strong

acute effects of light on mouse behaviour – suppressing activity and inducing sleep (e.g. [51]) –

the partitioning between inactive light and active dark phase would be at least as pronounced in

an animal facility under LD12:12.

*IRP loss affects the regulation of 5' and 3' IRE-containing transcripts in a strongly timepoint-*

*specific fashion*

Next, we assessed across genotypes and timepoints how 5' IRE-containing transcripts (*Ftl1*, *Fth1*,
*Alas2*) were affected in their translation efficiencies (Fig. 5A-C) and 3' IRE-containing *Tfrc* in its
RNA abundance (Fig. 5D). In the TE analyses, we observed changes that were not only IRP-
specific, but also strongly dependent on timepoint. The most striking observation was the
timepoint-specificity of translational derepression in *Ireb2$^{-/-}$* mice (Fig. 5A-C, right panels). Thus,
*Ftl1* and *Fth1* (and to a lower extent *Alas2*) showed strongly increased translation efficiencies
(i.e., derepression) in the absence of IREB2/IRP2 during the light phase of the day (ZT5), in line
with the model that IREB2/IRP2 is the main IRE-regulating IRP in liver [15-17]. By contrast, at the
beginning of the dark phase (ZT12), we observed that TEs were low and near-identical between
*Ireb2$^{-/-}$* and *Ireb2$^{+/+}$* animals (Fig. 5A-C, right panels). These findings indicated that a repressive
mechanism distinct from IREB2/IRP2 was acting on the translation of IRE-containing transcripts
specifically at ZT12. This mechanism was unable to compensate for IREB2/IRP2 loss at the light
phase timepoint.
Timepoint-specific effects in *Ireb2$^{-/-}$* vs. *Ireb2$^{+/+}$* animals could also be observed for *Tfrc* (Fig. 5D,
right panels). On the one hand, the tendency to reduced *Tfrc* levels in *Ireb2$^{-/-}$* livers (in line with
the model that IRP2 protects *Tfrc* mRNA from degradation) was visible at ZT5. However, at ZT12
there was no decrease in *Tfrc* mRNA in the absence of IREB2/IRP2 and rather a trend towards
increased abundance. Therefore, similarly to the TE repressive effects seen for 5' IREs, also the
mRNA stabilising effect for 3' IRE-containing *Tfrc* did not require IRP2 at the dark-phase
timepoint.
We next analysed the effects associated with the absence of IRP1 (*Aco1$^{-/-}$* vs. *Aco1$^{+/+}$* animals).
Previous work has demonstrated that FTL1 protein levels are unchanged in livers of IRP1-
deficient mice, including under low- and high-iron diets [15]. Indeed, *Ftl1* TE was overall similar
between *Aco1$^{-/-}$* and *Aco1$^{+/+}$* animals (Fig. 5A, left panel). For *Fth1* and *Alas2,* some translational
derepression was visible at ZT5 and, in particular for *Fth1*, also at ZT12 (Fig. 5B, C, left panels;
effects not statistically significant). As a result, the differences between TEs at peak (ZT5) and
trough (ZT12) were reduced in *Aco1$^{-/-}$* vs. *Aco1$^{+/+}$* animals, indicating slightly blunted amplitudes
of rhythmic *Fth1* and *Alas2* translation when ACO1/IRP1 was absent. Finally, changes in *Tfrc*
mRNA abundance were overall consistent with the effects seen for *Fth1* and *Alas2* TEs. Thus,
*Tfrc* mRNA levels were lower in the absence of IRP1 at both timepoints (Fig. 5D, left panels;
statistically significant at ZT5 but not at ZT12 due to the spread seen across biological replicates).
We thus concluded that despite the generally held model that IRP1 has little effect on IRE
regulation in liver, specific differences between knockouts and controls were apparent. However,

these did not reach statistical significance due to the relatively low sample size (3 animals per
group) and considerable variability across mice.

In summary, the above analyses uncovered a complex relationship between timepoint (ZT5,
ZT12) and genotype ($Aco1^{-/-}$ vs. $Aco1^{+/+}$; $Ireb2^{-/-}$ vs. $Ireb2^{+/+}$) in the regulation of different IRE-
containing mRNAs (*Ftl1, Fth1*, *Alas2, Tfrc*). Our data indicate that IRP2 is not solely responsible
for IRE regulation in liver. Especially at ZT12, IRP2 actually appears to be dispensable for
translational repression. They also reveal that absence of IRP1 affects the regulation of IRE-
containing mRNAs and, in several cases, the consequences of IRP1-loss appeared pronounced
at ZT12. Taken together, it is conceivable that for these rhythmically regulated transcripts, IRP
activities are partitioned according to time-of-day, with IRP2 dominating regulation at ZT5 (light
phase) and IRP1 able to act at ZT12 (dark phase).

*Distinct timepoint-dependent regulation and basal repressed states correlate for core iron*
*metabolic vs. other 5' IRE-containing transcripts*

Beyond the four transcripts analysed in the previous section, several other mRNAs contain
validated IREs (e.g. Fig. 1F, H), and many more have been predicted [52]. A recent study has
proposed that across different IRE-containing transcripts in rat liver, translational control through
IRPs follows a hierarchy based on different iron concentration set-points at which translational
repression occurs [53]. Thus, IRE-containing mRNAs encoding core iron storage factors (e.g.
Ferritins) are translationally repressed under normal physiological conditions and chow diet (i.e.,
"regular" cellular iron levels; our animals were fed normal chow diet throughout), whereas those
mRNAs that encode proteins with broader physiological roles are actively translated under these
conditions (e.g. *Aco2*, encoding the TCA cycle enzyme, mitochondrial aconitase). Under iron
overload, the former transcripts would undergo derepression, yet the latter only experience very
little additional stimulation of translation. In iron deficiency, by contrast, *Aco2* is highly responsive
and becomes strongly repressed. Interestingly, it was proposed that IRP1 contributes to this
setpoint mechanism through the kinetics of IRP1:IRE dissociation [53].

Conceivably, similar mechanisms may also underlie the differences that we observe in rhythmic
regulation. Indeed, in the control mice, *Ftl1* and *Fth1* (encoding core iron management proteins)
were generally more strongly repressed than *Alas2* (encoding a broader function protein involved
in heme biosynthesis) – see TE values in the wild-type animals in Fig. 5A-C that were in the range
of -4 to -1 for *Ftl1* and *Fth1*, but around -1.5 to -0.5 for *Alas2*. Please note that TE values are
derived from RPF/RNA ratios and can therefore be compared between different transcripts.

In order to investigate this phenomenon more globally, we set out to analyse the TEs of other IRE-containing mRNAs in our ribosome profiling datasets. First, we explored the group of transcripts with documented 5' IREs that only presented weak, or no, rhythmicity in our original dataset [6] and that are shown in Fig. 1F. Consistent with the validity of the "setpoint model", these transcripts all showed high basal translation efficiencies (-1 to +0.5) and IRP deficiency led, at most, to modest stimulation of translation (Fig. 5E-G). Thus, in *Ireb2*[-/-] animals, *Slc40a1/Ferroportin*, *Epas2/Hif2a* and *Aco2* were only weakly altered in their TE at either timepoint (Fig. 5E-G, right panels). In *Aco1/Irp1* knockouts, significant derepression was observable at both ZT5 and ZT12 for *Slc40a1/Ferroportin* (encoding a protein in the core iron metabolic machinery that functions as an iron exporter [53]; Fig. 5E, left panel). For *Epas1/Hif2a* a small effect was apparent at ZT12 (Fig. 5F, left panel; statistically not significant; please note that a previous report had suggested specific *Epas1/Hif2a* regulation through IRP1 in liver [17]). The effect size of de-repression for *Aco2* was quantitatively minimal as well (Fig. 5G, left panel). Finally, we also analysed possible gene expression changes for *Slc11a2/Dmt1*, which bears an IRE in its 3' UTR [21]. *Slc11a2* RNA levels changed in a similar fashion to what we had observed for *Tfrc* upon loss of IRP1 or IRP2 (compare Fig. 5H to Fig. 5D).

Next, we set out to classify transcripts in a comprehensive fashion, according to how their translation rates were affected across conditions in our ribo-seq datasets. To this end, we used k-means clustering to partition the transcriptome into overall 15 clusters according to the similarity of their z-score-transformed TE profiles (Fig. 5I and Table S1). Validated IRE-containing transcripts and predicted IRE-like mRNAs (list from [52]) were found in several clusters. The two most rhythmically translated IRE-containing mRNAs, *Ftl1* and *Fth1* – but no other (validated or predicted) IRE transcripts – were found together in a common cluster (cluster 8), where they were grouped in immediate neighbourhood (Fig. 5J, K), consistent with their distinct strongly rhythmic regulation. Another cluster – cluster 13 – contained several 5' IRE transcripts (*Aco2, Alas2, Slc40a1*, and predicted IRE-like candidates; see Fig. 5I, L), in line with common regulatory mechanisms and possible similar "set-points". Within cluster 13, the close vicinity of *Alas2* and *Slc40a1*, as compared to the more distant *Aco2* (Fig. 5L), recapitulated the differences in magnitude of derepression seen in the individual transcript analyses shown in Fig. 5C, E, G.

In summary, the above analyses support the idea that the observability of rhythmicity on IRE-containing transcripts depends on whether in the basal state, these mRNAs are strongly repressed, which allows for a timepoint-dependent derepression as a function of reduced IRP activity.

*Feeding-fasting cycles constitute the physiological basis of IRP/IRE rhythmicity*

Finally, we wished to understand whether rhythmic IRP/IRE regulation was a *bona fide* circadian clock-driven phenomenon that depended on functional clocks locally in hepatocytes and/or systemically in the whole organism, or if other rhythmicity cues were involved. Briefly, it is well established that various direct and indirect mechanisms, alone or in combination, are involved in driving gene expression oscillations in liver (reviewed in [54]). A considerable proportion of rhythmic mRNA expression is thus driven transcriptionally and directly through the cellular core clock. The transcription of these clock-controlled genes (CCGs) is dependent on the transcription factors that also generate the oscillations within the timekeeping mechanism. However, numerous cases of hepatic gene expression oscillations still persist in the absence of local liver clocks, as they are driven by systemic cues. The underlying mechanisms include, for example, circadian oscillations in body temperature that are generated by the master clock in the SCN and lead to changes in the activity of the transcription factor *Hsf1*, which in turn drives transcriptional oscillations without the obligatory involvement of local hepatocyte clocks [55-57]. Moreover, feeding/fasting cycles – that are a consequence of behavioural rhythmicity and hence an output from the master clock in the SCN – are a particularly important timing cue for liver and other metabolically active peripheral organs. Rhythmic feeding is a well-established *Zeitgeber* for the entrainment of hepatic clocks and, in addition, feeding-dependent signals also regulate gene expression directly, without the need of local hepatocyte clocks. For example, the nutrient-responsive mTOR pathway undergoes daily activation cycles in response to feeding, which regulates the translation of ribosomal protein mRNAs containing regulatory 5' TOP motifs [6, 58, 59].

In order to explore at which level in the organism a circadian clock was required for IRP/IRE rhythms, we used the oscillations of *Tfrc* mRNA as a proxy for rhythmic IRP/IRE regulation. This allowed us to take advantage of the numerous published around-the-clock RNA-seq datasets from wild-type and clock-deficient mice under a variety of feeding regimes. First, we analysed the RNA-seq data from a recent study [60] that used three different genetic animal models: wild-type, *Bmal1* knockout (i.e., behaviourally arrhythmic with no functional clocks in the entire body) and liver-specific *Bmal1* re-expression (i.e. behaviourally arrhythmic, with functional clock only in hepatocytes), either fed *ad libitum* or night-restricted. Night-restricted feeding exacerbated *Tfrc* rhythmicity in wild-type mice (Fig. 6A, upper). In *Bmal1* KO mice, *Tfrc* was arrhythmic under *ad libitum* feeding, yet rhythmicity was re-installed under night-feeding (Fig. 6A, middle) and in *Bmal1* re-expression mice, *Tfrc* regained rhythmicity under night-feeding (Fig. 6A, lower). This data was fully compatible with a model according to which local clocks (liver) or a circadian master clock

(SCN) were *per se* not essential for *Tfrc* rhythmicity, as long as feeding-fasting cycles persisted.

We confirmed this conclusion in two other datasets. Atger et al. [7] collected livers from night-fed

full-body *Bmal1* KO vs. wild-type mice and applied an RNA-seq methodology to quantify both

mRNA and pre-mRNA levels. In this experiment, *Tfrc* mRNA rhythms are indistinguishable

between the circadian clock-deficient *Bmal1* KO and the control animals under imposed night-

feeding (Fig. 6B, upper panel), and pre-mRNA (transcription) is identical and arrhythmic in both

genotypes (Fig. 6B, lower panel). We concluded that rhythmic feeding was sufficient to induce

post-transcriptionally generated *Tfrc* mRNA oscillations in the absence of an endogenous

circadian system. Next, we used the study by Greenwell et al. [61] to investigate the

complementary scenario, i.e. how *Tfrc* rhythmicity in liver fares in the presence of a wild-type

circadian system, but in the absence of feeding/fasting cycles (Fig. 6C). The analysis of four

different feeding regimens from night-restricted (100% food intake during the dark phase) to *ad*

*libitum* (~76% during dark), "damped"/almost entirely arrhythmic feeding (~56% during dark) and

arrhythmic feeding (~51% of food intake during dark vs. ~49% during light phase), revealed a

gradual disappearance of *Tfrc* rhythmicity that scaled with the loss of feeding/fasting cycles (Fig.

6C). Finally, we analysed data comprised in the study by Manella et al. [62] in which wild-type

mice were subjected to *ad libitum* or to day-restricted feeding under conditions that had little

influence on the phase of the liver circadian clock, which revealed that the peak in *Tfrc* mRNA

accumulation was inverted and followed feeding rhythms rather than the light-dark cycle (Fig. 6D).

From the above analyses that probe the relative importance of the circadian system vs.

feeding/fasting cycles for the rhythmic accumulation of *Tfrc*, we can conclude that it is via the

latter mechanism – feeding – that *Tfrc* mRNA oscillations are engendered. By extension, we

propose that feeding-related timing cues constitute a physiological basis for rhythmicity of the

IRP/IRE system in liver tissue.

**Discussion**

Several decades of research have led to a remarkable level of understanding of the IRP/IRE

system, which has become a paradigm, and literally a "textbook example" [63], of post-

transcriptional gene regulatory control, and which has caught particular attention certainly also

due to its relevance for human diseases such as iron storage disorders [44]. Surprisingly, the

regulatory feature that we have explored in detail in the current study – differences that are

associated with the time-of-day – has nevertheless gone largely unnoticed so far with the

exception of a recent preprint that touches on this topic as well [30]. One of our key findings is

that high-amplitude rhythmic regulation in the liver is restricted precisely to the subset of

transcripts that are archetypes of IRE regulation, namely the Ferritins (*Fth1, Fth2*), *Alas2* and *Tfrc*,

which were also historically the first to be discovered [8-10, 20]. Over the years, IREs in many

other transcripts have been reported, validated, or predicted – yet for none of them have we

observed comparably high peak-to-trough amplitudes across the various datasets that we have

analysed. In line with the "set-point model" that has recently been put forward to explain how IRP

activity thresholds lead to a hierarchy of translational repression across 5' IRE-containing

transcripts [53], the most rhythmically regulated mRNAs are also the most sensitive to repression

and generally exhibit lower translation rates (Fig. 5A-C) compared with the non- (or less-)

rhythmically regulated transcripts, which are overall more highly translated (Fig. 5E-G). Therefore,

we deem it likely that the appearance of rhythmicity for some but not for other IRE-containing

transcripts is not necessarily the reflection of principally different molecular mechanism of

regulation across these mRNAs, but rather of whether the transcript/IRE-specific thresholds

match the physiological dynamic range of regulatory cues that modulate the IRPs. This model

would also explain why rhythmic regulation for the same mRNAs is less prominent in the kidney;

conceivably, the dynamic range of the cues could simply be different in this organ.

An open question that we have not addressed in the framework of this study and that will require

future investigations, concerns the identity of the cue that is responsible for imparting rhythmicity.

Certainly, a prime candidate would be iron itself, more particularly iron in the so-called "labile iron

pool" or LIP, which consists of redox-active iron ions that are weakly liganded and therefore

engage in regulatory and metabolic processes [64]. Quantifying the relevant iron pool in tissues

is technically not a facile enterprise, given the large amounts of iron that is stored in ferritin or

otherwise bound in various Fe-S cluster- or cytochrome-containing proteins, plus the large

quantities of iron in the mitochondria and lysosomes. Moreover, in tissue analyses from liver, even

minute iron contaminations that can be released from lysed red blood cells (hemoglobin), would

likely by far exceed possible rhythmic differences in the relevant intracellular iron pool, making *in*

*vivo* measurements a particularly daunting task. Finally, there is ample evidence that additional

cues act to regulate IRPs as well, in particular oxygen, reactive oxygen species as well as nitric

oxide [42, 65-67]. Interestingly, daily rhythms in tissue $O_2$ [68] and $H_2O_2$ [69] levels have been

described. It will thus be important, yet not trivial, to disentangle to what extent these various cues

are relevant for rhythmic regulation of IRE-containing transcripts.

Another important aspect of our study is that it allows us to refine our understanding of the role of

IRP2 in hepatic IRE regulation. While there is indeed a large body of literature supporting the idea

that IRP2 is the principle actor in the regulation of Ferritin protein and *Tfrc* mRNA accumulation

in liver (e.g. [15-17]), we deem it likely that the vast majority of former studies analysed tissues

collected during the light phase of the day. Our examination of time-resolved ribo-seq data indeed
confirms that at the light phase timepoint ZT5, knockouts of IRP2 (but not of IRP1) present strong
translational depression of Ferritin mRNAs. However, at the dark phase timepoint, ZT12, low
translation efficiency is fully maintained even in the absence of IRP2 (Fig. 5A-B), indicating that a
different repressor is capable of acting on these IREs at this time-of-day. An obvious candidate is
IRP1, whose knockout indeed showed a tendency to derepression at ZT12, at least for *Fth1* (Fig.
5B). Since the double deficiency for both IRPs is not viable – even as a conditional, liver-specific
knockout [70] – we cannot test the hypothesis that at ZT12, it is the combined activity of IRP1 and
IRP2 that acts in repression. Still, based on our data we would propose a model according to
which IRP1 is inactive at ZT5, and thus all repression occurs through IRP2 for the light phase
timepoint, yet IRP1 and IRP2 are redundantly active at ZT12. In terms of possible molecular
mechanisms that could engender higher IRP1 activity at this time point, lower iron levels or
changes in one of the other relevant cues could play a role. For example, it is interesting to note
that liver $H_2O_2$ levels (a reflection of overall mitochondrial activity) are highest precisely at the
light-dark transition [69]. One may thus hypothesize that high $H_2O_2$ could engender the oxidation
and destabilisation of IRP1's Fe-S cluster, leading to apo-IRP1 that is capable of binding to IREs.
Experimental evidence will be needed to validate such speculations.
By leveraging publicly available datasets and using *Tfrc* oscillations as a readout for diurnal
IRP/IRE regulation, we also uncovered that rhythmic control does not require a local circadian
clock in hepatocytes, nor a central pacemaker in the SCN, as long as rhythmic feeding is
maintained (Fig. 6). In animals with a fully intact circadian system, night-restricted feeding readily
exacerbated the peak-to-trough amplitude of *Tfrc* abundance oscillations, and it led to some
phase delay as compared to *ad libitum* feeding. These observations are in line with the idea of
limited direct influence and interference from the circadian clock onto the observed rhythm. The
fact that daytime-fed mice have inverted hepatic *Tfrc* rhythms further underscores the dominant
role of feeding (Fig. 6D). In this aspect, IRP/IRE rhythmicity distinguishes itself from the rhythmic
mechanism acting on ribosomal protein (RP) mRNAs whose translational upregulation occurs
also around the light-dark transition in mice fed *ad libitum* or at night [6, 7, 58]. However, in day-
fed animals, the rhythm in newly synthesised RPs does not invert but becomes flat [59]. The RP
mRNAs contain 5' TOP motifs that are regulated via the mTOR pathway, which shows multiple
interactions with clock regulation [71]. A plausible model has emerged according to which the
translational upregulation of RP mRNAs by feeding is gated in a circadian fashion, i.e. the feeding
signal can exert its stimulatory effect only at a specific time of day. No such circadian gating
appears to be in place for the IRP/IRE system, as judged by *Tfrc* mRNA profiles.

Conceptually, it is interesting to note that our findings reinforce the idea that rhythmic gene

expression is governed by at least two distinct mechanisms. On the one hand, these are truly

circadian clock-driven oscillations that are engendered at the transcriptional level and can thus

be considered "hard-wired" within the DNA sequence. On the other hand, most post-

transcriptional/translational oscillations reported so far appear to reflect the activity of other, acute

cues and signals – in particular from feeding. Indeed, translation can be considered an ideal step

where such information can be integrated to directly modulate protein output in a rapid and

efficient manner. The reactivity of the IRP/IRE system to rhythmic feeding would be a further

example for this mechanism.

**Materials and methods**

**Animals**

All animal experiments were performed according to the cantonal guidelines of the Canton of

Vaud, Switzerland, license VD3402. Healthy adult male mice of age 12-24 weeks were used. All

mouse lines were maintained on a C57BL/6J background. The knockout mice for IRP1 (allele

*Aco1*$^{tm1.1Mwh}$) and IRP2 (allele *Ireb2*$^{tm1.2Mwh}$) have been described in [72]. Strains kindly provided

by Matthias W. Hentze, EMBL Heidelberg.

**Protein extraction and Western Blots**

Total liver protein lysates (*Aco1* and *Ireb2* strains) were prepared as follows: Livers were

homogenized in 3 volumes (w/v) of RIPA buffer (150 mM NaCl, 50 mM Tris pH 8.0, 1% NP40,

0.5% Na-deoxycholate, 0.1% SDS and Roche complete protease inhibitors) using a Teflon

homogenizer, incubated for 30 min on ice and centrifuged for 10 min at 16000 x g and 4°C.

Supernatants were stored at -80°C. Protein concentrations were quantified using a BCA protein

assay kit (Pierce). For the around-the-clock wild-type and *Smg6* mutant livers, total lysates from

our previous study were used that had been prepared by the NUN (NaCl, Urea, Nonidet P-40)

procedure, see [29].

Immunoblotting to PVDF membrane was carried out using iBlot 2 dry blotting system (Invitrogen).

For the blots in Fig 4G and J, bi-directional diffusion transfer that results in two mirror image

membranes per gel was carried out (protocol Ulrich Laemmli, personal communication). Briefly,

PVDF membranes were placed on both sides of the gel and further sandwiched by 2 pre-wet

Whatman filter papers and fibre pads. The stack was placed inside a transfer stack holder and

submerged in a tank filled with transfer buffer (Tris-HCl 10 mM pH7.5, EDTA 2 mM, NaCl 50 mM

and DTT 0.1 mM) and tightly sealed. The container was then placed in an incubator at 50°C for

overnight diffusion transfer. Following transfer, membranes were blocked and incubated with

primary and secondary antibodies in 5% milk/TBS-T. For IRP1 and IRP2 Western blots, undiluted

crude hybridoma supernatants were used in primary antibody incubation. The dilutions for the

remaining antibodies are as follows: anti-Transferrin Receptor (ThermoFisher 13-6800, 1:1000),

anti-FBXL5 (ThermoFisher PA5-113529, 1:1000), anti-VINCULIN (Abcam 129002, 1:10000), rat

monoclonal anti-ACO1 and anti-IREB2 (undiluted hybridoma supernatant from own preparation);

secondary antibodies: anti-rabbit HRP or anti-mouse HRP (Promega, 1:10000).

**RNA extraction and library preparation**

*Library preparation from footprints and total RNA*. The libraries for ribosome profiling were

generated as [6] with few modifications. Briefly, frozen liver tissues from individual mice were

homogenized with 5-6 strokes using a Teflon homogenizer in 3 volumes of polysome buffer (150

mM NaCl, 20 mM Tris-HCl pH7.4, 5 mM $MgCl_2$, 5 mM DTT, complete EDTA-free protease

inhibitors (Roche) and 40 U/ml RNasin plus (Promega)) supplemented with 1% Triton X-100 and

0.5% Na deoxycholate. Liver lysates were incubated on ice for 10 min and cleared by

centrifugation at 10000 g, 4°C for 10 min. Supernatants were flash-frozen and stored in liquid

nitrogen. For each sample, 15 $OD_{260}$ of liver lysate was digested with RNase I (Ambion) for 45

min at RT under gentle agitation. Digested samples were passed through pre-washed (4x with

700 µl polysome buffer for 1 min at 450 x g) Sephacryl S-400 HR spin columns (GE healthcare,

currently Cytiva), and centrifuged for 2 min at 450 x g at 4°C. The resulting flow-through was

mixed with Qiazol and ribosome-protected fragments (RPFs) were purified using miRNeasy RNA

extraction kit (Qiagen) according to the manufacturer's instructions. Total RNA was extracted from

undigested liver lysates using Qiazol and miRNeasy kit in parallel to generate total RNA libraries.

1 µg of RPF and total RNA samples were depleted of ribosomal RNA using RiboCop rRNA

depletion kit V1.3 (Lexogen), followed by purification with RNA Clean & Concentrator-5 (Zymo

research). All the subsequent steps were performed according to Illumina's TruSeq Ribo-Profile

protocol with minor modifications as described in [73]. Size-selection of fragments at different

steps was as follows: ribosome-protected fragments (RPFs) were size-separated on 15% urea-

polyacrylamide gels and excised at a size range of 24-36 nt (monosome footprint). The cDNA

fragments generated from RPFs and total RNA were separated on 10% urea-polyacrylamide gels

and excised at a size range of 70-80 nt and 70-100 nt, respectively. The PCR-amplified libraries

were size-selected on 8% native polyacrylamide gels. Monosome libraries were at ~150 bp and

total RNA libraries were at ~150 to 170 nt. All libraries were sequenced on Illumina HiSeq 2500.

*Ribosome profiling data analysis.* Sequencing reads were mapped following overall the
strategy described in [6]. Adapter sequences were removed from the sequenced reads using
cutadapt (v3.5) with the following options: --match-read-wildcards --overlap 8 --discard-untrimmed
--minimum-length 30 and reads were quality filtered using fastx_toolkit (-Q33 -q 30 -p 90).
Trimmed read sequences were filtered by their size using an in-house Python script with following
inclusive ranges: [26,35] for footprints, [21,60] for total RNA. Reads were then mapped to rRNA
and tRNA databases to remove contaminants using bowtie2 (-p 2 -L 15 -k 20 --no-unal -q). Only
reads that did not map to the above databases were mapped to mouse genome v38.100 using
STAR (version 2.70f). Read counts of total RNA and RPF were normalized with upper quantile
method of R package edgeR v3.4.2 [74]. Translational efficiencies (TE) were calculated on
normalised reads as the ratio of RPF / mRNA for each gene per sample. TEs were $\log_2$-
transformed and plotted for comparing means of replicates per time point for each genotype. K-
Means cluster analysis was performed on the different conditions (averaged by replicates). Based
on the Within-Cluster-Sum of Squared Errors, the number of clusters was fixed to 15 sets.
Hierarchical clustering was then performed within individual clusters to assess proximity between
genes of a given cluster.
696
**Liver explant preparation and bioluminescence recording**
For explant preparation, male IRP1 or IRP2 KO and matched control animals from the same
breedings (i.e., same genetic background) were euthanized following deep anaesthesia by
isoflurane inhalation. Liver tissue was excised and placed immediately in ice-cold Hank's buffer
(Thermo Fisher Scientific). The outermost edges of the tissue were carefully excised in a sterile
cabinet, and immediately placed on a 0.4 micron Millicell cell culture inserts (PICMORG50) in a
35 mm dish with phenol-free DMEM (Thermo Fisher Scientific, 11880028) containing 5% FBS, 2
mM glutamine, 100 U/ml penicillin, 100 µg/ml streptomycin and 0.1 mM luciferin. The parafilm-
sealed plates were placed for recording in the LumiCycler (Actimetrics) at 37°C and 5% CO2.
706
*In vivo* **recording in the RT-Biolumicorder**
Adult male mice, 12-20 weeks of age, carrying the genetically encoded circadian reporter allele
*mPer2::Luc* [50] and deficient for *Aco1* or *Ireb2* were used for the RT-Biolumicorder experiments,
along with controls, i.e. wild-type male mice from the same breedings/with the same genetic
background and of the same age. The experimental procedure followed our previously published
protocols [49]. Briefly, ALZET mini-osmotic pumps were filled with D-luciferin sodium salt and
closed with blue-coloured flow moderators (ALZET) under sterile conditions. The pumps were

activated at 37°C according to the manufacturer's instructions, followed by subcutaneous, dorsal implantation. Before implantation, the dorsal area of the mouse at the site where the liver is positioned was shaved using an electric razor. The RT-Biolumicorder (Lesa-Technology) consists of a cylindrical cage equipped with a photomultiplier tube (PMT) allowing for recording of real-time longitudinal bioluminescence from single-housed mice. The data, which also contain locomotor activity traces, light and food access information, were saved as text files and analysed using the MATLAB-based "Osiris" software according to [49].

**Acknowledgements**

We thank Matthias W. Hentze for knockout mouse strains; the Lausanne Genomic Technologies Facility for library preparation and sequencing; and Paul Franken and members of the Gatfield lab for comments on the manuscript.

**Funding**

Swiss National Science Foundation (SNSF), NCCR RNA & Disease grant 141735 (DG)

Swiss National Science Foundation (SNSF) individual grant 179190 (DG)

**Competing interests:** Authors declare that they have no competing interests.

**Data and Materials Availability:** All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary Materials. Sequencing data have been deposited at GEO (GSE243134) with reviewer token (suppressed).

**References**

1.  Cox, K.H. and J.S. Takahashi, *Circadian clock genes and the transcriptional architecture of the clock mechanism.* J Mol Endocrinol, 2019. **63**(4): p. R93-R102.

2.  Partch, C.L., C.B. Green, and J.S. Takahashi, *Molecular architecture of the mammalian circadian clock.* Trends Cell Biol, 2014. **24**(2): p. 90-9.

3.  Bolshette, N., et al., *Circadian regulation of liver function: from molecular mechanisms to disease pathophysiology.* Nat Rev Gastroenterol Hepatol, 2023.

4.  Crosby, P. and C.L. Partch, *New insights into non-transcriptional regulation of mammalian core clock proteins.* J Cell Sci, 2020. **133**(18).

5.  Castelo-Szekely, V. and D. Gatfield, *Emerging Roles of Translational Control in Circadian Timekeeping.* J Mol Biol, 2020. **432**(12): p. 3483-3497.

6.  Janich, P., et al., *Ribosome profiling reveals the rhythmic liver translatome and circadian*
*clock regulation by upstream open reading frames.* Genome Res, 2015. **25**(12): p. 1848-
59.
7.  Atger, F., et al., *Circadian and feeding rhythms differentially affect rhythmic mRNA*
*transcription and translation in mouse liver.* Proc Natl Acad Sci U S A, 2015. **112**(47): p.
E6579-88.
8.  Hentze, M.W., et al., *Identification of the iron-responsive element for the translational*
*regulation of human ferritin mRNA.* Science, 1987. **238**(4833): p. 1570-3.
9.  Aziz, N. and H.N. Munro, *Both subunits of rat liver ferritin are regulated at a translational*
*level by iron induction.* Nucleic Acids Res, 1986. **14**(2): p. 915-27.
10. Dandekar, T., et al., *Identification of a novel iron-responsive element in murine and human*
*erythroid delta-aminolevulinic acid synthase mRNA.* EMBO J, 1991. **10**(7): p. 1903-9.
11. Kaasik, K. and C.C. Lee, *Reciprocal regulation of haem biosynthesis and the circadian*
*clock in mammals.* Nature, 2004. **430**(6998): p. 467-71.
12. Anderson, C.P., et al., *Mammalian iron metabolism and its control by iron regulatory*
*proteins.* Biochim Biophys Acta, 2012. **1823**(9): p. 1468-83.
13. Muckenthaler, M.U., et al., *A Red Carpet for Iron Metabolism.* Cell, 2017. **168**(3): p. 344-
361.
14. Galy, B., M. Conrad, and M. Muckenthaler, *Mechanisms controlling cellular and systemic*
*iron homeostasis.* Nat Rev Mol Cell Biol, 2023.
15. Meyron-Holtz, E.G., et al., *Genetic ablations of iron regulatory proteins 1 and 2 reveal why*
*iron regulatory protein 2 dominates iron homeostasis.* EMBO J, 2004. **23**(2): p. 386-95.
16. Cooperman, S.S., et al., *Microcytic anemia, erythropoietic protoporphyria, and*
*neurodegeneration in mice with targeted deletion of iron-regulatory protein 2.* Blood, 2005.
**106**(3): p. 1084-91.
17. Anderson, S.A., et al., *The IRP1-HIF-2alpha axis coordinates iron and oxygen sensing*
*with erythropoiesis and iron absorption.* Cell Metab, 2013. **17**(2): p. 282-90.
18. Campillos, M., et al., *SIREs: searching for iron-responsive elements.* Nucleic Acids Res,
2010. **38**(Web Server issue): p. W360-7.
19. Sanchez, M., et al., *Iron regulatory protein-1 and -2: transcriptome-wide definition of*
*binding mRNAs and shaping of the cellular proteome by iron regulatory proteins.* Blood,
2011. **118**(22): p. e168-79.

20.    Casey, J.L., et al., *Iron regulation of transferrin receptor mRNA levels requires iron-*
*responsive elements and a rapid turnover determinant in the 3' untranslated region of the*
*mRNA.* EMBO J, 1989. **8**(12): p. 3693-9.

21.    Gunshin, H., et al., *Iron-dependent regulation of the divalent metal ion transporter.* FEBS
Lett, 2001. **509**(2): p. 309-16.

22.    Luscieti, S., et al., *The actin-binding protein profilin 2 is a novel regulator of iron*
*homeostasis.* Blood, 2017. **130**(17): p. 1934-1945.

23.    Castelo-Szekely, V., et al., *Translational contributions to tissue specificity in rhythmic and*
*constitutive gene expression.* Genome Biol, 2017. **18**(1): p. 116.

24.    Lymboussaki, A., et al., *The role of the iron responsive element in the control of*
*ferroportin1/IREG1/MTP1 gene expression.* J Hepatol, 2003. **39**(5): p. 710-5.

25.    Gray, N.K., et al., *Translational regulation of mammalian and Drosophila citric acid cycle*
*enzymes via iron-responsive elements.* Proc Natl Acad Sci U S A, 1996. **93**(10): p. 4925-
30.

26.    Sanchez, M., et al., *Iron-regulatory proteins limit hypoxia-inducible factor-2alpha*
*expression in iron deficiency.* Nat Struct Mol Biol, 2007. **14**(5): p. 420-6.

27.    Le Martelot, G., et al., *Genome-wide RNA polymerase II profiles and RNA accumulation*
*reveal kinetics of transcription and associated epigenetic changes during diurnal cycles.*
PLoS biology, 2012. **10**(11): p. e1001442.

28.    Du, N.H., et al., *MicroRNAs shape circadian hepatic gene expression on a transcriptome-*
*wide scale.* Elife, 2014. **3**: p. e02510.

29.    Katsioudi, G., et al., *A conditional Smg6 mutant mouse model reveals circadian clock*
*regulation through the nonsense-mediated mRNA decay pathway.* Sci Adv, 2023. **9**(2): p.
eade2828.

30.    Bennett, C., et al., *Serum iron and hepatic transferrin receptor expression are circadian-*
*regulated.* bioRxiv, 2023: p. 2023.05.07.539729.

31.    Miyazawa, M., et al., *Regulation of transferrin receptor-1 mRNA by the interplay between*
*IRE-binding proteins and miR-7/miR-141 in the 3'-IRE stem-loops.* RNA, 2018. **24**(4): p.
468-479.

32.    Corral, V.M., et al., *Roquin is a major mediator of iron-regulated changes to transferrin*
*receptor-1 mRNA stability.* iScience, 2021. **24**(4): p. 102360.

33.    Yoshinaga, M., et al., *Regnase-1 Maintains Iron Homeostasis via the Degradation of*
*Transferrin Receptor 1 and Prolyl-Hydroxylase-Domain-Containing Protein 3 mRNAs.* Cell
Rep, 2017. **19**(8): p. 1614-1630.

34.    Mino, T., et al., *Translation-dependent unwinding of stem-loops by UPF1 licenses*
815           *Regnase-1 to degrade inflammatory mRNAs.* Nucleic Acids Res, 2019. **47**(16): p. 8838-
816           8859.

817    35.    Zhu, X., H. Zhang, and J.T. Mendell, *Ribosome Recycling by ABCE1 Links Lysosomal*
818           *Function and Iron Homeostasis to 3' UTR-Directed Regulation and Nonsense-Mediated*
819           *Decay.* Cell Rep, 2020. **32**(2): p. 107895.

820    36.    Fader, K.A., et al., *Corrigendum to "Convergence of hepcidin deficiency, systemic iron*
821           *overloading, heme accumulation, and REV-ERBalpha/beta activation in aryl hydrocarbon*
822           *receptor-elicited hepatotoxicity" [Toxicol. Appl. Pharmacol. 321 (2017) 1-17].* Toxicol Appl
823           Pharmacol, 2018. **344**: p. 74.

824    37.    Okazaki, F., et al., *Circadian Clock in a Mouse Colon Tumor Regulates Intracellular Iron*
825           *Levels to Promote Tumor Progression.* J Biol Chem, 2016. **291**(13): p. 7017-28.

826    38.    Ripperger, J.A. and U. Schibler, *Rhythmic CLOCK-BMAL1 binding to multiple E-box motifs*
827           *drives circadian Dbp transcription and chromatin transitions.* Nat Genet, 2006. **38**(3): p.
828           369-74.

829    39.    Vashisht, A.A., et al., *Control of iron homeostasis by an iron-regulated ubiquitin ligase.*
830           Science, 2009. **326**(5953): p. 718-21.

831    40.    Salahudeen, A.A., et al., *An E3 ligase possessing an iron-responsive hemerythrin domain*
832           *is a regulator of iron homeostasis.* Science, 2009. **326**(5953): p. 722-6.

833    41.    Terzi, E.M., et al., *Iron-sulfur cluster deficiency can be sensed by IRP2 and regulates iron*
834           *homeostasis and sensitivity to ferroptosis independent of IRP1 and FBXL5.* Sci Adv, 2021.
835           **7**(22).

836    42.    Meyron-Holtz, E.G., M.C. Ghosh, and T.A. Rouault, *Mammalian tissue oxygen levels*
837           *modulate iron-regulatory protein activities in vivo.* Science, 2004. **306**(5704): p. 2087-90.

838    43.    Galy, B., D. Ferring, and M.W. Hentze, *Generation of conditional alleles of the murine Iron*
839           *Regulatory Protein (IRP)-1 and -2 genes.* Genesis, 2005. **43**(4): p. 181-8.

840    44.    Wilkinson, N. and K. Pantopoulos, *The IRP/IRE system in vivo: insights from mouse*
841           *models.* Front Pharmacol, 2014. **5**: p. 176.

842    45.    Yin, L., et al., *Rev-erbalpha, a heme sensor that coordinates metabolic and circadian*
843           *pathways.* Science, 2007. **318**(5857): p. 1786-9.

844    46.    Andreini, C., et al., *The human iron-proteome.* Metallomics, 2018. **10**(9): p. 1223-1231.

845    47.    Liu, J., et al., *Iron accumulation with age alters metabolic pattern and circadian clock gene*
846           *expression through the reduction of AMP-modulated histone methylation.* J Biol Chem,
847           2022. **298**(6): p. 101968.

48.    Saini, C., et al., *Real-time recording of circadian liver gene expression in freely moving mice reveals the phase-setting behavior of hepatocyte clocks.* Genes Dev, 2013. **27**(13): p. 1526-36.

49.    Katsioudi, G., et al., *Recording of Diurnal Gene Expression in Peripheral Organs of Mice Using the RT-Biolumicorder.* Methods Mol Biol, 2022. **2482**: p. 217-242.

50.    Yoo, S.H., et al., *PERIOD2::LUCIFERASE real-time reporting of circadian dynamics reveals persistent circadian oscillations in mouse peripheral tissues.* Proc Natl Acad Sci U S A, 2004. **101**(15): p. 5339-46.

51.    Tsai, J.W., et al., *Melanopsin as a sleep modulator: circadian gating of the direct effects of light on sleep and altered sleep homeostasis in Opn4(-/-) mice.* PLoS Biol, 2009. **7**(6): p. e1000125.

52.    Volz, K., *Conservation in the Iron Responsive Element Family.* Genes (Basel), 2021. **12**(9).

53.    Garza, K.R., et al., *Differential translational control of 5' IRE-containing mRNA in response to dietary iron deficiency and acute iron overload.* Metallomics, 2020. **12**(12): p. 2186-2198.

54.    Dibner, C., U. Schibler, and U. Albrecht, *The mammalian circadian timing system: organization and coordination of central and peripheral clocks.* Annu Rev Physiol, 2010. **72**: p. 517-49.

55.    Reinke, H., et al., *Differential display of DNA-binding proteins reveals heat-shock factor 1 as a circadian transcription factor.* Genes Dev, 2008. **22**(3): p. 331-45.

56.    Buhr, E.D., S.H. Yoo, and J.S. Takahashi, *Temperature as a universal resetting cue for mammalian circadian oscillators.* Science, 2010. **330**(6002): p. 379-85.

57.    Saini, C., et al., *Simulated body temperature rhythms reveal the phase-shifting behavior and plasticity of mammalian circadian oscillators.* Genes Dev, 2012. **26**(6): p. 567-80.

58.    Jouffe, C., et al., *The circadian clock coordinates ribosome biogenesis.* PLoS Biol, 2013. **11**(1): p. e1001455.

59.    Sinturel, F., et al., *Diurnal Oscillations in Liver Mass and Cell Size Accompany Ribosome Assembly Cycles.* Cell, 2017. **169**(4): p. 651-663 e14.

60.    Greco, C.M., et al., *Integration of feeding behavior by the liver circadian clock reveals network dependency of metabolic rhythms.* Sci Adv, 2021. **7**(39): p. eabi7828.

61.    Greenwell, B.J., et al., *Rhythmic Food Intake Drives Rhythmic Gene Expression More Potently than the Hepatic Circadian Clock in Mice.* Cell Rep, 2019. **27**(3): p. 649-657 e5.

62.    Manella, G., et al., *The liver-clock coordinates rhythmicity of peripheral tissues in response*
*to feeding.* Nat Metab, 2021. **3**(6): p. 829-842.

63.    Lodish, H.F., *Molecular Cell Biology*. 2000: W.H. Freeman.

64.    Kakhlon, O. and Z.I. Cabantchik, *The labile iron pool: characterization, measurement, and*
*participation in cellular processes(1).* Free Radic Biol Med, 2002. **33**(8): p. 1037-46.

65.    Weiss, G., et al., *Translational regulation via iron-responsive elements by the nitric*
*oxide/NO-synthase pathway.* EMBO J, 1993. **12**(9): p. 3651-7.

66.    Pantopoulos, K., G. Weiss, and M.W. Hentze, *Nitric oxide and oxidative stress (H2O2)*
*control mammalian iron metabolism by different pathways.* Mol Cell Biol, 1996. **16**(7): p.
3781-8.

67.    Pantopoulos, K. and M.W. Hentze, *Rapid responses to oxidative stress mediated by iron*
*regulatory protein.* EMBO J, 1995. **14**(12): p. 2917-24.

68.    Adamovich, Y., et al., *Rhythmic Oxygen Levels Reset Circadian Clocks through*
*HIF1alpha.* Cell Metab, 2017. **25**(1): p. 93-101.

69.    Pei, J.F., et al., *Diurnal oscillations of endogenous H(2)O(2) sustained by p66(Shc)*
*regulate circadian clocks.* Nat Cell Biol, 2019. **21**(12): p. 1553-1564.

70.    Galy, B., et al., *Iron regulatory proteins secure mitochondrial iron sufficiency and function.*
Cell metabolism, 2010. **12**(2): p. 194-201.

71.    Cao, R., *mTOR Signaling, Translational Control, and the Circadian Clock.* Front Genet,
2018. **9**: p. 367.

72.    Galy, B., et al., *Targeted mutagenesis of the murine IRP1 and IRP2 genes reveals context-*
*dependent RNA processing differences in vivo.* RNA, 2004. **10**(7): p. 1019-25.

73.    Arpat, A.B., et al., *Transcriptome-wide sites of collided ribosomes reveal principles of*
*translational pausing.* Genome Res, 2020. **30**(7): p. 985-999.

74.    Robinson, M.D. and A. Oshlack, *A scaling normalization method for differential expression*
*analysis of RNA-seq data.* Genome Biol, 2010. **11**(3): p. R25.

 **Figure Legends**

**Figure 1. Translational rhythmicity is detected on IRE-containing mRNAs in a transcript-**

**and tissue-specific fashion.**

(**A**) Schematic of the dataset collection carried out for the studies [6, 23] for the liver and kidney

gene expression graphs shown in this Figure.

(**B**) Schematic model of IRP activity on IREs. Thus, IRP1 can bind to IRE hairpins under conditions

of low iron availability. When iron is abundant, it assembles an Fe-S cluster and the protein

becomes cytoplasmic Aconitase (Aco1). In this state, it does not bind IREs anymore. For IRP2,

there is an iron-dependent degradation pathway that relies on an iron-sensor F-box protein,

FBXL5.

(**C**) IREs in 5' UTRs and 3' UTRs act by different mechanisms (schematic): When in the 5' UTR,

IRP-bound IREs inhibit scanning ribosomes from accessing the initiation codon, thus leading to

translational repression. After removal of IRP binding, ribosomes can translate the coding

sequence (upper). For 3' UTR IREs, the binding of IRPs inhibits access to endonucleases

(depicted as red scissors) that would otherwise initiate transcript degradation. When IRPs

dissociate, the mRNA is thus destabilised and becomes less abundant (lower).

(**D**) Schematic of the IRE hairpin consensus sequence and structure. IREs are about 35 nt in

length land are characterized by an apical 6-nt loop motif 5'-CAGWGH-3' (W = A or T and H = A,

C or T), and two stem regions (N5 upper stem and variable lower stem) that are separated by a

unpaired C bulge. Colour code depicts sequence conservation across IREs from high (blue) to

medium (orange) and low (olive).

(**E**) RPKM values of CDS-mapping ribosome protected fragment (RPF) data (blue) and RNA-seq

data (orange) from the datasets of [6, 23] for the transcripts *Ftl1, Fth1* and *Alas2* around the 24-

h daily cycle in liver (upper) and kidney (lower). All three genes show high amplitude translation

in liver. Means per timepoint are plotted; "error bars"/vertical lines connect the two biological

replicates. Dashed lines represent rhythmic curve fittings to the data, as described in [6].

(**F**) As in panel (E), but for the non-/less-rhythmically translated IRE-containing transcripts

*Slc40a1/Ferroportin, Aco2, Epas1/Hif2-alpha*.

(**G**) Sequence and predicted structure, as well as relative position within 5' UTR, of the IREs of

the murine transcripts shown in (E). Colour code as in (D).

(**H**) As in (G), but for the transcripts that are shown in panel (F).

**Figure 2. Correspondence between *Tfrc* mRNA rhythms and translational oscillations are consistent with diurnal IRP activity.**

(**A**) Sequence and predicted hairpin structures of the five IREs annotated within the mouse *Tfrc* 3' UTR. Nucleotide distances from the TAA stop codon, between individual haipins, and to the annotated transcript end are noted in grey.

(**B**) RPKM values of CDS-mapping ribosome protected fragment (RPF) data (blue) and RNA-seq data (orange) from the datasets of [6, 23] for the *Tfrc* transcript around the 24-h daily cycle in liver (upper) and kidney (lower). Means per timepoint are plotted; "error bars"/vertical lines connect the two biological replicates. Dashed lines represent rhythmic curve fittings to the data, as described in [6].

(**C**) Expression plots (RPKM) showing around-the-clock data from [28] for *Tfrc* mRNA (exonic reads; upper panel) and pre-mRNA (intronic reads; lower panel) in control (blue) and miRNA-deficient (*Dicer* KO) livers (yellow). Means per timepoint are plotted; "error bars"/vertical lines connect the two biological replicates. The two time series are plotted together in the same graph (vertical lines connect the data points from the two series). Rhythms are seen for mRNA but not for pre-mRNA, indicating that oscillations are post-transcriptionally generated (RNA stability). Dashed lines represent rhythmic curve fittings to the data, as described in [28].

(**D**) Expression plots for around-the-clock RNA-seq data from [29] for *Tfrc* showing mRNA (top panel; exonic reads) and pre-mRNA (bottom panel; intronic reads) for *Smg6* mutants (yellow) and controls (blue). RPKM values (reads per kilobase of transcript per million mapped reads) of individual mouse livers are shown as dots with solid lines connecting the means across time points. Dashed lines represent rhythmic curve fittings to the data, as described in [29].

(**E**) Western blot analysis of total liver proteins (pool of three mice per sample), for TFRC and VINCULIN (loading control), from control mice (blue; on left side) and *Smg6* mutant mice (yellow; on right side). Same samples as in [29] were used.

(**F**) Quantification of Western blot from (E). TFRC itensities were normalised to control band intensities (VINCULIN). The 1.4x upregulation refers to the mean signal in *Smg6* mutants vs. controls.

(**G**) Schematic representation of the model that can be deduced from the gene expression data from Fig. 1E (rhythmic translation of *Ftl1, Fth1* and *Alas2*) and Fig. 2B-D (rhythmic mRNA abundance of *Tfrc*). The observed rhythms are thus consistent with IRP activity being minimal and maximal at the light-phase timepoint ZT5 and the light-dark transition timepoint ZT12, respectively.

(**H**) As in panel (B), RPKM values of CDS-mapping ribosome protected fragment (RPF) data (blue) and RNA-seq data (orange) from the datasets of [6, 23] for the transcripts *Aco1* (encoding IRP1) and *Ireb2* (encoding IRP2) around the 24-h daily cycle in liver (upper) and kidney (lower). Means per timepoint are plotted; "error bars"/vertical lines connect the two biological replicates. Dashed lines represent rhythmic curve fittings to the data, as described in [6].

(**I**) Expression plots (RPKM) showing around-the-clock data from [28] for the indicated transcripts, *Aco1* and *Ireb2,* for mRNA (exonic reads; blue) and pre-mRNA (intronic reads; magenta) in control livers. The data from the miRNA-deficient (*Dicer* KO) livers were ommitted from these analyses. Means per timepoint are plotted; "error bars"/vertical lines connect the two biological replicates. Dashed lines represent rhythmic curve fittings to the data, as described in [28].

(**J**) As in (I), but for transcripts *Arntl/Bmal1* and *Dbp.*

(**K**) Western blot analysis of total liver proteins as in (E), for ACO1/IRP1 and VINCULIN (loading control).

(**L**) Quantification of Western blot from (K). ACO1/IRP1 itensities were normalised to control band intensities (VINCULIN).

(**M**) Western blot analysis of total liver proteins from individual animals for IREB2/IRP2; the same extracts that constituted the pools ZT4 and ZT12 (control series) in panel N were used.

(**N**) Western blot analysis of total liver proteins as in (E), for IREB2/IRPs and VINCULIN (loading control).

(**O**) Quantification of Western blot from (N). IREB2/IRP2 itensities were normalised to control band intensities (VINCULIN).

(**P**) Quantification of Western blot from individual livers at ZT4 and ZT12 from (M). IREB2/IRP2 itensities were normalised to control band intensities (VINCULIN). Statistical significance evaluated via Student's t-test.

(**Q**) Western blot analysis for FBXL5 and VINCULIN (loading control) in total liver proteins from the control series samples from [29]. Left side of blot: around-the-clock, pool of three mice per sample; right side: individiual mice, as in panel (M).

(**R**) Quantification of around-the-clock data from Western blot from (Q). FBXL5 itensities were normalised to control band intensities (VINCULIN).

(**S**) Quantification of Western blot from individual livers at ZT4 and ZT12 from (Q), right part of blot. FBXL5 itensities were normalised to control band intensities (VINCULIN). Statistical significance evaluated via Student's t-test.

**Figure 3. *In vivo* bioluminescence recoding shows that entrained circadian rhythms in liver**
**are unperturbed in mice that are deficient for *Aco1* or *Ireb2*.**
(**A**) Cartoon depicting the *in vivo* recording setup (RT-Biolumicorder) used to measure diurnal
bioluminescence rhythms in freely moving mice. The mouse is implanted with a luciferin-charged
osmotic minipump and dorsally shaved to ensure detectability of photons emitted from the liver.
It is then placed in the arena of the RT-Biolumicorder that records photons via a photomultiplier
tube (PMT), activity via infrared detection, and also offers the possibility to program food
availability and illumination.
(**B**) Schematic representation of the applied recording protocol in the RT-Biolumicorder setup.
Mice expressing the *Per2::Luc* reporter and implanted with the minipump are placed in the RT-
Biolumicorder where they are first subject to LD12:12 schedule and *ad libitum* feeding for 1-2
days, allowing for acclimatisation to the setup. Bioluminescence recording only occurs during the
dark phase. Then, the illumination schedule is changed to skeleton photoperiod (i.e. two 30 min
light pulses applied at times corresponding to the beginning and to the end of the light phase in a
12h-light-12h-dark (LD12:12) cycle). Recording occurs throughout the day with the exception of
the duration of the 30 min light pulses. The skeleton photoperiod is important as it keeps the SCN
clock entrained to a defined 24-hour T-cycle, allowing to specifically query potential changes in
the entrained liver clock,
(**C**) Bioluminescence rhythms and activity traces recorded with the protocol shown in (B) from
mice deficient for *Aco1* (ochre, N=2) and controls (grey, N=2). Mean signal (solid trace) and SEM
(shaded) over the whole course of the experiment are shown.
(**D**) Compiled data of (C), averaging all cycles from day 3. No change in PER2::LUC phase was
detectable between genotypes.
(**E**) Bioluminescence rhythms and activity traces recorded with the protocol shown in (B) from
mice deficient for *Ireb2* (purple, N=4) and controls (grey, N=4). Mean signal (solid trace) and SEM
(shaded) over the whole course of the experiment are shown.
(**F**) Compiled data of (E), averaging all cycles from day 3, For *Ireb2*$^{-/-}$ mice, PER2::LUC peak and
trough phases were slightly delayed as compared to the control mice with the same genetic
background.
(**G**) Schematic of liver explant rhythm measurements from the two genotypes, *Ireb2*$^{-/-}$ and
controls, using male mice from the same breedings. Mice carried the *Per2::Luc* reporter allele.
(**H**) Liver explant bioluminescence recording from *Ireb2*$^{-/-}$ (purple) and control (grey) mice. Free-
running circadian traces show mean signal and SEM for the two genotype over a recording period
of 96 hours. N=10 per genotype.

(**I**) Period length quantification of experiments shown in (H). No significant change in period length

was detectable. *Ireb2* knockout period lengths appeared more variable than controls.

**Figure 4. Time-resolved datasets allow analysis of IRP1 vs. IRP2-dependence of rhythmic**

**regulation.**

(**A**) Schematic representation of the conclusions drawn from the rhythmic analyses of Fig. 1-2

regarding the timing of IRP activity and the selection of timepoints for the ribo-seq experiment.

Thus, high *Ftl1, Fth1* and *Alas2* translation efficiency and low *Tfrc* abundance, conceivably as a

result of mRNA instability, occur in the light phase of the day, allowing us to predict generally low

IRP activity around ZT5. By contrast, low *Ftl1, Fth1* and *Alas2* translation efficiency and the rise

in *Tfrc* abundance at the light-dark transition suggests high IRP activity at ZT12.

(**B**) Schematic of the experiment carried out to assess timepoint- and genotype-specific ribosome

profiling in the two IRP-deficient mouse strains and matched controls. Thus, 3 animals each were

sacrificed at ZT5 and ZT12, for genotypes *Aco1*$^{+/+}$ (dark grey), *Aco1*$^{-/-}$ (ochre), *Ireb2*$^{+/+}$ (light grey),

*Ireb2*$^{-/-}$ (purple). Livers were further processed for ribo-seq and RNA-seq.

(**C**) Quantification of ribosome protected fragments (RPF, top) and RNA abundance (middle) and

relative translation efficiency (log$_2$ ratio of RPF/RNA) for the *Aco1* transcript in *Aco1* knockout

mice and their matched controls, for the two timepoints ZT5 and ZT12. Read count is normalised

to library depth. Statistical significance evaluated by unpaired t-test and indicated with asterisks

(*** $P < 0.001$, ** $P < 0.01$, * $P < 0.05$).

(**D**) As in (C), for the *Aco1* transcript in *Ireb2* knockout mice and their matched controls.

(**E**) Quantification of ribosome protected fragments (RPF, top) and RNA abundance (middle) and

relative translation efficiency (log$_2$ ratio of RPF/RNA) for the *Ireb2* transcript in *Aco1* knockout

mice and their matched controls, for the two timepoints ZT5 and ZT12. Read count is normalised

to library depth. Statistical significance evaluated by unpaired t-test and indicated with asterisks

(** $P < 0.001$, ** $P < 0.01$, * $P < 0.05$).

(**F**) As in (E), for the *Ireb2* transcript in *Ireb2* knockout mice and their matched controls.

(**G**) Western blot analysis of total protein extracts from the livers used for the gene expression

profiling in panels (C) and (D), i.e. the *Aco1*$^{-/-}$ and matched *Aco1*$^{+/+}$ mice. VINCULIN was used as

loading control to normalise ACO1/IRP1 and IREB2/IRP2 signals.

(**H**) Quantification of Western blot signals for ACO1/IRP1 (normalised for loading via VINCULIN)

from panel (G). No ACO1 is detectable in the *Aco1*$^{-/-}$ animals, as expected. ACO1 abundance is

identical in the control animals at ZT5 and ZT12.

(**I**) Quantification of Western blot signals for IREB2/IRP2 (normalised for loading via VINCULIN)

from panel (G). IREB2 is slightly upregulated in the *Aco1*$^{-/-}$ vs. *Aco1*$^{+/+}$ animals (not significant).

IREB2 levels are increased at ZT12 vs. ZT5 (** $P < 0.01$, One-way ANOVA with Šídák's multiple

comparisons test).

(**J**) As in (G), but for the *Ireb2*$^{-/-}$ and matched *Ireb2*$^{+/+}$ animals whose livers were used for gene

expression analysis in panels (E) and (F).

(**K**) Quantification of Western blot signals for ACO1/IRP1 (normalised for loading via VINCULIN)

from panel (J). Differences beween timepoints or genotypes (*Ireb2*$^{-/-}$ vs. *Ireb2*$^{+/+}$) were statistically

not significant (One-way ANOVA with Šídák's multiple comparisons test).

(**L**) Quantification of Western blot signals for IREB2/IRP2 (normalised for loading via VINCULIN)

from panel (J). IREB2 is undetectable in *Ireb2*$^{-/-}$ mice as expected. The upregulation from ZT5 to

ZT12 is statistically significant (* $P < 0.05$, Student's t-test).

**Figure 5. Strong timepoint- and transcript-dependence of derepression upon loss of ACO1/IRP1 vs. IREB2/IRP2.**

(**A**) Translation efficiency analysis for *Ftl1* at ZT5 and ZT12 in the *Aco1* line (*Aco1$^{-/-}$* vs. *Aco1$^{+/+}$* in ochre and grey, respectively) in the left panel and for *Ireb2* line (*Ireb2$^{-/-}$* vs. *Ireb2$^{+/+}$* in purple and grey, respectively) in the right panel. Statistical significance evaluated by unpaired t-test and indicated with asterisks (*** $P < 0.001$, ** $P < 0.01$, * $P < 0.05$).

(**B**) As in (A), for *Fth1*.

(**C**) As in (A), for *Alas2*.

(**D**) RPF (upper) and RNA (lower) normalised reads for *Tfrc* at ZT5 and ZT12 in the *Aco1* line (*Aco1$^{-/-}$* vs. *Aco1$^{+/+}$* in ochre and grey, respectively) in the left panel and for *Ireb2* line (*Ireb2$^{-/-}$* vs. *Ireb2$^{+/+}$* in purple and grey, respectively) in the right panel. Statistical significance evaluated by unpaired t-test and indicated with asterisks (*** $P < 0.001$, ** $P < 0.01$, * $P < 0.05$).

(**E**) As in (A), for *Slc40a1*.

(**F**) As in (A), for *Epas1*.

(**G**) As in (A), for *Aco2*.

(**H**) As in (D) for *Slc11a2/Dmt1*.

(**I**) K-means clustering analysis on z-score-transformed TE profiles across all eight conditions (2 timepoints, 4 genotypes; replicate means were used). The numbering of the 15 clusters and the gene count are given below the heatmap. Below the clusters, the known/validated IRE-containing transcripts are highlighted in red, with (*Tfrc* and *Slc11a2* in parantheses, given that their IRE-regulation is not at the level of TE, but of RNA stability). Predicted IRE-like mRNAs from a list in [52] are highlighted in orange. Note that the two most rhythmically translated IRE-containing mRNAs, *Ftl1* and *Fth1* are found together in cluster 8. Several other IRE transcipts are enriched in cluster 13.

(**J**) Hierarchical clustering according to the similarity in TE profiles for transcripts in cluster 8 from panel (I). *Ftl1* and *Fth1* cluster together (marked in red).

(**K**) Zoom into the part of cluster 8 that contains *Ftl1* and *Fth1* in direct vicinity (marked in red). The unsupervised clustering thus confirms that these two transcript indeed have a distinct mode of (rhythmic) regulation that sets them apart from other transcripts, even IRE-containing ones.

(**L**) Hierarchical clustering according to the similarity in TE profiles for transcripts belonging to cluster 13 from panel (I), which contains several validated (red) and predicted (orange) IRE-containing transcripts, possibly indicating common regulatory mechanisms and, in particular, similar "set-points".

**Figure 6. Phase of *Tfrc* oscillations across feeding paradigms and clock models suggests that food intake is the major driver of rhythmicity in the IRP/IRE system.**

(**A**) Plotting of liver *Tfrc* expression data from the study by Greco et al. [60], which included two feeding paradigms (*ad libitum* vs. night-restricted) and three genotypes (WT, full-body clock-deficient *Arntl/Bmal1* KO, liver re-expression *Arntl/Bmal1*). The upper panel indicates that night-restricted feeding exacerbates the rhythm. The middle panel shows that night-restricted feeding restores rhythmicity in an otherwise circadian clock-deficient animal. A similar outcome is seen in the *Bmal1* re-expression animals shown in the lower panel.

(**B**) Analogous to (A) using the data from Atger et al. [7] that contains both mRNA and pre-mRNA reads. Thus, under night-restricted feeding, wild-type and clock-deficient *Arntl/Bmal1* knockout animals show near-identical *Tfrc* oscillations (upper panel). These oscillations are post-transcriptionally generated, given that *Tfrc* pre-mRNA levels are not rhythmic (lower panel).

(**C**) Analogous to (A) and (B) using the data from Greenwell et al. [61]. The upper panel compares *ad libitum* and night-restricted feeding effect on *Tfrc*, with the latter showing higher amplitude rhythms. The lower panel shows the loss of *Tfrc* rhythms in wild-type animals (i.e., containing a functional clock) when feeding becomes gradually less rhythmic.

(**D**) Analysis of dataset from Manella et al. [62] which compares wild-type animals (i.e. with a functional circadian clock) that are *ad libitum*-fed vs. exclusively day-fed. Day-feeding inverts the *Tfrc* rhythmicity.

1135 **Figures**

**A**

Dataset:
Greco et al., Science Advances 2021
*Tfrc* RNA-seq data (mRNA)
feeding: *ad libitum* & night-restricted
mice: WT, *Bmal1* KO,
liver-specific *Bmal1* re-expression

**B**

Dataset:
Atger et al., PNAS 2015
*Tfrc* RNA-seq data (mRNA & pre-mRNA)
feeding: night-restricted
mice: WT, *Bmal1* KO

**C**

Dataset:
Greenwell et al., Cell Reports 2019
*Tfrc* RNA-seq data (mRNA)
feeding: *ad libitum*, night-restricted,
arrhythmic feeding & damped rhythm
mice: WT

**D**

Dataset:
Manella et al., Nature Metabolism 2021
*Tfrc* RNA-seq data (mRNA)
feeding: *ad libitum* & day-restricted
mice: WT