Mapping Alzheimer's Molecular Pathologies in Large-Scale Connectomics Data: A Publicly Accessible Correlative Microscopy Resource

Xiaomeng Han[1*], Peter H. Li[2], Shuohong Wang[1], Tim Blakely[2], Sneha Aggarwal[3], Bhavika Gopalani[3], Morgan Sanchez[4], Richard Schalek[1], Yaron Meirovitch[1], Zudi Lin[5], Daniel Berger[1], Yuelong Wu[1], Fatima Aly[6], Sylvie Bay [7], Benoît Delatour [8], Pierre Lafaye[9], Hanspeter Pfister[5], Donglai Wei[3], Viren Jain[2], Hidde Ploegh[10], Jeff Lichtman[1*]

[1] Department of Molecular and Cellular Biology, Harvard University, Cambridge, MA

[2] Google Research, Mountain View, CA

[3] Computer Science Department, Boston College, Chestnut Hill, MA

[4] Department of Biomedical Informatics, Harvard Medical School, MA

[5] School of Engineering and Applied Sciences, Harvard University, Cambridge, MA

[6] Harvard College, MA

[7] Institut Pasteur, Université Paris Cité, CNRS UMR 3523, Unité de Chimie des Biomolécules; Paris, France

[8] Paris Brain Institute (ICM), CNRS UMR 7225, INSERM U1127, Sorbonne Université, Hôpital de la Pitié-Salpêtrière; Paris, France

[9] Institut Pasteur, Université Paris Cité, CNRS UMR 3528, Antibody Engineering Platform; Paris, France

[10] Program of Cellular and Molecular Medicine, Boston Children's Hospital, Boston, MA

* Corresponding Authors: xiaomenghan@fas.harvard.edu; jeff@mcb.harvard.edu

Abstract

Connectomics using volume-electron-microscopy enables mapping and analysis of neuronal networks, revealing insights into neural circuit function and dysfunction. In Alzheimer's disease (AD), where amyloid-β (Aβ) and hyperphosphorylated-Tau (pTau) are implicated, connectomics offers an approach to unravel how these molecules contribute to circuit alterations by enabling the study of these molecules within the context of the complete local neuronal and glial milieu. We present a volumetric-correlated-light-and-electron microscopy (vCLEM) protocol using fluorescent nanobodies to localize Aβ and pTau within a large-scale connectomics dataset from the hippocampus of the 3xTg AD mouse model. A key outcome of this work is a publicly accessible vCLEM dataset, featuring fluorescent labeling of Aβ and pTau in the ultrastructural context with segmented neurons, glia, and synapses. This dataset provides a unique resource for exploring AD pathology in the context of connectomics and fosters collaborative opportunities in neurodegenerative disease research. As a proof-of-principle, we uncovered new localizations of Aβ and pTau, including pTau-positive spine-like protrusions at the axon initial segment and changes in the number and size of synapses near Aβ plaques. Our vCLEM approach facilitates the discovery of both molecular and structural alterations within large-scale EM data, advancing connectomics research in Alzheimer's and other neurodegenerative diseases.

Keywords

Alzheimer's disease, Aβ, tau, nanobody, volume electron microscopy, connectomics, correlated light and electron microscopy

Introduction

Introduction

Alzheimer's disease (AD) and related dementias are escalating global health crises, affecting

over 6 million people in the U.S., with care costs projected to surpass $1 trillion annually by 2050 ("2024

Alzheimer's Disease Facts and Figures" 2024). The annual incidence of dementia is expected to nearly

double by 2060, underscoring the urgent need to uncover mechanisms underlying AD pathology and

develop more effective treatments (M. Fang et al. 2025). Despite decades of research targeting

pathological markers such as amyloid-β (Aβ) and hyperphosphorylated tau (pTau), most therapeutic

efforts have achieved only modest efficacy (Cummings et al. 2024).

Traditional studies of AD often focus on isolated aspects—gene expression changes, protein

alterations, or synaptic density—without capturing how these abnormalities interact within neuronal

networks, a key driver of cognitive impairments (Yu, Sporns, and Saykin 2021). The field has been asking

for integrative approaches that can access perturbation of the diseases at molecular, cellular as well as

anatomical levels (Urbanski et al. 2019; Wood, Winslow, and Strasser 2015). Volume electron microscopy

(volume EM) (Peddie et al. 2022)-based connectomics (Swanson and Lichtman 2016) offers a

transformative framework for studying these interactions by providing nanoscale insights into neuronal

and synaptic organization while simultaneously revealing cellular and subcellular pathology. This

approach, while not yet extensively applied to AD, has been used in human tissue studies (Loomba et al.

2022; Shapson-Coe et al. 2024; Karlupia et al. 2023) and holds promise for advancing AD research.

Volume EM techniques, such as focused ion beam-scanning electron microscopy (FIB-SEM)

(Knott et al. 2008), have provided valuable insights into synapse morphology and density in AD samples

(Montero-Crespo et al. 2021; Domínguez-Álvaro et al. 2018; Blazquez-Llorca et al. 2013). However, their

limited imaging volume (~10 µm³) restricts the study of larger networks. Advances in ATUM-SEM

(Hayworth et al. 2006; Kasthuri et al. 2015) and TEMCA (Lee et al. 2016) have enabled datasets on the

millimeter scale, encompassing thousands of neurons and more comprehensive circuits (MICrONS

Consortium et al. 2021; Loomba et al. 2022; Motta et al. 2019; Sievers et al. 2024; Nguyen et al. 2022;

Kuan et al. 2024; Shapson-Coe et al. 2024). Yet, datasets specific to AD remain small and fragmented,

hindering a holistic understanding of the disease. A hybrid method, ATUM-FIB, combines large-scale

context information with local high-resolution FIB volumes and has been applied to samples from the AD

mouse model 5xFAD (Kislinger, Fabig, et al. 2023). Despite these advancements, existing Alzheimer's

datasets remain small and fragmented, with the largest ATUM-SEM dataset for AD models covering only

~45,000 cubic microns (Jiang et al. 2022; Pang et al. 2022; Kislinger, Niemann, et al. 2023). The scarcity

of larger-scale vEM datasets restricts the ability to comprehensively analyze and interpret alterations in

neuronal networks and cellular structures, which are critical for understanding the mechanisms of AD and

related dementias.

In addition to neuronal connectivity disruptions, AD is characterized by pathological protein

aggregates, including amyloid-β (Aβ) (Taylor, Hardy, and Fischbeck 2002; G.-F. Chen et al. 2017) and

hyperphosphorylated tau (pTau) (Mandelkow and Mandelkow 1998; Noble et al. 2013) in the brain

parenchyma. Despite their importance, there is very limited knowledge of how these hallmark pathological

proteins exhibit distinct and identifiable ultrastructural features in EM datasets, making additional

molecular labeling essential for their reliable detection. Extracellular Aβ accumulates into dense-core or

diffuse plaques, which are detectable by EM (Blazquez-Llorca et al. 2013), while intracellular Aβ—

potentially more neurotoxic than plaques—has also been identified (Gouras et al. 2010; LaFerla, Green,

and Oddo 2007; Kelly et al. 2021). However, the precise manifestation of intracellular Aβ in EM images

remains elusive, underscoring the importance of complementary molecular techniques. Similarly, pTau

aggreagtes, identifiable by TEM as either straight or paired helical filaments (Crowther 1991), localize to

cytoplasm, dendrites, and axons. Soluble forms of pTau are considered more neurotoxic than aggregates

(Noble et al. 2013), yet whether either form exhibits specific ultrastructural characteristics detectable in

ATUM-SEM data remains uncertain. These gaps highlight the critical need for innovative molecular

labeling strategies to advance the study of AD pathology within large-scale EM datasets. Traditional

methods for detecting these proteins often rely on detergents, which compromise sample ultrastructure

(Im et al. 2019). Advanced labeling techniques are necessary to localize these pathological proteins

reliably while preserving ultrastructure.

To address these challenges, we employed volumetric correlated light and electron microscopy

(vCLEM), combining detergent-free immuno-probe (nanobody or single-chain variable fragment, scFv) -

assisted immunofluorescence with large-scale ATUM-SEM imaging (T. Fang et al. 2018; Han et al. 2024).

Nanobodies and scFvs, being only one-tenth or one-fifth the size of full-length IgG antibodies (see Figure

1 a for nanobody), exhibit excellent diffusion properties, enabling efficient labeling of intracellular proteins

without requiring detergents. This approach preserves the ultrastructure of the sample (compare Figures

1 b and 1 c) and allows the labeling of multiple proteins in thick, unpermeabilized tissue. The labeled

protein localizations can then be superimposed onto EM data generated by ATUM-SEM from the same

sample, providing a powerful tool to directly link protein alterations to connectomic data. Despite its

potential, vCLEM techniques have not yet been applied to AD or other dementia-related samples to

facilitate research into these diseases.

In this study, we leveraged the vCLEM technique to localize Aβ and pTau in the hippocampal

CA1 region of a female 3xTg AD model mouse. We developed two fluorescent nanobody probes

targeting Aβ and pTau, which proved effective for detergent-free immunofluorescence. Alongside

additional probes for microglia/macrophages (CD11b), astrocytes (GFAP), and blood vessels (Ly-6C/6G),

these enabled precise molecular labeling of thick, unpermeabilized brain tissue. Using this approach, we

generated and share [a large-volume ATUM-SEM-based connectomics dataset](#) with preserved

ultrastructure, integrating molecular and structural information. In the dataset, we were able to localize

extracellular Aβ plaques, intracellular Aβ and pTau at various novel locations.

Machine learning-based segmentation, including style transfer (Michał Januszewski et al. 2018;

Michal Januszewski and Jain 2019) and supervised learning (Lin et al. 2021), facilitated automatic

segmentation of neurons, dendrites, axons, glial processes, and synapses. The fully segmented dataset,

now publicly accessible, supports 3D reconstruction of neurons, glia, their compartments, and synapses.

Importantly, it is designed to be user-friendly for researchers in the AD field who may not be familiar with

volume EM-based connectomics, enabling detailed analyses of cellular and network alterations. As a

proof of principle, we identified previously unreported pTau localizations in enlarged, spine-like structures

at the axon initial segment and observed synaptic changes surrounding an Aβ plaque. These findings

highlight the potential of our dataset to uncover novel pathologies at molecular, cellular, and circuit levels.

Results

Generation of fluorescent nanobody probes for detergent-free immunolabeling of pTau and Aβ

To verify that detergent-free immunolabeling using nanobodies can localize pTau and Aβ in a

vCLEM-compatible manner, we adapted two previously developed and validated nanobodies— A2, which

specifically binds tau protein phosphorylated at multiple sites including S422; R3VQ, which selectively

binds Aβ40/42 without cross-reacting with amyloid precursor protein (APP), the precursor from which Aβ

is derived (T. Li et al. 2016; Lafaye et al. 2019, 2020)—into fluorescent probes (see Table 1). To facilitate

broader use, we designed expression plasmids of these two nanobodies, optimized for easy, high-yield

purification by Expi 293 cells and direct fluorescent dye conjugation (see Methods). These plasmids have

been deposited at Addgene, and we provide a step-by-step protocol (Sup. File 1) to enable the

generation of these probes.

147   **Table 1. Fluorescent nanobody immuno-probes generated in this work.**

Target	Clone No.	Source	Fluorescent dyes conjugated	Link for expression plasmid at Addgene
pTau	A2	(Li et al. 2016)	5-TAMRA, Alexa Fluor 594, Alexa Fluor 647	https://www.addgene.org/233218/
Aβ 40/42	R3VQ	(Li et al. 2016)	5-TAMRA, Alexa Fluor 647	https://www.addgene.org/233217/

We tested two red fluorescent (5-TAMRA-conjugated) A2 and R3VQ probes using our detergent-

free immunofluorescence protocol (see Methods) on tissue sections from the hippocampus CA1 region of

3xTg mice that preserves fine ultrastructure and is compatible with vCLEM (Compare Figure 1 b and c).

This AD mouse model shows neurofibrillary tangles formed by pTau, Aβ extracellular plaques and

intraneuronal Aβ deposits (Oddo et al. 2003).The A2 probe selectively labeled neurofibrillary tangle

(NFT)-like structures in the cell bodies and dendrites of certain neurons (Figure 1 d-1, d-2; Sup. Fig. 1), as

well as the dystrophic neurites nearby plaque material (Figure 1 e-1, e-2; ; Sup. Fig. 1). The R3VQ probe

selectively labeled extracellular plaques (Figure 1 f-1, f-2; ; Sup. Fig. 1) and the cell bodies of certain

neurons (Figure 1 g-1, g-2; ; Sup. Fig. 1).

Interestingly, we also observed R3VQ labeling in neuronal nuclei within neurons showing

intracellular R3VQ signal (Figure 1 g-1, g-2; ; Sup. Fig. 1 e). While nuclear localization of Aβ is not widely

established, a few reports suggest it may occur under certain conditions (Barucker et al. 2014; D'Andrea

et al. 2021). However, given that prolonged fixation with glutaraldehyde was used in our protocol to

preserve ultrastructural, this nuclear signal could also be an artifact of fixation-induced autofluorescence

or redistribution of Aβ. Further investigation, such as alternative fixation protocol and colabeling

approaches, would be required to clarify this observation.

Figure 1. Characterization of fluorescent nanobody probes for pTau and Aβ.

a, Schematic representations of a full-length IgG antibody, a camelid antibody, and a nanobody probe
with a conjugated fluorescent dye. **b**-**c**, Comparison of ultrastructure between a sample (**b**) treated by
Triton X-100 and a sample (**c**) labeled by nanobody probes, untreated with Triton X-100. Blue arrow in **c**
indicates a synapse. **d-1** to **e-2**, Confocal images from the hippocampus of a 3xTg mouse labeled with a
pTau-specific nanobody probe (A2) conjugated with the red dye 5-TAMRA. The panels **d-1**, **e-1** only
show the nanobody labeling. Arrows in **d-1** and **d-2** show the labeled neuronal cell body. Arrowheads in
d-1 and **d-2** show the labeled apical dendrite. Arrows in **e-1** and **e-2** show the labeled neurites. Asterisks
in **e-2** show extracellular plaque material labeled by Hoechst, as reported before (Uchida and Takahashi
2008). **f-1** to **g-2**, Confocal images from the hippocampus of a 3xTg mouse labeled with an Aβ-specific
nanobody probe (R3VQ) conjugated with the red dye 5-TAMRA. The panels **f-1**, **g-1** only show the
nanobody labeling. Asterisks in **f-1** and **f-2** show the labeled plaque material. Asterisks in **g-1** and **g-2**
show the labeled neuronal cell bodies. Arrows in **g-1** and **g-2** show the autofluorescent lipofuscin.

Double immunofluorescence using A2 and anti-pTau mAb AT8, and R3VQ and anti-Aβ mAb 4G8

confirmed colocalization of signals (Sup. Figure a-1 to a-3, b-1 to b-3). No labeling was detected with A2

or R3VQ in sections from age-matched control animals (Sup. Figure 1 d and f). These results

demonstrate that these fluorescent nanobody probes can localize Aβ and pTau in a detergent-free

immunolabeling approach compatible with vCLEM.

Leveraging the flexibility of our protocol for easy conjugation of different fluorophores to the

nanobodies, a broader application of these probes is for multiplex detergent-free immunolabeling. As a

proof-of-principle, we combined these two probes with three additional nanobody probes for multiplex

detergent-free immunolabeling. In one set of experiments, We used three fluorescent nanobody probes

(A2, R3VQ, and anti-CD11b) to label two 120-μm thick brain sections in the absence of detergent (Figure

2 a, b; see Sup. Figure 2 for the anatomical details of the labeled sections) of a one-year-old female 3xTg

mouse. CD11b is a marker for myeloid cells, including microglia and macrophages in the brain

parenchyma (Bisht et al. 2016; Sedgwick et al. 1991; Martin et al. 2017). We acquired a four-color

fluorescence 57-μm image stack (Figure 2 a) from the first section and a 21-μm stack from the second

section (Figure 2 b; see Sup. Figure 2 for the anatomical details of the imaged regions of interest) in the

stratum pyramidale in the CA1 region of the hippocampus. In both image stacks, NFT-like structures in

the cell bodies and dendrites of certain pyramidal neurons and some dystrophic neurites were labeled by

the anti-pTau probe (yellow signals in Figure 2 c, d). In addition, intraneuronal Aβ of certain pyramidal

neurons were labeled by the anti-Aβ probe (magenta signals in Figure 2 c, d). Microglia/macrophages

were labeled by the anti-CD11b probe (green signals in Figure 2 c, d). We noted that these labeled

structures/cells were more abundant, and the labeled apical dendrites appeared to be more tortuous in

the image stack acquired from the more caudal section (Figure 2 c, d). Only in this image stack (Figure 2

a, c) did we find five extracellular Aβ plaques labeled by the anti-Aβ 40/42 probe (magenta signals in

Figure 2 a, c). These findings are consistent with the fact that the pathologies of neurofibrillary tangles

and Aβ plaques progress from caudal to rostral in the hippocampus in the 3xTg mice (Belfiore et al.

2019).

Figure 2. Four-color detergent-free immunofluorescence enabled by nanobody probes.

Maximum intensity projection of the multi-color fluorescence image stacks acquired from the hippocampus at the Bregma -3.78 mm (**a**) or the Bregma -3.28 mm (**b**) of a 3xTg mouse labeled with a CD11b-specific nanobody probe conjugated with Alexa Fluor 488, a pTau-specific nanobody probe conjugated with 5-TAMRA, and an Aβ-specific nanobody probe conjugated with Alexa Fluor 647. Red boxes in the coronal section illustrations show where the image stacks were acquired. The stratum pyramidale of the hippocampal CA1 region is labeled in blue in the illustrations. **c-d**, enlarged boxed insets from **a-b**. Because the red fluorescence signal from the anti-pTau probe was far stronger than the green signal from the anti-CD11b probe, we saw crosstalk in the green channel. This meant that in the overlay, the pTau label appears yellow whereas the microglia appear green, but the two labels were nonetheless unambiguous. Another issue was that aged mice have lipofuscin particles in cell bodies of neurons and glia. This non-specific fluorescence localized to inclusion bodies that were easily differentiated from the membrane labeling of microglia.

In another experiment, we used four fluorescent nanobody probes (A2, anti-CD11b, anti-GFAP,

and anti-Ly-6C/6G) to label a 120-μm thick brain section at Bregma -3.78 mm of another one-year-old

female 3xTg mouse in the absence of detergent. Ly-6C is a lymphocyte antigen expressed by circulating

monocytes recruited to sites of inflammation and endothelial cells of blood vessels (Martínez-Carmona et

al. 2021; Mildner et al. 2007). We acquired a five-color fluorescence image stack of 23 μm (Figure 3).

Numerous activated astrocytes, labeled by the anti-GFAP nanobody probe, were observed nearby

extracellular plaque material and neurons/neurites containing pTau (yellow signals in Figure 3 a, b).

Interestingly, the anti-Ly-6C/6G nanobody labeled not only blood vessels but also some, though not all,

plaque material (white arrows in Figure 3 d), which was surrounded by microglia/macrophages marked by

the anti-CD11b probe (green signals in Figure 3 d). Since Ly-6C labels monocytes recruited to

inflammation sites, which differentiate into macrophages also expressing CD11b (Mildner et al. 2007), it is

likely that the CD11b-positive cells surrounding plaques are macrophages derived from monocytes, rather

than resident microglia.

Figure 3. Five-color detergent-free immunofluorescence enabled by nanobody probes.

a, Maximum intensity projection of the multi-color fluorescence image stacks acquired from the hippocampus of a 3xTg mouse labeled with a CD11b-specific nanobody probe conjugated with Alexa Fluor 488, a GFAP-specific nanobody probe conjugated with 5-TAMRA, a pTau-specific nanobody probe conjugated with Alexa Fluor 594, and a Ly-6C/6G-specific nanobody probe conjugated with Alexa Fluor 647. The signal of each fluorescent dye was pseudo-colored for better visualization. **b-d**, Three-channel maximum intensity projection images of the multi-color fluorescence image stack. White arrows in **d** show the labeled plaque material by the anti-Ly-6C/6G nanobody probe. The arrowhead shows plaque material not labeled by the anti-Ly-6C/6G nanobody probe. Asterisks show the crosstalk signals from the anti-pTau labeling.

Multi-color, large-scale vCLEM dataset with nanobody labeling, segmented neurons, glia and

synapses

From the section shown in Figure 2 a, we acquired a high-resolution vEM dataset measuing 245

µm x 238 µm x 23 µm at a resolution of 4 nm x 4 nm x 30 nm (Figure 4 a) through ATUM-multi-SEM (see

Methods). To the best of our knowledge, this represents the largest vEM volume obtained from an AD

mouse model. Notably, the ultrastructure was well preserved across the entire volume, attributed to the

exclusion of detergents during labeling, enabling the clear identification of various types of synapses

(Figure 4 b, blue arrows).

Figure 4. The four-color vCLEM dataset reveals ultrastructural abnormalities that colocalize with nanobody labeling.

a, The high-resolution vEM volume acquired from the hippocampus CA1 region with four-color immunofluorescence from nanobody probes. The four-color fluorescence data was co-registered with the high-resolution vEM data. **b-1** to **b-4**, the preserved ultrastructure from this vEM volume. The blue arrows

show synapses on a dendritic spine head (**b-1**), a dendritic shaft (**b-2**), a neuronal cell body (**b-3**), and an
axon initial segment (**b-4**). **c**, Visualization of the vCLEM dataset in the Neuroglancer interface, overlaying
fluorescence and EM ultrastructure. The asterisk indicates the cell body of a pyramidal neuron labeled by
the anti-pTau nanobody. The arrow indicates the dystrophic neurite labeled by the anti-pTau nanobody.
The arrowhead indicates the plaque material labeled by the anti-Aβ nanobody. The obelisk indicates the
cell body of a pyramidal neuron labeled by the anti-Aβ nanobody. The hash mark indicates the microglia
labeled by the anti-CD11b nanobody. This region corresponds to the one shown in Figure 2 c.
267
To enable the precise co-registration between the confocal fluorescence volume and the vEM

dataset for generating a vCLEM dataset, the heavy-metal-stained, resin-embedded section was scanned

by X-ray CT (microCT) (see Methods) prior to ATUM sectioning. The microCT image volume revealed

regions with Aβ plaques and neuronal degeneration consistent with previous report (Barbone et al.

2022)(Sup. Figure 3). High-precision co-registration between the fluorescence volume and the vEM

dataset was achieved by identifying corresponding landmarks, such as blood vessels and cell nuclei,

across both volumes. We employed the FIJI/ImageJ plugin BigWarp (Bogovic et al. 2016) to perform a 3D

transformation of the fluorescence volume via thin-plate spline interpolation to co-register the volumes.

The transformed fluorescence volume, overlaid onto more than one-third of the high-resolution EM

volume, is shown in Figure 4 a. This co-registration allows for the precise localization of fluorescence

signals from each nanobody labeling within EM ultrastructure. In Figure 4 c, a representative region

demonstrates that pTau (yellow) overlaps with the cell body of a pyramidal neuron and a dystrophic

neurite, Aβ (magenta) associates with an Aβ plaque and the cell body of another pyramidal neuron, and

CD11b (green) localized to a microglial cell/macrophage. We ingested this vCLEM dataset into

Neuroglancer (Jeremy Maitin-Shepard, et al.), providing a publicly accessible link for users to visualize

the fluorescence and the EM channels, either separately or in combination, and navigate through slices at

different resolution levels (Figure 4 c; Video 1).

Efficient connectomic analysis of large vEM dataset critically depends on the rapid and accurate

segmentation of image data. We employed flood-filling networks (FFNs) (Michał Januszewski et al. 2018)

trained on a human cortex dataset (Shapson-Coe et al. 2024). While the direct application of the pre-

trained FFNs was sufficient for segmenting neuronal cell bodies and dendrites, it performed suboptimally

for segmenting axonal and glial processes (Sup. Figure 4 a). To address this, we utilized Segmentation

Enhanced CycleGANs (Michal Januszewski and Jain 2019) to adjust the visual appearance of this data to

more closely match the H01 dataset prior to segmentation. This approach substantially improved the

segmentation of axonal and glial processes (Sup. Figure 4 b; Figure 5 a). The updated segmentation

results are also publicly available as a segmentation layer ("Segmentation_SECGAN_improved") through

the previously mentioned link. Users can perform 3D reconstructions of neuronal or glial cells by manually

agglomerating segments, simply by selecting the automatically generated segments associated with the

target cell (Figure 5 c; Video1).

Figure 5. Efficient 3D reconstruction and synapse identification in the vCLEM dataset using
segmentation and synapse prediction layers.

a, Visualization of the 3D segmentation layer ("Segmentation_SECGAN_improved") in the Neuroglancer
interface. Each segmented object is uniquely color-coded, and users can click on these objects to
reconstruct them in 3D. **b**,Synapse prediction layer ("Synapse_supervised_learning") visualized in
Neuroglancer. Synapses are labeled with distinct colors for pre-synaptic (e.g., purple) and post-synaptic
(e.g., orange) appositions. The blue arrow highlights a synapse located on a dendritic shaft. Users can
click on these appositions to view their 3D structure. **c**, Example 3D reconstruction of a dendrite (orange)
and six axonal profiles (numbered 1–6, each with a distinct color) using the segmentation and synapse
prediction layers. Synapses formed by these axons are shown in purple. The blue arrow points to
synapses made by axon No. 4.
309
Another key requirement for connectomic analysis is the comprehensive annotation and

segmentation of all synapses. This is particularly crucial in AD studies, given the well-documented

synaptic alterations (Scheff and Price 2006; Montero-Crespo et al. 2021, 2020) and the observed

correlation between cognitive decline and reduced synapse density (DeKosky and Scheff 1990; Davies et

al. 1987; Tzioras et al. 2023). Our vCLEM dataset offers the necessary resolution to precisely locate and

analyze synapses at the ultrastructural level (Figure 4 b-1 to b-4). While manual annotation for synapse

analysis is straightforward, it is not feasible for large-scale studies. Machine-learning-based automated

synapse detection (Çiçek et al. 2016; Parag et al. 2019; Lin et al. 2021; Buhmann et al. 2021) has been

successfully applied to large-scale connectomics datasets (Shapson-Coe et al. 2024; Turner et al. 2022;

Scheffer et al. 2020), demonstrating efficiency, objectivity, and accuracy.

For automatic synapse prediction, we employed the U-Net deep learning model (Parag et al.

2019) to identify the pre- and postsynaptic components of each synapse. Two approaches were explored:

a supervised learning approach and a transfer learning approach, utilizing two existing synapse prediction

frameworks (Lin et al. 2021; Shapson-Coe et al. 2024). For evaluation, we manually annotated all

synapses within a small subvolume measuring 6 μm × 6 μm × 1.5 μm.

In the supervised learning approach, synapses were annotated within a randomly selected

subvolume measuring 6 μm × 6 μm × 6 μm, distinct from the evaluation subvolume. This training dataset

primarily consisted of neuropil and excluded regions containing blood vessels or cell bodies. Following

the guidelines of an online tutorial (Lin et al. 2021), we trained the synapse prediction model and applied

the watershed transform to convert the predictions into synapse instances (see Methods). On the

evaluation subvolume, this model achieved a precision of 86.06% and a recall of 98.88%. Applied to the

entire dataset, the model identified approximately 3.5 million synapses. This first version of synapse

predictions is publicly available via the aforementioned repository as an independent data layer

("Synapse_supervised_learning") (Figure 5 b). In this layer, each synapse is labeled with one presynaptic

segment and one postsynaptic segment, assigned consecutive segment IDs (Figure 5 b). Users can

identify synapses in EM images and visualize them in 3D by interacting with the corresponding segments

(Figure 5 c; Video 1). Additionally, these 3D visualizations can be integrated with cell reconstructions for

advanced analysis (Figure 5 c; Video 1).

While effective, the supervised approach missed many synapses due to the limited diversity in

the small training dataset. To address this limitation, we adopted a transfer learning approach using a

pretrained synapse prediction model developed for a human cortex dataset (Shapson-Coe et al. 2024).

To adapt it to this dataset, we first ran the synapse models trained on the human cortex dataset on this

data with SECGAN style-transfer applied to mimic the human dataset. This produced 0.225 million

synaptic masks with low false positive rates, but with a significant number of false negatives (estimated at

about 50%), particularly within neuronal processes (Sup Figure 5 a-1 to a-3) which require filtering. We

then used these masks as model-produced ground truth to retrain the synapse model from scratch on this

dataset without SECGAN. The retrained model produced synapses with significantly improved precision

of 46% and recall of 52%. Applied to the entire dataset, the model's predictions were filtered to retain only

synapses associated with neuronal membranes, resulting in the identification of approximately 0.7 million

synapses. This second version of synapse predictions is also publicly available as a separate data layer

("Synapse_transfer_learning") (Sup Figure 5 b-2). Like the first version, it enables users to identify,

visualize, and reconstruct synapses in 3D. This version proved particularly effective in detecting synapses

missed by the supervised approach (Sup Figure 5 b-2), leveraging the pretrained model's knowledge of

synapse morphology in the human cortex. Since both synapse detection layers are accessible in

Neuroglancer, users can combine them, resulting in a union that achieves greater precision than either

model alone. This dual-layer approach serves as a complementary resource for synapse identification

and analysis, ensuring a more comprehensive and accurate mapping of synaptic structures.

The publicly accessible dataset is among the largest vEM datasets generated for an AD model,

offering unique opportunities for the scientific community. With fully segmented neuronal and glial cells,

as well as synapses, it provides an invaluable resource for researchers, particularly those studying AD

and other dementias who may be less familiar with volume EM-based connectomics. This innovative

dataset enables the following: (1) pinpointing cells and cellular structures where Aβ and pTau are

localized using overlapping fluorescence signal layers; (2) reconstructing affected neurons and glial cells

at nanometer resolution with the 3D segmentation layer to understand how AD-related abnormalities

manifest in cellular morphology; (3) exploring how the localization of Aβ and pTau impacts synaptic

connectivity and neuronal networks using the synapse segmentation layer; (4) generating new

hypotheses based on findings from the dataset to be tested with other methodologies; and (5) comparing

findings with those reported in the literature or observed in other animal models or human tissues. By

lowering the barrier to entry for researchers unfamiliar with volume EM or segmentation tools, this dataset

fosters broader participation in connectomics-driven AD research and encourages cross-disciplinary

collaboration.

vCLEM dataset uncovers ultrastructural pathologies that colocalize with nanobody labeling

To demonstrate the potential of this dataset for advancing AD research, we conducted an

analysis of sites labeled with each of the three molecular markers (pTau, Aβ, CD11b). This analysis

revealed an array of previously uncharacterized ultrastructural abnormalities colocalize with each marker,

providing insights into molecular and structural pathology at the nanoscale. These findings, along with

their spatial coordinates in the dataset, are compiled in Table 2 and discussed in detail in the following

paragraphs, providing a valuable resource for further investigation. Researchers can leverage this dataset

for in-depth 3D reconstruction of affected cells and subcellular structures, enabling new avenues for

identifying and understanding AD-related pathology.

Table 2. Ultrastructural Pathologies that Colocalize with Nanobody Labeling

Label	Location Type	Subtype	Other Ultrastructural Property	Coordinates
pTau				
	Neuronal soma			
		Only pTau		
			pTau with straight filament	(26467, 24370, 155)

			pTau with no straight filament, but short-filamanet	(28449, 33128, 151), (35520, 32879, 177), (19278, 27572, 51), (20141, 28463, 466)
		Both pTau and A-beta		(30106, 31592, 365), (18693, 21208, 510)
	Dendrite			
		Straight		(25988, 27500, 121), (28382, 36337, 121), (35212, 35235, 142)
		Tortuous		(21209, 33200, 220), (25698, 33034, 266), (21539, 35309, 235)
	Axon			
		Bleb-like structure from AIS		(35464, 30205, 439), (34864, 30905, 460)
		Bleb-like structure from myelinated axon		(28612, 18697, 157)
		Swelling		(28301, 30383, 77)
	Glia			(32846, 20364, 661)
Aβ	Plaque			(34331, 29527, 215), (32240, 12443, 266), (36268, 27060, 438), (32071, 16340, 110), (39797, 13403, 748)
	Neuronal soma	Only Aβ		(23970, 30423, 166), (33496, 33578, 365)
CD11b	Cell associated with A-beta plaques			(32878, 30971, 183), (34740, 27790, 255), (33988, 28188, 7), (37528, 35367, 55), (36983, 26638, 404)
	Cell associated with dystrophic neurites			(24013, 20988, 186), (31741, 19684, 293), (37100, 25861, 659), (29528, 12771, 599)
	Cell adjacent to neuron with intraneuronal A-beta			(23113, 30303, 100)

*The coordinates can be entered into Neuroglancer as in Sup. Figure 6 to move to the location.

pTau: Our analysis revealed a diverse array of pTau labeling patterns across different

ultrastructural contexts, highlighting a complex landscape of pTau-associated abnormalities. We found

NFT-like structures from the labeling of the anti-pTau nanobody localized within cell bodies of seven

pyramidal neurons and one glial cell (red fluorescence, see Figure 6 a-1 to d-2 for four examples for

neurons, and j-1 and j-2 for the glial cell) among the ~100 neuronal and glial cells in the volume where the

fluorescence signals were co-registered. Only one of the seven pTau positive pyramidal cells (which did

not show intraneuronal Aβ), showed pathological straight filament-like structures (Figure 6 a-1, a-2; Sup.

Figure 7 a), which is one of the ultrastructural manifestations of neurofibrillary tangles (Crowther 1991).

The other pTau positive pyramidal cells showed short filament-like structures in their cell bodies (Figure 6

b-2, c-2). We suspect that these are not pathological because in other pyramidal neurons free of pTau

labeling, similar structures were seen (Sup. Figure 7 c-1, c-2). Two of the pyramidal cells also showed

intraneuronal Aβ (magenta fluorescence, see Figure 6 d-1, d-2 for one example). These two neurons had

the same morphological features as neurons that only had intraneuronal Aβ (see below). We also found

NFT-like structures localized in dendrites that were either straight or tortuous (Figure 6 e-1 to f-2). Again,

we did not find specific ultrastructural manifestations of the neurofibrillary tangle structures in these

dendrites despite them being pTau positive (Sup. Figure 7 d-1, d-2). We examined four axonal sites with

pTau labeling. Two of these sites consisted of pTau positive, bleb-like structures each filled with straight

filaments and stemmed from the axon initial segment of a pTau positive pyramidal neuron (see Figure 6

g-1, g-2; see Sup. Figure 7 b for one of them). Another site showed an enlarged pTau positive swelling

filled with lipofuscin granules near the axon initial segment of a different pTau positive pyramidal neuron

(Figure 6 h-1, h-2). The fourth side contained a large pTau positive bleb-like structure filled with

degenerating organelles, emerging from a myelinated axon (Figure 6 i-1, i-2), similar to what has been

seen previously (Gowrishankar et al. 2015). The labeled glial cell had glial inclusions known as "coiled

bodies" (Ferrer et al. 2024; T. Li et al. 2016) which were specifically labeled by the anti-pTau probe

(Figure 6 j-1, j-2). Based on our observation that not all pTau-labeled sites exhibited characteristic

ultrastructural features of neurofibrillary tangles (NFTs), we propose that certain NFT-like structures or

pTau aggregates may lack visible ultrastructural representations. This underscores the importance of

using correlative microscopy and labeling technique like ours to identify neurons harboring these

aggregates in EM micrographs, as they may otherwise go undetected.

Figure 6. Ultrastructural abnormalities that colocalize with the labeling of pTau.

a-1, red fluorescence from the anti-pTau nanobody overlaps with the cell body of a pyramidal neuron. **a-2**,
enlarged inset from **a-1**. The blue arrow indicates the straight filament-like structure. **b-1** and **c-1**, red
fluorescence from the anti-pTau nanobody overlaps with the cell bodies of two pyramidal neurons. **b-2**
and **c-2**, enlarged insets from **b-1** and **c-1**. The red arrows indicate the short filament-like structures that
were also seen in neurons that were not labeled by the anti-pTau nanobody (see Sup. Figure 4 c-1, c-2).
d-1 and **d-2**, the pyramidal neuron that was labeled by both the anti-pTau nanobody (red fluorescence in
d-1) and the anti-Aβ nanobody (magenta fluorescence in **d-2**). **e-1** and **f-1**, red fluorescence from the anti-
pTau nanobody overlaps with a straight apical dendrite and a tortuous apical dendrite of two pyramidal
neurons. **e-2** and **f-2**, enlarged insets from **e-1** and **f-1**. The blue arrow in **e-2** indicates the short filament-
like structure that was also seen in the apical dendrite that was not labeled by the anti-pTau nanobody
(see Sup. Figure 4 d-1, d-2). **g-1**, red fluorescence from the anti-pTau nanobody overlaps with a bleb-
structure connected to an axon. **g-2**, enlarged inset from **g-1**. The blue arrow indicates the straight
filament-like structure. **h-1**, red fluorescence from the anti-pTau nanobody overlaps with an enlarged
swelling of an axon filled with lipofuscin granules. **h-2**, enlarged inset from **h-1**. The asterisk shows the
lipofuscin granules. **i-1**, red fluorescence from the anti-pTau nanobody overlaps with a bleb-like structure
that emerged from a myelinated axon. This bleb was filled with degenerating organelles. The blue arrow
indicates the end of the myelination. **i-2**, enlarged inset from showing the degenerating organelles. **j-1**
and **j-2**, red fluorescence from the anti-pTau nanobody overlaps with a glial cell that had coiled bodies
(red fluorescence in **j-1**).
433
Aβ: We found extracellular Aβ plaque structures from the labeling of the anti-Aβ nanobody

(magenta fluorescence in Figure 7) localized within five separate spots of plaque material (see Figure 7 a-

1 to b-2 for two examples) which showed the typical ultrastructure of dense-cored plaques (Blazquez-

Llorca et al., 2013). These plaques had dense cores of Aβ deposits in the center, Aβ fibrils, glial

processes and dystrophic neurites surrounding them. Intraneuronal Aβ localized within neurons that had

very tortuous apical dendrites, shrunken cell bodies, dark osmiophilic cytoplasm and degenerating

organelles (Figure 7 c-1 to d-2). These neurons with intraneuronal Aβ appeared to be more abnormal

than neurons that only had pTau.

Figure 7. Ultrastructural abnormalities that colocalize with the labeling of Aβ.

a-1, magenta fluorescence from the anti-Aβ nanobody overlaps with plaque material. **a-2**, enlarged inset
from **a-1**. **b-1**, magenta fluorescence from the anti-Aβ nanobody overlaps with another plaque material. **b-2**, enlarged inset from **b-1**. **c-1**, magenta fluorescence from the anti-Aβ nanobody overlaps with the cell
2, enlarged inset from **b-1**. **c-1**, magenta fluorescence from the anti-Aβ nanobody overlaps with the cell
body of a pyramidal neuron. **c-2**, enlarged inset from **c-1** showing the degenerating organelles in this
neuron. **d-1**, magenta fluorescence from the anti-Aβ nanobody overlaps with the cell body of another
pyramidal neuron. **d-2**, enlarged inset from **d-1** showing the dilated endoplasmic reticula in this neuron.

CD11b: CD11b is a myeloid cell marker that labels both resident microglia and infiltrating

macrophages within the brain parenchyma (Sedgwick et al. 1991; Martin et al. 2017). As a component of

the phagocytic receptor CR3, CD11b is present on the plasma membrane of glial processes that surround

synaptic elements (Bisht et al. 2016). Among these cells, some microglia, characterized by a condensed,

electron-dense cytoplasm and nucleoplasm, were referred to as 'dark microglia' in a previous electron

microscopy studies (Bisht et al. 2016; St-Pierre et al. 2022). In our vCLEM dataset, we identified ten

labeled cells based on anti-CD11b nanobody labeling (green fluorescence in Figure 8; five examples

shown), although we could not distinguish between microglia and macrophages. Among them, five were

associated with Aβ plaques (Figure 8 a-1, c-3, c-4), extending ramified processes to encircle and perhaps

phagocytose plaque material (Figure 8 b); four however were associated with dystrophic neurites where

there was no Aβ plaque (Figure 8 c-1); one was adjacent to a neuron that had intraneuronal Aβ (Figure 8

c-2). These cells appeared stressed: they had a large amount of lipofuscin granules, osmiophilic

cytoplasm, altered mitochondria, condensed nucleoplasm, and dilated endoplasmic reticulum (Figure 8 a-

2, a-3), which may be attributed to the activation of the integrated stress response (ISR) as previously

described (Flury et al. 2024).

Figure 8. Ultrastructural abnormalities that colocalize with the labeling of CD11b.

a-1, green fluorescence from the anti-CD11b nanobody overlaps with a microglial cell. **a-2**, the nucleus of
the microglial cell in **a-1**. Asterisk indicates the condensed nucleoplasm. **a-3**, enlarged inset from **a-1**.
Asterisk indicates the altered mitochondria. **b**, green fluorescence from the anti-CD11b nanobody
overlaps with the glial processes encircling plaque material, which shows overlapped magenta
fluorescence from the anti-Aβ nanobody. **c-1**, green fluorescence from the anti-CD11b nanobody
overlaps with a microglial cell nearby dystrophic neurites (asterisks). **c-2**, green fluorescence from the
anti-CD11b nanobody overlaps with a microglial cell next to a pyramidal neuron (the same neuron in
Figure 6 c-1). **c-3** and **c-4**, green fluorescence from the anti-CD11b nanobody overlaps with two

microglial cells nearby plaque material, which shows overlapped magenta fluorescence from the anti-Aβ
nanobody.
478
3D Reconstructions Reveal Detailed Morphological Features of Neurons with Intraneuronal Aβ or

pTau

As a proof-of-principle demonstration of the dataset's potential, we conducted 3D reconstructions

of neurons with intraneuronal Aβ or pTau, showcasing the dataset's ability to uncover detailed,

unreported morphological features associated with these pathologies. Using the automatic segmentation

followed by manual proofreading, a pyramidal neuron with intraneuronal Aβ (Figure 7 c-1) and two

pyramidal neurons with intraneuronal pTau (Figure 6 b-1, c-1) were reconstructed.

Figure 9. 3D reconstruction of a neuron with intracellular Aβ and two neurons with intracellular pTau.

a, 3D reconstruction of three pyramidal neurons. The insets pointing to the cell bodies show the labeling

of Aβ or pTau by nanobody probes. Insets 1 to 3 show segments of spiny dendrites of Neuron 1 to 3. **b**
shows the AIS of Neuron 2 as labeled in **a**. White arrow, the swelling filled with pTau. White arrow heads,
spines on AIS. **c** shows the AIS of Neuron 3 as labeled in **a**. White arrows, two pTau positive blebs
connected to the AIS. White arrows, thin stalks connecting the blebs to the AIS. **d-1** and **d-2**, 2D
segmentations of the two pTau positive blebs filled with straight filaments. **e-1** and **e-2** show that each of
the blebs was adjacent to an axon profile filled with synaptic vesicles.
496
The cell body of neuron 1 with intraneuronal Aβ labeling appeared abnormal as if it were in the

process of degenerating (Figure 7 c-1, c-2). A microglial cell was immediately adjacent raising the

possibility that the neuron would soon be phagocytosed (Figure 8 c-2). Surprisingly, this neuron still

retained multiple spiny dendrites (Figure 9 a, inset 1) and its axon projected normally for ~100 μm without

myelin and then showed normal myelination (Figure 9 a). Neuron 2 showed intraneuronal pTau labeling in

its soma (Figure 6 b-1, b-2). Nonetheless it also had multiple normal looking spiny dendrites (Figure 9 a,

inset 2). This neuron's axon ran normally ~80 μm before becoming myelinated (Figure 9 a). As was the

case for the other two neurons, there were several spine-like structures innervated by vesicle-filled axonal

profiles (Figure 9 b) along this neuron's axon initial segment (AIS). This AIS also had an abnormally

enlarged swelling that was positive for anti-pTau nanobody labeling. This swelling was filled with

lipofuscin granules (see Figure 6 h-1, h-2). The third neuron (Neuron 3) with intraneuronal pTau labeling

(Figure 6 c-1, c-2) was located such that its soma was adjacent to an Aβ plaque (Figure 9 a). This neuron

also possessed multiple normal looking spiny dendrites (Figure 9 a inset 3) and an axon with normal

appearance that ran ~60 μm before becoming myelinated (Figure 9 a). Two spine-like swollen

protrusions emerged from its AIS via narrow stalks (Figure 9 a box c, d-1, d-2; Figure 9 c). Each of these

was filled with straight filaments, similar to those described above (Figure 9 d-1, d-2; also see Figure 6 g-

1, g-2 and Sup. Figure 7 b). We suspect that these objects are abnormally enlarged postsynaptic spine

heads because they were innervated by vesicle-filled axonal profiles (Figure 9 e-1, e-2). The protrusion

stalks are extremely thin (less than 1 μm in diameter), suggesting the possibility that these structures may

eventually detach from the AIS. Supporting this hypothesis, we identified a nearby pTau-filled bleb that

appeared disconnected from the AIS or any adjacent structures (Sup. Figure 8).

Analysis of synapse nearby an Aβ plaque reveals alterations in synapse density and size

As another proof-of-principle demonstration of the robustness and utility of our dataset, we

applied the synapse prediction by supervised learning that has high precision and recall to investigate

synaptic alterations in proximity to an Aβ plaque—an area of critical interest given the central role of

synaptic pathology in the etiology, manifestation, and progression of AD (Tzioras et al. 2023; Koffie,

Hyman, and Spires-Jones 2011). We randomly selected nine 6 x 6 x 6 µm regions predominantly

comprising neuropil. These regions included three representative areas located at distances of 10 µm, 20

µm, and 30 µm from the plaque center (Sup. Figure 9).

Figure 10. Changes in synapse density and total synapse volume with relationship to the distance
from an extracellular Aβ plaque.

a-1, A representative image slice with detected synapses from the box at 10 μm away from the center of
an Aβ plaque. **a-2** to **a-4**, 3D rendering of all detected synapses in the box as in **a-1** with the dimensions
of 6 μm x 6 μm x 6 μm at 10 μm away from the center of an Aβ plaque. White block is of the dimension of
1 μm x 1 μm x 1 μm. **b-1**, A representative image slice with detected synapses from the box at 20 μm
away from the center of an Aβ plaque. The image slice is of the dimension of 6 μm x 6 μm. **b-2** to **b-4**, 3D
rendering of all detected synapses in the box as in **b-1** with the dimensions of 6 μm x 6 μm x 6 μm at 20
μm away from the center of an Aβ plaque. White block is of the dimension of 1 μm x 1 μm x 1 μm. **c-1**, A
representative image slice with detected synapses from the box at 30 μm away from the center of an Aβ
plaque. The image slice is of the dimension of 6 μm x 6 μm. **c-2** to **c-4**, 3D rendering of all detected
synapses in the box as in **c-1** with the dimensions of 6 μm x 6 μm x 6 μm at 30 μm away from the center
of an Aβ plaque. White block is of the dimension of 1 μm x 1 μm x 1 μm. **d**, Synapse density, **e**, Average
synapse volume, **f**, Total synapse volume were measured for each of the three box at 10 μm away from
the plaque, the three box at 20 μm away from the plaque, and the three box at 30 μm away from the
plaque. Two-tailed, unpaired t-test was performed. Synapse density, 10 μm vs. 20 μm, p = 0.0004,
t=11.29, df=4; 20 μm vs. 30 μm, p = 0.0051, t=5.553, df=4; 10 μm vs. 30 μm, p = 0.0007, t=9.597, df=4.
Average synapse volume, 10 μm vs. 20 μm, p = 0.4356, t=0.8655, df=4; 20 μm vs. 30 μm, p = 0.6104,
t=0.5519, df=4; 10 μm vs. 30 μm, p = 0.3864, t=0.9713, df=4. Total synapse volume, 10 μm vs. 20 μm,
0.0009, t=8.789, df=4; 20 μm vs. 30 μm, p = 0.0210, t=3.690, df=4; 10 μm vs. 30 μm, p = 0.0029, t=6.489,
df=4.
548
Synapse detection on all nine boxes (Figure 10 a-1 to a-3, b-1 to b-3, c-1 to c-3) showed that a

decrease in synapse density as regions progressively closer to the plaque: synapse density was 0.181 ±

0.068 per μm³ at 10 μm, 0.653 ± 0.025 per μm³ at 20 μm, 1.216 ± 0.174 per μm³ at 30 μm (Figure 10 d).

There was a statistically significant trend as the distance to the plaques dropped from 30 μm to 10 μm

(two-tailed, unpaired t-test. 10 μm vs. 20 μm, p = 0.0004; 20 μm vs. 30 μm, p = 0.0051; 10 μm vs. 30 μm,

p = 0.0007).

The average synapse volume (pre-synaptic side) was 0.00436 ± 0.00007 μm³ at 10 μm, 0.00452

± 0.00030 μm³ at 20 μm, 0.00476 ± 0.00071 μm³ at 30 μm (Figure 10 e). There was no statistically

significant change in average synapse volume as the distance to the plaques dropped from 30 μm to 10

μm (two-tailed, unpaired t-test. 10 μm vs. 20 μm, p = 0.4356; 20 μm vs. 30 μm, p = 0.6104; 10 μm vs. 30

μm, p = 0.3864).

The total synaptic volume (pre-synaptic side) for each was 0.170 ± 0.065 μm³ at 10 μm, 0.638 ±

0.066 μm³ at 20 μm, 1.255 ± 0.282 μm³ at 30 μm (Figure 10 f). There was a statistically significant

decrease in total synaptic volume as the distance to the plaques dropped from 30 μm to 10 μm (two-

tailed, unpaired t-test. 10 μm vs. 20 μm, p = 0.0009; 20 μm vs. 30 μm, p = 0.0210; 10 μm vs. 30 μm, p =

0.0029).

To investigate whether smaller synapses are more susceptible to elimination, suggested by

previous studies (John and Reddy 2021; Meftah and Gan 2023)—we combined synapse volume data

(pre-synaptic + post-synaptic) from the three boxes at each distance and analyzed the frequency

distributions (Sup. Figure 10). We performed two-sample Kolmogorov–Smirnov tests and found there was

no significant difference between any two of the distributions (10 μm vs. 20 μm: p = 0.8498; 20 μm vs. 30

μm: p = 0.1231; 10 μm vs. 30 μm: p = 0.09247). This result suggests that there was synapse elimination

closer to the plaque, but the elimination did not preferentially target smaller nor larger synapses.

Synapses of all volumes were possibly eliminated equally when closer to the plaque. To further test

whether one of the pre-synaptic site or post-synaptic is affected more, we plotted the frequency

distributions of pre- or post-synaptic volumes (pre-synaptic, Sup. Figure 11 a; post-synaptic, Sup. Figure

11 b) for all nine boxes. We found that the pre-synaptic volumes of all boxes were slightly larger than the

post-synaptic volumes (Sup. Figure 11 c).

Conclusions and Discussion

This work introduces a volumetric correlated light and electron microscopy (vCLEM) technique

using fluorescent nanobody immunolabeling to localize Alzheimer's disease (AD)-related molecules,

including phosphorylated tau (pTau), amyloid-β (Aβ), and CD11b, in a large-scale volume EM dataset

from an AD mouse model. By preserving ultrastructure through detergent-free labeling, this method

addresses a major limitation of conventional volume EM-based connectomics, which lacks molecular

specificity, and overcomes the challenges posed by traditional molecular labeling techniques that often

compromise sample integrity. The resulting dataset enables precise localization of pathological proteins

within their ultrastructural context, allowing for integrated analyses at molecular, cellular, and circuit levels

across a significantly larger volume than previous vEM datasets.

The publicly accessible dataset, featuring advanced machine-learning-based segmentations of

neurons, dendrites, axons, glial processes, and synapses, offers a valuable resource for studying AD

pathology. Researchers can pinpoint cellular and subcellular structures where Aβ and pTau localize,

reconstruct affected cells at nanometer resolution, and investigate how these pathological markers impact

synaptic connectivity and neuronal networks. This dataset lowers the entry barrier for researchers less

familiar with volume EM or segmentation tools, fostering broader participation in connectomics-driven AD

research.

Following the above framework, as a proof-of-principle, we used our dataset to identify novel

pTau-associated abnormalities at the axon initial segment (AIS), including enlarged blebs filled with

straight filaments and pTau-positive swellings containing lipofuscin granules (Figure 6 g-1, g-2). These

findings highlight the complex landscape of pTau pathology, where certain aggregates lack visible

neurofibrillary tangle-like features in EM micrographs. We also observed spine-like structures at the AIS

with thin stalks (Figure 9 c; less than 1 µm in diameter), potentially prone to detachment, raising the

intriguing hypothesis that these structures may contribute to exosome formation, a mechanism implicated

in pTau release in AD (Saman et al. 2012; Miyoshi et al. 2021). These abnormalities suggest disrupted

molecular mechanisms at the AIS, likely involving ankyrin-G (Sobotzik et al. 2009; Zempel et al. 2017) or

diffusion barriers affected by pTau or Aβ (X. Li et al. 2011), leading to tau missorting and synaptic

dysfunction (Blazquez-Llorca et al. 2011). The AIS's established role as a barrier for regulating tau

trafficking underscores the potential of these disruptions to impact synaptic transmission significantly. For

Aβ, in addition to extracellular deposits in plaques, we identified neurons with intraneuronal Aβ deposits

exhibiting severe morphological abnormalities, including tortuous dendrites, shrunken cell bodies, and

degenerating organelles. These features indicate profound cellular dysfunction and are consistent with

Aβ's known neurotoxicity. Interestingly, 3D reconstructions enabled by machine-learning segmentation

provided a detailed view of these affected neurons and revealed novel morphologies not easily captured

by prior FIB-SEM and ATUM-SEM datasets, which were limited to smaller subvolumes (Montero-Crespo

et al. 2021; Domínguez-Álvaro et al. 2018; Jiang et al. 2022; Pang et al. 2022; Blazquez-Llorca et al.

2013).

Our synapse analysis further revealed significant alterations near Aβ plaques. Synapse density

declined sharply with proximity to plaques, reflecting Aβ's synaptotoxic effects (Hampel et al. 2021; Koffie

et al. 2009; Urbanc et al. 2002) and glial-mediated mechanisms (Chung et al. 2015; Rajendran and

Paolicelli 2018; Vilalta and Brown 2018). Reactive glial processes from microglia and astrocytes were

notably abundant near plaques (Figure 8 b, also see Sup. Figure 11 for astrocytic processes), supporting

their role in synaptic pruning and dysfunction. While synapse density decreased, the average synapse

volume remained constant, resulting in a reduction in total synapse volume near plaques. This challenges

earlier findings suggesting compensatory synaptic enlargement with declining synapse density (Scheff

and Price 2001; Neuman et al. 2015; Androuin et al. 2018; Scheff and Price 2003). Instead, our results

align with more recent volume EM studies showing no evidence of synaptic enlargement (Martínez-Serra

et al. 2022). Future classification of synapses by type (e.g., excitatory vs. inhibitory) or location (e.g.,

somatic vs. dendritic) could clarify whether specific subpopulations respond differently to Aβ toxicity.

While the vCLEM approach provides transformative insights, it has limitations. We focused on

two canonical markers—fibrillar Aβ40/42 and pTau phosphorylated at specific sites—without addressing

other molecular forms known to elicit more severe toxicity. Additionally, the limited availability of

nanobody probes constrains the range of molecules that can be studied. The resolution gap between

confocal fluorescence microscopy and EM may introduce slight inaccuracies in registering molecular

signals onto EM micrographs. Furthermore, findings from this single-sample study require validation in

additional datasets, including other animal models and human tissues, to ensure broader applicability.

A key future direction involves expanding the repertoire of fluorescent nanobody probes to target

a broader range of pathological molecules. For instance, probes could be developed for specific Aβ

species (e.g., Aβ1-40 and Aβ1-42) and conformations (e.g., oligomers, protofibrils, and fibrils) (Hampel et

al. 2021), as well as tau isoforms or site-specific phosphorylations (Arendt, Stieler, and Holzer 2016).

These advancements would enable more detailed analyses of molecular variants with distinct toxicities

affecting neurons and synapses. Furthermore, nanobody probes for other neurodegeneration-related

molecules, such as α-synuclein (Twohig and Nielsen 2019), TDP-43 (transactive response DNA-binding

protein of 42 kDa) (Wilson et al. 2011), PLD3 (Nackenoff et al. 2021; Hooli et al. 2015), and SORL1

(sortilin-related receptor 1), which plays a crucial role in neuronal endosome trafficking (Yin, Yu, and Tan

2015; Knupp et al. 2020), as well as huntingtin (Bates et al. 2015), could significantly broaden the utility of

this technique. These additions would expand its application beyond Alzheimer's disease to other forms

of dementia and neurodegenerative disorders. Many nanobodies for these targets already exist (Danis et

al. 2022; De Genst et al. 2010; Habicht et al. 2007; Lafaye et al. 2009) and could be easily adapted into

fluorescent probes. Beyond pathological proteins, this approach can also be extended to cell-type-specific

markers, allowing researchers to investigate how toxic molecules impact distinct cell populations. For

example, nanobody or single-chain variable fragment (scFv)-based probes could target markers for glial

cells or inhibitory neuron subtypes, providing insights into cell-type-specific vulnerabilities and

mechanisms in Alzheimer's disease and other neurodegenerative conditions.

Recent large-scale proteomic and transcriptomic studies (Johnson et al. 2022, 2020; Magistri et

al. 2015; Mathys et al. 2019) have showed alterations at the levels of protein and gene expression in AD.

To localize these new molecular abnormalities in connectomics datasets, a large collection of nanobody

probes that target common molecular markers will be needed. Multiple nanobody databases are now

available where thousands of nanobody sequences are being deposited (Deszyński et al. 2021; Wilton et

al. 2018). Researchers can query sequences of nanobodies that target the molecular abnormalities of

interest, and then, as we have done, convert these nanobodies into fluorescent probes. If no nanobody

for a desired target is available, there are efficient ways to generate new nanobodies from immune

libraries (Muyldermans 2021) or synthetic libraries (McMahon et al., 2018; Zimmermann et al.,

2020)(McMahon et al. 2018; Zimmermann et al. 2020).

With a growing collection of nanobody probes, integrating vCLEM with super-resolution

microscopy (Hoffman et al., n.d.; Tuijtel et al. 2019) could enhance the precision of molecular target

localization within EM micrographs. Combining vCLEM with advanced multiplex imaging techniques, such

as super-multicolor fluorescence imaging enabled by spectral unmixing (Seo et al. 2022) or multiplexed

Raman vibrational imaging (Wei et al. 2017), could allow for the simultaneous labeling of numerous

markers in a single sample. This would unlock unprecedented opportunities for integrative molecular and

structural studies, enabling a deeper understanding of complex pathological processes.

Notably, this technique is well-suited for application to human samples, as it does not rely on

transgenic models and can directly target molecules in human tissues using nanobody probes. The

nanobody probes tested in this study (A2 and R3VQ) have already been shown to effectively detect their

targets in brain samples from AD patients (T. Li et al. 2016). By integrating this approach with recently

developed methods for processing large surgical brain samples (Karlupia et al. 2023; Loomba et al.

2022), vCLEM offers a powerful tool for investigating AD connectomics directly in human tissues,

providing critical insights into disease mechanisms and pathology.

This study highlights the transformative potential of vCLEM for bridging molecular and structural

data in large-scale volume EM datasets, offering a publicly accessible resource to advance integrative

research into AD and other neurodegenerative diseases.

Material and Methods

Animals

682   Animals used in the study were female one-year old 3xTg (B6;129-Tg(APPSwe,tauP301L)1Lfa

683 Psen1$^{tm1Mpm/Mmjax}$) mice (Jackson Laboratory) and female one-year old B6129SF2/J mice (as age-

684 matched control animal, Jackson Laboratory). All experiments using animals were conducted according

685 to US National Institutes of Health guidelines and approved by the Committee on Animal Care at Harvard

686 University.

Nanobody immuno-probe production

A signal peptide (MDWTWRILFLVAAATGAHS) was added to the N'-terminus of the nanobody. A

sortase tag (SLPETGG) and a 6 x His tag was added to the C'-terminus. The amino acid sequence was

reverse translated into a DNA sequence with codon optimization for human cells. The DNA sequence

was synthesized and cloned into pcDNA 3.1 vectors.

Nanobody expression was performed with Expi 293 cells (ThermoFisher). Detailed expression,

purification, and dye conjugation protocols can be found in Sup. File.

Perfusion and fixation

The mouse was anesthetized by isoflurane until there was no toe-pinch reflex. Mice were then

transcardially perfused with aCSF (125 mM NaCl, 26 mM NaHCO$_3$, 1.25 mM NaH$_2$PO$_4$, 2.5 mM KCl, 26

mM glucose, 1 mM MgCl$_2$ and 2 mM CaCl$_2$ (all chemicals from Sigma-Aldrich) at the flow rate of 10

ml/min for 2 min to remove blood, followed with 4% paraformaldehyde (Electron Microscopy Sciences),

0.1% glutaraldehyde (Electron Microscopy Sciences) in 1 x PBS for 3 min for fixation. Brains were

dissected and then post-fixed in the same fixative on a rotator overnight at 4°C. Brains were sectioned

into 50-μm or 120-μm coronal sections using a Leica VT1000 S vibratome and stored in the same fixative

at 4°C.

Immunofluorescence

Detergent-free immunofluorescence labeling was performed with nanobody probes (see Sup.

Table 1 for the nanobody probes and the final concentration of each nanobody probe used in this study).

The detailed protocol was described in (Han et al. 2024). For the 120-µm coronal sections used for

vCLEM, the section was first washed with 1 x PBS for 3 x 10 min, and then blocked in glycine blocking

buffer (0.1 M glycine, 0.05% NaN$_3$ in 1 x PBS) for 1 h on a rotator at 4 °C. The labeling solution was

prepared by diluting nanobody-dye conjugates in glycine blocking buffer (see Sup. Table 1 for the final

concentration used). The labeling solution and any remaining steps were protected from light. The section

was incubated with the labeling solution on a rotator at 4 °C for three to seven days. The 120-µm sections

for Figure 2 were first incubated with the labeling solution with only the anti-pTau nanobody for three

days, and then incubated with the labeling solution with the anti-Aβ and the anti-CD11b nanobodies for

another three days. The 120-µm section for Figure 3 was incubated with the labeling solution with all four

nanobodies for seven days. After the incubation, the section was washed with 1 x PBS for 3 x 10 min,

and then stained with Hoechst 33342 (Invitrogen, diluted 1:5000 in 1 x PBS) for 1 h on a rotator at 4 °C.

The section was washed with 1 x PBS for 3 x 10 min, and then mounted onto glass slide (VWR).

718   To validate the labeling of the nanobody A2, double immunofluorescence labeling with detergent

was performed on 50-µm sections with the anti-pTau mAb AT8 (ThermoFisher) plus the anti-mouse IgG

H + L secondary antibody conjugated with Alexa Fluor 488 (ThermoFisher), and A2 conjugated with Alexa

Fluor 594 (see Sup. Figure 1 a) (see Sup. Table 2 and 3 for the final concentrations used for each probe).

To validate the labeling of the nanobody R3VQ, double immunofluorescence labeling with detergent was

performed with the anti-Aβ mAb 4G8 conjugated with Alexa Fluor 647 (BioLegend) and R3VQ conjugated

with 5-TAMRA (see Sup. Figure 2) (see Sup. Table 2 and 3 for the final concentrations used for each

probe). The detailed protocol is available on the NeuroMab website ("Welcome to NeuroMab!," n.d.). In

brief, 50-µm coronal sections were first washed with 1 x PBS for 3 x 10 min, and then blocked in vehicle

(10% normal goat serum, 0.3% Triton X-100 in 1 x PBS) overnight on a rotator at 4 °C. Subsequent steps

were protected from light. For the experiment with AT8 and A2, the sections were then incubated with

primary antibody solution plus nanobody probes overnight on a rotator at 4 °C. After the incubation,

sections were washed with vehicle for 3 x 10 min, and then incubated with the secondary antibody

solution for 1 hour on a rotator at 4 °C. For the experiment with 4G8 and R3VQ, the sections were then

incubated with primary antibody solution plus nanobody probes for 3 days on a rotator at 4 °C. After the

incubation, sections were washed with 1 x PBS for 3 x 10 min, and then stained with Hoechst (diluted

1:5000 in 1 x PBS) for 1 h on a rotator at 4 °C. Sections were washed with 1 x PBS for 3 x 10 min, and

then mounted onto glass slides.

Fluorescence confocal microscopy

Fluorescence confocal microscopy was performed as described in (Han et al. 2024). In brief, 50-

μm sections were mounted in Vectashield H-1000 (Vector Laboratories) with a #1 coverslip (Electron

Microscopy Sciences) on top sealed with clear nail polish. 120-μm sections (for vCLEM) were mounted in

1 x PBS inside 120-um spacer (Invitrogen) with a #1 coverslip on top sealed with the spacer. Sections

were imaged with a Zeiss LSM 880 confocal laser scanning microscope equipped with either a 20x/0.8

NA air-objective or a 40x/1.1 NA water immersion objective. The 120-μm section for vCLEM was imaged

with a 40x/1.1 NA water immersion objective. Acquisition of double or triple color fluorescent images was

done with appropriate band pass filters for the specific fluorescent dyes to avoid crosstalk.

The brightness, contrast, and gamma of all fluorescent images were adjusted. Fluorescence

image volumes were projected to a single plane by maximum intensity for visualization in 2D.

EM preparation

Sample preparation for electron microscopy was performed as described in (Han et al. 2024).

After the 120-μm sections was imaged, it was transferred into 3.7 ml shell vials with 1 ml secondary

fixative (2% PFA, 2,5% glutaraldehyde in 0.15M sodium cacodylate buffer with 4mM Ca^{2+} and 0.4 mM

Mg^{2+}) and incubated for at least one week on a rotator at 4 °C. A modified ROTO (Reduced Osmium-

Thiocarbohydrazide-Osmium) protocol was used to stain the section. The section was embedded in LX-

112 resin.

X-ray microCT scanning

X-ray micro-computed tomography (μCT) images of the resin-embedded sections were acquired

using a Zeiss Xradia 510 Versa system and Zeiss' Scout and Scan software. Detailed protocol was

described in (Han et al. 2024).

EM imaging

The resin-embedded sections were cut into 30 nm serial ultrathin sections using automated tape-

collecting ultramicrotome (ATUM) (Kasthuri et al. 2015). Serial sections were collected onto carbon-

coated and plasma-treated Kapton tape. The tape was cut into strips and affixed onto 150 mm silicon

wafers (University Wafer). A Zeiss Sigma scanning electron microscope was used to acquire overview

images from the serial sections. Detailed protocol was described in (Han et al. 2024).

Prior to acquiring high-resolution images, the serial section sections on wafers were post-stained

for 4 min with a 3% lead citrate solution. After staining, the sections were degassed for a minimum of 24 h

at 1×10^{-6} Torr. A Zeiss MultiSEM 505 scanning electron microscope equipped with 61 electron beams

was used to acquire high-resolution images from the serial sections. Images were collected using a 1.5-

kV landing energy, 4-nm image pixel, and a 400-ns dwell time.

High-resolution EM image processing

The preparation of the vEM data before it could be segmented/analyzed includes two steps:

affine stitching and elastic alignment. Detailed protocol was described in (Han et al. 2024). We excluded

20 slices that had sub-optimal imaging quality (e.g. very thin slices or contains tears and wrinkles) and

copied the adjacent sections in the places of these excluded slices. The aligned stack was rendered at

full resolution (4 x 4 x 30 nm) and each section was cut into 4k x 4k .png tiles, imported into VAST as a

.vsvi file, and ingested into Neuroglancer for further analysis.

The initial alignment of the vEM dataset showed several jumps through the z-axis. A second

attempt at alignment was performed after discarding four more slices (we also copied the adjacent

sections in the places of these discarded slices) that had suboptimal image quality, which successfully

eliminated the jumps. The 3D reconstruction of Neuron 3, the automatic Aβ plaque detection, and the

synapse detection using a U-Net classifier was performed on the first version of alignment. The 3D

reconstruction of Neuron 3 and the automatic Aβ plaque detection were later transferred to the second

version of the alignment based on matching coordinates. The other reconstructions and the synapse

detection by PyTorch were performed on the second version of the alignment. The publicly accessible

vCLEM dataset shows the second version of the alignment.

Co-registration of fluorescence and EM volumes

The co-registration was performed using the BigWarp plugin in FIJI. Detailed description can be

found in (Han et al. 2024).

Automatic segmentation

We segmented a central cutout of the dataset using flood-filling networks (Michał Januszewski et

al. 2018). The FFN segmentation model was trained at 16 x 16 x 33 nm nominal resolution on the H01

dataset (Shapson-Coe et al. 2024), and run here on CLAHE intensity normalized data (Zuiderveld 1994)

downsampled to 16 x 16 x 30 nm. Direct application of the H01-trained FFN to the AD dataset did not

perform well, particularly on thinner objects. Therefore we used Segmentation Enhanced CycleGANs

(Michal Januszewski and Jain 2019) to modify the visual appearance (photometrics, noise statistics,

staining, etc.) of the AD dataset at 16 x 16 x 30 nm to mimic H01, prior to running the FFN. We trained

the SECGAN style transfer on a central cutout of the AD dataset totaling 240 x 160 x 15 um and a cutout

of H01 totaling 1600 x 760 x 130 um for 8 million steps at batch size 1. The SECGAN generator

architecture was an 8-layer fully convolutional stack with residual connections and no downsampling, and

the discriminator was a ResNet18 (He et al., n.d.).

2D membrane detection at 4 x 4 x 30 nm resolution was performed on the vEM dataset using a

method developed in our lab. A description of this approach was given in (Karlupia et al. 2023; Meirovitch

et al. 2018; Pavarino et al. 2023). In brief, an algorithm (Pavarino et al. 2023) pre-trained on a mouse

cerebellar dataset was used to generate 2D membrane detection. 2D membrane detection was use for

filtering false positive in synapse prediction.

Synapse detection by supervised leaning

We sampled nine EM image subvolumes with dimensions of 6 μm x 6 μm x 6 μm at three

distances (10 μm, 20 μm, 30 μm) from the center of an Aβ plaque at the original resolution. We took a

semi-automatic approach to annotate all the synapses in these subvolumes. First, one human annotator

labeled all the pre- and post-synaptic appositions in 1/4 of one of the nine EM image subvolumes. Then,

we trained a synapse detection model (Parag et al. 2019) using the manual annotations with the PyTorch

Connectomics deep learning package. To refine the model, we tested it on all slices of this subvolume,

manually proofread its results, and fine-tuned the model with all the annotations, which led to a desirable

detection accuracy and recall result. Lastly, we used the improved model to detect synapses in the entire

dataset and identified 3.5 million synapses..

Synapse detection by transfer learning

The original synapse annotation model was a UNet (3-down and 3-up modules) trained at 8 x 8 x 33 nm nominal resolution on the H01 dataset (Shapson-Coe et al. 2024). For the initial synapse predictions, we used the SECGAN model trained for 683k steps to style-transfer an ROI of the AD data to mimic H01, and then ran a synapse model that was trained for 786k steps on H01. This produced 0.255 million synapse masks, with low rates of precision and recall. We then used the model-inferred synapse masks as automated ground truth to train a new UNet from scratch directly on the AD data (8 x 8 x 30 nm resolution, without SECGAN style transfer, but with CLAHE). We trained this model for 38M steps and biased it against false negatives by weighting positively labeled pre- and post-synaptic voxels in the training data by 2x. The new synapse annotation model produced 2.1 million raw synapse masks, which resulted in 0.732 million total synapses after filtering for synapses above a 100 voxel volume threshold, removing any pairs that had identical pre- and post-synaptic parents (i.e. self-synapses) and masking out synaptic sites residing on blood vessels, somas, or myelin. We estimated the final precision and recall as 46% and 52% based on evaluation against the manualy annotated subvolume in the first method of synapse detection.

Automatic Aβ plaque segmentation

We optimized a DeepLab-v3 semantic segmentation model (L.-C. Chen et al. 2018) on a cropped stack of images with manually identified Aβ plaque. The model was pre-trained on another vEM dataset from the Alzheimer's disease mouse model Tg2576 (Dr. Olga Morozova, personal communication) to decrease the requirements of manually annotated training data in the target volume. We also oversample patches near the Aβ plaque border during training to improve segmentation quality as the mask structures in those patches are more challenging. The training and inference are implemented with the open-source PyTorch Connectomics codebase.

Statistical analysis

Two-tailed, unpaired t-tests on synaptic density, synaptic volume and total synaptic volume of the three categories defined in the section of synapse detection (The PyTorch method) were performed in Prism-GraphPad. Two-sample Kolmogorov–Smirnov test was performed using Python.

Acknowledgement

We thank the Harvard Center for Biological Imaging (RRID:SCR_018673) for infrastructure and support, the Biopolymers and Proteomics Core Facility at the Koch Institute at MIT for processing the peptide-fluorescent dye conjugates, the Bauer Core Facility at Harvard for infrastructure and support, Dr. Olga Morozova for helpful discussion on the Alzheimer's disease mouse model and immunolabeling of Aβ plaques, and Dr. Marta Montero-Crespo for helpful discussion on synapse annotation. This work was supported by NIH grants U19 NS104653, UG3 MH123386, and P50 MH094271 (J. W. L.); NSF grants NSF-CAREER-2239688 (D. W.), NCS-FO-2124179, and NCS-FO-1835231 (H. Pfister.), NLM T15LM007092-31 (M. S.); Office of Naval Research grant N00014-20-1-2828 (J. W. L). X. H. was supported by the Edward R. and Anne G. Lefler predoctoral fellowship from the Lefler Center for Neurodegenerative Disorders at Harvard Medical School and the Simmons Awards from the Harvard Center for Biological Imaging.

Author Contribution

X. H. and J. W. L conceived this study. Nanobody A2 and R3VQ were originally generated in the labs of P. L., S. B., and B. D.. X. H. generated the fluorescent nanobody probes A2 and R3VQ. H. Ploegh provided the fluorescent anti-CD11b, anti-GFAP, and anti-Ly6C/6G nanobodies. X. H. performed the LM experiments. X. H. and R. S. performed the EM experiments. S. W. performed the imaging processing of the vEM dataset. P. H. L. and V. J. performed 3D segmentation and curated the vCLEM dataset at Neuroglancer. X. H., M. S., and F. A. performed 3D reconstruction of neurons. S. A., B. G., D. W., and X. H. performed synapse prediction through supervised learning.T. B. performed synapse detection through transfer learning. Y. M. (yaron.mr@gmail.com) performed 2D membrane detection. M. S. performed the filtering of the synapse detection results using a U-Net classifier. Z. L., and H. Pfister performed automatic plaque detection. D. B. provided the reconstruction tool VAST and advice on 3D reconstruction and rendering. Y. W. helped with LM and EM co-registration. X. H. and J. W. L wrote the paper with inputs from H. Ploegh, B. D., D. B., and R. S.

Competing interests

The authors declare no competing interests.

References

"2024 Alzheimer's Disease Facts and Figures." 2024. *Alzheimer's & Dementia: The Journal of the Alzheimer's Association* 20 (5): 3708–3821.

Androuin, Alexandre, Brigitte Potier, U. Valentin Nägerl, Daniel Cattaert, Lydia Danglot, Manon Thierry, Ihsen Youssef, et al. 2018. "Evidence for Altered Dendritic Spine Compartmentalization in Alzheimer's Disease and Functional Effects in a Mouse Model." *Acta Neuropathologica* 135 (6): 839–54.

Arendt, Thomas, Jens T. Stieler, and Max Holzer. 2016. "Tau and Tauopathies." *Brain Research Bulletin* 126 (Pt 3): 238–92.

Barbone, Giacomo E., Alberto Bravin, Alberto Mittone, Alexandra Pacureanu, Giada Mascio, Paola Di Pietro, Markus J. Kraiger, et al. 2022. "X-Ray Multiscale 3D Neuroimaging to Quantify Cellular Aging and Neurodegeneration Postmortem in a Model of Alzheimer's Disease." *European Journal of Nuclear Medicine and Molecular Imaging* 49 (13): 4338–57.

Barucker, Christian, Anja Harmeier, Joerg Weiske, Beatrix Fauler, Kai Frederik Albring, Stefan Prokop, Peter Hildebrand, et al. 2014. "Nuclear Translocation Uncovers the Amyloid Peptide Aβ42 as a Regulator of Gene Transcription." *The Journal of Biological Chemistry* 289 (29): 20182–91.

Bates, Gillian P., Ray Dorsey, James F. Gusella, Michael R. Hayden, Chris Kay, Blair R. Leavitt, Martha Nance, et al. 2015. "Huntington Disease." *Nature Reviews. Disease Primers* 1 (April):15005.

Belfiore, Ramona, Alexis Rodin, Eric Ferreira, Ramon Velazquez, Caterina Branca, Antonella Caccamo, and Salvatore Oddo. 2019. "Temporal and Regional Progression of Alzheimer's Disease-like Pathology in 3xTg-AD Mice." *Aging Cell* 18 (1): e12873.

Bisht, Kanchan, Kaushik P. Sharma, Cynthia Lecours, Maria Gabriela Sánchez, Hassan El Hajj, Giampaolo Milior, Adrián Olmos-Alonso, et al. 2016. "Dark Microglia: A New Phenotype Predominantly Associated with Pathological States." *Glia* 64 (5): 826–39.

Blazquez-Llorca, Lidia, Virginia Garcia-Marin, Paula Merino-Serrais, Jesús Ávila, and Javier DeFelipe. 2011. "Abnormal Tau Phosphorylation in the Thorny Excrescences of CA3 Hippocampal Neurons in Patients with Alzheimer's Disease." *Journal of Alzheimer's Disease: JAD* 26 (4): 683–98.

Blazquez-Llorca, Lidia, Ángel Merchán-Pérez, José-Rodrigo Rodríguez, Jorge Gascón, and Javier DeFelipe. 2013. "FIB/SEM Technology and Alzheimer's Disease: Three-Dimensional Analysis of Human Cortical Synapses." *Journal of Alzheimer's Disease: JAD* 34 (4): 995–1013.

Bogovic, John A., Philipp Hanslovsky, Allan Wong, and Stephan Saalfeld. 2016. "Robust Registration of Calcium Images by Learned Contrast Synthesis." In *2016 IEEE 13th International Symposium on Biomedical Imaging (ISBI)*, 1123–26.

Buhmann, Julia, Arlo Sheridan, Caroline Malin-Mayor, Philipp Schlegel, Stephan Gerhard, Tom

Kazimiers, Renate Krause, et al. 2021. "Automatic Detection of Synaptic Partners in a
Whole-Brain Drosophila Electron Microscopy Data Set." *Nature Methods* 18 (7): 771–74.
Chen, Guo-Fang, Ting-Hai Xu, Yan Yan, Yu-Ren Zhou, Yi Jiang, Karsten Melcher, and H. Eric
Xu. 2017. "Amyloid Beta: Structure, Biology and Structure-Based Therapeutic
Development." *Acta Pharmacologica Sinica* 38 (9): 1205–35.
Chen, Liang-Chieh, Yukun Zhu, George Papandreou, Florian Schroff, and Hartwig Adam. 2018.
"Encoder-Decoder with Atrous Separable Convolution for Semantic Image Segmentation."
In *Computer Vision – ECCV 2018*, 833–51. Lecture Notes in Computer Science. Cham:
Springer International Publishing.
Chung, Won-Suk, Christina A. Welsh, Ben A. Barres, and Beth Stevens. 2015. "Do Glia Drive
Synaptic and Cognitive Impairment in Disease?" *Nature Neuroscience* 18 (11): 1539–45.
Çiçek, Özgün, Ahmed Abdulkadir, Soeren S. Lienkamp, Thomas Brox, and Olaf Ronneberger.
2016. "3D U-Net: Learning Dense Volumetric Segmentation from Sparse Annotation." In
Medical Image Computing and Computer-Assisted Intervention – MICCAI 2016, 424–32.
Springer International Publishing.
Creators Jeremy Maitin-Shepard Alex Baden1 William Silversmith2 Eric Perlman3 Forrest
Collman4 Tim Blakely5 Jan Funke Chris Jordan Ben Falk6 Nico Kemnitz7 tingzhao Chris
Roat Manuel Castro Sridhar Jagannathan8 moenigin Jody Clements Austin Hoag9 Bill
Katz10 Dave Parsons Jingpeng Wu11 Lee Kamentsky Pavel Chervakov Philip Hubbard12
Stuart Berg John Hoffer13 Akhilesh Halageri Christian Machacek Kevin Mader Lutz
Roeder14 Peter H. Li Show affiliations 1. @omnisci 2. Seung Lab @ Princeton University 3.
Yikes LLC 4. Allen Institute for Brain Science 5. @google 6. @stsci 7. @seung-lab
@zettaai 8. Charité Universitätsmedizin Berlin 9. Princeton University 10. Janelia Research
Center 11. @flatironinstitute 12. Howard Hughes Medical Institute, Janelia Research
Campus 13. Harvard 14. @microsoft. n.d. *Google/neuroglancer:*
https://doi.org/10.5281/zenodo.5573294.
Crowther, R. A. 1991. "Straight and Paired Helical Filaments in Alzheimer Disease Have a
Common Structural Unit." *Proceedings of the National Academy of Sciences of the United
States of America* 88 (6): 2288–92.
Cummings, Jeffrey, Yadi Zhou, Garam Lee, Kate Zhong, Jorge Fonseca, and Feixiong Cheng.
2024. "e12465 10.1002/trc2.12465 New therapies to prevent or delay the onset of
symptoms, slow progression, or improve cognitive and behavioral symptoms of Alzheimer's
disease (AD) are needed. We interrogated clinicaltrials.gov including all clinical trials
assessing pharmaceutical therapies for AD active in on January 1, 2024. We used the
Common Alzheimer's Disease Research Ontology (CADRO) to classify the targets of
therapies in the pipeline. There are 164 trials assessing 127 drugs across the 2024 AD
pipeline. There were 48 trials in Phase 3 testing 32 drugs, 90 trials in Phase 2 assessing 81
drugs, and 26 trials in Phase 1 testing 25 agents. Of the 164 trials, 34% (N = 56) assess
disease-modifying biological agents, 41% (N = 68) test disease-modifying small molecule
drugs, 10% (N = 17) evaluate cognitive enhancing agents, and 14% (N = 23) test drugs for
the treatment of neuropsychiatric symptoms. Compared to the 2023 pipeline, there are
fewer trials (164 vs. 187), fewer drugs (127 vs. 141), fewer new chemical entities (88 vs.
101), and a similar number of repurposed agents (39 vs. 40). In the 2024 Alzheimer's
disease drug development pipeline, there are 164 clinical trials assessing 127 drugs.The
2024 Alzheimer's disease drug development pipeline has contracted compared to the 2023
Alzheimer pipeline with fewer trials, fewer drugs, and fewer new chemical entities.Drugs in
the Alzheimer's disease drug development pipeline target a wide array of targets; the most
common processes targeted include neurotransmitter receptors, inflammation, amyloid, and
synaptic plasticity.The total development time for a potential Alzheimer's disease therapy to
progress from nonclinical studies to FDA review is approximately 13 years. © 2024 The

Authors. Alzheimer's & Dementia: Translational Research & Clinical Interventions published by Wiley Periodicals LLC on behalf of Alzheimer's Association. Cummings Jeffrey J Chambers-Grundy Center for Transformative Neuroscience Department of Brain Health School of Integrated Health Sciences University of Nevada, Las Vegas (UNLV) Las Vegas Nevada USA. Zhou Yadi Y Genomic Medicine Institute Lerner Research Institute, Cleveland Clinic Cleveland Ohio USA. Lee Garam G Chambers-Grundy Center for Transformative Neuroscience Department of Brain Health School of Integrated Health Sciences University of Nevada, Las Vegas (UNLV) Las Vegas Nevada USA. Zhong Kate K Chambers-Grundy Center for Transformative Neuroscience Department of Brain Health School of Integrated Health Sciences University of Nevada, Las Vegas (UNLV) Las Vegas Nevada USA. Fonseca Jorge J Howard R Hughes College of Engineering Department of Computer Science University of Nevada, Las Vegas (UNLV) Las Vegas Nevada USA. Cheng Feixiong F Genomic Medicine Institute Lerner Research Institute, Cleveland Clinic Cleveland Ohio USA. Department of Molecular Medicine Cleveland Clinic Lerner College of Medicine Case Western Reserve University Cleveland Ohio USA. Case Comprehensive Cancer Center Case Western Reserve University School of Medicine Cleveland Ohio USA. Cleveland Clinic Genome Center Lerner Research Institute, Cleveland Clinic Cleveland Ohio USA. dut R21 AG083003 AG NIA NIH HHS United States R01 AG082118 AG NIA NIH HHS United States P20 AG068053 AG NIA NIH HHS United States R01 AG084250 AG NIA NIH HHS United States RF1 NS133812 NS NINDS NIH HHS United States U01 AG073323 AG NIA NIH HHS United States R01 AG066707 AG NIA NIH HHS United States R01 AG076448 AG NIA NIH HHS United States P20 GM109025 GM NIGMS NIH HHS United States R35 AG071476 AG NIA NIH HHS United States RF1 AG082211 AG NIA NIH HHS United States English Abstract Journal Article Review Alzheimer's disease drug development pipeline: 2024. 2024 04 24." *Alzheimer's & dementia (New York, N. Y.)* 10 (2): e12465.

D'Andrea, Laura, Ramona Stringhi, Monica Di Luca, and Elena Marcello. 2021. "Looking at Alzheimer's Disease Pathogenesis from the Nuclear Side." *Biomolecules* 11 (9): 1261.

Danis, Clément, Elian Dupré, Orgeta Zejneli, Raphaëlle Caillierez, Alexis Arrial, Séverine Bégard, Justine Mortelecque, et al. 2022. "Inhibition of Tau Seeding by Targeting Tau Nucleation Core within Neurons with a Single Domain Antibody Fragment." *Molecular Therapy: The Journal of the American Society of Gene Therapy* 30 (4): 1484–99.

Davies, C. A., D. M. Mann, P. Q. Sumpter, and P. O. Yates. 1987. "A Quantitative Morphometric Analysis of the Neuronal and Synaptic Content of the Frontal and Temporal Cortex in Patients with Alzheimer's Disease." *Journal of the Neurological Sciences* 78 (2): 151–64.

De Genst, Erwin J., Tim Guilliams, Joke Wellens, Elizabeth M. O'Day, Christopher A. Waudby, Sarah Meehan, Mireille Dumoulin, et al. 2010. "Structure and Properties of a Complex of α-Synuclein and a Single-Domain Camelid Antibody." *Journal of Molecular Biology* 402 (2): 326–43.

DeKosky, S. T., and S. W. Scheff. 1990. "Synapse Loss in Frontal Cortex Biopsies in Alzheimer's Disease: Correlation with Cognitive Severity." *Annals of Neurology* 27 (5): 457–64.

Deszyński, Piotr, Jakub Młokosiewicz, Adam Volanakis, Igor Jaszczyszyn, Natalie Castellana, Stefano Bonissone, Rajkumar Ganesan, and Konrad Krawczyk. 2021. "INDI—integrated Nanobody Database for Immunoinformatics." *Nucleic Acids Research* 50 (D1): D1273–81.

Domínguez-Álvaro, M., M. Montero-Crespo, L. Blazquez-Llorca, R. Insausti, J. DeFelipe, and L. Alonso-Nanclares. 2018. "Three-Dimensional Analysis of Synapses in the Transentorhinal Cortex of Alzheimer's Disease Patients." *Acta Neuropathologica Communications* 6 (1). https://doi.org/10.1186/s40478-018-0520-6.

Fang, Michael, Jiaqi Hu, Jordan Weiss, David S. Knopman, Marilyn Albert, B. Gwen Windham,

Keenan A. Walker, et al. 2025. "Lifetime Risk and Projected Burden of Dementia." *Nature Medicine*, January, 1–5.

Fang, Tao, Xiaotang Lu, Daniel Berger, Christina Gmeiner, Julia Cho, Richard Schalek, Hidde Ploegh, and Jeff Lichtman. 2018. "Nanobody Immunostaining for Correlated Light and Electron Microscopy with Preservation of Ultrastructure." *Nature Methods* 15 (12): 1029–32.

Ferrer, Ismael, Chadni Sanyal, Marie-Jo Moutin, and Damaris N. Lorenzo. 2024. "Putting the Brakes on Axonal Branching." *Trends in Neurosciences*, May. https://doi.org/10.1016/j.tins.2024.05.001.

Flury, Anna, Leen Aljayousi, Hye-Jin Park, Mohammadparsa Khakpour, Jack Mechler, Siaresh Aziz, Jackson D. McGrath, et al. 2024. "A Neurodegenerative Cellular Stress Response Linked to Dark Microglia and Toxic Lipid Secretion." *Neuron* 0 (0). https://doi.org/10.1016/j.neuron.2024.11.018.

Gouras, Gunnar K., Davide Tampellini, Reisuke H. Takahashi, and Estibaliz Capetillo-Zarate. 2010. "Intraneuronal Beta-Amyloid Accumulation and Synapse Pathology in Alzheimer's Disease." *Acta Neuropathologica* 119 (5): 523–41.

Gowrishankar, Swetha, Peng Yuan, Yumei Wu, Matthew Schrag, Summer Paradise, Jaime Grutzendler, Pietro De Camilli, and Shawn M. Ferguson. 2015. "Massive Accumulation of Luminal Protease-Deficient Axonal Lysosomes at Alzheimer's Disease Amyloid Plaques." *Proceedings of the National Academy of Sciences of the United States of America* 112 (28): E3699–3708.

Habicht, Gernot, Christian Haupt, Ralf P. Friedrich, Peter Hortschansky, Carsten Sachse, Jessica Meinhardt, Karin Wieligmann, et al. 2007. "Directed Selection of a Conformational Antibody Domain That Prevents Mature Amyloid Fibril Formation by Stabilizing Abeta Protofibrils." *Proceedings of the National Academy of Sciences of the United States of America* 104 (49): 19232–37.

Hampel, Harald, John Hardy, Kaj Blennow, Christopher Chen, George Perry, Seung Hyun Kim, Victor L. Villemagne, et al. 2021. "The Amyloid-β Pathway in Alzheimer's Disease." *Molecular Psychiatry* 26 (10): 5481–5503.

Han, Xiaomeng, Xiaotang Lu, Peter H. Li, Shuohong Wang, Richard Schalek, Yaron Meirovitch, Zudi Lin, et al. 2024. "Multiplexed Volumetric CLEM Enabled by scFvs Provides Insights into the Cytology of Cerebellar Cortex." *Nature Communications* 15 (1): 6648.

Hayworth, K. J., N. Kasthuri, R. Schalek, and J. W. Lichtman. 2006. "Automating the Collection of Ultrathin Serial Sections for Large Volume TEM Reconstructions." *Microscopy and Microanalysis: The Official Journal of Microscopy Society of America, Microbeam Analysis Society, Microscopical Society of Canada* 12 (S02): 86–87.

He, K., X. Zhang, S. Ren, and J. Sun. n.d. "Deep Residual Learning for Image Recognition." http://openaccess.thecvf.com/content_cvpr_2016/html/He_Deep_Residual_Learning_CVPR_2016_paper.html.

Hoffman, David P., Gleb Shtengel, C. Shan Xu, Kirby R. Campbell, Melanie Freeman, Lei Wang, Daniel E. Milkie, et al. n.d. "Correlative Three-Dimensional Super-Resolution and Block Face Electron Microscopy of Whole Vitreously Frozen Cells." https://doi.org/10.1101/773986.

Hooli, B. V., C. M. Lill, K. Mullin, D. Qiao, C. Lange, and L. Bertram. 2015. "PLD3 Gene Variants and Alzheimer's Disease." *Nature*. https://www.nature.com/articles/nature14040.

Im, Kyuseok, Sergey Mareninov, M. Fernando Palma Diaz, and William H. Yong. 2019. "An Introduction to Performing Immunofluorescence Staining." *Methods in Molecular Biology* 1897:299–311.

Januszewski, Michal, and Viren Jain. 2019. "Segmentation-Enhanced CycleGAN." *bioRxiv*. bioRxiv. https://doi.org/10.1101/548081.

Januszewski, Michał, Jörgen Kornfeld, Peter H. Li, Art Pope, Tim Blakely, Larry Lindsey,

Jeremy Maitin-Shepard, Mike Tyka, Winfried Denk, and Viren Jain. 2018. "High-Precision Automated Reconstruction of Neurons with Flood-Filling Networks." *Nature Methods* 15 (8): 605–10.

Jiang, Yi, Linlin Li, Keliang Pang, Jiazheng Liu, Bohao Chen, Jingbin Yuan, Lijun Shen, Xi Chen, Bai Lu, and Hua Han. 2022. "Synaptic Degeneration in the Prefrontal Cortex of a Rat AD Model Revealed by Volume Electron Microscopy." *Journal of Molecular Cell Biology* 14 (3). https://doi.org/10.1093/jmcb/mjac012.

John, Albin, and P. Hemachandra Reddy. 2021. "Synaptic Basis of Alzheimer's Disease: Focus on Synaptic Amyloid Beta, P-Tau and Mitochondria." *Ageing Research Reviews* 65 (101208): 101208.

Johnson, Erik C. B., E. Kathleen Carter, Eric B. Dammer, Duc M. Duong, Ekaterina S. Gerasimov, Yue Liu, Jiaqi Liu, et al. 2022. "Large-Scale Deep Multi-Layer Analysis of Alzheimer's Disease Brain Reveals Strong Proteomic Disease-Related Changes Not Observed at the RNA Level." *Nature Neuroscience* 25 (2): 213–25.

Johnson, Erik C. B., Eric B. Dammer, Duc M. Duong, Lingyan Ping, Maotian Zhou, Luming Yin, Lenora A. Higginbotham, et al. 2020. "Large-Scale Proteomic Analysis of Alzheimer's Disease Brain and Cerebrospinal Fluid Reveals Early Changes in Energy Metabolism Associated with Microglia and Astrocyte Activation." *Nature Medicine* 26 (5): 769–80.

Karlupia, Neha, Richard L. Schalek, Yuelong Wu, Yaron Meirovitch, Donglai Wei, Alexander W. Charney, Brian H. Kopell, and Jeff W. Lichtman. 2023. "Immersion Fixation and Staining of Multi-Cubic Millimeter Volumes for Electron Microscopy-Based Connectomics of Human Brain Biopsies." *Biological Psychiatry*, February. https://doi.org/10.1016/j.biopsych.2023.01.025.

Kasthuri, Narayanan, Kenneth Jeffrey Hayworth, Daniel Raimund Berger, Richard Lee Schalek, José Angel Conchello, Seymour Knowles-Barley, Dongil Lee, et al. 2015. "Saturated Reconstruction of a Volume of Neocortex." *Cell* 162 (3): 648–61.

Kelly, Louise, Mohsen Seifi, Ruolin Ma, Scott J. Mitchell, Uwe Rudolph, Kirsten L. Viola, William L. Klein, Jeremy J. Lambert, and Jerome D. Swinny. 2021. "Identification of Intraneuronal Amyloid Beta Oligomers in Locus Coeruleus Neurons of Alzheimer's Patients and Their Potential Impact on Inhibitory Neurotransmitter Receptors and Neuronal Excitability." *Neuropathology and Applied Neurobiology* 47 (4): 488–505.

Kislinger, Georg, Gunar Fabig, Antonia Wehn, Lucia Rodriguez, Hanyi Jiang, Cornelia Niemann, Andrey S. Klymchenko, et al. 2023. "ATUM-Tomo: A Multi-Scale Approach to Cellular Ultrastructure by Combined Volume Scanning Electron Microscopy and Electron Tomography." *eLife*. https://doi.org/10.7554/elife.90565.1.

Kislinger, Georg, Cornelia Niemann, Lucia Rodriguez, Hanyi Jiang, Maryam K. Fard, Nicolas Snaidero, Adrian-Minh Schumacher, Martin Kerschensteiner, Thomas Misgeld, and Martina Schifferer. 2023. "Neurons on Tape: Automated Tape Collecting Ultramicrotomy-Mediated Volume EM for Targeting Neuropathology." Edited by Kedar Narayan, Lucy Collinson, and Paul Verkade. *Methods in Cell Biology* 177 (February):125–70.

Knott, G., H. Marchman, D. Wall, and B. Lich. 2008. "Serial Section Scanning Electron Microscopy of Adult Brain Tissue Using Focused Ion Beam Milling." *Journal of Neuroscience*. https://doi.org/10.1523/jneurosci.3189-07.2008.

Knupp, Allison, Swati Mishra, Refugio Martinez, Jacquelyn E. Braggin, Marcell Szabo, Chizuru Kinoshita, Dale W. Hailey, Scott A. Small, Suman Jayadev, and Jessica E. Young. 2020. "Depletion of the AD Risk Gene SORL1 Selectively Impairs Neuronal Endosomal Traffic Independent of Amyloidogenic APP Processing." *Cell Reports* 31 (9): 107719.

Koffie, Robert M., Bradley T. Hyman, and Tara L. Spires-Jones. 2011. "Alzheimer's Disease: Synapses Gone Cold." *Molecular Neurodegeneration* 6 (1): 63.

Koffie, Robert M., Melanie Meyer-Luehmann, Tadafumi Hashimoto, Kenneth W. Adams,

Matthew L. Mielke, Monica Garcia-Alloza, Kristina D. Micheva, et al. 2009. "Oligomeric Amyloid β Associates with Postsynaptic Densities and Correlates with Excitatory Synapse Loss near Senile Plaques." *Proceedings of the National Academy of Sciences* 106 (10): 4012–17.

Kuan, Aaron T., Giulio Bondanelli, Laura N. Driscoll, Julie Han, Minsu Kim, David G. C. Hildebrand, Brett J. Graham, et al. 2024. "Synaptic Wiring Motifs in Posterior Parietal Cortex Support Decision-Making." *Nature* 627 (8003): 367–73.

Lafaye, Pierre, Ikbel Achour, Patrick England, Charles Duyckaerts, and François Rougeon. 2009. "Single-Domain Antibodies Recognize Selectively Small Oligomeric Forms of Amyloid β, Prevent Aβ-Induced Neurotoxicity and Inhibit Fibril Formation." *Molecular Immunology* 46 (4): 695–704.

Lafaye, Pierre, Sylvie Bay, Benoît Delatour, Marc Dhenain, Charles Duyckaerts, Tengfei Li, Matthias Vandesquille, Christian Czech, and Fiona Grueninger. 2019. Camelid single-domain antibody directed against amyloid bêta and methods for producing conjugates thereof. EPO 3068799:B1. *European Patent*, filed November 13, 2014, and issued August 28, 2019. https://patentimages.storage.googleapis.com/06/2d/dd/30cf8941c20327/EP3068799B1.pdf.

Lafaye, Pierre, Sylvie Bay, Benoit Delatour, Marc Dhenain, Charles Duyckaerts, Tengfei Li, Matthias Vandesquille, Christian Czech, and Fiona Grueninger. 2020. Camelid single-domain antibody directed against phosphorylated tau proteins and methods for producing conjugates thereof. USPTO 20200165329:A1. *US Patent*, filed December 2, 2019, and issued May 28, 2020. https://patentimages.storage.googleapis.com/88/2f/7a/dd7804c87c4ce3/US20200165329A1.pdf.

LaFerla, Frank M., Kim N. Green, and Salvatore Oddo. 2007. "Intracellular Amyloid-Beta in Alzheimer's Disease." *Nature Reviews. Neuroscience* 8 (7): 499–509.

Lee, Wei-Chung Allen, Vincent Bonin, Michael Reed, Brett J. Graham, Greg Hood, Katie Glattfelder, and R. Clay Reid. 2016. "Anatomy and Function of an Excitatory Network in the Visual Cortex." *Nature* 532 (7599): 370–74.

Lin, Zudi, Donglai Wei, Jeff Lichtman, and Hanspeter Pfister. 2021. "PyTorch Connectomics: A Scalable and Flexible Segmentation Framework for EM Connectomics." *arXiv [eess.IV]*. arXiv. http://arxiv.org/abs/2112.05754.

Li, Tengfei, Matthias Vandesquille, Fani Koukouli, Clémence Dudeffant, Ihsen Youssef, Pascal Lenormand, Christelle Ganneau, et al. 2016. "Camelid Single-Domain Antibodies: A Versatile Tool for in Vivo Imaging of Extracellular and Intracellular Brain Targets." *Journal of Controlled Release: Official Journal of the Controlled Release Society* 243 (December):1–10.

Li, Xiaoyu, Yatender Kumar, Hans Zempel, Eva-Maria Mandelkow, Jacek Biernat, and Eckhard Mandelkow. 2011. "Novel Diffusion Barrier for Axonal Retention of Tau in Neurons and Its Failure in Neurodegeneration." *The EMBO Journal* 30 (23): 4825–37.

Loomba, Sahil, Jakob Straehle, Vijayan Gangadharan, Natalie Heike, Abdelrahman Khalifa, Alessandro Motta, Niansheng Ju, et al. 2022. "Connectomic Comparison of Mouse and Human Cortex." *Science* 377 (6602): eabo0924.

Magistri, Marco, Dmitry Velmeshev, Madina Makhmutova, and Mohammad Ali Faghihi. 2015. "Transcriptomics Profiling of Alzheimer's Disease Reveal Neurovascular Defects, Altered Amyloid-β Homeostasis, and Deregulated Expression of Long Noncoding RNAs." *Journal of Alzheimer's Disease: JAD* 48 (3): 647–65.

Mandelkow, E. M., and E. Mandelkow. 1998. "Tau in Alzheimer's Disease." *Trends in Cell Biology* 8 (11): 425–27.

Martin, Elodie, Mohamed El-Behi, Bertrand Fontaine, and Cecile Delarasse. 2017. "Analysis of

Microglia and Monocyte-Derived Macrophages from the Central Nervous System by Flow Cytometry." *Journal of Visualized Experiments*, no. 124 (June). https://doi.org/10.3791/55781.

Martínez-Carmona, Marina, Fernando Lucas-Ruiz, Alejandro Gallego-Ortega, Caridad Galindo-Romero, María Norte-Muñoz, María José González-Riquelme, Francisco J. Valiente-Soriano, Manuel Vidal-Sanz, and Marta Agudo-Barriuso. 2021. "Ly6c as a New Marker of Mouse Blood Vessels: Qualitative and Quantitative Analyses on Intact and Ischemic Retinas." *International Journal of Molecular Sciences* 23 (1): 19.

Martínez-Serra, Raquel, Lidia Alonso-Nanclares, Kwangwook Cho, and K. Peter Giese. 2022. "Emerging Insights into Synapse Dysregulation in Alzheimer's Disease." *Brain Communications* 4 (2): fcac083.

Mathys, Hansruedi, Jose Davila-Velderrain, Zhuyu Peng, Fan Gao, Shahin Mohammadi, Jennie Z. Young, Madhvi Menon, et al. 2019. "Single-Cell Transcriptomic Analysis of Alzheimer's Disease." *Nature* 570 (7761): 332–37.

McMahon, Conor, Alexander S. Baier, Roberta Pascolutti, Marcin Wegrecki, Sanduo Zheng, Janice X. Ong, Sarah C. Erlandson, et al. 2018. "Yeast Surface Display Platform for Rapid Discovery of Conformationally Selective Nanobodies." *Nature Structural & Molecular Biology* 25 (3): 289–96.

Meftah, Soraya, and Jian Gan. 2023. "Alzheimer's Disease as a Synaptopathy: Evidence for Dysfunction of Synapses during Disease Progression." *Frontiers in Synaptic Neuroscience* 15 (March):1129036.

Meirovitch, Yaron, Lu Mi, Hayk Saribekyan, Alexander Matveev, David Rolnick, and Nir Shavit. 2018. "Cross-Classification Clustering: An Efficient Multi-Object Tracking Technique for 3-D Instance Segmentation in Connectomics." *arXiv [cs.CV]*. arXiv. http://openaccess.thecvf.com/content_CVPR_2019/html/Meirovitch_Cross-Classification_Clustering_An_Efficient_Multi-Object_Tracking_Technique_for_3-D_Instance_CVPR_2019_paper.html.

MICrONS Consortium, J. Alexander Bae, Mahaly Baptiste, Agnes L. Bodor, Derrick Brittain, Joann Buchanan, Daniel J. Bumbarger, et al. 2021. "Functional Connectomics Spanning Multiple Areas of Mouse Visual Cortex." *bioRxiv*. https://doi.org/10.1101/2021.07.28.454025.

Mildner, Alexander, Hauke Schmidt, Mirko Nitsche, Doron Merkler, Uwe-Karsten Hanisch, Matthias Mack, Mathias Heikenwalder, Wolfgang Brück, Josef Priller, and Marco Prinz. 2007. "Microglia in the Adult Brain Arise from Ly-6ChiCCR2+ Monocytes Only under Defined Host Conditions." *Nature Neuroscience* 10 (12): 1544–53.

Miyoshi, Emily, Tina Bilousova, Mikhail Melnik, Danyl Fakhrutdinov, Wayne W. Poon, Harry V. Vinters, Carol A. Miller, et al. 2021. "Exosomal Tau with Seeding Activity Is Released from Alzheimer's Disease Synapses, and Seeding Potential Is Associated with Amyloid Beta." *Laboratory Investigation; a Journal of Technical Methods and Pathology* 101 (12): 1605–17.

Montero-Crespo, Marta, Marta Domínguez-Álvaro, Lidia Alonso-Nanclares, Javier DeFelipe, and Lidia Blazquez-Llorca. 2021. "Three-Dimensional Analysis of Synaptic Organization in the Hippocampal CA1 Field in Alzheimer's Disease." *Brain: A Journal of Neurology* 144 (2): 553–73.

Montero-Crespo, Marta, Marta Dominguez-Alvaro, Patricia Rondon-Carrillo, Lidia Alonso-Nanclares, Javier DeFelipe, and Lidia Blazquez-Llorca. 2020. "Three-Dimensional Synaptic Organization of the Human Hippocampal CA1 Field." *eLife* 9 (July):e57013.

Motta, Alessandro, Manuel Berning, Kevin M. Boergens, Benedikt Staffler, Marcel Beining, Sahil Loomba, Philipp Hennig, Heiko Wissler, and Moritz Helmstaedter. 2019. "Dense Connectomic Reconstruction in Layer 4 of the Somatosensory Cortex." *Science* 366 (6469). https://doi.org/10.1126/science.aay3134.

Muyldermans, Serge. 2021. "A Guide to: Generation and Design of Nanobodies." *The FEBS Journal* 288 (7): 2084–2102.

Nackenoff, Alex G., Timothy J. Hohman, Sarah M. Neuner, Carolyn S. Akers, Nicole C. Weitzel, Alena Shostak, Shawn M. Ferguson, et al. 2021. "PLD3 Is a Neuronal Lysosomal Phospholipase D Associated with β-Amyloid Plaques and Cognitive Function in Alzheimer's Disease." *PLoS Genetics* 17 (4): e1009406.

Neuman, Krystina M., Elizabeth Molina-Campos, Timothy F. Musial, Andrea L. Price, Kwang-Jin Oh, Malerie L. Wolke, Eric W. Buss, Stephen W. Scheff, Elliott J. Mufson, and Daniel A. Nicholson. 2015. "Evidence for Alzheimer's Disease-Linked Synapse Loss and Compensation in Mouse and Human Hippocampal CA1 Pyramidal Neurons." *Brain Structure & Function* 220 (6): 3143–65.

Nguyen, Tri M., Logan A. Thomas, Jeff L. Rhoades, Ilaria Ricchi, Xintong Cindy Yuan, Arlo Sheridan, David G. C. Hildebrand, Jan Funke, Wade G. Regehr, and Wei-Chung Allen Lee. 2022. "Structured Cerebellar Connectivity Supports Resilient Pattern Separation." *Nature*, November. https://doi.org/10.1038/s41586-022-05471-w.

Noble, Wendy, Diane P. Hanger, Christopher C. J. Miller, and Simon Lovestone. 2013. "The Importance of Tau Phosphorylation for Neurodegenerative Diseases." *Frontiers in Neurology* 4 (July):83.

Oddo, Salvatore, Antonella Caccamo, Jason D. Shepherd, M. Paul Murphy, Todd E. Golde, Rakez Kayed, Raju Metherate, Mark P. Mattson, Yama Akbari, and Frank M. LaFerla. 2003. "Triple-Transgenic Model of Alzheimer's Disease with Plaques and Tangles: Intracellular Abeta and Synaptic Dysfunction." *Neuron* 39 (3): 409–21.

Pang, Keliang, Richeng Jiang, Wei Zhang, Zhengyi Yang, Lin-Lin Li, Makoto Shimozawa, Simone Tambaro, et al. 2022. "An App Knock-in Rat Model for Alzheimer's Disease Exhibiting Aβ and Tau Pathologies, Neuronal Death and Cognitive Impairments." *Cell Research* 32 (2): 157–75.

Parag, Toufiq, Daniel Berger, Lee Kamentsky, Benedikt Staffler, Donglai Wei, Moritz Helmstaedter, Jeff W. Lichtman, and Hanspeter Pfister. 2019. "Detecting Synapse Location and Connectivity by Signed Proximity Estimation and Pruning with Deep Nets." In *Lecture Notes in Computer Science*, 354–64. Lecture Notes in Computer Science. Cham: Springer International Publishing.

Pavarino, Elisa C., Emma Yang, Nagaraju Dhanyasi, Mona D. Wang, Flavie Bidel, Xiaotang Lu, Fuming Yang, et al. 2023. "mEMbrain: An Interactive Deep Learning MATLAB Tool for Connectomic Segmentation on Commodity Desktops." *Frontiers in Neural Circuits* 17 (June):952921.

Peddie, Christopher J., Christel Genoud, Anna Kreshuk, Kimberly Meechan, Kristina D. Micheva, Kedar Narayan, Constantin Pape, et al. 2022. "Volume Electron Microscopy." *Nature Reviews. Methods Primers* 2 (July):51.

Rajendran, Lawrence, and Rosa C. Paolicelli. 2018. "Microglia-Mediated Synapse Loss in Alzheimer's Disease." *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience* 38 (12): 2911–19.

Saman, Sudad, Wonhee Kim, Mario Raya, Yvonne Visnick, Suhad Miro, Sarmad Saman, Bruce Jackson, et al. 2012. "Exosome-Associated Tau Is Secreted in Tauopathy Models and Is Selectively Phosphorylated in Cerebrospinal Fluid in Early Alzheimer Disease." *The Journal of Biological Chemistry* 287 (6): 3842–49.

Scheffer, Louis K., C. Shan Xu, Michal Januszewski, Zhiyuan Lu, Shin-Ya Takemura, Kenneth J. Hayworth, Gary B. Huang, et al. 2020. "A Connectome and Analysis of the Adult Drosophila Central Brain." *eLife* 9 (September). https://doi.org/10.7554/eLife.57443.

Scheff, Stephen W., and Douglas A. Price. 2001. "Alzheimer's Disease-Related Synapse Loss in the Cingulate Cortex." *Journal of Alzheimer's Disease: JAD* 3 (5): 495–505.

———. 2003. "Synaptic Pathology in Alzheimer's Disease: A Review of Ultrastructural Studies." *Neurobiology of Aging* 24 (8): 1029–46.

———. 2006. "Alzheimer's Disease-Related Alterations in Synaptic Density: Neocortex and Hippocampus." *Journal of Alzheimer's Disease: JAD* 9 (3 Suppl): 101–15.

Sedgwick, J. D., S. Schwender, H. Imrich, R. Dörries, G. W. Butcher, and V. ter Meulen. 1991. "Isolation and Direct Characterization of Resident Microglial Cells from the Normal and Inflamed Central Nervous System." *Proceedings of the National Academy of Sciences of the United States of America* 88 (16): 7438–42.

Seo, Junyoung, Yeonbo Sim, Jeewon Kim, Hyunwoo Kim, In Cho, Hoyeon Nam, Young-Gyu Yoon, and Jae-Byum Chang. 2022. "PICASSO Allows Ultra-Multiplexed Fluorescence Imaging of Spatially Overlapping Proteins without Reference Spectra Measurements." *Nature Communications* 13 (1): 2475.

Shapson-Coe, Alexander, Michał Januszewski, Daniel R. Berger, Art Pope, Yuelong Wu, Tim Blakely, Richard L. Schalek, et al. 2024. "A Petavoxel Fragment of Human Cerebral Cortex Reconstructed at Nanoscale Resolution." *Science* 384 (6696): eadk4858.

Sievers, Meike, Alessandro Motta, Martin Schmidt, Yagmur Yener, Sahil Loomba, Kun Song, Johannes Bruett, and Moritz Helmstaedter. 2024. "Connectomic Reconstruction of a Cortical Column." *bioRxiv*. https://doi.org/10.1101/2024.03.22.586254.

Sobotzik, Jürgen-Markus, Jana Maria Sie, Chrisoula Politi, Domenico Del Turco, Vann Bennett, Thomas Deller, and Christian Schultz. 2009. "AnkyrinG Is Required to Maintain Axo-Dendritic Polarity in Vivo." *Proceedings of the National Academy of Sciences of the United States of America* 106 (41): 17564–69.

St-Pierre, Marie-Kim, Micaël Carrier, Fernando González Ibáñez, Eva Šimončičová, Marie-Josée Wallman, Luc Vallières, Martin Parent, and Marie-Ève Tremblay. 2022. "Ultrastructural Characterization of Dark Microglia during Aging in a Mouse Model of Alzheimer's Disease Pathology and in Human Post-Mortem Brain Samples." *Journal of Neuroinflammation* 19 (1): 235.

Swanson, Larry W., and Jeff W. Lichtman. 2016. "From Cajal to Connectome and Beyond." *Annual Review of Neuroscience* 39 (July):197–216.

Taylor, J. Paul, John Hardy, and Kenneth H. Fischbeck. 2002. "Toxic Proteins in Neurodegenerative Disease." *Science* 296 (5575): 1991–95.

Tuijtel, Maarten W., A. Koster, S. Jakobs, F. Faas, and T. Sharp. 2019. "Correlative Cryo Super-Resolution Light and Electron Microscopy on Mammalian Cells Using Fluorescent Proteins." *Scientific Reports* 9 (February). https://doi.org/10.1038/s41598-018-37728-8.

Turner, Nicholas L., Thomas Macrina, J. Alexander Bae, Runzhe Yang, Alyssa M. Wilson, Casey Schneider-Mizell, Kisuk Lee, et al. 2022. "Reconstruction of Neocortex: Organelles, Compartments, Cells, Circuits, and Activity." *Cell* 185 (6): 1082–1100.e24.

Twohig, Daniel, and Henrietta M. Nielsen. 2019. "α-Synuclein in the Pathophysiology of Alzheimer's Disease." *Molecular Neurodegeneration* 14 (1): 23.

Tzioras, Makis, Robert I. McGeachan, Claire S. Durrant, and Tara L. Spires-Jones. 2023. "Synaptic Degeneration in Alzheimer Disease." *Nature Reviews. Neurology* 19 (1): 19–38.

Urbanc, B., L. Cruz, R. Le, J. Sanders, K. Hsiao Ashe, K. Duff, H. E. Stanley, M. C. Irizarry, and B. T. Hyman. 2002. "Neurotoxic Effects of Thioflavin S-Positive Amyloid Deposits in Transgenic Mice and Alzheimer's Disease." *Proceedings of the National Academy of Sciences of the United States of America* 99 (22): 13990–95.

Urbanski, Alysson H., Department of Clinical and Toxicological Analyses, School of Pharmaceutical Sciences, University of Sao Paulo, Sao Paulo, Brazil, José D. Araujo, Rachel Creighton, Helder I. Nakaya, Department of Clinical and Toxicological Analyses, School of Pharmaceutical Sciences, University of Sao Paulo, Sao Paulo, Brazil, Department of Bioengineering, University of Washington, Seattle, WA, USA, Department of

Clinical and Toxicological Analyses, School of Pharmaceutical Sciences, University of Sao Paulo, Sao Paulo, Brazil, and Scientific Platform Pasteur/USP, University of Sao Paulo, Sao Paulo, Brazil. 2019. "Integrative Biology Approaches Applied to Human Diseases." In *Computational Biology*, 19–36. Codon Publications.

Vilalta, Anna, and Guy C. Brown. 2018. "Neurophagy, the Phagocytosis of Live Neurons and Synapses by Glia, Contributes to Brain Development and Disease." *The FEBS Journal* 285 (19): 3566–75.

Wei, Lu, Zhixing Chen, Lixue Shi, Rong Long, Andrew V. Anzalone, Luyuan Zhang, Fanghao Hu, Rafael Yuste, Virginia W. Cornish, and Wei Min. 2017. "Super-Multiplex Vibrational Imaging." *Nature* 544 (7651): 465–70.

"Welcome to NeuroMab!" n.d. Accessed April 13, 2024. https://neuromab.ucdavis.edu/neuromabs.cfm.

Wilson, Andrea C., Brittany N. Dugger, Dennis W. Dickson, and Deng-Shun Wang. 2011. "TDP-43 in Aging and Alzheimer's Disease - a Review." *International Journal of Clinical and Experimental Pathology* 4 (2): 147–55.

Wilton, Emily E., Michael P. Opyr, Senthilkumar Kailasam, Ronja F. Kothe, and Hans-Joachim Wieden. 2018. "sdAb-DB: The Single Domain Antibody Database." *ACS Synthetic Biology* 7 (11): 2480–84.

Wood, Levi B., Ashley R. Winslow, and Samantha Dale Strasser. 2015. "Systems Biology of Neurodegenerative Diseases." *Integrative Biology: Quantitative Biosciences from Nano to Macro* 7 (7): 758–75.

Yin, Rui-Hua, Jin-Tai Yu, and Lan Tan. 2015. "The Role of SORL1 in Alzheimer's Disease." *Molecular Neurobiology* 51 (3): 909–18.

Yu, Meichen, Olaf Sporns, and Andrew J. Saykin. 2021. "The Human Connectome in Alzheimer Disease - Relationship to Biomarkers and Genetics." *Nature Reviews. Neurology* 17 (9): 545–63.

Zempel, Hans, Frank J. A. Dennissen, Yatender Kumar, Julia Luedtke, Jacek Biernat, Eva-Maria Mandelkow, and Eckhard Mandelkow. 2017. "Axodendritic Sorting and Pathological Missorting of Tau Are Isoform-Specific and Determined by Axon Initial Segment Architecture." *The Journal of Biological Chemistry* 292 (29): 12192–207.

Zimmermann, Iwan, Pascal Egloff, Cedric A. J. Hutter, Benedikt T. Kuhn, Philipp Bräuer, Simon Newstead, Roger J. P. Dawson, Eric R. Geertsma, and Markus A. Seeger. 2020. "Generation of Synthetic Nanobodies against Delicate Proteins." *Nature Protocols* 15 (5): 1707–41.

Zuiderveld, K. 1994. "Contrast Limited Adaptive Histogram Equalization." *Graphics Gems* 0:474–85.