Unexpected delayed incursion of highly pathogenic avian influenza H5N1 (clade 2.3.4.4b) in the Antarctic region

Simeon Lisovski[1], Anne Günther[2], Meagan Dewar[3], David Ainley[4], Fabián Aldunate[5,6], Rodrigo Arce[5,6], Grant Ballard[7], Silke Bauer [8], Josabel Belliure[9], Ashley C. Banyard[10,11], Thierry Boulinier[12], Ashley Bennison[13], Christina Braun[14], Craig Cary[15, †], Paulo Catry[16], Augustin Clessin[12], Maelle Connan[16], Edna Correia[17], Aidan Cox[7], Juan Cristina[5], Megan Elrod[7], Julia Emerit[18,19], Irene Ferreiro[5,6], Zoe Fowler[20], Amandine Gamble[18,19], José P. Granadeiro[17], Joaquin Hurtado[5,6], Dennis Jongsomjit[7], Célia Lesage[17], Mathilde Lejeune[12], Amanda Kuepfer[21], Amélie Lescroël[7], Amy Li[7], Ian R. McDonald[15], Javier Menéndez-Blázquez[22], Virginia Morandini[23], Gonzalo Moratorio[5,6], Teresa Militão[16], Pilar Moreno[5,6], Paula Perbolianachis[5,6], Jean Pennycook[4], Maryam Raslan[24], Scott M. Reid[10], Roanna Richards-Babbage[15], Annie E. Schmidt[7], Martha Maria Sander[1], Lucy Smyth[20], Alvaro Soutullo[24], Andrew Stanworth[21], Léo Streith[12,25], Jérémy Tornos[12], Arvind Varsani[26], Ulrike Herzschuh[1,27,28], Martin Beer[2], Michelle Wille[29,30]

[1] Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Polar Terrestrial Environmental Systems, 14473 Potsdam, Germany
[2] Friedrich-Loeffler-Institut, Greifswald, Isle of Riems, Germany
[3] Future Regions Research Centre, Federation University Australia
[4] H.T. Harvey & Associates Ecological Consultants, Los Gatos, CA 95032, USA
[5] Laboratorio de Virología Molecular, Centro de Investigaciones Nucleares, Facultad de Ciencias, Universidad de la República, Igua 4225, Montevideo 11400, Uruguay
[6] Laboratorio de Evolución Experimental de Virus, Institut Pasteur de Montevideo, Mataojo 2020, Montevideo 11400, Uruguay
[7] Point Blue Conservation Science. Petaluma, CA 94924, USA
[8] Federal Research Institute WSL, Birmensdorf 8903, Switzerland
[9] Universidad de Alcalá, A.P. 20 Campus Universitario, 28805 Alcalá de Henares, Madrid, España
[10] Department of Virology, Animal and Plant Health Agency-Weybridge, Woodham Lane, New Haw, Addlestone, Surrey, KT15 3NB, United Kingdom
[11] WOAH/FAO International Reference laboratory for Avian Influenza, Swine influenza and Newcastle disease virus, Animal and Plant Health Agency-Weybridge, Woodham Lane, New Haw, Addlestone, Surrey, KT15 3NB, United Kingdom
[12] Centre d'Ecologie Fonctionnelle et Evolutive (CEFE), CNRS, Université Montpellier, EPHE, IRD, Montpellier, France
[13] British Antarctic Survey, High Cross, Madingley, Cambridge, United Kingdom
[14] Friedrich Schiller University Jena, Institute for Ecology, Jena, Germany
[15] Thermophile Research Unit, Te Aka Mātuatua-School of Science, Te Whare Wānanga o Waikato, University of Waikato, Hamilton, Aotearoa-New Zealand
[16] MARE – Marine and Environmental Sciences Centre / ARNET - Aquatic Research Network, Ispa – Instituto Universitário, Portugal
[17] Centro de Estudos do Ambiente e do Mar (CESAM), Departamento de Biologia Animal, Faculdade de Ciências da Universidade de Lisboa, 1749 016 Lisbon, Portugal
[18] Cornell University, Department of Public and Ecosystem Health, Ithaca NY 14853, USA
[19] Ecole Normale Superieure de Lyon, Lyon 69000, France
[20] Department of Zoology, Marine Apex Predator Research Unit (MAPRU), Institute for Coastal and Marine Research, Nelson Mandela University, Gqeberha 6001, South Africa
[21] Falklands Conservation, Stanley, FIQQ 1ZZ, Falkland Islands
[22] Cavanilles Institute of Biodiversity and Evolutionary Biology, University of Valencia, Valencia, Spain
[23] Department of Public & Ecosystem Health, Cornell University, New York, United States
[24] Departamento de Ecologia y Gestion Ambiental, Centro Universitario Regional del Este, Universidad de la Republica, Maldonado, Uruguay
[25] Centre d'Ecologie Fonctionnelle et Evolutive (CEFE), CNRS, Université Montpellier, EPHE, IRD, Montpellier, France
[26] Biodesign Center for Fundamental and Applied Microbiomics, Center for Evolution and Medicine, School of Life Sciences, Arizona State University, Tempe, AZ, USA
[27] Institute of Biochemistry and Biology, University of Potsdam, Potsdam, Germany
[28] Institute of Environmental Science and Geography, University of Potsdam, Potsdam, Germany
[29] Centre for Pathogen Genomics, Department of Microbiology and Immunology, at the Peter Doherty Institute for Infection and Immunity, The University of Melbourne, Melbourne, Victoria, Australia
[30] WHO Collaborating Centre for Reference and Research on Influenza, at the Peter Doherty Institute for Infection and Immunity, Melbourne, Victoria, Australia.

Abstract

The current highly pathogenic avian influenza H5N1 panzootic has substantial impacts on wild birds and marine mammals. Although major outbreaks occurred in South America, incursion to Antarctica emerged late in the breeding season of 2023/2024 and was confined the wider region of the Antarctic Peninsula. To infer potential underlying processes, we compiled H5N1 surveillance from Antarctica and Sub-Antarctic Islands prior to the first confirmed cases.

Main text

The increasing intensity of highly pathogenic avian influenza virus (HPAIV) H5N1 clade 2.3.4.4b outbreaks have had a substantial impact on poultry and wildlife [1]. Wild bird movements have underpinned the rapid spread of this virus that swept across most continents, except for Australia and Antarctica, within two years [2]. Compared to previous HPAIV subtypes and clades, H5N1 2.3.4.4b has significantly improved replication in wild birds [3], and increased fitness through continuous reassortments [4] which has likely contributed to a shift in infection dynamics leading to the infection of a broader range of avian species [1]. In addition to their role as viral spreaders, wild birds are suffering huge losses following mass mortality events, and the scale of mortality amongst wild birds is likely in the millions rather than tens of thousands reported [5]. Thus, the recent panzootic is a serious conservation concern for a large range of wild bird species.

Due to the absence of waterfowl species that migrate to the Antarctic and sub-Antarctic islands, the incursion risk of HPAIV in these southernmost regions had been considered low prior to 2021. However, waterfowl are present in northern fringes of the Southern Ocean, and millions of known migration and post-breeding dispersal routes establish links and thereby substantial global connectivity, including with regions of recent HPAI H5N1 outbreaks involving seabirds and marine mammals [2]. Despite the perceived remoteness, low pathogenicity avian influenza viruses and antibodies against these viruses have previously been detected in various seabird species nesting at sites along the Antarctic Peninsula and South Shetland Islands, with viral genomes illustrating phylogenetic connectivity to viruses circulating on other continents [6,7]. As a result, the Scientific Committee on Antarctic Research (SCAR) Antarctic Wildlife Health Network (AWHN) had considered the risk of incursion of the recent panzootic HPAIV H5 into the Antarctic region in 2022/23 summer season to be high [8], and considerably higher in 2023/24 following virus spread to the southernmost regions of South America [9], infecting

seabirds including Magellanic penguins (*Spheniscus magellanicus*) and Humbold

penguins (*Spheniscus humboldti*), and several species of marine mammals (9).

To identify possible incursions of H5N1 into the Antarctic region during the

summer season 2022/23 and the early season 2023/24, we sampled migratory seabirds

at different locations across Antarctica and in sub-Antarctic areas (Figure 1), and

collated a range of observation data. Here, we define Antarctica as the region south of

the Antarctic Convergence and sub-Antarctic areas include adjacent Islands within the

Southern Ocean. In particular, we aimed to collect information pertaining to suspicious

signs of unusual mortality and known clinical signs of HPAIV infection including loss of

coordination and balance, trembling head and body, lethargy, respiratory distress, and

conjunctivitis [8]. Across all locations, sample collection was done in accordance with

institutional animal ethics approval and sample testing was performed with national

frameworks, with details available in the technical annex.

64
Figure 1: Sampling locations for qPCR analysis and the detection of H5N1 2.3.4.4b, as well as
locations with intensive observational efforts to identify signs of HPAI infections within breeding bird
communities for the breeding season 2022/23 (left) and 2023/24 (right). In addition, locations of
confirmed cases of infection in 2023/24 (left) are included. Numbers refer to the following references,
(a) technical annex, (b) [10], (c, e) [11], (d, f) [12]. Maps created with Natural Earth.

Overall, sampling and observational efforts were conducted from early November

2022 to late March 2023, and from October 2023 onwards until the end of February in

2024. Surveillance efforts included a large range of species (*i.e.,* penguins, gulls, skuas,

and petrels; see technical annex for more information) and locations. In 2022/23,

samples for HPAIV testing were collected from apparently healthy birds from 20

locations in the sub-Antarctic and Antarctic Region. There were several suspicious

observations of dead wild birds on the Falkland Islands (Gentoo penguin *Pygoscelis*

papua, Cattle egret *Bubulcus ibis*), and South Georgia (Wandering albatrosses

Diomedea exulans). However, all swab samples collected from these animals, in

addition to apparently healthy wild birds in other locations were negative for HPAI (see

technical annex for details on location and species). Together, this strongly suggests

that HPAIV H5N1 clade 2.3.4.4b did not enter the Antarctic region during the austral

summer 2022/23, and that the lack of detection was unlikely due to lack of surveillance,

testing or disease investigations. This is in contrast to the seabird breeding season

2023/24. In October 2023 the first confirmed H5N1 cases were detected on the Falkland

(Malvinas) Islands, and in November on South Georgia Island in the sub-Antarctic [10,12]

(Figure 1). Given the overlap of species breeding and migrating via the Falkland Island

and South Georgia towards the Antarctic Peninsula and its offshore Islands (e.g., the

South Shetland Islands), researchers in the region and the tourist industry have been

very diligent in identifying unusual bird behaviour and mortality events. Despite active

cases in the Falkland Islands and South Georgia Island, sample collection and

observations from 16 locations between November 2023 – early February 2024 in the

Antarctic Peninsula and related island were negative for HPAIV. Data from the SCAR

monitoring project did, however, report suspected cases in the Antarctic region starting

in December 2023 [12]. These include Brown Skuas on the South Orkney Islands in

December 2023 (no samples collected), a mortality in Brown Skuas on Heronia Island in

December 2023 (samples collected, HPAIV negative). Since mid-February several

positive cases were reported from the Antarctic Peninsula (see Figure 1). This suggests

that H5N1 was spread among colonies in the later breeding season, however, so far

there is no evidence for large outbreaks on the Antarctic Peninsula. Further, based on

observation data, the strain did not appear to have reached the Indian Ocean sub-

Antarctic islands as of February 2024 (see technical annex).

Obviously, incursion risk, and successful establishment of HPAIV is contingent on

a combination of factors. Most importantly, that (i) host species get in contact with

HPAIV before travelling into the Antarctic regions, (ii) can migrate with an infection, and

(iii) have contact and transmit the virus to susceptible species which could be the

starting point of a new epizootic. Most species occupying the Antarctic region are

pelagic seabirds with little to no contact with terrestrial birds such as waterfowl,

significantly reducing their exposure to outbreaks on land (e.g. South America).

However, some species like the Brown skua (*Stercorarius antarcticus*) and the giant

petrel species (*Macronectes sp.)* are known scavengers (Figure 2), leading to high risks

of exposure to HPAIV via the consumption of infected carcasses. It is thus no surprise

that Brown skuas where often among the first confirmed cases both on South Georgia
and the Antarctic Peninsula [10]. This species, which can be observed at shorelines of
South America, the Falkland Islands and South Georgia [13], is likely to be an important
player in spreading the virus. Yet, it seems that the connectivity established by the
animals' movements from South America and South Georgia over the Drake Passage to
Antarctica is rather limited during the breeding season but might increase again towards
the end when the breeding activities terminate, and the movement range of both adults
and first juveniles are becoming larger again. Together with the increasing number of
naïve juveniles and concomitant changes in densities, this may explain the delay
between initial outbreaks in the Falklands/South Georgia and the first confirmed cases
on the Antarctic peninsula.
124

125
126
127   *Figure 2: Northern giant petrels and Brown skuas scavenging on an Antarctic fur seal carcass,*
128   *showing inter-species interactions with the potential for HPAI virus transmission (photo taken on*
129   *South Georgia by Paulo Catry).*
130
Still, the consequences of viral incursion(s) into Southern Ocean wildlife are
unclear but based on observations from other regions, will likely have devastating
effects. Critically, densities of seabird colonies are very high, facilitating the
transmissions between individuals [14]. Further prospecting movements of potential
recruits, predator-prey interactions between bird species (e.g., skuas, penguins, and
sheathbills), as well as species scavenging on dead seabirds and mammals, may
promote rapid spread of the virus between colonies [15]. Once the virus has been
established in the region, interaction between seabirds and marine mammals may also
result in further transmissions and facilitate the adaptation of the virus in mammalian

species [14]. Finally, most animals of the Southern Ocean are endemic to the region, such

that mass mortality events in Antarctica due to HPAIV H5 will cause a very real

conservation concern for many species.

Detecting H5N1 incursion(s) and describing the infection dynamics into and

within the sub-Antarctic and Antarctic regions is highly relevant and standardized

surveys for mortality and sampling should therefore be prioritized. These activities

should be undertaken with consideration of the potentially zoonotic risks of (emerging)

HPAIV H5 [8] and require strict hygiene measures to prevent the spread of the virus

through human activities. Sampling and detailed analysis of lineages and virus

phenotype will provide crucial information needed to assess risks and respond to future

wild bird outbreaks.

Acknowledgements

We thank the following institutions and agencies for financial and logistical support: The

Uruguayan Antarctic Institute; Government South Georgia and South Sandwich Islands;

Antarctica New Zealand; French Polar Institute (ECOPATH-1151); National Nature

Reserve of Terres Australes et Antarctique Françaises; The South African Departments of

Science and Innovation (National Research Foundation - South African National Antarctic

Programme project no. 129226; MC) and of Forestry, Fisheries and the Environment;

Falklands Conservation, the Wildlife Conservation Society; Falkland Islands

Government; The Antarctic Research Trust; The South Atlantic Environmental Research

Institute; Commission for the Conservation of Antarctic Marine Living Resources

(CCAMLR); Spanish Research Agency (VM). The study was supported by the following

research grants: ANR (ECOPATHS ANR-21-CE35-0016 and REMOVE_DISEASE ANR-

21-BIRE-006-01; TB, ML, JT); Biodiversa and Water JPI under the BiodivRestore ERA-

NET Cofund (GA N°101003777; TB, MC, ML, AC, PC); National Science Foundation

award ANT 1935870 and ANT 2040199 (GB, DA, ,ME, AC AS, AL, AL, AV, JP); Royal

Society (Newton International Fellowship NIF\R1\211869; AGa); Biodiversity Challenge

Fund (Darwin Plus Grant DPLUS167; AGa); Fundação para a Ciência e a Tecnologia

(FTC) – Portugal (UIDB/04292/2020 (MARE) and LA/P/0069/2020 (ARNET) and

DivRestore/0012/2020 (REMOVE_DISEASE project), Agencia Nacional de Investigación

e Innovación's (ANII) Clemente Estable Fund (project FCE_1_2021_1_166587); Ecos-

Sud Program (project PU20B01/U20B03); General Capacity Building Fund; the

Department for Environment, Food and Rural Affairs (Defra, UK) and the devolved

administrations of Scotland and Wales under grant numbers SV3006, SE2213 and

SV3045; Biotechnology and Biological Sciences Research Council (BBSRC) and Defra

funded research initiative 'FluMAP' [grant number BB/X006204/1, ACB, SMR].

European Union Horizon Europe research program (grant agreement "Kappa-Flu No.

101084171", with the Schweizer Staatssekretariat, AGü, MB, SB). In addition, we thank

Corisande Abiven, Katharina Reusch, Louise Pole-Evans, Edna Correia, Jose P

Granadeiro, Michelle Risi and Christopher Jones for fieldwork support, and all the

inhabitants and visitors of the Falkland Islands for reporting suspected cases and

facilitating research on their lands, in particular Kicki Ericson and Thies Matzen, Sarah

Crofts and Micky Reeves, and the Rendell, Pole-Evans, Gould, Delignieres and Hazel

families.

References

1. Wille M, Barr IG. Resurgence of avian influenza virus. Science 2022;376:459-460.

2. Adlhoch C, Fusaro A, Gnzales JL, et al. Avian influenza overview March - April 2023.
Efsa Journal 2023;21.

3. James J, Billington E, Warren CJ, et al. Clade 2.3.4.4b H5N1 high pathogenicity avian
influenza virus (HPAIV) from the 2021/22 epizootic is highly duck adapted and poorly
adapted to chickens. Journal of General Virology 2023;104.

4. Kandeil A, Patton C, Jones JC, et al. Rapid evolution of A(H5N1) influenza viruses after
intercontinental spread to North America. Nature Communications 2023;14:3082.

5. WOAH. WAHIS: World Animal Health Information System. Available at
https://wahis.woah.org/.

6. Hurt Aeron C., Vijaykrishna Dhanasekaran, Butler Jeffrey, et al. Detection of
Evolutionarily Distinct Avian Influenza A Viruses in Antarctica. mBio
2014;5:10.1128/mbio.01098-14.

7. de Seixas MMM, de Araújo J, Krauss S, et al. H6N8 avian influenza virus in Antarctic
seabirds demonstrates connectivity between South America and Antarctica.
Transboundary and Emerging Diseases 2022;69:e3436-e3446.

8. Dewar M, Wille M, Gamble A, et al. The Risk of Avian Influenza in the Southern Ocean:
A practical guide. 2022.

9. Godoy M, de Oca MM, Caro D, Pontigo JP, Kibenge M, Kibenge F. Evolution and
Current Status of Influenza A Virus in Chile: A Review. Pathogens 2023;12:1252.

10. Ashley Bennison, Alexander M. P. Byrne, Scott M. Reid, et al. Detection and spread of
high pathogenicity avian influenza virus H5N1 in the Antarctic Region. bioRxiv January
2023:2023.11.23.568045.

11. Fabiola León, Céline Le Bohec, Eduardo J. Pizarro, et al. Highly Pathogenic Avian
Influenza A (H5N1) Suspected in penguins and shags on the Antarctic Peninsula and
West Antarctic Coast. bioRxiv January 2024:2024.03.16.585360.

12. SCAR. Sub-Antarctic and Antarctic Highly Pathogenic Avian Influenza H5N1 Monitoring
Project | SCAR. Available at https://scar.org/library-data/avian-flu. Accessed 30 March
2024.

13. Krietsch J, Hahn S, Kopp M, Phillips RA, Peter HU, Lisovski S. Consistent variation in
individual migration strategies of brown skuas. Marine Ecology Progress Series
2017;578:213-225.

14. Dewar M, Wille M, Gamble A, et al. Risk Assessment of Highly Pathogenic Avian
Influenza in the Southern Ocean Report prepared for Scientific Committee on Antarctic
Research. 2023.

15. Boulinier T. Avian influenza spread and seabird movements between colonies. Trends in
Ecology & Evolution 2023;38:391-395.

223