Adipokinetic hormone signaling mediates the enhanced fecundity of *Diaphorina citri* infected by '*Candidatus Liberibacter asiaticus*'

Jiayun Li[1], Paul Holford[3], George Andrew Charles Beattie[3], Shujie Wu[1], Jielan He[1], Shijian Tan[1], Desen Wang[1], Yurong He[1], Yijing Cen[1]*, Xiaoge Nian[1,2]*

[1]National Key Laboratory of Green Pesticide, Department of Entomology, College of Plant Protection, South China Agricultural University, Guangzhou, 510642, China；

[2]Henry Fok School of Biology and Agriculture, Shaoguan University, Shaoguan, 512005, China；

[3]School of Science, Western Sydney University, Locked Bag 1797, Penrith, NSW 2751, Australia;

*Corresponding author: Yijing Cen and Xiaoge Nian

Email: cenyj@scau.edu.cn; nianxiaoge@scau.edu.cn

Abstract

Diaphorina citri serves as the primary vector for '*Candidatus* Liberibacter asiaticus' (*C*Las), the bacterium associated with the severe Asian form of huanglongbing. *C*Las-positive *D. citri* are more fecund than their *C*Las-negative counterparts and require extra energy expenditure. Therefore, understanding the molecular mechanisms linking metabolism and reproduction is of particular importance. In this study, we found adipokinetic hormone (*DcAKH*) and its receptor (*DcAKHR*) were essential for increasing lipid metabolism and fecundity in response to *C*Las infection in *D. citri.* Knockdown of *DcAKH* and *DcAKHR* not only resulted in the accumulation of triacylglycerol and a decline of glycogen, but also significantly decreased fecundity and *C*Las titer in ovaries. Combined *in vivo* and *in vitro* experiments showed that miR-34 suppresses *DcAKHR* expression by binding to its 3' untranslated region, whilst overexpression of miR-34 resulted in a decline of *DcAKHR* expression and *C*Las titer in ovaries and caused defects that mimicked *DcAKHR* knockdown phenotypes. Additionally, knockdown of *DcAKH* and *DcAKHR* significantly reduced juvenile hormone (JH) titer and JH signaling pathway genes in fat bodies and ovaries, including the JH receptor, *methoprene-tolerant* (*DcMet*), and the transcription factor, *Krüppel homolog 1 (DcKr-h1)*, that acts downstream of it, as well as the egg development related genes *vitellogenin 1-like* (*DcVg-1-like*), *vitellogenin A1-like* (*DcVg-A1-like*) and the vitellogenin receptor (*DcVgR*). As a result, *C*Las hijacks AKH/AKHR-miR-34-JH signaling to improve *D. citri* lipid metabolism and fecundity, while simultaneously increasing the replication of *C*Las, suggesting a mutualistic interaction between *C*Las and *D. citri* ovaries.

Key words: huanglongbing; *Diaphorina citri*; *DcAKH*/*DcAKHR*; miR-34; mutualism

Introduction

The majority of devastating plant pathogens, including viruses and bacteria, are transmitted by insect vectors [1]. In recent decades, the interactions between vectors and pathogens have been widely studied [1-4]. However, studies on the mechanisms underlying these interactions have mainly focused on vector-virus or vector-fungus interactions; less research has been performed on vector-bacteria interactions. Knowledge of the relationships between vectors and pathogens is of great significance for understanding the epidemiology of plant pathogens, and this knowledge may help develop new strategies for controlling both vectors and pathogens. The severe Asian form of huanglongbing (HLB) is currently the most destructive citrus disease in Asia and the Americas where it is associated with the phloem-restricted, Gram-negative, α-Proteobacteria 'Candidatus Liberibacter asiaticus' (CLas) and transmitted by the Asian citrus psyllid, Diaphorina citri Kuwayama (Hemiptera: Psyllidae) [5-9]. However, since CLas has not been cultured in vitro, Koch's postulates have not been proven. Control of D. citri is the most effective way to prevent HLB epidemics [6,7] but experience in Asia [10] and the United States of America [11,12], and recent press reports from Brazil, indicate that it is impossible to suppress psyllid populations with insecticides to levels that prevent spread of CLas.

Previous studies have reported that infection with CLas significantly increases the fecundity of D. citri [13-15]. Reproduction is a costly process in terms of energy usage in the adult life of female insects [16], and there is an inevitable trade-off between lipid storage and use, because animals mobilize their lipid reserves during reproduction. However, the mechanism of how D. citri maintains a balance between lipid metabolism and increased fecundity after infection with CLas remains unknown. In insects, energy mobilization is under control of adipokinetic hormone (AKH), which was the first insect neurohormone to be identified; it was shown to stimulate lipid mobilization for locomotor activity in locusts [17]. To date, more than 80 different forms of AKH have been identified or predicted in arthropod species due to similar structural characteristics of the enzyme [18]. The mechanisms of synthesis and biological activity of AKH have also been investigated. Insect store lipids in the form of triacylglycerides (TAG) and as carbohydrates in the form of glycogen in their fat bodies [19]. Under conditions of energy demand, AKHs are synthesized in the corpora cardiaca,

released into the hemolymph and bind to receptors (AKHRs) located on the plasma membrane

of fat body cells. AKHR binding and activation triggers the release of energy-rich substrates

after which various cellular signaling pathway components come into play [20]. AKHRs are a

class of G protein-coupled receptors and were first identified in *Drosophila melanogaster* [21]

and *Bombyx mori* [22]; subsequently, they were identified in a large number of insect species

[23-26].

While the AKH pathway is recognized for its role in mobilizing stored lipids, its involvement

in vector-pathogen interactions remains understudied. For example, in *Anopheles*

gambiae-*Plasmodium falciparum* interactions. *P. falciparum* infection influences *A. gambiae*

AKH signaling, lipid metabolism, and mobilization of energy reserves to support the parasite's

metabolic and structural needs [27]. In another example, infection of cockroaches with the

fungus *Isaria fumosorosea* result in a significant rise in AKH levels in the central nervous

system in response to stress responses [28]. The functions of the AKH/AKHR in

insect-pathogen interactions need further investigation. Recent studies have shown that

AKH/AKHR signaling plays a regulatory role during female reproduction in insects. For

example, depletion of AKHR in *Bactrocera dorsalis* resulted in TAG accumulation, decreased

sexual courtship activity, and fecundity [29]. AKHR knockdown in *Nilaparvata lugens* interferes

with trehalose homeostasis and vitellogenin uptake by oocytes, which causes delayed oocyte

development and reduced fecundity [30]. AKH/AKHR signaling also regulates vitellogenesis

and egg development in *Locusta migratoria* via triacylglycerol mobilization and trehalose

homeostasis [31]. As outlined above, the fecundity of *D. citri* is increased by *C*Las. Whether

AKH/AKHR signaling participates in this increase in fecundity due to infection via alterations in

lipid metabolism is not currently known.

Although numerous studies have been conducted on the AKH/AKHR signaling pathway,

miRNA-mediated regulation of the AKH signaling pathway at the post-transcriptional level has

been little studied. MicroRNAs (miRNAs) are small, non-coding RNAs containing

approximately 22 nucleotides that act mostly on the 3' untranslated region (UTR) of target

mRNAs, resulting in translation repression or mRNA degradation [32]. miRNAs play important

roles in regulating cellular events at the post-transcriptional level during biological processes

[32,33,34]. For example, in *D. citri*-*C*Las interaction, *C*Las hijacks the JH signaling pathway

and host miR-275 that targets the *vitellogenin receptor* (*DcVgR*) to improve *D. citri* fecundity,

while simultaneously increasing the replication of *C*Las itself, suggesting a mutualistic

interaction in *D. citri* ovaries with *C*Las [35]. In term of host-virus interactions, miR-8 and

miR-429 target *Broad isoform Z2*, the gene involved in the hormonal regulation of single

nucleopolyhedrovirus (HaSNPV)-mediated climbing behavior in *Helicoverpa armigera* [36]. In

relation to reproduction, in *Aedes aegypti*, miR-275 is essential for egg development [37],

miR-309 for ovarian development [38] and miR-8 for other reproductive processes [39]. As for

lipid metabolism, knockdown of miR-277 in *A. aegypti* activates insulin/FOXO signaling by

increasing the expression of ilp7 and ilp8, which reduces TAG levels and inhibits ovarian

development [40]. Nevertheless, the functions of the miRNAs in insect-pathogen interactions

need further investigation. To date, there are no reports that miRNAs participate in *D.*

citri-*C*Las interactions or are hijacked by *C*Las to affect lipid metabolism and the fecundity of *D.*

citri. As a consequence, in this study, we used the *D. citri*-*C*Las system as a model to study the

molecular events associated with AKH/AKHR signaling that affect lipid metabolism and

increase fecundity of *D. citri* induced by *C*Las. Our study contributes to our understanding of

the interactions between vectors and pathogens and may also provide new ways for

controlling *D. citri* and HLB.

Results

Effects of *C*Las infection on the reproduction and metabolism of adult *D. citri*.

Total TAG and glycogen levels in fat bodies of *C*Las-negative and *C*Las-positive psyllids were determined. After infection with *C*Las, the TAG and glycogen levels of *C*Las-positive psyllids significantly increased compared with *C*Las-negative individuals (Figures 1A-B). Since TAG is mainly stored in fat bodies as lipid droplets, we next evaluated the changes in the lipid droplet size using Nile red staining. As shown in Figure 1C, the size of lipid droplets in the *C*Las-positive group was significantly larger than those of the *C*Las-negative group. In order to evaluate the effects of *C*Las on the reproduction of *D. citri*, the preoviposition period, oviposition period and fecundity of both *C*Las-negative and *C*Las-positive psyllids were examined. After infection with *C*Las, the preoviposition period was significantly shortened, the oviposition period was extended, and the fecundity was markedly increased (Figures 1D-E).

Figure 1. Effects of *C*Las on lipid metabolism and reproductive behavior of *D. citri*.

(A) Comparison of TAG levels in the fat bodies of *C*Las-positive (*C*Las+) and *C*Las-negative

(*C*Las-) females 5, 9, 13 days after emergence (DAE). (B) Comparison of glycogen levels in

the fat bodies of *C*Las+ and *C*Las- females 5, 9, 13 DAE. (C) Lipid droplets in fat bodies

dissected from *C*Las+ and *C*Las- females 9 DAE stained with Nile red. Scale bar = 40 μm.

(D-F) Comparison of the preoviposition period, oviposition period and the fecundity of *C*Las+

and *C*Las- adults. Data are shown as means ± SEM. The significant differences between

*C*Las-positive and *C*Las-negative psyllids were determined using Student's *t*-tests (**$p < 0.01$,

***$p < 0.001$).

140 *DcAKH* is involved in changes in metabolism and fecundity in *D. citri* mediated by *C*Las

Using our transcriptome database, cDNA sequences of *DcAKH* were identified and

cloned. The cDNA sequence of *DcAKH* is 225 bp in length, encodes a deduced polypeptide of

74 amino acid residues and has the conserved structural characteristics of a neuropeptide.

The conserved structure was "signal peptide + mature peptide + related peptide", and the

mature peptide sequence of *Dc*AKH is "QVNFSPNW" (Figure S1A). A phylogenetic analysis

was conducted to evaluate the association of *Dc*AKH with other insect AKHs; *Dc*AKH was

most closely related to the AKHs of three hemipteran species, *Nilaparvata lugens*, *Sitobion*

avenae and *Acyrthosiphon pisum* (Figure S1B).

Tissue expression profiles showed that *Dc*AKH was expressed at the highest level in the

brain, significantly higher than in the ovaries, fat bodies, and midgut in *D. citri* (Figure S2). The

levels of *DcAKH* in ovaries of *C*Las-negative and -positive psyllids decreased during the

assessment period. However, levels in the ovaries of *C*Las-positive psyllids were significantly

higher than those that were *C*Las-negative at all assessment times (Figure 2A). To investigate

the role of *DcAKH* in metabolic and reproductive changes induced by *C*Las, we knocked down

DcAKH expression using RNAi. When fed with ds*DcAKH*, the mRNA level of *DcAKH* in

*C*Las-negative and *C*Las-positive ovaries significantly decreased by 89% and 66%,

respectively, compared with psyllids fed with ds*GFP* (Figure 2B). Knockdown of *DcAKH*

resulted in TAG accumulation and a significant decrease in glycogen (Figures 2C-D).

Correspondingly, depletion of *DcAKH* led to a significant increase the size of the lipid droplets

(Figure 2E). Compared with the controls, *DcAKH* RNAi slowed ovarian development (Figure

2F), extended the preoviposition period (from 13.3 to 14.2 days in *C*Las-negative psyllids and

from 10.7 to 13.4 days in *C*Las-positive psyllids), shortened the oviposition period (from 7.1 to

4.3 days in *C*Las-negative psyllids and from 10.1 to 5.7 days in *C*Las-positive psyllids), and

decreased fecundity (from 138.0 to 94.7 days in *C*Las-negative psyllids and from 230.0 to

118.1 days in *C*Las-positive psyllids) (Figures 2G-I). Furthermore, the *C*Las signal and relative

titer in the ovaries of *DcAKH* knockdown psyllids significantly decreased (Figures 2J-K). Taken

together, it can be concluded that *DcAKH* is involved in *C*Las-induced metabolic and

reproductive changes in *D.citri.*

Figure 2. *DcAKH* is involved in the mutualistic relationship between *C*Las and *D. citri* resulting in increased fecundity. (A) Comparison of temporal expression patterns of *DcAKH* between the ovaries of *C*Las- and *C*Las+ psyllids. (B) Efficiency of RNAi of *DcAKH* in *C*Las- and *C*Las+ females treated with ds*DcAKH* for 48 h. (C) Comparison of TAG levels in fat bodies of *C*Las- and *C*Las+ females treated with ds*DcAKH* for 48 h. (D) Comparison of glycogen levels in fat bodies of *C*Las- and *C*Las+ females treated with ds*DcAKH* for 48 h. (E) Lipid droplets stained with Nile red in fat bodies dissected from *C*Las+ females treated with ds*DcAKH* for 48 h. Scale bar = 40 μm. (F) Ovary phenotypes of *C*Las+ females treated with

ds*DcAKH* for 48 h. Scale bar = 200 μm. o: ovary, s: spermathecae. (G-I) Comparison of the

preoviposition period, oviposition period and the fecundity between *C*Las- and *C*Las+ adults

treated with ds*DcAKH*. (J) The *C*Las titer in ovaries of *C*Las+ females treated with ds*DcAKH*

for 48 h. (K) Representative confocal images of *C*Las in the reproductive system of *C*Las+

females treated with ds*DcAKH* for 48 h. Scale bar = 200 μm. DAPI: the cell nuclei were stained

with DAPI and visualized in blue. *C*Las-Cy3: the *C*Las signal visualized in red by staining with

Cy3. Merge: merged imaging of co-localization of cell nuclei and *C*Las. Data are shown as

means ± SEM. The significant differences between treatment and controls are indicated by

asterisks (Student's *t*-test, $*p < 0.05$, $**p < 0.01$, $***p < 0.001$).

***DcAKHR* respond to *DcAKH* and participates in *D. citri*-*C*Las mutualism**

The protein tertiary structure of *DcAKHR* was predicted; *DcAKHR* has seven conserved

transmembrane (TM1-TM7) domains and is a typical G protein coupled receptor (Figure S3A).

A phylogenetic analysis was conducted to evaluate the association of *DcAKHR* with other

insect *AKHRs*; *DcAKHR* was most closely related to the *AKHR* of *Nilaparvata lugens* (Figure

S3B). To further confirm *Dc*AKH can activate *DcAKHR*, we performed a cell-based calcium

mobilization assay. As shown in Figure S3C, *DcAKHR* was strongly activated by *Dc*AKH in a

dose-dependent manner, with an EC_{50} value of 2.20×10^{-5} mM, but there were no responses

when challenged with an empty vector or other evolutionarily-related peptides. In addition,

after *DcAKH* knockdown, *DcAKHR* expression in the ovaries significantly decreased (Figure

S3D). *DcAKHR* was cloned, sequenced, designated and then used for further studies.

Sequence analysis showed that *DcAKHR* consisted of an open reading frame (ORF) of 1290

bp and a 185 bp 3' UTR. It encodes a putative protein consisting of 429 amino acids with a

predicted molecular weight of 48.32 kDa. Tissue expression patterns showed that *DcAKHR*

had the highest expression in the midgut, follow by the head, fat bodies, and ovaries (Figure

S4). In the ovaries of *C*Las-positive and -negative psyllids, the transcript and protein levels of

DcAKHR increased over the assessment period with levels of both being higher in

*C*Las-positive psyllids than in *C*Las-negative psyllids (Figure 3A). To investigate the role of

DcAKHR in metabolic and reproductive changes induced by *C*Las, we knocked down

DcAKHR expression by RNAi. When fed with ds*DcAKHR*, the mRNA levels of *DcAKHR* in the

ovaries of *C*Las-negative and *C*Las-positive were significantly decreased by 59.4 and 66.0%,

respectively, compared with psyllids fed with ds*GFP*; western blot analysis confirmed that

*Dc*AKHR protein levels were also significantly reduced (Figure 3B). Knockdown of *DcAKHR*

resulted in TAG accumulation and a significant decrease in glycogen (Figures 3C-D).

Correspondingly, depletion of *DcAKHR* led to a significant increase in the size of lipid droplets

(Figure 3E). Compared with the controls, *DcAKHR* RNAi interfered with ovarian development

(Figure 3F): the preoviposition period was extended (from 13.3 to 16.2 days in *C*Las-negative

psyllids and from 10.7 to 12.9 days in *C*Las-positive psyllids); oviposition period was shortened

(from 7.1 to 4.6 days in *C*Las-negative psyllids and from 10.1 to 5.3 days in *C*Las-positive

psyllids); and fecundity decreased (from 138.0 to 90.6 eggs per female in *C*Las-negative

psyllids and from 230.0 to 98.2 eggs per female in *C*Las-positive psyllids). These effects are

similar to those caused by *DcAKH* knockdown (Figures 3G-I). Furthermore, the *C*Las signal

and relative titer in the ovaries of *DcAKH* knockdown psyllids were significantly decreased

(Figures 3J-K). Taken together, *DcAKHR* responds to *DcAKH* and is involved in the

*C*Las-induced metabolic and reproductive changes in *D. citri.*

Figure 3. *DcAKHR* is involved in the mutualistic relationship between *C*Las and *D. citri* resulting in increased fecundity. (A) Comparison of temporal expression patterns of *DcAKHR* between the ovaries of *C*Las- and *C*Las+ psyllids. (B) The efficiency of RNAi of *DcAKHR* in *C*Las- and *C*Las+ psyllids treated with ds*DcAKHR* for 48 h. (C) Comparison of TAG levels in fat bodies of *C*Las- and *C*Las+ females treated with ds*DcAKHR* for 48 h. (D) Comparison of glycogen levels in fat bodies of *C*Las- and *C*Las+ females treated with ds*DcAKHR* for 48 h. (E) Lipid droplets stained with Nile red in fat bodies dissected from *C*Las-positive females treated with ds*DcAKHR* for 48 h. Scale bar = 40 µm. (F) Ovary

phenotypes of CLas+ females treated with dsDcAKHR for 48 h. Scale bar = 200 μm. o: ovary,

s: spermathecae. (G-I) Comparison of the preoviposition period, oviposition period and the

fecundity of CLas- and CLas+ adults treated with dsDcAKHR. (J) The CLas titer in the ovaries

of CLas+ females treated with dsDcAKHR for 48 h. (K) Representative confocal images of the

reproductive system of CLas+ females treated with dsDcAKHR for 48 h. Scale bar = 200 μm.

DAPI: the cell nuclei were stained with DAPI and visualized in blue. CLas-Cy3: the CLas signal

visualized in red by staining with Cy3. Merge: merged imaging of co-localization of cell nuclei

and CLas. Data are shown as means ± SEM. The significant differences between treatment

and controls are indicated by asterisks (Student's t-test, ***p < 0.001).

miR-34 targeted *DcAKHR*

Based on the small RNA libraries of *D. citri*, we used miRanda and RNAhybrid software to

predict the potential miRNAs targeting *DcAKHR*; three putative miRNAs, namely miR-34,

miR-2 and miR-14 (Figure 4A), were identified. To confirm the binding activity of these

miRNAs with *DcAKHR*, several dual-luciferase reporter assays were conducted. Compared

with the controls, the luciferase activity was markedly reduced when miR-34 agomir was

co-transfected with the recombinant plasmid containing the full 3'UTR sequence of *DcAKHR*,

while miR-2 and miR-14 activities were not significantly changed (Figure 4B). In addition, the

reporter activity was recovered when the binding sites of miR-34 were mutated in the *DcAKHR*

3'-UTR (Figure 4C). To determine whether miR-34 specifically targets *DcAKHR*, the following

experiments were conducted. Firstly. tissue expression patterns showed that the miR-34 was

highly expressed in the head, followed by the fat bodies, ovaries, and midgut (Figure 4D).

Secondly, miR-34 has the opposite expression pattern to *DcAKHR* (Fig. 3A), and transcript

levels decreased over the assessment period with the levels of both being lower in

CLas-positive psyllids than in CLas-negative psyllids (Figure 4E). Thirdly, after feeding with

agomir-34, the relative levels of miR-34 mRNA in the ovaries of CLas-negative and

CLas-positive psyllids increased 1.9- and 2.3-fold, respectively (Figure S5). The mRNA and

protein levels of *DcAKHR* were correspondingly increased or decreased after separately

feeding psyllids with either antagomir-34 or agomir-34 (Figure 4F). Fourthly, an RNA

immunoprecipitation (RIP) assay was performed; *DcAKHR* mRNA was significantly enriched in

the anti-AGO-immunoprecipitated RNAs from ovaries of agomir-34-fed female psyllids relative

to the control samples (Figure 4G). Taken together, these results strongly suggest that

DcAKHR is a direct target of miR-34.

Figure 4. Identification and validation of the target relationship between miR-34 and

DcAKHR. (A) The putative binding sites of miRNAs in the *DcAKHR* 3'-UTR as predicted by

miRanda and RNAhybrid. (B) Dual-luciferase reporter assays using HEK293T cells

co-transfected with miRNA agomir and recombinant pmirGLO vectors containing the predicted

binding sites for miR-2, miR-14 and miR-34 in the CDS of *DcAKHR.* (C) Dual-luciferase

reporter assays using HEK293T cells co-transfected with miR-34 agomir plus recombinant

pmirGLO vectors containing *DcAKHR*-3'UTR or mutated *DcAKHR*-3'UTR. (D) Tissue

expression pattern of miR-34 in *C*Las+ female adults at 7 DAE in the head, ovary, fat body,

and midgut. (E) Comparison of temporal expression patterns of miR-34 in ovaries of *C*Las-

and *C*Las+ females. (F) Effect of miR-34 agomir and antagomir treatments on *Dc*AKHR mRNA

expression and protein level in ovaries of *C*Las- and *C*Las+ psyllids after 48 h. (G) Relative

expression of miR-34 targeted *DcAKHR in vivo* as demonstrated by an RNA

immunoprecipitation assay. Data are shown as mean ± SEM. For B-D, significant differences

among the different treatments are indicated by lowercase letters above the bars (one-way

ANOVA followed by Tukey's Honestly Significant Difference test at $P < 0.05$). The significant

differences between treatment and control are indicated by asterisks in E-G (Student's *t*-test,

**$p < 0.01$, ***$p < 0.001$).

miR34 participates in *D. citri-C*Las mutualism in ovaries

To investigate the roles of miR-34 in the interaction between *D. citri* and *C*Las,

*C*Las-positive and –negative females were fed with agomir-negative control (NC) or with

miR-34 agomir. After feeding with miR-34 agomir, there was a significant accumulation of TAG

and a decrease in glycogen levels in the two populations (Figures 5A-B). The size of lipid

droplets in the fat bodies of *C*Las-positive psyllids were bigger than those of the control (Figure

5C). After treatment with miR-34 agomir, ovarian development of *C*Las-positive psyllids was

reduced (Figure 5D). In addition, when *C*Las-negative and *C*Las-positive psyllids were fed with

miR-34 agomir, the oviposition period of both psyllid types was shortened, preoviposition

period was markedly extended, and fecundity significantly decreased compared with the

relevant control groups (Figures 5E-G); these phenotypes mimic those expressed after

treatment with ds*DcAKHR*. Moreover, the *C*Las signals and relative titer in *C*Las-positive

ovaries were significantly reduced (Figures 5H-I). All the results indicate that miR-34

suppresses *DcAKHR* expression and is involved in the mutualistic interaction.

Figure 5. miR-34 participation in mutualistic interactions between *D. citri* and *C*Las.

(A) Comparison of TAG levels in fat bodies of *C*Las- and *C*Las+ females treated with agomir-34 for 48 h. (B) Comparison of glycogen levels in the fat bodies of *C*Las- and *C*Las+ females treated with agomir-34 for 48 h. (C) Lipid droplets stained with Nile red in fat bodies

dissected from CLas+ females treated with agomir-34 for 48 h. Scale bar = 40 μm. (D) Ovary

phenotypes of CLas+ female treated with agomir-34 for 48 h. Scale bar = 200 μm. o: ovary, s:

spermathecae. (E-G) Comparison of the preoviposition period, oviposition period and the

fecundity between CLas- and CLas+ adults treated with agomir-34. (H) CLas titer in ovaries of

CLas+ females treated with agomir-34 for 48 h. (I) Representative confocal images of CLas in

the reproductive system of CLas+ females treated with agomir-34 for 48 h. Scale bar = 200 μm.

The signals of DAPI and CLas-Cy3 are same as described in Figure 2. Data are shown as

means ± SEM. The significant differences between treatment and controls are indicated by

asterisks (Student's t-test, ***$p < 0.001$).

The JH signaling pathway is regulated by the AKH pathway and is involved in the

increase in fecundity induced by CLas

To evaluate whether the JH signaling pathway is regulated by the AKH signaling pathway,

we assayed the relative JH titers and the expression levels of several JH pathway-related

genes in DcAKH-deficient, CLas-positive females. Compared to control females, the relative

JH titer in the abdomen decreased significantly after dsDcAKH treatment for 48 h (Figure 6A).

After feeding with dsDcAKH for 48 h, the expression levels of the JH receptor gene, DcMet,

and the downstream transcription factor, DcKr-h1, of the JH signaling pathway were

significantly reduced in the fat bodies and ovaries (Figures 6B-C). In addition, the expression

levels of DcVg1-like, DcVgA1-like and DcVgR, three important downstream genes of the JH

signaling pathway related to ovarian development were lower in DcAKH-deficient

CLas-positive females than in controls after dsRNA feeding for 48 h (Figures 6B-C). Similar

results were observed in the DcAKHR-deficient CLas-positive females (Figures 6D-F) as well

as CLas-positive females fed with agomir-34 (Figures 6G-I). These results suggest that JH

signaling pathway is regulated by the AKH-AKHR-miR-34 signaling pathway and is involved in

the increased fecundity of D. citri induced by CLas.

Figure 6. The JH signaling pathway is regulated by AKH signaling pathway and is involved in the increase in fecundity of *D. citri* induced by *C*Las. (A) JH titer in the abdomen of *C*Las+ females treated with ds*DcAKH* for 48 h. (B) Effects of ds*DcAKH* treatment on mRNA level of JH signaling pathway in fat bodies of *C*Las+ females. (C) Effects of ds*DcAKH* treatment on mRNA levels of components of the JH signaling pathway in the ovaries of *C*Las+ females. (D) JH titers in the abdomens of *C*las+ females treated with ds*DcAKHR* for 48 h. (E) Effects of ds*DcAKHR* treatment on mRNA level of JH signaling pathway in fat bodies of *C*Las+ females. (F) Effects of ds*DcAKHR* treatment on mRNA levels of components of the

339   JH signaling pathway in ovaries of *C*Las+ females. (G) JH titer in abdomen of *C*Las+ females

340   treated with agomir-34 for 48 h. (H) Effects of agomir-34 treatment on mRNA levels of

341   components of the JH signaling pathway in fat bodies of *C*Las+ females. (I) Effects of

342   agomir-34 treatment on mRNA levels of components of the JH signaling pathway in the

343   ovaries of *C*Las+ females.

Discussion

Discussion

An increasing number of studies have focused on the effects of vector-virus interactions

on reproduction. For example, rice gall dwarf virus is transmitted vertically by hitchhiking on

insect sperm, thereby promoting long-term viral epidemics and persistence in nature [2].

Berasategui et al. (2022) [3] found a mutualistic relationship between *Chelymorpha alternans*

and *Fusarium oxysporum*, where the pupae were protected in exchange for the dissemination

and propagation of the fungus. Although there is limited research on the mechanisms

underlying vector-bacteria interactions. In *D. citri*-*C*Las interaction, *C*Las operates host

hormone signaling and miRNA to mediate the mutualistic interactio between *D. citri* fecundity

and its replication [35]. Singh and Linksvayer (2020) [41] found that *Wolbachia*-infected

colonies of *Monomorium pharaonis* exhibited increased colony-level growth, accelerated

colony reproduction, and shortened colony life cycles compared to uninfected colonies.

The fat body is the major insect storage organ that produces the majority of

hemolymph-born vitellogenic proteins and critically contributes to insect ovary development

[16]. In this study, after infection with *C*Las, the preoviposition period of *D. citri* was shortened,

oviposition period was prolonged, and fecundity was significantly increased. These results

suggests that there is a mutualistic interaction in *D. citri* ovaries with *C*Las. *C*Las accelerates

ovarian development and prolongs the oviposition period thereby enhancing the fecundity of *D.*

citri; in turn, *D. citri* provides sites and nutrients for *C*Las replication during ovarian

development. *C*Las-positive *D. citri* require more energy to support this increased fecundity.

Hence, the TAG and glycogen levels of both *C*Las-positive and *C*Las-negative were assayed.

After infection with *C*Las, the TAG and glycogen levels significantly increased, and the

average size of lipid droplets in the fat bodies were distinctly larger, indicating that the increase

in lipid mobilization may prepare for an increase in fecundity.

The total amount of lipids present in fat bodies is a balance between lipogenesis and

lipolysis [42], and both processes are essential for lipid homeostasis. Disruption of either

lipogenesis or lipolysis could interfere with nutritional and physiological conditions in fat bodies

and affect vitellogenesis. AKH/AKHR mediated lipolytic systems are essential for lipid

provision and nutrient transfer during reproduction. In *Gryllus bimaculatus* [43,44] and

Caenorhabditis elegans [45], injections of AKH interfered with egg production by inhibiting the

production of vitellogenins and the accumulation of lipid stores. In *Bactrocera dorsalis*, AKHR

knockdown led to decreased lipolytic activity and delayed oocyte maturation [29] and these

authors suggested the decline in fecundity may be due to the inability to utilize the lipid stores

of fat bodies to promote oocyte maturation. In *Glossina morsitans*, the silencing of *AKHR* and

brummer led to the inability to mobilize lipid reserves in fat bodies during pregnancy, slowed

down the development of oocytes, and caused embryogenesis to fail, indicating that female

fertility depends on lipid metabolism regulated by AKHR [24]. These results suggest that lipid

homeostasis appears critical for insect fecundity, as its disruption substantially suppresses

egg production and negatively affects vitellogenesis.

In the study, *DcAKH* and *DcAKHR* played important roles in the increased lipid

metabolism and fecundity of *D. citri* mediated by *C*Las. The cDNA sequences encoding AKH

and AKHR from *D. citri* were cloned and identified, and phylogenetic analysis showed that

DcAKH and *DcAKHR* were orthologous to other hemipteran AKHs and AKHRs. Functional

analysis showed that *DcAKHR* shares properties found with other AKHRs.

DcAKHR-transfected CHO cells could be activated by AKH peptide, while it did not response

to Crz peptide, despite the sequences and gene structures of AKH and Crz being closely

related [21]. *DcAKH* and *DcAKHR* were more highly expressed in the ovaries of *C*Las-positive

psyllids than those of *C*Las-negative individuals. Knockdown of *DcAKH* and *DcAKHR* resulted

in TAG accumulation and a significant decrease in glycogen levels in the fat bodies. Most

importantly, the *C*Las titer in the ovaries of *C*Las-positive psyllids and fecundity significantly

reduced when feeding with *dsDcAKH* and *dsDcAKHR*. Our results support the idea that

reproductive disruption from AKH/AKHR knockdown is due to the inability of female to mobilize

lipid reserves in the fat bodies required for oocyte maturation during vitellogenesis.

To date, there are no reports on the post-transcriptional regulation of AKHR. As important

post-transcriptional regulators, miRNAs generally suppress their target genes by triggering

mRNA degradation or translational repression [32]. There is increasing evidence implicating

miRNAs in the metabolic processes of insects, particularly in relation to reproduction. In

addition to functions of miR-277 described in the Introduction, this miRNA targets ilp7 and ilp8

thereby controlling lipid metabolism and reproduction in *A. aegypti* [40]. miR-8 depletion in

mosquitoes resulted in an increase of its target gene, *secreted wingless-interacting molecule*,

and in lipid accumulation in developing oocytes [46]. Depletion of miR-8 in mosquitoes by

antagomir injection upregulates *secreted wingless-interacting molecule* and decreases the

level of lipid in ovaries to block vitellogenesis post blood meals [47]. A switch from catabolic to

anabolic amino acid metabolism in fat bodies mediated by miR-276 restricts mosquito

investment into oogenesis and benefits the development of *Plasmodium falciparum* [48]. In the

current study, based on *in vitro* and *in vivo* experiments, we found that host miR-34 targets

DcAKHR. miR-34 was observed to have an opposite expression trend to *DcAKHR*, and its

expression in the ovaries of *C*Las-positive psyllids was lower than in those of CLas-negative

individuals. Treatment of miR-34 agomir and antagomir, respectively, resulted in a significant

decrease and increase in *DcAKHR* mRNA and protein levels. In addition, miR-34

overexpression not only led to the accumulation of TAG and the reduction of glycogen levels in

fat bodies, but also decreased fecundity and *C*Las titers in the ovaries similar to the

phenotypes caused by silencing of *DcAKHR*. This indicates that *C*Las inhibited host miR-34 to

enhance *DcAKHR* expression and improve the lipid metabolism and fecundity. This study is

the first to report the manipulation of host miRNAs and target genes by a bacterium to affect

lipid metabolism and reproduction of their vector.

Vitellogenesis is strongly linked with insect JH production [49]. The fat body is an

important tissue that senses and integrates various nutritional and hormonal signals required

for the regulation of vitellogenesis [16]. Moreover, JH is also involved in the nutrient-dependent

regulation of insect reproduction, but the mechanism of action remains obscure. Lu et al.

(2016a, b) [50,51] have demonstrated that nutritional signaling during female reproduction

induces JH biosynthesis that in turn stimulates vitellogenin expression in fat bodies and egg

production in *N. lugens.* In order to evaluate the possible relationship between

AKH/AKHR-mediated lipolysis and JH-dependent vitellogenesis, the expressions of JH

signaling pathway-related genes in fat bodies and ovaries were determined after knockdown of

DcAKH and *DcAKHR*. We found that reducing *DcAKH* or *DcAKHR* by RNAi in *C*Las-positive *D.*

citri not only decreased JH titer but also resulted in a decrease in the expression of *DcMet* and

DcKr-h1 in the fat bodies and ovaries*,* as well as the egg development related genes,

DcVg-1-like, *DcVg-A1-like* and *DcVgR*. Similar results were detected in the agomir-34

treatment. Our results suggest that AKH/AKHR-based lipolysis plays an important regulatory

role in the JH signaling pathway during *D. citri-C*Las interactions.

In conclusion, we proposed the scheme of events following *C*Las infection presented in

Figure 7. Upon infection with *C*Las, *D. citri* exhibits enhanced fecundity compared to

uninfected individuals.The interaction between *C*Las and *D. citri* affecting reproduction is a

win-win strategy; the increased offspring of *D. citri* contributes to a higher presence of *C*Las in

the field. *C*Las upregulates the AKH/AKHR signaling and downregulates miR-34 to increase

lipid metabolism and activate JH-dependent vitellogenesis, thereby improving the fecundity of

*C*Las-positive females. Overall, our results not only broaden our understanding of the

molecular mechanism underlying the *D. citri-C*Las interaction, but also provide valuable

insights for further integrated management of *D. citri* and HLB by targeting the

DcAKH/DcAKHR signaling pathway.

Figure 7. Mechanisms linking metabolism and reproduction of *D. citri* induced by *C*Las.
After infection with *C*Las, the TAG and glycogen levels in fat bodies of *C*Las-positive psyllids
significantly increased as well as the size of lipid droplets. In ovaries, *C*Las upregulates the
AKH/AKHR signaling and downregulates miR-34 to increase lipid metabolism and activate
JH-dependent vitellogenesis, thereby improving the fecundity of *C*Las-positive females. *D. citri*
are more fecund than their uninfected counterparts. The interaction between *C*Las and *D. citri*
affecting reproduction is a win-win strategy; the more offspring of *D. citri*, the more *C*Las in the
field.

Materials and Methods

Host plants and insect colonies

The CLas-negative (healthy) and CLas-positive *D. citri* colonies were obtained from a laboratory culture continuously reared on the healthy and CLas-infected lemon (*Citrus ×limon* (L.) Osbeck) plants. Monthly monitoring of the CLas infection in both the lemon plants and psyllids was conducted using quantitative polymerase chain reaction (qPCR). The two populations were raised in different incubators with same conditions (26 ± 1 ℃, 65 ± 5% relative humidity (RH) and a 14-h light: 10-h dark cycle). CLas-negative and CLas-positive lemon plants were grown in two separate glasshouses.

Bioinformatics and phylogenetic analyses

The physicochemical properties of *DcAKH* (GenBank accession number: MG550150.1) and *DcAKHR* (GenBank accession number: OR259432) were analyzed using the online bioinformatics ProtParam tool (http://web.expasy.org/protparam/). Search and download amino acid sequences of other species in the NCBI database. Homologous protein sequences of AKH and AKHR were aligned with those of other insects using DNAman6.0.3 software. Phylogenetic trees were constructed with the neighbor-joining method in MEGA5.10 software, with 1000 bootstrap replicates; bootstrap values > 50% shown on the tree.

Assays of reproductive parameters

Tested females from two colonies were selected and paired with healthy males. One female and one male were placed onto young vegetative shoots (flush) of healthy lemon plants to promote oviposition; the flush and insects were sealed with tied white mesh bag (15 × 20 cm). These lemon plants were held in an incubator (25 ± 1 ℃, 65 ± 5% RH, 14L: 10D photoperiod). After 24h, the preoviposition period and oviposition period were recorded, the number of eggs laid per female were counted, and this pair of psyllids were transferred to new flush to continue egg laying. Eggs were counted daily until all females died. All the experiments were performed using three replications, and about 15 pairs of *D. citri* for each replication.

Quantitative RT-PCR (qRT-PCR)

For mRNA expression analysis, total RNA was extracted using TRIzol reagent (Invitrogen, Carlsbad, CA, United States), and the synthesis of first-strand cDNA was performed using a PrimeScript™ II 1st Strand cDNA Synthesis Kit (Takara, Beijing, China) in accordance with the manufacturer's instructions. qRT-PCR was carried out by using the TB Green® Premix Ex Taq™ II (Takara, Beijing, China) on an ABI PRISM® 7500 Real-Time System (Applied Biosystems, Foster City, CA, USA). The beta-actin (*Dcβ-ACT*, GenBank XM_026823249.1) gene was used as an internal control to normalize gene expression levels. For miRNA expression analysis, miRNA was extracted using a miRcute miRNA Isolation Kit (TIANGEN, Beijing, China), synthesized using a miRcute Plus miRNA First-Strand cDNA Kit (TIANGEN, Beijing, China), and quantified using the miRcute Plus miRNA qPCR Kit (SYBR Green) (TIANGEN, Beijing, China). U6 snRNA was used as an internal control to normalize the expression of miRNA. All the primers used are listed in Supplementary Table 1.

Luciferase activity assay

The 185 bp sequence of the 3'UTR containing the predicted target sites for miR-34 in *DcAKHR* was cloned into the pmirGLO vector (Promega, Wisconsin, USA) downstream of the luciferase gene to construct the recombinant plasmid, *DcAKHR*-3'UTR-pmirGLO, using the pEASY®-Basic Seamless Cloning and Assembly Kit. The mutated 3'UTR sequence was amplified and cloned into the pmirGLO vector to generate the *DcAKHR*-3'UTR mutant-pmirGLO plasmid. The agomir and antagomir of miRNAs were chemically synthesized and modified by GenePharma (Shanghai, China) with chemically modified RNA oligos of the same sequence or anti-sense oligonucleotides of the miRNA. The negative control for agomir and antagomir was provided by the manufacturer. According to the manufacturer's instructions, the constructed vector (500 ng) and miRNA agomir (275 nM) were co-transferred into HEK293T cells in a 24-well plate using a Calcium Phosphate Cell Transfection Kit (Beyotime, Nanjing, China). After co-transfection for 24 h, the activities of firefly and Renilla luciferase were detected using the Dual-Glo Luciferase Assay System (Promega, Madison, WI, USA), and the average luciferase activity was calculated.

Fluorescence in situ hybridization (FISH)

FISH using a *C*Las probe was carried out as previously described [52]. Under a dissecting microscope, the ovaries of *C*Las-negative and *C*Las-postive *D. citri* were dissected in 1 × phosphate-buffered saline (PBS) (Beyotime Biotechnology, Shanghai, China). Firstly, the isolated ovaries were fixed in Carnoy's fixative (glacial acetic acid-ethanol-chloroform, 1: 3: 6, vol/vol) for 12 h at 25 ℃, washed four times (5 min each time) with 6% H_2O_2 in 80% ethanol and then three times (10-15 min per time) with PBST (1 × PBS: TritonX-100, 99.7: 0.3, (vol/vol)). Secondly, the samples were pre-incubated three times (10 min per time) in hybridization buffer (20 mM Tris-HCl, pH 8.0, 0.9 M NaCl, 30% formamide (vol/vol), 0.01% sodium dodecyl sulfate (wt/wol)) containing 10 pmol/mL of each probe for 24 h at 25 ℃, then washed three times with PBST. Thirdly, nuclei were stained with 0.1 mg/mL of 40, 60-diamidino-2-phenylindole (DAPI) (Sigma-Aldrich, St Louis, MO, USA) for 15 min, then washed once with PBST. Finally, the stained ovaries were mounted in mounting medium. The slides were scanned under a Leica TCS-SP8 (Leica Microsystems Exton, PA USA) confocal microscope; excitation lasers emitting at 405 nm and 550 nm were used to detect the DAPI and Cy3 signals, respectively, and sequential scanning was used to avoid the signal overlap of probes. Image processing was completed using Leica LAS-AF software (v2.6.0). Specificity of detection was carried out using without the probe and *C*Las-negative controls. Three FISH tests were performed and at least 20 ovaries were viewed under the microscope to confirm the repeatability. The probe sequences used in the current study are listed in Supplementary Table 1.

Western blotting

The total proteins from *D. citri* ovaries were extracted using RIPA protein lysis buffer (50 mM Tris pH 7.4, 1% Triton X-100, 150 mM NaCl, 1% sodium deoxycholate and 0.1% SDS) with 1 mM PMSF. Proteins concentrations were quantified using a BCA protein assay kit (Beyotime, Jiangsu, China). Aliquots of 60 μg protein were separated by 12% SDS-PAGE and transferred to polyvinylidene fluoride membranes (Millipore). Subsequently, the membranes were incubated with the primary antibody against *DcAKHR* (1: 1000 dilution, ABclonal Technology Co., Ltd., Wuhan, China) for 12 h at 4 °C and then with the secondary antibody

(goat anti-rabbit IgG conjugated with HRP, 1: 10000 dilution) for 2 h at 25 °C. A mouse

monoclonal antibody against β-actin (TransGen Biotech, Beijing, China) was used as a control.

Immunoreactivity was imaged with the enhanced chemiluminescence with Azure C600

multifunctional molecular imaging system.

Nile red staining

The fat bodies were dissected from adult females 5 DAE, fixed in 4% paraformaldehyde

for 30 min at 25 ℃, and washed twice with 1 × PBS. Lipid droplets were incubated for 30 min

within the mixture of Nile red (0.1 μg/μL, Beijing Coolaber Technology Co., Ltd) and DAPI

(0.05 μg/μL), and then washed twice with 1 × PBS. The samples were imaged using a laser

scanning microscopy (TCS-SP8, Leica Microsystems Exton, PA USA).

Determination of TAG levels

TAG levels were measured using a Triglycerides Colorimetric Assay (Cayman) in

accordance with the manufacturer's instructions. In short, thirty fat bodies were homogenized

in 100 μL of Diluent Assay Reagent. Then 10 μL of supernatant was incubated with Enzyme

Mixture solution, the TAG contents from the measurements were normalized to protein levels

in the supernatant of the samples determined using a BCA protein assay (Thermo Fisher

Scientific). Three independent biological replicates were analyzed for each treatment, and

each treatment included three technical replicates.

Determination of glycogen levels

Glycogen levels were assayed using a Glycogen Assay Kit (Cayman) in accordance with

the manufacturer's instructions. In brief, thirty fat bodies were homogenized in 100 μL of

Diluent Assay Reagent, and 10 μL of supernatant was incubated with Enzyme Mixture solution

and Developer Mixture. The glycogen levels from the measurements were normalized to

protein levels in the supernatant of the sample determined using a BCA protein assay (Thermo

Fisher Scientific). Three independent biological replicates were analyzed for each treatment,

and each treatment included three technical replicates.

dsRNA synthesis and RNAi assay

dsRNAs of *DcAKH* and *DcAKHR* were transcribed by a Transcript Aid T7 High Yield kit (Thermo Scientific, Wilmington, DE, United states) and purified using the GeneJET RNA Purification kit (Thermo Scientific). miRNA agomir, agomir negative control, miRNA antagomir, and antagomir negative control were synthesized in the Shanghai GenePharma Co. Ltd. (Shanghai, China). The RNAi and miRNA treatments were performed by feeding dsRNA or miRNA antagomir/agomir through an artificial diet as described previously [53]. Briefly, twenty females at 7 DAE were placed into a glass cylinder (25 × 75 mm) and sealed with two stretched paramembranes. 200 µL of 20% (w:v) sucrose mixed with dsRNA was placed between two paramembranes for feeding. The final ingestion concentrations of ds*DcAKH* and ds*DcAKHR* were 200 ng/µL and 300 ng/µL, respectively. After feeding with dsRNA for 48 h, the treated females were assayed as follows. The ovaries were dissected or qRT-PCR and western blot analysis. Reproductive parameters including the pre-oviposition period, oviposition period, and fecundity were recorded. *C*Las titers in the ovaries were assayed. Ovary morphologies were observed using a Ultra-Depth Three-Dimensional Microscope (VHX-500). Fat bodies were stained with Nile red. TAG levels and glycogen levels were determined. Green fluorescent protein (GFP) was used as a reporter gene. Feeding ds*GFP* was used as the control when psyllids were fed with dsRNA. All the experiments were performed in three replications, and each replication included 15-20 pairs of *D. citri*.

Heterologous expression and calcium mobilization assay

The complete ORF of *DcAKHR* was cloned into the expression vector pcDNA3.1$^+$; cloning was confirmed by sequencing (TSINGKE Bio). Endotoxin-free plasmid DNA of the vector was extracted, and all tested peptides were synthesized by Sangon Biotech (Shanghai, China) at a purity >95 % (see Fig. 1B for peptide structures). Chinese hamster ovary (CHO-WTA11) cells supplemented with Gα16 subunit and aequorin were used [54] and transfected with pcDNA3.1$^+$-*DcAKHR* or empty pcDNA3.1$^+$ (negative control) vectors using Lipofectamine 2000 (Thermo Fisher Scientifc, Waltham, MA USA). The calcium mobilization assay was performed with slight modification as described by Gui et al. (2017) [55] and Shi et al. (2017) [56]. In brief, the cells were incubated for 3 h in the dark with coelenterazine h (Promega,

Madison, WI, USA) 48 h after the transfection. Then, serial dilutions of peptide ligands (10^{-6} to

10^1 µM) were loaded into opaque 96-well plates. Luminescence was measured over 15 s

using a SpectraMax i3x Multi Mode Microplate Reader (Molecular Devices). Medium alone

was used as a blank control and 100 µM ATP was served as a positive control. Each

experiment was repeated three times.

JH titer detection

JH titers were assessed using the Insect JH Ⅲ ELISA kit (Shanghai Enzyme-linked

Biotechnology Co., Ltd. Shanghai, China) following the manufacturer's protocol.

Statistical analysis

All statistical values are presented as means ± SEM. Statistical analyses were performed

by GraphPad Prism 8.0 software. Student's *t*-test was used to determine the pairwise

comparisons at the following significance levels (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$). For

multiple comparisons, one-way ANOVA with Tukey's Honest Significant Difference tests used

to separate means at $p < 0.05$.

Acknowledgments

This research was supported by the National Natural Science Foundation of China, grant

number 32102193, and the Open Competition Program of Ten Major Directions of Agricultural

Science and Technology Innovation for the 14th Five-Year Plan of Guangdong Province, grant

number 2022SDZG07.

Author Contributions

Conceptualization: Jiayun Li, Paul Holford, George Andrew Charles Beattie, Yurong He, Yijing

Cen, Xiaoge Nian.

Data curation: Jiayun Li, Shujie Wu, Jielan He, Xiaoge Nian.

Formal analysis: Jiayun Li, Xiaoge Nian, Desen Wang.

Funding acquisition: Xiaoge Nian, Yurong He, Yijing Cen.

Investigation: Jiayun Li, Shujie Wu, Jielan He, Shijian Tan, Xiaoge Nian.

Methodology: Jiayun Li, Paul Holford, George Andrew Charles Beattie, Yurong He, Xiaoge

Nian, Yijing Cen.

Project administration: Xiaoge Nian, Yijing Cen.

Resources: Jiayun Li, Shujie Wu, Jielan He, Xiaoge Nian, Yijing Cen.

Software: Jiayun Li, Xiaoge Nian, Desen Wang.

Supervision: Yurong He, Yijing Cen.

Validation: Jiayun Li, Paul Holford, George Andrew Charles Beattie, Xiaoge Nian.

Visualization: Jiayun Li, Paul Holford, George Andrew Charles Beattie, Yurong He, Xiaoge

Nian, Yijing Cen.

Writing-original draft: Jiayun Li, Xiaoge Nian, Yijing Cen.

Writing-review & editing: Jiayun Li, Paul Holford, George Andrew Charles Beattie, Xiaoge Nian,

Yijing Cen.

Competing interests

651 The authors declare no conflict of interest.

Data Availability Statement

654 The published article includes all data generated or analyzed during this study. The full

655 sequences were submitted to NCBI, accession number: MG550150.1 for *DcAKH*, OR259432

656 for *DcAKHR*, OP251123 for *DcMet*, XM_026820026.1 for *DcKr-h1*, XM_008488883.3 for

657 *DcVg-1-like*, XM_026832896.1 for *DcVg-A1-like* and OP251122 for *DcVgR*.

References

[1] Eigenbrode SD, Bosque-Pérez NA, Davis TS. Insect-borne plant pathogens and their

vectors: Ecology, evolution, and complex interactions. Annu Rev Entomol. 2018;

63:169-191.

[2] Mao QZ, Wu Wei, Liao ZF, Li JJ, Jia DS, Zhang XF, et al. Viral pathogens hitchhike with

insect sperm for paternal transmission. Nat Commun. 2019; 10:955.

[3] Berasategui A, Breitenbach N, García-Lozano M, Pons I, Sailer B, Lanz C, et al. The leaf

beetle *Chelymorpha alternans* propagates a plant pathogen in exchange for pupal

protection. Curr Biol. 2022; 32:4114-4127.

[4]   Ray S, Casteel CL. Effector-mediated plant-virus-vector interactions. Plant Cell. 2022;

34:1514-1531.

[5]   Capoor SP, Rao DG, Viswanath SM. *Diaphorina citri* Kuwayama. A vector of the greening

disease of citrus in India. Indian J. Agric. Sci, 1967; 37:572-576.

[6]   Gottwald TR. Current epidemiological understanding of citrus Huanglongbing. Annu Rev

Phytopathol. 2010; 48:119-139.

[7]   Bové JM. Huanglongbing or yellow shoot, a disease of Gondwanan origin: Will it destroy

citrus worldwide? Phytoparasitica. 2014; 42:579-583.

[8]   da Graça JV, Douhan GW, Halbert SE, Keremane ML, Lee RF, Vidalakis G, et al.

Huanglongbing: An overview of a complex pathosystem ravaging the world's citrus. J

Integr Plant Biol. 2016; 58:373-387.

[9]   Fuentes A, Braswell WE, Ruiz-Arce R, Racelis A. Genetic variation and population

structure of *Diaphorina citri* using cytochrome oxidase I sequencing. PLoS ONE. 2018;

13:e0198399.

[10] Beattie GAC. Management of the Asian citrus psyllid in Asia. In: Qureshi JA, Stansly PA

(eds) Asian citrus psyllid: biology, ecology and management of the huanglongbing vector.

CAB International, Wallingford, UK. 2020; 179-209.

[11] Graham J, Gottwald T, Setamou M. Status of huanglongbing (HLB) outbreaks in Florida,

California and Texas. Tropical Plant Pathology. 2020; 45:265-278.

[12] Li S, Wu F, Duan Y, Singerman A, Guan Z. Citrus greening: Management strategies and

their economic impact. HortScience. 2020; 55:604-612.

[13] Pelz-Stelinski KS, Killiny N. Better together: Association with '*Candidatus* Liberibacter

asiaticus' increases the reproductive fitness of its insect vector, *Diaphorina citri*

(Hemiptera: Liviidae). Ann Entomol Soc Am. 2016; 109:371-376.

[14] Ren SL, Li YH, Zhou YT, Xu WM, Cuthbertson AGS, Guo YJ et al. Effects of *Candidatus*

Liberibacter asiaticus on the fitness of the vector *Diaphorina citri*. J Appl Microbiol. 2016;

121:1718-1726.

[15] Wu FN, Qureshi JA, Huang JQ, Fox EGP, Deng XL, Wan FH, et al. Host plant-mediated

interactions between '*Candidatus* Liberibacter asiaticus' and its vector *Diaphorina citri*

Kuwayama (Hemiptera: Liviidae). J Econ Entomol. 2018; 111:2038-2045.

[16] Arrese EL, Soulages JL. Insect fat body: energy, metabolism, and regulation. Annu Rev

Entomol. 2010; 55:207-225.

[17] Stone JV, Mordue W, Batley KE, Morris HR. Structure of locust adipokinetic hormone, a

neurohormone that regulates lipid utilisation during flight. Nature. 1976; 263:207-211.

[18] Toprak U. The role of peptide hormones in insect lipid metabolism. Front Physiol. 2020;

11:434.

[19] Azeez OI, Meintjes R, Chamunorwa JP. Fat body, fat pad and adipose tissues in

invertebrates and vertebrates: the nexus. Lipids Health Dis. 2014; 13:71.

[20] Gáliková M, Diesner M, Klepsatel P, Hehlert P, Xu Y, Bickmeyer I, et al. Energy

homeostasis control in *Drosophila* adipokinetic hormone mutants. Genetics. 2015;

201:665-683.

[21] Park Y, Kim YJ, Adams ME. Identification of G protein-coupled receptors for *Drosophila*

PRXamide peptides, CCAP, corazonin, and AKH supports a theory of ligand-receptor

coevolution. Proc Natl Acad Sci U S A. 2002; 99:11423-11428.

[22] Staubli F, Jorgensen TJ, Cazzamali G, Williamson M, Lenz C, Sondergaard L, et al.

Molecular identification of the insect adipokinetic hormone receptors. Proc Natl Acad Sci

U S A. 2002; 99:3446-3451.

[23] Ziegler R, Isoe J, Moore W, Riehle MA, Wells MA. The putative AKH receptor of the

tobacco hornworm, *Manduca sexta*, and its expression. J Insect Sci. 2011; 11:40.

[24] Attardo GM, Benoit JB, Michalkova V, Yang G, Roller L, Bohova J, et al. Analysis of

lipolysis underlying lactation in the tsetse fly, *Glossina morsitans*. Insect Biochem Mol Biol.

2012; 42:360-370.

[25] Konuma T, Morooka N, Nagasawa H, Nagata S. Knockdown of the adipokinetic hormone

receptor increases feeding frequency in the two-spotted cricket *Gryllus bimaculatus*.

Endocrinology. 2012; 153:3111-3122.

[26] Jedlička P, Ernst UR, Votavová A, Hanus R, Valterová I. Gene expression dynamics in

major endocrine regulatory pathways along the transition from solitary to social life in a

bumblebee, *Bombus terrestris*. Front Physiol. 2016; 7:574.

[27] Nyasembe VO, Hamerly T, López-Gutiérrez B, Leyte-Vidal AM, Coatsworth H, Dinglasan

RR. Adipokinetic hormone signaling in the malaria vector Anopheles gambiae facilitates

Plasmodium falciparum sporogony. Commun Biol. 2023; 6(1):171.

[28] Karbusová N, Gautam UK, Kodrík D. Effect of natural toxins and adipokinetic hormones

on the activity of digestive enzymes in the midgut of the cockroach Periplaneta americana.

Arch Insect Biochem Physiol. 2019;101(4):e21586.

[29] Hou QL, Chen EH, Jiang HB, Wei DD, Gui SH, Wang JJ, et al. Adipokinetic hormone

receptor gene identification and its role in triacylglycerol mobilization and sexual behavior

in the oriental fruit fly (*Bactrocera dorsalis*). Insect Biochem Mol Biol. 2017; 90:1-13.

[30] Lu K, Wang Y, Chen X, Zhang X, Li W, Cheng Y, et al. Adipokinetic Hormone receptor

mediates trehalose homeostasis to promote vitellogenin uptake by oocytes in *Nilaparvata*

lugens. Front Physiol. 2019; 9:1904.

[31] Zheng H, Chen C, Liu C, Song Q, Zhou S. Rhythmic change of adipokinetic hormones

diurnally regulates locust vitellogenesis and egg development. Insect Mol Biol. 2020;

29:283-292.

[32] Bartel DP. Metazoan microRNAs. Cell. 2018; 173:20-51.

[33] Asgari S. Role of microRNAs in insect host-microorganism interactions. Front Physiol.

2011; 2:48.

[34] Asgari S. Role of microRNAs in arbovirus/vector interactions. Viruses. 2014;

6:3514-3534.

[35] Nian X, Luo Y, He X, Wu S, Li J, Wang D, et al. Infection with '*Candidatus* Liberibacter

asiaticus' improves the fecundity of *Diaphorina citri* aiding its proliferation: A win-win

strategy. Mol Ecol. 2024; 33:e17214.

[36] Zhang SD, An SH, Hoover K, Li Z, Li XR, Liu, XM, et al. Host miRNAs are involved in

hormonal regulation of HaSNPV-triggered climbing behaviour in *Helicoverpa armigera*.

Mol Ecol. 2018; 27:459-475.

[37] Bryant B, Macdonald W, Raikhel AS. microRNA miR-275 is indispensable for blood

digestion and egg development in the mosquito *Aedes aegypti*. Proc Natl Acad Sci U S A.

2010; 107:22391-22398.

[38] Zhang Y, Zhao Bo, Roy S, Saha TT, Kokoza VA, Li M, et al. microRNA-309 targets the

Homeobox gene *SIX4* and controls ovarian development in the mosquito *Aedes aegypti*.

Proc Natl Acad Sci U S A. 2016; 113:E4828-4836.

[39] Song JS, Zhou ST. Post-transcriptional regulation of insect metamorphosis and
oogenesis. Cell Mol Life Sci. 2020; 77:1893-1909.

[40] Ling L, Kokoza VA, Zhang C, Aksoy E, Raikhel AS. MicroRNA-277 targets *insulin-like*
pseptide 7 and *8* to control lipid metabolism and reproduction in *Aedes aegypti*
mosquitoes. Proc Natl Acad Sci U S A. 2017; 114:E8017-E8024.

[41] Singh R, Linksvayer TA. *Wolbachia*-infected ant colonies have increased reproductive
investment and an accelerated life cycle. J Exp Biol. 2020; 223:jeb220079.

[42] Grönke S, Mildner A, Fellert S, Tennagels N, Petry S, Müller G, et al. Brummer lipase is
an evolutionary conserved fat storage regulator in *Drosophila*. Cell Metab. 2005;
1:323-330.

[43] Lorenz MW. Adipokinetic hormone inhibits the formation of energy stores and egg
production in the cricket *Gryllus bimaculatus*. Comp Biochem Physiol B Biochem Mol Biol.
2003; 136:197-206.

[44] Lorenz MW, Gäde G. Hormonal regulation of energy metabolism in insects as a driving
force for performance. Integr Comp Biol. 2009; 49:380-392.

[45] Lindemans M, Liu F, Janssen T, Husson SJ, Mertens I, Gäde G, et al. Adipokinetic
hormone signaling through the gonadotropin-releasing hormone receptor modulates
egg-laying in *Caenorhabditis elegans*. Proc Natl Acad Sci U S A. 2009; 106:1642-1647.

[46] Lucas KJ, Roy S, Ha J, Gervaise AL, Kokoza VA, Raikhel AS. MicroRNA-8 targets the
Wingless signaling pathway in the female mosquito fat body to regulate reproductive
processes. Proc Natl Acad Sci U S A. 2015; 112:1440-1445.

[47] Page RE Jr, Rueppell O, Amdam GV. Genetics of reproduction and regulation of
honeybee (*Apis mellifera* L.) social behavior. Annu Rev Genet. 2012; 46:97-119.

[48] Lampe L, Jentzsch M, Kierszniowska S, Levashina EA. Metabolic balancing by miR-276
shapes the mosquito reproductive cycle and *Plasmodium falciparum* development. Nat
Commun. 2019; 10:5634.

[49] Roy S, Saha TT, Zou Z, Raikhel AS. Regulatory pathways controlling female insect
reproduction. Annu Rev Entomol. 2018; 63:489-511.

[50] Lu K, Chen X, Liu WT, Zhang XY, Chen MX, Zhou Q. Nutritional signaling regulates

vitellogenin synthesis and egg development through juvenile hormone in *Nilaparvata*

lugens (Stål). Int J Mol Sci. 2016; 17:269.

[51] Lu K, Chen X, Liu WT, Zhou Q. TOR pathway-mediated juvenile hormone synthesis

regulates nutrient-dependent female reproduction in *Nilaparvata lugens* (Stål). Int J Mol

Sci. 2016; 17:438.

[52] Hosseinzadeh S, Shams-Bakhsh M, Mann M, Fattah-Hosseini S, Bagheri A, Mehrabadi

M, et al. Distribution and variation of bacterial endosymbiont and "*Candidatus*

Liberibacter asiaticus" titer in the Huanglongbing insect vector, *Diaphorina citri*

Kuwayama. Microb Ecol. 2019; 78:206-222.

[53] Yu XD, Killiny N. RNA interference of two glutathione S-transferase genes, *Diaphorina*

citri DcGSTe2 and *DcGSTd1*, increases the susceptibility of Asian citrus psyllid

(Hemiptera: Liviidae) to the pesticides fenpropathrin and thiamethoxam. Pest Manag Sci.

2018; 74:638-647.

[54] Jiang HB, Gui SH, Xu L, Pei YX, Smagghe G, Wang JJ. The short neuropeptide F

modulates olfactory sensitivity of *Bactrocera dorsalis* upon starvation. J Insect Physiol.

2017; 99:78-85.

[55] Gui SH, Jiang HB, Liu XQ, Xu L, Wang JJ. Molecular characterizations of natalisin and its

roles in modulating mating in the oriental fruit fly, *Bactrocera dorsalis* (Hendel). Insect Mol

Biol. 2017; 26:103-112.

[56] Shi Y, Jiang HB, Gui SH, Liu XQ, Pei YX, Xu L, et al. Ecdysis triggering hormone

signaling (ETH/ETHR-A) is required for the larva-larva ecdysis in *Bactrocera dorsalis*

(Diptera: Tephritidae). Front Physiol. 2017; 8:587.

     Supplemental Information

Figure S1. Sequence characterization of AKH from different insect species (data are related to Figure 2). (A) Alignment of the amino acid sequences of the *Dc*AKH transmembrane domain with homologs from other insect species. The sequences in the green box represent the mature peptides (QVNFSPNW). (B) Phylogenetic analysis of *Dc*AKH and its homologs in other insect species. The phylogenetic tree was constructed using neighbor-joining with 1000 bootstrap replicates; values > 50% are shown on the tree. The scale bar indicates the number of amino acid substitutions per site.

Figure S2. Tissue expression of *DcAKH* in *C*Las-positive adult females 9 days after emergence in the head, ovary, fat body, and midgut (data are related to Figure 2).

**Figure S3. *In vivo* and *in vitro* studies validating the DcAKH-DcAKHR interaction (data
are related to Figure 3).** The predicted tertiary structure of the *DcAKHR* protein. The seven
conserved transmembrane ion channels are shown as TM1-TM7. (B) Phylogenetic analysis
the protein sequences of AKHR. The phylogenetic tree was constructed using neighbor-joining
with 1000 bootstrap replicates; values > 50% are shown on the tree. The scale bar indicates
the number of amino acid substitutions per site. (C) Concentration-response curves for the
effect of Ca^{2+} on *DcAKHR*-expression in CHO cells. (D) The expression levels of *DcAKHR* in
*C*Las-negative and *C*Las-positive females treated with ds*DcAKH* for 48 h.

Figure S4. Tissue expression of *DcAKHR* in *C*Las-positive female adults 9 DAE in the head, ovary, fat body, and midgut (the data are related to Figure 3).

Figure S5. The expression levels of miR-34 in *C*Las-negative and *C*Las-positive females treated with agomir-34 for 48 h (the data are related to Figure 4).

841 **Figure S6. (A-E) Melting curve for qRT-PCR primers of *DcAKH*, *DcAKHR*, *Dcβ-ACT*,**

842 **miR-34 and U6.**

843 **Table S1**. **Primer list used for 3'RACE, qRT-PCR, and RNAi analysis. Related to Method details.**

Gene name	Accession number	primer name	Sequences of primers (5'-3')	Product size (bp)	Purpose
DcAKH	MG550150.1	*DcAKH*-Full-F	ATGGAGCGACACCATGTGAG	222	Validation of full length sequence
		DcAKH-Full-R	CTGATTGAATTTTTCACATT		
		DcAKH-qF	TGGACTCTCTCAAGTACATTT	54	qRT-PCR
		DcAKH-qR	ACAATTTTTGGGCTTCACTC		
		DcAKH-RNAi-F	taatacgactcactatagggATGGAGCGACACCATGTGAG	138	dsRNA synthesis
		DcAKH-RNAi-R	taatacgactcactatagggGCGACATTCATCAGGAGGAT		
		DcAKH mature peptide	pQVNFSPNW-NH2	-	mature peptide
		DcCrZ mature peptide	pQTFQYSRGWTN-NH2	-	
DcAKHR	OR259432	*DcAKHR*-Full-F	CACATCACCCGGTATGATT	1452	Validation of full length sequence
		DcAKHR-Full-R	CTCCCCAAGATTCAACCTT		
		DcAKHR-3UTR-F	CTGATACGGCCAATAACGTG	185	3'UTR amplification
		3'RACE-Outer Primer	TACCGTCGTTCCACTAGTGATTT		
		DcAKHR-qF	TCAGTTGGAAGAAGGACAAG	116	qRT-PCR
		DcAKHR-qR	GGTACTCAGGCTACACTCTA		

		DcAKHR-RNAi-F	<u>taatacgactcactataggg</u>CTGAGTGGCCTCTGCCTAAC	399	dsRNA synthesis
		DcAKHR-RNAi-R	<u>taatacgactcactataggg</u>TTAACGGCTTCAAAACTGCC		
		DcAKHR-3UTR-Full-F	CTAGTTGTTTAAACGAGCTC TAAGAGAAGAGACTGGC CGG	166	Full sequence of 3'UTR
		DcAKHR-3UTR-Full-R	TGCATGCCTGCAGGTCGACTCTAGA CACAGGAGTTTC AATCTATT		
		DcAKHR-3UTR-mutant-F	CTAGTTGTTTAAACGAGCTC GCCACATTGCGTTAAACC GT	166	Mutant sequence of 3'UTR
		DcAKHR-3UTR-mutant-R	TGCATGCCTGCAGGTCGACTCTAGA CACAGGAGTTTC AATCTATT		
		DcAKHR-probe	FAM-CACGTTATTGGCCGTATCAG	-	Labled with FAM for FISH
GFP	ACY56286	GFP-RNAi-F	<u>taatacgactcactataggg</u>ACTCCAGCAGGACCATGTGATC	596	dsRNA synthesis
		GFP-RNAi-R	<u>taatacgactcactataggg</u>ACCTGAAGTTCATCTGCACCAC		
Dcβ-ACT	DQ675553.1	β-ACT-qF	TGTTCCAACCTTCCTTCCTG	109	qRT-PCR
		β-ACT-qR	GTGTTGGCGTACAGGTCCTT		
DcMet	OP251123	DcMet-qF	AAGCTCAAGGGCCAAGTCAT	257	qRT-PCR

Gene	Accession	Primer	Sequence (5'-3')	Product (bp)	Application
		DcMet-qR	TAATCAATACCAGGGGCGGC		
DcKr-h1	XM_026820026.1	*DcKr-h1-qF*	CTCCAGTGCTGAGTCCACAA	103	qRT-PCR
		DcKr-h1-qR	ATCTCCCGGAGGTTTCTGTT		
DcVg-1-like	XM_008488883.3	*DcVg-1-like-qF*	CACCTACTCCTTGTCCTCTA	166	qRT-PCR
		DcVg-1-like-qR	GAAAAATCCCCAGAGTCCTT		
DcVg-A1-like	XM_026832896.1	*DcVg-A1-like-qF*	CTCCTCAGAAAGTGGAAGTT	132	qRT-PCR
		DcVg-A1-like-qR	TTGTTTCCGATGAAGTAGGG		
DcVgR	OP251122	*DcVgR-qF*	AGCAGCTGGATATACATGTG	186	qRT-PCR
		DcVgR-qR	CTCCACAGTACTGATTACCG		
*C*Las 16s rRNA	L22532.1	*C*Las 16s-probe	Cy3-CATTATCTTCTCCGGCG	-	Labled with Cy3 for FISH
		*C*Las 16s-qF	TCGAGCGCGTATGCAATACG	-	qRT-PCR
		*C*Las 16s-qR	GCGTTATCCCGTAGAAAAAGGTAG		
miR-34	-	miR34-qF	TGGCAGTGTGGTTAGCTGGTTG		qRT-PCR
		miR-34-probe	Cy3-CACAACCAGCTAACCACACTGCCA	-	Labled with Cy3 for FISH
U6	-	U6-qF	AGGATGACACGCAAAATCGT	-	qRT-PCR

844   Note: The lowercase with underline indicated the T7 promoter sequences in the primers of dsRNA synthesis. The sequences with black boxes displayed the

845   restriction enzyme cutting sites.