1 **A tuneable minimal cell membrane reveals that two lipids suffice for life**

3 **Authors: Isaac Justice[a] and James P Saenz[a, b]***

[a] B CUBE Center for Molecular Bioengineering, Technische Universität Dresden, 01062 Dresden, Germany

[b] Medical Faculty, Technische Universität Dresden, 01062 Dresden, Germany

*Corresponding author and lead contact: james.saenz@tu-dresden.de

Abstract

All cells are encapsulated by a lipid membrane which facilitates the interaction between life and

its environment. How life exploits the diverse mixtures of lipids that dictate membrane property

and function has been experimentally challenging to address. We introduce an approach to tune

and minimize lipidomes in *Mycoplasma mycoides* and the Minimal Cell (JCVI-Syn3A) revealing

that a 2-component lipidome can support life. Systematically reintroducing phospholipid features

demonstrated that acyl chain diversity is more critical for growth than head group diversity. By

tuning lipid chirality, we explored the lipid divide between Archaea and the rest of life, showing

that ancestral lipidomes could have been heterochiral. Our approach offers a tuneable minimal

membrane system to explore the role of lipid complexity, opening new directions in

bioengineering.

Introduction

Cell membranes are complex and responsive systems that serve to protect and mediate

interactions of life with its environment. A large part of this molecular complexity is due to the

diverse panel of lipids that make up the lipidome and which ultimately determine the form and

function of the membrane. The complexity of cellular lipidomes can be staggering, from tens of

unique structures in bacteria[1], to hundreds in eukaryotic organisms[2]. How life has evolved to

utilize such complex mixtures of lipids to build cellular membranes remains an active area of

exploration[3,4]. One approach to studying the role of lipidome complexity is to experimentally

tune lipidome composition and observe effects on cell fitness (e.g. growth). A recent study in

bacteria, for example, revealed the importance of homeoviscous adaptation for electron

transport by tuning acyl chain unsaturation in *Escherichia coli*[5]. However, for interpreting

lipidome flexibility, *E. coli*, like many bacteria, is complicated by the fact that it has multiple

membranes (e.g. inner and outer). Changes in a single membrane are obfuscated by whole cell

lipid extracts, and purifying specific membrane types is laborious, and can lead to substantial

experimental error due to varying purity. Gram positive organisms such as *Bacillus subtilis*, that

have only a single membrane are better models in this regard[6, 7]. However, traditional genetic

approaches to tuning lipidome composition have so far not afforded complete control over both

lipid class composition, and phospholipid acyl chain composition. Thus, a cellular model system

in which lipidome complexity can be reduced in a systematic fashion has not yet been

established.

Mycoplasma are a class of genomically simple bacterial pathogens possessing several

features that are promising as tunable living model membrane systems. Mycoplasmas have a

single plasma membrane, and lack a cell wall[8]. Therefore, their membrane composition and

biophysical properties can be examined in situ without the need for laborious membrane

isolation. Having evolved to a parasitic lifestyle, mycoplasmas have lost many pathways for

biomolecular synthesis, including most of their lipid synthesis pathways, which they take up from

their hosts, or the growth media[8, 9, 10, 11, 12]. This affords the possibility to control their lipidome

composition by controlling what lipids are provided in the media. Further, because of their

relatively small genomes, the number of components involved in managing lipidome composition

and membrane adaptation is within reach of being completely characterized and modeled[13].

Recently, a minimal cell, JCVI-Syn3.0, was engineered from *Mycoplasma mycoides subsp capri*

strain *GM12* by systematically removing genes from *M. mycoides* to achieve an organism where

every remaining gene is essential or quasi-essential[14]. To restore normal division and remove

pleomorphism, 19 genes were added back to JCVI-Syn3.0 to create JCVI-Syn3A; a regularly

dividing quasi minimal cell which offers a platform to study the role of lipids for the most

fundamental requirements of life in a minimal genetic system[15].

In this study we establish an approach to modulate lipidome composition and reduce its

complexity in *M. mycoides* and JCVI-Syn3A. By tuning the lipid composition of the growth medium

and introducing diether phospholipids we introduce an approach to bypass cellular lipid

remodeling, achieving a lipidome with only two lipids. Using these lipidomically minimal living

membranes, we compare the relative importance of phospholipid head group vs. acyl chain

complexity. Additionally, we observed profound effects resulting from subtle changes in the lipid

chirality that distinguishes Archaea from the rest of life. By developing approaches to tune and

minimize mycoplasma lipidome composition, we hope to introduce a new tool for deciphering

the principles of living membranes and to introduce a new paradigm for understanding why life

has evolved to utilize so many lipids.

Results

Mycoplasmas as minimal model membrane systems

Mycoplasma mycoides is a pathogen of mammals that has historically been used as a

simple model membrane system. There are a number of reasons why *M. mycoides* makes a good

model membrane system. First, its small genome (~1,100,000 bp) limits the complexity of its

genetic regulation, and has allowed researchers to fully sequence and annotate the genome

(although a number of genes still have unknown or only putative functional assignments)[8,11].

Second, as a pathogen, the primary source of lipids for *M. mycoides* is through environmental

uptake from the host rather than synthesis[8, 9, 10, 11, 16]. In a laboratory setting using growth

medium as the lipid source, this feature provides the experimenter with direct control over the

lipid components available for synthesizing and maintaining the membrane; by adding or

removing specific lipids from the growth medium of *M. mycoides*, the composition of the

membrane can be altered. Third, *M. mycoides* has a single plasma membrane and no cell wall or

organelles, making it easy to interrogate the membrane and ensuring that membrane targeting

probes in *M. mycoides* are acting on the membrane and not another structure[8]. Fourth, and

perhaps most importantly for this research, *M. mycoides* is unable to synthesize or alter fatty

acid composition[8, 9, 10, 11]. This means that, while it can alter the acyl chain composition of its

membrane lipids, it is limited in its ability to do so by the pool of fatty acids it has access to. These

factors combine to make *M. mycoides* one of the model systems in which membrane remodeling

is both the simplest and most controllable, and as such it is an excellent model system to study

membrane remodeling in living organisms[17].

In *M. mycoides* membranes phospholipids, sterols, and free fatty acids are taken up from

the environment and either incorporated into the membrane or taken up into the cytoplasm (Fig.

1; 1)[8, 18, 19, 20]. *M. mycoides* can cleave acyl chains from exogenous phospholipids (Fig. 1; 2)[21, 22, 23]

When a pool of free fatty acids is present, *M. mycoides* can use those fatty acids to modify the

acyl chain composition of phospholipids taken up from the media, or can synthesize the

phosphotidylglycerol (PG) class of lipids which can in turn be modified and used to synthesize

cardiolipins (Fig. 1; 3, 4, 5)[8, 20, 21, 24, 25, 26, 27]. These newly synthesized or modified lipids can be

broken down to replenish the fatty acid pool, or reinserted into the membrane (Fig. 1; 6)[25]. With

the exception of the modification of PG headgroups to make cardiolipins, all lipid remodeling in

M. mycoides is acyl chain remodeling–that is, it relies on having access to a pool of free fatty acids

to modify the existing acyl chain composition of phospholipids or synthesize *de novo* PGs[8, 9].

Cholesterol is essential for growth of *M. mycoides* (Fig. 1; 8)[28, 29, 30]. For the first time, we report

the feeding of *M. mycoides* on a defined lipid diet which contains no exogenous source of free

fatty acids. When fed such a diet, *M. mycoides* is forced to rely completely on acyl chain

scavenging from exogenous phospholipids as its only source of acyl chains for which to remodel

its membrane lipidome.

Minimizing the lipidome

Our first goal was to determine the minimal lipidome that can support growth of *M.*

mycoides. Sterols, preferably cholesterol, are required for growth, and must be included in any

lipid diet. *M. mycoides* can synthesize several phospholipids (e.g. PG and CL) when provided free

fatty acids[20, 29, 30]. Therefore, to minimize the phospholipid diversity, we removed free fatty acids

from the lipid diet and instead provided a single phospholipid, POPC, along with cholesterol,

yielding a minimal 2-component lipid diet. When transferred from growth medium containing

fetal bovine serum (FBS, a complex undefined lipid source), to the two-component lipid diet, cells

initially grew poorly and tended to aggregate in clumps (not shown). Thin layer chromatography

(TLC) of cell lipid extracts showed that, initially, PG and CL (both internally synthesized lipids)

disappeared, and only traces of SM remained, presumably carry over from growth on FBS (Fig.

2a). After adaptation, sphingomyelin was no longer visible, however, PG and CL reappeared along

with lyso-PC (a PC lipid with one of its acyl chains removed). Since *M. mycoides* cannot synthesize

phospholipids in the absence of fatty acids, these observations indicate that acyl chains were

being scavenged from POPC for the synthesis of PG and CL, presumably as a result of lipase

activity (Fig. 2a). A systematic analysis of PG and CL production on minimal lipid diets composed

of varying phospholipid species demonstrated that *M. mycoides* is capable of scavenging acyl

chains from a broad range of phospholipid head groups (Fig. S1, TableS1), demonstrating a robust

capacity for *M. mycoides* to procure acyl chains for internal PG and CL synthesis from nearly any

phospholipid source.

The capacity for *M. mycoides* to scavenge acyl chains from exogenous phospholipids

provided a hurdle to our goal of minimizing the lipidome. Shotgun mass spectrometry of cells

grown on the minimal 2-component lipid diet revealed 28 lipid structures generated from the

remodeling of POPC, and presence of phospholipids including PG, CL, PA, and DAG (Fig. 2b). Since

the source of acyl chains for internal phospholipid synthesis presumably came from exogenous

POPC, we reasoned that internal lipid synthesis could be eliminated by blocking acyl chain

scavenging. To do this, we replaced POPC with an analogous lipid containing ether-linked 16:0

and 18:1 hydrocarbon chains, which are inert to lipase activity[31, 32]. Following the transfer of cells

from FBS or minimal POPC-cholesterol diet to a minimal diether PC-diet, TLC analysis of lipid

extracts over several passages showed the disappearance of CL and PG, and presence of only two

bands corresponding to cholesterol and diether PC (Fig. 2c). Shotgun lipidomic analysis

demonstrated that cholesterol and diether PC accounted for greater than 99 mol% of the

detected lipids. The remaining very low abundance lipids are derived from impurities in the media

and the diether PC stock. It is unlikely that these trace lipids play a significant role physically or

chemically, however this highlights the value of establishing a defined growth medium for the

continued development of mycoplasmas as minimal model membrane systems[26]. Thus, by

introducing an enzymatically inert phospholipid, diether PC, to the minimal lipid diet, we could

achieve a minimal lipidome composed predominantly of only two lipid structures, cholesterol

and diether PC.

Tuning lipidome composition

Minimizing the lipidome of *M. mycoides* to two lipids resulted in a two-fold decrease in

growth rates. We next asked which components of the lipidome are critical for growth. By feeding

cells with diether PC, we had reduced lipidome composition in several ways. First, we eliminated

the presence of internally synthesized phospholipids such as PG and CL. Therefore, minimizing

the diversity of phospholipid head groups could have contributed to impaired growth. Second,

we eliminated phospholipid hydrocarbon chain diversity by limiting the cell to one configuration

(16:0/18:1). Thus, it is also possible that restricting the diversity of phospholipid hydrocarbon

chain configurations could have impaired growth.

To test whether head group diversity was a factor in determining growth rate we provided

cells with lipid diets containing diether PG and a mixture of diether PG with diether PC. When

lipid extracts of cells grown on diether PG were analyzed by TLC, we observed bands

corresponding to both PG and CL, indicating that CL synthesis can proceed from diether PG (Fig.

3a). When grown on diether PG and PC, TLC analysis showed three bands corresponding to PC,

PG, and CL (Fig. 3b). Therefore, cells grown on diether PG generate lipidomes containing two

phospholipid head groups and when grown on diether PG and PC generate lipidomes with three

head phospholipid head groups. Surprisingly, growth rates for both diether PG and diether PG+PC

diets were lower than for diether PC alone (Fig. 3f). Further, growth of cells on POPC prior to

adaptation and ability to synthesize PG and CL, show only slightly higher growth rates than

diether PC diets, indicating that reduced growth is not solely due to the introduction of diether

phospholipids. Phospholipid headgroup diversity alone, therefore, is not sufficient to rescue

growth.

We next asked whether phospholipid acyl chain diversity could account for impaired

growth. To this end, we grew cells on a diet of cholesterol and two fatty acids (palmitate and

oleate) designated as '2FA'. On this diet, the phospholipidome is composed predominantly of PG

and CL (Fig. 3c) and cells can synthesize phospholipids with a variety of acyl chain configurations

(e.g. 16:0/18:1, 16:0/16:0, etc...). Thus, headgroup diversity is effectively two, comparable to

growth on diether PG, however, diversity of acyl chain configurations is higher. Growth rates on

the 2FA diet were more than twice growth on diether PG, and approached growth rates of cells

adapted to growth on POPC (Fig. 3f). Therefore, for a lipidome with two phospholipid
headgroups, increased diversity of acyl chain configurations rescues growth, demonstrating the
importance of acyl chains for growth.

Finally, we asked whether growth could also be rescued by providing the full suite of ~28
lipids in cells adapted to growth on POPC. It is possible, for example, that growth is impaired by
the disruption of internal phospholipid synthesis or remodeling pathways due to coupling of lipid
synthesis with cellular growth. To test this, we took advantage of the fact that when cells grown
on the minimal lipid diet are switched to a diet containing ester phospholipids (e.g. POPC +
cholesterol) there is a period of adaptation similar to that seen in Fig. 2a. After the first passage,
before adaptation, the cells do no yet undergo significant acyl chain scavenging, and the
membrane composition remains predominantly composed of the two lipids POPC and
cholesterol (Fig. 3d). This delayed adaptation to acyl chain scavenging allows us a brief window
to study simplified membranes composed of lipids *M. mycoides* could normally scavenge acyl
chains from. The simplified membranes only persist until *M. mycoides* is adapted to the new diet
after several passages. We thus transferred cells grown on the minimal diet (Diether PC +
cholesterol) to a lipid diet derived from lipid extracts of cells adapted to growth on POPC (Fig.
3e), and measured growth in the first passage, before cells adapted to scavenge acyl chains from
the dietary phospholipids. Growth rates on this transplanted lipidome diet resulted in a nearly
complete rescue to levels observed in POPC cells after adaptation (Fig. 3f). These results indicate
that internal phospholipid synthesis and remodeling is not required for optimal growth, and
demonstrates a proof-of-principle that functional lipidomes can be engineered and transplanted
to living membranes to support growth.

Minimizing the lipidome of the Minimal Cell

JCVI-Syn3A is a synthetic cell created by the J. Craig Venter Institute (JCVI) by removing
every non-essential gene from *M. mycoides*[14]. To do this, Hutchison et al. conceptually divided
the genome of *M. mycoides* into 8 segments and systematically went through each, removing
genes to see which were essential or non-essential (Fig. 4a). As such, JCVI-Syn3A is an even
simpler (genetically) model system than *M. mycoides*, while still possessing all of the previously

expressed characteristics of *M. mycoides* that make it a valuable model membrane system[33]. In

this study we used a strain that expresses an additional gene for the fluorescent mCherry protein,

to create JCVI-Syn3A-*mCherry*[15, 34]*,* which for simplicity we subsequently refer to as JCVI-Syn3A.

JCVI-Syn3A provides a good comparison organism to *M. mycoides*, and an experimental platform

to examine the role of lipidome composition in supporting the minimal requirements for life. We

therefore asked whether the JCVI-Syn3A lipidome could also be minimized.

Since JCVI-Syn3A has a truncated set of genes compared to *M. mycoides*, we first tested

whether they retained the capacity to scavenge acyl chains from exogenous phospholipids for

the internal synthesis of PG and CL. Unexpectedly we observed that JCVI-Syn3A could still

synthesize PC and CL when fed with a minimal 2-component diet of cholesterol and POPC, as well

as with a range of other phospholipid head groups (Fig. 4b). This suggests that an undiscovered

lipase remains in the JCVI-Syn3A genome. As a more minimal system, JCVI-Syn3A is a more fragile

organism, and not all cells for a given lipid diet survived the growth protocol. To not bias the

growth rates, these cells were removed from the growth rate analysis. An interesting feature was

noticed, however, that a much higher fraction of cells that could not undergo acyl chain

scavenging were non-viable than cells that could (Fig. 4b) further illustrating the decreased

fitness of cells unable to remodel acyl chain composition.

Next, we assayed growth of JCVI-Syn3A on a 2-component lipid diet of cholesterol +

diether PC (16:0/18:1). Growth for JCVI-Syn3A and *M. mycoides* on the Diether POPC +

Cholesterol diet are commensurate, but on the POPC + Cholesterol diet *M. mycoides* has

significantly improved growth (Fig. 4c). Shotgun lipidomic analysis confirmed that JCVI-Syn3A

cells grown on a 2-component diether PC + Cholesterol diet yielded a lipidome with over 99 mol%

composed of only two lipid species (Fig. 4d). As such, this work has developed the simplest known

living membrane in one of the simplest known living organisms, thereby extending the concept

of minimal life from the genome to the lipidome.

Tuning lipid chirality

Having explored the minimal requirements for lipidome complexity in *M. mycoides* and

JCVI-Syn3A, we next sought to leverage these model systems to probe another fundamental

aspect of lipid biology: chirality. Glycerolipids (including phospholipids) have a chiral center in the

glycerol backbone leading to enantiomeric lipids that are mirror images of each other (Fig. 5a).

In bacteria and eukaryotes, glycerolipids are synthesized with acyl chains at the sn-1 and sn-2

positions, and the phosphate head group at the sn-3 position (Fig. 5a: G3P enantiomer).

Conversely, archaea synthesize glycerolipids with the phosphate at the sn-1 position and the acyl

chains at the sn-2 and sn-3 positions (G1P enantiomer)[35, 36]. Known as the 'Lipid Divide', this

difference in stereochemistry between lipids of archaea and the rest of life has long stood as an

unexplained enigma[37, 38, 39, 40]. Did a last common ancestor have membranes with both

enantiomers? What are the consequences of having a racemic mixture of phospholipids in a living

membrane? To date, no naturally occurring organism has been found that has comparable

amounts of both enantiomers in its membrane. Furthermore, experimentally modulating

phospholipid chirality in the lipidome through genetic approaches presents significant

challenges, as shown by the work of Caforio et al[41]. However, the ability of mycoplasma to uptake

exogenous lipids makes them an exceptionally well-suited model for unraveling this elusive

problem in membrane biology.

To establish whether *M. mycoides* and JCVI-Syn3A can grow on a G1P phospholipid

enantiomer, we prepared a lipid diet consisting of cholesterol and enantiomeric POPC (entPOPC),

as well as a racemic mixture of cholesterol and POPC:entPOPC (1:1 mol%). We compared the two

diets containing entPOPC against growth on cholesterol + POPC (Fig. 5b, 5c). Growth rates of both

organisms show that the introduction of entPOPC to the lipid diet results in impaired growth,

with the racemic diet yielding the most pronounced decrease in growth rate. Thus, we

demonstrate that cells are viable when fed enantiomeric phospholipids.

We next asked how the introduction of entPOPC affected the mechanical robustness and

permeability of the membrane. To assay membrane robustness, we measured sensitivity of cells

to hypoosmotic shock, by determining what fraction of cells are lysed following a shock. Since *M.*

mycoides and JCVI-Syn3A lack a cell wall or cytoskeleton, cell lysis during rapid hypoosmotic shock

is indicative of membrane rupture strength. Hypoosmotic sensitivity increased significantly for

cells grown on entPOPC diets, demonstrating that the enantiomeric lipids affect the mechanical

robustness of the membrane and, consequently, the whole cell (Fig. 5b, c). To assay permeability,

we measured the rate of permeation of fluorescein diacetate (FDA), a non-chiral molecule, across

the cell membrane[42, 43]. Membrane permeability, as measured by the permeability coefficient of

FDA increased significantly upon the introduction of enantiomeric lipids, but was the highest for

both *M. mycoides* and JCVI-Syn3A for the racemic lipid diet (Fig. 5b, c). When plotted against

growth rate, there is an apparent correlation of lower growth with higher permeability (Fig 5d,

e), possibly indicating membrane leakiness as one of the factors underlying impaired growth. The

found permeability coefficients were similar to those previously reported in mammalian cells[43].

Increased permeability and reduced membrane robustness could be due either to

changes in the property of the lipid bilayer, or through changes in lipid-protein interactions.

Previous work on phosphatidylcholine enantiomers in model membranes revealed that modest

changes in permeability to calcein occur in membranes composed of both enantiomers, in

particular scalemic (not 1:1) mixtures. However, such small changes would not account for the

large change in permeability we observe. To determine if introducing enantiomeric POPC

affected membrane bilayer robustness or permeability we reconstituted cellular lipid extracts

from *M. mycoides* into liposomes and evaluated C-Laurdan fluorescence (Fig. 6). The C-Laurdan

General Polarization index (GP) reports bilayer hydration, which is closely coupled with

permeability and mechanical robustness of the membrane[44, 45, 46]. Surprisingly, GP values did not

vary significantly across all three POPC diets, indicating that the lipid bilayer itself was not

disrupted by changes in chirality, and implicating an effect on lipid-protein interactions. Indeed,

several studies have demonstrated an effect of varying lipid chirality on lipid-peptide

interactions, and the permeability of membranes to chiral amino acids[47, 48]. Thus, disrupted lipid-

protein interactions are the most likely basis for the observed phenotypes, setting the stage for

future work employing *M. mycoides* and JCVI-Syn3A as model systems to explore the significance

of lipid chirality on lipid-protein interactions in a living membrane.

Discussion

In this study, we introduce *M. mycoides* and the Minimal Cell JCVI-Syn3A as simple model

organisms with tunable lipidomes for studying the role of lipid complexity. By choosing a model

membrane system incapable of synthesizing or modifying fatty acids, and developing a set of

defined lipid diets for that system, we have demonstrated the creation of a platform in which the

lipidome can be tuned in terms of phospholipid head group, acyl chain composition or lipid

chirality. Using this platform, we created the simplest living membrane, and one incapable of

undergoing acyl chain remodeling. These tunable living membranes allowed us to quantitatively

examine the contribution of lipidomic features to the fitness of a minimal living system.

Lipid scavenging

Our observations broadly demonstrate how lipidome composition is crucial even for

relatively simple microorganisms. This is anecdotally illustrated by the capacity we observe for

mycoplasma to produce a complex lipidome from a single exogenous phospholipid, which is an

impressive evolutionary adaptation to their pathogenic lifestyle. Indeed, this acyl chain

scavenging activity, which confounded our initial attempt to simplify the lipidome, suggests a

potential target for treating mycoplasma infection. Although mycoplasma can survive in pure

culture with a reduced lipidome, they may fare much worse in the context of a host immune

system. Furthermore, the ability for the Minimal Cell to grow with a lipidome of two lipids implies

the possibility for further minimization of the genome, through the deletion of pathways involved

in the scavenging of phospholipid acyl chains and the internal synthesis of phospholipids.

Minimal lipidomes

A long-standing challenge in membrane biology has been to understand why life has

evolved such complex lipidomes, and to identify the essential features of lipidomes required for

optimal membrane function and cellular fitness. The simplest lipidome so far reported was from

a Gram-negative bacterium and was composed of 27 lipid species, excluding outer membrane

lipopolysaccharides that were not analyzed. By reducing the lipidome of *M. mycoides* and JCVI-

Syn3A down to 2 lipids, we show that a complex lipidome is not essential for life, but that two

lipids are far from optimal. By systematically reintroducing lipidomic features to cells with a

minimal lipidome, we were able to show that diversity of acyl chain configurations is more crucial

than diversity of phospholipid head groups. This observation is consistent with previous work

showing that reducing the head group diversity of *B. subtilis* had relatively little effect on

growth[49]. This is not to say that having a diversity of head groups is not important. There are well-
characterized examples of head group chemistry influencing bioactivity, for example the
dependence of bacterial peptide secretion machinery on phophatidylethanolamine[50]. We
previously showed how headgroup structure influences the biophysical effect of acyl chain
composition[4]. Thus, having the right phospholipid head group may be as important as having
diverse headgroups. However, acyl chain structure directly influences the properties of the
bilayer and the activity of integral membrane proteins. Indeed, the critical role of phospholipid
acyl chain unsaturation was recently demonstrated for the assembly and function of the nuclear
pore complex in Eukaryotes[51]. Our results demonstrate that the diversity of acyl chain
configurations is one of the minimal parameters required for a functional lipidome.

One factor limiting the minimization of the lipidome to one lipid is that cholesterol (or an
analogue) is required for growth, but also cannot form a bilayer alone. So, it is possible that even
one bilayer forming lipid could support life in a cell that has not evolved to require sterols. It is
also, however, possible that having a membrane reinforcing sterol or sterol analogue is critical
for building a stable cell membrane with only one major phospholipid. A mycoplasma-like
organism such as Mesoplasma that does not require sterols for growth would provide a means
to test this hypothesis[52].

Lipid chirality and the lipid divide

The lipid divide represents another major problem in membrane biology. Homochirality
is a fundamental feature of biomolecular chemistry. Biomolecules exhibit enantioselectivity for
chirally compatible interaction partners. Consequently, life has evolved to rely on homochiral
molecules (e.g. L-amino acids and nucleic acids). An interesting twist is in the divide between the
chirality of phospholipids made by Archaea and the rest of life[35]. The divide raises questions
about whether the last universal common ancestor (LUCA) possessed a heterochiral lipidome, or
whether the divide happened after the divergence of Bacteria and Archaea[53, 54]. Further, current
theories that eukaryotes emerged from an Archaeal ancestor[55], create a conundrum in explaining
why Eukaryotes don't have Archaeal lipid enantiomers, and whether a gradual transition

involving heterochiral lipidomes could have occurred. At the crux of these conundrums is
whether heterochiral lipidomes can support stable membranes and optimal cellular fitness.

We took advantage of the tuneability of *M. mycoides* and JCVI-Syn3A lipidomes to
observe how heterochiral lipidomes influence the membrane and cellular growth. Previous work
in living systems has achieved lipidomes with a mixture of structurally diverse bacterial and
archaeal lipids that differed not only by chirality, but also in a variety of other ways[56]. A unique
feature of this study is that we were able to introduce two enantiomers of a single lipid structure
(POPC) into a living membrane. Our results demonstrate that a heterochiral lipidome results in a
leakier membrane and impaired cellular robustness and growth. In contrast, previous work in *E.*
coli indicated that heterochiral lipidomes did not affect growth[56]. E. coli has an outer membrane
and cell wall that might compensate for lack of stability caused by heterochirality. The lipids in
the E. coli study were not stereoisomers, but rather entirely different lipid structures (ether vs.
ester acyl chain linkages, isoprenoid tails vs. fatty acid tails). It is possible that these structural
differences somehow obscured the destabilizing effects of heterochirality. Ultimately, the basis
for these diverging observations will provide insight into the biological significance of lipid
chirality. Nonetheless, our work shows that in a minimal organism with a single membrane,
heterochiral lipidomes can support growth, but lead to impaired robustness and fitness. This
demonstrates that ancestral heterochiral lipidomes could have supported life.

The changes in membrane robustness and permeability that we observed do not seem to
be explained by changes in lipid order of pure lipid vesicles reconstituted from cell lipid extracts
of homo- and heterochiral lipidomes. This implies that lipid-protein interactions may be affected
by lipid chirality in ways that impair membrane robustness and function. Numerous in vitro
studies have demonstrated the effect of lipid chirality on lipid-protein interactions, consistent
with the possibility that the phenotypes we observed are rooted in perturbed lipid-protein
interactions. Looking forward, *M. mycoides* and JCVI-Syn3A will be excellent model system to
explore the functional consequences of lipid chirality on lipid-protein interactions.

Outlook

Our approach to employ *M. mycoides* and JCVI-Syn3A as minimal living model membrane

systems paves a new path towards unraveling the role of lipidome diversity and complexity. Our

observations reveal that life does not require complex lipidomes. However, minimization comes

with clear trade-offs in cellular fitness. We further demonstrate the capacity of these model

systems to serve as chassis for exploring fundamental questions in membrane biology. The ability

of this system as a testing platform for lipid diets with a variety of features (including class and

acyl chain composition) makes it a useful and simple in vivo model for design-test-build

applications of membrane composition experiments. Furthermore, we demonstrated the ability

to force *M. mycoides* and JCVI-Syn3A to take up and incorporate enantiomeric lipids in their

membranes, the first time this has been shown in eukaryotic or prokaryotic organisms, and an

exciting first step to allow us to probe questions about the lipid divide and the role of chirality in

membrane stability and cell fitness. Overall, we have demonstrated the creation of a simple,

tunable, living, model membrane system that can be used as a novel platform for probing the

design principles of living membranes.

Acknowledgements

The authors wish to thank the Saenz lab; Nataliya Safronova, Lisa Junghans, Ha Ngoc Anh
Nguyen, and Tomasz Czerniak for discussions and feedback through the process, and for help
with experimental designs; the JCVI, and specifically John Glass and Kim Wise, for providing *M.*
mycoides and JCVI-Syn3A and for further feedback, including manuscript comments; Helena
Jambor for assisting with figure design; and Jennifer Katz-Brandoli for assisting with robot runs
and figure construction. We thank Telesis Bio, Inc. for allowing us to use JCVI-syn3A and
Lipotype GmbH for lipidomic analyses. This work was supported by the B CUBE of the TU
Dresden, a German Federal Ministry of Education and Research BMBF grant (to J.S., project
03Z22EN12), and a VW Foundation ''Life ''grant (to J.S., project 93090).

Materials and Methods

Cell Culture:

M. mycoides and JCVI-Syn3A were grown in liquid culture at 37ºC on a modified SP4 growth

medium with Phenol Red[57]. The lipid source was provided either by Fetal Bovine Serum (FBS),

complex natural source of mammalian lipids, or by complexing lipids with delipidated bovine

serum albumin (BSA). Lipids were dissolved in ethanol and added to the medium at 37ºC

immediately before the addition of cells, in the concentrations given in Supplementary Table 2.

M. mycoides was passaged to new growth medium by adding 50 μL of cell culture at OD600 0.4-

0.8 into 7 mL of freshly prepared media in a T25 Flask (Stand., Vent. Cap) or 10mL of media in a

Duran 100mL glass flask. This corresponded to roughly one passage per day. JCVI-Syn3A was

passaged to new medium by adding 200 μL of cell culture at OD600 0.2-0.6 into 7 mL of freshly

prepared media. This corresponded to one passage every two to four days, depending on cell

growth rates. Cells were grown at 37ºC and 40RPM in a KuhnerShakerX incubator.

Growth Rates:

Metabolic Growth Index:

Due to differences in cell size and behavior between *M. mycoides* and JCVI-Syn3A, measuring

growth curves by relying on taking the absorbance at 600nm proved to be an unreliable method

to achieve growth estimates that were robustly comparable between the two organisms. Phenol

red is a pH indicator that is commonly used as a readout for growth in mycoplasma cell culture[58]

as it detects changes in pH as metabolic activity acidifies the culture media. To obtain the rate of

pH change over time cells were grown on the liquid handler system Biomek i7 Automated

Workstation, and the absorbance at 562 nm was recorded hourly to record changes in phenol

red absorbance as cell growth acidifies the media[13, 58, 59]. All absorbances were then plotted with

respect to time, and then fitted to the logistic function $Nt = K (1 + (K-No)/No) e^{-rt}$) where Nt is

population at time t, K is the carrying capacity, N0 is the initial population size, and r is the growth

rate; or, in this case, the rate of pH change. r was determined by calculating the slope of change

at the steepest point of the logarithmic curve, corresponding to mid exponential growth. To

validate the relevance of this method as a readout for cell growth, growth rates for a

representative set of *M. mycoides* diets were calculated using the absorbance of the cells at

600nm and were compared to the rates calculated with the phenol red method. The result was

a strong linear correlation between the two methods (Supplementary figure 5a). Representative

curves of the entire period of growth for *M. mycoides* on one diet and generated by the two

methods is also shown (Supplementary figure 5b).

Cellular growth data was obtained by growing cells on the liquid handler system Biomek i7
Automated Workstation with absorbance readings taken hourly at OD 562 nm to record changes
in phenol red absorbance as cell growth acidifies the media[13, 58, 59]. Growth rates were
determined by calculating the slope of change at the steepest point of the exponential growth
curve, corresponding to mid exponential growth.

Cell growth data measured by optical density at 600nm was obtained through hourly manual
measurements from cultures grown in 30mL in a Duran 100 mL glass flask at 37ºC using a DeNovix
Spectrophotometer (Supplementary figure 6a,b).

Lipid Extraction and Thin Layer Chromatography:
Lipids were extracted using a Bligh-Dyer lipid extraction protocol[60]. When appropriate, lipid
concentration was determined with a phosphate assay. TLC plates with Silica Gel and a 10x
concentrating zone were pre-washed with chloroform to remove debris and other artifacts, and
were then dried at 60ºC for > 30 minutes to ensure complete solvent evaporation. Lipid samples
were then loaded on the plate and were dried at 60ºC for > 10 minutes to again ensure complete
solvent evaporation. The plate was then run in a closed glass chamber using a running solution
of Chloroform:Methanol:Acetic Acid:Water (85:25:5:4), and were subsequently dried again at
60ºC for > 30 minutes. TLC plates were then rinsed with a 3% Copper acetate and 8% phosphoric
acid solution and heated by a handheld heat gun at 280ºC to char lipid spots.

Liposome Preparation: Lipids of a known concentration (either in stock solutions or extracted
from cells and validated with a phosphate assay) were prepared in a 2mL glass by having their
solvents evaporated overnight under a 10-17mbar vacuum. Lipids were then resuspended in
the appropriate volume of Liposome buffer (LiB) (10mM HEPES, 100 mM NaCl) to achieve 100mM
liposome solution and incubated for 30 minutes at 37ºC. Subsequently, liposomes were homogenized
with ten freeze thaw cycles (1 minute in liquid nitrogen followed by 5 minutes at 60C) and 7 extrusion
cycles in a Hamilton syringe setup through a 100nm filter.

Propidium Iodide Osmotic Shock Assay: Harvest 3X 0.4 ODU of cells with OD 0.2-0.4 (1 ODU

= 1 mL of cells at OD 1). Spin down in pre-warmed centrifuge at 5000g, 7minutes, 37ºC, slow

acceleration slow deceleration. Add 1 µL of 1mM Propidium Iodide dye to appropriate wells

on a black or clear bottom 96well plate. Aspirate medium and resuspend each cell pellet in

400 µL of either H2O or 5X diluted mycoplasma wash buffer (MWB) (original MWB: 20mM

HEPES, 200 mM NaCl, 1% W/V Glucose). Set up an extra tube with cells resuspended in 400 µL

H2O to be boiled as a positive control. Boil control tube in 95ºC thermoshaker at 1000 rpm

for 10 minutes. During that time, incubate all other tubes on the 37ºC thermoshaker at

600rpm. Add 100 µL of the cells in the appropriate buffer to each of their three analytical

wells on the 96 well plate. Incubate 30 minutes with shaking at 37ºC in a TECAN platereader

and subsequently measure the fluorescent signal at excitation 539nm emission 619nm.

Fluorescein Diacetate Permeability Assay: Harvest 3X 0.4 ODU of cells with OD 0.2-0.4 (1 ODU

= 1mL of cells at OD 1). Spin down in pre-warmed centrifuge at 5000g, 7 minutes, 37ºC, slow

acceleration slow deceleration. Wash 1x in MWB, resuspend in 400 µL MWB. Prepare a

Fluorescein Standard curve in plate: 0, 0.25 ,0.5, 1.2, 2 µM Fluorescein Add 100 µL

resuspended cells to proper wells (in triplicate), and 100 µL MWB to Fluorescein wells. Add

FDA to cell wells to achieve final FDA concentration of 5 µM. Incubate and measure on TECAN

(37ºC, 150 minutes, excitation 485 emission 525, measurements every 30 minutes). The

permeability coefficient (P) of FDA was calculated using Fick's Law ($Q = P*A*(C_{out} - C_{in})$) where

Q is the flux across the membrane, and is a constant as the slope of fluorescein increase over

time was linear; A is the area of the membrane, calculated using the assumption that the

cross sectional area of POPC + Cholesterol in a bilayer is 45.1 Å^2 [61], the ratio of POPC :

Cholesterol is roughly 1:1, the area of POPC + cholesterol is equivalent to the area of

Enantiomeric POPC + Cholesterol, and the number of POPC molecules can be calculated from

the phosphate assay described below; C_{out} and C_{in} are the concentrations of FDA outside and

inside the cell (respectively); and C_{in} is 0 as FDA is immediately converted to fluorescein upon

entering the cell.

C-Laurdan General Polarization Assay: Incubate liposomes at 37ºc and 1000rpm for 10 minutes on a tabletop thermoshaker. While incubating, remove C-Laurdan from freezer and let warm to temperature on bench. Add 1 μL C-Laurdan (Dilute 1mM stock 4x in EtOH. Add 1uL of diluted stock for final molarity 0.5 μM C-Laurdan) to 500 μL of liposomes. Incubate samples at 37ºc and 1000rpm for 10 minutes on a tabletop thermoshaker.Add 100 μL of each sample to a flat bottom 96 well plate well, repeating three times in order to achieve analytical triplicates. Measure on TECAN (2 channel fluorescence reading with excitation at 385nm and emission at 440 and 490 nm respectively). General polarization (GP) was calculated using the formula: GP = (I440 - I490) / (I440 + I490) where I is the fluorescence emission intensity at the respective wavelength.

Phosphate Assay: To estimate the concentration of phospholipids in a certain amount of cells, the amount of phosphate in the lipid extraction of 1ODU of cells was measured. To measure phosphate amount a modified version of the method of Chen et al. was used[62]. The lipid extraction was added to glass pyrex tubes with both biological and analytical triplicates (3 vials per biological replicate), and the solvent was evaporated by a brief (<5 minute) incubation at 200ºC. 50 μL of water were added to each sample. To prepare a standard curve of phosphate amounts, an ICP Phosphorus standard was diluted in water to give 5, 10, 20, 50 and 100, and 200 nmols of phosphate. 500 μL of 70% perchloric acid was added to each vial and, after brief vortexing, samples were incubated at 200ºC for 120 minutes. Tubes were cooled down in ice water, and 1mL of 10% w/v ascorbic acid followed immediately by 1 mL of 2.5% w/v ammonium heptamolybdate were added (with a brief vortex after adding each reagent). Samples were incubated at 37ºC for 30 minutes, and then absorbance at 820nm was measured on a TECAN plate reader by adding 200 μL of each tube to a clear bottom 96 well plate . A standard curve of absorbance at 820 vs phosphate amount was calculated using the phosphate standard, and lipid amount of each sample was calculated based on the absorbance value, standard curve, and assumption that one phosphate molecule equals one phospholipid.

Lipidomic Analysis: Lipids extracted from cells using the aforementioned Bligh-Dyer protocol were submitted to Lipotype for mass spectrometry based analysis[63]. The general procedure is described in Sampaio et.al, 2011[64]. For the analysis, samples were spiked with internal lipid standard mixture containing: cardiolipin 14:0/14:0/14:0/14:0, ceramide 18:1;2/17:0, diacylglycerol 17:0/17:0, hexosylceramide 18:1;2/12:0, lyso-phosphatidate 17:0, lyso-phosphatidylcholine 12:0, lyso-phosphatidylethanolamine 17:1, lyso-phosphatidylglycerol 17:1, lyso-phosphatidylinositol 17:1, lyso-phosphatidylserine 17:1, phosphatidate 17:0/17:0, phosphatidylcholine 17:0/17:0, diether phosphatidylcholine 16:0/18:1, phosphatidylethanolamine 17:0/17:0, phosphatidylglycerol 17:0/17:0, phosphatidylinositol 16:0/16:0, phosphatidylserine 17:0/17:0, cholesterol ester 20:0, sphingomyelin 18:1;2/12:0;0, triacylglycerol 17:0/17:0/17:0, and cholesterol D6. After extraction, the organic phase was transferred to an infusion plate and dried in a speed vacuum concentrator. It was then resuspended in 7.5 mM ammonium acetate in chloroform/methanol/propanol (1:2:4, V:V:V) and a 33% ethanol solution of methylamine in chloroform/methanol (0.003:5:1; V:V:V). All liquid handling steps were performed using Hamilton Robotics STARlet robotic platform with the Anti Droplet Control feature for organic solvents pipetting. Samples were analyzed by direct infusion on a QExactive mass spectrometer (Thermo Scientific) equipped with a TriVersa NanoMate ion source (Advion Biosciences). Samples were analyzed in both positive and negative ion modes with a resolution of Rm/z=200=280000 for MS and Rm/z=200=17500 for MSMS experiments, in a single acquisition. MSMS was triggered by an inclusion list encompassing corresponding MS mass ranges scanned in 1 Da increments. Both MS and MSMS data were combined to monitor CE, DAG and TAG ions as ammonium adducts; PC, PC O-, as acetate adducts; and CL, PA, PE, PE O-, PG, PI and PS as deprotonated anions. MS only was used to monitor LPA, LPE, LPE O-, LPI and LPS as deprotonated anions; ceramide, hexosylceramide, sphingomyelin, LPC and LPC O- as acetate adducts and cholesterol as an ammonium adduct of an acetylated derivative.

Materials:

552   SP4 (For 1L):

553   PPLO (3.5g) (Becton, Dickinson, and Company product no. 255420)

554   Tryptone (10g) (Sigma product no 70169)

555   Peptone (5g) (Sigma product no 70176)

556   20% Glucose (25mL) (Ross product no. X997.2)

557   20% Yeastolate (10mL) (Becton, Dickinson, and Company product no. 255772)

558   15% Yeast Extract (35mL) (Roth product no. 2904.3)

559   70g/L BSA (85mL) (Sigma product no. A7030) or FBS (170mL) (Biowest Product no.
560   5181H-500)

561   400,000U/mL Penicillin G-sodium salt (2.5mL) (Roth product no. HP48.1)

562   10mg/mL L-Glutamate (5mL) (Roth product no. HN08.2)

563   Sodium Bicarbonate (1.04g) (Honeywell product no. 71630)

564   CMRL (4.9g) (US Biological Lifesciences product no. C5900)

565   Phenol Red (11mg) (Sigma product no. P3532)

567   Lipids:

568   Cholesterol (Avanti product no. 700100)

569   Diether POPC (Diether 1-palmitoyl-2-oleoyl-glycero-3-phosphocholine) (Avanti
570   product no. 999983)

571   Diether DPPC (Diether 1-palmitoyl-2-palmitoyl-glycero-3-phosphocholine) (Avanti
572   product no. 999992)

573   Diether DOPC (Diether 1-oleoyl-2-oleoyl-glycero-3-phosphocholine) (Avanti
574   product no. 999991)

575   Diether POPG (Diether 1-palmitoyl-2-oleoyl-glycero-3-phosphotdylglycerol)
576   (Avanti product no. 999973)

577   POPC (1-palmitoyl-2-oleoyl-glycero-3-phosphocholine) (Avanti product no.
578   850457)

579   DPPC (1-palmitoyl-2-palmitoyl-glycero-3-phosphocholine) (Avanti product no.
580   850355)

581     DOPC (1-oleoyl-2-oleoyl-glycero-3-phosphocholine) (Avanti product no. 850375)

582     POPG (1-palmitoyl-2-oleoyl-glycero-3-phosphotdylglycerol) (Avanti product no.
583 840457)

584     DPPG (1-palmitoyl-2-palmitoyl-glycero-3-phosphoglycerol) (Avanti product no.
585 840455)

586     DOPG (1-oleoyl-2-oleoyl-glycero-3-phosphoglycerol) (Avanti product no. 850475)

587     POPE (1-palmitoyl-2-oleoyl-glycero-3-phosphoethanolamine) (Avanti product no.
588  850757)

590     DPPE (1-palmitoyl-2-palmitoyl-glycero-3-phosphoethanolamine) (Avanti product
591 no. 850705)

592     DOPE (1-oleoyl-2-oleoyl-glycero-3-phosphoethanolamine) (Avanti product no.
593 850725)

594     Oleic Acid (*cis*-9-Octadecenoic acid) (Sigma product no. 01383)

595     Palmitic Acid (1-Pentadecanecarboxylic acid) (Sigma P0500)

596     Enantiomeric POPC (3-palmitoyl-2-oleoyl-glycero-1-phosphocholine) (Avanti
597 product no. 850855)

598     16-0 Cardiolipin (1',3'-bis[1,2-dipalmitoyl-sn-glycero-3-phospho]-glycerol) (Avanti
599 product no. 710333)

601  Assays:

602     FDA (Sigma product no. F7378)

603     Fluorescein (Fluka product no. 28803)

604     Propidium Iodide (Sigma product no. 537059)

605     C-Laurdan (Stratech Scientific Ltd. product no. T0001-SFC-1)

607  Cells:

608     JCVI-Syn3A-*mCherry (*From Telesis Bio)

609     *Mycoplasma mycoides subspecies capri strain GM12 (*From Telesis Bio)

611   Machines:

612   TECAN

613   Biomek i7 Automated Workstation

614   DeNovix DS-11 FX +

615   PicoQuant FluoTime 300 High Performance Fluorescence Lifetime and Steady

616   State Spectrometer

617
618

Citations

1.   van Meer, G., Voelker, D. R. & Feigenson, G. W. Membrane lipids: where they are and
how they behave. Nat. Rev. Mol. Cell Biol. 9, 112–124 (2008).
2.   Membrane lipidome of an epithelial cell line.doi:10.1073/pnas.1019267108.
3.   Klose, C. et al. Flexibility of a Eukaryotic Lipidome – Insights from Yeast Lipidomics. PLOS
ONE 7, e35063 (2012).
4.   Chwastek, G. et al. Principles of Membrane Adaptation Revealed through Environmentally
Induced Bacterial Lipidome Remodeling. Cell Rep. 32, 108165 (2020).
5.   Viscous   control   of   cellular   respiration   by   membrane   lipid   composition.
https://www.science.org/doi/10.1126/science.aat7925 doi:10.1126/science.aat7925.
6.   Willdigg, J. R. & Helmann, J. D. Mini Review: Bacterial Membrane Composition and Its
Modulation in Response to Stress. Front. Mol. Biosci. 8, (2021).
7.   Mascher, T., Margulis, N. G., Wang, T., Ye, R. W. & Helmann, J. D. Cell wall stress
responses in Bacillus subtilis: the regulatory network of the bacitracin stimulon. Mol. Microbiol.
50, 1591–1604 (2003).
8.   Mycoplasma Cell Membranes. vol. 20 (Springer US, 1993).
9.   Kornspan, J. D. & Rottem, S. The Phospholipid Profile of Mycoplasmas. J. Lipids 2012,
1–8 (2012).
10.   McElhaney, R. N. Physical Studies of Lipid Organization and Dynamics in Mycoplasma
Membranes. in Mycoplasma Cell Membranes (eds. Rottem, S. & Kahane, I.) 53–108 (Springer
US, 1993). doi:10.1007/978-1-4615-2924-8_3.
11.   Bittman, R. Mycoplasma Membrane Lipids Chemical Composition and Transbilayer
Distribution. in Mycoplasma Cell Membranes (eds. Rottem, S. & Kahane, I.) 29–52 (Springer US,
1993). doi:10.1007/978-1-4615-2924-8_2.
12.   Razin S, Yogev D, Naot Y. Molecular biology and pathogenicity of mycoplasmas. Microbiol
Mol Biol Rev. 1998 Dec;62(4):1094-156. doi: 10.1128/MMBR.62.4.1094-1156.1998. PMID:
9841667; PMCID: PMC98941.
13.   Gaspari, E. et al. Model-driven design allows growth of Mycoplasma pneumoniae on
serum-free media. Npj Syst. Biol. Appl. 6, 33 (2020).
14.   Hutchison, C. A. et al. Design and synthesis of a minimal bacterial genome. Science 351,
aad6253 (2016).

15.   James F. Pelletier, Lijie Sun, Kim S. Wise, Nacyra Assad-Garcia, Bogumil J. Karas,
Thomas J. Deerinck, Mark H. Ellisman, Andreas Mershin, Neil Gershenfeld, Ray-Yuan Chuang,
John I. Glass, Elizabeth A. Strychalski, Genetic requirements for cell division in a genomically
minimal cell, Cell, Volume 184, Issue 9, 2021, https://doi.org/10.1016/j.cell.2021.03.008.
16.   Rottem S, Yashouv J, Ne'eman Z, Razin S. Cholesterol in mycoplasma membranes.
Composition, ultrastructure and biological properties of membranes from Mycoplasma mycoides
var. capri cells adapted to grow with low cholesterol concentrations. Biochim Biophys Acta. 1973
Nov 16;323(4):495-508. doi: 10.1016/0005-2736(73)90158-2. PMID: 4357439.
17.   Archer, D. B. Y. 1975. Modification of the Membrane Composition of *Mycoplasma*
mycoides subsp. capri by the Growth Medium. Microbiology 88, 329–338.
18.   Dahl J. Uptake of fatty acids by Mycoplasma capricolum. J Bacteriol. 1988
May;170(5):2022-6.   doi: 10.1128/jb.170.5.2022-2026.1988.   PMID: 2834319;   PMCID:
PMC211080.
19.   Razin, S., Efrati, H., Kutner, S. & Rottem, S. Cholesterol and phospholipid uptake by
mycoplasmas. Rev. Infect. Dis. 4 Suppl, S85-92 (1982).
20.   Plackett, P. The Synthesis of Polar Lipids by Mycoplasma. Ann. N. Y. Acad. Sci. 143, 158–
164 (1967).
21.   Salman M, Rottem S. The cell membrane of Mycoplasma penetrans: lipid composition
and phospholipase A1 activity. Biochim Biophys Acta. 1995 May 4;1235(2):369-77. doi:
10.1016/0005-2736(95)80026-c. PMID: 7756347.
22.   Bhandari, S., Asnani, P.J. Characterization of phospholipase A2 of mycoplasma
species. Folia Microbiol 34, 294–301 (1989). https://doi.org/10.1007/BF02814471
23.   Rottem S, Adar L, Gross Z, Ne'eman Z, Davis PJ. Incorporation and modification of
exogenous phosphatidylcholines by mycoplasmas. J Bacteriol. 1986 Jul;167(1):299-304. doi:
10.1128/jb.167.1.299-304.1986. PMID: 3087959; PMCID: PMC212875.
24.   Rottem S. Membrane lipids of mycoplasmas. Biochim Biophys Acta. 1980 May
27;604(1):65-90. doi: 10.1016/0005-2736(80)90585-4. PMID: 6992872
25.   Rottem S, Markowitz O. Membrane lipids of Mycoplasma gallisepticum: a disaturated
phosphatidylcholine and a phosphatidylglycerol with an unusual positional distribution of fatty
acids. Biochemistry. 1979 Jul 10;18(14):2930-5. doi: 10.1021/bi00581a002. PMID: 465448.
26.   Thornburg ZR, Bianchi DM, Brier TA, Gilbert BR, Earnest TM, Melo MCR, Safronova N,
Sáenz JP, Cook AT, Wise KS, Hutchison CA 3rd, Smith HO, Glass JI, Luthey-Schulten Z.
Fundamental behaviors emerge from simulations of a living minimal cell. Cell. 2022 Jan
20;185(2):345-360.e28.   doi: 10.1016/j.cell.2021.12.025.   PMID: 35063075;   PMCID:
PMC9985924.
27.   Gross Z, Rottem S. Lipid interconversions in aging Mycoplasma capricolum cultures. J
Bacteriol. 1986 Sep;167(3):986-91. doi: 10.1128/jb.167.3.986-991.1986. PMID: 3745126;
PMCID: PMC215969.
28.   Shmuel Razin, Sterols in Mycoplasma Membranes, Current Topics in Membranes and
Transport, Academic Press, Volume 17, 1982, Pages 183-205,https://doi.org/10.1016/S0070-
2161(08)60311-1.
29.   Dahl, J. The Role of Cholesterol in Mycoplasma Membranes. in Mycoplasma Cell
Membranes (eds. Rottem, S. & Kahane, I.) 167–188 (Springer US, 1993). doi:10.1007/978-1-
4615-2924-8_5.

30.   Razin, S., Efrati, H., Kutner, S. & Rottem, S. Cholesterol and phospholipid uptake by
mycoplasmas. Rev. Infect. Dis. 4 Suppl, S85-92 (1982).

31.   Rangholia, N., Leisner, T. M. & Holly, S. P. Bioactive Ether Lipids: Primordial Modulators
of Cellular Signaling. Metabolites 11, 41 (2021).

32.   Choquet, C. G., Patel, G. B., Beveridge, T. J. & Sprott, G. D. Stability of pressure-extruded
liposomes made from archaeobacterial ether lipids. Appl. Microbiol. Biotechnol. 42, 375–384
(1994).

33.   Breuer, M. et al. Essential metabolism for a minimal cell. eLife 8, e36842 (2019).

34.   Sakai A, Deich CR, Nelissen FHT, Jonker AJ, Bittencourt DMC, Kempes CP, Wise KS,
Heus HA, Huck WTS, Adamala KP, Glass JI. Traditional protocols and optimization methods lead
to absent expression in a mycoplasma cell-free gene expression platform. Synth Biol (Oxf). 2022
May 21;7(1):ysac008. doi: 10.1093/synbio/ysac008. PMID: 35774105; PMCID: PMC9239315.

35.   van Wolferen, M., Pulschen, A. A., Baum, B., Gribaldo, S. & Albers, S.-V. The cell biology
of archaea. Nat. Microbiol. 7, 1744–1755 (2022).

36.   Altamura, E. et al. Racemic Phospholipids for Origin of Life Studies. Symmetry 12, 1108
(2020).

37.   Villanueva, L. et al. Bridging the membrane lipid divide: bacteria of the FCB group
superphylum have the potential to synthesize archaeal ether lipids. ISME J. 15, 168–182 (2021).

38.   Martin, H. S., Podolsky, K. A. & Devaraj, N. K. Probing the Role of Chirality in Phospholipid
Membranes. ChemBioChem 22, 3148–3157 (2021).

39.   Sojo, V. Why the Lipid Divide? Membrane Proteins as Drivers of the Split between the
Lipids of the Three Domains of Life. BioEssays 41, 1800251 (2019).

40.   Villanueva, L. Engineering E. coli to Have a Hybrid Archaeal/Bacterial Membrane. Trends
Microbiol. 26, 559–560 (2018).

41.   Caforio, A. et al. Converting Escherichia coli into an archaebacterium with a hybrid
heterochiral membrane. Proc. Natl. Acad. Sci. 115, 3704–3709 (2018).

42.   Rizk, S. et al. Functional diversity of isoprenoid lipids in Methylobacterium extorquens
PA1. Mol. Microbiol. 116, 1064–1078 (2021).

43.   Levental, K. R. et al. Lipidomic and biophysical homeostasis of mammalian membranes
counteracts dietary lipid perturbations to maintain cellular fitness. Nat. Commun. 11, 1339 (2020).

44.   Parasassi, T., De Stasio, G., Ravagnan, G., Rusch, R. M. & Gratton, E. Quantitation of
lipid phases in phospholipid vesicles by the generalized polarization of Laurdan fluorescence.
Biophys. J. 60, 179–189 (1991).

45.   Ma, Y., Benda, A., Kwiatek, J., Owen, D. M. & Gaus, K. Time-Resolved Laurdan
Fluorescence Reveals Insights into Membrane Viscosity and Hydration Levels. Biophys. J. 115,
1498–1508 (2018).

46.   Steinkühler, J., Sezgin, E., Urbančič, I. et al. Mechanical properties of plasma membrane
vesicles correlate with lipid order, viscosity and cell density. Commun Biol 2, 337 (2019).

47.   Lei Chen, Wen Si, Liang Zhang, Gangfeng Tang, Zhan-Ting Li, and Jun-Li Hou
Journal of the American Chemical Society 2013 135 (6), 2152-2155

48.   Takaaki Ishigami, Keishi Suga, and Hiroshi Umakoshi ACS Applied Materials & Interfaces
2015 7 (38), 21065-21072

49.   Willdigg JR, Helmann JD. Mini Review: Bacterial Membrane Composition and Its
Modulation in Response to Stress. Front Mol Biosci. 2021 May 11;8:634438.

50.   van Klompenburg W, Paetzel M, de Jong JM, Dalbey RE, Demel RA, von Heijne G, de
Kruijff B. Phosphatidylethanolamine mediates insertion of the catalytic domain of leader peptidase
in membranes. FEBS Lett. 1998 Jul 10;431(1):75-9.
51.   Romanauska, A., Köhler, A. Lipid saturation controls nuclear envelope function. Nat Cell
Biol (2023). https://doi.org/10.1038/s41556-023-01207-8
52.   Matteau D, Lachance JC, Grenier F, Gauthier S, Daubenspeck JM, Dybvig K, Garneau D,
Knight TF, Jacques PÉ, Rodrigue S. Integrative characterization of the near-minimal bacterium
Mesoplasma florum. Mol Syst Biol. 2020 Dec;16(12):e9844. doi: 10.15252/msb.20209844. PMID:
33331123; PMCID: PMC7745072.
53.   Koga, Y. Early Evolution of Membrane Lipids: How did the Lipid Divide Occur?. J Mol Evol
72, 274–282 (2011).
54.   Rie Matsumi, Haruyuki Atomi, Arnold J.M. Driessen, John van der Oost, Isoprenoid
biosynthesis in Archaea – Biochemical and evolutionary implications, Research in Microbiology,
Volume 162, Issue 1, 2011, Pages 39-52, ISSN 0923-2508
55.   Koonin, E.V. Archaeal ancestors of eukaryotes: not so elusive any more. BMC Biol 13, 84
(2015).
56.   Caforio A, Siliakus MF, Exterkate M, Jain S, Jumde VR, Andringa RLH, Kengen SWM,
Minnaard AJ, Driessen AJM, van der Oost J. Converting Escherichia coli into an archaebacterium
with a hybrid heterochiral membrane. Proc Natl Acad Sci U S A. 2018 Apr 3;115(14):3704-3709.
57.   Fritz, B. A., Thomas, C. B., Van Ess, P. & Yuill, T. M. Comparison of a Modified Edward-
Type Medium and a Modified SP4-Type Medium for Primary Isolation of Mycoplasma
gallisepticum (MG) from Chickens Vaccinated with the F Strain of MG. Avian Dis. 35, 591–598
(1991).
58.   Low, I. E. & Eaton, M. D. Replication of Mycoplasma pneumoniae in Broth Culture. J.
Bacteriol. 89, 725–728 (1965).
59.   MEUR, S. K., SIKDAR, A., SRIVASTAVA, N. C. & SRIVASTAVA, S. K. Rapid photometric
assay of growth of *Mycoplasma mycoides* subsp. capri. J. Appl. Bacteriol. 66, 301–302 (1989).
60.   Bligh, E. G. & Dyer, W. J. A rapid method of total lipid extraction and purification. Can. J.
Biochem. Physiol. 37, 911–917 (1959).
61.   Leftin A, Molugu TR, Job C, Beyer K, Brown MF. Area per lipid and cholesterol interactions
in membranes from separated local-field (13)C NMR spectroscopy. Biophys J. 2014 Nov
18;107(10):2274-86. doi: 10.1016/j.bpj.2014.07.044. PMID: 25418296; PMCID: PMC4241443.
62.   Chen, P. S., Toriuaka, T. Y. and Warner, H. (1956). Microdetermination of phosphorus.
Anal. Chew., 28. 1756-1758.
63.   Ejsing CS, Sampaio JL, Surendranath V, Duchoslav E, Ekroos K, Klemm RW, Simons K,
Shevchenko A. Global analysis of the yeast lipidome by quantitative shotgun mass spectrometry.
Proc Natl Acad Sci U S A. 2009 Feb 17;106(7):2136-41. doi: 10.1073/pnas.0811700106. Epub
2009 Jan 27. PMID: 19174513; PMCID: PMC2650121.
64.   Sampaio JL, Gerl MJ, Klose C, Ejsing CS, Beug H, Simons K, Shevchenko A. Membrane
lipidome of an epithelial cell line. Proc Natl Acad Sci U S A. 2011 Feb 1;108(5):1903-7. doi:
10.1073/pnas.1019267108. Epub 2011 Jan 18. PMID: 21245337; PMCID: PMC3033259.

Figure 1: Membrane Remodeling in _M. Mycoides_ is Dependent on Acyl Chain Scavenging. 1 _Mycoplasma mycoides_ can take up phospholipids and free fatty acids from its environment.[8, 18, 19, 20] **2** Cleaving ester bonds from glycerophospholipids results in a pool of free fatty acids.[21, 22, 23] **3** With fatty acids, _M. mycoides_ is able to synthesize Phosphatidylglycerols (PG) with various fatty acid compositions, as well as remodel other phospholipid classes.[8, 20, 21, 24, 25] **4** Cardiolipins can be synthesized from two PGs, starting with cleavage of PG headgroups.[26, 27] **5** _M. mycoides_ can synthesize a diversity of Cardiolipins from PGs.[8, 24, 26, 27] **6** Cleaving acyl chains from remodeled and synthesized phospholipids replenishes the pool of free fatty acids.[25] **7** Glycerophospholipids are inserted into the membrane.[23] **8** Cholesterol is essential for _M. mycoides_ growth.[28, 29, 30]

Legend:

PC headgroup ○
PG headgroup ◇
CL headgroup ⬦⬦
Saturated fatty acid
Unsaturated fatty acid
PC phospholipid
PG phospholipid
CL phospholipid
Lyso-PC phospholipid
Cholesterol

2a Adaptation of cells to a defined diet

2b Minimal diet: POPC + Cholesterol

2c Minimal diet: D.PC + Cholesterol

Figure 2: A Defined Lipid Diet Results in a Living Membrane with Two Lipids.
a *M. mycoides* can adapt to defined lipid "diets" with diester or diether phospholipids; resulting in simpler membranes than when grown on a complex diet (e.g. FBS). Adaptation to new diets occours after 3 passages (p > 3). TLC aspect ratio modified for legibility; unmodified TLCs can be found in Figure S4. **b** *M. mycoides* can scavenge acyl chains from POPC, yielding a more complex membrane from a defined diet. Lipidomic analysis shows acyl chain scavenging leads to the synthesis of a diversity of lipids, resulting in a membrane with 28 lipids from a diet of only two. **c** 16:0-18:1 diether PC has ether-linked hydrocarbon chains that cannot be cleaved by *M. mycoides*, eliminating scavenging. Lipidomic analysis shows living cells with two lipids comprising 99.33 mol% of their lipidome.

3a Defined diet: D.PG + Cholesterol

3b Defined diet: D.PC + D.PG + Cholesterol

3c Defined diet: 2 Fatty Acids + Cholesterol

3d Defined diet: POPC + Cholesterol diet after first passage

3e Defined diet: Membrane Transplant

3f *M. mycoides* Growth Rates on Defined Diets

Figure 3: Living Membranes with Tunable Lipid Class and Acyl Chain Complexity. a 16:0-18:1 diether PG diets results in the synthesis of Cardiolipin with fully ether-linked hydrocarbon chains, increasing head group, but not acyl chain complexity. **b** A diet of both Diether PC and PG restores the headgroup, but not acyl chain, complexity of the POPC diet. **c** A diet of free fatty acids restores the acyl chain, but not head group, complexity of the POPC diet. **d** When switched from one diet to another, in this case from the D. PC diet to the POPC diet, there is an adaptation period before acyl chain scavenging occurs. **e** A diet derived from a total lipid extract of cells grown on the POPC diet (a "transplant" diet) restores the full complexity of the POPC. Growth on this transplanted diet was performed after the first passage (p = 1) to the new diet, before cells begin to scavenge acyl chains from exogenous phospholipids. **f** Reintroducing acyl chain complexity is more effective than headgroup complexity for rescuing growth (relative to cells adapted to a POPC diet), but, even when full complexity is restored, acyl chain scavenging capacity still improves growth.

4a

Mycoplasma mycoides subsp. capri strain GM12 genome:
1,084,586 bp, 910 genes

Segment 1: 140,739 bp
2: 125,388 bp
3: 133,208 bp
4: 131,623 bp
5: 101,708 bp
6: 189,357 bp
7: 124,976 bp
8: 137,887 bp

JCVI-Syn3A genome:
543,379 bp, 591 genes

Segment 1: 75,732 bp
2: 49,888 bp
3: 73,958 bp
4: 82,531 bp
5: 56,501 bp
6: 83,060 bp
7: 54,482 bp
8: 66,717 bp

4b

Condition	Phenol Red Growth Rate	Scavenging
POPG D.PC	0.0749	Yes
POPE D.PC	0.0457	No
POPC Ent POPC	0.0057	Yes
POPC	0.0523	Yes
OPPC	0.0382	Yes
Ent POPC	0.0246	Yes
DPPG+D.PC	0.0178	No
DPPE+D.PC	0.0307	No
DOPG	0.0016	Yes
DOPE+D.PC	0.0002	Yes
DOPC+DPPC+D.PC	0.0257	Yes
D.PG	0.0009	No
D.PC	0.0327	No
18:1-D.PC	0.0011	No
18:1+POPC	0.0479	Yes
18:1+DPPG	0.0321	Yes
18:1+DPPE	0.0557	Yes
18:1+DPPC+D.PC	0.0696	Yes
18:1+DOPC	0.0001	No
18:1+16.0-CL	0.0772	Yes
18:1+16:0-D.PC	0.0742	Yes
16:0-D.PC+18:1-D.PC	0.0003	No
16:0-D.PC	0.0001	No
16:0-CL	0.0001	No
16:0+POPC	0.0001	Yes
16:0+DPPC	0.0001	No
16:0+DOPG	0.0589	Yes
16:0+18:1-D.PC	0.0002	No
16:0+18:1	0.0852	Yes
16:0+DOPC+D.PC	0.0015	No
16:0	0.0019	No

■ Number of Growing Replicates
▨ Total Number of Replicates

4c

N.S. *** ***

scavenging
no scavenging

4d MEMBRANE

Diether PC
Cholesterol

Figure 4:JCVI-Syn3A is a Simpler Cell with a More Fragile Membrane.
a JCVI-Syn3A is minimal synthetic cell analog of *M. mycoides.* Adapted from Hutchison et. al, 2016. **b** Some JCVI- Syn3A replicates exhibit no growth on a number of lipid diets, especially where acyl chain scavenging is restricted.

c JCVI-Syn3A grows significantly slower than *M. mycoides* on the same diets.
d JCVI-Syn3A is viable grown on a Diether PC + Cholesterol diet with 99.16 mol% of the membrane comprised of only two lipids.

Figure 5: Enantiomeric Lipid Diets Negatively Affect Cell Growth and Membrane Properties. a Enantiomeric (Ent) POPC is a synthetic chiral POPC with the head group in the SN1 position, rather than SN3 position. **b** *M. mycoides* and **c** JCVI-Syn3A cells exhibit lower growth when grown on diets with enantiomeric POPC present; are more fragile to hypoosmotic shock when grown on diets with enantiomeric POPC present; and are more permeable to non-chiral fluorescein diacetate (FDA) when grown on diets with enantiomeric POPC present. **d** *M. mycoides* and **e** JCVI-Syn3A cell growth and membrane permeability are inversely correlated.

Figure 6: Enantiomeric Lipids Have no Significant Effect on Membrane Order.
Lipid vesicles reconstituted from total lipid extracts of *M. mycoides* cells had a similar membrane order (measured as the general polarization of the membrane using C-laurdan) for all three diets with POPC enantiomers.