A tuneable minimal cell membrane reveals that two lipid species suffice for life

Authors: Isaac Justice[a], Petra Kiesel[b], Nataliya Safronova[a], Alexander von Appen[b], and James P.

Saenz[a, c]*

[a] Technische Universität Dresden, B CUBE Center for Molecular Bioengineering, 01307 Dresden,

Germany

[b]Max Planck Institute of Molecular Cell Biology and Genetics, Pfotenhauerstrasse 107, 01307

Dresden

[c]Technische Universität Dresden, Faculty of Medicine, Dresden 01307, Germany

*corresponding author and lead contact: james.saenz@tu-dresden.de

Abstract

All cells are encapsulated by a lipid membrane which facilitates the interaction between

life and its environment. How life exploits the diverse mixtures of lipids that dictate membrane

property and function has been experimentally challenging to address. We introduce an

approach to tune and minimize lipidomes in *Mycoplasma mycoides* and the Minimal Cell (JCVI-

Syn3A) revealing that a 2-component lipidome can support life. Systematically reintroducing

phospholipid features demonstrated that acyl chain diversity is more critical for growth than

head group diversity. By tuning lipid chirality, we explored the lipid divide between Archaea and

the rest of life, showing that ancestral lipidomes could have been heterochiral. Our approach

offers a tunable minimal membrane system to explore the fundamental lipidomic requirements

for life, thereby extending the concept of minimal life from the genome to the lipidome.

Introduction

Cell membranes are complex and responsive systems that serve to protect and mediate interactions of life with its environment. A large part of this molecular complexity is due to the diverse panel of lipids that make up the lipidome and which ultimately determine the form and function of the membrane. The complexity of cellular lipidomes can be staggering, from tens of unique structures in bacteria[1], to hundreds in eukaryotic organisms[2]. How life has evolved to utilize such complex mixtures of lipids to build cellular membranes remains an active area of exploration[1,3–6]. While synthetic membranes can be constructed using a single lipid species, the minimal number of lipid species required for a functional cell membrane remains undetermined. Identifying a minimal viable lipidome would provide a critical starting point for elucidating the combinations of lipid structures that are essential for membrane integrity and function, offering insights into the fundamental chemical and physical requirements of cellular life.

One approach to studying lipidome complexity is to experimentally manipulate lipidome composition and observe effects on cell fitness (e.g. growth). Bacterial model organisms have proven to be excellent systems for tuning lipidome composition by genetically disrupting lipid biosynthesis[7]. For example, early work with *Escherichia coli* mutants deficient in fatty acid synthesis explored the role of acyl chain unsaturation for cell growth[8], and a similar approach with *Bacillus subtilis* mutants explored the role of branched acyl chains[9]. More recent work, employing approaches to tune lipid unsaturation revealed the importance of homeoviscous adaptation for electron transport[10] and demonstrated how low membrane fluidity can induce phase separation, impaired membrane potential, cell growth and division[11]. However, for interpreting how lipidome composition and flexibility are affected by perturbed lipid synthesis, *E. coli*, like many bacteria, is complicated by the fact that it has multiple membranes (e.g. inner and outer). Changes in a single membrane are obfuscated by whole cell lipid extracts, and purifying specific membrane types is laborious, and can lead to substantial experimental error due to varying purity. Gram positive organisms such as *B. subtilis*, that have only a single membrane are better models in this regard[12,13]. However, traditional genetic approaches to tuning lipidome composition have so far not afforded complete control over both lipid class

composition, and phospholipid acyl chain composition. Thus, a cellular model system in which

lipidome complexity can be reduced in a systematic fashion has not yet been established.

Mycoplasmas are a genus of genomically simple bacterial pathogens possessing several

features that are promising as tunable living model membrane systems. *Mycoplasmas* have a

single plasma membrane, and lack a cell wall[14]. Therefore, their membrane composition and

biophysical properties can be examined *in situ* without the need for laborious membrane

isolation. Having evolved to a parasitic lifestyle, *Mycoplasmas* have lost many pathways for

biomolecular synthesis, including most of their lipid synthesis pathways, , relying instead on lipids

acquired from their hosts or from the growth media[14–18]. This affords the possibility to control

their lipidome composition by controlling what lipids are provided in the media. Further, because

of their relatively small genomes, the number of components involved in managing lipidome

composition and membrane adaptation is within reach of being completely characterized and

modeled[19]. Recently, a minimal cell, JCVI-Syn3.0, was engineered from *Mycoplasma mycoides*

subsp. capri strain GM12 by systematically removing genes from *M. mycoides* to achieve an

organism where every remaining gene is essential or quasi-essential[20]. However, this organism

exhibited pleomorphism and irregular cell division[21]. To restore normal cell division, a new strain,

JCVI-Syn3A was created with 19 additional genes not present in JCVI-Syn3.0. The addition of

these genes resulted in a regularly dividing quasi minimal cell which offers a platform to study

the role of lipids for the most fundamental requirements of life in a genomically minimized

system.

In this study, we establish an approach to modulate lipidome composition and reduce its

complexity in *M. mycoides* and JCVI-Syn3A. By tuning the lipid composition of the growth medium

and introducing diether phospholipids we introduce an approach to bypass cellular lipid

remodeling, achieving a lipidome with only two lipids. Using these lipidomically minimal living

membranes, we compare the relative importance of phospholipid head group vs. acyl chain

complexity. Additionally, we observed profound effects resulting from subtle changes in the lipid

chirality that distinguishes Archaea from the rest of life. By developing approaches to tune and

minimize mycoplasma lipidome composition, we hope to introduce a new tool for deciphering

the principles of living membranes and to introduce a new paradigm for understanding why life

has evolved to utilize so many lipids.

Results

Mycoplasmas as minimal model membrane systems

Mycoplasma mycoides is a pathogen of mammals that has historically been used as a

simple model membrane system. There are a number of reasons why *M. mycoides* makes a good

model membrane system. First, its small genome (~1,100,000 bp) limits the complexity of its

genetic regulation, and has allowed researchers to fully sequence and annotate the genome

(although a number of genes still have unknown or only putative functional assignments)[22].

Second, as a pathogen, the primary source of lipids for *M. mycoides* is through environmental

uptake from the host rather than synthesis[14–16,23]. In a laboratory setting using growth medium

as the lipid source, this feature provides the experimenter with direct control over the lipid

components available for synthesizing and maintaining the membrane; by adding or removing

specific lipids from the growth medium of *M. mycoides*, the composition of the membrane can

be altered. Third, *M. mycoides* has a single plasma membrane and no cell wall or organelles,

making it easy to interrogate the membrane and ensuring that membrane targeting probes in *M.*

mycoides are acting on the membrane and not another structure[14]. Fourth, and perhaps most

importantly for this research, *M. mycoides* is unable to synthesize or alter fatty acid

composition[14–17]. This means that, while it can alter the acyl chain composition of its membrane

lipids, it is limited in its ability to do so by the pool of fatty acids it has access to. These factors

combine to make *M. mycoides* one of the model systems in which membrane remodeling is both

the simplest and most controllable, and as such it is an excellent model system to study

membrane remodeling in living organisms[24].

In *M. mycoides* membranes phospholipids, sterols, and free fatty acids are taken up from

the environment and either incorporated into the membrane or taken up into the cytoplasm (Fig.

1; 1)[14,25–27]. *M. mycoides* can cleave acyl chains from exogenous phospholipids (Fig. 1; 2)[28–30].

When a pool of free fatty acids is present, *M. mycoides* can use those fatty acids to modify the

acyl chain composition of phospholipids taken up from the media, or can synthesize the

phosphotidylglycerol (PG) class of lipids which can in turn be modified and used to synthesize

cardiolipin (CL) (Fig. 1; 3, 4, 5)[14,27,28,31–34]. These newly synthesized or modified lipids can be

broken down to replenish the fatty acid pool, or reinserted into the membrane (Fig. 1; 6, 7)[32].

With the exception of the modification of PG headgroups to make cardiolipin, all lipid remodeling

in *M. mycoides* is acyl chain remodeling–that is, it relies on having access to a pool of free fatty

acids to modify the existing acyl chain composition of phospholipids or synthesize *de novo*

PGs[14,15]. Cholesterol is essential for growth of *M. mycoides* (Fig. 1; 8)[26,35,36]. For the first time, we

report the feeding of *M. mycoides* on a defined lipid diet which contains no exogenous source of

free fatty acids. When fed such a diet, *M. mycoides* is forced to rely completely on acyl chain

scavenging from exogenous phospholipids as its only source of acyl chains for which to remodel

its membrane lipidome.

The diverse lipid structures that *M. mycoides* can take up or synthesize ultimately

determine the physical properties of their cell membranes[37,38]. Phospholipid acyl chains can vary

in terms of their length and degree of unsaturation, both influencing physical parameters such

as membrane fluidity, thickness, and permeability. Phospholipid headgroups such as PG and

cardiolipin introduce a negative surface charge to the membrane, which can influence their

interaction with peripheral membrane proteins. Conversely, PC is zwitterionic and introduces a

neutral charge to the membrane surface. The geometric shape of phospholipids is also important,

and determines whether a lipid spontaneously aggregates to form a bilayer or non-bilayer

structure. For example, cardiolipin has four acyl chains and a relatively small headgroup giving it

a conical profile. Cells have been shown to tune the abundance of such conical lipids to modulate

the curvature and bending rigidity of their membranes[39]. Sterols, which also do not form bilayers

by themselves, play an important biophysical role in the membrane, including in modulating

membrane fluidity, stability, facilitating liquid-liquid phase separation, and membrane

asymmetry[40,41]. By limiting the diversity of lipids that can be taken up or synthesized, we aimed

to identify a minimal viable lipidome that can be used as an experimental platform in which lipid

diversity can be systematically tuned in a living membrane.

Minimizing the lipidome

Our first goal was to determine the minimal lipidome that can support growth of *M.*
mycoides. Sterols, preferably cholesterol, are required for growth, and must be included in any
lipid diet. *M. mycoides* can synthesize several phospholipids (e.g. PG and cardiolipin; Full lipid
names and abbreviations can be found in Table 1) when provided free fatty acids[14,27]. For
example, when grown on a lipid diet consisting of cholesterol, and two fatty acids (palmitate -
C16:0 and oleate - C18:1), the lipidome of *M. mycoides* contains primarily cholesterol, PG,
cardiolipin and small amounts of diacylglycerol (DAG) and phosphatidic acid (PA), both precursors
of PG synthesis (Fig 2b, Supplementary Table S3). Therefore, to minimize the phospholipid
diversity, we removed free fatty acids from the lipid diet and instead provided a single
phospholipid, 16:0/18:1 phosphatidylcholine (POPC), along with cholesterol, yielding a minimal
2-component lipid diet. When transferred from growth medium containing fetal bovine serum
(FBS, a complex undefined lipid source), to the two-component lipid diet, cells initially grew
poorly and tended to aggregate in clumps (not shown). Thin layer chromatography (TLC) of cell
lipid extracts showed that, initially, PG and cardiolipin (both internally synthesized lipids)
disappeared, and only traces of sphingomyelin (SM) remained, presumably carry over from
growth on FBS (Fig. 2a; full TLCs are shown in Supplementary Fig. S2). After adaptation (> 3
passages in batch culture), SM was no longer visible, however, phosphatidylglycerol (PG) and
cardiolipin reappeared along with lyso-PC (a PC lipid with one of its acyl chains removed, which
appears below the SM band). Since *M. mycoides* cannot synthesize phospholipids in the absence
of fatty acids, these observations indicate that acyl chains were being scavenged from POPC for
the synthesis of PG and cardiolipin, presumably as a result of lipase activity (Fig. 2a). A systematic
analysis of PG and cardiolipin production on minimal lipid diets composed of varying
phospholipid species demonstrated that *M. mycoides* is capable of scavenging acyl chains from a
broad range of phospholipid headgroups (Fig. S1, Table S2), demonstrating a robust capacity for
M. mycoides to procure acyl chains for internal PG and cardiolipin synthesis from nearly any
phospholipid source.

The capacity for *M. mycoides* to scavenge acyl chains from exogenous phospholipids

provided a hurdle to our goal of minimizing the lipidome. Shotgun mass spectrometry of cells

grown on the minimal 2-component lipid diet revealed 31 lipid species generated from the

remodeling of POPC (with 18 lipid species comprising 99% of the lipidome), and presence of

phospholipids including PG, cardiolipin, with small amounts of the PG precursors PA and DAG

(Fig. 2c, Table S3). Since the source of acyl chains for internal phospholipid synthesis presumably

came from exogenous POPC, we reasoned that internal lipid synthesis could be eliminated by

blocking acyl chain scavenging. To do this, we replaced POPC with an analogous lipid containing

ether-linked 16:0 and 18:1 hydrocarbon chains, which are inert to lipase activity[42,43]. Following

the transfer of cells from FBS or minimal POPC-cholesterol diet to a minimal Diether PC-diet, TLC

analysis of lipid extracts over several passages showed the disappearance of cardiolipin and PG,

and presence of only two bands corresponding to cholesterol and Diether PC (D.PC; Fig. 2a).

Shotgun lipidomic analysis demonstrated that cholesterol and D.PC accounted for 99.9 mol% of

the detected lipids (Fig. 2d) (the remaining very low abundance lipids are derived from impurities

in the media, in particular from yeast extract, Table S3). Thus, by introducing an enzymatically

inert phospholipid, D.PC, to the minimal lipid diet, we could achieve a minimal lipidome

composed predominantly of only two lipid species, cholesterol and D.PC.

Tuning lipidome composition

Minimizing the lipidome of *M. mycoides* to two lipids resulted in a two-fold decrease in

growth rates (Figure. 3e, D.PC condition compared to POPC condition). We next asked which

components of the lipidome most effect growth. By feeding cells with D.PC, we had reduced

lipidome composition in several ways. First, we eliminated the presence of internally synthesized

phospholipids such as PG and cardiolipin. Therefore, minimizing the diversity of phospholipid

headgroups could have contributed to impaired growth. Second, we eliminated phospholipid

hydrocarbon chain diversity by limiting the cell to one configuration (16:0/18:1). Thus, it is also

possible that restricting the diversity of phospholipid hydrocarbon chain configurations could

have impaired growth.

To investigate the importance of head group diversity and structure on determining
growth rate we provided cells with lipid diets containing Diether PG (D.PG) and a mixture of D.PG
with D.PC. When lipid extracts of cells grown on D.PG were analyzed by TLC, we observed bands
corresponding to both PG and cardiolipin, indicating that cardiolipin synthesis can proceed from
D.PG (Fig. 3a). When grown on D.PG and D.PC, TLC analysis showed three bands corresponding
to PC, PG, and cardiolipin (Fig. 3b). Thus, cells grown on D.PC generate lipidomes with one
phospholipid headgroup, those on D.PG produce lipidomes with two headgroups, and cells
grown on both D.PG and D.PC yield lipidomes with three headgroups. Specifically, D.PC cells have
only a neutral, bilayer-forming phospholipid, while D.PG cells contain negatively charged
phospholipids, including both bilayer-forming (PG) and non-bilayer-forming (cardiolipin) types.
In D.PC + D.PG cells, both neutral and negatively charged phospholipids, as well as bilayer- and
non-bilayer-forming types, are present. Surprisingly, despite this increased lipidome complexity,
the introduction of negatively charged (PG) or conical non-bilayer-forming (cardiolipin)
phospholipids did not improve growth rates compared to D.PC cells (Fig. 3e; full growth curves
are shown in Supplementary Fig. S4a). To put these values in context, the growth rates we
estimate from all of the defined diets considered in this study are more than 10-fold lower than
for cells grown on a complex FBS lipid diet, highlighting the effect of reducing lipid diet complexity
on growth[44]. Indeed, growth rates for both D.PG and D.PG + D.PC diets were slightly lower than
for D.PC alone (Fig. 3e). It is possible that this result is due to the fact that the synthesis of
cardiolipin from D.PG results in a single cardiolipin species with hydrocarbon chains of fixed
length and saturation, meaning that the *M. mycoides* cannot remodel cardiolipin to optimize acyl
chain structure[6]. Further, growth of cells on POPC prior to adaptation and ability to synthesize
PG and cardiolipin, show only slightly higher growth rates than D.PC diets, indicating that reduced
growth is not solely due to the introduction of diether phospholipids. Phospholipid headgroup
diversity alone, therefore, is not sufficient to rescue growth.

To understand the impact of phospholipid acyl chain diversity on cell growth, we grew
cells on a diet of cholesterol and two fatty acids (palmitate and oleate), designated as '2FA'. On
this diet, the phospholipidome is predominantly composed of PG and cardiolipin (Fig. 2b),
allowing for the synthesis of phospholipids with various acyl chain configurations (e.g., 16:0/18:1

POPG, 16:0/16:0 DPPG, 18:1/18:1 DOPG for PG, and different permutations of PGs as substrates

for cardiolipin synthesis). Although DAG and PA are present in low abundances (1.8 mol%, and

0.3 mol%, respectively, Supplemental Table S3), they are also both conical-shaped lipids, similar

to cardiolipin. In that respect, their contribution to the membrane's physical properties is

relatively small and does not introduce significant differences compared to what is already

provided by the far more abundant cardiolipin. Therefore, while the 2FA diet exhibits comparable

headgroup diversity to cells grown on D.PG, they exhibit greater acyl chain diversity. Growth rates

on the 2FA diet were more than double those on D.PG and approached the growth rates of cells

adapted to POPC (Fig. 3e). This suggests that for a lipidome with predominately two phospholipid

headgroups, increased acyl chain diversity can rescue growth. An important consideration in

comparing cells grown on the 2FA diet versus the D.PC + D.PG diet is that phospholipids with

diether-linked acyl chains are not natural for this organism and could introduce a growth deficit.

However, the similar growth rates observed in cells grown on POPC prior to adaptation, which

could not yet synthesize PG and cardiolipin, compared to D.PC diets, suggest that reduced growth

is not primarily due to the introduction of diether phospholipids. Acknowledging that differences

in natural ester linkages and unnatural ether linkages, and the presence of DAG and PA—albeit

in low abundance—may also influence the observed growth rates, these observations suggest

that acyl chain complexity is an important factor in rescuing growth.

Finally, we asked whether growth could also be rescued by providing the full suite of 31 lipids in

cells adapted to growth on POPC. It is possible, for example, that growth is impaired by the

disruption of internal phospholipid synthesis or remodeling pathways due to coupling of lipid

synthesis with cellular growth. To test this, we took advantage of the fact that when cells grown

on the minimal lipid diet are switched to a diet containing ester phospholipids (e.g. POPC +

cholesterol) there is a period of adaptation similar to that seen in Fig. 2a. After the first passage,

before adaptation, the cells do not yet undergo detectable acyl chain scavenging, and the

membrane composition remains predominantly composed of the two lipids POPC and

cholesterol (Fig. 3d). This delayed adaptation to acyl chain scavenging allows us a brief window

to study simplified membranes composed of lipids *M. mycoides* could normally scavenge acyl

chains from. The simplified membranes only persist until *M. mycoides* is adapted to the new diet

after several passages. We thus transferred cells grown on the minimal diet (D.PC + cholesterol)

to a lipid diet derived from lipid extracts of cells adapted to growth on POPC (Fig. 3e), and

measured growth in the first passage, before cells adapted to scavenge acyl chains from the

dietary phospholipids. Growth rates on this transplanted lipidome diet resulted in a nearly

complete rescue to levels observed in POPC cells after adaptation (Fig. 3e). These results indicate

that internal phospholipid synthesis and remodeling are not required for optimal growth, and

demonstrates a proof-of-principle that functional lipidomes can be engineered and transplanted

to living membranes to support growth.

Minimizing the lipidome of the Minimal Cell

JCVI-Syn3A is a synthetic cell created by the J. Craig Venter Institute (JCVI) by removing

every non-essential gene from *M. Mycoides*[20]. To do this, Hutchison et al. conceptually divided

the genome of *M. mycoides* into 8 segments and systematically went through each, removing

genes to see which were essential or non-essential. As such, JCVI-Syn3A is genomically an even

simpler model system than *M. mycoides*, while still possessing all of the previously described

characteristics of *M. mycoides* that make it a valuable model membrane system[45]. In this study,

we used a strain that expresses an additional gene for the fluorescent mCherry protein, to create

JCVI-Syn3A-*mCherry*[21,46], which for simplicity we subsequently refer to as JCVI-Syn3A. JCVI-Syn3A

provides a good comparison organism to *M. mycoides*, and an experimental platform to examine

the role of lipidome composition in supporting the minimal requirements for life. We therefore

asked whether the JCVI-Syn3A lipidome could also be minimized.

Since JCVI-Syn3A has a truncated set of genes compared to *M. mycoides*, we first tested

whether they retained the capacity to scavenge acyl chains from exogenous phospholipids for

the internal synthesis of PG and cardiolipin. Unexpectedly we observed that, like *M. mycoides*,

JCVI-Syn3A could still synthesize PG and cardiolipin when fed with a minimal 2-component diet

of cholesterol and POPC, as well as with a range of other phospholipid head groups

(Supplementary Table S2). This suggests that an undiscovered lipase remains in the JCVI-Syn3A

genome. The presence of such lipase activity in a genomically minimal cell could mean that it is

essential. Alternatively, it could also indicate that, while not essential itself, it is a secondary

"moonlighting" activity of an enzyme with an essential activity[47]. This implies that with the

appropriate complement of lipids, the genome could be further minimized by eliminating genes

involved in lipid remodeling activities.

Next, we assayed growth of JCVI-Syn3A on a 2-component lipid diet of cholesterol + D.PC.

Growth for JCVI-Syn3A and *M. mycoides* on the D.PC + Cholesterol diet are commensurate, but

on the POPC + Cholesterol diet *M. mycoides* has significantly improved growth (Fig. 4a). By

comparison, cells grown on FBS, a complex lipid diet, exhibit ~ 10-fold higher growth rates[44].

Although growth is exceptionally slow, the culture can be continuously passaged in batch, and

samples taken 24 hours post-inoculation consistently yield colony-forming units, confirming the

viability of the cells (Supplementary Fig S5). Shotgun lipidomic analysis confirmed that JCVI-Syn3A

cells grown on a 2-component D.PC + cholesterol diet yielded a lipidome with over 99 mol%

composed of only two lipid species (Fig. 4b, Table S3). As such, this work has developed the

simplest known living membrane in one of the simplest known living organisms.

Previous studies reported that genome minimization in JCVI-Syn3.0 caused pleomorphic

traits and abnormalities in cell division, which were rescued by reintroducing 19 genes, resulting

in the creation of JCVI-Syn3A[21]. Given that a reduced lipidome could also impair cellular functions,

we investigated whether lipidome minimization might lead to abnormal cell morphologies in

JCVI-Syn3A. Transmission electron microscopy (TEM) images of JCVI-Syn3A grown on three

different lipid diets—fetal bovine serum (FBS), POPC + cholesterol, and D.PC + cholesterol—

revealed mostly typical ovoid cells (Fig. 4c, TEM overview images provided in Supp. Fig. 6a-g).

However, two distinct morphological features were observed in subpopulations of cells: internal

membrane-encapsulated vesicles and tube-like membranous structures connecting cells.

The tubules observed in fewer than 20% of cells, were less frequent in cells grown on

POPC or D.PC compared to those grown on FBS. These structures are reminiscent of those seen

in wall-less L-form bacteria, where they are hypothesized to function as an FtsZ-independent

mechanism, possibly representing a primitive form of cell division[48]. Further investigation is

required to determine if the tubules observed in JCVI-Syn3A are functional or represent

incomplete cell division. Notably, their higher prevalence in FBS-grown cells suggests that they

are not directly related to lipidome minimization.

In contrast, the frequency of cells with membrane invaginations increased more than two-
fold from ~15% in FBS-grown cells to nearly 40% in those grown on D.PC. In some instances, these
invaginations appeared as distinct membrane-encapsulated vesicles within the cell, separate
from the cell surface membrane. To confirm the presence of these internalized membrane
vesicles, we used cryogenic electron microscopy (cryoEM) to construct whole-cell tomograms.
Tomograms of JCVI-Syn3A from all three diets confirmed the presence of internal membrane-
encapsulated vesicles (Fig. 4d, Supp. Fig. S6h-i). The lower electron density within these vesicles
(pixel brightness), compared to the cytoplasm, indicates they result from membrane
invagination, encapsulating extracellular fluid.

Interestingly, cells with reduced lipidomes were larger on average compared to those
grown on FBS (~0.75 um vs. 0.3 um diameter respectively), as estimated semi-quantitatively from
TEM images (Supplementary Fig. S6j). Cells with internal vesicles were particularly enlarged
across all conditions (up to 1.5 um average diameter). These results suggest that lipidome
minimization leads to larger cell sizes and a higher frequency of membrane invaginations,
indicative of impaired regulation of cell size and shape. The increased frequency of membrane
invaginations could result from non-optimal membrane bending rigidity or intrinsic curvature
from the loss of cardiolipin, and acyl chain diversity. Nonetheless, the fact that around half of the
observed cells maintained normal morphology shows that even with just two lipid species, JCVI-
Syn3A is capable of preserving typical cell morphology.

Tuning lipid chirality

Having explored the minimal requirements for lipidome complexity in *M. mycoides* and
JCVI-Syn3A, we next sought to leverage these model systems to probe another fundamental
aspect of lipid biology: chirality. Glycerolipids (including phospholipids) have a chiral center in the
glycerol backbone leading to enantiomeric lipids that are mirror images of each other (Fig. 5a).
In bacteria and eukaryotes, glycerolipids are synthesized with acyl chains at the sn-1 and sn-2
positions, and the phosphate head group at the sn-3 position (Fig. 5a: G3P enantiomer).
Conversely, archaea synthesize glycerolipids with the phosphate at the sn-1 position and the acyl
chains at the sn-2 and sn-3 positions (G1P enantiomer)[49,50]. Known as the ′Lipid Divide′, this

difference in stereochemistry between lipids of archaea and the rest of life has long stood as an

unexplained enigma[51–54]. Did a last common ancestor have membranes with both enantiomers?

What are the consequences of having a racemic mixture of phospholipids in a living membrane?

To date, no naturally occurring organism has been found that has comparable amounts of both

enantiomers in its membrane. Furthermore, experimentally modulating phospholipid chirality in

the lipidome through genetic approaches presents significant challenges, as shown by the work

of Caforio et al[54]. However, the ability of mycoplasma to uptake exogenous lipids makes them an

exceptionally well-suited model for unraveling this elusive problem in membrane biology.

To establish whether *M. mycoides* and JCVI-Syn3A can grow on a G1P phospholipid

enantiomer, we prepared a lipid diet consisting of cholesterol and enantiomeric POPC (entPOPC),

as well as a racemic mixture of cholesterol and POPC:entPOPC (1:1 mol%). We compared the two

diets containing entPOPC against growth on cholesterol + POPC (Fig. 5b, 5c). Growth rates

derived from Phenol Red absorbance of both organisms show that the introduction of entPOPC

to the lipid diet results in impaired growth, with the racemic diet yielding the most pronounced

decrease in growth rate. We also evaluated growth by measuring optical density, as a proxy for

cell density (Supplementary Figure S7), which confirmed a reduction in growth with the

introduction of entPOPC. However, in contrast, growth was similar for the entPOPC and racemic

lipid diets. This difference likely reflects the fact that Phenol Red growth rate estimates reflect

metabolic activity, which is not necessarily coupled with the production of cell biomass measured

by optical density. Thus, we demonstrate that cells are viable when fed enantiomeric

phospholipids.

We next asked how the introduction of entPOPC affected the mechanical robustness and

permeability of the membrane. To assay membrane robustness, we measured sensitivity of cells

to hypoosmotic shock, by determining what fraction of cells are lysed following a shock. Since *M.*

mycoides and JCVI-Syn3A lack a cell wall or cytoskeleton, cell lysis during rapid hypoosmotic shock

is indicative of membrane rupture strength. Furthermore, mechanosensitive ion channels which

can protect cells from hypoosmotic shock have, to the best of our knowledge, not been reported

in either organism, or annotated in the genomes, and are not present in all *Mycoplasmas*[47,55].

Therefore, it is reasonable to cautiously interpret the susceptibility to lysis from hypoosmotic

shock as indicative of membrane stability. However, we cannot rule out the possibility that there
are undiscovered mechanosensitive channels, and this would suggest that mechanosensitive
gating is sensitive to lipid chirality. Hypoosmotic sensitivity increased significantly for cells grown
on entPOPC diets, indicating that the enantiomeric lipids affect the mechanical robustness of the
membrane and, consequently, the whole cell (Fig. 5b, c). To assay permeability, we measured
the rate of permeation of fluorescein diacetate (FDA), a non-chiral molecule, across the cell
membrane[56,57]. Membrane permeability, as measured by the permeability coefficient of FDA
increased significantly upon the introduction of enantiomeric lipids, but was the highest for both
M. mycoides and JCVI-Syn3A for the racemic lipid diet (Fig. 5b, c). When plotted against growth
rate, there is an apparent correlation of lower growth with higher permeability (Fig 5d, e),
possibly indicating membrane leakiness as one of the factors underlying impaired growth. The
permeability coefficients were similar to those previously reported in mammalian cells[57].

Increased permeability and reduced membrane robustness could be due either to
changes in the property of the lipid bilayer, or through changes in lipid-protein interactions.
Previous work on PC enantiomers in model membranes revealed that modest changes in
permeability to calcein occur in membranes composed of both enantiomers, in particular
scalemic (not 1:1) mixtures. However, such small changes would not account for the large change
in permeability we observe. To determine if introducing enantiomeric POPC affected membrane
bilayer robustness or permeability we reconstituted cellular lipid extracts from *M. mycoides* into
liposomes and evaluated C-Laurdan fluorescence (Fig. 6). The C-Laurdan General Polarization
index (GP) reports bilayer hydration, which is closely coupled with permeability and mechanical
robustness of the membrane[58,59]. Surprisingly, GP values did not vary significantly across all three
POPC diets, indicating that the lipid bilayer itself was not disrupted by changes in chirality, and
implicating an effect on lipid-protein interactions. Indeed, several studies have demonstrated an
effect of varying lipid chirality on lipid-peptide interactions, and the permeability of membranes
to chiral amino acids[60,61]. Thus, disrupted lipid-protein interactions are the most likely basis for
the observed phenotypes, setting the stage for future work employing *M. mycoides* and JCVI-
Syn3A as model systems to explore the significance of lipid chirality on lipid-protein interactions
in a living membrane.

Discussion

In this study, we introduce *M. mycoides* and the Minimal Cell JCVI-Syn3A as simple model organisms with tunable lipidomes for studying the role of lipid complexity. By choosing a model membrane system incapable of synthesizing or modifying fatty acids, and developing a set of defined lipid diets for that system, we have demonstrated the creation of a platform in which the lipidome can be tuned in terms of phospholipid head group, acyl chain composition or lipid chirality. Using this platform, we created the simplest living membrane, and one incapable of undergoing acyl chain remodeling. These tunable living membranes allowed us to quantitatively examine the contribution of lipidomic features to the fitness of a minimal living system.

Lipid scavenging

Our observations broadly demonstrate how lipidome composition is crucial even for relatively simple microorganisms. This is anecdotally illustrated by the capacity we observe for mycoplasma to produce a complex lipidome from a single exogenous phospholipid, which is an impressive evolutionary adaptation to their pathogenic lifestyle. Indeed, this acyl chain scavenging activity, which confounded our initial attempt to simplify the lipidome, suggests a potential target for treating mycoplasma infection. Although mycoplasma can survive in pure culture with a reduced lipidome, they may fare much worse in the context of a host immune system. Furthermore, the ability for the Minimal Cell to grow with a lipidome of two lipids implies the possibility for further minimization of the genome, through the deletion of pathways involved in the scavenging of phospholipid acyl chains and the internal synthesis of phospholipids.

Minimal lipidomes

A long-standing challenge in membrane biology has been to understand why life has evolved such complex lipidomes and to identify the essential features of lipidomes required for optimal membrane function and cellular fitness. The simplest lipidome so far reported was from a Gram-negative bacterium, composed of 27 lipid species, excluding outer membrane lipopolysaccharides that were not analyzed[1]. By reducing the lipidome of M. mycoides and JCVI-Syn3A down to two lipids, we show that a complex lipidome is not essential for life, but that two lipids are far from optimal.

Through systematic reintroduction of lipidomic features into cells with a minimal

lipidome, we demonstrated that phospholipid headgroup diversity alone does not significantly

rescue growth. Consistently, minimizing headgroup diversity in *B. subtilis* has little effect on

growth[13]. Similarly, E. coli mutants lacking phosphatidylglycerol (PG) and cardiolipin can be viable

and often do not exhibit significant growth deficiencies[7,62]. It was perhaps most surprising that

M. mycoides grew comparably well in the absence of zwitterionic lipids (PC), which are essential

for *E. coli* to support proper membrane insertion and activity of membrane proteins[63]. These

observations emphasize the limited impact of headgroup diversity, at least in simple bacterial

organisms, in the absence of other critical lipid features.

Restoring acyl chain diversity, in contrast, did enhance growth rates. This finding is

consistent with recent work showing the importance of acyl chain unsaturation in various cellular

processes, including the assembly and function of the nuclear pore complex in eukaryotes[64] and

in neuronal membranes[65]. The enhanced growth observed when acyl chain diversity was restored

in *M. mycoides* suggests that while headgroup diversity alone is insufficient, it may contribute to

improved cellular fitness when coupled with the appropriate acyl chain composition. For

example, the introduction of cardiolipin or PG could potentially improve growth, but only if these

lipids are present with the correct acyl chain configurations that support optimal membrane

properties and lipid-protein interactions. The approach to tuning the lipidome that we introduce

here provides a platform to explore the synergistic effects of specific acyl chain-headgroup

combinations, to unravel how these lipidomic features can be optimized for membrane function

and cellular fitness in minimal systems.

By applying a targeted chemical approach to reduce the lipidome of JCVI-Syn3A,

we demonstrated the feasibility of further simplifying the molecular composition of a

genomically minimized organism. The observation that a minimal cell membrane can function

with only cholesterol and one species of PC demonstrates that the fundamental requirements

for life can be achieved with a remarkably simple lipid composition. For synthetic biology, this

insight simplifies the challenge of designing synthetic cells, revealing the potential to create

functional living membranes with minimal components. This work lays a foundation for future

efforts to understand how minimal lipidomes can be optimized in synthetic and engineered

biological systems

One limitation of this work is that the growth medium is not defined, and there is a very

small contribution of lipids from components of the growth medium, such as the yeast extract.

Thus, while the majority (>99%) of the minimal lipidome is composed of 2 lipids, there is a fraction

of a percent of very low abundance lipids derived from the media and it is possible, but unlikely,

that these trace lipids play a significant role. In this regard, development of a defined growth

medium will be essential in the continued development of mycoplasmas as minimal model

membrane systems[33]. Another factor limiting the minimization of the lipidome to one lipid is that

cholesterol (or an analogue) is required for growth, but also cannot form a bilayer alone. So, it is

possible that even one bilayer forming lipid could support life in a cell that has not evolved to

require sterols. It is also, however, possible that having a membrane reinforcing sterol or sterol

analogue is critical for building a stable cell membrane with only one phospholipid. A

mycoplasma-like organism such as *Mesoplasma* that does not require sterols for growth would

provide a means to test this hypothesis[66–68]. Nonetheless, a JCVI-Syn3A membrane with two lipids

comprising over 99% of the lipidome is currently the simplest living membrane that has been

reported.

Lipid chirality and the lipid divide

The lipid divide represents another major problem in membrane biology. Homochirality

is a fundamental feature of biomolecular chemistry. Biomolecules exhibit enantioselectivity for

chirally compatible interaction partners. Consequently, life has evolved to rely on homochiral

molecules (e.g. L-amino acids and nucleic acids). An interesting twist is in the divide between the

chirality of phospholipids made by Archaea and the rest of life[69]. The divide raises questions

about whether the last universal common ancestor (LUCA) possessed a heterochiral lipidome, or

whether the divide happened after the divergence of Bacteria and Archaea[70,71]. Further, current

theories that eukaryotes emerged from an Archaeal ancestor[72], create a conundrum in explaining

why Eukaryotes don't have Archaeal lipid enantiomers, and whether a gradual transition

involving heterochiral lipidomes could have occurred. At the crux of these conundrums is

whether heterochiral lipidomes can support stable membranes and optimal cellular fitness.

We took advantage of the tuneability of *M. mycoides* and JCVI-Syn3A lipidomes to

observe how heterochiral lipidomes influence the membrane and cellular growth. Previous work

in living systems has achieved lipidomes with a mixture of structurally diverse bacterial and

archaeal lipids that differed not only by chirality, but also in a variety of other ways[54]. A unique

feature of this study is that we were able to introduce two enantiomers of a single lipid structure

(POPC) into a living membrane. Our results demonstrate that a heterochiral lipidome results in a

leakier membrane and impaired cellular robustness and growth. In contrast, previous work in *E.*

coli indicated that heterochiral lipidomes did not affect growth[54]. *E. coli* has an outer membrane

and cell wall that might compensate for lack of stability caused by heterochirality. The lipids in

the *E. coli* study were not stereoisomers, but rather entirely different lipid structures (ether vs.

ester acyl chain linkages, isoprenoid tails vs. fatty acid tails). It is possible that these structural

differences somehow obscured the destabilizing effects of heterochirality. Ultimately, the basis

for these diverging observations will provide insight into the biological significance of lipid

chirality. Nonetheless, our work shows that in a minimal organism with a single membrane,

heterochiral lipidomes can support growth, but lead to impaired robustness and fitness.

Interestingly, this suggests that LUCA could have existed before the evolution of homochiral

cellular membranes. This possibility would eliminate the need to view the lipid divide between

bacteria and archaea as resulting from independent evolutionary events. Instead, it suggests the

ancestral lipidome could have been heterochiral, consistent with a simpler path for the evolution

of modern membranes. Additionally, the reduced fitness and increased membrane permeability

resulting from heterochiral lipidomes suggests selective pressure against such membranes in

ancestral organisms, favoring the evolution of the homochiral membranes characteristic of

modern life.

The changes in membrane robustness and permeability that we observed do not seem to

be explained by changes in lipid order of pure lipid vesicles reconstituted from cell lipid extracts

of homo- and heterochiral lipidomes. This implies that lipid-protein interactions are

enantioselective and may be affected by lipid chirality in ways that impair membrane robustness

and function. Numerous *in vitro* studies have demonstrated the effect of lipid chirality on lipid-

protein interactions, consistent with the possibility that the phenotypes we observed are rooted

in perturbed lipid-protein interactions. Looking forward, *M. mycoides* and JCVI-Syn3A will be
excellent model system to explore the functional consequences of lipid chirality on lipid-protein
interactions.

Outlook

Our approach to employ *M. mycoides* and JCVI-Syn3A as minimal living model membrane
systems paves a new path towards unraveling the role of lipidome diversity and complexity. Our
observations reveal that life does not require complex lipidomes. However, minimization comes
with clear trade-offs in cellular fitness. We further demonstrate the capacity of these model
systems to serve as chassis for exploring fundamental questions in membrane biology. The ability
of this system as a testing platform for lipid diets with a variety of features (including class and
acyl chain composition) makes it a useful and simple *in vivo* model for design-test-build
applications of membrane composition experiments. Furthermore, we demonstrated the ability
to force *M. mycoides* and JCVI-Syn3A to take up and incorporate enantiomeric lipids in their
membranes, the first time this has been shown in eukaryotic or prokaryotic organisms, and an
exciting first step to allow us to probe questions about the lipid divide and the role of chirality in
membrane stability and cell fitness. Overall, we have demonstrated the creation of a simple,
tunable, living, model membrane system that can be used as a novel platform for probing the
design principles of living membranes.

Acknowledgements

The authors wish to thank the Saenz lab; Lisa Junghans, Ha Ngoc Anh Nguyen, and Tomasz
Czerniak for discussions and feedback through the process, and for help with experimental
designs; the JCVI, and specifically John Glass and Kim Wise, for providing *M. mycoides* and JCVI-
Syn3A and for further feedback, including manuscript comments; Helena Jambor for assisting
with figure design; and Jennifer Katz-Brandoli for assisting with robot runs and figure
construction. We would like to thank Telesis Bio, Inc. for allowing us to use JCVI-syn3A and
Lipotype GmbH for lipidomic analyses. We would like to thank the Electron Microscopy Facility
of the MPI-CBG for their support. We additionally wish to thank Michaela Wilsch-Bräuninger from
the MPI-CBG for insights regarding EM images, Mareike Jordan from the Von Appen Lab for
assistance in reconstructing tomograms, as well as Adrian Nievergelt from the Gaia Pigino Group
at the Human Technopole Milano for his help processing cryo-EM tomograms with IsoNet. This
work was supported by the B CUBE of the TU Dresden, a German Federal Ministry of Education

and Research BMBF grant (to J.S., project 03Z22EN12), and a VW Foundation ''Life'' grant (to J.S.,
project 93090).
544
Materials and Methods

<u>Cell Culture:</u>

M. mycoides and JCVI-Syn3A were grown in liquid culture at 37ºC on a modified SP4 growth

medium with Phenol Red[73]. The lipid source was provided either by Fetal Bovine Serum (FBS),

complex natural source of mammalian lipids, or by complexing lipids with delipidated bovine

serum albumin (BSA). Lipids were dissolved in ethanol and added to the medium at 37ºC

immediately before the addition of cells, in the concentrations given in Table S1. *M. mycoides*

was passaged to new growth medium by adding 50 µL of cell culture at OD600 0.4-0.8 into 7 mL

of freshly prepared media in a T25 Flask (Stand., Vent. Cap) or 10mL of media in a Duran 100 mL

glass flask. This corresponded to roughly one passage per day. JCVI-Syn3A was passaged to new

medium by adding 200 µL of cell culture at OD600 0.2-0.6 into 7 mL of freshly prepared media.

This corresponded to one passage every two to four days, depending on cell growth rates. Cells

were grown at 37ºC and 40 RPM in a Kuhner shaker incubator.

<u>Growth Rates:</u>

Metabolic Growth Index

Due to differences in cell size and behavior between *M. mycoides* and JCVI-Syn3A, measuring

growth curves with optical density by taking the absorbance at 600 nm (OD_{600}) proved to be an

unreliable method to achieve growth estimates that were robustly comparable between the two

organisms. Phenol Red is a pH indicator that is commonly used as a readout for growth in

mycoplasma cell culture[74] as it detects changes in pH as metabolic activity acidifies the culture

media through absorbance at 562 nm (A_{562}). To obtain the rate of pH change (A_{562}) over time cells

were grown on the liquid handler system Biomek i7 Automated Workstation in 96-well plates

(square well, clear bottom, 400 uL media volume, 100 rpm shaking), and the absorbance was

recorded hourly to record changes in phenol red absorbance as cell growth acidifies the media

(Supplementary Fig. S3a)[19,73,74]. All absorbances were then plotted with respect to time, and then

fitted to the logistic function $N_t = K (1 + (K-N_0)/N_0) e^{-rt})$ where N_t is population at time t, K is the

carrying capacity, N_0 is the initial population size, and r is the growth rate; or, in this case, the

rate of pH change. Only the exponential part of the curve was considered to exclude lag and

stationary phase. To validate the relevance of this method as a readout for cell growth, growth

rates for a representative set of *M. mycoides* diets were calculated using OD_{600} and were

compared to the rates calculated with Phenol Red absorbance. The result was a strong linear

correlation between the two estimates (Supplementary Fig. S3b). Representative curves of the

entire period of growth for *M. Mycoides* and JCVI-Syn3B on three different diet and generated

by the two methods is also shown (Supplementary Fig S5). Cell growth data shown in

Supplementary Figure S5, measured by Phenol Red A_{562} and OD_{600} was obtained through hourly

manual measurements from cultures grown in 30 mL in a Duran 100 mL glass flask in a Kuhner

shaker incubator at 37ºC and 40 rpm shaking using a DeNovix Spectrophotometer.

<u>Lipid Extraction and Thin Layer Chromatography:</u>

Lipids were extracted using a Bligh-Dyer lipid extraction protocol[76]. When appropriate, lipid

concentration was determined with a phosphate assay. TLC plates with Silica Gel and a 10x

concentrating zone were pre-washed with chloroform to remove debris and other artifacts, and

were then dried at 60ºC for > 30 minutes to ensure complete solvent evaporation. Lipid samples

were then loaded on the plate and were dried at 60ºC for > 10 minutes to again ensure complete

solvent evaporation. The plate was then run in a closed glass chamber using a running solution

of Chloroform:Methanol:Acetic Acid:Water (85:25:5:4), and were subsequently dried again at

60ºC for > 30 minutes. TLC plates were then rinsed with a 3% Copper acetate and 8% phosphoric

acid solution and heated by a handheld heat gun at 280ºC to char lipid spots.

<u>Liposome Preparation:</u> Lipids of a known concentration (either in stock solutions or extracted

from cells and validated with a phosphate assay) were prepared in a 2 mL glass by having their

solvents evaporated overnight under a 10-17 mbar vacuum. Lipids were then resuspended in the

appropriate volume of Liposome buffer (LiB) (10 mM HEPES, 100 mM NaCl) to achieve 100 mM

liposome solution and incubated for 30 minutes at 37ºC. Subsequently, liposomes were

homogenized with ten freeze thaw cycles (1 minute in liquid nitrogen followed by 5 minutes at

60C) and 7 extrusion cycles in a Hamilton syringe setup through a 100 nm filter.

Propidium Iodide Osmotic Shock Assay: 3X 0.4 ODU of cells with OD 0.2-0.4 (1 ODU = 1 mL of cells

at OD 1) were harvested and spun down in pre-warmed centrifuge at 5000 g, 7minutes, 37ºC,

with slow acceleration slow deceleration. 1 µL of 1 mM Propidium Iodide dye was added to

appropriate wells on a black or clear bottom 96well plate. The supernatant was aspirated and

each cell pellet was resuspended in 400 µL of either H2O or 5X diluted mycoplasma wash buffer

(MWB) (original MWB: 20 mM HEPES, 200 mM NaCl, 1% W/V Glucose). An extra tube with cells

resuspended in 400 µL H2O and boiled at 95ºC in a thermoshaker at 1000 rpm for 10 minutes

was set up as a positive control. During that time, all other tubes were incubated on the 37ºC

thermoshaker at 600 rpm. 100 µL of the cells in the appropriate buffer was added to each of their

three analytical wells on the 96 well plate. Cells were incubated for 30 minutes with shaking at

37ºC in a TECAN Spark 20M plate reader and the fluorescent signal was subsequently measured

at excitation 539 nm emission 619 nm.

Fluorescein Diacetate Permeability Assay: 3X 0.4 ODU of cells with OD 0.2-0.4 were harvested

and spun down in pre-warmed centrifuge at 5000 g, 7minutes, 37ºC, with slow acceleration slow

deceleration. Cells were washed 1x in MWB and resuspended in 400 µL MWB. A Fluorescein

Standard curve was prepared on the plate in the following concentrations: 0, 0.25 ,0.5, 1.2, 2 µM

Fluorescein. 100 µL resuspended cells were added to the proper wells (in triplicate), and 100 µL

MWB was added to the Fluorescein standard wells. FDA was added to cell wells to achieve final

FDA concentration of 5 µM. Cells were incubated on the TECAN Spark 20M at 37ºC for 140

minutes and measurements were taken every 20 minutes at excitation 485 and emission 525.

The permeability coefficient (P) of FDA was calculated using Fick's Law ($Q = P*A*(C_{out} - C_{in})$) where

Q is the flux across the membrane, and is a constant as the slope of fluorescein increase over

time was linear; A is the area of the membrane, calculated using the assumption that the cross

sectional area of POPC + Cholesterol in a bilayer is 45.1 Å[77], the ratio of POPC : Cholesterol is

roughly 1:1, the area of POPC + cholesterol is equivalent to the area of Enantiomeric POPC +

Cholesterol, and the number of POPC molecules can be calculated from the phosphate assay

described below; C_{out} and C_{in} are the concentrations of FDA outside and inside the cell

(respectively); and Ci_n is 0 as FDA is immediately converted to fluorescein upon entering the cell.

C-Laurdan General Polarization Assay: Liposomes from whole cell lipid extracts were incubated

at 37ºc and 1000 rpm for 10 minutes on a tabletop thermoshaker. While cells were incubating,

c-Laurdan was removed from a -20ºc freezer and warmed to room temperature on the bench. 1

mM c-Laurdan stock in EtOH was diluted 4x and 1 uL of diluted stock for a final molarity 0.5 µM

C-Laurdan was added to 500 µL of liposomes. Samples were incubated at 37ºc and 1000 rpm

shaking for 10 minutes on a tabletop thermoshaker. 100 µL of each sample was added to a flat

bottom 96 well plate well with analytical triplicates for each biological replicate. The signal was

measured on a TECAN Spark 20M with a 2 channel fluorescence reading with excitation at 385nm

and emission at 440 and 490 nm respectively. General polarization (GP) was calculated using the

formula: GP = (I440 - I490) / (I440 + I490) where I is the fluorescence emission intensity at the

respective wavelength.

Phosphate Assay: To estimate the concentration of phospholipids in a certain amount of cells,

the amount of phosphate in the lipid extraction of 1 OD Unit (ODU: 1mL of OD_{600} = 1) of cells was

measured. To measure phosphate amount a modified version of the method of Chen et al. was

used[78]. The lipid extraction was added to glass pyrex tubes with both biological and analytical

triplicates (3 vials per biological replicate), and the solvent was evaporated by a brief (<5 minute)

incubation at 200ºC. 50 µL of water were added to each sample. To prepare a standard curve of

phosphate amounts, an ICP Phosphorus standard was diluted in water to give 5, 10, 20, 50 and

100, and 200 nmols of phosphate. 500 µL of 70% perchloric acid was added to each vial and, after

brief vortexing, samples were incubated at 200ºC for 120 minutes. Tubes were cooled down in

ice water, and 1mL of 10% w/v ascorbic acid followed immediately by 1 mL of 2.5% w/v

ammonium heptamolybdate were added (with a brief vortex after adding each reagent). Samples

were incubated at 37ºC for 30 minutes, and then absorbance at 820 nm was measured on a

TECAN Spark 20M plate reader by adding 200 µL of each tube to a clear bottom 96 well plate . A

standard curve of absorbance at 820 vs phosphate amount was calculated using the phosphate

standard, and lipid amount of each sample was calculated based on the absorbance value,

standard curve, and assumption that one phosphate molecule equals one phospholipid.

<u>Lipidomic Analysis:</u> Lipids extracted from cells using the aforementioned Bligh-Dyer protocol

were submitted to Lipotype for mass spectrometry based analysis[79]. The general procedure is

described in Sampaio et.al, 2011[2]. For the analysis, samples were spiked with internal lipid

standard mixture containing: cardiolipin 14:0/14:0/14:0/14:0, ceramide 18:1;2/17:0,

diacylglycerol 17:0/17:0, hexosylceramide 18:1;2/12:0, lyso-phosphatidate 17:0, lyso-

phosphatidylcholine 12:0, lyso-phosphatidylethanolamine 17:1, lyso-phosphatidylglycerol 17:1,

lyso-phosphatidylinositol 17:1, lyso-phosphatidylserine 17:1, phosphatidate 17:0/17:0,

phosphatidylcholine 17:0/17:0, phosphatidylethanolamine 17:0/17:0, phosphatidylglycerol

17:0/17:0, phosphatidylinositol 16:0/16:0, phosphatidylserine 17:0/17:0, cholesterol ester 20:0,

sphingomyelin 18:1;2/12:0;0, triacylglycerol 17:0/17:0/17:0, and cholesterol D6. After extraction,

the organic phase was transferred to an infusion plate and dried in a speed vacuum concentrator.

It was then resuspended in 7.5 mM ammonium acetate in chloroform/methanol/propanol (1:2:4,

V:V:V) and a 33% ethanol solution of methylamine in chloroform/methanol (0.003:5:1; V:V:V).

All liquid handling steps were performed using Hamilton Robotics STARlet robotic platform with

the Anti Droplet Control feature for organic solvents pipetting. Samples were analyzed by direct

infusion on a QExactive mass spectrometer (Thermo Scientific) equipped with a TriVersa

NanoMate ion source (Advion Biosciences). Samples were analyzed in both positive and negative

ion modes with a resolution of Rm/z=200=280000 for MS and Rm/z=200=17500 for MSMS

experiments, in a single acquisition. MSMS was triggered by an inclusion list encompassing

corresponding MS mass ranges scanned in 1 Da increments. Both MS and MSMS data were

combined to monitor CE, DAG and TAG ions as ammonium adducts; PC, PC O-, and Diether PC as

acetate adducts; and CL, PA, PE, PE O-, PG, PI and PS as deprotonated anions. MS only was used

to monitor LPA, LPE, LPE O-, LPI and LPS as deprotonated anions; ceramide, hexosylceramide,

sphingomyelin, LPC and LPC O- as acetate adducts and cholesterol as an ammonium adduct of an

acetylated derivative[80].

<u>Room temperature TEM</u>:

JCVI-Syn3A cells were adapted to FBS, POPC + Cholesterol, and D.PC + Cholesterol diets and

harvested at between OD_{600} 0.05-0.2 at volumes to achieve 1ODU of cells for each diet. Cells

were spun down and resuspended in 1mL Mycoplasma Wash Buffer without glucose (20mM

HEPES, 200 mM NaCl). Cells were pre-fixed 30 min at room temperature with a final

concentration of 0.5% glutaraldehyde (25% aqueous stock solution (EMS), directly added to the

culture medium). Then cells were spun down for 7 min @ 7000 g @ 37°C in a centrifuge (Heraeus

Biofuge PrimoR with swingout buckets). The cell pellet was sucked into cellulose capillary tubes

with an inner diameter of 200 μm and a permeability cut-off >5 KD (Leica). Filled capillaries were

chopped into small pieces with a scalpel blade, at the same time sealing the capillary ends.

Capillary pieces were rapidly frozen using Leica EM-ICE high pressure freezing device in 6 mm–

aluminum carriers with 200 μm deep depression (Leica) using hexadecene as a filler. Freeze

substitution was done in an AFS 2 freeze substitution device (Leica) in 1% osmium tetroxide in

acetone starting at -90 °C and raising the temperature to 0° C over 72 hrs. Samples were

infiltrated gradually with 25%, 50%, 75%, 100% EMbed 812 (Science Services) in acetone over

two days. Capillaries were infiltrated with pure resin over two days and embedded in double end

silicon molds (TED PELLA INC) and polymerized at 60 °C for 48 hrs. 70 nm–sections were cut with

a Leica EM UC6 and mounted on formvar-coated copper slot grids (EMS). Sections on the grids

were contrasted with uranyl acetate and lead citrate prior to imaging. Imaging was done on a

Tecnai T12 (Thermo Fisher Scientific (formerly FEI), Hillsboro, Oregon, USA) transmission electron

microscope at 100 kV acceleration voltage. The images were acquired with a F416 camera (Tietz

Video and Image Processing Systems GmbH, Gilching, Germany) at 4096x4096 pixels using

SerialEM software. Before taking an image, the sample was automatically exposed to the beam

for 0.5-2 seconds to reduce drift. The exposure time (between 0.5 and 1.5 seconds) was

subdivided into 3 subframes which were then automatically aligned by cross correlation and

added up in order to form the final image in order to reduce image blurring by sample drift.

<u>*Cryo TEM*</u>:

JCVI-Syn3A cells were adapted to FBS, POPC + Cholesterol, and D.PC + Cholesterol diets and
harvested at between OD_{600} 0.05-0.2 (mid-exponential growth phase) at volumes to achieve
1ODU of cells for each diet. Short before freezing, cells were spun down and resuspended in 1
mL Mycoplasma wash buffer without glucose (20 mM HEPES, 200 mM NaCl). Quantifoil 2.1
copper 200 mesh grids were cleaned with chloroform before usage. Glow discharging was
performed on a PELCO Easy glow for 30 seconds, 15 mA on both sides of the grid. 2 μl of sample
being added to both sides of the grid, 1 μl of gold (Protein A gold, PAG 10, pre-diluted 1:25 was
added to the carbon side only. Cryo fixation was performed by plunge freezing in liquid Ethane
using a Leica GP with humidity chamber set to 21 ˚C, a humidity of 98 % and a blotting time of 5
seconds from the back of the grid with Whatmann paper No 1. Frozen grids were stored in liquid
nitrogen[81].

Cryo-electron tomography was done on a Titan Halo transmission electron microscope with field
emission gun electron source and a Gatan K2 Summit direct electron detector at 300 kV with an
energy filter using a slit width of 20 eV. Full grid overview was acquired with SerialEM by
automatically acquiring and stitching low-magnification (×210) images. Tilt series was taken with
SerialEM on areas of interest at ×30,000 nominal image magnification, calibrated pixel size of
2.36 Å (super-resolution mode) and 2° increments with a bidirectional tilt scheme from 20° to -
58° and from 22° to 58°. The acquisition was done with a defocus target of 5 μm and the
accumulative dose was 80 - 90 e– per Å2 per tomogram. Images were acquired in dose
fractionation mode with frame times between 0.10 and 0.25 s. Correction of the sample motion
induced by the electron beam was done with MotionCor. Tomogram reconstruction was
performed using Etomo from IMOD 4.11.18 using weighted back projection. Contrast transfer
function curves were estimated with Ctfplotter and corrected by phase-flipping with the software
Ctfphaseflip, both implemented in IMOD. Dose-weighted filtering was performed by using the
mtffilter implemented in IMOD. To enhance the contrast of macromolecular structures and fill
up missing wedge information IsoNet was used. Visualization of tomograms and averaged
electron density maps was performed in 3dmod from IMOD[82]. Computer visualization of three-
dimensional image data using IMOD. J. Struct. Biol. 116:71-76.); rendering of isosurfaces and

structure fitting was performed using UCSF ChimeraX 1.8, developed by the Resource for

Biocomputing, Visualization, and Informatics at the University of California, San Francisco, with

support from National Institutes of Health R01-GM129325 and the Office of Cyber Infrastructure

and Computational Biology, National Institute of Allergy and Infectious Diseases.

<u>Statistical Analysis</u>

For growth rate calculations in Figures 3, 4, and 5 the number of biological and analytical

replicates is given in Table S1. For the propidium iodide assays in Figure 5 there are three

biological replicates each with three analytical replicates. For the FDA assay in Figure 5 there are

three biological replicates each with three analytical replicates. For the c-Laurdan assay in Figure

6 there are three biological replicates each with three analytical replicates. Statistical significance

was calculated with an unpaired t-test. All error bars indicate the mean +/- the standard

deviation.

<u>Materials:</u>

SP4 (For 1 L):

PPLO (3.5 g) (Becton, Dickinson, and Company product no. 255420)

Tryptone (10 g) (Sigma product no 70169)

Peptone (5 g) (Sigma product no 70176)

20% Glucose (25 mL) (Ross product no. X997.2)

20% Yeastolate (10 mL) (Becton, Dickinson, and Company product no. 255772)

15% Yeast Extract (35 mL) (Roth product no. 2904.3)

70g/L BSA (85 mL) (Sigma product no. A7030) or FBS (170 mL) (Biowest Product

no. 5181H-500)

400,000U/mL Penicillin G-sodium salt (2.5 mL) (Roth product no. HP48.1)

10 mg/mL L-Glutamate (5 mL) (Roth product no. HN08.2)

Sodium Bicarbonate (1.04 g) (Honeywell product no. 71630)

CMRL (4.9 g) (US Biological Lifesciences product no. C5900)

Phenol Red (11 mg) (Sigma product no. P3532)

Lipids:

776   Cholesterol (Avanti product no. 700100)

777   Diether POPC (Diether 1-palmitoyl-2-oleoyl-glycero-3-phosphocholine) (Avanti
778   product no. 999983)

779   Diether DPPC (Diether 1-palmitoyl-2-palmitoyl-glycero-3-phosphocholine) (Avanti
780   product no. 999992)

781   Diether DOPC (Diether 1-oleoyl-2-oleoyl-glycero-3-phosphocholine) (Avanti
782   product no. 999991)

783   Diether POPG   (Diether 1-palmitoyl-2-oleoyl-glycero-3-phosphotdylglycerol)
784   (Avanti product no. 999973)

785   POPC (1-palmitoyl-2-oleoyl-glycero-3-phosphocholine) (Avanti product no.
786   850457)

787   DPPC (1-palmitoyl-2-palmitoyl-glycero-3-phosphocholine) (Avanti product no.
788   850355)

789   DOPC (1-oleoyl-2-oleoyl-glycero-3-phosphocholine) (Avanti product no. 850375)

790   POPG (1-palmitoyl-2-oleoyl-glycero-3-phosphotdylglycerol) (Avanti product no.
791   840457)

792   DPPG (1-palmitoyl-2-palmitoyl-glycero-3-phosphoglycerol) (Avanti product no.
793   840455)

794   DOPG (1-oleoyl-2-oleoyl-glycero-3-phosphoglycerol) (Avanti product no. 850475)

795   POPE (1-palmitoyl-2-oleoyl-glycero-3-phosphoethanolamine) (Avanti product no.
796   850757)

798   DPPE (1-palmitoyl-2-palmitoyl-glycero-3-phosphoethanolamine) (Avanti product
799   no. 850705)

800   DOPE (1-oleoyl-2-oleoyl-glycero-3-phosphoethanolamine) (Avanti product no.
801   850725)

802   Oleic Acid (*cis*-9-Octadecenoic acid) (Sigma product no. 01383)

Palmitic Acid (1-Pentadecanecarboxylic acid) (Sigma P0500)

Enantiomeric POPC (3-palmitoyl-2-oleoyl-glycero-1-phosphocholine) (Avanti

product no. 850855)

16-0 Cardiolipin (1',3'-bis[1,2-dipalmitoyl-sn-glycero-3-phospho]-glycerol) (Avanti

product no. 710333)

Assays:

FDA (Sigma product no. F7378)

Fluorescein (Fluka product no. 28803)

Propidium Iodide (Sigma product no. 537059)

C-Laurdan (Stratech Scientific Ltd. product no. T0001-SFC-1)

Cells:

JCVI-Syn3A-*mCherry (*From Telesis Bio)

*Mycoplasma mycoides subspecies capri strain GM12 (*From Telesis Bio)

Machines:

TECAN Spark 20M with Te-Cool cooling module

Biomek i7 Automated Workstation

DeNovix DS-11 FX +

PicoQuant FluoTime 300 High Performance Fluorescence Lifetime and Steady

State Spectrometer

Data Availability Statement:

The lipidomic datasets generated in this study have been uploaded to the Zenodo repository

and can be found at the DOI: 10.5281/zenodo.13817894. Due to proprietary restrictions

imposed by the data provider, Lipotype GmbH, raw lipidomics instrument files cannot be

deposited in a standard metabolomics repository, as they contain commercially sensitive

information. The TEM data has been deposited in the EBI Biolibraries repository and can be

found at the DOI: 10.6019/S-BSST1651. The cryoEM data has been deposited in the Electron

Microscopy Data Bank (EMDB) within the pDBE database, and can be found with the accession

codes: EMD-51614, EMD-51606, EMD-51607, EMD-51608, EMD-51609, and EMD-51610. All other source data are provided with this paper.

Citations

1. Chwastek, G. *et al.* Principles of Membrane Adaptation Revealed through Environmentally Induced Bacterial Lipidome Remodeling. *Cell Rep.* **32**, 108165 (2020).
2. Sampaio, J. L. *et al.* Membrane lipidome of an epithelial cell line. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 1903–1907 (2011).
3. Melero, A. & Jiménez-Rojo, N. Cracking the membrane lipid code. *Curr. Opin. Cell Biol.* **83**, 102203 (2023).
4. Harayama, T. & Riezman, H. Understanding the diversity of membrane lipid composition. *Nat. Rev. Mol. Cell Biol.* **19**, 281–296 (2018).
5. Klose, C. *et al.* Flexibility of a Eukaryotic Lipidome – Insights from Yeast Lipidomics. *PLoS ONE* **7**, e35063 (2012).
6. Safronova, N., Junghans, L. & Saenz, J. P. Temperature change elicits lipidome adaptation in the simple organisms Mycoplasma mycoides and JCVI-syn3B. *Cell Rep.* **43**, (2024).
7. Dowhan, W. A retrospective: Use of Escherichia coli as a vehicle to study phospholipid synthesis and function. *Biochim. Biophys. Acta BBA - Mol. Cell Biol. Lipids* **1831**, 471–494 (2013).
8. Cronan, J. E. & Gelmann, E. P. An Estimate of the Minimum Amount of Unsaturated Fatty Acid Required for Growth of Escherichia coli. *J. Biol. Chem.* **248**, 1188–1195 (1973).
9. Boudreaux, D. P., Freese, E., Eisenstadt, E. & Iijima, T. Biochemical and Genetic Characterization of an Auxotroph of Bacillus subtilis Altered in the Acyl-CoA:Acyl-Carrier-Protein Transacylase. *Eur. J. Biochem.* **115**, 175–182 (1981).
10. Budin, I. *et al.* Viscous control of cellular respiration by membrane lipid composition. *Sci. N. Y. NY* **248**, eaat7925 (2018).
11. Gohrbandt, M. *et al.* Low membrane fluidity triggers lipid phase separation and protein segregation in living bacteria. *EMBO J.* **41**, e109800 (2022).
12. Willdigg, J. R. & Helmann, J. D. Mini Review: Bacterial Membrane Composition and Its Modulation in Response to Stress. *Front. Mol. Biosci.* **8**, (2021).
13. Salzberg, L. I. & Helmann, J. D. Phenotypic and Transcriptomic Characterization of Bacillus subtilis Mutants with Grossly Altered Membrane Composition▽ †. *J. Bacteriol.* **190**, 7797–7807 (2008).
14. Rottem, S. & Kahane, I. *Mycoplasma Cell Membranes*. (Springer Science & Business Media, 2012).
15. Kornspan, J. D. & Rottem, S. The Phospholipid Profile of Mycoplasmas. *J. Lipids* **2012**, 640762 (2012).
16. McElhaney, R. N. Physical Studies of Lipid Organization and Dynamics in Mycoplasma Membranes. in *Mycoplasma Cell Membranes* (eds. Rottem, S. & Kahane, I.) 53–108 (Springer US, Boston, MA, 1993). doi:10.1007/978-1-4615-2924-8_3.
17. Bittman, R. Mycoplasma Membrane Lipids Chemical Composition and Transbilayer Distribution. in *Mycoplasma Cell Membranes* (eds. Rottem, S. & Kahane, I.) 29–52 (Springer US, Boston, MA, 1993). doi:10.1007/978-1-4615-2924-8_2.

18.   Razin, S., Yogev, D. & Naot, Y. Molecular biology and pathogenicity of mycoplasmas.
Microbiol. Mol. Biol. Rev. **62**, 1094-+ (1998).
19.   Gaspari, E. *et al.* Model-driven design allows growth of Mycoplasma pneumoniae on
serum-free media. *Npj Syst. Biol. Appl.* **6**, 1–11 (2020).
20.   Hutchison, C. A. *et al.* Design and synthesis of a minimal bacterial genome. *Sci. N. Y. NY*
351, aad6253–aad6253 (2016).
21.   Pelletier, J. F. *et al.* Genetic requirements for cell division in a genomically minimal cell.
Cell **184**, 2430-2440.e16 (2021).
22.   Lartigue, C. *et al.* Creating bacterial strains from genomes that have been cloned and
engineered in yeast. *Science* **325**, 1693–1696 (2009).
23.   Rottem, S., Yashouv, J., Ne'eman, Z. & Razin, S. Cholesterol in mycoplasma membranes.
Composition, ultrastructure and biological properties of membranes from *Mycoplasma*
mycoides var. *capri* cells adapted to grow with low cholesterol concentrations. *Biochim.*
Biophys. Acta BBA - Biomembr. **323**, 495–508 (1973).
24.   Archer, D. B. Y. 1975. Modification of the Membrane Composition of Mycoplasma
mycoides subsp. capri by the Growth Medium. *Microbiology* **88**, 329–338.
25.   Dahl, J. Uptake of fatty acids by Mycoplasma capricolum. *J. Bacteriol.* **170**, 2022–2026
(1988).
26.   Razin, S., Efrati, H., Kutner, S. & Rottem, S. Cholesterol and Phospholipid Uptake by
Mycoplasmas. *Rev. Infect. Dis.* **4**, S85–S92 (1982).
27.   Plackett, P. The Synthesis of Polar Lipids by Mycoplasma. *Ann. N. Y. Acad. Sci.* **143**, 158–
164 (1967).
28.   Salman, M. & Rottem, S. The cell membrane of *Mycoplasma penetrans*: lipid
composition and phospholipase A1 activity. *Biochim. Biophys. Acta BBA - Biomembr.* **1235**, 369–
377 (1995).
29.   Bhandari, S. & Asnani, P. J. Characterization of phospholipase A2 of mycoplasma species.
Folia Microbiol. (Praha) **34**, 294–301 (1989).
30.   Rottem, S., Adar, L., Gross, Z., Ne'eman, Z. & Davis, P. J. Incorporation and modification
of exogenous phosphatidylcholines by mycoplasmas. *J. Bacteriol.* **167**, 299–304 (1986).
31.   Rottem, S. Membrane lipids of mycoplasmas. *Biochim. Biophys. Acta BBA - Biomembr.*
604, 65–90 (1980).
32.   Rottem, S. & Markowitz, O. Membrane lipids of Mycoplasma gallisepticum: a
disaturated phosphatidylcholine and a phosphatidylglycerol with an unusual positional
distribution of fatty acids. *Biochemistry* **18**, 2930–2935 (1979).
33.   Thornburg, Z. R. *et al.* Fundamental behaviors emerge from simulations of a living
minimal cell. *Cell* **185**, 345-360.e28 (2022).
34.   Gross, Z. & Rottem, S. Lipid interconversions in aging Mycoplasma capricolum cultures.
J. Bacteriol. **167**, 986–991 (1986).
35.   Razin, S. Sterols in Mycoplasma Membranes. in *Current Topics in Membranes and*
Transport (eds. Bronner, F. & Kleinteller, A.) vol. 17 183–205 (Academic Press, 1982).
36.   Dahl, J. The Role of Cholesterol in Mycoplasma Membranes. in *Mycoplasma Cell*
Membranes (eds. Rottem, S. & Kahane, I.) 167–188 (Springer US, Boston, MA, 1993).
doi:10.1007/978-1-4615-2924-8_5.

37.   van Meer, G., Voelker, D. R. & Feigenson, G. W. Membrane lipids: where they are and
how they behave. *Nat. Rev. Mol. Cell Biol.* **9**, 112–124 (2008).
38.   Janmey, P. A. & Kinnunen, P. K. J. Biophysical properties of lipids and dynamic
membranes. *Trends Cell Biol.* **16**, 538–546 (2006).
39.   Winnikoff, J. R. *et al.* Homeocurvature adaptation of phospholipids to pressure in deep-
sea invertebrates. *Science* **384**, 1482–1488 (2024).
40.   Mouritsen, O. G. & Zuckermann, M. J. What's so special about cholesterol? *Lipids* **39**,
1101–1113 (2004).
41.   Varma, M. & Deserno, M. Distribution of cholesterol in asymmetric membranes driven
by composition and differential stress. *Biophys. J.* **121**, 4001–4018 (2022).
42.   Rangholia, N., Leisner, T. M. & Holly, S. P. Bioactive Ether Lipids: Primordial Modulators
of Cellular Signaling. *Metabolites* **11**, 41 (2021).
43.   Choquet, C. G., Patel, G. B., Sprott, G. D. & Beveridge, T. J. Stability of pressure-extruded
liposomes made from archaeobacterial ether lipids. *Appl. Microbiol. Biotechnol.* **42**, 375–384
(1994).
44.   Safronova, N., Junghans, L. & Saenz, J. P. Temperature change elicits lipidome
adaptation in the simple organisms Mycoplasma mycoides and JCVI-syn3B. *Cell Rep.* **43**, (2024).
45.   Breuer, M. *et al.* Essential metabolism for a minimal cell. *eLife* **8**, e36842 (2019).
46.   Sakai, A. *et al.* Traditional protocols and optimization methods lead to absent expression
in a mycoplasma cell-free gene expression platform. *Synth. Biol.* **7**, ysac008 (2022).
47.   Bianchi, D. M., Pelletier, J. F., Hutchison, C. A., Glass, J. I. & Luthey-Schulten, Z. Toward
the Complete Functional Characterization of a Minimal Bacterial Proteome. *J. Phys. Chem. B*
126, 6820–6834 (2022).
48.   Studer, P. *et al.* Proliferation of Listeria monocytogenes L-form cells by formation of
internal and external vesicles. *Nat. Commun.* **7**, 13631 (2016).
49.   van Wolferen, M., Pulschen, A. A., Baum, B., Gribaldo, S. & Albers, S.-V. The cell biology
of archaea. *Nat. Microbiol.* **7**, 1744–1755 (2022).
50.   Altamura, E. *et al.* Racemic Phospholipids for Origin of Life Studies. *Symmetry* **12**, 1108
(2020).
51.   Villanueva, L. *et al.* Bridging the membrane lipid divide: bacteria of the FCB group
superphylum have the potential to synthesize archaeal ether lipids. *ISME J.* **15**, 168–182 (2021).
52.   Martin, H. S., Podolsky, K. A. & Devaraj, N. K. Probing the Role of Chirality in
Phospholipid Membranes. *ChemBioChem* **22**, 3148–3157 (2021).
53.   Sojo, V. Why the Lipid Divide? Membrane Proteins as Drivers of the Split between the
Lipids of the Three Domains of Life. *BioEssays* **41**, 1800251 (2019).
54.   Caforio, A. *et al.* Converting Escherichia coli into an archaebacterium with a hybrid
heterochiral membrane. *Proc. Natl. Acad. Sci.* **115**, 3704–3709 (2018).
55.   Brown, M. B., Stoll, M., Maxwell, J. & Senior, D. F. Survival of feline mycoplasmas in
urine. *J. Clin. Microbiol.* **29**, 1078–1080 (1991).
56.   Rizk, S. *et al.* Functional diversity of isoprenoid lipids in Methylobacterium extorquens
PA1. *Mol. Microbiol.* (2021) doi:10.1111/mmi.14794.
57.   Leventai, K. R. *et al.* Lipidomic and biophysical homeostasis of mammalian membranes
counteracts dietary lipid perturbations to maintain cellular fitness. *Nat. Commun.* **11**, 1339
(2020).

58.   Parasassi, T., Stasio, G. de, Ravagnan, G., Rusch, R. M. & Gratton, E. Quantitation of lipid
phases in phospholipid vesicles by the generalized polarization of Laurdan fluorescence.
Biophys. J. **60**, 179–189 (1991).
59.   Steinkühler, J., Sezgin, E., Urbančič, I., Eggeling, C. & Dimova, R. Mechanical properties
of plasma membrane vesicles correlate with lipid order, viscosity and cell density. *Commun.*
Biol. **2**, 337–8 (2019).
60.   Chen, L. *et al.* Chiral Selective Transmembrane Transport of Amino Acids through
Artificial Channels. *J. Am. Chem. Soc.* **135**, 2152–2155 (2013).
61.   Ishigami, T., Suga, K. & Umakoshi, H. Chiral Recognition of l-Amino Acids on Liposomes
Prepared with l-Phospholipid. *ACS Appl. Mater. Interfaces* **7**, 21065–21072 (2015).
62.   Dowhan, W. Molecular genetic approaches to defining lipid function. *J. Lipid Res.* **50**,
S305–S310 (2009).
63.   Dowhan, W. & Bogdanov, M. Molecular genetic and biochemical approaches for
defining lipid-dependent membrane protein folding. *Biochim. Biophys. Acta BBA - Biomembr.*
1818, 1097–1107 (2012).
64.   Romanauska, A. & Köhler, A. Lipid saturation controls nuclear envelope function. *Nat.*
Cell Biol. **25**, 1290–1302 (2023).
65.   Manni, M. M. *et al.* Acyl chain asymmetry and polyunsaturation of brain phospholipids
facilitate membrane vesiculation without leakage. *eLife* **7**, e34394 (2018).
66.   Nguyen, H. N. A., Sharp, L., Lyman, E. & Saenz, J. P. Varying the position of phospholipid
acyl chain unsaturation modulates hopanoid and sterol ordering. *Biophys. J.* **123**, 1896–1902
(2024).
67.   Mccoy, R. E. *et al.* Acholeplasma florum, a New Species Isolated from Plants†. *Int. J. Syst.*
Evol. Microbiol. **34**, 11–15 (1984).
68.   Matteau, D. *et al.* Integrative characterization of the near-minimal bacterium
Mesoplasma florum. *Mol. Syst. Biol.* **16**, e9844 (2020).
69.   Koga, Y., Kyuragi, T., Nishihara, M. & Sone, N. Did Archaeal and Bacterial Cells Arise
Independently from Noncellular Precursors? A Hypothesis Stating That the Advent of
Membrane Phospholipid with Enantiomeric Glycerophosphate Backbones Caused the
Separation of the Two Lines of Descent. *J. Mol. Evol.* **46**, 54–63 (1998).
70.   Koga, Y. Early Evolution of Membrane Lipids: How did the Lipid Divide Occur? *J. Mol.*
Evol. **72**, 274–282 (2011).
71.   Matsumi, R., Atomi, H., Driessen, A. J. M. & van der Oost, J. Isoprenoid biosynthesis in
Archaea – Biochemical and evolutionary implications. *Res. Microbiol.* **162**, 39–52 (2011).
72.   Koonin, E. V. Archaeal ancestors of eukaryotes: not so elusive any more. *BMC Biol.* **13**,
84 (2015).
73.   Fritz, B. A., Thomas, C. B., Van Ess, P. & Yuill, T. M. Comparison of a Modified Edward-
Type Medium and a Modified SP4-Type Medium for Primary Isolation of Mycoplasma
gallisepticum (MG) from Chickens Vaccinated with the F Strain of MG. *Avian Dis.* **35**, 591–598
(1991).
74.   Low, I. E. & Eaton, M. D. Replication of Mycoplasma pneumoniae in Broth Culture. *J.*
Bacteriol. **89**, 725–728 (1965).
75.   Meur, S. k., Sikdar, A., Srivastava, N. c. & Srivastava, S. k. Rapid photometric assay of
growth of Mycoplasma mycoides subsp. capri. *J. Appl. Bacteriol.* **66**, 301–302 (1989).

76.   Bligh, E. G. & Dyer, W. J. A Rapid Method of Total Lipid Extraction and Purification. *Can.*
J. Biochem. Physiol. **37**, 911–917 (1959).
77.   Leftin, A., Molugu, T. R., Job, C., Beyer, K. & Brown, M. F. Area per Lipid and Cholesterol
Interactions in Membranes from Separated Local-Field 13C NMR Spectroscopy. *Biophys. J.* **107**,
2274–2286 (2014).
78.   Chen, P. S., Toribara, T. Y. & Warner, H. Microdetermination of Phosphorus. *Analytical*
Chemistry **28**, 1756-1758 (1956).
79.   Ejsing, C. S. *et al.* Global analysis of the yeast lipidome by quantitative shotgun mass
spectrometry. *Proc. Natl. Acad. Sci.* **106**, 2136–2141 (2009).
80.   Liebisch, G. *et al.* High throughput quantification of cholesterol and cholesteryl ester by
electrospray ionization tandem mass spectrometry (ESI-MS/MS). *Biochim. Biophys. Acta BBA -*
Mol. Cell Biol. Lipids **1761**, 121–128 (2006).
81.   Gilbert, B. R. *et al.* Generating Chromosome Geometries in a Minimal Cell From Cryo-
Electron Tomograms and Chromosome Conformation Capture Maps. *Front. Mol. Biosci.* **8**,
(2021).
82.   Kremer, J. R., Mastronarde, D. N. & McIntosh, J. R. Computer Visualization of Three-
Dimensional Image Data Using IMOD. *J. Struct. Biol.* **116**, 71–76 (1996).
1025
1026

Table 1 - Lipid Names and Abbreviations			
Lipid Class Name	**Lipid Class Abbreviation**	**Lipid Species Name**	**Lipid Species Abbreviation**
Sterol	N/A	Cholesterol	N/A
Diether Phosphatidylcholine	diether PC	1-O-hexadecanyl-2-O-(9Z-octadecenyl)-sn-glycero-3-phosphocholine	diether PC
Diether Phosphatidylglycerol	diether PG	1-O-hexadecanyl-2-O-(9Z-octadecenyl)-sn-glycero-3-phospho-(1'-rac-glycerol)	diether PG
Diether Phosphatidylethanolamine	diether PE	1-O-hexadecanyl-2-O-(9Z-octadecenyl)-sn-glycero-3-phosphoethanolamine	diether PE
Phosphatidylcholine	PC	1-palmitoyl-2-oleoyl-glycero-3-phosphocholine	POPC
		1,2-dipalmitoyl-sn-glycero-3-phosphocholine	DPPC
		1,2-dioleoyl-sn-glycero-3-phosphocholine	DOPC
Phosphatidylglycerol	PG	1-palmitoyl-2-oleoyl-sn-glycero-3-phospho-(1'-rac-glycerol)	POPG
		1,2-dipalmitoyl-sn-glycero-3-phospho-(1'-rac-glycerol)	DPPG
		1,2-dioleoyl-sn-glycero-3-phospho-(1'-rac-glycerol)	DOPG
Phosphatidylethanolamine	PE	1-palmitoyl-2-oleoyl-sn-glycero-3-phosphoethanolamine	POPE
		1,2-dipalmitoyl-sn-glycero-3-phosphoethanolamine	DPPE
		1,2-dioleoyl-sn-glycero-3-phosphoethanolamine	DOPE
Fatty Acids	FA	(9Z)-Octadec-9-enoic acid	Oleic Acid
		Hexadecanoic acid	Palmitic Acid
Enantiomeric Phosphatidylcholine	Ent PC	3-palmitoyl-2-oleoyl-sn-glycero-1-phosphocholine	Enantiomeric POPC
Phosphatidic acid	PA	1-palmitoyl-2-oleoyl-sn-glycero-3-phosphate	POPA
Cardiolipin	CL	1',3'-bis[1,2-dipalmitoyl-sn-glycero-3-phospho]-glycerol	16:0 Cardiolipin
		1',3'-bis[1-palmitoyl-2-oleoyl-sn-glycero-3-phospho]-glycerol	16:0-18:1 Cardiolipin
Sphingomyelin	SM	N/A	N/A
Lyso-Phosphatidylcholine	Lyso PC	1-oleoyl-2-hydroxy-sn-glycero-3-phosphocholine	18:1-Lyso PC
		1-palmitoyl-2-hydroxy-sn-glycero-3-phosphocholine	16:0-Lyso PC
Diacylglycerol	DAG	N/A	N/A

1027

 Figure Legends:

 Figure 1: Membrane Remodeling in M. Mycoides is Dependent on Acyl Chain Scavenging

 1 *Mycoplasma mycoides* can take up phospholipids and free fatty acids from its environment. 2
 Cleaving ester bonds from glycerophospholipids results in a pool of free fatty acids. 3 With fatty
 acids, *M. mycoides* is able to synthesize Phosphatidylglycerol (PG) with various fatty acid
 compositions, as well as remodel other phospholipid classes. 4 Cardiolipin (CL) can be synthesized
 from two PGs, starting with cleavage of PG headgroups. 5 *M. mycoides* can synthesize a variety
 of Cardiolipins from PG. 6 Cleaving acyl chains from remodeled and synthesized phospholipids
 replenishes the pool of free fatty acids. 7 Glycerophospholipids are inserted into the membrane.
 8 Cholesterol is essential for *M. Mycoides* growth.

 Figure 2: A Defined Lipid Diet Results in a Living Membrane with Two Lipids

 a *M. mycoides* can adapt to defined lipid "diets" with diester or diether phospholipids; resulting
 in simpler membranes than when grown on a complex diet (e.g. FBS). Adaptation to new diets
 occurs after 3 passages (p > 3). TLC aspect ratio adjusted for legibility; unmodified TLCs can be
 found in Figure S2. *M. mycoides* can scavenge acyl chains from POPC, yielding a more complex
 membrane from a defined diet. b Lipidomic analysis of cells grown on only oleic and palmitic acid
 shows *M. mycoides* cells can synthesize Phosphatidylglycerol and Cardiolipin species when a
 source of fatty acids is present. c Lipidomic analysis of *M. mycoides* cells grown on a POPC +
 Cholesterol diet shows acyl chain scavenging leads to the synthesis of a diversity of lipids,
 resulting in a membrane with 28 lipids from a diet of only two. d 16:0-18:1 diether PC has ether-
 linked hydrocarbon chains that cannot be cleaved by *M. mycoides*, eliminating scavenging.
 Lipidomic analysis shows living cells with two lipids comprising 99.9 mol% of their lipidome.

 Figure 3: Living Membranes with Tunable Lipid Class and Acyl Chain Complexity

 a 16:0-18:1 Diether PG diet results in the synthesis of Cardiolipin with fully ether-linked
 hydrocarbon chains, increasing head group, but not acyl chain complexity. b A diet of both
 Diether PC and PG restores the headgroup, but not acyl chain, complexity of the POPC diet. c
 When switched from one diet to another, in this case from the D.PC diet to the POPC diet, there
 is an adaptation period before acyl chain scavenging occurs. d A diet derived from a total lipid
 extract of cells grown on the POPC diet (a "transplant" diet) restores the full complexity of the

POPC. Growth on this transplanted diet was performed after the first passage (p = 1) to the new
diet, before cells begin to scavenge acyl chains from exogenous phospholipids. e Reintroducing
acyl chain complexity is more effective than headgroup complexity for rescuing growth (relative
to cells adapted to a POPC diet), but, even when full complexity is restored, acyl chain scavenging
capacity still improves growth. P-values, calculated using a two-tailed Student's-t-test, are shown
when relevant. N, number of replicates, for all growth rates can be found in table S1.
1072
1073 Figure 4: Lipidome Minimization in JCVI-Syn3A: Achieving a Minimal Membrane in a Genomically
1074 Minimal Cell
1075
a JCVI-Syn3A exhibits growth rates similar to *M. mycoides* on a minimal lipid diet. b JCVI-Syn3A is
viable when grown on a Diether-PC + Cholesterol diet with 99.34 mol% of the membrane
comprised of only two lipids. c JCVI-Syn3A cells display three phenotypes when observed with
Transmission Electron Microscopy; a normal ovoid morphology, cells with internal membranes,
and cells with intercellular tubules. Cell counts reveal the Diether-PC diet yields significantly more
cells with the internal membranes than the other diets. Cell counts can be found in Figure S6a-
S6g d A tomogram of a JCVI-Syn3A cell on the Diether-PC diet with internal membranes reveals
a large cell with multiple membrane-bound internal vesicle-like structures that have a lower
electron density compared to the rest of the cell, suggesting the internal membrane bound
structures are vesicles occurring from membrane invagination. Modeling shows these vesicles
are completely enclosed and separate from the cell surface membrane. Images in this figure are
from a single biological replicate. P-values, calculated using a two-tailed Student's-t-test, are
shown when relevant. N, number of replicates for all growth rates, can be found in table S1.
1089
1090
1091 Figure 5: Enantiomeric Lipid Diets Negatively Affect Cell Growth and Membrane Properties
1092
a Enantiomeric (Ent) POPC is a synthetic chiral POPC with the head group in the SN1 position,
rather than SN3 position. b *M. mycoides* and c JCVI-Syn3A cells exhibit slower growth when grown
on diets with enantiomeric POPC present; are more fragile to hypoosmotic shock when grown on
diets with enantiomeric POPC present; and are more permeable to non-chiral fluorescein
diacetate (FDA) when grown on diets with enantiomeric POPC present. d M. mycoides and e JCVI-
Syn3A cell growth and membrane permeability are inversely correlated. P-values, calculated
using a two-tailed Student's-t-test, are shown when relevant. N, number of replicates for all
growth rates, can be found in table S1. N=3 biological replicates for osmotic sensitivity and
permeability subfigures.
1102
1103

1104

Figure 1: Membrane Remodeling in *M. Mycoides* is Dependent on Acyl Chain Scavenging

1 *Mycoplasma mycoides* can take up phospholipids and free fatty acids from its environment. **2** Cleaving ester bonds from glycerophospholipids results in a pool of free fatty acids. **3** With fatty acids, *M. mycoides* is able to synthesize Phosphatidylglycerol (PG) with various fatty acid compositions, as well as remodel other phospholipid classes. **4** Cardiolipin (CL) can be synthesized from two PGs, starting with cleavage of PG headgroups. **5** *M. mycoides* can synthesize a variety of Cardiolipins from PG. **6** Cleaving acyl chains from remodeled and synthesized phospholipids replenishes the pool of free fatty acids. **7** Glycerophospholipids are inserted into the membrane. **8** Cholesterol is essential for *M. Mycoides* growth.

Figure 2: A Defined Lipid Diet Results in a Living Membrane with Two Lipids

a *M. mycoides* can adapt to defined lipid "diets" with diester or diether phospholipids; resulting in simpler membranes than when grown on a complex diet (e.g. FBS). Adaptation to new diets occurs after 3 passages (p > 3). TLC aspect ratio adjusted for legibility; unmodified TLCs can be found in Figure S2. *M. mycoides* can scavenge acyl chains from POPC, yielding a more complex membrane from a defined diet. **b** Lipidomic analysis of cells grown on only oleic and palmitic acid shows *M. mycoides* cells can synthe- size Phosphatidylglycerol and Cardiolipin species when a source of fatty acids is present. **c** Lipidomic analysis of *M. mycoides* cells grown on a POPC + Cholesterol diet shows acyl chain scavenging leads to the synthesis of a diversity of lipids, resulting in a membrane with 28 lipids from a diet of only two. **d** 16:0-18:1 diether PC has ether-linked hydrocarbon chains that cannot be cleaved by *M. mycoides*, eliminating scavenging. Lipidomic analysis shows living cells with two lipids comprising 99.9 mol% of their lipidome.

3a Defined diet: D.PG + Cholesterol

MEMBRANE

Chol
CL

D.PG

No scavenging

3b Defined diet: D.PC + D.PG + Cholesterol

MEMBRANE

Chol
CL

D.PG
D.PC

No scavenging

3c Defined diet: POPC + Cholesterol diet after first passage

Chol

PC

No scavenging

3d Defined diet: Membrane Transplant

Scavenging   **No scavenging**

3e _M. mycoides_ Growth Rates on Defined Diets

Figure 3: Living Membranes with Tunable Lipid Class and Acyl Chain Complexity

a 16:0-18:1 Diether PG diet results in the synthesis of Cardiolipin with fully ether-linked hydrocarbon chains, increasing head group, but not acyl chain complexity. **b** A diet of both Diether PC and PG restores the headgroup, but not acyl chain, complexity of the POPC diet. **c** When switched from one diet to another, in this case from the D.PC diet to the POPC diet, there is an adaptation period before acyl chain scavenging occurs. **d** A diet derived from a total lipid extract of cells grown on the POPC diet (a "transplant" diet) restores the full complexity of the POPC. Growth on this transplanted diet

was performed after the first passage (p = 1) to the new diet, before cells begin to scavenge acyl chains from exogenous phospholipids. **e** Reintroducing acyl chain complexity is more effective than headgroup complexity for rescuing growth (relative to cells adapted to a POPC diet), but, even when full complexity is restored, acyl chain scavenging capacity still improves growth. Error bars are mean +/- SD. * equals p-value < 0.1; ** equals p-value < 0.05; *** equals p-value < 0.01. N, number of replicates, for all growth rates was 5 or greater and can be found in Supplementary Table S2.

Figure 4: Lipidome Minimization in JCVI-Syn3A: Achieving a Minimal Membrane in a Genomically Minimal Cell

a JCVI-Syn3A exhibits growth rates similar to *M. mycoides* on a minimal lipid diet. **b** JCVI-Syn3A is viable when grown on a Diether-PC + Cholesterol diet with 99.34 mol% of the membrane comprised of only two lipids. **c** JCVI-Syn-3A cells display three phenotypes when observed with Transmission Electron Microscopy; a normal ovoid morphology, cells with internal membranes, and cells with intracellular tubules. Cell counts reveal the Diether-PC diet yields significantly more cells with the internal membranes than the other diets. Cell counts can be found in Figure S6a-S6g **d** A tomogram of a JCVI-Syn3A cell

on the Diether-PC diet with internal membranes reveals a large cell with multiple membrane-bound internal vesicle-like structures that have a lower electron density compared to the rest of the cell, suggesting the internal membrane bound structures are vesicles occurring from membrane invagination. Modeling shows these vesicles are completely enclosed and separate from the cell surface membrane. Error bars are mean +/- SD. * equals p-value < 0.1; ** equals p-value < 0.05; *** equals p-value < 0.01. N, number of replicates for all growth rates, was 5 or greater and can be found in Supplementary Table S2.

Figure 5: Enantiomeric Lipid Diets Negatively Affect Cell Growth and Membrane Properties

a Enantiomeric (Ent) POPC is a synthetic chiral POPC with the head group in the SN1 position, rather than SN3 position. **b** *M. mycoides* and **c** JCVI-Syn3A cells exhibit slower growth when grown on diets with enantiomeric POPC present; are more fragile to hypoosmotic shock when grown on diets with enantiomeric POPC present; and are more permeable to non-chiral fluorescein diacetate (FDA) when grown on diets with enantiomeric POPC present. **d** *M. mycoides* and **e** JCVI-Syn3A cell growth and membrane permeability are inversely correlated. Error bars are mean +/- SD. * equals p-value < 0.1; ** equals p-value < 0.05; *** equals p-value < 0.01. N, number of replicates, for all growth rates was 5 or greater and can be found in Supplementary Table S2. N=3 for osmotic sensitivity and permeability subfigures.

Figure 6: Enantiomeric Lipids Have no Significant Effect on Lipid Order

Lipid vesicles reconstituted from total lipid extracts of *M. mycoides* cells had a similar lipid order (measured as the general polarization index of c-laurdan) for all three diets with POPC enantiomers. Error bars are mean +/- SD. * equals p-value < 0.1; ** equals p-value < 0.05; *** equals p-value < 0.01. N=3