Purine nucleosides replace cAMP in allosteric regulation of PKA in trypanosomatid pathogens

Veronica Ober[1#], George B. Githure[1#], Yuri Volpato Santos[1#], Sidney Becker[2, 3], Gabriel Moya[1], Jerôme Basquin[4], Frank Schwede[5], Esben Lorentzen[6], Michael Boshart[1]*

1 Faculty of Biology, Genetics, Ludwig-Maximilians University Munich (LMU), 82152 Martinsried, Germany

2 Max Planck Institute of Molecular Physiology, Dortmund, 44227 Germany

3 TU Dortmund, Department of Chemistry and Chemical Biology, 44227 Dortmund, Germany

4 Max Planck Institute for Biochemistry, 82152 Martinsried, Germany

5 BIOLOG Life Science Institute GmbH & Co. KG, 28199 Bremen, Germany

6 Department of Molecular Biology and Genetics, Aarhus University, Aarhus, Denmark

[#] equal contributions of authors

* corresponding author: Prof. Michael Boshart boshart@lmu.de

Abstract

Cyclic nucleotide binding domains (CNB) confer allosteric regulation by cAMP or cGMP to many signalling proteins, including PKA and PKG. PKA of phylogenetically distant *Trypanosoma* is the first exception as it is cyclic nucleotide-independent and responsive to nucleoside analogues (Bachmaier et al. 2019). Here we show that natural nucleosides inosine, guanosine and adenosine are nanomolar affinity CNB ligands and activators of PKA orthologs of the important tropical pathogens *T. brucei*, *T. cruzi* and *Leishmania.* The sequence and structural determinants of binding affinity, -specificity and kinase activation of PKAR were established by structure-activity relationship (SAR) analysis, co-crystal structures and mutagenesis. Substitution of 2-3 amino acids in the binding sites is sufficient for conversion of CNB domains from nucleoside to cyclic nucleotide specificity. In addition, a trypanosomatid-specific C-terminal helix (αD) is required for high affinity binding to CNB-B. The αD helix functions as a lid of the binding site that shields ligands from solvent. Selectivity of guanosine for CNB-B and of adenosine for CNB-A results in synergistic kinase activation at low nanomolar concentration. PKA pulldown from rapid lysis establishes guanosine as the predominant ligand *in vivo* in *T. brucei* bloodstream forms, whereas guanosine and adenosine seem to synergize in the procyclic developmental stage in the insect vector. We discuss the versatile use of CNB domains in evolution and recruitment of PKA for novel nucleoside-mediated signalling.

Introduction

Protein kinase A (PKA) is a prototype kinase first purified from rabbit skeletal muscle in 1968 (Walsh, Perkins, and Krebs 1968). More than 40 years of trailblazing biochemical and structural work elucidated the mechanism of allosteric activation by cAMP, providing a paradigm of allosteric regulation (Taylor et al. 2021). Inactive PKA is a dimeric or tetrameric complex of regulatory (R) and catalytic (C) subunits, depending on the species. Upon activation, two molecules of cAMP bind to two cyclic nucleotide binding domains (CNB) arranged in tandem in the C-terminal part of the regulatory subunit(s). Cyclic AMP binding to the C-terminal CNB-B initiates a conformation change that opens up the adjacent CNB-A for a second cAMP molecule whose binding completes the conformational transition that liberates the C-subunit from the holoenzyme complex (Kim et al. 2007). The free C subunit is thereby released from autoinhibition and activated (Kim et al. 2007; Taylor et al. 2012). PKA is a highly conserved kinase present in all eukaryotes except plants, functioning in diverse signalling processes ranging from metabolic regulation and hormone action to cell differentiation and synaptic long term potentiation underlying memory (Lee et al. 2021). In protists and fungi the predominant functions are response to carbon source changes and regulation of developmental transitions, infectivity or sexual dimorphism (Perrin et al. 2020; Hitz et al. 2021; Uboldi et al. 2018; Jia et al. 2017; Choi, Jung, and Kronstad 2015; Kim et al. 2021; Vaidyanathan et al. 2014). Regulation of PKA by cAMP was universally found (Rinaldi et al. 2010; Haste et al. 2012; Kurokawa et al. 2011; Taylor et al. 2012), which is why PKA and cAMP-dependent protein kinase are used as synonyms. Furthermore, cAMP dependence was assumed and widely cited in reviews for PKAs of species for which uncontested biochemical evidence is lacking, including protozoan flagellates of the phylogenetically distant order *Kinetoplastida*. It was therefore of greatest interest that PKA in *Trypanosoma brucei*, a parasitic and pathogenic member of the *Kinetoplastida* was found unresponsive to cAMP even at high intracellular concentrations (Bachmaier et al. 2019;

Bubis et al. 2018). Compound screening then identified 7-deazapurine nucleoside antibiotics

as nanomolar activators of PKA in *T. brucei* (Bachmaier et al. 2019).

Trypanosoma brucei, *Trypanosoma cruzi* and *Leishmania sp.* are related trypanosomatid

pathogens causing the deadly neglected tropical diseases sleeping sickness, Chagas

disease and leishmaniosis, respectively. In addition, *Trypanosoma* is responsible as animal

pathogen for important economic losses and impedes social development in affected

countries. These organisms are famous for discovery of many exotic biochemical and

genetic mechanisms (Matthews 2015), and not surprisingly, signaling systems diverge from

other model organisms and few pathways are on the way to be assembled (Matthews 2021).

Nevertheless, cAMP seems to play an important role: *T. brucei* encodes > 80 adenylate

cyclase genes (Salmon, Bachmaier, et al. 2012) that are important for host innate immunity

subversion (Salmon, Vanwalleghem, et al. 2012) and for development of the parasite in its

insect vector (Bachmaier et al. 2022). These pathways are obviously uncoupled from the

cAMP unresponsive PKA and seem to use alternative and novel cAMP effectors (Bachmaier

et al. 2023). PKA in these parasites is essential and important for cell division (Bachmaier et

al. 2019; Baker et al. 2021; Cayla et al. 2022). In *T. brucei* PKA has been identified as

candidate member of a quorum sensing pathway and implicated in stage development (Mony

et al. 2014; Toh et al. 2021). The genome of *T. brucei* encodes one regulatory and three

catalytic subunits that all have syntenic orthologs in *T. cruzi* and *Leishmania*. The upstream

pathway(s) regulating PKA in these organisms have not been identified. However, the high

degree of conservation of the CNBs in TbPKA and their high affinity binding to nucleoside

analogues (Bachmaier et al. 2019) suggested the existence of an alternative second

messenger. PKG, a related AGC kinase, is also subject to allosteric regulation by cyclic

nucleotides, responding to cyclic guanosine monophosphate (cGMP) instead of cAMP

(Huang, Kim, et al. 2014; Kim and Sharma 2021). The high structural similarity of the CNB

domains in PKA and PKG spurred attempts to define determinants of cyclic nucleotide

binding selectivity. Amino acids that contribute to selectivity have been identified (Lorenz et

al. 2017; Shabb et al. 1991; Shabb, Ng, and Corbin 1990; Huang, Gerlits, et al. 2014; Weber,

Shabb, and Corbin 1989; Corbin et al. 1986; Kim and Sharma 2021; Lorenz, Bertinetti, and

Herberg 2017), but establishing a consensus of key determinants has been challenging.

Differences in ligand specificity of PKAR between a pathogen and its host provides an

opportunity for much needed drug development to fight neglected vector borne diseases

caused by *Trypanosomatida*. Hence, identification of physiological ligands of trypanosomatid

PKA and definition of their binding selectivity were important goals.

Here we show that purine nucleosides exhibit nanomolar affinity for the PKA regulatory

subunits of these pathogens and activate the kinases. We define the minimal changes that

convert a nucleoside-specific CNB to cyclic nucleotide specificity. Furthermore, we see site-

selective binding and synergy between guanosine and adenosine, compatible with binding *in*

vivo to PKAR. Expansion of the ligand portfolio of CNBs in evolution has enabled

repurposing of PKA for a different signalling pathway, while maintaining the sophisticated

allosteric activation mechanism triggered by ligand binding to PKA.

Results

Nucleosides are direct activators of PKA in trypanosomes

We recently identified the nucleoside analogue 7-cyano-7-deaza-inosine (7-CN-7-C-Ino,

Jaspamycin) and related compounds like Toyocamycin as potent activators of the cAMP-

independent PKA of *Trypanosoma* (Bachmaier et al. 2019). Attempts to bioinformatically

detect pathways for synthesis of these nucleoside antibiotics in trypanosomatids have been

unsuccessful. Therefore, we considered unmodified purine nucleosides and studied the

structure-activity-relationship (SAR) for kinase activation (Fig. 1a and Table 1). Tagged R-

and C1-subunits of *T. brucei* PKA were co-expressed in *Leishmania tarentolae* and

stoichiometric holoenzyme complexes were tandem-affinity purified (Supplementary Fig. 1a).

The tandem-affinity purification to near homogeneity guaranteed removal of any

heterologous complexes formed with endogenous PKA subunits of the expression system.

EC_{50} values for kinase activation were determined from dose response assays (Fig. 1b,

Table 1, Supplementary Fig. 1a). Surprisingly, inosine was the most potent activator (EC_{50} 14

nM). We did not expect this result as the structure of TcPKAR bound to 7-CN-7-C-Ino (PDB:

6FTF) and computational docking of 7-deaza analogues had suggested an important role of

the cyano group at position 7 of the purine ring (Bachmaier et al. 2019). The SAR analysis

(Fig. 1a, Table 1, Supplementary Fig. 1a) showed the oxygen at position 6 in the purine ring

to be particularly important, as nebularine, lacking a 6-substitution, was 186-fold less potent

than inosine. An amino group substitution at position 6 (adenosine) resulted in a further 2-

fold drop in potency. An amino group at position 2 (guanosine) caused 11-fold and a keto

group in this position (xanthosine) a >4400-fold lower activation potency, respectively. A

structural isomer of inosine (allopurinol riboside) with restricted delocalized π-electron

system showed 120-fold reduced activation. The structure of TcPKAR bound to 7-CN-7-C-

Ino (Bachmaier et al. 2019) predicted an important role of the ribose moiety that is

accommodated deep in the binding pocket. The 2´-, 3´- and 5´-deoxy derivatives of

adenosine confirmed essential roles for all three hydroxyl groups of the ribose ring (Table 1,

Supplementary Fig. 1a). Inosine and guanosine 5′-monophosphates were >7700-fold less

potent than the respective nucleosides and AMP did not activate even at 5 mM. Cyclic GMP

activated the kinase in the upper micromolar range, whereas cAMP and cIMP were inactive

up to 5mM. Pyrimidine nucleosides uridine and cytidine were 3-4 orders of magnitude less

potent than purine nucleosides (Table 1, Supplementary Fig. 1a). In summary, the natural

nucleoside inosine is only 2-fold less potent as activator of TbPKAR than the nucleoside

analogue activator 7-CN-7-C-Ino (6.5 nM)(Bachmaier et al. 2019) but 5-fold more potent than

cAMP activation of the recombinant mammalian PKARIα2-2PKACα holoenzyme purified

from *E. coli* (Fig. 1c, Table 1). Therefore, purine nucleosides qualify as possible physiological

activators of TbPKA in trypanosomes.

Nucleoside activation of PKA in kinetoplastid pathogens

Next, we asked if activation by nucleosides and complete insensitivity to cAMP is a unique

feature of *T. brucei* PKA or a shared feature in the protozoan class of Kinetoplastida. We

selected the medically important *Trypanosoma cruzi* and *Leishmania donovani* as

representatives of this group. Orthologous regulatory subunits (TcPKAR and LdPKAR1) and

catalytic subunits (TcPKAC2 and LdPKAC1), respectively, were tagged and co-expressed in

Leishmania tarentolae, and holoenzyme complexes were tandem affinity purified

(Supplementary Fig. 1b, c). The kinase assay dose responses (Fig. 1c, Supplementary Fig.

1b, c) show that inosine is the most potent of the tested nucleosides for all analysed species,

whereas no activation was observed with cAMP, even at very high concentrations (Fig. 1c,

Table 1). Activation potency of inosine or guanosine was between 3-fold and 23-fold lower

for *Leishmania* and *T. cruzi*, compared to *T. brucei*. The mammalian RIα2:Cα2 holoenzyme,

included as control, was activated by cAMP with an EC_{50} of 75 nM in agreement with

Herberg, Taylor, and Dostmann (1996), but was completely insensitive to inosine. We

conclude that the PKAs of *T. cruzi* and *L. donovani* are also cAMP-independent nucleoside

activated kinases. The same order of potency was found among the 3 tested nucleosides,

adenosine being the weakest activator (EC_{50} ~6-8 µM) of PKA in the three parasite species

(Supplementary Fig. 1b-c, Table 1).

Binding of nucleosides to kinetoplastid PKAs

To further investigate purine nucleoside-specific allosteric regulation of kinetoplastid PKAs,

nucleoside binding parameters were determined for the isolated R-subunits. The N-terminally

truncated PKARs of the respective species containing the two tandem cyclic nucleotide

binding domains (CNBs) were expressed in *E. coli* and purified. Initially, binding assays with

natively purified PKAR were inconclusive and highly variable using several methods. We

concluded that the purified PKAR was at least partially bound by ligands or metabolites

derived from *E. coli*, similar to the mammalian PKAR subunit that tightly binds cAMP when

purified from bacteria (Buechler, Herberg, and Taylor 1993). To confirm this directly,

TbPKAR or HsPKARIα purified from *E. coli* were boiled to denature protein and were

separated by centrifugation. Supernatants containing released ligands were collected and

tested in kinase assays with purified holoenzyme as before (Supplementary Fig. 2). The

supernatant of the boiled HsPKARIα fully activated the mammalian holoenzyme, but not the

T. brucei holoenzyme, as would be expected for cAMP. In contrast, the supernatant from

boiled TbPKAR fully activated the *T. brucei* holoenzyme but not the mammalian one

(Supplementary Fig. 2b). In the HsPKARIα-derived supernatant only cAMP was detected by

LC-MS (Supplementary Fig. 2d), whereas in the TbPKAR-derived supernatant nucleosides

(predominantly inosine) were detected (Supplementary Fig. 2c). This experiment qualitatively

showed tight binding of nucleosides to TbPKAR in *E. coli*. Subsequently, we routinely

denatured the purified His-tagged regulatory subunits to remove any pre-bound ligands (see

Methods). Refolding conditions were optimized by a buffer screen and monitored by

differential scanning fluorimetry (nanoDSF) (Niesen, Berglund, and Vedadi 2007) and size

exclusion chromatography (Supplementary Fig. 3a-d). The thermal stability of proteins was

determined by nanoDSF that records changes of the ratio of intrinsic fluorescence at two

wavelengths (330 and 350nM). Natively purified TbPKAR(199-499) unfolded at T_m of 59,5°C.

When refolded in absence of ligand (APO form) the T_m was only 42,3°C (Supplementary Fig.

3b). This is interpreted as stabilization of the purified TbPKAR by its partial loading with

nucleosides from *E. coli*. Indeed, upon saturating the ligand-bound state of the refolded and

the natively purified protein preparations by addition of excess inosine, the T_m raised to 68°C

for both. The identical T_m strongly indicates correct folding after renaturation. Ligand-

depleted LdPKAR1(200-502) was prepared in the same way (Supplementary Fig. 3c-d),

whereas for TcPKAR the yield of refolded protein (≤ 2μM) was too low to carry out further

experiments. Isothermal titration calorimetry (ITC) measurements showed high affinity

binding of inosine and guanosine to both TbPKAR (Fig. 1d) and LdPKAR1 (Fig.1e) with

nanomolar K_D values (Supplementary Table 2), matching closely the EC_{50} values for kinase

activation (Table 1). The close match of binding K_D and activation EC_{50} values for inosine

and guanosine suggests that the K_D measured for PKAR expressed in *E. coli* is an excellent

proxy for the binding K_D to the holoenzyme. Adenosine, the weakest activator, is also the

weakest binder. Inosine did not bind at all to human PKARIα, which bound cAMP with a K_D of

23 nM (Supplementary Fig. 1d). LdPKAR1 did not bind to cAMP (Supplementary Fig. 1d) as

shown before for TbPKAR (Bachmaier et al. 2019). The binding data thus support

nucleoside-specificity and cAMP independence of the trypanosomatid PKAs. The

stoichiometry of purine nucleoside binding to TbPKAR as calculated from ITC data was N ≈1,

apparently lower than expected for the two binding sites occupied by inosine in the co-crystal

structures (see below). We cannot exclude the possibility that a fraction of the refolded

protein unfolds or aggregates after purification or is bound to remaining traces of the ligand

and therefore not available for binding at the time of ITC analysis (see Methods). The

thermodynamic signature of nucleosides binding to kinetoplastid PKA resembles that of

mammalian PKA bound by cAMP (Supplementary Fig. 5). The enthalpic contribution to

binding (ΔH), indicating strong hydrogen bonding, is counteracted by a relatively large loss of

entropy (TΔS), indicating bound-state conformational constrains.

Structure of the nucleoside-binding pockets

To evaluate the binding mode, we solved the crystal structures of *T. cruzi* PKAR(200-503)

and *T. brucei* PKAR(199-499) bound to inosine at 1.4 Å and 2.1 Å resolution, respectively

(Supplementary Fig. 3f and g, Supplementary Table 1). Attempts to crystallize

LdPKAR1(200-502) were unsuccessful. The structures of TbPKAR and TcPKAR show high

overall similarity. Calculated RMSD of Cα alignment was 0.796 Å for the entire proteins and

0.281/0.342 Å for CNB-A and CNB-B, respectively. Residues that contribute to high affinity

binding by interacting with the ribose moiety of inosine are identical in both structures (Fig. 2

a, b; Supplementary Fig. 3o; Supplementary Movie 1) and reside in a segment that we

denoted ribose binding cassette (RBC), in analogy to the phosphate binding cassette (PBC)

nomenclature for mammalian PKA (Canaves and Taylor 2002). These residues in site A

(308-320$^{\text{TbPKAR}}$, 309-321$^{\text{TcPKAR}}$) and site B (432-445$^{\text{TbPKAR}}$, 433-446$^{\text{TcPKAR}}$) engage in the same

interactions with inosine as in our previously described co-crystal structure of TcPKAR(200-

503) with 7-CN-7-C-Ino (PDB: 6FTF) (Bachmaier et al. 2019). Likewise, the "capping" by π-

stacking with the purine ring in both sites (Y371/483$^{\text{TcPKAR}}$, Y370/482$^{\text{TbPKAR}}$) and the

interacting residues in the lid-like αD helix are conserved. Thus, the binding mode of inosine

and 7-CN-7C-Ino is almost identical. Minor differences are compatible with similar affinities of

inosine and 7-CN-7-C-Ino. In site A amino acid K294 donates a hydrogen bond to the cyano

group of 7-CN-7-C-Ino (Fig. 2c). When bound to inosine, however, a different side chain

rotamer of K294 is preferred, and a hydrogen bond can now be formed with the keto group in

position 6 of the purine ring (Fig. 2c). In site B the bulkiness of the cyano group displaces the

side chain of Y485 by 1.5 Å, creating a small hydrophobic pocket able to fit C7-derivatives

(Bachmaier et al. 2019) (Fig. 2c). Comparison of mammalian cAMP-bound PKARIα

(PDB:1RGS) with nucleoside-bound kinetoplastid structures (PDB: 6FLO) clearly suggests

that residues A202/R209$^{\text{PKARIα}}$ in site A and A326/R333$^{\text{PKARIα}}$ in site B are key to explain

the altered ligand specificity of the kinetoplastid PKAR subunits (Fig. 2d). The arginine

residues 209/333$^{\text{PKARIα}}$ conserved in most PKARs are replaced by polar amino acids, and the

alanine residues 202/326$^{\text{PKARIα}}$ are replaced by glutamates highly conserved in the

kinetoplastid PKARs. The arginine in PKARIα neutralizes the negative charge of the

phosphate in cAMP and also donates hydrogen bonds to the exocyclic oxygens of the cyclic

phosphate. The glutamates in kinetoplastid RBCs interacts with the 3' and 5' OH groups of

ribose. Moreover, the arginines 209/333$^{PKARI\alpha}$ and the glutamates 311/435^{TbPKAR} occupy the

same spatial position in the structures (Fig. 2d). A superposition of the mammalian and *T.*

brucei structures shows a clash between the phosphate group of cAMP and the negatively

charged side chain of E311/435^{TbPKAR} (Fig. 2d insets). The high-resolution crystal structures

of *T. brucei* PKAR and *T. cruzi* PKAR thus provides a molecular rationale for absence of

binding and activation by cAMP.

Synthetic conversion of TbPKAR to cyclic nucleotide specificity

To identify the structural determinants of ligand specificity, we introduced site-directed amino

acid changes in TbPKAR to restore binding and activation by cyclic nucleotides. Three

residues in each of the binding sites were mutated: E311A, T318R, V319A in RBC-A and

E435A, N442R, V443A in RBC-B (mutant 1 in Table 2). In addition to the arginines and

glutamates discussed above, a third position (V319/V443) that differs in kinetoplastid PKA

compared to other eukaryotic PKAs (Mohanty et al. 2015; Bachmaier et al. 2019) was

changed to alanine. The consensus PBC sequence of cAMP dependent PKAs (Canaves and

Taylor 2002) was thereby restored. Mutant PKAR subunits were co-expressed with *T. brucei*

catalytic subunits in *Leishmania tarentolae*, and holoenzymes were tandem affinity purified.

Kinase activation by nucleosides and cyclic nucleotides was measured (Table 2,

Supplementary Fig. 4). Mutant 1 restored kinase activation by cIMP (EC$_{50}$ 340 nM) and

reduced activation potency of inosine >21.000-fold compared to WT. Thus, we confirmed

that replacing these key residues was sufficient for conversion to cyclic nucleotide specificity.

Activation by cAMP was also restored, but at lower activation potency (EC$_{50}$ 33 µM). This

corresponds to lower activation potency of adenosine compared to inosine for wild type

TbPKA. The same ranking is also seen for the very low potencies of IMP, GMP and AMP

(Tables 1 and 2). To confirm the binding mode of cAMP to the converted binding site, mutant

6 of TbPKAR(199-499) carrying the triple replacements in site A was expressed in *E. coli* and

co-crystallized with cAMP (Supplementary Fig. 3h, Supplementary Table 1). A molecule of

cAMP was bound to site A and an inosine molecule (captured during expression in *E. coli*) to

the unmodified site B (Fig. 3a). Structural similarity of mutant 6 to wild type TbPKAR was

very high (Cα RMSD = 0.430 Å). All hydrogen bonds to cAMP observed in the PKARIα

structure (PDB: 1RGS, Fig. 3b, right) were also present between cAMP and homologous

residues in the A-site pocket of TbPKAR mutant 6 (Fig. 3b, left and Supplementary Movie 2).

The only remarkable difference is that cAMP binds in an *anti*-conformation in mutant 6 and in

the *syn*-conformation in the mammalian PKAR. The E311A and V319A replacements created

additional space inside the pocket to accommodate the bulky phosphate group of cAMP. An

altered conformer of cysteine 278 and slight displacement of the loop between β-2 and β-3 in

site-A allowed R318 to be accommodated so that it can interact with an exocyclic oxygen of

cAMP (Supplementary Movie 2).

Simultaneous binding of cAMP and inosine to mutant 6 was supported by nanoDSF analysis.

The refolded protein (APO form) had a low T_m measured by nanoDSF, but T_m raised by 14°C

upon addition of cAMP and by 28°C upon addition of both cAMP and inosine (Fig. 3c). cAMP

also stabilized the refolded mutant 6 protein during purification (Supplementary Fig. 3i, j).

Correct refolding of this mutant was indicated by identical T_m after addition of cAMP plus

inosine to native and refolded protein preparations and was confirmed by circular dichroism

spectroscopy (Supplementary Fig. 3e). To evaluate the role of individual amino acids in the

"conversion set", single and double mutations were introduced at equivalent positions in RBC

A and RBC B of TbPKAR and co-expressed with the *T. brucei* catalytic subunit PKAC1 in

Leishmania tarentolae. The tandem affinity purified holoenzymes were used for kinase

assays to determine EC_{50} values (Table 2, Fig. 3d, Supplementary Fig. 4). Replacement of

positions 318/442[TbPKAR] by arginines in both RBCs (mutant 4) was sufficient for response to

cIMP (EC_{50} 24 µM). To achieve cyclic nucleotide activation in the upper nM range the

glutamates 311/435 needed to be replaced by alanines as well in mutant 5 (Fig. 3d, Table 2).

The potency of inosine was 1556-fold reduced by E311A/E435A[TbPKAR] (mutant 2) alone. In

contrast, substituting the non-conserved amino acids at positions 318/442 by arginine

(mutant 4) or a valine (mutant 3) did not have a significant effect on inosine response. The

adjacent valine 319/443^{TbPKAR} seems to contribute to activation by inosine 17-fold (compare

mutants 1 and 5, Tab. 2 and Fig. 3d). In the B site this valine engages in hydrophobic

interactions with the side chains of Y485 and K488, both belonging to the kinetoplastid-

specific αD helix that supports inosine binding by sealing the binding pocket (Supplementary

Movie 3). The EC$_{50}$ values of all mutants tested were almost 100-fold higher for cAMP than

for cIMP, not surprising as inosine is a much better activator of the WT protein compared to

adenosine (Tables 1, 2).

The αD helix is required for high affinity binding to the B-site

The binding and activation assays used so far average K$_D$ and EC$_{50}$ values over both the A-

site and B-site of TbPKAR. As these sites are structurally not identical, we considered a

kinetoplastid-specific feature of the B-site, the αD-helix (Bachmaier et al. 2019). This is a

helical extension of αC beyond the small loop containing the capping residue Y482 at the

end of αC that stacks with the purine ring of inosine (Fig. 4a). In the ligand bound structures

of TbPKAR and TcPKAR, this helix docks to the β-barrel of the B-site, covers the binding

pocket and shields the ligand from solvent (Fig 4a, Supplementary Movie 4). Only one water

molecule was found inside the binding pocket (Fig. 4a). Two tyrosine residues (Y484, Y485)

in αD are conserved in trypanosomatids, as are amino acids in the beta barrel of the B-site

that are linked to these two amino acids via hydrogen bonds (Fig. 4b, c). Y484 interacts with

the backbone of V443 and R413 while Y485 forms two hydrogen bonds to the side chains of

N438 and H440 (Fig. 4b). *In silico* structure relaxation (Fig. 4d) under an OPLS3 force field

(Maestro-Schrödinger^{TM}) showed two histidines (H440, H430) engaging in salt bridge

interactions with the ribose-binding glutamates (E433 and E435), together forming a stable

structure on which the αD-helix can dock (Fig. 4d). The beta factor representation of

TbPKAR suggests that proline 480 functions as a hinge between αC and αD, as the average

displacement of P480 is higher than that of the other residues around it, likely correlating with

higher mobility (Fig. 4e). The αD-helix might therefore function as a lid to close the pocket and determine ligand affinity. This hypothesis was confirmed by ITC measurements of inosine binding to the Y484A/Y485A double mutant (mutant 8). To selectively measure nucleoside binding to the mutated Y484A/Y485A B-site, the A-site was made unavailable for nucleoside binding by using TbPKAR mutant 6 as context and refolding of the protein in the presence of cAMP (Supplementary Fig. 3 k). Mutant 8 shows 82-fold decreased affinity for inosine (Fig. 4f, Supplementary Table 2). The αD is therefore important for high binding affinity of 6-oxopurine nucleosides to the B-site.

Site-selective binding and synergism of nucleosides

The specific role of the αD-helix in site B prompted us to investigate binding affinities and ligand specificity of both sites individually. As interdomain CNB-A to CNB-B contacts are important for the allosteric activation mechanism of mammalian PKAR (Akimoto et al. 2015; Berman et al. 2005; Kim et al. 2007; Malmstrom et al. 2015), we analysed the contribution of each site in the R-subunit context. Mutant 6 with site A converted to cyclic nucleotide specificity (E311A, T318R, V319A) was blocked by excess of cAMP during refolding and was used to measure nucleoside binding to the non-mutated site B (Fig. 4g, Supplementary Table 2). The corresponding mutant 7 with site B converted to cyclic nucleotide specificity was refolded in presence of cIMP to block this site and allow measurements of nucleoside binding affinity to the non-mutated A-site (Fig. 4h, Supplementary Table 2). Correct refolding of mutant 6 and 7 was monitored by size exclusion chromatography and comparison of native and refolded protein by nanoDSF (Fig. 3c; Supplementary Fig. 3j, l-n). Inosine bound to mutant 6 and 7, and to the wildtype protein with similar high affinity (K_D 9-18 nM). Adenosine had two orders of magnitude lower affinity for site B than inosine (K_D 1.1 μM) but displayed high affinity for site A (K_D 82 nM). In contrast, guanosine bound with highest affinity of all nucleosides to site B (K_D 6 nM), but with 36-fold lower affinity to site A (K_D 221 nM). The preference of mutant 7 for cIMP over cAMP also reflects the B-site specificity for 6-oxo purines. *In silico* docking of inosine, guanosine and adenosine to site A and B of TbPKAR

(PDB: 6FLO, chain B) using GLIDE (Friesner et al. 2004) provided an explanation for the much lower binding affinity of adenosine to the B-site. Differences in interaction of the respective purine bases with the αD helix (Supplementary Fig. 6) include the hydrogen bonds of guanosine and inosine via the keto group at the C6 position to the backbone nitrogen of Y485. Since adenosine has an amino group in place of the C6 keto group, this specific interaction cannot take place. On the other hand, the C2 amino group of adenosine interacts with the keto group of N481 but apparently this interaction is not equivalent. Docking also suggests that the αD helix connects to guanosine, inosine and adenosine via three, two and one hydrogen bonds, respectively (Supplementary Fig. 6) which is perfectly compatible with weaker binding of adenosine to the B-site (Fig. 4g). In the more solvent exposed site A, smaller differences in binding affinity of the three nucleosides (Fig. 4h) correspond to smaller differences in the Glide G scores (Supplementary Fig. 6). In summary, molecular docking is compatible with the ITC data showing 37-fold binding selectivity of guanosine over adenosine at the B-site and 13-fold binding selectivity of adenosine over guanosine at the A-site (Fig. 4g, h). An important implication of the site-selective binding of adenosine and guanosine is their possible synergism in kinase activation. This hypothesis was directly tested in kinase assays by determining the dose-response for adenosine in the presence of guanosine concentrations far below its EC_{50}. As seen in Fig. 5a, the dose response curves were clearly left shifted (up to 20-fold) by guanosine addition. Thus, adenosine can activate TbPKA in the nanomolar range upon co-stimulation by very low concentrations of guanosine.

Allosteric kinase activation

Comparing ligand binding data (K_D, Supplementary Table 2, Fig. 1d) and kinase activation EC_{50} (Table 2, Fig. 1b) of WT TbPKA we noticed that K_D and EC_{50} values matched very well for inosine and guanosine that bind with high affinity to the B-site, whereas a 6-fold weaker activation compared to binding is seen for adenosine that preferentially binds to the A-site. This indicates that the B-site is the gate keeper and that our data are compatible with the model of allosteric regulation established for mammalian PKA (Rehmann, Wittinghofer, and

Bos 2007; Kim et al. 2007), where a conformational change upon B-site binding gives access

to the ligand at the A-site. We then calculated the EC_{50}/K_D ratio (Fig. 5b) for mutants 6 and 7

with either the B-site or the A-site intact. Both mutants show a high (>100) EC_{50}/K_D ratio for

nucleosides. This indicates that both binding domains are required in a ligand-bound

conformation for efficient allosteric kinase activation by release of the catalytic subunit. The

extreme (> 4500) EC_{50}/K_D ratio for adenosine and mutant 6 corresponds to the low affinity of

adenosine to the B-site and confirms the role of that site in initiating the conformation

change. The EC_{50}/K_D value of only 11 for mutant 6 and inosine is interpreted as weak binding

of inosine to the mutated A binding site that we cannot exclude as the site is not blocked by

cAMP in the kinase assays. Together, analysis of single binding site mutants and different

nucleoside ligands in the context of the full length PKA provides strong support for

conservation of an allosteric activation mechanism triggered by cooperative binding in

hierarchical order, initiated by B-site binding.

Ligands of trypanosome PKA *in vivo*

The biochemical and structural characterization of purified TbPKAR did not address the

relative importance of the natural purine nucleosides for kinase activation *in vivo* in

trypanosomes. The initial focus on inosine was due to the abundance of this nucleoside in *E.*

coli and preloaded recombinant protein. We then quantified the loading of tagged PKAR with

ligands upon rapid pulldown from *T. brucei* lysates. We expected at least a fraction of PKAR

to be loaded with ligands due to the dynamic equilibrium between dissociated, ligand-bound

and C-subunit-bound (holoenzyme complex) state. Tagged TbPKAR, but not endogenous

TbPKAR was efficiently pulled down via the tag. Indeed, C subunits were co-purified,

indicating only partial holoenzyme dissociation (Supplementary Fig. 7). Blood stream forms

(BSF) and the procyclic fly vector stage of *T. brucei* (PCF) expressing tagged TbPKAR were

used in these experiments and compared to matched isogenic wild type and Δ*pkar/*Δ*pkar*

knock out controls. Nucleosides were released from PKAR bound to beads by boiling, then

quantified by mass spectrometry using stable isotope labelled internal standards (Fig. 6,

Supplementary Fig. 7). In the procyclic stage (PCF) the relative amounts of nucleosides

detected in the bead fraction were 63% adenosine, 30% guanosine and 7% inosine (Fig. 6a,

Supplementary Fig. 7a). In the bloodstream stage (BSF) 94% guanosine, 6% inosine and

only background level of adenosine were found (Fig. 6b, Supplementary Fig. 7b). A priori the

MS method did not exclude the additional presence of an unknown endogenous ligand of

TbPKAR in trypanosomes. However, careful searches of the MS data sets for all known

modified nucleosides detected in living systems from the MODOMICS database (Boccaletto

et al. 2018) did not return significant hits absent in the blank (Supplementary data 1). Thus,

we propose that the nucleosides guanosine, adenosine and possibly inosine are endogenous

ligands and likely activators of trypanosomatid PKA, probably acting synergistically.

Discussion

PKA was the first protein kinase studied at the structural and mechanistic level and became a paradigm for allosteric kinase regulation by ligands (Taylor et al. 2012; Taylor et al. 2021). It is highly conserved through evolution, including its activation by cAMP and present in most species, except plants. Here we show that in the phylogenetically distant protozoan group *Trypanosomatida* nucleosides have replaced cyclic nucleotides as ligands of PKA. Inosine, guanosine and adenosine bind with high affinity to the regulatory subunit PKAR and efficiently activate PKA of *Trypanosoma brucei*, *Trypanosoma cruzi* and *Leishmania spp*. Site-selective binding affinities and synergism of guanosine and adenosine suggest a new second messenger signalling pathway or nucleoside sensing mechanism in *Trypanosomatida.*

The CNB domain - a versatile ligand binding domain

In trypanosomatid PKAR orthologs, few residues in each CNB domain systematically deviate from the consensus of the cyclic nucleotide binding motif (Mohanty et al. 2015; Canaves and Taylor 2002). By mutagenesis of these residues (E311, T318, V319 in RBC A and E435, N442, V443 in RBC B of TbPKA) we were able to restore binding and kinase activation by cyclic nucleotides and structurally interpret the determinants of altered ligand specificity. Whereas the crystal structures predict that the most critical glutamates E311[RBC:A] and E435[RBC:B] required for nucleoside binding preference would clash with the cyclic phosphate of cAMP (Fig. 2d), kinase activation at high concentration suggests that this incompatibility is not absolute (mutant 4, Fig. 3d), as expected for a dynamic structure *in vivo*. Cyclic nucleotide binding domains (CNB) are present in most species and in a broad variety of proteins, reaching from protein kinases (Diller et al. 2001; Su et al. 1995) to ion channels (Zagotta et al. 2003) and transcription factors, such as the catabolite activator protein (CAP) in bacteria (Kannan et al. 2007; Passner and Steitz 1997). The ancient CNB fold has been described as core module for allosteric regulation by cyclic nucleotides (Berman et al. 2005;

Kannan et al. 2007). Here we propose that the CNB is a module for allosteric regulation by a

broader spectrum of small ligands. This is reminiscent of other families of ligand binding

proteins like the G-protein coupled receptors or steroid hormone binding domains in

transcription factors that have been initially characterized by a limited set of ligands. Diverse

ligands have later been identified for the "orphan" members of those families (Davenport et

al. 2013). Future investigations might identify additional CNB domain ligands also outside the

Trypanosomatida group.

In contrast to residues important for ligand specificity, the π-stacking interaction by the so-

called capping residues (Wu et al. 2004; Kim et al. 2007) with the purine ring of cAMP or

nucleosides are well conserved between mammalian and trypanosomatids. Single mutations

of the mammalian PKA capping residues W260 [RIα:CNB-A] or Y371[RIα:CNB-B] that interact with

cAMP reduced activation potency by 4.6 and 9-fold, respectively and influence the

cooperativity of the two binding sites (Kim et al. 2007). The importance of stacking

interactions with Y371 in site A and Y482 in site B of TbPKAR explains why allopurinol

riboside, a purine derivative very similar to inosine with reduced delocalized electron system,

is 120-fold less potent than inosine (Table 1).

The biochemical and structural evidence for a distinct ligand specificity of trypanosomatid

PKA that is provided here, will resolve a very controversial issue: whereas difficulties to

detect cAMP-dependent kinase activity in *T. brucei* and *Leishmania* were reported long ago

(Walter 1978; Banerjee and Sarkar 2001), cAMP regulation of *T. cruzi* and *Leishmania* PKA

activity and binding of cAMP to LdPKAR1 in the μM range have been proposed by others

(Huang et al. 2006; Bhattacharya, Biswas, and Das 2012). Our data contradict the latter

reports and possible technical reasons have been discussed previously (Bachmaier et al.

2019; Bachmaier and Boshart 2013).

The tail makes the affinity difference

The very high affinity of nucleosides to the B-site was surprising as the ionic interaction of the

cyclic phosphate deep in the pocket is important for strong binding of cAMP in mammalian

PKA (Su et al. 1995; Herberg, Taylor, and Dostmann 1996). The C-terminal extension (αD

helix) is so far only found in trypanosomatid PKAR and contains the conserved sequence

(K/N)YxYY. Our crystal structures show that this helix covers the B binding site in the ligand

bound state like a lid and shields the ligand from solvent (Supplementary Movie 4). Inside the

binding pocket, the capping residue (Y482) π-stacks with the purine ring that may

additionally engage in T-shape π-stacking interaction with Y484 and Y485 from the αD helix

(Fig. 4b). C-terminal extensions are found in some PKAR subunits e.g. RIIß (PDB: 1CX4)

(Diller et al. 2001) or *Plasmodium falciparum* PKAR (PDB: 5K8S) (Littler et al. 2016). These

differ from the αD helix in that they just seem to prolong the αC helix. In contrast, the αD helix

is separated from αC by a proline resulting in a kink that positions the αD helix on top of the

binding pocket. The helix is attached to the binding pocket by several hydrogen bonds

donated by the conserved tyrosines Y484 and Y485 (Fig. 4b). Consequently, the

Y484A/Y485A mutation reduced the binding affinity for inosine to site B drastically. The αD

helix lid mechanism may compensate for the weaker bonds of the ribose moiety of

nucleosides deep in the pocket, whereas cAMP binding is stabilized by a strong ionic

interaction with the phosphate in cAMP dependent PKARs.

Site selectivity and synergism of nucleosides

The binding data for trypanosomatid PKARs show a clear affinity ranking of the three purine

nucleosides: K_D(ino)<K_D(guano)<K_D(adeno). The preference of inosine and guanosine over

adenosine is most striking for the B-site. The structures and molecular docking show that the

αD-helix contributes to the high affinity of guanosine and inosine, as in contrast to the

adenine base, the 6-oxopurines can form several hydrogen bonds to the αD-helix. This

preference is also seen as a 97-fold difference in kinase activation between cIMP and cAMP

in mutant 1 (Tab. 2). Initially, binding of nucleosides to TbPKAR and LdPKAR1 was

measured with recombinant proteins containing the complete C-terminus with both CNB

domains. This averages over the affinity of two binding sites. To determine single binding site

affinities, previous work on mammalian PKA used individual expressed CNB domains (Moll

et al. 2007; Lorenz et al. 2017). Here we blocked either the A (mutant 6) or B site (mutant 7)

of TbPKAR by conversion to cyclic nucleotide specificity to measure binding to the other site

in the context of the intact protein. This strategy reduces the risk of protein truncation

artefacts, but we are aware of the limitation of measuring binding to a "primed" protein (the

other binding site is occupied). The true affinity to the B-site in the holoenzyme (APO) state

available *in vivo* cannot be easily determined. This in mind, a strong site preference of

adenosine for site A and guanosine for site B was observed, whereas inosine bound equally

to both sites (Tab. S2). The αD-helix seems important for guanosine preference in site B

(see above), but single amino acids contributing to adenosine preference in site A could not

be identified by *in-silico* docking experiments (Supplementary Fig. 6). A co-crystal structure

of TbPKAR with adenosine is not yet available. The site selective binding of adenosine and

guanosine to sites A and B, respectively, of *T. brucei* PKAR is reminiscent of site selectivity

of cyclic nucleotide analogues for mammalian PKAR. A synergistic effect on kinase activation

of these synthetic compounds was exploited for development of potential anti-proliferative

drugs (Schwede et al. 2000; Cho-Chung et al. 1989; Huseby et al. 2011; Gausdal et al.

2013). Priming mammalian RI and RII isoenzymes with B-site selective cyclic nucleotides led

to an increase of activation potency of A-site selective compounds (Ogreid et al. 1989;

Dostmann et al. 1990; Corbin et al. 1986). We observed a comparable synergistic effect

upon priming of TbPKA with 10-20 nM of the B-site selective guanosine, resulting in a 6-20-

fold shift in activation potency by the A-site selective adenosine. In contrast to

pharmacological synergism of drugs acting on mammalian PKA, the synergism of two

endogenous ligands present in trypanosomes may have *in vivo* relevance by providing a

logical AND switch to respond to and integrate over the two most important purine

nucleosides in the cell.

Allosteric regulation of PKA

A detailed model for the allosteric activation mechanism of mammalian PKA has been

elaborated over many years (Su et al. 1995; Kim et al. 2007; Taylor et al. 2021; Rehmann,

Wittinghofer, and Bos 2007). Is this activation mode also applicable to the nucleoside

dependent PKA of trypanosomatids? The crystal structures of inosine bound TbPKAR and

mammalian PKARIα (PDB: 1RGS) are highly homologous (rmds = 3.2 Å). Key sequence

features implicated in the allosteric regulation, such as the salt bridge (E371, R475^{TbPKAR})

keeping the B/C helix extended in the apo conformation, the capping residues (Y370 and

Y482) that participate in ligand binding by π stacking (Kim et al. 2007; Wu et al. 2004) as

well as residues involved in R-C interaction e. g. the inhibitory sequence (RRTTV, res. 201-

206 in TbPKAR) (Kim, Xuong, and Taylor 2005; Kim et al. 2007) are conserved. The binding

and kinase activation data for single site mutants of TbPKA (Fig. 5b) clearly show that

nucleoside binding to both sites is required for efficient kinase activation and suggests that

the B-site has a "gatekeeper" function that initiates the conformational change and leads to

accessibility of the A-site, like in mammalian PKA. Subsequent ligand binding to the A-site

then triggers dissociation of the catalytic subunit (Herberg, Taylor, and Dostmann 1996; Kim,

Xuong, and Taylor 2005; Kim et al. 2007) and thereby releases the kinase from

(auto)inhibition. The results of our single site mutant analysis of TbPKA (Fig. 5b) are perfectly

compatible with this model. Therefore, the basic allosteric mechanism seems to be

conserved and may predate in evolution the separation of PKA into different CNB ligand

specificities. More insight into the conformational detail of kinase activation by nucleosides

will require the structure of a trypanosomatid PKA holoenzyme complex.

Nucleosides as second messengers?

Inosine, guanosine and adenosine were shown to bind to TbPKAR *in vivo* (Fig. 6), whereas

no compounds from an exhaustive list of nucleoside analogues previously identified in living

organisms (MODOMICS database) (Dunin-Horkawicz et al. 2006; Boccaletto et al. 2022;

Boccaletto et al. 2018) did match MS spectra from pulled down bound material

(Supplementary data 1). We may have missed a low affinity or labile ligand not captured by

the pulldown procedure. However, in favour of a second messenger role of nucleosides, their

binding affinities and activation potencies match well the affinity and potency of cAMP for

mammalian PKA (Fig. 1, Table 1, Supplementary Table 2). Furthermore, an unidentified second messenger would compete with the nucleosides in the cell and therefore require even higher affinity. Obviously, PKA might also serve as an intracellular nucleoside gauge. Trypanosomes are purine auxotroph and require sensitive regulation of purine uptake (Rico-Jimenez et al. 2021). Reconciling a classical second messenger function of nucleosides with their role of metabolites in the cell seems less of a conceptual problem. First, an increasing number of metabolites with signalling function emerge (Baker and Rutter 2023) and second, subcellular compartmentalization of signal generation/degradation is well established for cAMP signalling and PKA that act mostly in microdomains (Buxton and Brunton 1983; Zaccolo, Zerio, and Lobo 2021; Paolocci and Zaccolo 2023; Musheshe, Schmidt, and Zaccolo 2018). Compartmentalization of cAMP signalling also characterizes the trypanosome flagellum that serves as sensory organelle (Bachmaier et al. 2022; Ooi and Bastin 2013; Oberholzer et al. 2010; Shaw et al. 2022). Microdomains enriched in nucleoside salvage pathway enzymes may facilitate kinase activation or signal termination by rapidly generating or degrading specific nucleosides. The compartmentalized nucleotide metabolism (Ginger, Portman, and McKean 2008) in the flagellum exemplifies such a scenario. In fact, PKA of the trypanosomatid species analysed here is predominantly localized in the flagellum (Bachmaier et al. 2016; Oberholzer et al. 2011; Billington et al. 2023), The subcellular distribution of nucleosides in trypanosomes is not known and even reliable values for total guanosine and inosine per cell are unavailable in the literature. Only adenosine was measured in the bloodstream stage of *T. brucei* in the range of 12-28 µM depending on the growth medium (Kim et al. 2015). Taking the binding affinities into account, we can use the relative amounts of bound nucleosides pulled down with TbPKAR from cell lysates (Fig. 6) as proxy for relative nucleoside concentrations in PKA containing compartments. Only a small fraction of PKAR bound inosine *in vivo* in both life cycle stages, indicating that this high affinity binder is not readily available for PKAR in the parasites. This is in contrast to *E. coli* where inosine seems more abundant (extrapolated from Bennett et al. (2009)) resulting in inosine bound to trypanosomatid PKAR when expressed in this heterologous system. In *T. brucei* bloodstream

forms guanosine was captured but adenosine remained below the level of detection (Fig. 6).

In procyclic forms guanosine and adenosine were both captured by PKAR, indicating a much

higher adenosine/guanosine ratio in this life cycle stage. Based on these estimations,

guanosine seems to be the primary physiological ligand in BSF, whereas adenosine and

guanosine likely synergize to activate PKA in PCF (Fig. 5a). Most PKAR pulled down from

PCF would then have adenosine in the A-site and guanosine in the B-site. Termination of

signalling may be supported by PKA subunit turnover since rapid degradation of the

TbPKAC1 was seen upon depletion of TbPKAR by RNAi (Bachmaier et al. 2019).

As alternatives to a classical second messenger role of nucleosides, physiological activation

of the PKA holoenzyme *in vivo* may be co-activated by nucleoside binding together with a

second trigger like a posttranslational modification, redox state, specific protein-protein

interaction or kinase regulation by liquid–liquid phase separation (Lopez-Palacios and

Andersen 2023; Hardy et al. 2023). These triggers may allosterically shift the affinity or may

be required for the final activating conformational change upon binding (Khamina et al.

2022), giving nucleosides a more auxiliary role in the allosteric kinase regulation. In

trypanosomatids abundant and regulated phosphorylation of PKA subunits has been

reported (Tsigankov et al. 2014; Urbaniak, Martin, and Ferguson 2013). Kinase regulation

may then even not require a signal-related change of the intracellular nucleoside

concentration. The reasoning is inspired by the role of specific phosphorylations of

mammalian PKA (Haushalter et al. 2018) and the phenomenon of allosteric pluripotency

described for the analogue Rp-cAMP, that acts as an inhibitor or activator depending on

MgATP concentrations (Dostmann and Taylor 1991; Byun, Akimoto, et al. 2020).

Mechanistically, the opposite effects of Rp-cAMP are explained by formation of energetically

stabilized mixed intermediate states of the kinase, in which CNB-A and CNB-B adopt

different conformational states (Byun, VanSchouwen, et al. 2020; Akimoto et al. 2015).

Similar intermediate states were also reported for *Plasmodium falciparum* PKG (Byun, Van,

et al. 2020). Models for activation of the kinetoplastid PKA remain speculative as long as the

upstream signalling that leads to PKA activation *in vivo* has not been elucidated in these

organism. Future research will use kinase activation as readout for genome-wide screening

to detect upstream pathway components regulating the nucleoside-dependent PKA.

Acknowledgements

We thank Thomas Carell, LMU Chemistry, for generous support of S.B., MS instrument time and discussions, Ralph Heermann, LMU Microbiology, and Michaela Smolle, LMU BMC, for advice, and Andreas Anger, LMU Gene Center for structural homology modelling in the early phase of the project. Eleni Polatoglou launched Y.V.S. in the laboratory. Access to nanoDSF instruments and advice was generously provided by NanoTemper Technologies GmbH (Munich, Germany). We are grateful to Oliver Plettenburg for discussions and Ricardo Biondi, IBIOBA, Buenos Aires for critical reading of the manuscript. The work was supported by the Bundesministerium für Bildung und Forschung (BMBF) grant 16GW0281-3 to M.B. and F.S. Y.V.S. was supported by a fellowship from the Brazilian Science Without Borders/CNPq program and by the Life Sciences Munich (LSM) graduate school.

Author Contributions

Conceptualization [MB, GG, YVS]; Methodology [JB, SB]; Formal analysis [VO, GG, YVS, SB, MB]; Investigation [VO, GG, YVS, SB, GM]; Resources [FS]; Data curation [JB]; Writing – original draft preparation [VO; YVS; GG; MB]; Writing- review & editing [VO; YVS; MB]; Supervision [EL, MB]; Funding acquisition [MB]

Materials and Method

Preparation of PKA holoenzymes and kinase assay

T. brucei PKAR/PKAC1(TriTrypDB: Tb927.11.4610 and Tb927.9.11100); *T. cruzi* PKAR1/PKAC2 (TriTrypDB: TcCLB.506227.150 and TcCLB.508461.280), *L. donovani* PKAR1/PKAC1 (TritrypDB: LdBPK_130160.1 and LINF_350045600) isoform ORFs were amplified by PCR from their respective gDNA and fused to a 6xHis tag and a TEV cleavage site. PKACs were N-terminally fused to a strep tag. Mutations were introduced by PCR site directed mutagenesis via overlap extensions as described previously in Ho et al. (1989). Primer sequences are available in Supplementary file 2. The fusion ORFs were cloned into 3rd generation pLEXSY® vectors: pLEXSY_I-ble3 for PKARs and pLEXSY_I-neo3 for PKACs (Jena Bioscience). Holoenzymes were reconstituted *in vivo* by co-expression in *Leishmania tarentolae* T7-TR according to the Jena Bioscience manual. The holoenzyme complexes were isolated using a tandem affinity purification protocol: Ni-NTA followed by Streptactin matrix. *L. tarentolae* cells were detergent lysed by vortex homogenization in Ni-NTA binding buffer (50 mM NaH$_2$PO$_4$ pH 7.4, 150 mM NaCl, 20 mM imidazole, 0.5% Triton-X 100, Complete Mini EDTA-free protease inhibitor cocktail (Roche)). The soluble fraction was loaded onto a gravity flow Ni-NTA column. After washing with Ni-NTA wash buffer (50 mM NaH$_2$PO$_4$ pH 7.4, 150 mM NaCl, 40 mM imidazole) the protein complex was eluted in Ni-NTA elution buffer (50 mM NaH$_2$PO$_4$ pH 7.4, 150 mM NaCl, 250 mM imidazole). The eluate was immediately loaded onto the gravity flow Streptactin column, washed with Streptactin wash buffer (50 mM NaH$_2$PO$_4$ pH 7.4, 150 mM NaCl) and eluted with Streptactin elution buffer (50 mM NaH$_2$PO$_4$ pH 7.4, 150 mM NaCl, 2.5 mM desthiobiotin). All purification steps were carried out at 4°C. The mammalian PKA holoenzyme: human RIα/mouse Cα was co-expressed and co-purified from *E. coli* strain APE304 as previously described in (Bachmaier et al. 2019). The kinase assays with radiolabelled [γ32P] ATP were set up and performed as described in Hastie, McLauchlan, and Cohen (2006). Briefly, a 50µl kinase reaction mix was prepared at 4 °C by the addition of 5µl of 10x reaction buffer (500 mM MOPS pH 7; 1 M NaCl; 10 mM EGTA; 10 mM DTT; 1mg/ml BSA; 100 mM MgCl$_2$), 5 µl kemptide (1mM), 5 µl

purified PKA holoenzyme. A test run using known activating ligands (Bachmaier et al. 2019)

was carried out and the kinase would then be diluted in 1x reaction buffer, in order to work

within the linear range of the assay. The Ligands were diluted in 30µl H_2O prior to addition.

After temperature equilibration to 30°C, the kinase reaction was started by addition of 5 µl

1mM ATP spiked with [γ 32P] ATP to give 200-400 cpm/pmole. The reaction was stopped

after 10 min by pipetting 40 µl onto a 2x2cm p81-phosphocellulose paper and immediate

immersion into 75 mM phosphoric acid. Measurements were carried out in triplicates. Dose

response curve fitting was performed with Graphpad® prism's non-linear regression for

calculation of half activation constants EC_{50} and 95% confidence interval.

Protein expression and purification for ligand binding studies

N-terminally truncated *T. brucei* PKAR(199-499) and *T. cruzi* PKAR(200-503) were cloned

into pETDuet1 (Novagen) with a N-terminal 6xHis-tag. TbPKAR(199-499) mutants 6 and 7

(for sequences see Table 2) were generated by site-directed mutagenesis and cloned into

pETDuet1. N-terminally truncated *L. donovani* PKAR(200-502) was fused to a

Sumo_Ubiquitin Tag in a pET_Sumo vector (ThermoFisher). Refolded (RfAPO) and native

(N) protein samples were subjected to nano differential scanning fluorimetry (nanoDSF),

whereas for isothermal titration calorimetry (ITC), only refolded protein was used. Native and

refolded proteins were prepared as reported by Bachmaier et al. (2019) with the following

modifications: native protein (N) eluted from affinity chromatography (Ni-NTA column) was

dialyzed overnight and then directly probed for thermal stability using nanoDSF. Purification

of LdPKAR1(200-502) by Ni-NTA affinity chromatography was followed by SenP2 protease

mediated cleavage of the N-terminal Sumo tag during dialysis of the protein in 50mM Hepes

pH 7.5 and 50mM NaCl (buffer B). After denaturation of TbPKAR(199-499) mutants 6 and 7,

refolding occurred in a dialysis bag in presence of 1mM cAMP or 1mM cIMP, respectively. To

mutant 7, additionally 6.5 moles of cIMP per mole of refolded protein were added before ITC

measurements. Final elution of all proteins after Size Exclusion Chromatography (SEC) was

in either 50mM HEPES pH 7.5, 50 mM NaCl, and 1% DMSO (buffer A) or 50mM HEPES pH

7.5 and 50 mM NaCl (buffer B). Preparation of cAMP-free human PKARIα was performed

according to Buechler, Herberg, and Taylor (1993).

Isothermal titration calorimetry (ITC)

ITC measurements were carried out on a MicroCal PEAQ-ITC (Malvern) instrument. Refolded

proteins were diluted to 10-20 µM in buffer A or B. 100-200 µM ligand were diluted in the same

buffer as the protein and DMSO concentration of protein and ligand samples was adjusted as

accurately as possible. As we observed that the molar ratio N decreased with time after final

purification, we did all ITC experiments within a day to avoid precipitation of the protein. 2-4

µl of ligand were injected in a series of 13-19 injections into the protein sample at 298K. The

Differential Power (DP) between reference and sample cell was maintained at 8-10 µcal s^{-1}

in all experiments. Data analysis was performed with the MicroCal PEAQ-ITC software

applying a model with one binding site.

Thermal shift analysis using Nano Differentiation Scanning Fluorimetry (nanoDSF)

For nano differential scanning fluorimetry (nanoDSF), a Prometheus NT.48 (Nanotemper

Technologies, Munich) equipped with high sensitivity glass capillaries (PR-C006,

Nanotemper) was used. The technique allows label-free monitoring of protein melting

temperatures (T_m). Upon heating 10µl of protein sample per capillary from 15^o to 90^o, at a

rate of 1-2°C per minute, intrinsic fluorescence at 330 and 350nm (F_{330}/F_{350}) was recorded

and the ratio of both or only the fluorescence at 330nm was plotted as function of

temperature. The melting temperature (T_m) was calculated from the first derivative of the

curve, using the instrument's built-in software. Native (N) and refolded (RfAPO) protein

preparations of TbPKAR(199-499), TbPKAR(199-499) mutant 6 and 7, and LdPKAR1(200-

502) were subjected to nanoDSF before and after incubation in 1mM of ligand(s). Accurate

protein refolding was assumed when the melting temperature (T_m) of refolded and native

samples, both loaded with an excess of 1 mM ligand, presented close matching values.

Circular dichroism spectroscopy

For circular dichroism (CD) spectroscopy, native and refolded TbPKAR(199-499) samples from mutant 6 were prepared as described above with the following modifications: Refolding occurred in presence of 1mM inosine and 1mM cAMP, followed by a SEC on a Superdex 200 Increase 10/20 GL column (GE Healthcare) and elution in CD buffer (20mM NaPi), free of chlorine. To ensure complete buffer change, the refolded protein was subsequently passed over a PD10 column (Ge Healthcare) and again eluted in CD buffer. Similarly, native protein was passed twice over PD10 columns with elution in CD buffer. Proteins were diluted to 2µM (0.1mAU absorbance) and incubated with 10µM inosine and 10µM cAMP before measurement. The CD spectra were recorded using a Jasco J-815-150S spectropolarimeter (Jasco®, Tokyo, Japan) connected to a PTC 343 peltier set up to maintain the system at a constant temperature of 20 °C. The protein (sample volume=in 200µL) was inserted into a rectangular quartz cell of 0.1 cm path length and the UV spectra recorded by averaging 20 scans in the wavelength of 185-260 nm. The CD signal was recorded in a window of -7 to 10 mdeg. The identification of the peaks in the spectra related to α-helices (193 nm) and ß-sheet (208 and 222 nm) enrichment were performed according to Greenfield and Fasman (1969).

Crystallization, X-ray diffraction data collection and structure determination of ligand bound PKARs

Protein purification for crystallization of TbPKAR(199-499), TbPKAR(199-499) mutant 6 and TcPKAR(200-503) was performed as described in Bachmaier et al. (2019) with the following modifications: Native protein eluted from a Ni-NTA column was cleaved by TEV protease for removal of the N-terminal 6xHis tag and then subjected to SEC. Protein freshly eluted from the Superdex 200 10/300 GL column was concentrated to at least 10 mg ml^{-1} and, in order to ensure homogeneous ligand binding, incubated with either 1mM inosine (TbPKAR(199-499) and TcPKAR(200-503)) or 1mM cAMP (TbPKAR(199-499) mutant 6). Crystals grew within 7-

10 days using sitting drops (100-500 nL) crystalizing via the vapor diffusion method (Davies

and Segal 1971). Crystals of TbPKAR(199-499) were obtained in 50 mM Tris pH 8.0, 4%

MPD, 0.2 M ammonium sulfate, 32% PEG 3350 at 4°C. Crystals of TcPKAR(200-503) were

obtained in 20% PEG 3350, 0.2M Magnesium acetate at 4°C. Crystals of TbPKAR(199-499)

mutant 6 were obtained in 50 mM Tris pH 8.0, 0.2 mM Magnesium Chloride, 30% PEG 3350

at 4°C. Prior to flash cooling in liquid nitrogen, the crystals were briefly soaked in a mother

liquor solution made of the reservoir buffer and 40% (v/v) of ethylene glycol. The X-ray

diffraction data were collected at the Swiss Light Source beamline PXIII and on a Bruker D8

venture Metaljet system, at 100K. The collected data were processed using XDS and scaled

using XSCALE (Kabsch 2010; Kabsch 2012). The structure of TbPKAR(199-499) with

inosine was solved using the Sulphur SAD (Single-wavelength Anomalous Diffraction)

phasing method (Doutch et al. 2012). All other structures were solved by Molecular

Replacement (MR) (McCoy 2007) using the structure of TbPKAR + inosine as a search

model in the software Phaser as implemented in PHENIX (Liebschner et al. 2019; Adams et

al. 2010). All the MR solutions presented a TFZ score (Translation Function Z-score) > 8

indicating correct solutions. The molecular models of the proteins were built using the 2Fo-Fc

electron density map while the ligands were built using the difference map. The final

structure was achieved by iterative cycles of manual building in Coot (Emsley et al. 2010)

and refinement using PHENIX. Data collection and refinement statistics are summarized in

Supplementary Table 1.

In silico docking of nucleosides to TbPKAR

In silico docking was performed using the software Glide (Friesner et al. 2004) as

implemented in Maestro (Schrödinger®). Ligands were built manually and prepared using

LigPrep (Schrödinger®). Ionization states and tautomers were not considered during ligand

preparation. Stereoisomers had their chirality determined from the 3D structure (max. 32 per

ligand). The docking mode chosen was SP (Standard Precision). Chain B of TbPKAR (PDB:

6FLO) was chosen as a template for docking, since it presented a better overall electron

density. For docking of inosine and guanosine to the A-site of TbPKAR the grid constrains used were E311 and water 96 (Match at least = 2). For docking of adenosine, the K293 conformer was changed to reach a hydrogen bond with the N7 of the purine ring. The grid restrains used were E311, K293 (Match at least = 2). For docking to the B-site the chosen grid constraints were A444(N), G432(N), E435 and water 273 (Match at least = 2) for all three nucleosides. Poses were analysed by visual inspection and ranked according to Glide G-score (GG), a mathematical prediction of Gibbs Free Energy.

Mass spectrometry analysis of ligands bound to TbPKAR *in vivo*

The TbPKAR ORF was N-terminally fused to a 6xHis tag by PCR and cloned into the pLEW82 expression vector and transfected into MITat1.2SM (single marker) blood stream form (BSF) cells and EATRO11252T7 insect stage form (PCF) cells, both of which expressed T7 polymerase and tetracycline repressor. Cell culture was exactly as reported before for BSF (Bachmaier et al. 2019) and PCF (Schenk et al. 2021). Transfected MiTat1.2SM blood stream forms were kept under constant selection with 1 µg/ml G418 and 2.5 µg/ml Bleomycin. Transfected EATRO11252T7 PCF cells were cultured under constant selection with 10 µg/ml G418, 25 µg/ml hygromycin and 2.5 µg/ml bleomycin. Selected clones were induced with 1 µg/ml tetracycline for 24 hours. The cells were harvested by centrifugation (1400 g for 10 min), washed once with PBS and then detergent lysed in Ni-NTA binding buffer. The soluble fraction was incubated with magnetic Ni-NTA beads for 1 hour, followed by quick single washes in Ni-NTA binding buffer, Ni-NTA wash buffer, Streptactin wash buffer (50 mM NaH$_2$PO$_4$ pH 7.4, 150 mM NaCl) and finally MS-Grade H$_2$O. The beads were suspended in MS-grade water and boiled at 95°C for 5 minutes. Beads and denatured protein precipitate were removed by centrifugation at 10,000g for 10 min. The supernatant was then transferred to a fresh tube and stored at -20°C until analysis. For LC-ESI-MS, the samples were chromatographed by a Dionex Ultimate 3000 HPLC system with a flow of 0.15 ml/min over an Interchim Uptisphere 120Å 3HDO C18 column (150 x 2 mm), while maintaining the column temperature at 30 °C. Elution was performed with buffer A (2

mM HCOONH$_4$ in H$_2$O, pH 5.5) and buffer B (2 mM HCOONH$_4$ in H2O/MeCN 20/80, pH 5.5),

with a linear gradient from 0% to 15% buffer B in 45 min. The elution was monitored at 260

nm (Dionex Ultimate 3000 Diode Array Detector). The chromatographic eluent was directly

injected into the ion source of a Thermo Finnigan LTQ Orbitrap XL without prior splitting. Ions

were scanned by use of a positive polarity mode over a full-scan range of m/z 80-500 with a

resolution of 30000. Parameters of the mass spectrometer were tuned with a freshly mixed

aqueous solution of inosine (5 µM). The synthetic ^{13}C$_5$-labeled internal isotope standards

with an isotope enrichment of >99% were procured from Omicron Biochemicals Inc. The

quantification of nucleosides was carried out, as described in Traube et al. (2019), with the

following amounts of the corresponding isotope labelled internal standards: 256.8 fmol [^{13}C$_5$]-

inosine, 152.8 fmol [^{13}C$_5$]-guanosine, 662.8 fmol [^{13}C$_5$]-adenosine.

References

Adams, Paul D, Pavel V Afonine, Gábor Bunkóczi, Vincent B Chen, Ian W Davis, Nathaniel Echols, Jeffrey J Headd, L-W Hung, Gary J Kapral, and Ralf W Grosse-Kunstleve. 2010. "PHENIX: a comprehensive Python-based system for macromolecular structure solution." *Acta Crystallographica Section D: Biological Crystallography* 66 (2):213-221.

Akimoto, M., E. T. McNicholl, A. Ramkissoon, K. Moleschi, S. S. Taylor, and G. Melacini. 2015. "Mapping the Free Energy Landscape of PKA Inhibition and Activation: A Double-Conformational Selection Model for the Tandem cAMP-Binding Domains of PKA RIalpha." *PLoS Biol* 13 (11):e1002305. doi: 10.1371/journal.pbio.1002305.

Bachmaier, S., G. Giacomelli, E. Calvo-Alvarez, L. R. Vieira, J. Van Den Abbeele, A. Aristodemou, E. Lorentzen, M. K. Gould, A. Brennand, J. W. Dupuy, I. Forne, A. Imhof, M. Bramkamp, D. Salmon, B. Rotureau, and M. Boshart. 2022. "A multi-adenylate cyclase regulator at the flagellar tip controls African trypanosome transmission." *Nat Commun* 13 (1):5445. doi: 10.1038/s41467-022-33108-z.

Bachmaier, S., M. K. Gould, E. Polatoglou, R. Omelianczyk, A. E. Brennand, M. A. Aloraini, J. C. Munday, D. Horn, M. Boshart, and H. P. de Koning. 2023. "Novel kinetoplastid-specific cAMP binding proteins identified by RNAi screening for cAMP resistance in Trypanosoma brucei." *Front Cell Infect Microbiol* 13:1204707. doi: 10.3389/fcimb.2023.1204707.

Bachmaier, S., Y. Volpato Santos, S. Kramer, G. B. Githure, T. Klockner, J. Pepperl, C. Baums, R. Schenk, F. Schwede, H. G. Genieser, J. W. Dupuy, I. Forne, A. Imhof, J. Basquin, E. Lorentzen, and M. Boshart. 2019. "Nucleoside analogue activators of cyclic AMP-independent protein kinase A of Trypanosoma." *Nat Commun* 10 (1):1421. doi: 10.1038/s41467-019-09338-z.

Bachmaier, S., R. Witztum, P. Tsigankov, R. Koren, M. Boshart, and D. Zilberstein. 2016. "Protein kinase A signaling during bidirectional axenic differentiation in Leishmania." *Int J Parasitol* 46 (2):75-82. doi: 10.1016/j.ijpara.2015.09.003.

Bachmaier, Sabine, and Michael Boshart. 2013. "Kinetoplastid AGC Kinases." In *Protein Phosphorylation in Parasites*, 99-122.

Baker, N., C. M. C. Catta-Preta, R. Neish, J. Sadlova, B. Powell, E. V. C. Alves-Ferreira, V. Geoghegan, J. B. T. Carnielli, K. Newling, C. Hughes, B. Vojtkova, J. Anand, A. Mihut, P. B. Walrad, L. G. Wilson, J. W. Pitchford, P. Volf, and J. C. Mottram. 2021. "Systematic functional analysis of Leishmania protein kinases identifies regulators of differentiation or survival." *Nat Commun* 12 (1):1244. doi: 10.1038/s41467-021-21360-8.

Baker, S. A., and J. Rutter. 2023. "Metabolites as signalling molecules." *Nat Rev Mol Cell Biol* 24 (5):355-374. doi: 10.1038/s41580-022-00572-w.

Banerjee, C., and D. Sarkar. 2001. "The cAMP-binding proteins of Leishmania are not the regulatory subunits of cAMP-dependent protein kinase." *Comp Biochem Physiol B Biochem Mol Biol* 130 (2):217-26. doi: 10.1016/s1096-4959(01)00232-9.

Bennett, B. D., E. H. Kimball, M. Gao, R. Osterhout, S. J. Van Dien, and J. D. Rabinowitz. 2009. "Absolute metabolite concentrations and implied enzyme active site occupancy in Escherichia coli." *Nat Chem Biol* 5 (8):593-9. doi: 10.1038/nchembio.186.

Berman, H. M., L. F. Ten Eyck, D. S. Goodsell, N. M. Haste, A. Kornev, and S. S. Taylor. 2005. "The cAMP binding domain: an ancient signaling module." *Proc Natl Acad Sci U S A* 102 (1):45-50. doi: 10.1073/pnas.0408579102.

Bhattacharya, A., A. Biswas, and P. K. Das. 2012. "Identification of a protein kinase A regulatory subunit from Leishmania having importance in metacyclogenesis through

induction of autophagy." *Mol Microbiol* 83 (3):548-64. doi: 10.1111/j.1365-
2958.2011.07950.x.
Billington, K., C. Halliday, R. Madden, P. Dyer, A. R. Barker, F. F. Moreira-Leite, M.
Carrington, S. Vaughan, C. Hertz-Fowler, S. Dean, J. D. Sunter, R. J. Wheeler, and K.
Gull. 2023. "Genome-wide subcellular protein map for the flagellate parasite
Trypanosoma brucei." *Nat Microbiol* 8 (3):533-547. doi: 10.1038/s41564-022-01295-
6.
Boccaletto, P., M. A. Machnicka, E. Purta, P. Piatkowski, B. Baginski, T. K. Wirecki, V. de
Crecy-Lagard, R. Ross, P. A. Limbach, A. Kotter, M. Helm, and J. M. Bujnicki. 2018.
"MODOMICS: a database of RNA modification pathways. 2017 update." *Nucleic*
Acids Res 46 (D1):D303-D307. doi: 10.1093/nar/gkx1030.
Boccaletto, P., F. Stefaniak, A. Ray, A. Cappannini, S. Mukherjee, E. Purta, M. Kurkowska,
N. Shirvanizadeh, E. Destefanis, P. Groza, G. Avsar, A. Romitelli, P. Pir, E. Dassi, S.
G. Conticello, F. Aguilo, and J. M. Bujnicki. 2022. "MODOMICS: a database of RNA
modification pathways. 2021 update." *Nucleic Acids Res* 50 (D1):D231-D235. doi:
10.1093/nar/gkab1083.
Bubis, J., J. C. Martinez, M. Calabokis, J. Ferreira, C. E. Sanz-Rodriguez, V. Navas, J. L.
Escalona, Y. Guo, and S. S. Taylor. 2018. "The gene product of a Trypanosoma
equiperdum ortholog of the cAMP-dependent protein kinase regulatory subunit is a
monomeric protein that is not capable of binding cyclic nucleotides." *Biochimie*
146:166-180. doi: 10.1016/j.biochi.2017.12.010.
Buechler, Y. J., F. W. Herberg, and S. S. Taylor. 1993. "Regulation-defective mutants of type
I cAMP-dependent protein kinase. Consequences of replacing arginine 94 and
arginine 95." *J Biol Chem* 268 (22):16495-503.
Buxton, I. L., and L. L. Brunton. 1983. "Compartments of cyclic AMP and protein kinase in
mammalian cardiomyocytes." *J Biol Chem* 258 (17):10233-9.
Byun, J. A., M. Akimoto, B. VanSchouwen, T. S. Lazarou, S. S. Taylor, and G. Melacini.
2020. "Allosteric pluripotency as revealed by protein kinase A." *Sci Adv* 6
(25):eabb1250. doi: 10.1126/sciadv.abb1250.
Byun, J. A., K. Van, J. Huang, P. Henning, E. Franz, M. Akimoto, F. W. Herberg, C. Kim,
and G. Melacini. 2020. "Mechanism of allosteric inhibition in the Plasmodium
falciparum cGMP-dependent protein kinase." *J Biol Chem* 295 (25):8480-8491. doi:
10.1074/jbc.RA120.013070.
Byun, J. A., B. VanSchouwen, M. Akimoto, and G. Melacini. 2020. "Allosteric inhibition
explained through conformational ensembles sampling distinct "mixed" states."
Comput Struct Biotechnol J 18:3803-3818. doi: 10.1016/j.csbj.2020.10.026.
Canaves, J. M., and S. S. Taylor. 2002. "Classification and phylogenetic analysis of the
cAMP-dependent protein kinase regulatory subunit family." *J Mol Evol* 54 (1):17-29.
doi: 10.1007/s00239-001-0013-1.
Cayla, M., Y. R. Nievas, K. R. Matthews, and J. C. Mottram. 2022. "Distinguishing functions
of trypanosomatid protein kinases." *Trends Parasitol* 38 (11):950-961. doi:
10.1016/j.pt.2022.08.009.
Cho-Chung, Y. S., T. Clair, P. Tagliaferri, S. Ally, D. Katsaros, G. Tortora, L. Neckers, T. L.
Avery, G. W. Crabtree, and R. K. Robins. 1989. "Site-selective cyclic AMP analogs as
new biological tools in growth control, differentiation, and proto-oncogene
regulation." *Cancer Invest* 7 (2):161-77. doi: 10.3109/07357908909038282.
Choi, J., W. H. Jung, and J. W. Kronstad. 2015. "The cAMP/protein kinase A signaling
pathway in pathogenic basidiomycete fungi: Connections with iron homeostasis." *J*
Microbiol 53 (9):579-87. doi: 10.1007/s12275-015-5247-5.

Corbin, J. D., D. Ogreid, J. P. Miller, R. H. Suva, B. Jastorff, and S. O. Doskeland. 1986. "Studies of cGMP analog specificity and function of the two intrasubunit binding sites of cGMP-dependent protein kinase." *J Biol Chem* 261 (3):1208-14.

Davenport, A. P., S. P. Alexander, J. L. Sharman, A. J. Pawson, H. E. Benson, A. E. Monaghan, W. C. Liew, C. P. Mpamhanga, T. I. Bonner, R. R. Neubig, J. P. Pin, M. Spedding, and A. J. Harmar. 2013. "International Union of Basic and Clinical Pharmacology. LXXXVIII. G protein-coupled receptor list: recommendations for new pairings with cognate ligands." *Pharmacol Rev* 65 (3):967-86. doi: 10.1124/pr.112.007179.

Davies, David R, and DM Segal. 1971. "[25] Protein crystallization: Micro techniques involving vapor diffusion." In *Methods in enzymology*, 266-269. Elsevier.

Diller, T. C., Madhusudan, N. H. Xuong, and S. S. Taylor. 2001. "Molecular basis for regulatory subunit diversity in cAMP-dependent protein kinase: crystal structure of the type II beta regulatory subunit." *Structure* 9 (1):73-82. doi: 10.1016/s0969-2126(00)00556-6.

Dostmann, W. R., and S. S. Taylor. 1991. "Identifying the molecular switches that determine whether (Rp)-cAMPS functions as an antagonist or an agonist in the activation of cAMP-dependent protein kinase I." *Biochemistry* 30 (35):8710-6. doi: 10.1021/bi00099a032.

Dostmann, W. R., S. S. Taylor, H. G. Genieser, B. Jastorff, S. O. Doskeland, and D. Ogreid. 1990. "Probing the cyclic nucleotide binding sites of cAMP-dependent protein kinases I and II with analogs of adenosine 3',5'-cyclic phosphorothioates." *J Biol Chem* 265 (18):10484-91.

Doutch, J., M. A. Hough, S. S. Hasnain, and R. W. Strange. 2012. "Challenges of sulfur SAD phasing as a routine method in macromolecular crystallography." *J Synchrotron Radiat* 19 (Pt 1):19-29. doi: 10.1107/S0909049511049004.

Dunin-Horkawicz, S., A. Czerwoniec, M. J. Gajda, M. Feder, H. Grosjean, and J. M. Bujnicki. 2006. "MODOMICS: a database of RNA modification pathways." *Nucleic Acids Res* 34 (Database issue):D145-9. doi: 10.1093/nar/gkj084.

Emsley, P., B. Lohkamp, W. G. Scott, and K. Cowtan. 2010. "Features and development of Coot." *Acta Crystallogr D Biol Crystallogr* 66 (Pt 4):486-501. doi: 10.1107/S0907444910007493.

Friesner, R. A., J. L. Banks, R. B. Murphy, T. A. Halgren, J. J. Klicic, D. T. Mainz, M. P. Repasky, E. H. Knoll, M. Shelley, J. K. Perry, D. E. Shaw, P. Francis, and P. S. Shenkin. 2004. "Glide: a new approach for rapid, accurate docking and scoring. 1. Method and assessment of docking accuracy." *J Med Chem* 47 (7):1739-49. doi: 10.1021/jm0306430.

Gausdal, G., A. Wergeland, J. Skavland, E. Nguyen, F. Pendino, N. Rouhee, E. McCormack, L. Herfindal, R. Kleppe, U. Havemann, F. Schwede, O. Bruserud, B. T. Gjertsen, M. Lanotte, E. Segal-Bendirdjian, and S. O. Doskeland. 2013. "Cyclic AMP can promote APL progression and protect myeloid leukemia cells against anthracycline-induced apoptosis." *Cell Death Dis* 4 (2):e516. doi: 10.1038/cddis.2013.39.

Ginger, M. L., N. Portman, and P. G. McKean. 2008. "Swimming with protists: perception, motility and flagellum assembly." *Nat Rev Microbiol* 6 (11):838-50. doi: 10.1038/nrmicro2009.

Greenfield, N., and G. D. Fasman. 1969. "Computed circular dichroism spectra for the evaluation of protein conformation." *Biochemistry* 8 (10):4108-16. doi: 10.1021/bi00838a031.

Hardy, J. C., E. H. Pool, J. G. H. Bruystens, X. Zhou, Q. Li, D. R. Zhou, M. Palay, G. Tan, L. Chen, J. L. C. Choi, H. N. Lee, S. Strack, D. Wang, S. S. Taylor, S. Mehta, and J.

Zhang. 2023. "Molecular Determinants and Signaling Effects of PKA RIalpha Phase Separation." *bioRxiv*. doi: 10.1101/2023.12.10.570836.

Haste, N. M., H. Talabani, A. Doo, A. Merckx, G. Langsley, and S. S. Taylor. 2012. "Exploring the Plasmodium falciparum cyclic-adenosine monophosphate (cAMP)-dependent protein kinase (PfPKA) as a therapeutic target." *Microbes Infect* 14 (10):838-50. doi: 10.1016/j.micinf.2012.05.004.

Hastie, C. J., H. J. McLauchlan, and P. Cohen. 2006. "Assay of protein kinases using radiolabeled ATP: a protocol." *Nat Protoc* 1 (2):968-71. doi: 10.1038/nprot.2006.149.

Haushalter, K. J., D. E. Casteel, A. Raffeiner, E. Stefan, H. H. Patel, and S. S. Taylor. 2018. "Phosphorylation of protein kinase A (PKA) regulatory subunit RIalpha by protein kinase G (PKG) primes PKA for catalytic activity in cells." *J Biol Chem* 293 (12):4411-4421. doi: 10.1074/jbc.M117.809988.

Herberg, F. W., S. S. Taylor, and W. R. Dostmann. 1996. "Active site mutations define the pathway for the cooperative activation of cAMP-dependent protein kinase." *Biochemistry* 35 (9):2934-42. doi: 10.1021/bi951647c.

Hitz, E., N. Wiedemar, A. Passecker, B. A. S. Graca, C. Scheurer, S. Wittlin, N. M. B. Brancucci, I. Vakonakis, P. Maser, and T. S. Voss. 2021. "The 3-phosphoinositide-dependent protein kinase 1 is an essential upstream activator of protein kinase A in malaria parasites." *PLoS Biol* 19 (12):e3001483. doi: 10.1371/journal.pbio.3001483.

Ho, S. N., H. D. Hunt, R. M. Horton, J. K. Pullen, and L. R. Pease. 1989. "Site-directed mutagenesis by overlap extension using the polymerase chain reaction." *Gene* 77 (1):51-9. doi: 10.1016/0378-1119(89)90358-2.

Huang, G. Y., O. O. Gerlits, M. P. Blakeley, B. Sankaran, A. Y. Kovalevsky, and C. Kim. 2014. "Neutron diffraction reveals hydrogen bonds critical for cGMP-selective activation: insights for cGMP-dependent protein kinase agonist design." *Biochemistry* 53 (43):6725-7. doi: 10.1021/bi501012v.

Huang, G. Y., J. J. Kim, A. S. Reger, R. Lorenz, E. W. Moon, C. Zhao, D. E. Casteel, D. Bertinetti, B. Vanschouwen, R. Selvaratnam, J. W. Pflugrath, B. Sankaran, G. Melacini, F. W. Herberg, and C. Kim. 2014. "Structural basis for cyclic-nucleotide selectivity and cGMP-selective activation of PKG I." *Structure* 22 (1):116-24. doi: 10.1016/j.str.2013.09.021.

Huang, H., L. M. Weiss, F. Nagajyothi, H. B. Tanowitz, M. Wittner, G. A. Orr, and Y. Bao. 2006. "Molecular cloning and characterization of the protein kinase A regulatory subunit of Trypanosoma cruzi." *Mol Biochem Parasitol* 149 (2):242-5. doi: 10.1016/j.molbiopara.2006.05.008.

Huseby, S., G. Gausdal, T. J. Keen, E. Kjaerland, C. Krakstad, L. Myhren, K. Bronstad, C. Kunick, F. Schwede, H. G. Genieser, R. Kleppe, and S. O. Doskeland. 2011. "Cyclic AMP induces IPC leukemia cell apoptosis via CRE-and CDK-dependent Bim transcription." *Cell Death Dis* 2 (12):e237. doi: 10.1038/cddis.2011.124.

Jia, Y., J. B. Marq, H. Bisio, D. Jacot, C. Mueller, L. Yu, J. Choudhary, M. Brochet, and D. Soldati-Favre. 2017. "Crosstalk between PKA and PKG controls pH-dependent host cell egress of Toxoplasma gondii." *EMBO J* 36 (21):3250-3267. doi: 10.15252/embj.201796794.

Kabsch, W. 2012. "XSCALE."

Kabsch, Wolfgang. 2010. "XDS." *Acta Crystallographica Section D* 66 (2):125-132. doi: doi:10.1107/S0907444909047337.

Kannan, N., J. Wu, G. S. Anand, S. Yooseph, A. F. Neuwald, J. C. Venter, and S. S. Taylor. 2007. "Evolution of allostery in the cyclic nucleotide binding module." *Genome Biol* 8 (12):R264. doi: 10.1186/gb-2007-8-12-r264.

Khamina, M., K. Martinez Pomier, M. Akimoto, B. VanSchouwen, and G. Melacini. 2022. "Non-Canonical Allostery in Cyclic Nucleotide Dependent Kinases." *J Mol Biol* 434 (17):167584. doi: 10.1016/j.jmb.2022.167584.

Kim, C., C. Y. Cheng, S. A. Saldanha, and S. S. Taylor. 2007. "PKA-I holoenzyme structure reveals a mechanism for cAMP-dependent activation." *Cell* 130 (6):1032-43. doi: 10.1016/j.cell.2007.07.018.

Kim, C., and R. Sharma. 2021. "Cyclic nucleotide selectivity of protein kinase G isozymes." *Protein Sci* 30 (2):316-327. doi: 10.1002/pro.4008.

Kim, C., N. H. Xuong, and S. S. Taylor. 2005. "Crystal structure of a complex between the catalytic and regulatory (RIalpha) subunits of PKA." *Science* 307 (5710):690-6. doi: 10.1126/science.1104607.

Kim, D. H., F. Achcar, R. Breitling, K. E. Burgess, and M. P. Barrett. 2015. "LC-MS-based absolute metabolite quantification: application to metabolic flux measurement in trypanosomes." *Metabolomics* 11 (6):1721-1732. doi: 10.1007/s11306-015-0827-2.

Kim, J. S., K. T. Lee, M. H. Lee, E. Cheong, and Y. S. Bahn. 2021. "Adenylyl Cyclase and Protein Kinase A Play Redundant and Distinct Roles in Growth, Differentiation, Antifungal Drug Resistance, and Pathogenicity of Candida auris." *mBio* 12 (5):e0272921. doi: 10.1128/mBio.02729-21.

Kurokawa, H., K. Kato, T. Iwanaga, T. Sugi, A. Sudo, K. Kobayashi, H. Gong, H. Takemae, F. C. Recuenco, T. Horimoto, and H. Akashi. 2011. "Identification of Toxoplasma gondii cAMP dependent protein kinase and its role in the tachyzoite growth." *PLoS One* 6 (7):e22492. doi: 10.1371/journal.pone.0022492.

Lee, S. J., B. Lodder, Y. Chen, T. Patriarchi, L. Tian, and B. L. Sabatini. 2021. "Cell-type-specific asynchronous modulation of PKA by dopamine in learning." *Nature* 590 (7846):451-456. doi: 10.1038/s41586-020-03050-5.

Liebschner, D., P. V. Afonine, M. L. Baker, G. Bunkoczi, V. B. Chen, T. I. Croll, B. Hintze, L. W. Hung, S. Jain, A. J. McCoy, N. W. Moriarty, R. D. Oeffner, B. K. Poon, M. G. Prisant, R. J. Read, J. S. Richardson, D. C. Richardson, M. D. Sammito, O. V. Sobolev, D. H. Stockwell, T. C. Terwilliger, A. G. Urzhumtsev, L. L. Videau, C. J. Williams, and P. D. Adams. 2019. "Macromolecular structure determination using X-rays, neutrons and electrons: recent developments in Phenix." *Acta Crystallogr D Struct Biol* 75 (Pt 10):861-877. doi: 10.1107/S2059798319011471.

Littler, D. R., H. E. Bullen, K. L. Harvey, T. Beddoe, B. S. Crabb, J. Rossjohn, and P. R. Gilson. 2016. "Disrupting the Allosteric Interaction between the Plasmodium falciparum cAMP-dependent Kinase and Its Regulatory Subunit." *J Biol Chem* 291 (49):25375-25386. doi: 10.1074/jbc.M116.750174.

Lopez-Palacios, T. P., and J. L. Andersen. 2023. "Kinase regulation by liquid-liquid phase separation." *Trends Cell Biol* 33 (8):649-666. doi: 10.1016/j.tcb.2022.11.009.

Lorenz, R., D. Bertinetti, and F. W. Herberg. 2017. "cAMP-Dependent Protein Kinase and cGMP-Dependent Protein Kinase as Cyclic Nucleotide Effectors." *Handb Exp Pharmacol* 238:105-122. doi: 10.1007/164_2015_36.

Lorenz, R., E. W. Moon, J. J. Kim, S. H. Schmidt, B. Sankaran, I. V. Pavlidis, C. Kim, and F. W. Herberg. 2017. "Mutations of PKA cyclic nucleotide-binding domains reveal novel aspects of cyclic nucleotide selectivity." *Biochem J* 474 (14):2389-2403. doi: 10.1042/BCJ20160969.

Malmstrom, R. D., A. P. Kornev, S. S. Taylor, and R. E. Amaro. 2015. "Allostery through the computational microscope: cAMP activation of a canonical signalling domain." *Nat Commun* 6:7588. doi: 10.1038/ncomms8588.

Matthews, K. R. 2015. "25 years of African trypanosome research: From description to molecular dissection and new drug discovery." *Mol Biochem Parasitol* 200 (1-2):30-40. doi: 10.1016/j.molbiopara.2015.01.006.

Matthews, K. R. 2021. "Trypanosome Signaling-Quorum Sensing." *Annu Rev Microbiol*
75:495-514. doi: 10.1146/annurev-micro-020321-115246.
McCoy, Airlie J. 2007. "Solving structures of protein complexes by molecular replacement
with Phaser." *Acta Crystallographica Section D: Biological Crystallography* 63
(1):32-41.
Mohanty, S., E. J. Kennedy, F. W. Herberg, R. Hui, S. S. Taylor, G. Langsley, and N.
Kannan. 2015. "Structural and evolutionary divergence of cyclic nucleotide binding
domains in eukaryotic pathogens: Implications for drug design." *Biochim Biophys*
Acta 1854 (10 Pt B):1575-85. doi: 10.1016/j.bbapap.2015.03.012.
Moll, D., S. Schweinsberg, C. Hammann, and F. W. Herberg. 2007. "Comparative
thermodynamic analysis of cyclic nucleotide binding to protein kinase A." *Biol Chem*
388 (2):163-72. doi: 10.1515/BC.2007.018.
Mony, B. M., P. MacGregor, A. Ivens, F. Rojas, A. Cowton, J. Young, D. Horn, and K.
Matthews. 2014. "Genome-wide dissection of the quorum sensing signalling pathway
in Trypanosoma brucei." *Nature* 505 (7485):681-685. doi: 10.1038/nature12864.
Musheshe, N., M. Schmidt, and M. Zaccolo. 2018. "cAMP: From Long-Range Second
Messenger to Nanodomain Signalling." *Trends Pharmacol Sci* 39 (2):209-222. doi:
10.1016/j.tips.2017.11.006.
Niesen, Frank H., Helena Berglund, and Masoud Vedadi. 2007. "The use of differential
scanning fluorimetry to detect ligand interactions that promote protein stability."
Nature Protocols 2 (9):2212-2221. doi: 10.1038/nprot.2007.321.
Oberholzer, M., G. Langousis, H. T. Nguyen, E. A. Saada, M. M. Shimogawa, Z. O. Jonsson,
S. M. Nguyen, J. A. Wohlschlegel, and K. L. Hill. 2011. "Independent analysis of the
flagellum surface and matrix proteomes provides insight into flagellum signaling in
mammalian-infectious Trypanosoma brucei." *Mol Cell Proteomics* 10 (10):M111
010538. doi: 10.1074/mcp.M111.010538.
Oberholzer, M., M. A. Lopez, B. T. McLelland, and K. L. Hill. 2010. "Social motility in
african trypanosomes." *PLoS Pathog* 6 (1):e1000739. doi:
10.1371/journal.ppat.1000739.
Ogreid, D., R. Ekanger, R. H. Suva, J. P. Miller, and S. O. Doskeland. 1989. "Comparison of
the two classes of binding sites (A and B) of type I and type II cyclic-AMP-dependent
protein kinases by using cyclic nucleotide analogs." *Eur J Biochem* 181 (1):19-31.
doi: 10.1111/j.1432-1033.1989.tb14689.x.
Ooi, C. P., and P. Bastin. 2013. "More than meets the eye: understanding Trypanosoma brucei
morphology in the tsetse." *Front Cell Infect Microbiol* 3:71. doi:
10.3389/fcimb.2013.00071.
Paolocci, E., and M. Zaccolo. 2023. "Compartmentalised cAMP signalling in the primary
cilium." *Front Physiol* 14:1187134. doi: 10.3389/fphys.2023.1187134.
Passner, J. M., and T. A. Steitz. 1997. "The structure of a CAP-DNA complex having two
cAMP molecules bound to each monomer." *Proc Natl Acad Sci U S A* 94 (7):2843-7.
doi: 10.1073/pnas.94.7.2843.
Perrin, A. J., A. Patel, C. Flueck, M. J. Blackman, and D. A. Baker. 2020. "cAMP signalling
and its role in host cell invasion by malaria parasites." *Curr Opin Microbiol* 58:69-74.
doi: 10.1016/j.mib.2020.09.003.
Rehmann, H., A. Wittinghofer, and J. L. Bos. 2007. "Capturing cyclic nucleotides in action:
snapshots from crystallographic studies." *Nat Rev Mol Cell Biol* 8 (1):63-73. doi:
10.1038/nrm2082.
Rico-Jimenez, M., G. Ceballos-Perez, C. Gomez-Linan, and A. M. Estevez. 2021. "An RNA-
binding protein complex regulates the purine-dependent expression of a nucleobase
transporter in trypanosomes." *Nucleic Acids Res* 49 (7):3814-3825. doi:
10.1093/nar/gkab181.

Rinaldi, J., J. Wu, J. Yang, C. Y. Ralston, B. Sankaran, S. Moreno, and S. S. Taylor. 2010. "Structure of yeast regulatory subunit: a glimpse into the evolution of PKA signaling." *Structure* 18 (11):1471-82. doi: 10.1016/j.str.2010.08.013.

Salmon, D., S. Bachmaier, C. Krumbholz, M. Kador, J. A. Gossmann, P. Uzureau, E. Pays, and M. Boshart. 2012. "Cytokinesis of Trypanosoma brucei bloodstream forms depends on expression of adenylyl cyclases of the ESAG4 or ESAG4-like subfamily." *Mol Microbiol* 84 (2):225-42. doi: 10.1111/j.1365-2958.2012.08013.x.

Salmon, D., G. Vanwalleghem, Y. Morias, J. Denoeud, C. Krumbholz, F. Lhomme, S. Bachmaier, M. Kador, J. Gossmann, F. B. Dias, G. De Muylder, P. Uzureau, S. Magez, M. Moser, P. De Baetselier, J. Van Den Abbeele, A. Beschin, M. Boshart, and E. Pays. 2012. "Adenylate cyclases of Trypanosoma brucei inhibit the innate immune response of the host." *Science* 337 (6093):463-6. doi: 10.1126/science.1222753.

Schenk, R., S. Bachmaier, F. Bringaud, and M. Boshart. 2021. "Efficient flavinylation of glycosomal fumarate reductase by its own ApbE domain in Trypanosoma brucei." *FEBS J*. doi: 10.1111/febs.15812.

Schwede, F., E. Maronde, H. Genieser, and B. Jastorff. 2000. "Cyclic nucleotide analogs as biochemical tools and prospective drugs." *Pharmacol Ther* 87 (2-3):199-226. doi: 10.1016/s0163-7258(00)00051-6.

Shabb, J. B., B. D. Buzzeo, L. Ng, and J. D. Corbin. 1991. "Mutating protein kinase cAMP-binding sites into cGMP-binding sites. Mechanism of cGMP selectivity." *J Biol Chem* 266 (36):24320-6.

Shabb, J. B., L. Ng, and J. D. Corbin. 1990. "One amino acid change produces a high affinity cGMP-binding site in cAMP-dependent protein kinase." *J Biol Chem* 265 (27):16031-4.

Shaw, S., S. Knusel, D. Abbuhl, A. Naguleswaran, R. Etzensperger, M. Benninger, and I. Roditi. 2022. "Cyclic AMP signalling and glucose metabolism mediate pH taxis by African trypanosomes." *Nat Commun* 13 (1):603. doi: 10.1038/s41467-022-28293-w.

Su, Y., W. R. Dostmann, F. W. Herberg, K. Durick, N. H. Xuong, L. Ten Eyck, S. S. Taylor, and K. I. Varughese. 1995. "Regulatory subunit of protein kinase A: structure of deletion mutant with cAMP binding domains." *Science* 269 (5225):807-13. doi: 10.1126/science.7638597.

Taylor, S. S., R. Ilouz, P. Zhang, and A. P. Kornev. 2012. "Assembly of allosteric macromolecular switches: lessons from PKA." *Nat Rev Mol Cell Biol* 13 (10):646-58. doi: 10.1038/nrm3432.

Taylor, S. S., J. Wu, J. G. H. Bruystens, J. C. Del Rio, T. W. Lu, A. P. Kornev, and L. F. Ten Eyck. 2021. "From structure to the dynamic regulation of a molecular switch: A journey over 3 decades." *J Biol Chem* 296:100746. doi: 10.1016/j.jbc.2021.100746.

Toh, J. Y., A. Nkouawa, S. R. Sanchez, H. Shi, N. G. Kolev, and C. Tschudi. 2021. "Identification of positive and negative regulators in the stepwise developmental progression towards infectivity in Trypanosoma brucei." *Sci Rep* 11 (1):5755. doi: 10.1038/s41598-021-85225-2.

Traube, F. R., S. Schiffers, K. Iwan, S. Kellner, F. Spada, M. Muller, and T. Carell. 2019. "Isotope-dilution mass spectrometry for exact quantification of noncanonical DNA nucleosides." *Nat Protoc* 14 (1):283-312. doi: 10.1038/s41596-018-0094-6.

Tsigankov, P., P. F. Gherardini, M. Helmer-Citterich, G. F. Spath, P. J. Myler, and D. Zilberstein. 2014. "Regulation dynamics of Leishmania differentiation: deconvoluting signals and identifying phosphorylation trends." *Mol Cell Proteomics* 13 (7):1787-99. doi: 10.1074/mcp.M114.037705.

Uboldi, A. D., M. L. Wilde, E. A. McRae, R. J. Stewart, L. F. Dagley, L. Yang, N. J. Katris, S. V. Hapuarachchi, M. J. Coffey, A. M. Lehane, C. Y. Botte, R. F. Waller, A. I. Webb, M. J. McConville, and C. J. Tonkin. 2018. "Protein kinase A negatively

regulates Ca2+ signalling in Toxoplasma gondii." *PLoS Biol* 16 (9):e2005642. doi: 10.1371/journal.pbio.2005642.

Urbaniak, M. D., D. M. Martin, and M. A. Ferguson. 2013. "Global quantitative SILAC phosphoproteomics reveals differential phosphorylation is widespread between the procyclic and bloodstream form lifecycle stages of Trypanosoma brucei." *J Proteome Res* 12 (5):2233-44. doi: 10.1021/pr400086y.

Vaidyanathan, P. P., B. Zinshteyn, M. K. Thompson, and W. V. Gilbert. 2014. "Protein kinase A regulates gene-specific translational adaptation in differentiating yeast." *RNA* 20 (6):912-22. doi: 10.1261/rna.044552.114.

Walsh, D. A., J. P. Perkins, and E. G. Krebs. 1968. "An adenosine 3',5'-monophosphate-dependant protein kinase from rabbit skeletal muscle." *J Biol Chem* 243 (13):3763-5.

Walter, R. D. 1978. "Multiple protein kinases from Trypanosoma gambiense." *Hoppe Seylers Z Physiol Chem* 359 (5):601-6. doi: 10.1515/bchm.1978.359.1.601.

Weber, I. T., J. B. Shabb, and J. D. Corbin. 1989. "Predicted structures of the cGMP binding domains of the cGMP-dependent protein kinase: a key alanine/threonine difference in evolutionary divergence of cAMP and cGMP binding sites." *Biochemistry* 28 (14):6122-7. doi: 10.1021/bi00440a059.

Wu, J., S. Brown, N. H. Xuong, and S. S. Taylor. 2004. "RIalpha subunit of PKA: a cAMP-free structure reveals a hydrophobic capping mechanism for docking cAMP into site B." *Structure* 12 (6):1057-65. doi: 10.1016/j.str.2004.03.022.

Zaccolo, M., A. Zerio, and M. J. Lobo. 2021. "Subcellular Organization of the cAMP Signaling Pathway." *Pharmacol Rev* 73 (1):278-309. doi: 10.1124/pharmrev.120.000086.

Zagotta, W. N., N. B. Olivier, K. D. Black, E. C. Young, R. Olson, and E. Gouaux. 2003. "Structural basis for modulation and agonist specificity of HCN pacemaker channels." *Nature* 425 (6954):200-5. doi: 10.1038/nature01922.

 Figure Legends

 Figure 1. Trypanosomatid PKA binds to and is selectively activated by purine

 nucleosides.

a Structure-Activity Relationship (SAR) analysis for TbPKA kinase activation by nucleoside derivatives.

Chemical structures and the corresponding EC_{50} values are taken from Table 1. For representative

dose-response curves see Supplementary Fig. 1.

b Representative dose-response curves for activation of *T. brucei* PKAR-PKAC1 holoenzyme by

inosine, guanosine or adenosine (*in vitro* kinase assay, n ≥ 3 biological replicates). The calculated EC_{50}

values are displayed next to the graph and in Table 1, error bars indicate SD of technical triplicates.

Purity of PKA enzymes is shown in Supplementary Fig. 1a.

c Representative dose-response curves for activation of *L. donovani*, *T. cruzi* and mammalian (human

RIα/mouse Cα) holoenzymes by purine nucleosides and cAMP, as in A. The calculated EC_{50} values are

displayed next to the curve and in Table 1, error bars indicate SD of technical triplicates; Purity of PKA

holoenzyme is shown in Supplementary Fig. 1b, c.

d Binding isotherms (ITC) of nucleoside-depleted (APO) *T. brucei* PKAR(199-499) upon titration with

purine nucleosides. The graphs give the difference power (DP) between the reference and sample cells

upon ligand injection as a function of time (upper panel). In the lower panel, the total heat exchange per

mole of injectant (integrated peak areas from upper panel) is plotted against the molar ratio of ligand to

protein. A representative curve out of ≥ 3 independent replicates is shown. The final given K_D (as in

Table S2) was calculated as the mean (± SD) of at least 3 independent experiment (see Source Data

File). For purity of R subunit eluted from SEC see Supplementary Fig. S3a.

e Binding isotherms (ITC) of nucleoside-depleted (APO) *L. donovani* PKAR1(200-502) upon titration

with purine nucleosides, as in D. Purity, aggregation state and thermal stability of protein sample prior

to binding assays is shown in Supplementary Fig. 3c, d.

1221

Figure 2. Crystal structures of *T. cruzi* and *T. brucei* PKAR bound to inosine.

a Electron density (ED) maps of site A (left) and site B (right) of *T. cruzi* PKAR(200-503) and

corresponding ball and stick models of the hydrogen bond network around the bound inosine molecule.

The inosine molecule was modelled into the omit map (Fo-Fc, 3σ, green) in each binding site. The

surrounding protein atoms are shown together with a 2Fo-Fc map (1σ, dark blue). The black dashed

lines represent hydrogen bonds (≤ 3 Å cutoff). Residues G309; E310; E312 and A321 belong to Ribose

Binding Cassette A (RBC-A), while G433; E434 and E436 are part of Ribose Binding Cassette B (RBC-

B). Capping residues (Y371 and Y483) taking part in a π-stacking interaction with the hypoxanthine

ring of inosine are marked with an asterisk. Purine ring nomenclature is shown in the middle. PDB:

6HYI.

b *T. brucei* PKAR(199-499) displayed as in A, Residues G308, E309, E311 and A320 are part of RBC-

A while G432, E433, and E435 belong to RBC-B. Capping residues (Y370 and Y482) are marked with

an asterisk. PDB: 6FLO.

c Structural alignment of inosine-bound *T. cruzi* PKAR (PDB: 6HYI; protein grey, inosine yellow) and 7-

CN-7C-Ino-bound *T. cruzi* PKAR (PDB: 6FTF; protein and ligand in purple). The different ligand binding

to K294 (A-site, left) and a 1.5 Å displacement of Y485 due to the bulky cyano group of 7-CN-7-C-Ino

(B-site, right) are shown at two magnifications.

d Structural alignment of TbPKAR (PDB: 6FLO; protein grey, inosine yellow) and *B. taurus* PKARIα

(PDB: 1RGS; protein magenta, cAMP cyan) for binding sites A (left) and B (right). In the blow-up

panels, ligand-protein interactions are highlighted for the mammalian PKARIα (upper panel), TbPKAR

(middle panel), and TbPKAR overlayed with the cAMP ligand of the aligned PKARIα structure. A clash

between the exocyclic oxygens of cAMP and the side chain of glutamate residues (faded sphere-

representation) is seen in both binding sites.

1245

Figure 3. Conversion of TbPKAR to cyclic nucleotide specificity

a Structure of ligand binding sites of TbPKAR(199-499) mutant 6 crystallized in presence of 1mM

cAMP (Supplementary Fig. 3h, PDB: 6H4G). The scheme above the electron density map highlights

binding of cAMP to site A and inosine to site B. Below, the electron densities show the protein atoms

inside the 2Fo-Fc (1σ, dark blue) map and ligands inside the Fo-Fc omit map (3σ, green).

b Structural comparison between the A pocket of TbPKAR(199-499) mutant 6 (left) and BtPKARIα

(right, PDB: 1RGS). The point mutations in mutant 6 are coloured in purple (E311A), green (T318R)

and orange (V319A). The same colour code was used for the corresponding amino acids in BtPKARIα.

Hydrogen bonds (3Å) are indicated as dashed lines.

c Thermal denaturation profiles (nanoDSF) of refolded APO (upper panel) and native mutant 6

TbPKAR(199-499) (middle panel) in the absence and presence of 1 mM ligands as indicated. The

lower panel is a superposition of the thermal denaturation profiles of the two protein preparations

(native and refolded APO) both incubated with 1mM cAMP plus 1mM inosine.

d Mutational analysis of TbPKAR nucleoside binding sites. Relative kinase activation potency by

inosine (orange) and cIMP (blue) is displayed as $\log_2$ of the EC_{50}[Wildtype]/EC_{50} [$Mutant_n$] ratio on the

x-axis. Since up to 5mM cIMP did not activate the WT, this value was taken as minimal estimate of

EC_{50}[WT] for cIMP. This uncertainty propagating into the calculated ratio is indicated by a colour

gradient at the right end of the columns. All data are taken from Table 2. Missing columns are not

determined (n. d.). The sequences of RBC-A and RBC-B of mutants 1-5, with mutated amino acids

highlighted in red, are shown on the left to the respective columns.

1266

Figure 4. The αD helix of TbPKAR determines high binding affinity and ligand selectivity of the

B-site

a C-terminal helices αB (grey), αC (purple) and αD (green) in CNB-B of TbPKAR, illustrating the lid-like

position of the αD helix. The ribose binding cassette (RBC, residues 432 to 445) is shown in surface

representation (grey). Inosine (yellow) and one water molecule (blue) are sandwiched between RBC-B

and the αD helix (enlarged view in the blow-up panel). The 90° rotations of the structure show that the

αC and αD helices are orthogonally positioned to each other.

b The hydrogen-bond network formed between Y484 and Y485 in αD helix and amino acids in the beta

barrel of site B. Hydrogen bonds (3Å) are displayed as black dashed lines.

c Sequence alignment of C-terminal domains from *T. brucei*, *T. cruzi* and *L. donovani* PKARs. Only

TbPKAR numbering is shown. Red arrows mark residues involved in the hydrogen bond network with

Y484, black arrows mark residues involved in the hydrogen bond network with Y485.

d Extended network of hydrogen bonds and salt bridges (purple dashed lines) between two conserved

histidines (H430 and H440) and the ribose-binding glutamates E433 and E435.

e Beta factor representation of TbPKAR site B showing a higher overall atom displacement in the

crystal structure of the αD helix and in particular of P480. The Beta factor value increases from blue to

red and from thin to thick, indicating an increase of atom displacement in the crystal.

f Representative binding isotherm (ITC) for inosine binding to TbPKAR mutant 8 (mutant 6 with

additional substitutions Y484A, Y485A). Data representation as in Fig. 1d, the K_D value is taken from

Supplementary Table 2.

g, h Representative binding isotherms (ITC) for mutant 6 refolded in presence of 1mM cAMP (g) and

mutant 7 refolded in presence of 1mM cIMP (h). Data representation as in Fig. 1d, the K_D values are

taken from Supplementary Table 2. For sequences of mutants see Table 2, for purity and non-

aggregated state of R subunits see Supplementary Fig. 3j-l.

1291

Figure 5. Binding site selectivity and synergism of nucleosides

a Dose-response curves for kinase activation of TbPKA by adenosine in presence of 10nM or 20nM

guanosine. Error bars are m±SD of technical triplicates, the calculated EC_{50} values are given next to

the respective curve. Basal kinase activity in the absence of any ligand is indicated by a horizontal

dashed line. A green square (placed outside the log scale) represents the control with guanosine (10 or

20nM) only.

b Ratio of kinase activation over binding affinity (EC_{50}/K_D) for different purine nucleosides and

individual binding sites A and B. Unavailable binding sites in mutants 6 and 7 are indicated by red

crosses. Data are taken from Tables 2 and S2.

1301

Figure 6. Quantification of ligands bound to TbPKAR in lysed cells

HPLC-MS based quantification of nucleoside amounts released from boiled His-tagged TbPKAR pulled

down from lysed *T. brucei* (see Supplementary Fig. 7). Inosine (red), guanosine (green) and adenosine

(blue) were quantified using stable isotope-labelled internal standards. Error bars indicate SD from

three biological replicates. Note the different Y-axis scales. Pie charts on the right show the relative

amounts of nucleosides detected and quantified.

a Procyclic stage *T. brucei* strain EATRO1125 expressing His-TbPKAR and parental control cells.

Pulled down nucleosides from the control cell line were in the range of water blanks.

b Bloodstream stage *T. brucei* MITat 1.2 single marker line expressing His-TbPKAR, parental control

cells and isogenic *Δtbpkar/Δtbpkar* cells devoid of endogenous PKAR. The limit of quantification

(LOQ), defined by the linear part of the standard curves for stable isotope-labelled nucleoside

references, is given by a dashed line. Adenosine was below the LOQ.

Supplementary figures

1315

Supplementary Figure 1. Kinetoplastid PKA activation.

a Representative dose-response curves for activation of *T. brucei* PKAR-PKAC1 holoenzyme by

compounds (structural formula of some in Fig. 1a). The calculated EC_{50} values (taken from Table 1) are

displayed next to the curve, error bars indicate SD of technical triplicates. Purity of recombinant TbPKA

(6xHis-TbPKAR/strep-TbPKAC1) was confirmed by SDS-PAGE (left).

b, c as in A for LdPKA (6xHis-LdPKAR1/strep-LdPKAC1) and TcPKA (6xHis-TcPKAR/strep-TcPKAC2).

Purity of recombinant proteins was confirmed by SDS-PAGE (left).

d Binding isotherms of refolded APO *H. sapiens* PKARIα(1-381) in response to cAMP (left) and inosine

(centre). On the right, binding isotherm for *L. donovani* PKAR1(200-502) in response to cAMP. Data

representation as in Fig. 1d, the calculated K_D value for *H. sapiens* PKARIα(1-381) binding of cAMP

(mean± SD) is taken from Supplementary Table 2.

1327

Supplementary Figure 2. Mass spectrometry identification of ligands bound to recombinant PKAR.

a Coomassie staining and quantification relative to a BSA standard of purified 6xHis-TbPKAR (TbR, 800

μg) and *H. sapiens* 6xHis-PKARIα (hRIα, 2400 μg) expressed in *E. coli*.

b Proteins from A were boiled in MS grade H_2O to release bound ligands and supernatants were tested for

activation of mammalian PKAIα (left) and *T. brucei* PKA (right): (I) basal kinase activity (buffer control) (II)

supernatant from mock purification from *E. coli* harboring an empty vector (III) supernatant from boiled

hRIα (IV) supernatant from boiled TbPKAR (V) 10 µM of positive control ligand: inosine for TbPKA

holoenzyme and cAMP for human PKAIα holoenzyme.

c UV chromatogram (260 nm) from high-resolution HPLC-MS analysis of ligands from ≈ 3 nmoles of boiled

TbPKAR. The amounts of inosine (16.57) and guanosine (17.19) are indicated next to the respective peak

in the UV. Quantification has been performed by UV signal integration according to a standard curve.

Amounts of adenosine were below the range of reliable UV quantification.

d As in C for ligands released from human RIα; only cAMP was identified without quantification.

e MS chromatogram from the HPLC-MS analysis described in c and d. The selected mass range

([M + H$^+$] +/- 0.003 m/z) for inosine, guanosine, adenosine and cAMP is individually presented (from left to

right) as indicated. The retention time (Rt) is given for each identified peak.

1344

Supplementary Figure 3. Protein purification and quality controls for ITC and crystallization

experiments

a, c Size Exclusion Chromatogram (SEC) of refolded APO TbPKAR(199-499) and LdPKAR1(200-502),

respectively, used for ITC. Purity and expected molecular mass of protein samples are confirmed by SDS-

PAGE (inset).

b Thermal denaturation profiles (nanoDSF) of refolded APO (RfAPO) and native (N) *T. brucei* PKAR(199-

499) with and without ligands.

d Same as b for LdPKAR1(200-502) in absence or upon addition of 1mM inosine.

e Circular dichroism spectra (average of 20 scans) of TbPKAR mutant 6 native (dark blue) and refolded

APO (green) preparations.

f-h Size exclusion chromatograms of purified TcPKAR(200-503), TbPKAR(199- 499) and TbPKAR mutant

6, as indicated. Ligands added for co-crystallization are given next to the image of a representative crystal.

i-j Same as A for TbPKAR mutant 6 refolded without cAMP (i) and with 5 mM cAMP (j). Green arrows

indicate protein monomers, red arrows indicate aggregated or misfolded protein.

k Same as A for TbPKAR mutant 8.

l Same as A for TbPKAR mutant 7 refolded in presence of 1mM cIMP.

m Thermal denaturation profiles (nanoDSF) of native (N) TbPKAR mutant 7 in absence (light blue) and

upon incubation with 1mM cAMP (green) or 1mM cIMP (dark blue).

n Same as M for native TbPKAR mutant 7 (red) and refolded APO TbPKAR mutant 7(RfAPO, black) in

presence of 1mM cIMP and 1mM inosine.

o Sequence alignment of RBC-A/PBC-A(left) and RBC-B/PBC-B (right) motifs within CNB-A (left) and CNB-

B (right) of PKAR from *T. brucei (T.b.), T. cruzi (T.c.), L. donovani (L.d.) and B. taurus (B.t)*. Numbering

refers to sequence of *T. brucei* (top, green) and *B. taurus* (bottom, red). Degree of sequence conservation

is indicated in a colour code from red (high conservation) to blue (low conservation).

1369

1370 **Supplementary Figure 4. Activation of mutant TbPKA holoenzymes by different ligands.**

1371 Representative dose-response curves for activation of *T. brucei* PKA holoenzyme mutants 1-7 by

1372 purine nucleosides and cyclic nucleotides as in Fig. 1a. The sequences of RBC-A and RBC-B of

1373 mutants 1-7 are placed above the graphs with mutated amino acids highlighted in red. Calculated EC_{50}

1374 values are taken from Table 2.

1375

1376 **Supplementary Figure 5. Thermodynamic signatures of ligand binding from ITC experiments.**

1377 Direct comparison of the thermodynamic signatures from all ITC experiments summarized in

1378 Supplementary Table 2. Binding enthalpy is shown in green (ΔH), Gibbs free energy in blue (ΔG) and

1379 entropy in red (-TΔS).

1380

Supplementary Figure 6. Docking of nucleosides to A and B site of TbPKAR.

a Purine nucleosides (guanosine and adenosine) were docked into site A in the *T. brucei* PKAR crystal

structure (PDB: 6FLO, chain B,) using GLIDE (Friesner et al. 2004), as implemented in Maestro

(Schrödinger[TM]). Best poses were chosen according to the Glide G score (GG), given in the figure.

b Same as A for site B.

As a control, re-docking of inosine (magenta) was performed for comparison with the ligand in the

crystal structures (yellow). RMSD values are 0.035 Å for site A and 0.036 Å for site B.

1388

Supplementary Figure 7. HPLC-MS quantification of ligands bound to TbPKAR in parasite

lysates.

The left panels show western analysis of soluble fraction from lysed parasites expressing tagged

TbPKAR or of control parasites (input), pulled down beads and supernatants (all in triplicate). Anti-

PKAR, anti-PKAC1/2 and anti-His antibodies were used. Bound ligands were released by boiling of Ni-

NTA beads in water and the aqueous fraction subjected to HPLC-MS analysis. Graphs on the right

show chromatograms for the selected mass [M+H[+]] ± 0.003 m/z (given in parenthesis) for inosine,

guanosine and adenosine (all red coloured), and the matched stable isotope-labelled internal

standards (all blue coloured). Where peak quantification was in the linear range of the standard curves,

m ± SD pmole of nucleoside per biological replicate of 8 x 10[8] or 2 x 10[8] cells is given for PCF or BSF,

respectively (same values in Fig. 6). For graphical reasons the MS-signals were smoothened by

Boxcar algorithm.

a Procyclic stage (PCF) *T. brucei* strain EATRO1125 wild type cells and cells expressing His-TbPKAR,

as indicated. Nucleoside amounts pulled down from wild type PCF were in the range of blanks.

b Same as A for bloodstream stage *T. brucei* MITat 1.2 single marker *Δtbpkar/Δtbpkar* cells, wild type

cells and cells expressing His-TbPKAR, as indicated. Only trace amounts of adenosine were detected.

1405

1406 **Data availability**

1407

The coordinates of the crystal structures of *T. cruzi* PKAR bound to inosine, *T. brucei* PKAR

bound to inosine and *T. brucei* PKAR (mutant 6) bound to cAMP and inosine have been

deposited in the Protein Data Bank under the accession codes 6HYI, 6FLO, 6H4G,

respectively. Genome sequence and annotation information was obtained from TritrypDB

(http://www.tritrypdb.org). Results from a search of the MODOMICS database (Boccaletto et

al. 2022) to identify nucleoside analogues identified in living organisms is provided as

supplementary file 1. The source data underlying figures, tables, and Supplementary Figures

are provided as Source Data file.

1416

1417

Description of Additional Supplementary Files

File Name: Supplementary Data 1

Description: Results of searching the mass spectrometry data set (Fig. 6, S7) for nucleosides

matches in the MODOMICS (https://iimcb.genesilico.pl/modomics/) RNA modifications

database. Red cross: peak/mass not detected. Red: peak Rf/mass: detected but not

significant over background. Green mass: peak Rf/mass: detected but not confirmed.

1424

1425 File Name: Supplementary File 2

1426 Description: List of primers used in this study

1427

1428 File Name: Supplementary Movie 1

1429 Description: Alignment between TcPKAR (PDB: 6HYI, light blue) and TbPKAR (PDB:6FLO,

1430 chain B, light gray) displaying an RMSD of 0.909 Å calculated by PyMOL. Inosine is

1431 displayed in green and magenta in TbPKAR and TcPKAR, respectively.

1432

File Name: Supplementary Movie 2

Description: Alignment between A-site of PKARIα (PDB: 1RGS, gray, aa: 152-225) and A-

site from mutant 6 (PDB: 6H4G, light green, aa: 259-332). In mammalian PKAR, cAMP binds

in the *syn*-conformation, while in Mutant 6 it binds in the *anti*-conformation. Of particular note

is Cys278 in TbPKAR mutant 6, which is significantly displaced to the newly inserted R318.

Hydrogen bonds < 3Å are shown as black dashes.

1439

1440 File Name: Supplementary Movie 3

1441 Description: Sphere representation of the B-site from TbPKAR (PDB:6FLO, chain B, aa: 378

1442 to 490) showing residues Y484, Y485 and K488 in the αD helix in purple, V443 in green,

1443 inosine in blue and the rest of the protein in yellow. V443 is sandwiched between the alpha-D

1444 helix and the beta barrel, taking part in hydrophobic contacts to both sides.

1445

1446 File Name: Supplementary Movie 4

1447 Description: Surface representation of the B-site from TbPKAR (PDB:6FLO, chain B, aa: 378

1448 to 490) showing an inosine molecule (blue spheres) locked inside the protein with no access

1449 to solvent. The αD helix (N481 to end) is depicted in dark purple.

1450 **Table 1: Structure activity relation (SAR) analysis for PKA holoenzyme activation**
1451

PKA holoenzyme complex	Ligand	EC_{50} (95% CI)[1]
T. brucei PKAR/PKAC1	Inosine	14 (13 - 15) nM
	Guanosine	152 (132 - 172) nM
	Adenosine	7.0 (6.9 - 8.4) µM
	cAMP	-[2]
	cGMP	0.36 (0.33 - 0.41) mM
	cIMP	-[2]
	AMP	-[2]
	GMP	1.1 (0.9 - 1.3) mM
	IMP	108 (83 - 135) µM
	2'-deoxyadenosine	-[2]
	3'-deoxyadenosine	-[2]
	5'-deoxyadenosine	-[2]
	Nebularine	2.6 (2.2 - 3.2) µM
	Allopurinol riboside	1.7 (1.5 - 1.9) µM
	Xanthosine	62 (51 - 72) µM
	Uridine	40 (35 - 47) µM
	Cytidine	≥350 µM
Leishmania PKAR1/PKAC1	Inosine	47 (33 - 63) nM
	Guanosine	1.7 (1.4 - 2.1) µM
	Adenosine	6.5 (5.7 - 7.6) µM
	cAMP	-[2]
T. cruzi PKAR/PKAC2	Inosine	150 (110 - 200) nM
	Guanosine	3.5 (2.8 - 4.5) µM
	Adenosine	8.3 (5.2 - 12.4) µM
	cAMP	-[2]
human RIα/mouse Cα	Inosine	-[2]
	Guanosine	-[2]
	Adenosine	-[2]
	cAMP	75 (59 - 93) nM

1452

1453 [1] mean half activation constants (EC_{50}) and 95% confidence interval (95% CI) determined from
1454 Supplementary Fig. 1 using Graphpad prism 7.0 for technical triplicates.
1455 [2] no activation was detected up to a maximum concentration of 5mM

1456

1457 **Table 2: Activation of mutant TbPKA holoenzymes by different ligands**

1458

	TbPKA holoenzyme	EC_{50} (95% Cl)				
		Inosine	cIMP	Adenosine	Guanosine	cAMP
WT	RBC-A -GELELMYQTPTVA- RBC-B -GELEFLNNHANVA-	18 (13-22) nM[1]	-[3]	5 (3-7) µM[1]	0.14 (0.08-0.19) µM[1]	-[3]
Mut[5]						
1	RBC-A -GEL**A**LMYQTP**R**AA- RBC-B -GEL**A**FLNNHA**R**AA-	300 (160-600) µM[2]	0.34 (0.2-0.7) µM[2]	-[3]	370 (130-1000) µM[2]	33 (26-45) µM[2]
2	RBC-A -GEL**A**LMYQTPTVA- RBC-B -GEL**A**FLNNHANVA-	28 (23-35) µM[2]	nd[6]	nd	nd	-[4]
3	RBC-A -GELELMYQTP**V**VA- RBC-B -GELEFLNNHA**V**VA-	28 (15-37) nM[2]	nd	nd	nd	nd
4	RBC-A -GELELMYQTP**R**VA- RBC-B -GELEFLNNHA**R**VA-	36 (29-44) nM[2]	24 (22-26) µM[2]	nd	nd	-[4]
5	RBC-A -GEL**A**LMYQTP**R**VA- RBC-B -GEL**A**FLNNHA**R**VA-	18 (15-20) µM[2]	0.5 (0.3-0.6) µM[2]	nd	nd	90 (71-115) µM[2]
6	RBC-A -GEL**A**LMYQTP**R**AA- RBC-B -GELEFLNNHANVA-	0.2 (0.16-0.25) µM[2]	14 (10-25) µM[2]	-[3]	1.2 (0.6-2.6) µM[2]	25 (19-38) µM[2]
7	RBC-A -GELELMYQTPTVA- RBC-B -GEL**A**FLNNHA**R**AA-	1.1 (0.9-1.4) µM[2]	7 (5-11) µM[2]	21 (13-30) µM[2]	23 (17-32) µM[2]	-[4]

1459
[1] mean half activation constants (EC_{50}) and 95% confidence interval (95% Cl) for ≥ 3 biological replicates
[2] mean half activation constants (EC_{50}) and 95% confidence interval (95% Cl) for technical triplicate of a single biological
experiments
[3] no activation was detected up to a maximum concentration of 5mM
[4] no activation was detected up to a maximum concentration of 2mM
[5] number given to the respective mutant; site-directed mutations indicated in red
[6] not determined

1467

Supplementary Table 1: Data collection and refinement statistics for the crystal structures

	T.brucei PKAR (inosine)	*T.brucei* PKAR: E311A,T318R,V319A (cAMP – siteA inosine - siteB)	*T.cruzi* PKAR (inosine)
PDB	6FLO	6H4G	6HYI
Cell dimensions			
a, b, c (Å)	67.46, 71.55,122.82	64.67, 71.20, 121.40	40.74, 88.53,46.02
α, β, γ (°)	90, 90, 90	90, 90, 90	90, 98.28, 90
Data collection			
Resolution Range (Å)	49.08 - 2.14 (2.22 - 2.14)	30.21 - 2.14 (2.22 - 2.14)	44.26 - 1.40 (1.45 - 1.40)
Space group	P 21 21 21	P 21 21 21	P 1 21 1
X-ray wavelength (Å)	2.075	1.34	0.9792
Unique reflections	30009 (1997)	31336 (3027)	63354 (6238)
Reflections used for R-free	1998 (119)	2000 (194)	3536 (340)
Multiplicity	50	3.7	6.5
Completeness (%)	97.4	98.9	99.3
Mean I/σ (I)	25.7 (1.0)	10.4 (1.2)	11.7 (1.0)
Copies per ASU	2	2	1
R_{merge}	0.15 (2.03)	0.20 (1.78)	0.069 (1.70)
CC1/2	99.5 (49.1)	99.5 (47.2)	99.7 (48.2)
Refinement statistics			
R_{work} (%)	0.2354 (0.5243)	0.2304 (0.3318)	0.1803 (0.3538)
R_{free} (%)	0.2676 (0.4982)	0.2987 (0.3907)	0.2011 (0.3614)
Average B factor (A^2)			
Protein atoms	34.04	21.66	24.26
Solvent atoms	37.72	25.34	38.09
Ligand atoms	33.36	18.06	18.57
Ramachandran plot (%)			
Favored/Allowed	98/2	99/1	98/2
Outliers	0	0	0

1470

1471 *Values in parentheses are for highest-resolution shell.

1472

1473 **Supplementary Table 2: Binding parameters from ITC measurements**

1474

PKAR subunit	Ligand	K_D [nM][1]	Molar ratio
human PKARIα	cAMP	23 ± 15	1.6 ± 0.3
	Inosine[3]	>10[7]	
TbPKAR(199-499)	Inosine	18 ± 7	0.9 ± 0.2
	Guanosine	150 ± 51	1.0 ± 0.1
	Adenosine	825 ± 321	0.8 ± 0.1
	cAMP[2,3]	>10[7]	
LdPKAR1(200-502)	Inosine	59 ± 17	1.6 ± 0.4
	Guanosine	173 ± 58	1.5 ± 0.3
	Adenosine	1157 ± 226	2.0 ± 0.1
	cAMP[3]	>10[7]	
TbPKAR(199-499) E311A, T318R, V319A (mutant 6)	Inosine	18 ± 10	0.5 ± 0.1
	Guanosine	6 ± 3	0.7 ± 0.1
	Adenosine	1100 ± 200	0.6 ± 0.1
TbPKAR(199-499) E435A, N442R, V443A (mutant 7)	Inosine	9 ± 4	0.6 ± 0.1
	Guanosine	221 ± 27	0.7 ± 0.1
	Adenosine	82 ± 11	0.6 ± 0.1
TbPKAR(199-499) E311A, T318R, V319A **Y484A, Y485A** (mutant 8)	Inosine	1476 ± 125	0.6 ± 0.1

1475
1476 [1] mean ± SD of ≥ three independent measurements
1477 [2] taken from Bachmaier et al. 2019
1478 [3] technical limit of MicroCal PeaQ ITC - interpreted as no binding
1479

Figure 1. Trypanosomatid PKA binds to and is selectively activated by purine nucleosides.
a Structure-Activity Relationship (SAR) analysis for TbPKA kinase activation by nucleoside derivatives. Chemical structures and the corresponding EC_{50} values are taken from Table 1. For representative dose-response curves see Supplementary Fig. 1. **b** Representative dose response curves for activation of *T. brucei* PKAR-PKAC1 holoenzyme by inosine, guanosine or adenosine (*in vitro* kinase assay, n ≥ 3 biological replicates). The calculated EC_{50} values are displayed next to the graph and in Table 1, error bars indicate SD of technical triplicates. Purity of PKA enzymes is shown in Supplementary Fig. 1a. **c** Representative dose-response curves for activation of *L. donovani*, *T. cruzi* and mammalian (human RIα/mouse Cα) holoenzymes by purine nucleosides and cAMP, as in A. The calculated EC_{50} values are displayed next to the curve and in Table 1, error bars indicate SD of technical triplicates; Purity of PKA holoenzyme is shown in Supplementary Fig. 1b, c. **d** Binding isotherms (ITC) of nucleoside-depleted (APO) *T. brucei* PKAR(199-499) upon titration with purine nucleosides. The graphs give the difference power (DP) between the reference and sample cells upon ligand injection as a function of time (upper panel). In the lower panel, the total heat exchange per mole of injectant (integrated peak areas from upper panel) is plotted against the molar ratio of ligand to protein. A representative curve out of ≥ 3 independent replicates is shown. The final given K_D (as in Table S2) was calculated as the mean (± SD) of at least 3 independent experiment (see Source Data File). For purity of R subunit eluted from SEC see Supplementary Fig. S3a. **e** Binding isotherms (ITC) of nucleoside-depleted (APO) *L. donovani* PKAR1(200-502) upon titration with purine nucleosides, as in D. Purity, aggregation state and thermal stability of protein sample prior to binding assays is shown in Supplementary Fig. 3c, d.

Figure 2. Crystal structures of *T. cruzi* and *T. brucei* PKAR bound to inosine.
a Electron density (ED) maps of site A (left) and site B (right) of *T. cruzi* PKAR(200-503) and corresponding ball and stick models of the hydrogen bond network around the bound inosine molecule. The inosine molecule was modelled into the omit map (Fo-Fc, 3σ, green) in each binding site. The surrounding protein atoms are shown together with a 2Fo-Fc map (1σ, dark blue). The black dashed lines represent hydrogen bonds (≤ 3 Å cutoff). Residues G309; E310; E312 and A321 belong to Ribose Binding Cassette A (RBC-A), while G433; E434 and E436 are part of Ribose Binding Cassette B (RBC-B). Capping residues (Y371 and Y483) taking part in a π-stacking interaction with the hypoxanthine ring of inosine are marked with an asterisk. Purine ring nomenclature is shown in the middle. PDB: 6HYI. **b** *T. brucei* PKAR(199-499) displayed as in A, Residues G308, E309, E311 and A320 are part of RBC-A while G432, E433, and E435 belong to RBC-B. Capping residues (Y370 and Y482) are marked with an asterisk. PDB: 6FLO. **c** Structural alignment of inosine-bound *T. cruzi* PKAR (PDB: 6HYI; protein grey, inosine yellow) and 7-CN-7C-Ino-bound *T. cruzi* PKAR (PDB: 6FTF; protein and ligand in purple). The different ligand binding to K294 (A-site, left) and a 1.5 Å displacement of Y485 due to the bulky cyano group of 7-CN-7-C-Ino (B-site, right) are shown at two magnifications. **d** Structural alignment of TbPKAR (PDB: 6FLO; protein grey, inosine yellow) and *B. taurus* PKARIα (PDB: 1RGS; protein magenta, cAMP cyan) for sites A (left) and B (right). In the blow-up panels, ligand-protein interactions are highlighted for the mammalian PKARIα (upper panel), TbPKAR (middle panel), and TbPKAR overlayed with the cAMP ligand of the aligned PKARIα structure. A clash between the exocyclic oxygens of cAMP and the side chain of glutamate residues (faded sphere-representation) is seen in both binding sites.

Figure 3. Conversion of TbPKAR to cyclic nucleotide specificity.
a Structure of ligand binding sites of TbPKAR(199-499) mutant 6 crystallized in presence of 1mM cAMP (Supplementary Fig. 3h, PDB: 6H4G). The scheme above the electron density map highlights binding of cAMP to site A and inosine to site B. Below, the electron densities show the protein atoms inside the 2Fo-Fc (1σ, dark blue) map and ligands inside the Fo-Fc omit map (3σ, green). **b** Structural comparison between the A pocket of TbPKAR(199-499) mutant 6 (left) and BtPKARIα (right, PDB: 1RGS). The point mutations in mutant 6 are coloured in purple (E311A), green (T318R) and orange (V319A). The same colour code was used for the corresponding amino acids in BtPKARIα. Hydrogen bonds (3Å) are indicated as dashed lines. **c** Thermal denaturation profiles (nanoDSF) of refolded APO (upper panel) and native mutant 6 TbPKAR(199-499) (middle panel) in the absence and presence of 1 mM ligands as indicated. The lower panel is a superposition of the thermal denaturation profiles of the two protein preparations (native and refolded APO) both incubated with 1mM cAMP plus 1mM inosine.
d Mutational analysis of TbPKAR nucleoside binding sites. Relative kinase activation potency by inosine (orange) and cIMP (blue) is displayed as $\log_2$ of the EC50$_{50}$[Wildtype]/EC$_{50}$ [Mutant$_n$] ratio on the x-axis. Since up to 5mM cIMP did not activate the WT, this value was taken as minimal estimate of EC$_{50}$[WT] for cIMP. This uncertainty propagating into the calculated ratio is indicated by a colour gradient at the right end of the columns. All data are taken from Table 2. Missing columns are not determined (n. d.). The sequences of RBC-A and RBC-B of mutants 1-5, with mutated amino acids highlighted in red, are shown on the left to the respective columns.

Figure 4. The αD helix of TbPKAR determines high binding affinity and ligand selectivity of the B-site.
a C-terminal helices αB (grey), αC (purple) and αD (green) in CNB-B of TbPKAR, illustrating the lid-like position of the αD helix. The ribose binding cassette (RBC, residues 432 to 445) is shown in surface representation (grey). Inosine (yellow) and one water molecule (blue) are sandwiched between RBC-B and the αD helix (enlarged view in the blow-up panel). The 90° rotations of the structure show that the αC and αD helices are orthogonally positioned to each other. **b** The hydrogen-bond network formed between Y484 and Y485 in αD helix and amino acids in the beta barrel of site B. Hydrogen bonds (3Å) are displayed as black dashed lines. **c** Sequence alignment of C-terminal domains from *T. brucei*, *T. cruzi* and *L. donovani* PKARs. Only TbPKAR numbering is shown. Red arrows mark residues involved in the hydrogen bond network with Y484, black arrows mark residues involved in the hydrogen bond network with Y485. **d** Extended network of hydrogen bonds and salt bridges (purple dashed lines) between two conserved histidines (H430 and H440) and the ribose-binding glutamates E433 and E435. **e** Beta factor representation of TbPKAR site B showing a higher overall atom displacement in the crystal structure of the αD helix and in particular of P480. The Beta factor value increases from blue to red and from thin to thick, indicating an increase of atom displacement in the crystal.
f Representative binding isotherm (ITC) for inosine binding to TbPKAR mutant 8 (mutant 6 with additional substitutions Y484A, Y485A). Data representation as in Fig. 1d, the K_D value is taken from Supplementary Table 2. **g, h** Representative binding isotherms (ITC) for mutant 6 refolded in presence of 1mM cAMP (g) and mutant 7 refolded in presence of 1mM cIMP (h). Data representation as in Fig. 1d, the K_D values are taken from Supplementary Table 2. For sequences of mutants see Table 2, for purity and non-aggregated state of R subunits see Supplementary Fig. 3j-l.

Figure 5. Binding site selectivity and synergism of nucleosides.
a Dose-response curves for kinase activation of TbPKA by adenosine in presence of 10nM or 20nM guanosine. Error bars are m±SD of technical triplicates, the calculated EC_{50} values are given next to the respective curve. Basal kinase activity in the absence of any ligand is indicated by a horizontal dashed line. A green square (placed outside the log scale) represents the control with guanosine (10 or 20nM) only.
b Ratio of kinase activation over binding affinity (EC_{50}/K_D) for different purine nucleosides and individual binding sites A and B. Unavailable binding sites in mutants 6 and 7 are indicated by red crosses. Data are taken from Tables 2 and S2.

Figure 6. Quantification of ligands bound to TbPKAR in lysed cells
HPLC-MS based quantification of nucleoside amounts released from boiled His-tagged TbPKAR pulled down from lysed *T. brucei* (see Supplementary Fig. 7). Inosine (red), guanosine (green) and adenosine (blue) were quantified using stable isotope-labelled internal standards. Error bars indicate SD from three biological replicates. Note the different Y-axis scales. Pie charts on the right show the relative amounts of nucleosides detected and quantified. **a** Procyclic stage *T. brucei* strain EATRO1125 expressing His-TbPKAR and parental control cells. Pulled down nucleosides from the control cell line were in the range of water blanks. **b** Bloodstream stage *T. brucei* MITat 1.2 single marker line expressing His-TbPKAR, parental control cells and isogenic *Δtbpkar/Δtbpkar* cells devoid of endogenous PKAR. The limit of quantification (LOQ), defined by the linear part of the standard curves for stable isotope-labelled nucleoside references, is given by a dashed line. Adenosine was below the LOQ.

Supplementary Figure 1. Kinetoplastid PKA activation.
a Representative dose-response curves for activation of *T. brucei* PKAR-PKAC1 holoenzyme by compounds (structural formula of some in Fig. 1a). The calculated EC$_{50}$ values (taken from Table 1) are displayed next to the curve, error bars indicate SD of technical triplicates. Purity of recombinant TbPKA (6xHis-TbPKAR/strep-TbPKAC1) was confirmed by SDS-PAGE (left). **b, c** as in A for LdPKA (6xHis-LdPKAR1/strep LdPKAC1) and TcPKA (6xHis-TcPKAR/strep-TcPKAC2). Purity of recombinant proteins was confirmed by SDS-PAGE (left). **d** Binding isotherms of refolded APO *H. sapiens* PKARIα(1-381) in response to cAMP (left) and inosine (centre). On the right, binding isotherm for *L. donovani* PKAR1(200-502) in response to cAMP. Data representation as in Fig. 1d, the calculated K$_D$ value for *H. sapiens* PKARIα(1-381) binding of cAMP (mean± SD) is taken from Supplementary Table 2.

Supplementary Figure 2. Mass spectrometry identification of ligands bound to recombinant PKAR.
a Coomassie staining and quantification relative to a BSA standard of purified 6xHis-TbPKAR (TbR, 800 µg) and *H. sapiens* 6xHis-PKARIα (hRIα, 2400 µg) expressed in *E. coli*. **b** Proteins from A were boiled in MS grade H_2O to release bound ligands and supernatants were tested for activation of mammalian PKAIα (left) and *T. brucei* PKA (right): (I) basal kinase activity (buffer control) (II) supernatant from mock purification from *E. coli* harboring an empty vector (III) supernatant from boiled hRIα (IV) supernatant from boiled TbPKAR (V) 10 µM of positive control ligand: inosine for TbPKA holoenzyme and cAMP for human PKAIα holoenzyme. **c** UV chromatogram (260 nm) from high-resolution HPLC-MS analysis of ligands from ≈ 3 nmoles of boiled TbPKAR. The amounts of inosine (16.57) and guanosine (17.19) are indicated next to the respective peak in the UV. Quantification has been performed by UV signal integration according to a standard curve. Amounts of adenosine were below the range of reliable UV quantification. **d** As in C for ligands released from human RIα; only cAMP was identified without quantification. **e** MS chromatogram from the HPLC-MS analysis described in c and d. The selected mass range ([M + H$^+$] +/- 0.003 m/z) for inosine, guanosine, adenosine and cAMP is individually presented (from left to right) as indicated. The retention time (Rt) is given for each identified peak.

Supplementary Figure 3. Protein purification and quality controls for ITC and crystallization experiments
a, c Size Exclusion Chromatogram (SEC) of refolded APO TbPKAR(199-499) and LdPKAR1(200-502), respectively, used for ITC. Purity and expected molecular mass of protein samples are confirmed by SDS-PAGE (inset).**b** Thermal denaturation profiles (nanoDSF) of refolded APO (RfAPO) and native (N) *T. brucei* PKAR(199-499) with and without ligands. **d** Same as b for LdPKAR1(200-502) in absence or upon addition of 1mM inosine. **e** Circular dichroism spectra (average of 20 scans) of TbPKAR mutant 6 native (dark blue) and refolded APO (green) preparations. **f-h** Size exclusion chromatograms of purified TcPKAR(200-503), TbPKAR(199- 499) and TbPKAR mutant 6, as indicated. Ligands added for co-crystallization are given next to the image of a representative crystal. **i-j** Same as A for TbPKAR mutant 6 refolded without cAMP (i) and with 5 mM cAMP (j). Green arrows indicate protein monomers, red arrows indicate aggregated or misfolded protein. **k** Same as A for TbPKAR mutant 8. **l** Same as A for TbPKAR mutant 7 refolded in presence of 1mM cIMP. **m** Thermal denaturation profiles (nanoDSF) of native (N) TbPKAR mutant 7 in absence (light blue) and upon incubation with 1mM cAMP (green) or 1mM cIMP (dark blue). **n** Same as M for native TbPKAR mutant 7 (red) and refolded APO TbPKAR mutant 7(RfAPO, black) in presence of 1mM cIMP and 1mM inosine. **o** Sequence alignment of RBC-A/PBC-A(left) and RBC-B/PBC-B (right) motifs within CNB-A (left) and CNB-B (right) of PKAR from *T. brucei (T.b.), T. cruzi (T.c.), L. donovani (L.d.) and B. taurus (B.t)*. Numbering refers to sequence of *T. brucei* (top, green) and *B. taurus* (bottom, red). Degree of sequence conservation is indicated in a colour code from red (high conservation) to blue (low conservation).

Supplementary Figure 4. Activation of mutant TbPKA holoenzymes by different ligands.
Representative dose-response curves for activation of *T. brucei* PKA holoenzyme mutants 1-7 by purine nucleosides and cyclic nucleotides as in Fig. 1a. The sequences of RBC-A and RBC-B of mutants 1-7 are placed above the graphs with mutated amino acids highlighted in red. Calculated EC_{50} values are taken from Table 2.

	TbPKAR(199-499)			TbPKAR mutant 6				TbPKAR mutant 7			LdPKAR1			HsPKAR1α[2]
							Mutant 8 (Y484A, Y485A)							
	Ino	Guano	Adeno	Ino	Guano	Adeno	Ino	Ino	Guano	Adeno	Ino	Guano	Adeno	cAMP
ΔH[1]	-30±1	-25±1	-33±2	-30±1	-27±2	-26±3	-22±1	-24±2	-17±1	-27±0.3	-29±7	-22±4	-16±3	-19±1
ΔG[1]	-10.6±0.2	-9.3±0.2	-8.4±0.3	-10.6±0.4	-11.3±0.3	-8.1±0.1	-8.0±0.1	-11±0.3	-9.1±0.1	-9.7±0.1	-9.9±0.2	-9.3±0.2	-8.1±0.1	-9.9±0.4
-TΔS[1]	19±1	16±1	25±2	19±2	16±2	18±3	14±1	13±1	8±1	17±0.3	19±7	13±4	8±3	9±1

[1] mean ± SD of ≥ 3 independent measurements

[2] taken from Bachmaier et al. 2019

Supplementary Figure 5. Thermodynamic signatures of ligand binding from ITC experiments.
Direct comparison of the thermodynamic signatures from all ITC experiments summarized in Supplementary Table 2. Binding enthalpy is shown in green (ΔH), Gibbs free energy in blue (ΔG) and entropy in red (–TΔS).

a

Inosine
GG: -8.400

Guanosine
GG: -8.224

Adenosine
GG: -8.226

b

Inosine
GG: -9.961

Guanosine
GG: - 9.550

Adenosine
GG: - 7.814

Supplementary Figure 6. Docking of nucleosides to A and B site of TbPKAR.
a Purine nucleosides (guanosine and adenosine) were docked into site A in the *T. brucei* PKAR crystal structure (PDB: 6FLO, chain B,) using GLIDE (Friesner et al., 2004), as implemented in Maestro (Schrödinger™). Best poses were chosen according to the Glide G score (GG), given in the figure. **b** Same as A for site B.
As a control, re-docking of inosine (magenta) was performed for comparison with the ligand in the crystal structures (yellow). RMSD values are 0.035 Å for site A and 0.036 Å for site B.

Supplementary Figure 7. HPLC-MS quantification of ligands bound to TbPKAR in parasite lysates.
The left panels show western analysis of soluble fraction from lysed parasites expressing tagged TbPKAR or of control parasites (input), pulled down beads and supernatants (all in triplicate). Anti-PKAR, anti-PKAC1/2 and anti-His antibodies were used. Bound ligands were released by boiling of Ni-NTA beads in water and the aqueous fraction subjected to HPLC-MS analysis. Graphs on the right show chromatograms for the selected mass [M+H$^+$] 0.003 m/z (given in parenthesis) for inosine, guanosine and adenosine (all red coloured), and the matched stable isotope labelled internal standards (all blue coloured). Where peak quantification was in the linear range of the standard curves, m ± SD pmole of nucleoside per biological replicate of 8 x 10^8 or 2 x 10^8 cells is given for PCF or BSF, respectively (same values in Fig. 6). For graphical reasons the MS-signals were smoothened by Boxcar algorithm. **a** Procyclic stage (PCF) *T. brucei* strain EATRO1125 wild type cells and cells expressing His-TbPKAR, as indicated. Nucleoside amounts pulled down from wild type PCF were in the range of blanks. **b** Same as A for bloodstream stage *T. brucei* MITat 1.2 single marker Δtbpkar/Δtbpkar cells, wild type cells and cells expressing His-TbPKAR, as indicated. Only trace amounts of adenosine were detected.