1

The Hao-Fountain syndrome protein USP7 regulates neuronal connectivity in the brain via a novel p53-independent ubiquitin signaling pathway

5
6

Hao Chen[1,2], Cole J. Ferguson[2,3], Dylan C. Mitchell[4], Amanda Titus[5], Joao A. Paulo[4], Andrew Hwang[1], Tsen-Hsuan Lin[6], Hiroko Yano[1,7,8], Wei Gu[9], Sheng-Kwei Song[6], Carla M. Yuede[5], Steven P. Gygi[4], Azad Bonni[2,10]*, Albert H. Kim[1,7,8,11]*

[1]Department of Neurological Surgery, Washington University School of Medicine, St. Louis, MO 63110, USA

[2]Department of Neuroscience, Washington University School of Medicine, St. Louis, MO 63110, USA

[3]Department of Pathology, University of California, San Diego, La Jolla, CA 92093, USA

[4]Department of Cell Biology, Harvard Medical School, Boston, MA 02138, USA

[5]Department of Psychiatry, Washington University School of Medicine, St. Louis, MO 63130, USA

[6]Department of Radiology, Washington University School of Medicine, St. Louis, MO 63110, USA

[7]Department of Neurology, Washington University School of Medicine, St. Louis, MO 63110, USA

[8]Department of Genetics, Washington University School of Medicine, St. Louis, MO 63110, USA

[9]Institute for Cancer Genetics, Department of Pathology and Cell Biology, Columbia University, New York, NY 10032, USA

[10]Roche Pharma Research and Early Development, Neuroscience and Rare Disease discovery and translational area, Roche Innovation Center, Basel 4070, Switzerland

[11]The Brain Tumor Center, Siteman Cancer Center, Washington University School of Medicine, St. Louis, MO 63110, USA

*To whom correspondence should be addressed:

Albert H. Kim, alberthkim@wustl.edu

Azad Bonni, bonni@wustl.edu

Abstract

Precise control of protein ubiquitination is essential for brain development, and hence, disruption of ubiquitin signaling networks can lead to neurological disorders. Mutations of the deubiquitinase USP7 cause the Hao-Fountain syndrome (HAFOUS), characterized by developmental delay, intellectual disability, autism, and aggressive behavior. Here, we report that conditional deletion of USP7 in excitatory neurons in the mouse forebrain triggers diverse phenotypes including sensorimotor deficits, learning and memory impairment, and aggressive behavior, resembling clinical features of HAFOUS. USP7 deletion induces neuronal apoptosis in a manner dependent of the tumor suppressor p53. However, most behavioral abnormalities in USP7 conditional mice persist despite p53 loss. Strikingly, USP7 deletion in the brain perturbs the synaptic proteome and dendritic spine morphogenesis independently of p53. Integrated proteomics analysis reveals that the neuronal USP7 interactome is enriched for proteins implicated in neurodevelopmental disorders and specifically identifies the RNA splicing factor Ppil4 as a novel neuronal substrate of USP7. Knockdown of Ppil4 in cortical neurons impairs dendritic spine morphogenesis, phenocopying the effect of USP7 loss on dendritic spines. These findings reveal a novel USP7-Ppil4 ubiquitin signaling link that regulates neuronal connectivity in the developing brain, with implications for our understanding of the pathogenesis of HAFOUS and other neurodevelopmental disorders.

Keywords

ubiquitin, deubiquitinase, USP7, the Hao-Fountain syndrome, p53, brain development, glutamatergic neuron, synapse, TMT proteomics, Ppil4

Introduction

As a fundamental post-translational modification, protein ubiquitination plays critical roles in neuronal development and function, and deregulation of ubiquitination is thought to contribute to the pathogenesis of diverse neurological conditions including neurogenetic disorders. For instance, perturbation of ubiquitin signaling has been implicated in distinct genetic disorders of brain development including Angelman[1], Gordon-Holmes[2,3] and Ferguson-Bonni[4] syndromes. Substrate ubiquitination is regulated by both ubiquitin ligases, which catalyze the covalent attachment of the 76 amino acid-small ubiquitin protein to specific substrates, and deubiquitinases, which catalyze the removal of ubiquitin from specific substrates. Although ubiquitin ligases have been extensively studied for their roles in neurodevelopment[5], the functions of most deubiquitinases in the nervous system have remained largely unexplored[6], despite their link with neurodevelopmental disorders in genetic association studies[7]. We herein characterized the role of the deubiquitinase ubiquitin specific protease 7 (USP7) in the mammalian brain. Deletion or mutation of *USP7* causes the Hao-Fountain syndrome, which is characterized by developmental delay, intellectual disability, autism, and aggressive behavior[8,9]. However, the neuropathogenesis of the Hao-Fountain syndrome has remained poorly understood.

Studies in cancer biology have revealed that USP7 induces the turnover of the tumor suppressor protein p53[10,11], and inhibition of USP7 may hold therapeutic promise for suppressing tumor growth[12,13]. A key target of USP7 is the ubiquitin ligase Mdm2, which ubiquitinates p53 and thereby promotes p53 degradation. USP7 counteracts the effects of Mdm2 auto-ubiquitination to stabilize Mdm2 and promote Mdm2-dependent ubiquitination and consequent degradation of p53[11]. In mice, global mutation of *Usp7* leads to early embryonic lethality, indicating the essential role of USP7 in embryogenesis[14]. Loss of USP7 in neural stem cells leads to smaller brain size, perinatal lethality, and p53 accumulation[15]. Although mutation of *Trp53*, the gene encoding p53 in mice, rescues brain size in *Usp7* mutant mice, co-deletion of p53 fails to improve newborn survival, indicating that USP7 plays essential roles in the nervous system through unknown p53-independent mechanisms[15]. Importantly, whereas USP7 function in proliferating cells has been the subject of intense scrutiny[16–18], the roles and mechanisms of USP7 in post-mitotic neurons that might be relevant to the pathogenesis of the Hao-Fountain syndrome have remained largely unknown.

In this study, we conditionally knock out *Usp7* in post-mitotic glutamatergic neurons of the forebrain in mice. We find that USP7 mutant mice display distinct behavioral deficits that recapitulate many clinical features of the Hao-Fountain syndrome. We also find that USP7 loss induces neuronal apoptosis in the cerebral cortex during brain development in a p53-dependent manner. However, genetic deletion of p53 fails to rescue most behavioral deficits in USP7 mutant mice, suggesting that USP7 regulates brain development via p53-independent mechanisms. Accordingly, proteomic and dendritic spine morphological analyses reveal marked reduction in synapse formation in USP7 mutant mice independently of p53. An unbiased protein interaction and quantitative TMT-proteomic approach reveals, for the first time, the landscape of potential USP7 substrates in neurons, including the the SWI/SNF chromatin remodeling and spliceosomal complexes, implicating USP7 in fundamental aspects of chromatin and gene regulation in neurons. Specifically, we identify the RNA splicing factor Ppil4 as a novel functional substrate of USP7. Importantly, knockdown of Ppil4 recapitulates the dendritic spine phenotype observed in USP7 mutant mice. These findings define a novel ubiquitin signaling pathway, which links the deubiquitinase USP7 with the RNA splicing factor Ppil4 and drives neuronal connectivity in the developing brain, shedding light on the roles and mechanisms of USP7 in post-mitotic neurons, with important implications for our understanding of the neuropathogenesis of the Hao-Fountain syndrome.

111
112
113 **Results**
114
Conditional deletion of USP7 in glutamatergic excitatory neurons alters mouse behavior
To characterize the role of USP7 in post-mitotic glutamatergic neurons of the forebrain,
we crossed mice harboring a floxed allele of *Usp7* (*Usp7*$^{fl/fl}$) with mice expressing the
recombinase Cre under the control of the regulatory sequence of the mouse *NEX* gene (*NEX-*
cre)[19] (Figure 1A). Immunofluorescence (IF) analyses demonstrated loss of USP7 in NeuN-
positive cortical neurons of *Usp7*$^{fl/fl}$; *NEX-cre* mice (hereafter *Usp7* cKO) (Figure 1B). USP7 loss
occurred in post-mitotic neurons of the cortical plate but not in the ventricular zone, confirming
that USP7 is specifically deleted in post-mitotic neurons (Figure S1A).
123
Patients with *USP7* mutation or deletion exhibit a range of deficits early in life that include
growth retardation, hypotonia, visual impairment, gait abnormalities, aggression, and autism
with intellectual disability[9] (Figure 1A). To ascertain the effects of USP7 deletion on the
organism level, we subjected *Usp7* cKO (*Usp7*$^{fl/fl}$; *NEX-cre*), *Usp7* cHet (*Usp7*$^{fl/+}$; *NEX-cre*), and
Usp7 WT (*Usp7*$^{fl/fl}$) mice to a battery of behavioral tests and biometric measurements (Figure
1C-K and S1D-K). Although *Usp7* cKO mice exhibited small size during the juvenile period, the
body weight of adult *Usp7* cKO mice was similar to controls (Figure S1B and S1C). This pattern
of temporary growth delay is commonly seen in genetic mouse models of neurodevelopmental
disorders[20] and indicates that loss of USP7 in the brain is sufficient to generate phenotypes
throughout the body.
134
Visual deficits are common in the Hao-Fountain syndrome[9]. To examine visual function in *Usp7*
cKO mice, we employed the virtual optomotor system (VOS). In this experiment, mice make
reflexive head movements to track the rotation of an alternating pattern of black and white
vertical stripes, the spatial frequency of which can be experimentally varied. We found that *Usp7*
cKO mice stopped responding at a much lower spatial frequency, suggesting reduced visual
acuity (Figure 1C). This result shows that USP7 depletion in glutamatergic neurons is sufficient
to compromise vision in mice.
142
Hypotonia is a motor impairment associated with the Hao-Fountain syndrome[8,9]. To assess
motor performance in *Usp7* cKO mice, we began by performing the inverted screen test. In this
strength test, *Usp7* cKO mice exhibited a significant reduction in the latency to fall (Figure 1D).
We evaluated motor coordination using the ledge, platform, and pole tests. *Usp7* cKO mice also
displayed a reduction in the latency to fall in the ledge and platform tests (Figure S1D and S1E)
and were slower to turn around on the pole. (Figure S1G). However, mice of all genotypes spent
similar amounts of time climbing down the pole (Figure S1F). These observations indicate that
USP7 loss in the forebrain impairs motor strength and coordination without effects on simple
motor tasks.
152
In the course of characterizing *Usp7* cKO mice, we noticed that they exhibited obvious gait
abnormalities, another feature of USP7 mutation/deletion in humans[9]. During voluntary
locomotion, *Usp7* cKO mice exhibited abnormal splaying of the hindlimbs (Video S1). To
systematically evaluate this phenotype in *Usp7* cKO mice, we employed the CatWalk XT system
to deeply analyze gait and detect subtle abnormalities (Video S2). These experiments
demonstrated that *Usp7* cKO mice place extra weight onto their hind paws (Figure 1E). This
analysis also showed an unexpected increase in the swing speed of the limbs of *Usp7* cKO
mice (Figure 1F, S1H and S1I). Together, these results demonstrate that USP7 depletion in

glutamatergic neurons of the forebrain leads to diverse but specific motor deficits in mice that
mimic the motor phenotypes associated with the Hao-Fountain syndrome.
163
Having characterized the effect of conditional USP7 knockout on sensorimotor function, we next
evaluated whether *Usp7* cKO mice exhibit deficits in tests of learning and memory. To avoid the
potential confounding effect of visual impairment in *Usp7* cKO mice, we chose the fear
conditioning paradigm using an auditory cue. We confirmed that hearing was not significantly
impaired in *Usp7* cKO mice using the acoustic startle test (Figure S1J). On day 1, mice were
exposed to an auditory tone (a cue) in a peppermint-scented chamber (the context) prior to the
delivery of a mild electrical shock to the paw (Figure 1G). On day 2, during re-exposure to the
context environment, *Usp7* cKO mice exhibited reduced freezing behavior (the conditioned
response) (Figure 1H), suggesting diminished association between the context (the conditioned
stimulus) and the shock (the unconditioned stimulus) in *Usp7* cKO mice. On day 3, upon re-
exposure to the auditory cue (the conditioned stimulus) in a distinct context environment, *Usp7*
cKO also exhibited diminished freezing behavior (the conditioned response) (Figure 1I). In
contrast, in spite of their visual impairment, *Usp7* cKO mice showed no deficits in the Y-maze
test of spatial working memory (Figure S1K). Collectively, these results demonstrate that USP7
depletion in glutamatergic neurons of the forebrain causes specific deficits in behavioral tests of
associative learning.
180
Finally, *Usp7* cKO mice displayed striking forms of aggressive behavior characterized mainly by
violent attacks on cage mates that resulted in wounds and occasionally amputation of the other
animal's tail (Video S3). Between the ages of 3 and 12 weeks, 7 out of 21 *Usp7* cKO mice
inflicted bite marks centered around the tails of co-caged mice (Figure 1J and 1K). This
behavior is highly unusual and distinct from the commonly observed biting that is associated
with social aggression, which occurs after sexual maturity and typically targets the lower back
and abdomen[21].
188
In summary, this combination of observational and hypothesis-driven behavioral analyses
demonstrates that loss of USP7 in glutamatergic neurons of the forebrain causes a suite of
deficits that closely mimics the clinical hallmarks of the Hao-Fountain syndrome, including the
unusual phenotype of aggression[8,9]. These results suggest that USP7 may play a conserved
role in the brain in humans and mice. Therefore, the investigation of *Usp7* cKO mice should
advance our understanding of ubiquitin signaling during brain development and the
pathogenesis of neurodevelopmental disorders.
196
USP7 prevents p53-dependent neuronal apoptosis
After determining that loss of USP7 in the forebrain triggers diverse behavioral deficits in
mice, we sought to determine how USP7 regulates neuronal development in the cerebral cortex.
Therefore, we performed histological analyses at different stages of neurodevelopment,
beginning at embryonic day 12 (E12) to coincide with the onset of *NEX-Cre* expression[19]. The
earliest phenotype that we detected was a marked increase in the number of cells that were
positive for the apoptosis marker protein cleaved caspase 3 (c-cas 3) by immunofluorescence in
the cerebral cortex in *Usp7* cKO mice (Figure 2A). Importantly, this phenotype was temporally
restricted in the cerebral cortex of *Usp7* cKO mice to the postnatal day 0 (P0) timepoint, and
little difference in c-cas 3 was present at P5, P10 or E15 in control and *Usp7* cKO mice. These
results show that USP7 loss in cortical glutamatergic neurons leads to a dramatic wave of
apoptosis during the perinatal period.
209
To determine whether neuronal apoptosis in the cerebral cortex of *Usp7* cKO mice is cell-
autonomous, we used immunofluorescence to simultaneously detect Cre and c-cas 3. This

analysis demonstrated considerable overlap between Cre and c-cas 3 (Figure S2A), suggesting
that apoptosis occurs specifically in Cre-expressing glutamatergic neurons. In other experiments,
RNA interference (RNAi)-induced knockdown of USP7 sparsely in primary cortical neurons
using calcium phosphate transfection robustly induced apoptosis (Figure S2B-D). Together,
these results suggest that USP7 loss triggers neuronal apoptosis in a cell-autonomous manner.
217
Because USP7 regulates the Mdm2-p53 pathway in cancer and proliferating cells[9,10,14] (Figure
2B), we asked whether the apoptosis in post-mitotic neurons upon USP7 knockout in the mouse
cerebral cortex is mediated by p53. We crossed the *Usp7* cKO mice with mice harboring a null
allele in *Trp53*, the gene that encodes p53 in mice. Notably, the loss of just one *Trp53* allele in
Usp7 cKO mice reduced the c-cas 3 signal to a level that was comparable to control mice
(Figure 2C and 2D). Homozygous knockout of *Trp53* rescued apoptosis in the cortex of *Usp7*
cKO mice to a comparable degree as for heterozygous loss of p53 (Figure S3A). Together,
these results demonstrate that p53 is required for neuronal apoptosis in *Usp7* cKO animals
(Diagram as Figure 2B).
227
To test whether the rescue of apoptosis in *Usp7* cKO neurons by p53 loss is cell-autonomous,
we sought to assess the effect of p53 depletion on apoptosis in primary cultures of cortical
neurons. We began by isolating neurons from E15 mouse embryos harboring floxed *Usp7*
(*Usp7*$^{fl/fl}$). Following transduction with a lentivirus encoding Cre, immunocytochemical analyses
confirmed loss of USP7 upon Cre transduction (Figure S3B). In contrast, neurons transduced
with a catalytically dead Cre-Y331F mutant (Cre dead) failed to show depletion of USP7. In
cortical neurons dissociated from *Usp7*$^{fl/fl}$; *Trp53*$^{+/+}$ embryos, loss of USP7 following wild-type
Cre (Cre WT) transduction substantially reduced cell viability in the MTS assay. In contrast,
neurons from *Usp7*$^{fl/fl}$; *Trp53*$^{+/-}$ and *Usp7*$^{fl/fl}$; *Trp53*$^{-/-}$ embryos demonstrated no significant
decrease in viability upon Cre WT expression (Figure S3C). These results suggest that p53
deletion rescues USP7-dependent cell death in a cell-autonomous manner.
239
Our observation that deletion of USP7 triggers neuronal apoptosis led us to ask whether this
cellular phenotype might translate to a reduction in brain volume. Using diffusion-weighted *ex*
vivo MRI, we found that the volumes of both the cerebral cortex and the hippocampal formation
were dramatically reduced by ~40% in *Usp7* cKO mice (Table 1, Figure S3D and S3E).
Surprisingly, although p53 loss to rescued apoptosis in the brains of *Usp7* cKO mice, co-
deletion of p53 in *Usp7*$^{fl/fl}$; *NEX-cre*; *Trp53*$^{+/-}$ and *Usp7*$^{fl/fl}$; *NEX-cre*; *Trp53*$^{-/-}$ mice had minimal
effect on the volumes of the cerebral cortex or hippocampus (Table 1, Figure S3D and S3E). In
agreement with these observations in the forebrain, total brain volume was reduced by ~20% in
both *Usp7*$^{fl/fl}$; *NEX-cre*; *Trp53*$^{+/+}$ and *Usp7*$^{fl/fl}$; *NEX-cre*; *Trp53*$^{+/-}$ mice (Table 1, Figure S3D and
S3E). These results suggest that p53-mediated apoptosis does not significantly contribute to the
reduction of brain volume in USP7 cKO mice, indicating the importance of p53-independent
pathways to this phenotype.
252
p53 reduction rescues a subset of behavioral abnormalities in *Usp7* cKO mice
To determine whether dysregulation of p53 contributes to behavioral deficits in *Usp7*
cKO mice, we measured biometrics and tested motor performance and memory formation in
mice of the genotype *Usp7*$^{fl/fl}$; *NEX-cre*; *Trp53*$^{+/-}$ (hereafter *Usp7* cKO; *Trp53*$^{+/-}$) (Figure 2E-M
and S4A-I). The growth of *Usp7* cKO; *Trp53*$^{+/-}$ juvenile mice remained delayed compared to
control mice of the genotypes *Usp7*$^{+/+}$; *NEX-cre*; *Trp53*$^{+/-}$ (hereafter *Usp7* WT; *Trp53*$^{+/-}$) and
Usp7$^{fl/+}$; *NEX-cre*; *Trp53*$^{+/-}$ (hereafter *Usp7* cHet; *Trp53*$^{+/-}$) (Figure S4A and S4B). These findings
suggest that juvenile growth delay that occurs upon USP7 loss in the forebrain does not require
p53 as an intermediary. In contrast, *Usp7* cKO; *Trp53*$^{+/-}$ mice demonstrated comparable visual
acuity to *Usp7* WT; *Trp53*$^{+/-}$ mice, suggesting that visual impairment observed in *Usp7* cKO

mice requires p53 (Figure 2E). Assessment of motor function in *Usp7* cKO; *Trp53*$^{+/-}$ mice
demonstrated persistent impairment in the inverted screen test (Figure 2F), suggesting that
motor weakness associated with USP7 depletion in the brain persists despite p53 depletion.
Similarly, *Usp7* cKO; *Trp53*$^{+/-}$ mice were also impaired in the ledge, platform, and pole tests
(Figure S4C-F). p53 deletion also failed to normalize the performance of *Usp7* cKO mice in the
CatWalk, as *Usp7* cKO; *Trp53*$^{+/-}$ also improperly distributed their body weight and accelerated
the swing speed of their limbs during voluntary locomotion (Figure 2G, 2H, S4G-H and Video
S4). Tail biting behavior was also present in *Usp7* cKO; *Trp53*$^{+/-}$ mice (Figure 2L and 2M).
These results indicate that the motor impairment and aggression phenotypes in *Usp7* cKO mice
are largely independent of p53 loss.

Finally, we analyzed the effect of p53 loss on the performance of *Usp7* cKO mice during
associative learning paradigms. Similar to *Usp7* cKO mice, *Usp7* cKO; *Trp53*$^{+/-}$ mice exhibited
decreased freezing during contextual conditioning (Figure 2I and 2J). Prior to performing cued
conditioning, we used the acoustic startle test to confirm that hearing was unimpaired in *Usp7*
cKO; *Trp53*$^{+/-}$ mice (Figure S4I). During cued fear conditioning, the impairment in the freezing
response to the auditory cue in *Usp7* cKO mice was corrected in mice of the genotype *Usp7*
cKO; *Trp53*$^{+/-}$ (Figure 2K). Altogether, our behavioral analyses demonstrate that whereas p53
contributes to USP7 loss-induced impairment of vision and cued conditioning, most behavioral
deficits in *Usp7* cKO mice are independent of p53 reduction and its effects on apoptosis (Table
1).

Loss of USP7 reduces synapses in cortical neurons
After observing that most developmental phenotypes associated with USP7 mutation did
not require p53, we attempted to identify USP7-dependent phenotypes in neurons beyond p53-
mediated apoptosis. To probe cellular and molecular abnormalities upon USP7 depletion in the
developing brain, we profiled the proteome of the cerebral cortex of *Usp7* cKO mice using
unbiased tandem mass tag-mass spectrometry (TMT-MS)-based proteomic methods (Figure
3A). We included cortices from 5 pairs of sex-matched *Usp7* cKO mice and their control
littermates at age P17. Protein lysates were trypsinized and labeled with individual tandem
mass tag prior to quantitation of relative protein abundance using MS3. We sorted the
differentially abundant proteins based on fold change and p-value in the unpaired t-test (Figure
3B). Although more proteins were up-regulated than down-regulated by loss of USP7, gene set
enrichment analysis (GSEA) showed the most significantly enriched gene sets were among the
down-regulated proteins (Figure 3C and Table S1). Importantly, synapse-related gene sets were
prominent among the enriched gene sets (Figure 3C and Table S1). Additional GO analysis
using the GOrilla algorithm[22] also identified synapse-related terms among the ranked list of
proteins whose abundance was reduced by USP7 loss (fold change < 0.75) (Figure 3D). Further
SynGO analysis[66] revealed that both postsynapse and presynapse terms are enriched in the
down-regulated proteins by USP7 loss (Figure S5A&B). The down-regulated presynaptic
proteins include Cplx3, Kcnj11, Il1rapl2 and Rpl35, and down-regulated postsynaptic proteins
include Arc, Rtn4, Rnf220 and Dclk1 (Figure S5C). These results suggest that USP7 may
regulate synaptic proteins in the cerebral cortex.

To test whether USP7 loss impairs synapse development, we next characterized the effects of
USP7 loss on dendritic spines in glutamatergic neurons *in vivo*. We employed adeno-associated
virus (AAV)-PHP.eB harboring a syn-GFP-P2A-GFPf construct to drive expression of green
fluorescent protein (GFP) in neurons directed by the synapsin promoter. AAV was injected into
the ventricles of newborn *Usp7* WT and *Usp7* cKO mouse pups at titers that produced sparse
labeling of neurons, highlighting their dendritic spine morphology with cytoplasmic GFP and
membrane bound GFPf (Diagram as Figure 3E). To avoid the influence of p53-mediated

apoptosis in neurons, we analyzed the morphologic effects of USP7 loss on the *Trp53*[+/-]
background. Spine density within both apical and basal dendrites of pyramidal neurons in the
motor cortex of *Usp7* cKO; *Trp53*[+/-] mice at P18 was significantly reduced compared to that of
Usp7 WT; *Trp53*[+/-] mice (Figure 3F-I). In spine shape analyses, filopodial spines were increased
in *Usp7* cKO; *Trp53*[+/-] neurons as compared to *Usp7* cKO; *Trp53*[+/-] neurons (Figure S5D-J). We
also found that dendrite length was unaffected in Cre-transduced *Usp7*[fl/fl]; *Trp53*[+/-] primary
cortical neurons, and there was no effect on dendrite arborization by Sholl analysis upon USP7
loss (Figure S6A-F). These results demonstrate that the loss of USP7 specifically reduces
dendritic spines of pyramidal neurons in the developing mouse brain.
323
Defining the interactome and candidate substrates of USP7 in neurons
Having identified critical p53-independent roles for USP7 in regulating dendritic spines in
the developing mouse cerebral cortex, we next characterized the mechanism by which USP7
controls synapse formation in post-mitotic neurons. Because USP7 substrates in cancer cells
frequently bind to the TRAF domain of USP7[18,23] (Figure 4A), we reasoned that neuronal USP7
substrates might also interact with the TRAF domain of USP7 and thus characterized the
interactors of the TRAF domain in primary cortical neurons. In co-immunoprecipitation (co-IP)
analyses, we confirmed that the truncated USP7 TRAF domain interacts with the known USP7
substrate Mdm2. Alanine substitution mutations at key residues of the TRAF domain including
Arg104 (R104A), Asp165 (D165A) and Trp166 (W166A), collectively termed the RDW mutant,
disrupted the interaction between the TRAF domain and Mdm2 (Figure S7A).
Immunocytochemical analyses in primary cortical neurons confirmed the mostly nuclear
localization of both the wild-type TRAF domain and the RDW mutant TRAF domain (Figure S6B
and S7C). Silver staining of lysates from primary cortical neurons expressing lentiviral FLAG-
tagged TRAF domain showed a large number of proteins that co-immunoprecipitated with the
wild-type TRAF domain. Most of these putative interactions were lost by RDW mutations of the
TRAF domain. (Figure 4B). Using the RDW mutant as a negative control, we performed mass
spectrometry analysis of interactors of the USP7-TRAF domain (IP-MS) in 3 biological replicates
of primary cortical neurons. We ranked interactors with the wild-type TRAF domain in neurons
according to protein length-normalized spectral counts and generated an association network
of the top 50 interactors based on the STRING (Search Tool for the Retrieval of Interacting
Genes/Proteins) database. GO terms related to cellular components revealed a few densely
interconnected structures on the STRING network that included ribonucleoprotein complexes,
the RNA spliceosome, and the SWI/SNF (BAF) chromatin remodeling complex (Figure 4C and
4D). These results indicate that USP7 might interact with and/or target substrates involved in
critical processes related to chromatin and RNA splicing in a TRAF domain-dependent manner.
350
Among interactors of wild-type USP7 TRAF domain, a number of proteins are genetically
associated with neurodevelopmental diseases (Ddx3x, Nbea, Smarcc2 and Map1b). Therefore,
we compared curated neurodevelopmental disease gene sets[24] with interactors of the USP7
TRAF domain. We found that USP7 TRAF interactors are most significantly enriched in the
gene sets for intellectual disability and autism (Figure 4E), which are also the most common
comorbilities in the Hao-Fountain syndrome[9]. Interestingly, USP7 TRAF interactors also had
strong enrichment in the high-confidence mRNA targets of Fragile X syndrome protein FMRP[68]
(Figure 4E). These observations suggest that USP7 may interact with and target proteins
implicated in neurodevelopmental disorders.
360
USP7 stabilizes its substrates by removing ubiquitin tags and preventing subsequent
degradation. Therefore, we also searched for candidate USP7 substrate by profiling protein
abundance using quantitative TMT-MS (Figure 4F). To minimize the influence of secondary
effects that could occur in the setting of extended loss of USP7, we acutely inactivated USP7 by

transducing primary cortical neurons isolated from mouse embryos of the genotype *Usp7*[fl/fl];
Trp53[-/-] with a lentivirus expressing the recombinase Cre. Immunoblotting analyses at different
time points after transduction confirmed depletion of USP7 in lysates of primary cortical neurons
(Figure 4G). Transduction and expression of catalytically dead Cre failed to induce USP7 loss
(Figure 4G). Based upon the timing of USP7 depletion in this system, we selected DIV6 and
DIV8 for TMT proteomics analysis. Consistent with the immunoblotting analyses, the TMT
abundance of USP7 protein decreased at both DIV6 and DIV8 following transduction of wild-
type Cre (Figure 4H). We identified and quantitated >8000 proteins, and principal component
analysis (PCA) over these proteins showed clear segregation into the different experimental
groups (Figure S8A). The first principal component accounted for ~20% of the variance and
separated samples at DIV6 from samples at DIV8. The second principal component accounted
for ~10% of the variance and separated the wild-type Cre versus dead Cre samples. Importantly,
the distance between wild-type Cre versus dead Cre samples increased from DIV6 to DIV8,
consistent with more complete depletion of USP7 at DIV8. These results demonstrate that acute
USP7 loss induces a progressive proteomic change over time in primary cortical neurons.

Because USP7 stabilizes its substrates, we reasoned that the abundance of USP7 should
correlate with the abundance of its substrates. Using this rationale to identify putative USP7
substrates, we calculated the Euclidian distance between USP7 and each protein detected in all
16 samples. We then analyzed the p-value of Cre-dependent effects for each protein using two-
way ANOVA. Based on these analyses, we sorted out candidate substrates with the shortest
distances to USP7 and the smallest p-values (Figure 4H). The abundance of each candidate
substrate was down-regulated by expression of wild-type Cre at both DIV6 and DIV8. As proof
of principle, this approach identified the known USP7 substrate Trim27[8]. Among proteins that
were down-regulated by USP7 loss was the SWI/SNF subunit Smarcal1, the RNA spliceosome
proteins Ppil4 and Fam172a, and several ubiquitin ligases including Trip12 and Ring1.

We next asked if the protein levels of these candidate substrates also decrease upon USP7
knockout *in vivo*. In our *in vivo* TMT analysis in the cerebral cortex of *Usp7* cKO mice, most of
the candidate substrates that we found to be down-regulated upon USP7 loss in primary
neurons also showed varying degrees of reduction in the cortex of P17 *Usp7* cKO mice (Figure
S8B). To characterize these candidate substrates, we performed immunoblotting analysis of the
P0 cererbal cortex of *Usp7* cKO; *Trp53*[+/-] mice. Performing these assays in the setting of *Trp53*
depletion minimized the effects of apoptosis on our analysis. This analysis confirmed that
several candidate substrates, including Ppil4, Ubxn7, Sub1, Smarcal1, Ring1, Fam172a and
Trip12, were less abundant in the cortex of *Usp7* cKO; *Trp53*[+/-] mice (Figure 4I and S9A-I). RT-
qPCR showed no evidence that the abundance of any of the corresponding mRNAs was
reduced by USP7 loss (Figure 4J), suggesting that USP7 regulates the abundances of these
candidate substrates at the protein level. Altogether, these results demonstrate that USP7
regulates the abundance of specific candidate substrates in primary neurons and in the brain.

To better understand the global molecular network that is regulated by USP7, we used weighted
correlation network analysis (WGCNA) to cluster proteins that become differentially abundant
upon USP7 loss. We used a cutoff p-value of 0.05 for determining both Cre-dependent and Cre-
time interaction effects using two-way ANOVA. WGCNA analysis identified 10 distinct network
modules (Figure S8C). The yellow module, which contained most of the candidate substrates,
showed down-regulation of its eigengene upon USP7 loss (Figure S8D and S8E). GO analysis
of proteins in the yellow module revealed an enrichment in ubiquitin ligases (Figure S8F).
Interestingly, ribonucleoprotein and splicing-related GO terms were overrepresented in the
turquoise module, which consisted of the largest set of differentially abundant proteins among
all the modules (Figure S8C and S8F). Collectively, findings from these unbiased analyses

suggest that USP7 transforms the neuronal proteome by regulating a diverse network of
proteins.
418
419   **USP7 targets Ppil4 to promote synaptogenesis**
420   After profiling the interactome and putative targets of USP7 in neurons, we sought to
further use these datasets to identify the biologically relevant USP7 targets in neurons. TMT
proteomics followed by immunoblotting validation *in vivo* revealed that USP7 stabilizes Ppil4,
Ubxn7, Sub1, Smarcal1, Ring1, Fam172a and Trip12, and IP-MS experiments identified a
number of these proteins and their assembled complexes (Ppil4, Smarcal1 and Trip12) as
interactors of USP7 TRAF domain. To confirm that these candidate substrates interact with full-
length USP7, we used HEK293T cells to express USP7 tagged with streptavidin-binding peptide
and FLAG (SFB) prior to pull-down with streptavidin beads (Figure 5A). Immunoblotting
detection of endogenous candidate substrates showed that Trip12, Smarcal1, Ubxn7, Ppil4,
Ring1 and Fam172a co-precipitated with full-length USP7. Trip12 and Ppil4 displayed the
strongest interaction with USP7, as demonstrated by their enrichment in the pull-down fraction
compared to in the input lysate.
432
Reduction in dendritic spine density was a prominent phenotype in neurons with USP7 loss
(Figure 3). Therefore, we asked whether select candidate substrates might regulate the
morphogenesis of dendritic spines. We tested the effect of knockdown of the candidate
substrates at DIV4 using lentiviral shRNA in wild-type primary cortical neurons (validated in
Figure S10A-E) before assessing dendritic spine formation at DIV18 (Figure 5B-D). We found
that knockdown of Ppil4 using two independent shRNAs significantly decreased the density of
dendritic spines (Figure 5B-D and 5H). Further immunofluorescence analysis of PSD95, a major
postsynaptic component of dendritic spines, confirmed the reduction of PSD95-positive spines
in neurons subjected to Ppil4 knockdown (Figure 5E-H and S10F-H). These observations
suggest that Ppil4 is required for the formation of mature, stable synapses. Surprisingly,
knockdown of Trip12, a ubiquitin ligase associated with intellectual disability[25,26], led to an
increase in spine density (Figure 5B-H and Figure S10F-I). Knockdown of other candidate USP7
substrates, including the SWI/SNF component Smarcal1 had little to no effect on spine density
(Figure 5B-H and Figure S10F-I). Taken together, these results suggest that Ppil4 promotes
synaptogenesis in cortical neurons, and depletion of Ppil4 is sufficient to phenocopy the effects
of USP7 loss.
449
We next determined whether Ppil4 is a direct target of USP7-mediated deubiquitination. First,
we found that endogenous Ppil4 interacts with GFP-tagged USP7 in HEK293T cells by co-
immmunoprecipitation (Figure 5I). Using the Tandem Ubiquitin Binding Entities (TUBE) assay[27],
we purified endogenous polyubiquitinated proteins at DIV5 from lysates of primary cortical
neurons of genotype $Usp7^{fl/fl}$; $Trp53^{+/-}$ transduced with Cre lentivirus at DIV3. Subsequent
immunoblotting analyses showed that, compared to catalytically dead Cre, wild-type Cre
lentiviral transduction triggered a dramatic increase of high-molecular weight forms of Ppil4,
indicative of ubiquitination of Ppil4 upon USP7 loss (Figure 5J). Finally, we found that
recombinant USP7 was sufficient to deubiquitinate FLAG-Ppil4 immunopurified from HEK293T
cells in an *in vitro* cell-free system (Figure 5K & 5L). Collectively, these observations
demonstrate that Ppil4 is a bona fide target of USP7 deubiquitinase activity, and that USP7 and
Ppil4 regulate the formation of dendritic spines and synapses of glutamatergic neurons
(Schematic as Figure 5P).
463
Lastly, to examine the relevance of Ppil4 as a USP7 substrate in the context of the Hao-
Fountain syndrome, we asked whether reported patient mutations in USP7 disrupt the USP7-
Ppil4 signaling pathway. Among the 5 USP7 patient mutations[28–31] (T415A, K420E, K657A,

Y749fs and P1006L) we tested, four were stably expressed in HEK293T cells (Figure 5M).
Subsequent immunoblotting analyses showed that the protein abundance of endogenous Ppil4
was reduced in HEK293T cells expressing either patient USP7 mutants or catalytically dead
USP7 (C223A), compared to cells expressing wild-type USP7 (Figure 5N & 5O). These results
suggest that USP7 mutants in the Hao-Fountain syndrome patients lead to destablization of
Ppil4, similar to the catalytically inactive USP7 mutant.

Discussion

In this study, we have discovered novel roles and underlying mechanisms for the Hao-Fountain
syndrome protein USP7 in brain development. Conditional knockout of the deubiquitinase USP7
in post-mitotic glutamatergic neurons of the forebrain leads to deficits in mice, from
sensorimotor dysfunction to abnormalities of cognition and social behavior, reminiscent of
phenotypes in the Hao-Fountain syndrome. In developmental studies, conditional USP7 loss
leads to perinatal apoptosis in post-mitotic glutamatergic neurons, and genetic deletion of p53
rescues the USP7 loss-induced apoptosis. Deletion of p53 also rescues impairments in visual
acuity and auditory cue fear conditioning in *Usp7* cKO mice. Strikingly, however, loss of p53
fails to rescue most behavioral deficits or the brain volume reduction upon loss of USP7,
indicating that USP7 regulates brain development and function predominantly in a p53-
independent manner. Accordingly, USP7 loss reduces the amounts of synaptic proteins in the
cerebral cortex and density of dendritic spines in cortical pyramidal neurons in the background
of p53 deficiency. In immunoprecipitation followed by mass spectrometry analyses in primary
neurons, we uncover the interaction of the USP7 TRAF domain, which binds substrates in
proliferating cells, with ribonucleoprotein and RNA spliceosome complex proteins in neurons.
Importantly, in quantitative TMT-proteomics analyses, the abundance of the ribonucleoprotein
and spliceosome proteins covaries with USP7 in control and USP7-deleted primary cortical
neurons. Among the spliceosome proteins, we identify the RNA splicing factor Ppil4 as a novel
substrate of USP7 that is deubiquitinated by USP7. Knockdown of Ppil4 phenocopies the effect
of USP7 loss on dendritic spine morphogenesis in cortical neurons. Our findings define a novel
ubiquitin signaling link between the deubiquitinase USP7 and the splicing factor Ppil4 that
operates independently of p53 to drive synapse morphogenesis in the developing brain, with
ramifications for our understanding of the Hao-Fountain syndrome.

Our findings have significant implications for studies of ubiquitin signaling networks as well as
neurodevelopmental disorders of cognition. The USP7-p53 pathway has been extensively
characterized in the context of cancer biology[10,11,32]. Our finding that USP7 loss induces p53-
dependent apoptosis in glutamatergic neurons broadens the relevance of the p53 pathway to
post-mitotic cells. However, despite the requirement for p53 in cortical neuron apoptosis in *Usp7*
cKO mice, the majority of the behavioral deficits in *Usp7* cKO mice persist independently of p53
deletion. Consistently, impairment of dendritic spine morphogenesis upon USP7 loss occurs
independently of p53 loss. The loss of p53 also fails to reverse the reduction in brain volume in
Usp7 cKO mice, suggesting that USP7 may regulate brain size in a p53-independent manner.
Because USP7 loss has little effect on soma size and dendrites (Figure S6), our data suggest
that the reduction of dendritic spines in USP7-deficient neurons may contribute to the reduced
volume of the cerebral cortex[33]. Whereas dendritic spines house excitatory, but not inhibitory,
synapses[34], our proteomic analyses in the cerebral cortex reveal that down-regulation of both
excitatory and inhibitory synapse proteins occurs upon USP7 depletion. This observation raises
the question of whether USP7 also regulates the development of inhibitory synapses in the
brain. Because abnormalities of synapse development are thought to underlie the pathogenesis

of neurodevelopmental disorders of cognition[35–37], USP7 regulation of dendritic spine
morphogenesis in glutamatergic neurons may play a key role in the pathogenesis of the Hao-
Fountain syndrome.
520
One key finding in this study is the identification of the RNA splicing factor Ppil4 as a novel
substrate of USP7 in post-mitotic neurons, which bears broad implications for studies of the
major deubiquitinase USP7. First, although much of what has been learned about USP7 has
been in the context of cancer biology where USP7 is thought to promote tumorigenesis through
the deubiquitination of the ubiquitin ligase Mdm2 and consequent degradation of p53[10,11], the
identification of Ppil4 as a novel substrate elucidates a novel mechanism in post-mitotic neurons
in the developing brain with relevance to our understanding of the Hao-Fountain syndrome.
Second, by deubiquitinating and consequent regulation of the levels of the protein Ppil4, our
findings suggest that USP7 may regulate RNA splicing in neurons, which will be the subject of
future studies. Our findings in interaction and quantitative TMT-proteomics analyses revealing
that USP7 interacts with and regulates the abundance of small nuclear ribonucleoproteins, the
RNA helicase protein Dhx, and the RNA regulatory protein Fam172a[38,39], further support a role
for USP7 in the control of RNA processing. Recent studies suggest that Ppil4 is recruited to
active spliceosomes to facilitate catalytic activation of the spliceosome by positioning the key
RNA helicase Prp2 downstream of the branching site adenosine of introns[40,41]. It will be
important to determine how USP7 regulation of Ppil4 ubiquitination might influence the catalysis
of RNA splicing in neurons. Third, the observation that Ppil4 is required for dendritic spine
morphogenesis, phenocopying the effect of USP7 loss in cortical neurons, raises the important
question of whether the impairment of RNA splicing might contribute to the pathogenesis of the
Hao-Fountain syndrome. Because alternative splicing in neurons is frequently associated with
the regulation of synaptic proteins including adhesion molecules and scaffold proteins[42–44], it
may be interesting in future studies to test whether the USP7-Ppil4 ubiquitin signaling pathway
regulates the splicing of these molecules to influence dendritic spinogenesis. Finally, whereas
the discovery of the USP7-Ppil4 ubiquitin signaling link plays a critical role in the brain, our
findings also raise the question of whether this mechanism operates in proliferating cells and in
the context of cancer.
547
Besides Ppil4, our mass spectrometry analyses also reveal the interactome of USP7 and
proteomic dynamics upon acute USP7 loss for the first time in post-mitotic neurons. To our
knowledge, this represents the first comprehensive characterization of potential substrates of a
deubiquitinase in neurons. Notably, we observed that multiple ubiquitin ligases interact with
USP7, and the protein abundance of these ubiquitin ligases is reduced upon USP7 loss.
Ubiquitin ligases characteristically undergo self-ubiquitination and self-degradation[11,45,46].
Therefore, USP7 might function as a positive regulator of these ubiquitin ligases by limiting their
self-ubiquitination and degradation. Interestingly, among these ubiquitin ligases, Trip12 and
Huwe1 are both associated with intellectual disability[25,26,47,48], thus suggesting that the regulation
of these ubiquitin ligases by USP7 may bear potential relevance to the Hao-Fountain syndrome.
Furthermore, our proteomic and biochemical analyses have also identified several chromatin
and transcription regulators as novel candidate substrates of USP7 in neurons. In particular, the
chromatin remodeling SWI/SNF class enzyme Smarcal1 interacts with USP7 and is stabilized
by USP7, suggesting a role for USP7 in regulating chromatin accessibility.
562
The characterization of the behavioral deficits in *Usp7* cKO mice reveals that these mice display
several phenotypes reminiscent of the clinical features of patients with the Hao-Fountain
syndrome, including impairment in visual function, motor strength, gait, cognition, and
aggression. Although patients with the Hao-Fountain syndrome typically exhibit germline
heterozygous USP7 deletion or mutation[8,9], the phenotypic similarities between patients and our

Usp7 cKO mice suggest a shared biology for USP7 in both humans and mice. Our data suggest
that USP7 dysfunction in glutamatergic excitatory neurons might be relevant to the
pathogenesis of the Hao-Fountain syndrome. In particular, *Usp7* cKO mice display a tail-biting
behavior, not seen to our knowledge in other mouse models, which may bear implications for
understanding of aggression in the Hao-Fountain syndrome. Furthermore, our results that both
USP7 loss and USP7 patient mutations reduce the level of the substrate Ppil4 suggest that
USP7 loss of function may represent a shared mechanism between *Usp7* cKO mice and
patients with the Hao-Fountain syndrome. Collectively, our findings suggest that loss of USP7
function in excitatory glutamatergic neurons in the forebrain triggers phenotypes that might be
relevant to the Hao-Fountain syndrome. In future studies, it will also be crucial to determine the
effect of USP7 deletion in other cell types, including interneurons and glial cells in the brain, and
their contributions to the pathogenesis of the Hao-Fountain syndrome.

In conclusion, we have elucidated the roles and mechanisms of the Hao-Fountain syndrome
protein USP7 in the mammalian brain. Loss of USP7 in glutamatergic excitatory neurons in the
forebrain triggers mouse phenotypes with similarities to phenotypes in the Hao-Fountain
syndrome patients. We have found that USP7 regulates the development and function of the
brain via p53-dependent and predominantly p53-independent mechanisms, leading to the
discovery of the USP7-Ppil4 ubiquitin signaling link that plays a key role in dendritic spine
morphogenesis in the developing brain. Beyond advancing our understanding of the major
deubiquitinase USP7 in normal brain development and function, our study provides clues for the
identification of biomarkers and potential treatment of patients with the Hao-Fountain syndrome.

Limitations of the study
Although our study provides fundamental new insights into the functions and mechanisms of
USP7 in the developing mammalian brain, a number of limitations should be considered.
Whereas mouse genetics provides a rigorous approach to defining functions and mechanisms
of the Hao-Fountain syndrome protein USP7 in a living organism, the relevance of mice to
studies of human disorders remains a limitation. In the future, it will be worthwhile to extend
these studies to human model systems including induced pluripotent stem cell (iPSC)-derived
cells and tissues. In our mechanistic studies, although we focused on Ppil4 as a key target of
USP7, in the future, it will be important to extend our studies to the investigation of additional
candidate substrates. Importantly, ubiquitination influences protein degradation as well as other
aspects of protein function independently of protein turnover. Our TMT proteomics analyses and
follow-up biochemical experiments focused on the degradation pathway. Hence, it remains to be
determined whether USP7 impacts signaling pathways in a manner independent of its effects on
protein turnover.

Acknowledgements
The authors would like to thank the Kim lab and Bonni lab members for critical feedback. The
authors also thank A. Yen for the help with AAV injection and N. Mosammaparast and N. Tsao
for the help with TUBE assay. This work was supported by: NIH/NINDS R01NS051255 (to A.B.
and A.H.K), R01 NS111014 (to H.Y.), NIH/NIGMS R01 GM132129 (J.A.P.) and GM67945
(S.P.G.), a McDonnell Center for Systems Neuroscience grant award (to A.H.K.), the
Christopher Davidson and Knight Family Fund (to A.H.K.), and the Duesenberg Research Fund
(to A.H.K.). This work was also supported by the Washington University Center for Cellular
Imaging (S10 OD21629-01A1-WUCCI) and Hope Center Viral Vectors Core at Washington
University School of Medicine.

Author contributions

HC, CJF, AB and AHK designed the project and all the experiments. DM and SPG performed
the TMT experiments. JP and SPG performed the IP-MS experiment. AT and CMY performed
and analyzed the behavioral experiments. AH analyzed the *in vivo* apoptosis and dendritic spine
shape experiment. THL and SKS performed the *ex vivo* MRI and volumetric measurement. HY
contributed to DNA cloning. WG provided the USP7$^{fl/fl}$ mouse line. HC performed and analyzed
the rest of the experiments and wrote the manuscript. CJF, AB and AHK edited the manuscript.

Declarations of interest

The authors declare no competing interests. A.H.K. is a consultant for Monteris Medical and has
received a research grant from Stryker to study a dural substitute, both of which have no direct
relation to this study. A.B. is a full-time employee and shareholder of F. Hoffmann-La Roche Ltd.

628
629
630

631 **STAR Methods**
632
633 **Key resources table**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Rabbit anti-USP7	Bethyl labs	Cat# A300-033A
Rabbit anti-USP7	Bethyl labs	Cat# IHC-00018
Mouse anti-NeuN	Millipore	Cat# MAB377
Rat anti-Ki67 (SolA15)	Thermo Fisher Scientific	Cat# 13-9910-82
Rabbit anti-Cleaved caspase 3	Cell signaling	Cat# 9661
Chicken anti-GFP	Abcam	Cat# ab13970
Rabbit anti-GFP	Homemade	N/A
Mouse anti-mCherry (1C51)	Novus Biological	Cat# NBP1-96752
Mouse anti-Cre (2D8)	Millipore	Cat# MAB3120
Rabbit anti-Cre	Millipore	Cat# 69050
Mouse anti-Myc (9e10)	Santa Cruz Biotechnology	Cat# sc-40
Mouse anti-FLAG (M2)	Sigma	Cat# F3165
Mouse anti-FLAG M2 agarose beads	Sigma	Cat# A2220
Rabbit anti-HA (C29F4)	Cell Signaling	Cat# 3724
Rabbit anti-Map2	Abcam	Cat# ab32454
Rabbit anti-Phip	Bethyl labs	Cat# A302-054A
Rabbit anti-Xiap	Bioss Antibodies	Cat# bs-1281R
Rabbit anti-Fto	Abclonal	Cat# A1438
Rabbit anti-Rnf220	Abclonal	Cat# A15853
Rabbit anti-Cfdp1	Bethyl labs	Cat# A305-624A
Rabbit anti-Phc2	Abclonal	Cat# A14727
Rabbit anti-Ppil4	Bethyl labs	Cat# A304-965A
Rabbit anti-Ppil4	Sigma	Cat# HPA031600
Rabbit anti-Gpd1	Abclonal	Cat# A5715
Rabbit anti-Smarcal1	Millipore	Cat# ABE1836
Rabbit anti-Ubxn7	Bethyl labs	Cat# A303-865A
Rabbit anti-Rnf2	Abclonal	Cat# A5563
Rabbit anti-Trip12	Abclonal	Cat# A9958
Rabbit anti-Fam172a	Novus Biological	Cat# NBP1-86774
Rabbit anti-Sub1	Cohesion Biosciences	Cat# CQA1533
Rabbit anti-Chfr	Abclonal	Cat# A10447
Mouse anti-Ruvbl2	Santa Cruz	Cat# sc-374135
Mouse anti-Ring1 (8C12F4)	Novus Biological	Cat# NBP2-37370
Mouse anti-Gfap	Millipore	Cat# MAB3402
Mouse anti-14-3-3	Santa Cruz Biotechnology	Cat# sc-25276
Mouse anti-Lamin A/C (4C11)	Cell signaling	Cat# 4777
Mouse anti-PSD95 (K28/43)	NeuroMab	Cat# 75-028
Normal Rabbit IgG	Cell signaling	Cat# 2729
Bacterial and virus strains		
Lentivirus	This study	N/A
AAV PHP.eB	Hope Center Viral Vectors	N/A

	Core at Washington University	
Chemicals, peptides, and recombinant proteins		
MG132	Sigma	Cat# 474790
Bradford reagent	Bio-Rad	Cat# 5000006
Protease inhibitor cocktail	Sigma	Cat# P8340
Phosphatase inhibitor mini-tablet	Thermo Fisher Scientific	Cat# A32957
Hoechst 33258 solution	Sigma	Cat# 94403
3x FLAG peptide	APExBio	Cat# A6001
Dynabeads™ MyOne™ Streptavidin T1	Invitrogen	Cat# 65601
TUBE 2 magnetic beads	LifeSensors	Cat# UM-0402M
Recombinant His-USP7	R&D Systems	Cat# E-519
Deposited data		
IP mass spectrometry data	ProteomeXchange	PXD043225
TMT proteomics data	ProteomeXchange	PXD043848
TMT proteomics data	ProteomeXchange	PXD043849
Experimental models: cell lines		
HEK293T/HEK293LE	Clontech	Cat# 632180
Experimental models: organisms/strains		
NEX-cre mice on C57BL/6J	Goebbels et al., 2006	N/A
USP7 fl/fl mice on hybrid C57BL/6J and 129Sv	Kon et al., 2011	N/A
USP7 fl/fl mice on C57BL/6J	This study	N/A
p53 KO mice on C57BL/6J	The Jackson Laboratory	Cat# 002101
Wild-type outbred mice on CD1	Charles River	Cat# 022
Wild-type inbred mice on C57BL/6J	The Jackson Laboratory	Cat# 000664
Wild-type outbred Long-Evans rats	Charles River	Cat# 006
Oligonucleotides		
See Table S2	N/A	N/A
Recombinant DNA		
pCMV-myc3-Mdm2	Addgene	Cat# 20935
pAAV-hSyn1-EGFP-P2A-EGFPf-WPRE-HGHpA	Addgene	Cat# 74513
pMH-SFB-USP7	Addgene	Cat# 99393
pEGFP-USP7	Anckar et al., 2015	N/A
pEGFP-USP7 1-209	This study	N/A
pEGFP-USP7 55-209 (TRAF)	This study	N/A
pEGFP-USP7 C223A (catalytically dead)	This study	N/A
pEGFP-USP7 T415A	This study	N/A
pEGFP-USP7 K420E	This study	N/A
pEGFP-USP7 P657A	This study	N/A
pEGFP-USP7 Y749fs	This study	N/A
pEGFP-USP7 P1006L	This study	N/A

pHAGE-hSyn-FLAG-USP7 55-209 (TRAF)	This study	N/A
FCIV.1-Cre-IRES-mCherry	This study	N/A
FCIV.1-Cre Y331F-IRES-mCherry (Cre dead)	This study	N/A
pcDNA3.1-FLAG-Ppil4	This study	N/A
Software and algorithm		
ImageJ-Fiji	National Institutes of Health, USA	https://github.com/fiji/fiji
Zen	Zeiss	N/A
Imaris	Oxford Instruments	N/A
Image Studio	Li-Cor	N/A
Proteomics software pipeline #1	SEQUEST	http://fields.scripps.edu/yates/wp/?page_id=17
Proteomics software pipeline #2	Comet	https://uwpr.github.io/Comet/
GSEA (v4.3.2)	UC San Diego, Broad Institute	https://www.gsea-msigdb.org/gsea/index.jsp
Cytoscape (v3.9.1)	National Institute of General Medical Sciences	https://cytoscape.org/
ITK-SNAP (v3.8)	NYU, Tanden School of Engineering	www.itksnap.org
Prism (v.9)	GraphPad	N/A
SPSS (v.26)	IBM	N/A
R	The R project	N/A

Data and materials availability

All mouse lines, cell lines, purified protein or any other reagent generated or used in this study will be made available to researchers who are interested. The proteomics mass spec datasets (IP-MS, *in vivo* TMT and *in vitro* TMT) were deposited at ProteomeXchange via the PRIDE database and will be publicly available upon publication. Accession numbers for these datasets are listed in the Key resources table. Any additional information required to reanalyze the data reported in this paper is available from the corresponding authors upon request.

Experimental model and subject details

Mice

Animals were cared for in accordance with NIH guidelines. All experimental methods were approved by Washington University Institutional Committee on the Use and Care of Animals under the protocol number 20180303 and 210083. Animals were housed in a 12:12 light:dark cycle. We employed trio breeding and only used unaffected animals as parents. Pups were weaned from P21 to P28 and genotyped by phenol/chloroform DNA extraction of tail or toe (< P7 pups) biopsy followed by PCR. Weaned mice were separated by sex and relocated for breeding from 6 to 24 weeks' old. Animals were allocated to experimental versus control groups based on genotype. Sex matched-littermate controls were used in all experiments, and the age at which animals were used is reported in figure legends.

The *Usp7* floxed mice were first obtained on C57BL/6J and 129Sv mixed background[15] and then backcrossed to C57BL/6J background with speed congenics (last generation SNP typing > 99% determined by Dartmouse). The *NEX-cre* mice on C57BL/6J background were obtained

from Dr. Hui-Chen Lu's lab in Indiana University Bloomington, under the permission of Dr.
Sandra Goebbels[19]. Only the videotaped *Usp7* cKO mice were generated from the *Usp7* floxed
mice on C57BL/6J and 129Sv mixed background while the mice in all experiments were
generated from C57BL/6J mice and maintained on C57BL/6J background. *Usp7* cKO mice were
separated and single-caged once they were found to bite their cage mates.

The *Trp53* mutant mice on C57BL/6J background were purchased from the Jackson Laboratory
(Cat#002101). Consistent with the description from the Jackson Laboratory, female *Trp53* [-/-]
mice were born under the Mendelian ratio and exhibited young adult lethality. Therefore, only
male mice with *Trp53* [-/-] were used for breeding and *in vivo* experiments. For cortical neuron
culture, *Usp7* [fl/fl]; *Trp53* [-/-] male mice crossed with either *Usp7* [fl/fl]; *Trp53* [-/-] or *Usp7* [fl/fl]; *Trp53* [+/-]
female mice to produce embryos with *Usp7* [fl/fl]; *Trp53* [-/-] or *Usp7* [fl/fl]; *Trp53* [+/-]. As p53 deficiency
increases the propensity to grow tumor, mice and embryos with observed tumor bumps were
checked and excluded from all experiments.

The wild-type CD1 mice with timed E16 pregnancy for the *in vitro* apoptosis analysis, and the
Long-Evans rat with timed E17 pregnancy for the IP-MS analysis were both purchased from
Charles River. No breeding were made with these animals.

HEK293T cell line
Human embryonic kidney 293T (HEK293T/HEK293LE; Clontech 632180) cells were cultured in
DMEM with 10% fetal bovine serum (Fisher #10082-147) and penicillin/streptomycin (Life
Technologies #15140122). For lentiviral packaging, DNA/PEI (Polysciences) complexes were
suspended in Opti-MEM medium (Gibco) and transfected into HEK293T cells with about 80%
confluency. The transfer plasmid was mixed with packaging plasmid psPAX2 and envelope
plasmid pCMV-VSVG. The medium was replaced on the next day and the new medium
containing lentivirus was collected after 3 days. Cell debris was filtered from the collected
medium and Lenti-X Concentrator (Clontech) was then used to precipitate the lentivirus in the
medium. After incubation at 4°C overnight to 3 days, virus-containing pellets were collected by
centrifuging at 1500g for 45 min at 4°C, resuspended in phosphate-buffered saline (PBS), and
stored at –80°C in aliquots.

Cortical neuron culture
For IP-MS and immunocytochemistry of USP7 TRAF domain, cortical neurons were cultured
from E17 Long-Evans rat embryos. For *in vitro* apoptosis analysis, cortical neurons were
cultured wild-type E16 CD1 mouse embryos. For dendrite morphology analysis and TUBE
assay, cortical neurons were cultured from E15 *Usp7* [fl/fl]; *Trp53* [+/-] embryos on C57BL/6J
background. For *in vitro* TMT proteomics, cortical neurons were cultured from E15 *Usp7* [fl/fl];
Trp53 [-/-] embryos on C57BL/6J background. For spine analysis with RNAi knockdown, cortical
neurons were cultured from E15 C57BL/6J wild-type embryos.

To prepare cortical neurons (previously described as Spiegel et al. 2014[49]), embryonic cortices
were dissected in 1X HBSS and digested with 2.5% Trypsin (Gibco) at 37°C. DNase (Roche)
was added to digest free-floating DNA from dead cells. Trypsin-digested tissue was further
triturated with pipette acid to fully dissociate cells. Dissociated neurons were seeded onto tissue
culture dishes pre-coated with poly-L-lysine (0.1mg/mL) at a density of 1.2×10^5 cells/cm^2.
Neurons were first cultured with neurobasal medium with 1% fetal bovine serum (2% for rat
neuron), 0.5mM glutamine, B27 Supplement (Gibco) and penicillin/streptomycin (Life
Technologies #15140122) at 37°C with 5% CO_2. On DIV4, DIV8 and DIV12, neurons were fed
with half fresh medium without serum. Every day after DIV14, one fifth of the old medium was

replaced with fresh medium without serum to maintain stable medium composition. Neuron
viability was confirmed at DIV11 with MTS assay.
711
712 **Method details**
713
General design of behavioral tests
All behavior tests were conducted in the Washington University Animal Behavior Core and are
approved by the Washington University Institutional Animal Care and Use Committee. Sex and
litter-matched mice with and without p53 heterozygosity were tested separately due to limit of
cohort size for each batch of phenotypical analysis. After the event that one genotype of mice
were all dead or severely wounded, all mice in the litter were removed from the analysis to
reassure all animals are litter-matched. All tests were performed during the light phase of the
light/dark cycle by a female experimenter blind to the genotypes. Mice were acclimated to the
test room for 30 min prior to testing.
723
Sensorimotor battery
Ledge, platform, pole, and inclined and inverted screen tests were performed to assess
sensorimotor function. Time in each task was manually recorded. The average for two trials was
used for analyses. Test duration was 60s, except for the pole test, which was extended to 120s.
For walking initiation, time for an animal to leave a 21x21cm square on a flat surface was
recorded. For ledge and platform tests, the time the animal was able to balance on an acrylic
ledge (0.75cm wide and 30cm high), and on a wooden platform (1.0cm thick, 3.0cm in diameter
and elevated 47cm) was recorded, respectively. The pole test was used to evaluate fine motor
coordination by quantifying time to turn 180° and climb down a vertical pole. The screen tests
assessed a combination of coordination and strength by quantifying time to climb up or hang
onto a mesh wire grid measuring 16 squares per 10cm, elevated 47cm and inclined (60° or 90°)
or inverted.
736
Catwalk
Natural gait parameters were measured using the CatWalk XT gait analysis system (Noldus,
Inc). Mice were habituated to the apparatus and testing room the day before testing by placing
each mouse on the glass walkway for 5 min to help facilitate movement during the test session.
Testing was conducted with the room lights off. The following day, mice were placed on the 8-
cm wide glass walkway and allowed to walk naturally until 3 recordings were collected meeting
compliance criteria of less than 60% speed variability. A camera below the glass surface
recorded the paw placement and intensity of each paw as the mouse moved across the
walkway. Run speed, swing speed, stride length, and mean intensity of paw placement were
analyzed to assess differences in gait.
747
Virtual Optomotor System (VOS)
Visual acuity was assessed by recording Spatial frequency thresholds, in cycles per degree
(c/d) using the mouse OptoMotry HD System (Cerebral Mechanics Inc.). This is a virtual-reality
system designed to quantify visuomotor behavior[50]. Mice were allowed to acclimate to
procedure room 30 minutes prior to testing. Each mouse was placed onto a platform in the
center of the VOS chamber, and the experimenter tracked the movement of the animal's head
during the course of the test. VOS chamber consists of four computer screens surrounding the
platform with a camera to capture the OMR from above. Drift speed was set at 12.5, contrast
100 degree. The OptoMotry software uses a staircase method to determine at which threshold
the optomotor reflex (OMR) is no longer detected by the experimenter. Experimenter remained
blind to the direction of virtual movement.
759

Spontaneous alternation Y-maze

Testing was conducted according to our previously published procedures[51]. Briefly, this involved placing a mouse in the center of a Y-maze that contained three arms that were 10.5 cm wide, 40 cm long and 20.5 cm deep where an arm was oriented at 120° with respect to each successive other arm. Mice were allowed to explore the maze for 8 min and entry into an arm was scored only when the hindlimbs had completely entered the arm. An alternation was defined as any three consecutive choices of three different arms without re-exploration of a previously visited arm. The number of alternations and arm entries along with the percentage of alternations, which was determined by dividing the total number of alternations by the total number of entries minus 2, then multiplying by 100 was analyzed as a measure of spontaneous alternation.

Acoustic startle response/auditory function

Acoustic startle responses were measured using Kinder Scientific Startle Reflex equipment and software (Kinder Scientific, Inc) Auditory function was assessed by measuring startle response to auditory stimuli pulses (40 ms broadband burst) presented at 80dB, 90dB, 100dB, 110dB, and 120dB above background noise (65dB). Each trial type was presented 10 times in a random order over a 20 min session, with inter-trial-intervals ranging from 5-15 sec between trials. Beginning at stimulus onset, 1 ms force readings were obtained to record an animal's startle amplitude. The average force response to each sound pressure level was analyzed as a measure of auditory function.

Conditioned fear

Two distinct chambers were used to train and test mice (26×18×18 cm high) (Med-Associates, St. Albans, VT) which were easily distinguished by different olfactory, visual, and tactile cues present in each chamber. On day 1, each mouse was placed into the conditioning chamber (peppermint odor) for 5 min and freezing behavior was quantified during a 2 min baseline period. Freezing (no movement except that associated with respiration) was quantified using FreezeFrame image analysis software (Actimetrics, Evanston, IL) which allows for simultaneous visualization of behavior while adjusting for a "freezing threshold" during 0.75 s intervals. After baseline measurements, a conditioned stimulus (CS) consisting of an 80 dB tone (white noise) was presented for 20 sec followed by an unconditioned stimulus (US) consisting of a 1 s, 1.0 mA continuous foot shock. This tone-shock (T/S) pairing was repeated each minute over the next 2 min, and freezing was quantified after each of the three tone-shock pairings. Twenty-four hours after training, each mouse was placed back into the original conditioning chamber to test for fear conditioning to the contextual cues in the chamber. This involved quantifying freezing over an 8 min period without the tone or shock being present. Twenty-four hours later, the mice were evaluated on the auditory cue component of the conditioned fear procedure, which included placing each mouse into the other chamber containing distinctly different cues (coconut odor). Freezing was quantified during a 2 min "altered context" baseline period as well as over a subsequent 8 min period during which the auditory cue (CS) was presented. Shock sensitivity was evaluated following completion of the conditioned fear test.

Biting behavior

To quantitate biting behavior, Usp7 cKO or Usp7 cKO; Tp53 +/- mice were weaned after 3 weeks old and co-housed with their littermates. Male and female mice were separated and housed in a regular 12:12 light:dark cycle. Tail wounds were checked every week until 12 weeks old. Once tail wounds were found, the aggressors were marked and separated to avoid more biting. Victim mice with severe tail wound were euthanized and photographed. To video record the biting behavior, P33 aggressor and littermate were placed onto an ice cream box lid for free exploration.

811
Calcium phosphate sparse transfection
Primary cortical neurons were transfected as previously described[52]. Briefly, transfection mix was prepared by mixing 50ng/μL plasmid DNA, 0.25mM $CaCl_2$ and Hepes Buffered Saline (Invitrogen) and the DNA-calcium phosphate precipitate was made by pipetting up and down 10 times. Neurons were washed 3 times and starved with fresh serum-free medium at 37°C for 30min, and then 40μL of the transfection mix was dropped into each well in a 24-well plate (80μL for 12-well plate) and incubate at 37°C for 20-30 min. To remove DNA-calcium phosphate precipitate and avoid toxicity, neurons were then washed with fresh serum-free medium 3 times again and put back conditioned medium.

Immunofluorescence and immunocytochemistry
For immunofluorescence, mice were anesthetized with isoflurane, perfused with 40□m/mL heparin in PBS and 4% paraformaldehyde (> P5) or decapitated (< P5) before brain dissection and immersion fixation of the whole brain in 4% paraformaldehyde overnight at 4°C. Fixed brains were then placed in 30% sucrose in PBS for cryopreservation. After sinking in the sucrose solution, the brains were embedded with mixture of O.C.T. Compound (Sakura Finetek) and 30% sucrose solution (3:7), flash-frozen using liquid nitrogen. With a Leica cryostat, the frozen brains were sectioned at 16μm thickness and placed onto positively charged slides. For spine analysis, the brains were sectioned at 100μm thickness and placed into FD section storage solution (FD NeuroTechnologies). For immunocytochemistry, cells were initially seeded onto HNO_3-treated coverslips in 12 or 24 well-plates, washed with warm PBS and fixed in 4% paraformaldehyde for 30min on ice. For staining, brain sections or fixed cells were washed with PBS, permeabilized with 0.4% Triton X-100 and blocked with 3% Bovine serum alumin (BSA) and 10% goat serum. The permeabilized and blocked sections or cells was incubated with primary antibody overnight at 4°C, followed by incubation with fluorescence-conjugated secondary antibodies (Invitrogen) for 2 hours at room temperature and Hoechst diluted in PBS at 1:2000. The stained sections or cells were mounted onto slides with Fluoromount-G (SouthernBiotech) or Prolong™ Glass Antifade Mountant (Invitrogen) for spine imaging. Antigen retrieval with citrate acid (0.1 M, pH 6.8 with 0.1% Tween-20) for 15min boiling prior to blocking was required for labeling with anti-Ki67 antibody.

In vivo apoptosis analysis
Cryosections of P0 mouse brains were immunolabeled with cleaved-caspase 3 (c-cas 3) antibody (Cell Signaling #9961, 1:4000) for apoptosis analysis. Image acquisition and quantitation of the c-cas 3 intensity were blind to mouse genotypes at all steps below. 6-8 consecutive coronal sections per mouse, 3-5 mice per genotype were imaged with Leica DMi8 scanning microscope with instant computational clearing (THUNDER). The imaged sections were selected based on matched hippocampal morphologies across all mice. Segmentation of the cortex to analyze was performed based on Hoechst nuclei counterstain. C-cas 3 integral intensity within the segmented region of each section was measured with ImageJ and normalized to area.

In vitro apoptosis analysis
Primary cortical neurons from E16 wild-type CD1 mouse embryos were sparsely transfected at DIV3 with U6 empty vector, or U6-driven plasmids of shRNAs targeting USP7 or irrelevant USP30 along with EGFP plasmids. The transfected neurons were fixed at DIV5, 6 and 7, immunolabeled with anti-c-cas 3 (Cell Signaling #9961) and anti-GFP antibody (Abcam #ab13970) and imaged with Zeiss Axio Imager Z2 with ApoTome 2. Image acquisition and quantitation were blind to groups at all steps. Apoptotic cells are counted as GFP+ cells with c-

861 cas3 signal in the cell body and the intensity of apoptosis was quantitated by the percentage of
862 apoptotic cells over all GFP+ cells. The analysis was repeated with three batches of neuron
863 culture and 2-3 wells in 12-well plates were used per group for each culture.
864
Western blot
Cells or cortical tissues were lysed in RIPA buffer (Cell Signaling #9806) or TNE buffer (50mM
Tris pH 7.4, 100mM NaCl, 0.1mM EDTA, 1% NP40) mixed with protease inhibitor cocktail
(Sigma #P8340). Total protein was quantified with Bradford reagent (Bio-Rad). Lysate samples
were separated by SDS-PAGE or LDS-NuPAGE and transferred to 0.2µm nitrocellulose
membrane (GE Healthcare) via tank immersion. Membranes were blocked in 5% non-fat Milk in
Tris-buffered saline with 0.05% Tween-20 (TBST) at 4°C overnight, incubated with primary
antibodies at 4°C overnight, and subsequently with infrared fluorescent secondary antibodies
(LI-COR) for 2h at room temperature. Washes with TBST three times were applied to
membranes after every antibody incubation step. LI-COR Odyssey CLx imager and LI-COR
Image Studio software were used to image the blot at 600 dpi and to determine fluorescence
densitometry to quantitate relative protein abundance. The relative abundance of each protein
was normalized to loading control proteins (14-3-3 or Lamin A/C) in the same lane. Exposure
time was adjusted to avoid pixel saturation.
879
Immunoprecipitation
For immunoprecipitation of GFP-TRAF domain, we generated total cell lysates from HEK293T
cells with TNE lysis buffer as described above. 5µM proteasome inhibitor MG132 was applied to
enhance the expression of Myc3-Mdm2 in HEK293T cells. 1.5µg rabbit anti-GFP antibody
(homemade) was added to 200µL lysate at 2µg/µL total protein concentration determined by
Bradford reagent followed by rotation at 4°C overnight. After antibody binding, 18µL of PBS-
washed protein G agarose beads (GE Healthcare) were added per immunoprecipitation. Bead
binding proceeded for 1 hour at 4°C before 3 times wash with TNE lysis buffer and 3 times wash
with cold PBS. 30µL LDS sample buffer was added to beads per immunoprecipitation and
heated at 70°C for elution. The precipitated proteins were proceeded to western blot analysis.
890
For immunoprecipitation of endogenous Ppil4, we generated total cell lysates from HEK293T
cells with TNE lysis buffer. 4µg Rabbit anti-Ppil4 antibody (Bethyl) or Rabbit normal IgG control
was added to 200µL lysate at 4µg/µL total protein concentration determined by Bradford
reagent followed by rotation at 4°C overnight. After antibody binding, 16µL of PBS-washed
protein A agarose beads (GE Healthcare) were added per immunoprecipitation. Bead binding
proceeded for 1 hour at 4°C before 3 times wash with TNE lysis buffer and 3 times wash with
cold PBS. 30µL LDS sample buffer was added to beads per immunoprecipitation and heated at
70°C for elution. The precipitated proteins were proceeded to western blot analysis.
899
For immunoprecipitation of FLAG-TRAF domain, we generated total neuron lysates from
primary rat neurons with the TNE lysis buffer. pHAGE-hSyn-TRAF WT/RDW were transduced
into neurons at DIV7 and protein lysates were harvested at DIV10. 20µL PBS-washed M2 anti-
FLAG agarose beads (Sigma) were added to 2.5mL lysate at 2µg/µL total protein concentration
for 3 hours incubation at 4°C. The beads were washed 3 times with TNE lysis buffer, 3 times
with cold PBS, followed by 3 times wash with 200ng/µL 3x FLAG peptides (APExBio) for elution.
1.5% eluted samples were used for LDS-NuPAGE followed by silver stain (Invitrogen) while the
rest was proceeded to tricarboxylic acid (TCA) precipitation and LC-MS/MS analysis. 3
biological replicates of neuron plates with TRAF-WT and RDW mutant were used.
909
LC-MS/MS identification of interactors of the USP7 TRAF domain

We resuspended the sample in 20 µL of 8 M urea, 100 mM EPPS pH 8.5. We added 5mM
TCEP and incubated the mixture for 15 min at room temperature. We then added 10 mM of
iodoacetamide for 15min at room temperature in the dark. We added 15 mM DTT to consume
any unreacted iodoacetamide. We added 180µl of 100 mM EPPS pH 8.5. to reduce the urea
concentration to <1 M, 1 µg of trypsin, and incubated at 37□C for 6 hrs. The solution was
acidified with 2% formic acid and the digested peptides were desalted via StageTip, dried via
vacuum centrifugation, and reconstituted in 5% acetonitrile, 5% formic acid for LC-MS/MS
processing.

Mass spectrometry data were collected using a Q Exactive mass spectrometer (Thermo Fisher
Scientific, San Jose, CA) coupled with a Famos Autosampler (LC Packings) and an Accela600
liquid chromatography (LC) pump (Thermo Fisher Scientific). Peptides were separated on a 100
µm inner diameter microcapillary column packed with ~20 cm of Accucore C18 resin (2.6 µm,
150 Å, Thermo Fisher Scientific). For each analysis, we loaded ~1 µg onto the column. Peptides
were separated using a 75 min method from 2 to 25% acetonitrile in 0.125% formic acid with a
flow rate of ~300 nL/min. The scan sequence began with an Orbitrap MS^1 spectrum with the
following parameters: resolution 70,000, scan range 300–1500 Th, automatic gain control (AGC)
target 1×10^5, maximum injection time 250 ms, and centroid spectrum data type. We selected
the top twenty precursors for MS^2 analysis which consisted of HCD high-energy collision
dissociation with the following parameters: resolution 17,500, AGC 1×10^5, maximum injection
time 90 ms, isolation window 1.6 Th, normalized collision energy (NCE) 27%, and centroid
spectrum data type. The underfill ratio was set at 9%, which corresponds to a 1.5×10^5 intensity
threshold. In addition, unassigned and singly charged species were excluded from MS^2 analysis
and dynamic exclusion was set to automatic.

Mass spectra were processed using a SEQUEST-based in-house software pipeline. MS spectra
were converted to mzXML using a modified version of ReAdW.exe. Database searching
included all entries from the *rat* uniprot database which was concatenated with a reverse
database composed of all protein sequences in reversed order. Searches were performed using
a 50 ppm precursor ion tolerance. Product ion tolerance was set to 0.03 Th.
Carbamidomethylation of cysteine residues (+57.0215Da) were set as static modifications, while
oxidation of methionine residues (+15.9949 Da) was set as a variable modification. Peptide
spectral matches (PSMs) were altered to a 1% FDR[53,54]. PSM filtering was performed using a
linear discriminant analysis, as described previously[55], while considering the following
parameters: XCorr, ΔCn, missed cleavages, peptide length, charge state, and precursor mass
accuracy. Peptide-spectral matches were identified, quantified, and collapsed to a 1% FDR and
then further collapsed to a final protein-level FDR of 1%. Furthermore, protein assembly was
guided by principles of parsimony to produce the smallest set of proteins necessary to account
for all observed peptides.

TMT mass spectrometry
Sample preparation
Preparation of samples prior to mass spectrometry was performed as previously described[56]. In
brief, 5 murine cortices were harvested and flash-frozen at P17 and then lysed by bead-beating
in lysis buffer (200mM EPPS, pH 8.5 and 8 M urea with protease and phosphatase inhibitors) at
maximum speed for four cycles of 30 s each, with 1 min pauses on ice between cycles. For 16-
plex experiment, primary mouse cortical neurons were dissociated from E15 USP7 fl/fl; Tp53 -/-
embryos on C57BL/6J background. USP7 depletion was induced by lentiviral transduction of
wild-type Cre recombinase or catalytically dead Cre as control at DIV3. Four biological
replicates of neurons transduced with wild-type Cre and catalytically dead Cre were harvested

at DIV6 and DIV8, respectively. For cultured neurons the lysate was passed 10 times through a
21-gauge (1.25 inches long) needle. An equal quantity of protein (100 µg) for each condition (10
cortex slices, 16 samples of transduced neurons) was reduced with 5 mM tris (2-carboxyethyl)
phosphine (room temperature, 15 min) before proteins were alkylated with 10 mM
iodoacetamide (room temperature, 30 min in the dark). Finally, excess iodoacetamide was
quenched with 10 mM dithiothreitol (room temperature, 15 min). Alkylated proteins were isolated
by methanol-chloroform precipitation prior to protease digestion. The samples were diluted in
lysis buffer to a volume of 100 µL before adding methanol (400 µL), chloroform (100 µL) and
water (300 µL) stepwise with vortexing after addition of each solvent. The samples were
centrifuged for 1 min at 14,000 g and the aqueous and organic phases were removed, leaving
only the protein disk, which was washed with methanol (500 µL) and centrifuged at 21,000 g for
2 min at room temperature. After removal of methanol supernatant, the samples were
resuspended in 200 mM EPPS (pH 8.5) containing Lys-C protease (Wako) at a 100:1 protein-to-
protease ratio then incubated overnight at room temperature while shaking. Trypsin was then
added at a 100:1 protein-to-protease ratio, and the reaction was incubated 6 hr at 37°C.
Acetonitrile (30 µL) was added to the digested peptides and vortexed before adding 200 µg
TMT reagent (TMT10 or TMTpro16) for a 2:1 TMT-to-peptide ratio. The samples were incubated
at room temperature for 1 h and briefly vortexed every 10 min. Once labeling efficiency was
confirmed, the reaction was quenched with hydroxylamine to a final concentration of 0.3% (v/v)
for 15 min. The TMT-labeled samples were pooled, vacuum-centrifuged long enough to
evaporate the acetonitrile, then desalted with a C18 solid-phase extraction (SPE) (Sep-Pak,
Waters) and vacuum centrifuged to dryness.
983
Basic reversed-phase fractionation
For basic reversed-phase fractionation, desalted peptides were resuspended in Buffer A (10
mM ammonium bicarbonate, 5% ACN, pH 8), then subjected to a linear gradient from 13% to 42%
of Buffer B (10 mM ammonium bicarbonate, 90% ACN, pH 8) on an Agilent 300Extend C18
column (3.5 µm particles, 4.6 mm ID and 250 mm in length). The peptide mixture was
fractionated into a total of 96 fractions which were consolidated into 24 super fractions, of which,
12 non-adjacent samples were analyzed by the mass spectrometer. These 12 fractions were
vacuum centrifuged until dry, then desalted via StageTip for LC-MS analysis.
992
LC-MS analysis
The LC-MS analysis and protein quantification was slightly different for proteins extracted from
the mouse cortex slices versus those from the cultured neurons, reflecting changes in
instrumentation over the course of these experiments. Eluted peptides were introduced to an
Orbitrap Fusion with FAIMS pro (mouse cortex slices) or Orbitrap Fusion Lumos (cultured
neurons) mass spectrometer coupled to a Proxeon EASY-nLC 1200 LC pump (both from
ThermoFisher Scientific). Peptides were separated over a 90 min (cortex) or 120 min (neurons)
gradient on a 35-40 cm long 100 µm inner diameter column packed with Accucore (2.6 µm,
150Å) resin (ThermoFisher Scientific). For cortex slices the FAIMS voltages were set at -40, -60,
and -80V using a top speed method with 1s per CV. Full MS spectra were acquired in the
Orbitrap at a resolution of 120,000 and the 10 most intense MS1 ions were selected for MS2
analysis in the ion trap. The isolation width was set to 0.7 Da and isolated precursors were
fragmented by CID at a normalized collision energy (NCE) of 35%, and a max injection time of
35 ms (cortex) or 50 ms (neurons). Following acquisition of each MS2 spectrum, for the murine
cortex experiment, a synchronous precursor selection (SPS) MS3 scan was collected on the top
10 most intense ions in the MS2 spectrum. For the experiment from cultured neurons, a real-
time search (RTS-MS3) strategy was used, where only those SPS ions (up to 10) matching a
theoretical peptide spectrum were isolated for MS3 analysis. These SPS-MS3 precursors were
fragmented by higher energy collision-induced dissociation (HCD) at an NCE of 65% (TMT10)

or 50% (TMTpro) and analyzed using the Orbitrap at 50,000 resolution, with a max injection
time of 100 ms (cortex) or 200 ms (neurons).
1014
Database searching
Mass spectra were processed using SEQUEST (cortex) or Comet (neurons). Database
searching included all entries from the murine Uniprot database downloaded 07/2014 (cortex) or
07/2020 (neurons). Searches were performed using a 50-ppm precursor ion tolerance for total
protein-level analysis. The product ion tolerance was set to 0.9 Da (SEQUEST) or 1.005
(Comet). TMT tags (+229.163 Da) or TMTpro tags (+304.2071) on lysine residues and peptide
N termini and carbamidomethylation of cysteine residues (+57.021 Da) were set as static
modifications, while oxidation of methionine residues (+15.995 Da) was set as a variable
modification. Peptide-spectrum matches (PSMs) were identified, quantified, and filtered to a 1%
peptide false discovery rate (FDR) and then collapsed further to a final protein-level FDR of 1%.
The total summed signal-to-noise (S:N) measurements of the peptides in each condition were
used to calculate column normalization factors to account for small differences in sample
abundance. Then, for each protein, signal-to-noise (S:N) measurements of the peptides were
summed and then normalized to 100 across the all samples to yield a relative abundance
measurement.
1030
RNA extraction and RT-qPCR
Total RNA from mouse cortical tissues and primary cortical neurons were isolated using
QIAshredder and RNeasy Plus Mini Kit (QIAGEN) according to the manufacturer's instructions.
Cortical tissues were homogenized with mortar and pestle on dry ice during lysis. The extracted
RNA was reversely transcribed into cDNA by a High-Capacity cDNA Reverse Transcription Kit
with RNase Inhibitor (Applied Biosystems). Mixed cDNA, designed primers and Power SYBR
Green PCR Master Mix (Applied Biosystems) were loaded to qPCR on a CFX Connect Real-
Time PCR Detection System (Bio-Rad) with the program below: hot start, 95°C for 10 min;
amplification 40 cycles including denaturation (95°C for 15s), annealing (57°C for 30s), and
extension (72°C for 30s). A melting curve analysis (65°C for 5s and then 5s each at 0.5°C
increments between 65° and 95°C) was conducted to ensure single amplicon. Glyceraldehyde-
3-phosphate dehydrogenase (GAPDH) was used as the internal reference gene and the
conversion from cycle number to relative RNA level is performed with Excel 2019 (Microsoft).
1044
Streptavidin pull-down
For assessment of the interaction between USP7 and endogenously expressed proteins, a
Streptavidin-binding peptide-FLAG (SFB)-tagged USP7 construct[57] (Addgene plasmid #99393)
was transfected into HEK293T cells via PEI transfection. The lysate containing SFB-USP7 was
generated with TNE lysis buffer as described above and incubated with PBS-washed
Dynabeads[TM] MyOne[TM] Streptavidin T1 beads (Invitrogen #65601) for 1 hour at 4 °C with
rotation. Next, the beads were washed 3 times with TNE lysis buffer and 4 times with cold PBS,
followed by LDS sample buffer elution. The eluted proteins were analyzed by western blot along
with 2% input lysates in the same blot.
1054
Tandem ubiquitin binding entities (TUBE) assay
The TUBE assay was performed according to Tsao et al 2021 with modification[58]. Briefly,
primary cortical neurons from E15 mouse embryos were transduced with Cre wild-type (WT) or
catalytically dead (CD) lentivirus at DIV3 and treated with 10μM MG132 at DIV5. Neuron lysates
were generated with TNE lysis buffer 16 hours after MG132 treatment. Cold TBST-washed pan-
selective TUBE 2 beads (LifeSensors #UM-0402M) were incubated with neuron lysates
overnight at 4°C with rotation. Next, the beads were washed 5 times with cold TBST and 2 times

with cold PBS, followed by LDS sample buffer elution. The eluted proteins were analyzed by
western blot along with 2% input lysates in the same blot.
1064
***In vitro* deubiquitination assay**
For *in vitro* ubiquitination assay, HEK293T cells were transfected with pcDNA3.1-FLAG-Ppil4
plasmid. After the treatment of 10μM proteasome inhibitor MG132 for 16 hours, the cells were
harvested, lysed with 1% NP40 TNE lysis buffer, and the FLAG-Ppil4 was purified by
immunoprecipitation with M2 anti-FLAG agarose beads (Sigma) as described above. 20μL PBS-
washed M2 beads were added to 2.5mL lysate at 2μg/μL total protein concentration for 3 hours
incubation at 4°C. The beads were washed 3 times with TNE lysis buffer, 2 times with high-salt
PBS buffer (0.5M NaCl) to reduce non-specific binding, and then 3 times with regular PBS.
Then we resuspended the washed beads with deubiquitination buffer (50mM Tris-HCl, 150mM
NaCl, 5mM $MgCl_2$ and 10mM DTT, final pH 7.3) and distributed 9μL beads for each
deubiquitination reaction. Either 1μL 500nM recombinant His-USP7 (R&D system) or 1μL
deubiquitination buffer as vehicle control was added to each reaction and incubated for 1 hour
at 37 °C. LDS sample buffer was then added to stop the reaction and ubiquitinated Ppil4 was
analyzed with western blot. Densitometric analysis was performed with FLAG-Ppil4 purified from
3 biological replicates of HEK293T lysates. Ubiquitinated Ppil4 was quantified as the fraction of
Ppil4 integral fluorescent intensity above the molecular weight of the lowest full-length Ppil4
(~65kD).
1082
Diffusion-weighted *ex vivo* MRI and volumetric measurements
The skull of the fixed brain was removed before transferring to a 3-mL syringe containing fresh
PBS. Diffusion-weighted MRI experiment was performed on 9.4-T Bruker scanner with a
CryoProbe™. Fifteen sagittal slices, centered at the midline (slice 8), were acquired to cover the
entire brain. Twenty-five-direction diffusion-weighted measurements were performed using a
spin-echo diffusion-weighting sequence with the following parameters: repetition time (TR) = 1.5
s; echo time (TE) = 32 ms; field of view (FOV) = 20 × 20 mm2; matrix size = 192 × 192 (zero-
filled to 384 × 384), slice thickness = 0.6 mm, max; b value = 3,000 s/mm2, Δ = 20 ms; δ = 5 ms;
partial Fourier factor = 1.33; total scan time = 93.6 min. Diffusion-weighted data were converted
to nifiti format followed by analysis using the lab-developed DBSI software package to perform
DBSI multi-tensor analysis[59]. DBSI-derived fiber fraction was used to reflect apparent
axonal[60]/dendritic[61] density. DBSI-derived restricted and non-restricted fractions were used to
estimate the extent of cellularity and vasogenic edema[61]. For volumetric measurement,
segmentation was performed manually employing ITK-SNAP (version 3.8, www.itksnap.org)[62].
The Allen Mouse Brain Atlas was used as a reference to identify cortex and hippocampus. The
three parasagittal slices at each side of midline (slices 4 -6 and 10 -12) were used to segment
cortex and hippocampus.
1100
Adeno-associated virus (AAV) injection
The AAV transfer plasmid pAAV-hSyn1-EGFP-P2A-EGFPf-WPRE-HGHpA was a gift from Dr.
Guoping Feng (Addgene # 74513). The AAV PHP.eB was packaged at Hope Center Viral
Vector Core of Washington University School of Medicine and diluted to 1.5 x 10^7 vg/μL with 5%
D-sorbitol in PBS. For intraventricular injection, P0/1 pups were anesthetized with hypothermia.
1uL of the diluted AAV was injected into each hemisphere (0.8-1mm away from the middle point
between Lambda and Bregma, 2mm deep) of the pups. The injected mice were placed onto
heat pad for body temperature recovery before returned to dam in the cage. We then perfused
the injected mice at P18 and their brains were proceeded to cryosectioning and
immunofluorescence of GFP for spine imaging and analysis.
1111
Airyscan imaging and spine analysis

We used Zeiss 880 LSM2 confocal microscope with Airyscan hexagonal detector array and oil
immersion to image z-stacks of dendritic spines at near-super resolution. For *in vivo* spine
analysis, coronal sections at 100um thickness were float-stained for GFP. Based on Allen
mouse reference atlas, motor cortices (M1 and M2) were imaged and 3-8 pyramidal neurons
per section, 4 consecutive sections per animal, 3 animals per genotype were analyzed. The
analysis was restricted to secondary apical dendritic segment and secondary or tertiary basal
dendritic segment, starting from less than 40um distance to soma and for over 30um length.
Representative images of dendritic spines of four shapes (mushroom, thin, stubby and filopodial)
were pre-selected before quantification and used as references to compare with all other
dendritic spines to call their shape in a blinded manner. For *in vitro* spine analysis, primary
cortical neurons were seeded onto #1.5 coverslips. The neurons were transduced with lentivirus
of MISSION TRC1-shRNA (Sigma) knocking down candidate substrates at DIV4, sparsely
transfected with EGFP-expressing plasmid at DIV10, fixed at DIV18, and proceeded to
immunocytochemistry of GFP and PSD95. Dendritic spines of 4-12 neurons were imaged per
well, 3 wells per group, from 2 batches of neuron culture. The analysis was restricted to primary,
2^{nd} and 3^{rd} dendrites starting from less than 40µm distance to soma and for over 30µm length.
The spines and PSD95 puncta were counted with Zeiss ZEN software in three dimensional view
on a blinded basis. We used Imaris (Oxford Instruments) filament tracing to measure dendrite
length in three dimensional view and divide counted number of spines and PSD95 puncta to get
spine and PSD95 puncta density, respectively.

Dendrite tracing and morphological analysis
To assess dendrite morphology, primary cortical neurons from *Usp7* [fl/fl]; *Trp53* [+/-] embryos were
seeded onto #1 coverslips and transduced with Cre lentivirus at DIV3. The neurons was
sparsely transfected with EGFP-expressing plasmid at DIV9, fixed at DIV11 for
immunocytochemistry of GFP. The z-stacks of GFP signal were imaged with Zeiss 880 LSM2
confocal microscope and regular 20x objective. Based on the GFP signal, we used Imaris
(Oxford Instruments) to model soma shape and trace dendrite morphology with filament-
Autopath algorithm. Physically crossing dendrites were manually separated. Statistics of soma
volumes, dendrite lengths and Sholl intersections were all determined and exported with Imaris
filament. Complete dendrites of 9-15 neurons per well, 3 wells per group, from 2 batches of
neuron culture were imaged and analyzed. All image acquisition and analysis were performed
on a blinded basis.

MTS assay
Viability of the primary cortical neurons was assessed according to our previous publication[63].
Briefly, MTS reagent (Promega) was added to neurons 1:5 ratio to medium volume at DIV11
according to manufacturer's instructions. The treated neurons were incubated at 37 °C with 5%
CO_2 for 2 hours before measurement of absorbance at 490nm. The measured MTS absorbance
was subtracted with background signal in empty wells and normalized to control group (Cre
dead or untreated neurons).

Quantification and statistical analysis

Gene set enrichment analysis (GSEA)
GSEA software 4.3.2[64,65] was used to identify the significantly differential enrichment of
annotated gene sets between proteins in cortices of USP7 WT and Usp7 cKO mice. Gene sets
were from M5 (gene ontology) of Molecular signatures database. NES values were calculated
and gene sets with p-value < 0.05 were loaded into Cytoscape 3.9.1 for network construction.
The enrichment map was constructed with yfiles Organic Layout, q-value < 0.01 as node cutoff,
and similarity < 0.5 as edge cutoff. All gene sets shown in the enrichment map were enriched in

USP7 WT mice and down-regulated in Usp7 cKO mice, as indicated by the green color (color
scheme: PiYG-3).
1166
Gene ontology analysis
GOrilla[22] (http://cbl-gorilla.cs.technion.ac.il) was used to identify significantly enriched gene
ontology terms in ranked gene list. For *in vivo* TMT protein abundance, n=131 proteins down-
regulated in Usp7 cKO mice (Fold Change < 0.75) were ranked by Fold Change and loaded to
GOrilla with a background of total proteins detected in all mice. 3 uncharacterized proteins
without gene name were not detected by GOrilla. No GO term was significantly enriched in
Usp7 cKO up-regulated proteins.
1174
SynGO analysis
1176 For SynGO analysis, unranked down-regulated proteins in *Usp7* cKO; *Trp53*[+/-] cortex in TMT-
1177 mass spectrometry experiment were identified with cutoff of Fold Change < 0.75 and 2-tail
1178 unpaired t-test p-value < 0.01. All proteins detected in all mice were used as background. The
1179 SynGO enrichment analysis was performed with the SynGO (v1.2) online tool for both cellular
1180 components and biological processes (https://syngoportal.org)[66].
1181
STRING network analysis
The String network was built with StringApp 2.0.1[67] in Cytoscape 3.9.1. To ensure specificity,
we calculated the enrichment percentage of the spectral counts in USP7 TRAF WT group, and
only interactors with over 95% enrichment were proceeded for the following analysis. Top 50
interactors based on spectral counts normalized to protein length were converted into
homologous mouse proteins and loaded into StringApp to create the association map. Two
uncharacterized rat proteins were not included. Full STRING network with edges of confidence
over 0.4 was built. Enriched molecular components (Gene ontology) were assigned to subunits
of the complex in the association map. FDR of each enriched molecular components was
calculated using the Benjamini-Hochberg procedure.
1192
Disease gene enrichment analysis
Interactors of USP7 TRAF domain used in the disease gene enrichment analysis are the
detected proteins in all 3 biological replicates of IP-mass spectrometry experiment. The
interactors were filtered with 95% enrichment in USP7 TRAF WT group, as described in the
String network analysis. All proteins in primary neuron culture detected by TMT-mass
spectometry were used as background. Curated pathogenic gene sets for autism, intellectual
disability (ID), Epilepsy, ADHD, schizophrenia, bipolar disorder and cerebral palsy were from
Geisinger developmental brain disorder gene database (https://dbd.geisingeradmi.org/)[24]. High-
confidence FMRP targets were from FMRP enriched genes in HITS-CLIP[68]. USP7 TRAF
interactors, FMRP targets, and the background proteins were all converted into human
orthologs entrez ID with biomaRt R package before the analysis. p-values of the disease gene
set enrichment were calculated with one-sided hypergeometric test, and then adjusted by
Benjamini-Hochberg procedure.
1206
Principal component analysis (PCA)
1208 PCA of all 16 samples of the *in vitro* TMT experiment was performed with function prcom() of R.
1209 All 8304 detected proteins were loaded into the analysis and principal component 1 and 2 were
1210 mapped with variance percentage.
1211
Selection of candidate substrates
1213 To filter out proteins with little to no expression in glutamatergic neurons, we used Allen Institute
1214 Smart-Seq database[69] to calculate the total CPM of the corresponding gene in the clade of

glutamatergic neuron clusters. Proteins with total CPM level in glutamatergic neurons below that of the astrocyte marker aldolase C were excluded from the following analyses. To select candidate substrates of USP7, we first calculated the Euclidean distance of each protein to USP7 over all 16 samples and sorted out the 25 proteins closest to USP7 as majority of the candidate substrates. Because the sorting based only on Euclidean distance overlooks the proteins with differential abundances between the two time points (DIV6 and DIV8), a 2x2 two-way ANOVA was also applied to all proteins to calculate the p-values of Cre and Cre-time interaction effect on the abundance of each protein. Proteins down-regulated upon wild-type Cre with p-values $< 10^{-7}$ were added to the candidate substrate list.

Weighted gene correlation network analysis (WGCNA)

The R package WGCNA[70] was used to construct a co-expression network for all the proteins with significant Cre effect (n=2307) or Cre-time interaction effect (n=320) (p<0.05 by two-way ANOVA). All 16 TMT samples were used to calculate Pearson's correlation matrices representing co-expression similarity of proteins, and different Cre type (Cre WT and Cre dead) and time points (DIV6 and DIV8) were loaded as traits. The weighted adjacency matrix was created using Pearson correlation coefficient test of all differentially expressed proteins and transformed into a topological overlap measure (TOM) matrix and average linkage hierarchical clustering of the TOM matrix was used to construct a clustering dendrogram of the genes with thresholding power 12. 10 Modules in different colors were generated with minimal gene module size of 30 proteins and minimal height to merge modules of 0.15. GO term enrichment analyses for the modules were performed by DAVID 2021 Functional Annotation Tool[71,72] with a background of all module-assigned proteins. For module-trait relationship, Cre WT, Cre dead, DIV6 and DIV8 were assigned as 1, 0, 6 and 8 respectively. The correlation coefficient and correlation p-value of the first principal component of the module gene expression profile (eigengene) and traits were calculated to identify trait-correlated modules.

Statistics

All statistical analyses were performed using IBM SPSS Statistic software (v.26), GraphPad Prism (v.9) and R. For experiments to compare only two groups, student t-test was used. Analysis of variance (ANOVA), (one-way, two-way, factorial, repeated measures) were used analyze data with multiple groups. With a statistically significant interaction between main factors, simple main effects were calculated to provide clarification of statistically significant between-group and within-group differences. Where appropriate, the Huynh-Feldt adjustment was used to protect against violations of sphericity. For multiple pairwise comparisons and post-hoc tests, Tukey correction was applied to compare every two means, and Bonferroni correction was applied to compare selected means when independence cannot be assumed. For three groups with non-normal distributions (sensorimotor battery), the nonparametric Kruskal-Wallis test was used followed by post hoc test with the Dunn correction to compare every mean to a control mean. All data in bar graphs were presented as mean ± SEM and probability value for all analyses was *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001, unless otherwise stated.

References

1.   Kishino, T., Lalande, M., and Wagstaff, J. (1997). UBE3A/E6-AP mutations cause Angelman syndrome. Nat. Genet. *15*, 70–73. 10.1038/ng0197-70.

2.   Margolin, D.H., Kousi, M., Chan, Y.-M., Lim, E.T., Schmahmann, J.D., Hadjivassiliou, M., Hall, J.E., Adam, I., Dwyer, A., Plummer, L., et al. (2013). Ataxia, dementia, and hypogonadotropism caused by disordered ubiquitination. N. Engl. J. Med. *368*, 1992–2003. 10.1056/NEJMoa1215993.

3.   Alqwaifly, M., and Bohlega, S. (2016). Ataxia and Hypogonadotropic Hypogonadism with Intrafamilial Variability Caused by RNF216 Mutation. Neurol. Int. *8*, 6444. 10.4081/ni.2016.6444.

4.   Ferguson, C.J., Urso, O., Bodrug, T., Gassaway, B.M., Watson, E.R., Prabu, J.R., Lara-Gonzalez, P., Martinez-Chacin, R.C., Wu, D.Y., Brigatti, K.W., et al. (2022). APC7 mediates ubiquitin signaling in constitutive heterochromatin in the developing mammalian brain. Mol. Cell *82*, 90-105.e13. 10.1016/j.molcel.2021.11.031.

5.   Yamada, T., Yang, Y., and Bonni, A. (2013). Spatial organization of ubiquitin ligase pathways orchestrates neuronal connectivity. Trends Neurosci. *36*, 218–226. 10.1016/j.tins.2012.12.004.

6.   Anckar, J., and Bonni, A. (2015). Regulation of Neuronal Morphogenesis and Positioning by Ubiquitin-Specific Proteases in the Cerebellum. PLoS One *10*, e0117076. 10.1371/journal.pone.0117076.

7.   Jolly, L.A., Kumar, R., Penzes, P., Piper, M., and Gecz, J. (2022). The DUB Club: Deubiquitinating Enzymes and Neurodevelopmental Disorders. Biol. Psychiatry *92*, 614–625. 10.1016/j.biopsych.2022.03.022.

8.   Hao, Y.-H., Fountain, M., Fon?Tacer, K., Xia, F., Bi, W., Kang, S.-H., Patel, A., Rosenfeld, J., Le?Caignec, C., Isidor, B., et al. (2015). USP7 Acts as a Molecular Rheostat to Promote WASH-Dependent Endosomal Protein Recycling and Is Mutated in a Human Neurodevelopmental Disorder. Mol. Cell *59*, 956–969. 10.1016/j.molcel.2015.07.033.

9.   Fountain, M.D., Oleson, D.S., Rech, M.E., Segebrecht, L., Hunter, J. V., McCarthy, J.M., Lupo, P.J., Holtgrewe, M., Moran, R., Rosenfeld, J.A., et al. (2019). Pathogenic variants in USP7 cause a neurodevelopmental disorder with speech delays, altered behavior, and neurologic anomalies. Genet. Med. *21*, 1797–1807. 10.1038/s41436-019-0433-1.

10. Cummins, J.M., Rago, C., Kohli, M., Kinzler, K.W., Lengauer, C., and Vogelstein, B. (2004). Tumour suppression: disruption of HAUSP gene stabilizes p53. Nature *428*, 6–7. 10.1038/nature02501.

11. Li, M., Brooks, C.L., Kon, N., and Gu, W. (2004). A Dynamic Role of HAUSP in the p53-Mdm2 Pathway. Mol. Cell *13*, 879–886. 10.1016/S1097-2765(04)00157-1.

12. Turnbull, A.P., Ioannidis, S., Krajewski, W.W., Pinto-Fernandez, A., Heride, C., Martin, A.C.L., Tonkin, L.M., Townsend, E.C., Buker, S.M., Lancia, D.R., et al. (2017). Molecular basis of USP7 inhibition by selective small-molecule inhibitors. Nature *550*, 481–486. 10.1038/nature24451.

13. Kategaya, L., Di Lello, P., Rougé, L., Pastor, R., Clark, K.R., Drummond, J., Kleinheinz, T., Lin, E., Upton, J.P., Prakash, S., et al. (2017). USP7 small-molecule inhibitors interfere with ubiquitin binding (Nature Publishing Group) 10.1038/nature24006.

14. Kon, N., Kobayashi, Y., Li, M., Brooks, C.L., Ludwig, T., and Gu, W. (2010). Inactivation of HAUSP in vivo modulates p53 function. Oncogene *29*, 1270–1279. 10.1038/onc.2009.427.

15. Kon, N., Zhong, J., Kobayashi, Y., Li, M., Szabolcs, M., Ludwig, T., Canoll, P.D., and Gu, W. (2011). Roles of HAUSP-mediated p53 regulation in central nervous system development. Cell Death Differ. *18*, 1366–1375. 10.1038/cdd.2011.12.

16. Huang, Z., Wu, Q., Guryanova, O.A., Cheng, L., Shou, W., Rich, J.N., and Bao, S. (2011). Deubiquitylase HAUSP stabilizes REST and promotes maintenance of neural progenitor

1310   cells. Nat. Cell Biol. *13*, 142–152. 10.1038/ncb2153.

1311   17.   Nicklas, S., Hillje, A.L., Okawa, S., Rudolph, I.M., Collmann, F.M., van Wuellen, T., del
1312   Sol, A., and Schwamborn, J.C. (2019). A complex of the ubiquitin ligase TRIM32 and the
1313   deubiquitinase USP7 balances the level of c-Myc ubiquitination and thereby determines
1314   neural stem cell fate specification. Cell Death Differ. *26*, 728–740. 10.1038/s41418-018-
1315   0144-1.

1316   18.   Kim, R.Q., and Sixma, T.K. (2017). Regulation of USP7: A High Incidence of E3
1317   Complexes. J. Mol. Biol. *429*, 3395–3408. 10.1016/J.JMB.2017.05.028.

1318   19.   Goebbels, S., Bormuth, I., Bode, U., Hermanson, O., Schwab, M.H., and Nave, K.-A.
1319   (2006). Genetic targeting of principal neurons in neocortex and hippocampus of NEX-Cre
1320   mice. genesis *44*, 611–621. https://doi.org/10.1002/dvg.20256.

1321   20.   Ogden, K.K., Ozkan, E.D., and Rumbaugh, G. (2016). Prioritizing the development of
1322   mouse models for childhood brain disorders. Neuropharmacology *100*, 2–16.
1323   10.1016/J.NEUROPHARM.2015.07.029.

1324   21.   Hashikawa, K., Hashikawa, Y., Lischinsky, J., and Lin, D. (2018). The Neural
1325   Mechanisms of Sexually Dimorphic Aggressive Behaviors. Trends Genet. *34*, 755–776.
1326   10.1016/j.tig.2018.07.001.

1327   22.   Eden, E., Navon, R., Steinfeld, I., Lipson, D., and Yakhini, Z. (2009). GOrilla: A tool for
1328   discovery and visualization of enriched GO terms in ranked gene lists. BMC
1329   Bioinformatics *10*, 1–7. 10.1186/1471-2105-10-48.

1330   23.   Sheng, Y., Saridakis, V., Sarkari, F., Duan, S., Wu, T., Arrowsmith, C.H., and Frappier, L.
1331   (2006). Molecular recognition of p53 and MDM2 by USP7/HAUSP. Nat. Struct. Mol. Biol.
1332   *13*, 285–291. 10.1038/nsmb1067.

1333   24.   Gonzalez-Mantilla, A.J., Moreno-De-Luca, A., Ledbetter, D.H., and Martin, C.L. (2016). A
1334   cross-disorder method to identify novel candidate genes for developmental brain
1335   disorders. JAMA Psychiatry *73*, 275–283. 10.1001/jamapsychiatry.2015.2692.

1336   25.   Chahrour, M.H., Yu, T.W., Lim, E.T., Ataman, B., Coulter, M.E., Hill, R.S., Stevens, C.R.,
1337   Schubert, C.R., Collaboration, A.A.S., Greenberg, M.E., et al. (2012). Whole-Exome
1338   Sequencing and Homozygosity Analysis Implicate Depolarization-Regulated Neuronal
1339   Genes in Autism. PLOS Genet. *8*, e1002635.

1340   26.   Zhang, J., Gambin, T., Yuan, B., Szafranski, P., Rosenfeld, J.A., Balwi, M. Al, Alswaid, A.,
1341   Al-Gazali, L., Shamsi, A.M. Al, Komara, M., et al. (2017). Haploinsufficiency of the E3
1342   ubiquitin-protein ligase gene TRIP12 causes intellectual disability with or without autism
1343   spectrum disorders, speech delay, and dysmorphic features. Hum. Genet. *136*, 377–386.
1344   10.1007/s00439-017-1763-1.

1345   27.   Hjerpe, R., Aillet, F., Lopitz-Otsoa, F., Lang, V., England, P., and Rodriguez, M.S. (2009).
1346   Efficient protection and isolation of ubiquitylated proteins using tandem ubiquitin-binding
1347   entities. EMBO Rep. *10*, 1250–1258. https://doi.org/10.1038/embor.2009.192.

1348   28.   Lelieveld, S.H., Reijnders, M.R.F., Pfundt, R., Yntema, H.G., Kamsteeg, E.-J., de Vries,
1349   P., de Vries, B.B.A., Willemsen, M.H., Kleefstra, T., Löhner, K., et al. (2016). Meta-
1350   analysis of 2,104 trios provides support for 10 new genes for intellectual disability. Nat.
1351   Neurosci. *19*, 1194–1196. 10.1038/nn.4352.

1352   29.   Vissers, L.E.L.M., Van Nimwegen, K.J.M., Schieving, J.H., Kamsteeg, E.J., Kleefstra, T.,
1353   Yntema, H.G., Pfundt, R., Van Der Wilt, G.J., Krabbenborg, L., Brunner, H.G., et al.
1354   (2017). A clinical utility study of exome sequencing versus conventional genetic testing in
1355   pediatric neurology. Genet. Med. *19*, 1055–1063. 10.1038/GIM.2017.1.

1356   30.   Iossifov, I., O'Roak, B.J., Sanders, S.J., Ronemus, M., Krumm, N., Levy, D., Stessman,
1357   H.A., Witherspoon, K.T., Vives, L., Patterson, K.E., et al. (2014). The contribution of de
1358   novo coding mutations to autism spectrum disorder. Nature *515*, 216–221.
1359   10.1038/nature13908.

1360   31.   De Rubeis, S., He, X., Goldberg, A.P., Poultney, C.S., Samocha, K., Ercument Cicek, A.,

Kou, Y., Liu, L., Fromer, M., Walker, S., et al. (2014). Synaptic, transcriptional and chromatin genes disrupted in autism. Nature *515*, 209–215. 10.1038/nature13772.

32. Tavana, O., and Gu, W. (2017). Modulation of the p53/MDM2 interplay by HAUSP inhibitors. J. Mol. Cell Biol. *9*, 45–52. 10.1093/jmcb/mjw049.

33. Woods, C.G. (2004). Human microcephaly. Curr. Opin. Neurobiol. *14*, 112–117. 10.1016/J.CONB.2004.01.003.

34. Harris, K.M., Jensen, F.E., and Tsao, B. (1992). Three-dimensional structure of dendritic spines and synapses in rat hippocampus (CA1) at postnatal day 15 and adult ages: implications for the maturation of synaptic physiology and long-term potentiation [published erratum appears in J Neurosci 1992 Aug;1. J. Neurosci. *12*, 2685 LP – 2705. 10.1523/JNEUROSCI.12-07-02685.1992.

35. Zoghbi, H.Y., and Bear, M.F. (2012). Synaptic dysfunction in neurodevelopmental disorders associated with autism and intellectual disabilities. Cold Spring Harb. Perspect. Biol. *4*. 10.1101/cshperspect.a009886.

36. Volk, L., Chiu, S.L., Sharma, K., and Huganir, R.L. (2015). Glutamate Synapses in Human Cognitive Disorders. Annu. Rev. Neurosci. *38*, 127–149. 10.1146/annurev-neuro-071714-033821.

37. Bagni, C., and Zukin, R.S. (2019). A Synaptic Perspective of Fragile X Syndrome and Autism Spectrum Disorders. Neuron *101*, 1070–1088. 10.1016/J.NEURON.2019.02.041.

38. Bélanger, C., Cardinal, T., Leduc, E., Viger, R.S., and Pilon, N. (2022). CHARGE syndrome-associated proteins FAM172A and CHD7 influence male sex determination and differentiation through transcriptional and alternative splicing mechanisms. FASEB J. *36*, e22176. https://doi.org/10.1096/fj.202100837RR.

39. Bélanger, C., Bérubé-Simard, F.-A., Leduc, E., Bernas, G., Campeau, P.M., Lalani, S.R., Martin, D.M., Bielas, S., Moccia, A., Srivastava, A., et al. (2018). Dysregulation of cotranscriptional alternative splicing underlies CHARGE syndrome. Proc. Natl. Acad. Sci. U. S. A. *115*, E620–E629. 10.1073/pnas.1715378115.

40. Schmitzová, J., Cretu, C., Dienemann, C., Urlaub, H., and Pena, V. (2023). Structural basis of catalytic activation in human splicing. Nature *617*. 10.1038/s41586-023-06049-w.

41. Bertram, K., El Ayoubi, L., Dybkov, O., Agafonov, D.E., Will, C.L., Hartmuth, K., Urlaub, H., Kastner, B., Stark, H., and Lührmann, R. (2020). Structural Insights into the Roles of Metazoan-Specific Splicing Factors in the Human Step 1 Spliceosome. Mol. Cell *80*, 127-139.e6. 10.1016/j.molcel.2020.09.012.

42. Vuong, C.K., Black, D.L., and Zheng, S. (2016). The neurogenetics of alternative splicing. Nat. Rev. Neurosci. *17*, 265–281. 10.1038/nrn.2016.27.

43. Furlanis, E., Traunmüller, L., Fucile, G., and Scheiffele, P. (2019). Landscape of ribosome-engaged transcript isoforms reveals extensive neuronal-cell-class-specific alternative splicing programs. Nat. Neurosci. *22*, 1709–1717. 10.1038/s41593-019-0465-5.

44. Ule, J., Ule, A., Spencer, J., Williams, A., Hu, J.-S., Cline, M., Wang, H., Clark, T., Fraser, C., Ruggiu, M., et al. (2005). Nova regulates brain-specific splicing to shape the synapse. Nat. Genet. *37*, 844–852. 10.1038/ng1610.

45. Nuber, U., Schwarz, S.E., and Scheffner, M. (1998). The ubiquitin-protein ligase E6-associated protein (E6-AP) serves as its own substrate. Eur. J. Biochem. *254*, 643–649. 10.1046/j.1432-1327.1998.2540643.x.

46. De Bie, P., and Ciechanover, A. (2011). Ubiquitination of E3 ligases: Self-regulation of the ubiquitin system via proteolytic and non-proteolytic mechanisms. Cell Death Differ. *18*, 1393–1402. 10.1038/cdd.2011.16.

47. Froyen, G., Corbett, M., Vandewalle, J., Jarvela, I., Lawrence, O., Meldrum, C., Bauters, M., Govaerts, K., Vandeleur, L., Van Esch, H., et al. (2008). Submicroscopic Duplications of the Hydroxysteroid Dehydrogenase HSD17B10 and the E3 Ubiquitin Ligase HUWE1

Are Associated with Mental Retardation. Am. J. Hum. Genet. *82*, 432–443.
10.1016/j.ajhg.2007.11.002.

48. Moortgat, S., Berland, S., Aukrust, I., Maystadt, I., Baker, L., Benoit, V., Caro-Llopis, A.,
Cooper, N.S., Debray, F.G., Faivre, L., et al. (2018). HUWE1 variants cause dominant X-
linked intellectual disability: A clinical study of 21 patients. Eur. J. Hum. Genet. *26*, 64–74.
10.1038/s41431-017-0038-6.

49. Spiegel, I., Mardinly, A.R., Gabel, H.W., Bazinet, J.E., Couch, C.H., Tzeng, C.P., Harmin,
D.A., and Greenberg, M.E. (2014). Npas4 Regulates Excitatory-Inhibitory Balance within
Neural Circuits through Cell-Type-Specific Gene Programs. Cell *157*, 1216–1229.
10.1016/J.CELL.2014.03.058.

50. Prusky, G.T., Alam, N.M., Beekman, S., and Douglas, R.M. (2004). Rapid quantification
of adult and developing mouse spatial vision using a virtual optomotor system. Investig.
Ophthalmol. Vis. Sci. *45*, 4611–4616. 10.1167/iovs.04-0541.

51. Wozniak, D.F., Diggs-Andrews, K.A., Conyers, S., Yuede, C.M., Dearborn, J.T., Brown,
J.A., Tokuda, K., Izumi, Y., Zorumski, C.F., and Gutmann, D.H. (2013). Motivational
Disturbances and Effects of L-dopa Administration in Neurofibromatosis-1 Model Mice.
PLoS One *8*, e66024.

52. Konishi, Y., Lehtinen, M., Donovan, N., and Bonni, A. (2002). Cdc2 Phosphorylation of
BAD Links the Cell Cycle to the Cell Death Machinery. Mol. Cell *9*, 1005–1016.
10.1016/S1097-2765(02)00524-5.

53. Elias, J.E., and Gygi, S.P. (2007). Target-decoy search strategy for increased confidence
in large-scale protein identifications by mass spectrometry. Nat. Methods *4*, 207–214.
10.1038/nmeth1019.

54. Elias, J.E., and Gygi, S.P. (2010). Target-decoy search strategy for mass spectrometry-
based proteomics. Methods Mol. Biol. *604*, 55–71. 10.1007/978-1-60761-444-9_5.

55. Huttlin, E.L., Jedrychowski, M.P., Elias, J.E., Goswami, T., Rad, R., Beausoleil, S.A.,
Villén, J., Haas, W., Sowa, M.E., and Gygi, S.P. (2010). A Tissue-Specific Atlas of Mouse
Protein Phosphorylation and Expression. Cell *143*, 1174–1189.
10.1016/J.CELL.2010.12.001.

56. Navarrete-Perea, J., Yu, Q., Gygi, S.P., and Paulo, J.A. (2018). Streamlined Tandem
Mass Tag (SL-TMT) Protocol: An Efficient Strategy for Quantitative (Phospho)proteome
Profiling Using Tandem Mass Tag-Synchronous Precursor Selection-MS3. J. Proteome
Res. *17*, 2226–2236. 10.1021/acs.jproteome.8b00217.

57. An, L., Jiang, Y., Ng, H.H.W., Man, E.P.S., Chen, J., Khoo, U.-S., Gong, Q., and Huen,
M.S.Y. (2017). Dual-utility NLS drives RNF169-dependent DNA damage responses. Proc.
Natl. Acad. Sci. *114*, E2872–E2881. 10.1073/pnas.1616602114.

58. Tsao, N., Brickner, J.R., Rodell, R., Ganguly, A., Wood, M., Oyeniran, C., Ahmad, T., Sun,
H., Bacolla, A., Zhang, L., et al. (2021). Aberrant RNA methylation triggers recruitment of
an alkylation repair complex. Mol. Cell *81*, 4228-4242.e8. 10.1016/j.molcel.2021.09.024.

59. Wang, Y., Wang, Q., Haldar, J.P., Yeh, F.-C., Xie, M., Sun, P., Tu, T.-W., Trinkaus, K.,
Klein, R.S., Cross, A.H., et al. (2011). Quantification of increased cellularity during
inflammatory demyelination. Brain *134*, 3590–3601. 10.1093/brain/awr307.

60. Lin, T.H., Zhan, J., Song, C., Wallendorf, M., Sun, P., Niu, X., Yang, R., Cross, A.H., and
Song, S.K. (2021). Diffusion Basis Spectrum Imaging Detects Axonal Loss After
Transient Dexamethasone Treatment in Optic Neuritis Mice. Front. Neurosci. *14*, 1–12.
10.3389/fnins.2020.592063.

61. Zhan, J., Lin, T.H., Libbey, J.E., Sun, P., Ye, Z., Song, C., Wallendorf, M., Gong, H.,
Fujinami, R.S., and Song, S.K. (2018). Diffusion basis spectrum and diffusion tensor
imaging detect hippocampal inflammation and dendritic injury in a virus-induced mouse
model of epilepsy. Front. Neurosci. *12*, 1–9. 10.3389/fnins.2018.00077.

62. Yushkevich, P.A., Piven, J., Hazlett, H.C., Smith, R.G., Ho, S., Gee, J.C., and Gerig, G.

(2006). User-guided 3D active contour segmentation of anatomical structures:
Significantly improved efficiency and reliability. Neuroimage *31*, 1116–1128.
10.1016/J.NEUROIMAGE.2006.01.015.

63. Mao, D.D., Gujar, A.D., Mahlokozera, T., Chen, I., Pan, Y., Luo, J., Brost, T., Thompson,
E.A., Turski, A., Leuthardt, E.C., et al. (2015). A CDC20-APC/SOX2 Signaling Axis
Regulates Human Glioblastoma Stem-like Cells. Cell Rep. *11*, 1809–1821.
10.1016/J.CELREP.2015.05.027.

64. Daly, M.J., Patterson, N., Mesirov, J.P., Golub, T.R., Tamayo, P., and Spiegelman, B.
(2003). PGC-1α-responsive genes involved in oxidative phosphorylation are coordinately
downregulated in human diabetes. Nat. Genet. *34*, 267–273.

65. Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A.,
Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S., et al. (2005). Gene set
enrichment analysis: A knowledge-based approach for interpreting genome-wide
expression profiles. Proc. Natl. Acad. Sci. U. S. A. *102*, 15545–15550.
10.1073/pnas.0506580102.

66. Koopmans, F., van Nierop, P., Andres-Alonso, M., Byrnes, A., Cijsouw, T., Coba, M.P.,
Cornelisse, L.N., Farrell, R.J., Goldschmidt, H.L., Howrigan, D.P., et al. (2019). SynGO:
An Evidence-Based, Expert-Curated Knowledge Base for the Synapse. Neuron *103*, 217-
234.e4. 10.1016/J.NEURON.2019.05.002.

67. Szklarczyk, D., Franceschini, A., Wyder, S., Forslund, K., Heller, D., Huerta-Cepas, J.,
Simonovic, M., Roth, A., Santos, A., Tsafou, K.P., et al. (2015). STRING v10: protein–
protein interaction networks, integrated over the tree of life. Nucleic Acids Res. *43*, D447–
D452. 10.1093/nar/gku1003.

68. Darnell, J.C., Van Driesche, S.J., Zhang, C., Hung, K.Y.S., Mele, A., Fraser, C.E., Stone,
E.F., Chen, C., Fak, J.J., Chi, S.W., et al. (2011). FMRP stalls ribosomal translocation on
mRNAs linked to synaptic function and autism. Cell *146*, 247–261.
10.1016/j.cell.2011.06.013.

69. Tasic, B., Yao, Z., Graybuck, L.T., Smith, K.A., Nguyen, T.N., Bertagnolli, D., Goldy, J.,
Garren, E., Economo, M.N., Viswanathan, S., et al. (2018). Shared and distinct
transcriptomic cell types across neocortical areas. Nature *563*, 72–78. 10.1038/s41586-
018-0654-5.

70. Langfelder, P., and Horvath, S. (2008). WGCNA: an R package for weighted correlation
network analysis. BMC Bioinformatics *9*, 559. 10.1186/1471-2105-9-559.

71. Huang, D.W., Sherman, B.T., and Lempicki, R.A. (2009). Systematic and integrative
analysis of large gene lists using DAVID bioinformatics resources. Nat. Protoc. *4*, 44–57.
10.1038/nprot.2008.211.

72. Sherman, B.T., Hao, M., Qiu, J., Jiao, X., Baseler, M.W., Lane, H.C., Imamichi, T., and
Chang, W. (2022). DAVID: a web server for functional enrichment analysis and functional
annotation of gene lists (2021 update). Nucleic Acids Res. *50*, W216–W221.
10.1093/nar/gkac194.

1503

Figure and table legends

Figure 1. Loss of USP7 in glutamatergic excitatory neurons leads to multiple disease-relevant behavioral deficits in mice.

(A)   Patients with the Hao-Fountain syndrome and *Usp7* conditional knockout (cKO) mice exhibit phenotypes in similar behavioral domains.

(B)   Double immunofluorescence of USP7 and neuron nuclear protein (NeuN) in cerebral cortex at postnatal day 42 (P42). USP7 protein was absent in most neuronal nuclei in *Usp7* cKO cortex. Scale bar 100μm.

(C)   Virtual optomotor system (VOS) test: Spatial frequency threshold for head turn. *Usp7* cKO mice (n = 6) showed reduced visual acuity compared to *Usp7* WT (n = 5) and cHet mice (n = 8). **$p<0.01$ by Bonferroni's multiple comparison test.

(D)   Motor strength test: Time to grip onto inverted screen. *Usp7* cKO mice (n = 8) showed shorter latency to fall than *Usp7* WT (n = 8) and cHet mice (n = 11). *$p<0.05$, **$p<0.01$ by Dunn's multiple comparison test.

(E)   Catwalk 1: Relative mean intensity distributed onto each paw. *Usp7* cKO mice (n = 8) put less weight onto front paw and more onto hind paw, compared to *Usp7* WT (n = 7) and cHet mice (n = 9). **$p<0.01$ by Tukey's multiple comparison test. See also Video S2.

(F)   Catwalk 2: Limb swing speed. *Usp7* cKO mice (n = 8) showed higher swing speed of both front and hind limb than *Usp7* WT (n = 7) and cHet mice (n = 9). **$p<0.01$, ***$p<0.001$ by Tukey's multiple comparison test. See also Video S2.

(G)   Fear conditioning day 1: Time spent in freezing for tone/shock pairing (n = 8, 11 & 6 for *Usp7* WT, *Usp7* cHet and *Usp7* cKO mice). No significant genotype effect by two-way ANOVA.

(H)   Fear conditioning day 2: Time spent in freezing for contextual conditioning. *Usp7* cKO mice spent less time in freezing than *Usp7* WT mice. Numbers of mice are the same as in (G). *$p<0.05$, **$p<0.01$ by Bonferroni's multiple comparison test (*Usp7* WT vs. cKO).

(I)   Fear conditioning day 3: Time spent in freezing for auditory cue conditioning. *Usp7* cKO mice spent less time in freezing after the tone than *Usp7* WT mice. Numbers of mice are the same as in (G). *$p<0.05$ by Bonferroni's multiple comparison test (*Usp7* WT vs. cKO).

(J)   Percentage of tail biters among all monitored *Usp7* cKO mice from 3 to 12 weeks old. Tail biters were identified if their co-caged mice had biting marks on the tails.

(K)   Representative biting injury on mice co-caged with *Usp7* cKO mice. See also Video S3.

Data are presented as mean ± SEM. See also Figure S1.

Figure 2. Deletion of p53 rescues apoptosis, impairment of visual acuity and auditory cue conditioning in *Usp7* cKO mice but not other behavioral deficits.

(A)   Immunofluorescence of apoptosis marker cleaved caspase 3 (c-cas 3) in cerebral cortex at different developmental time points. *Usp7* cKO cortex showed large scale of apoptosis at P0, whereas apoptosis is low at E15, P5 and P10. Scale bars all 200μm.

(B)   Schematic of USP7-Mdm2-p53 signaling pathway regulating apoptosis.

(C)   Immunofluorescence of c-cas 3 in P0 cortex of different *Trp53* and *Usp7* genotypes. *Trp53* heterozygosity diminished USP7 loss-induced apoptosis. Scale bar 200μm.

(D)   Quantification of c-cas 3 fluorescent signal, as shown in (C). Number of mice = 3, 3, 4 & 3 for *Usp7* WT, *Usp7* cKO, *Usp7* cKO; *Trp53* $^{+/-}$ & *Usp7* WT; *Trp53* $^{+/-}$, respectively. *p<0.05 by Tukey's multiple comparison test.

(E)   Virtual optomotor system (VOS) test: Spatial frequency threshold for head turn. *Usp7* cKO; *Trp53*$^{+/-}$ mice (n = 15) did not have impaired visual acuity, compared to *Usp7* WT; *Trp53* $^{+/-}$ (n = 14) and *Usp7* cHet; *Trp53* $^{+/-}$ mice (n = 9). No significant genotype effect by one-way ANOVA.

(F)   Motor strength test: Time to grip onto inverted screen. *Usp7* cKO mice (n = 10) showed shorter latency to fall than *Usp7* WT; *Trp53*$^{+/-}$ (n = 10) and *Usp7* cHet; *Trp53*$^{+/-}$ mice (n = 14). *p<0.05, **p<0.01 by Dunn's multiple comparison test.

(G)   Catwalk 1: Relative mean intensity distributed onto each paw. *Usp7* cKO; *Trp53*$^{+/-}$ mice (n = 5) put less weight onto front paw and more onto hind paw than *Usp7* WT; *Trp53*$^{+/-}$ (n = 7) and *Usp7* cHet; *Trp53*$^{+/-}$ mice (n = 8). ***p<0.001, ****p<0.0001 by Tukey's multiple comparison test. See also Video S4.

(H)   Catwalk 2: Limb swing speed. *Usp7* cKO; *Trp53*$^{+/-}$ mice showed slightly higher swing speed than *Usp7* cHet; *Trp53*$^{+/-}$ mice. Numbers of mice are the same as in (G). *p<0.05 by Tukey's multiple comparison test. See also Video S4.

(I)   Fear conditioning day 1: Time spent in freezing for tone/shock pairing (n = 12, 6 & 11 for *Usp7* WT, *Usp7* cHet and *Usp7* cKO mice). No significant genotype effect by two-way ANOVA.

(J)   Fear conditioning day 2: Time spent in freezing for contextual conditioning. *Usp7* cKO; *Trp53*$^{+/-}$ mice spent less time in freezing than *Usp7* WT; *Trp53*$^{+/-}$ mice. Numbers of mice are the same as in (H). *p<0.05, by Bonferroni's multiple comparison test (*Usp7* WT vs. cKO).

(K)   Fear conditioning day 3: Time spent in freezing for auditory cue conditioning. *Usp7* cKO; *Trp53*$^{+/-}$ mice had no difference in freezing after the tone, compared to *Usp7* WT; *Trp53*$^{+/-}$ and *Usp7* cHet; *Trp53*$^{+/-}$ mice. Numbers of mice are the same as in (H). No significant genotype effect by two-way ANOVA.

(L)   Percentage of tail biters among all monitored *Usp7* cKO; *Trp53*$^{+/-}$ mice from 3 to 12 weeks old. Tail biters were counted if their co-caged mice had biting marks on their tails.

(M)   Representative biting injury on mice co-caged with *Usp7* cKO; *Trp53*$^{+/-}$ mice.

Data are presented as mean ± SEM. See also Figure S2, S3 and S4.

 Table 1 Phenotype comparison between *Usp7* cKO mice and *Usp7* cKO; *Trp53*$^{+/-}$ mice.
 Phenotypes significantly different from *Usp7* WT or *Usp7* WT; *Trp53*$^{+/-}$ mice are in bold.
 The whole brain volumes exclude olfactory bulbs. Hippo = hippocampus. na = not
 available.

Figure 3. Loss of USP7 reduces synapses in cortex.

(A)   Flowchart to profile the proteome of the cerebral cortex of *Usp7* cKO mice with tandem mass tag-mass spectrometry (TMT-MS).

(B)   Volcano plot of all profiled proteins in TMT-MS. Up-regulated and down-regulated proteins (*Usp7* cKO vs. WT) are labeled in red and blue, respectively. P-value were calculated by 2-tail unpaired t-test.

(C)   Organic layout of enriched GO gene sets based on TMT protein abundance in (B). Nodes represent gene sets (q-value < 0.01) and size of nodes represents how many genes contained in gene sets. Edges represent GO-defined relations (similarity > 0.5) and thickness of edges represents similarity between gene sets. Most enriched gene sets are related to synapse and circled in red. All gene sets shown are enriched in *Usp7* WT and de-enriched in *Usp7* cKO.

(D)   GO terms enriched from ranked list of down-regulated proteins (*Usp7* cKO vs. WT fold change < 0.75).

(E)   Ventricular injection of AAV PHP.eB Syn-GFP-P2A-GFPf at P0/1 to sparsely label pyramidal neurons in cerebral cortex and dendritic spine analysis at P18.

(F)   Representative GFP micrographs of dendritic spines on apical and basal dendrites. Scale bar 5μm.

(G, H, I) Spine density on apical, basal and all dendrites of pyramidal neurons in motor cortex. *Usp7* cKO; *Trp53*$^{+/-}$ neurons (n = 39 & 39 neurons for apical dendrites & basal dendrites from 3 mice) showed reduced spine density than *Usp7* WT; *Trp53*$^{+/-}$ neurons (n = 40 & 50 neurons for apical dendrites & basal dendrites from 3 mice). ****p<0.0001 by 2-tail unpaired t-test.

Data are presented as mean ± SEM. See also Figure S5&6.

1612 **Figure 4. Unbiased proteomic characterizations of USP7 in cortical neurons.**
1613 (A)   Domain structure of mouse USP7 protein. R105, D165 and W165 on the TRAF domain
1614   are three amino acids critical for substrate binding[18,23]. USP: Ubiquitin specific protease
1615   domain. Ubl: Ubiquitin like domain.
1616 (B)   Silver stain of FLAG-immunoprecipitated USP7 TRAF domain (arrow) with and without
1617   RDW mutations (R105A + D165A + W166A).1.5% of all eluted sample were stained while
1618   rest of the sample were analyzed by liquid chromatography-mass spectrometry (LC-MS).
1619 (C)   Protein-protein association map of USP7 TRAF interactors (full STRING network,
1620   confidence > 0.4). Edge thickness represents confidence in association between proteins.
1621   Shown are only 50 interactors with highest gene length-normalized spectral counts.
1622   Ribonucleoprotein complex is colored in red. Spliceosomal complex is colored in green.
1623   SWI/SNF complex is colored in blue. Mitochondrial pyruvate dehydrogenase complex is
1624   colored in yellow.
1625 (D)   Molecular component GO enrichments of top 50 interactors of the USP7 TRAF domain.
1626   Different molecular components are colored as in (C).
1627 (E)   Disease gene enrichment analysis of the interactors of USP7 TRAF domain. ID =
1628   intellectual disability. SCZ = schizophrenia. ns. = not significant.
1629 (F)   Flowchart to profile proteomic change with tandem mass tag-mass spectrometry (TMT-
1630   MS) in response to acute *Usp7* knockout in primary cortical neurons. Neurons were
1631   dissociated from *Usp7*$^{fl/fl}$; *Trp53*$^{-/-}$ embryos at E15 and subject to Cre lentiviral
1632   transduction at DIV3. Cre harboring Y331F mutation is catalytically dead (Cre dead) and
1633   served as negative control.
1634 (G)   Western blot validation of USP7 depletion and Cre expression in Cre WT-transduced
1635   neurons over time. 14-3-3 is loading control.
1636 (H)   Heatmap of TMT protein abundance of USP7 and its candidate substrates. Candidate
1637   substrates were sorted out by smallest p-value of Cre main effect (two-way ANOVA) and
1638   shortest Euclidean distance to USP7.
1639 (I)   Western blot quantitation of candidate substrates with the cerebral cortex of *Usp7* cKO;
1640   *Trp53*$^{+/-}$ (n = 3 mice) vs. *Usp7* WT; *Trp53*$^{+/-}$ (n = 3 mice) at P0. Western blot images are in
1641   Figure S9.
1642 (J)   RT-qPCR of candidate substrates with the cerebral cortex of *Usp7* cKO; *Trp53*$^{+/-}$ (n = 5
1643   mice) vs. *Usp7* WT; *Trp53*$^{+/-}$ (n = 3 mice) at P0. No candidate substrate in the analysis
1644   showed RNA reduction with the loss of USP7.
1645 Data are presented as mean ± SEM. See also Figure S7, S8 and S9.
1646
1647

Figure 5. USP7 targets and deubiquitinates Ppil4 to promote synaptogenesis.
(A)   Streptavidin pull-down of HEK293T lysate with overexpression of Streptavidin-binding
peptide-FLAG (SFB)-tagged USP7 followed by western blot of multiple endogenous
candidate substrates. 2% input lysate is shown on the left.
(B, C, D) Density of dendritic spine on apical, basal and all dendrites of pyramidal-shaped
neurons in (H). Two independent lentiviral shRNA knocking down Ppil4 reduced spine
density compared with control shRNA targeting Luciferase (Luci). Number of neurons
(apical dendrties) = 24, 27, 23, 20, 23, 18, 16 & 17 for Luci, Ppil4i.1, Ppil4i.2, Smarcal1i.1,
Smarcal1i.2, Trip12i, Ring1i & Sub1i. Number of neurons (basal dendrties) = 22, 24, 26,
20, 24, 17, 15 & 16 for Luci, Ppil4i.1, Ppil4i.2, Smarcal1i.1, Smarcal1i.2, Trip12i, Ring1i &
Sub1i. *$p<0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$ by Dunnett's multiple comparison test
(compared to Luci).
(E, F, G) Density of PSD95+ dendritic spine on apical, basal and all dendrites of cortical
neurons in (H) and Figure S10I. Two independent lentiviral shRNA knocking down Ppil4
reduced PSD95+ spine density compared with control shRNA targeting Luciferase (Luci).
Number of neurons are the same as in (B, C, D). *$p<0.05$, **$p<0.01$, ***$p<0.001$,
****$p<0.0001$ by Dunnett's multiple comparison test (compared to Luci).
(H)   Representative images show the effect of Ppil4 knockdown on dendritic spines of cortical
neurons at DIV18. PSD95 puncta decorating the dendritic spines were immunostained
along with GFP. Arrows show dendritic spines. Scale bar 5μm.
(I)   Immunoprecipitation of endogenous Ppil4 in HEK293T lysates followed by western blot of
GFP-USP7 and Ppil4. 2% Input lysate is shown on the left. GFP-USP7 was co-
immunoprecipiated with Ppil4.
(J)   Tandem Ubiquitin Binding Entities (TUBE) pull-down of neuronal lysate with USP7 loss
driven by lentiviral Cre, followed by immunoblotting of Ppil4. Input lysates are shown at
the bottom. Endogenous Poly-Ubiquitinated Ppil4 increased with USP7 loss (Cre WT).
A.U. = aribitary unit.
(K)   *In vitro* deubiquitination assay using FLAG-immunoprecipitated Ppil4 from HEK293T cells.
Incubation with recombinant His-USP7 at 37°C for 1 hour converted high-molecular
weight ubiquitinated Ppil4 (dashed rectangle) into low-molecular weight Ppil4 (arrow).
(L)   Densitometric quantification of high-molecular weight Ppil4 over all Ppil4 signal in (K)
over 3 biological replicates. **$p<0.01$ by 2-tail unpaired t-test.
(M)   Western blot of USP7 with mutations from the Hao-Fountain syndrome patients in
HEK293T cells. USP7 C223A is catalytically dead.
(N)   Western blot of endogenous Ppil4 in HEK293T cells expressing different USP7 variants.
(O)   Densitometric quantification of Ppil4 signal as in (N) over 4 biological replicates.
Repeated measurement one way ANOVA. *$p<0.05$ by Dunnett's multiple comparison test.
(P)   Schematic of USP7-Ppil4 signaling pathway regulating dendritic spine density in
glutamatergic cortical neuron.
Data are presented as mean ± SEM. See also Figure S10.
1688

1689

Figure S1. Additional phenotypic characterization of *Usp7* cKO mice. Related to Figure 1 and Table 1.

(A) Immunofluorescence of USP7 in cerebral cortex at embryonic day 14 (E14). USP7 is localized to both Ki-67+ ventricular-subventricular zone (V-SVZ) and cortical plate (CP) in USP7 WT cortex. But in *Usp7* cKO cortex, USP7 is depleted only in post-mitotic cortical plate (dashed outline). Scale bar 100μm.

(B) Body weight curve from P6 to P48. *Usp7* cKO mice (n = 7) exhibited body weight growth delay, compared to *Usp7* WT (n = 10) and cHet mice (n = 7).

(C) Body weight of adult mice. Number of mice = 8, 11 & 9 for *Usp7* WT, cHet & cKO, respectively. No significant genotype effect by one-way ANOVA.

(D) Motor coordination test 1: Ledge test. *Usp7* cKO mice (n = 8) showed shorter latency to fall from ledge than *Usp7* WT (n = 7) and cHet mice (n = 9). **$p<0.01$ by Dunn's multiple comparison test.

(E) Motor coordination test 2: Platform test. *Usp7* cKO mice showed shorter latency to fall from platform than *Usp7* WT mice. Numbers of mice are the same as in (D). **$p<0.01$ by Dunn's multiple comparison test.

(F) Motor coordination test 3: Pole test, time to climb down the pole. Numbers of mice are the same as in (D). No significant genotype effect by one-way ANOVA.

(G) Motor coordination test 4: Pole test, time to turn at the tip of the pole. *Usp7* cKO mice spent more time in turning than *Usp7* WT and cHet mice. Numbers of mice are the same as in (D). *$p<0.05$, **$p<0.01$ by Dunn's multiple comparison test.

(H) Catwalk 3: Stride length. *Usp7* cKO mice showed slightly shorter average stride length than control. Numbers of mice are the same as in (D). *$p<0.05$ by Tukey's multiple comparison test.

(I) Catwalk 4: Swing time. *Usp7* cKO mice used less time swinging front and hind legs for each step, compared to *Usp7* WT and cHet mice. Numbers of mice are the same as in (D). *$p<0.05$, ***$p<0.001$, ****$p<0.0001$ by Tukey's multiple comparison test.

(J) Hearing test: Magnitude of response to acoustic startle at different dB level. Number of mice = 5, 8 & 6 for *Usp7* WT, *Usp7* cHet & *Usp7* cKO. No significant genotype effect by two-way ANOVA.

(K) Y maze: Alternative rate in choosing arm to enter. Number of mice = 4, 6 & 5 for *Usp7* WT, *Usp7* cHet & *Usp7* cKO. No significant genotype effect by one-way ANOVA.

Data are presented as mean ± SEM.

1723
1724
1725

Figure S2. Loss of USP7 in glutamatergic neurons induces apoptosis in a cell autonomous manner. Related to Figure 2.

(A)   Double immunofluorescence of cleaved caspase 3 (c-cas 3) and Cre. An expanded region shows overlap (arrows) of cleaved caspase 3 and Cre, supporting that the apoptotic cells are glutamatergic neurons. Scale bar 200µm.

(B)   Western blot validation of USP7 RNAi in HEK293Tcells. Both USP7i.1 and USP7i.2 diminished expression of GFP-USP7. 14-3-3 is loading control.

(C)   Immunocytochemistry of c-cas 3 in primary cortical neurons dissociated from wild-type E14 cortices and transfected with a GFP plasmid. Scale bar 100µm. Cells were fixed at DIV5, 6 and 7. At the bottom are close-up of neurons with and without c-cas 3 signal.

(D)   Quantification of apoptotic percentage, defined as the percentage of c-cas 3 and GFP double positive neurons among all GFP positive neurons. Cortical neurons were transfected with USP7 shRNA plasmids at 3 days *in vitro* (DIV3) and analyzed at DIV5, 6 and 7. Data were collected from three independent batches of neuron culture, three wells per condition within each batch. USP7 knockdown by two independent shRNA led to a progressive elevation in apoptotic percentage. Empty vector U6 and knockdown of an irrelevant deubiquitinase, USP30, served as negative controls. ***$p<0.001$, ****$p<0.0001$ by Tukey's multiple comparison test.

Data are presented as mean ± SEM.

Figure S3. Deletion of p53 rescues USP7 loss-induced apoptosis but minimally restores the brain volume reduction in *Usp7* cKO mice. Related to Figure 2 and Table 1.

(A)   Immunofluorescence of c-cas 3 in the cerebral cortex of *Usp7* cKO mice at P0 with deletion of one or both allele of *Trp53*. C-cas 3 signal remained sparse with either *Trp53* heterozygosity or homozygosity. Scale bar 200μm.

(B)   Immunofluorescence of USP7 in primary cortical neurons dissociated from *Usp7*$^{fl/fl}$; *Trp53*$^{+/-}$ embryos. Neurons transduced with Cre WT showed loss of USP7 protein at DIV14. Scale bar 25μm.

(C)   MTS assay of *Usp7*$^{fl/fl}$ cortical neurons transduced by lentiviral Cre with different *Trp53* genotypes. USP7 knockout in *Trp53*$^{+/+}$ neurons (n = 5 wells) strongly reduced cell viability, indicating of neuronal death. No significant viability decrease in *Trp53*$^{+/-}$ (n = 3 wells) and *Trp53*$^{-/-}$ neurons (n = 5 wells). Puromycin (2μg/mL, 24hrs) treated neurons serve as positive control. ***$p<0.001$ by Šídák's multiple comparison test between Cre WT and Cre dead.

(D)   Representative segmentation on FA map and DWI for volumetric measurement with diffusion-weighted *ex vivo* MRI. Cerbral cortex and hippocampus are shown in blue and red, respectively.

(E)   Volumetric measurement with *ex vivo* MRI. *Usp7* cKO brains (P30-32) are smaller than *Usp7 WT*. Deletion of p53 minimally stored the volume of cortex and hippocampus. OB = olfactory bulbs. Number of mice = 5, 4, 5 & 5 for *Usp7* WT, *Usp7* cKO, *Usp7* WT; *Trp53* $^{+/-}$ & *Usp7* cKO; *Trp53* $^{+/-}$. *$p<0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$ by Tukey's multiple comparison test.

Data are presented as mean ± SEM.

Figure S4. Additional phenotypic characterization of *Usp7* cKO; *Trp53*$^{+/-}$ mice. Related to
Figure 2 and Table 1.
(A)   Body weight curve from P6 to P48. *Usp7* cKO; *Trp53*$^{+/-}$ mice (n = 6) exhibited body
weight growth delay, compared to *Usp7* WT; *Trp53*$^{+/-}$ (n = 10) and *Usp7* cHet; *Trp53*$^{+/-}$
mice (n = 5).
(B)   Body weight of adult mice. Number of mice = 10, 14 & 10 for *Usp7* WT, cHet & cKO;
Trp53 $^{+/-}$, respectively. No significant genotype effect by one-way ANOVA.
(C)   Motor coordination test 1: Ledge test. *Usp7* cKO; *Trp53*+/- mice showed slightly shorter
latency to fall from ledge than control mice. Numbers of mice are the same as in (B).
*p<0.05 by Dunn's multiple comparison test.
(D)   Motor coordination test 2: Platform test. *Usp7* cKO; *Trp53*$^{+/-}$ mice showed shorter latency
to fall from platform than control mice. Numbers of mice are the same as in (B). **p<0.01
by Dunn's multiple comparison test.
(E)   Motor coordination test 3: Pole test, time to climb down the pole. Numbers of mice are
the same as in (B). No significant genotype effect by one-way ANOVA.
(F)   Motor coordination test 4: Pole test, time to turn at the tip of the pole. *Usp7* cKO; *Trp53*$^{+/-}$
mice spent more time in turning than control mice. Numbers of mice are the same as in
(B). **p<0.01, ***p<0.001 by Dunn's multiple comparison test.
(G)   Catwalk 3: Stride length. *Usp7* cKO; *Trp53*$^{+/-}$ mice showed shorter average stride length
than control. Number of mice = 7, 8 & 5 for *Usp7* WT, cHet & cKO; *Trp53* $^{+/-}$. *p<0.05,
**p<0.01 by Tukey's multiple comparison test.
(H)   Catwalk 4: Swing time. *Usp7* cKO; *Trp53*$^{+/-}$ mice used less time swinging front and hind
legs for each step, compared to control mice. Numbers of mice are the same as in (G).
*p<0.05, ***p<0.001 by Tukey's multiple comparison test.
(I)   Hearing test: Magnitude of response to acoustic startle at different dB level. Number of
mice = 10, 14 & 10 for *Usp7* WT, cHet & cKO; *Trp53* $^{+/-}$. No significant genotype effect by
two-way ANOVA.
Data are presented as mean ± SEM.
1799
1800

Figure S5. Additional analyses of reduced synaptic proteins and dendritic spine
morphology in *Usp7* cKO mice. Related to Figure 3.
(A&B) Sunburst plot of SynGO terms enriched in down-regulated proteins of the TMT proteome
of *Usp7* cKO cortex. Cellular component SynGO terms in (A), biological process SynGO
terms in (B).
(C)   TMT relative abundances of down-regulated synaptic proteins in *Usp7* cKO cortex. The
individual proteins within SynGO: Postsynapse, SynGO: Presynapse and in both were
segregated into different panels.
(D)   Representative GFP micrographs of dendritic spines of different shapes, including
mushroom, thin, stubby and filopodial spines. Scale bar 1μm.
(E-J) Percentages of dendritic spines of different shapes in *Usp7* cKO and WT cortex. (E-G)
shows total percentages of dendritic spines of different shapes. (H-J) shows by neuron
percentages of dendritic spines of different shapes. There was an increase in the
proportion of filopodial dendritic spines in *Usp7* cKO pyramidal neurons (n = 25 & 22
neurons for apical dendrites & basal dendrites from 3 mice), compared to *Usp7* WT
pyramidal neurons (n = 20 & 19 neurons for apical dendrites & basal dendrites from 3
mice). $*p < 0.05$, $**p < 0.01$ by Šídák's multiple comparison test.
Data are presented as mean ± SEM.
1819
1820

1821

Figure S6. USP7 loss does not cause significant change in dendritic morphology in
primary cortical neurons with Trp53 heterzygosity. Related to Figure 3.
(A)   Representative DIV11 images of cortical pyramidal neurons and their Imaris dendritic
traces. For the Imaris dendritic traces, apical dendrties are highlighted in yellow. The
cortical neurons were dissociated from $Usp7^{fl/fl}$; $Trp53^{+/-}$ E15 embryos and transduced
with Cre WT or Cre dead lentivirus at DIV3.
(B)   Sholl profile of pyramidal neurons (n = 40 & 32 for Cre dead & WT) and all neurons (n =
55 & 44 for Cre dead & WT). Shaded area represents standard deviation. Profiles of
neurons transduced with Cre dead and Cre WT are colored in red and green, respectively.
(C)   Sum of dendrite length in pyramidal neurons, apical dendrites vs. basal dendrites (n = 40
& 32 for Cre dead & WT). No significant Cre effect by two-way ANOVA.
(D)   Sum of dendrite length in pyramidal neurons, across 1-4 branch levels. Numbers of
neurons are the same as in (C). No significant Cre effect by two-way ANOVA.
(E)   Average length of primary dendrites in pyramidal neurons. Numbers of neurons are the
same as in (C). No significance by 2-tail unpaired t-test.
(F)   Number of primary dendrites in pyramidal neurons. Numbers of neurons are the same as
in (C). No significance by 2-tail unpaired t-test.
(G)   Soma 3D-volumes of pyramidal neurons (n = 65 & 57 for Cre dead & WT). No
significance by 2-tail unpaired t-test.
Data are presented as mean ± SEM.

1842

1843

Figure S7. Validation of USP7 TRAF domain expression and interaction for IP-MS. Related
to Figure 4.
(A) Immunoprecipitation of GFP-TRAF domain followed by western blot of myc3-Mdm2 and
GFP. Lysates were prepared from HEK293T cells transfected with myc3-Mdm2 and GFP-
tagged USP7-TRAF domain. DW double and RDW triple mutations (All mutated into
Alanine) on the USP7-TRAF domain disrupted the interaction with MDM2. FT as flow-
through lysates.
(B) Mostly nuclear localization of truncated GFP-USP7 in rat cortical neurons at DIV9,
including USP7-TRAF domain (USP7 amino acid 55-209). Neurons were co-transfected
with pCAG-mCherry with Ca phosphate. USP7 fl as USP7 full length. Scale bar 50µm.
(C) Immunocytochemistry of FLAG showed mostly nuclear expression of FLAG-USP7 TRAF
domain in rat cortical neurons with and without RDW mutations at DIV10.
Immunocytochemistry of Map2 shows neuronal morphology. Scale bar 100µm.
1857
1858

Figure S8. Additional analyses of the TMT proteome with acute USP7 loss in cortical
neurons. Related to Figure 4.
(A)   PCA of 16 samples based on TMT abundance of all 8304 proteins detected by TMT-MS.
Principal component 1 (PC1) separates DIV6 and DIV8 samples whereas principal
component 2 (PC2) separates Cre WT and Cre dead samples.
(B)   TMT protein abundance of candidate substrates in previous *in vivo* TMT-MS (Figure 3A-
D). Most candidate substrates showed various degrees of reduction in the cerebral cortex
of *Usp7* cKO mice at P17. n = 5 for both *Usp7* WT; *Trp53* $^{+/-}$ and *Usp7* cKO; *Trp53* $^{+/-}$
mice.
(C)   Dendrogram and heatmap of TMT protein adjacencies by weighted gene co-expression
network analysis (WGCNA). WGCNA segragates 2627 differentially expressed proteins
(Cre effect $p<0.05$ or Cre time interaction effect $p<0.05$ by two-way ANOVA) into 10
modules in different colors.
(D)   Module-trait relationships. Correlations and p-values between module eigengene (ME)
and Cre (Cre WT as 1, Cre dead as 0) or time (DIV6 or DIV8) are denoted and color-
coded according to the color legend.
(E)   Membership of candidate substrates in each module.
(F)   GO annotation for each module. Turquiose module is enriched in ribonucleoprotein
complex and splicing-related terms. Brown module is enriched in microtubule and axon-
related terms. Black module is enriched in mitochondrion-related terms. Yellow module is
enriched in E3 ligase-related terms.
Data are presented as mean ± SEM.
1881
1882
1883

1884   **Figure S9. Western blot of candidate substrates in the cerebral cortex of P0 *Usp7* cKO;**
1885   ***Trp53*^{+/-} mice.** Related to Figure 4(H). Three biological replicates of each genotype were
1886   denoted in numbers. (A) Phip, USP7 and Xiap. (B) Ppil4, Ring1 and Gpd1. (C) Ubxn7,
1887   Ruvbl2 and Rnf2. (D) Chfr and Sub1. (E) Fto, Gfap and Cre. (F) Rnf220, Cfdp1 and Phc.
1888   (G) Fam172a. (H) Trip12. (I) Smarcal1. 14-3-3 and Lamin A/C are loading controls.
1889
1890

Figure S10. Additional characterization of the effect of candidate substrate knockdown on dendritic spine morphogenesis. Related to Figure 5.

(A-E)   RT-qPCR validation of candidate substrates RNAi (Ppil4, Smarcal1, Trip12, Ring1 and Sub1) in primary cortical neurons. Neurons were transduced with shRNA lentivirus at DIV4, lysed and harvested at DIV7. n = 3 biological replicates for all groups.

(F, G, H) Density of PSD95 puncta on apical, basal and all dendrites of pyramidal-shaped neurons in (Figure 5H). Two independent lentiviral shRNA knocking down Ppil4 reduced PSD95 puncta density compared with control shRNA targeting Luciferase (Luci). Number of neurons (apical dendrties) = 24, 27, 23, 20, 23, 18, 16 & 17 for Luci, Ppil4i.1, Ppil4i.2, Smarcal1i.1, Smarcal1i.2, Trip12i, Ring1i & Sub1i. Number of neurons (basal dendrties) = 22, 24, 26, 20, 24, 17, 15 & 16 for Luci, Ppil4i.1, Ppil4i.2, Smarcal1i.1, Smarcal1i.2, Trip12i, Ring1i & Sub1i. $**p<0.01$, $***p<0.001$, $****p<0.0001$ by Dunnett's multiple comparison test (compared to Luci).

(I)   Representative images show the effect of knockdown of candidate substrates on dendritic spines of cortical neurons at DIV18. PSD95 puncta decorating the dendritic spines were immunostained along with GFP. Arrows show dendritic spines. Scale bar 5µm.

Data are presented as mean ± SEM.

1910 **Other Supplemental Materials:**
1911
1912 Table S1. GSEA statistic report of TMT proteomics of *Usp7* cKO and control cortex.
1913 Table S2. Oligonucleotides used in this study.
1914
1915
1916 Video S1 Abnormal gait of *Usp7* cKO mouse. Mouse age P33.
1917 Video S2 *Usp7* cKO vs WT mouse CatWalk with 10x slow speed.
1918 Video S3 Aggressive biting in *Usp7* cKO mouse. Mouse age P33.
1919 Video S4 *Usp7* cKO; *Trp53*$^{+/-}$ vs *Usp7* WT; *Trp53*$^{+/-}$ mouse CatWalk with 10x slow speed.

Figure 1

Figure 2

Table 1

Test	Phenotype	*Usp7* cKO mice	*Usp7* cKO; *Trp53*[+/-] mice
c-cas 3 immunofluorescence	**Apoptosis**	**Ectopically induced**	**Normal**
Ex vivo MRI	Brain volume	-37.4% (cortex) -39.1% (hippo) -20.0% (whole brain)	-26.3% (cortex) -30.9% (hippo) -21.3% (whole brain)
Body weight measurement	Body weight	Growth delay	Growth delay
VOS test	**Visual acuity**	**Impaired**	**Normal**
Inverted screen test	Motor strength	Impaired	Impaired
Ledge, platform and pole tests	Motor coordination	Impaired	Impaired
Catwalk	Gait	Impaired	Impaired
Acoustic startle response	Hearing ability	Normal	Normal
Fear conditioning	Contextual conditioning	Impaired	Impaired
	Cued conditioning	**Impaired**	**Normal**
Y maze	Spatial working memory	Normal	na
Tail biting	Aggression	Observed in 33.3% mice	Observed in 34.62% mice

Figure 3

Figure 4

Figure 5

Figure S1

Figure S2

Figure S3

A

Trp53 $^{+/-}$ *Trp53* $^{-/-}$

Usp7 WT

Usp7 cKO

Hoechst C-cas 3

B

USP7   Hoechst

Cre dead

Cre WT

C

MTS assay

Cell viability (OD490nm) normalized to Cre dead

- *Usp7* $^{fl/fl}$
- *Usp7* $^{fl/fl}$; *Trp53* $^{+/-}$
- *Usp7* $^{fl/fl}$; *Trp53* $^{-/-}$

***   ns.   ns.

Cre WT Cre dead   Cre WT Cre dead Puromycin   Cre WT Cre dead Puromycin

D

FA Map   DWI

E

Ex vivo volumetric measurement

Volume (A.U.) normalized to USP7 WT littermates

- *Usp7* WT
- *Usp7* cKO
- *Usp7* WT; *Trp53* $^{+/-}$
- *Usp7* cKO; *Trp53* $^{+/-}$

****   **   ****   ***   *   **

Cortex   Hippocampus   Whole brain w/o OB

Figure S4

Figure S5

Figure S6

Figure S7

Figure S8

Figure S9

Figure S10

PSD95 GFP