Force-sensing LPHN2 is indispensable for normal hearing and mediates MET in cochleae through Gs and CNGA3 coupling

Zhao Yang[1,3#], Ming-Wei Wang[4#], Shu-Hua Zhou[1,2#], Zhi-Chen Song[2#], Jia-Rui Gao[2#], Wen-Wen Liu[8], Qi-Yue Zhang[1], Ying Guan[1,3], Lei Xu[8], Chuan Wang[7*], Zhi-Gang Xu[6*], Xiao Yu[2*], Jin-Peng Sun[1,3,4,5*]

Affiliations:

[1]NHC Key Laboratory of Otorhinolaryngology, Qilu Hospital of Shandong University, and Department of Biochemistry and Molecular Biology, School of Basic Medical Sciences, Cheeloo College of Medicine, Shandong University, Jinan 250012, China.

[2]Key Laboratory of Experimental Teratology of the Ministry of Education, Department of Physiology, School of Basic Medical Sciences, Cheeloo College of Medicine, Shandong University, Jinan 250012, China.

[3]Advanced Medical Research Institute, Cheeloo College of Medicine, Shandong University, Jinan 250012, China.

[4]Department of Clinical Laboratory, The Second Hospital, Cheeloo College of Medicine, Shandong University, Jinan 250012, China.

[5]Key Laboratory of Molecular Cardiovascular Science of the Ministry of Education, Department of Physiology and Pathophysiology, School of Basic Medical Sciences, Peking University, Beijing 100191, China.

[6]Shandong Provincial Key Laboratory of Animal Cells and Developmental Biology, Shandong University School of Life Sciences, Qingdao 266237, China

[7]Department of Pharmacology, Hebei Medical University, Shijiazhuang 050000, China

[8]Department of Otolaryngology-Head and Neck Surgery, Shandong Provincial ENT Hospital, Cheeloo College of Medicine, Shandong University, Jinan 250014, China.

# These authors contributed equally to this work.

* Correspondence: Jin-Peng Sun, E-mail: sunjinpeng@sdu.edu.cn

                Xiao Yu, E-mail: yuxiao@sdu.edu.cn

31   Zhi-Gang Xu, E-mail: xuzg@sdu.edu.cn

32   Chuan Wang, E-mail: wangchuan@hebmu.edu.cn

Abstract

Auditory perception enables acoustic experiences, including listening to melodious

music and engaging in linguistic communication. Conversion of the force sensation into

electrical signals via mechanoelectrical transduction (MET) in cochlear hair cells is the key

step for auditory perception; however, the sound receptor for auditory perception at the

molecular level is not clear. Here, we found that hair cell-specific deficiency of the G protein-

coupled receptor LPHN2 in *Pou4f3-CreER$^{+/-}$Lphn2$^{fl/fl}$* mice or pharmacological blockade of

CNG ion channels in mice severely impaired hearing. Importantly, sensation of force by

LPHN2 not only increased intracellular cAMP and calcium levels but also elicited rapid

membrane depolarization of cochlear hair cells via Gs-dependent CNGA3 coupling. Both

LPHN2 and CNGA3 were expressed at the stereocilia and cuticular plates and associated with

each other. Notably, hearing loss in LPHN2-deficient mice was almost fully reversed by re-

expression of LPHN2-GAIN in cochlear hair cells. Therefore, we propose that LPHN2 acts as

a direct sound sensor in cochlear hair cells by mediating MET through Gs and CNGA3

coupling. Furthermore, both GPCR members and ion channels modulated by second

messengers might actively participate in the MET process during auditory perception.

Introduction

Auditory perception is one of our basic senses and provides us with the opportunity to

experience the features of an acoustic world, including melodious music, the chirps of birds

and the tinkling of spring water. It is also our main gateway for communicating with each

other via speech and alerts us to possible dangers. By converting mechanical force into

electrical signals, mechanoelectrical transduction (MET) in cochlear hair cells acts as the

central event during auditory perception (Zheng and Holt, 2021). Importantly, recent studies

have identified a series of candidate components of the MET channel complex, which include

the transmembrane membrane proteins TMC1/2, TMIE, and LHFPL5 and auxiliary subunits

such as calcium (Ca^{2+}) and integrin binding protein 2 (CIB2) and ankyrin (Cunningham et al.,

2020; Liang et al., 2021; Pan et al., 2013; Tang et al., 2020; Xiong et al., 2012). Despite

several findings suggesting that TMC1 and TMC2 are probable pore-forming subunits of the

MET channel complex, direct evidence for force sensation by TMC1/2 is lacking (Jia et al.,

2020; Pan et al., 2018). In addition to these MET components concentrated at the tips of the

stereocilia, another mechanosensitive ion channel, Piezo2, which is expressed at the cuticular

plates of cochlear hair cells, has also been suggested to be a MET regulator (Wu et al., 2017).

Despite the significant progress in research, the molecular mechanisms underlying MET in

cochlear hair cells are unclear, and the components of MET have not been fully elucidated.

In addition to ion channels, G protein-coupled receptors (GPCRs) are well known to

function as important sensory receptors that mediate the sensation of light, odour, itch and at

least three qualities of taste (Buck and Axel, 1991; Yang et al., 2021; Yau and Hardie, 2009;

Zhang et al., 2003). Importantly, several GPCRs expressed in the cochlea have been shown to

be essential for hearing. For instance, whereas CELSR1 and Fzd3/6 are required for the

maintenance of cochlear planar cell polarity, LGR5 regulates the self-renewal abilities of

cochlear hair cells (Duncan et al., 2017; Shi et al., 2013; Wang et al., 2006). Moreover, many

mutations of VLGR1 are responsible for severe hearing loss (Guan et al., 2023). Although

GPCRs play important regulatory roles in cochlear development, it is still unknown whether

GPCRs directly participate in the MET process and regulate auditory perception.

In our parallel work described in two companion papers, we found that two adhesion

GPCRs, GPR133 and LPHN2, act as equilibrioceptors by directly sensing force and

mediating MET in vestibular hair cells. In our present study, we further found that

mechanosensitive LPHN2 showed a unique distribution in cochlear hair cells, being

immunostained at both stereocilia and cuticular plates. Global knockout or hair-cell-specific

knockout of *Lphn2* in mice impaired hearing. Mechanistically, force sensation by LPHN2

induced rapid Ca^{2+} signalling and acute membrane current changes by activating the Gs-

cAMP-CNGA3 signalling axis. Furthermore, pharmacological inhibition of CNGA3 activity

abrogated LPHN2-mediated MET and caused hearing impairment in mice. Overall, our

findings identify a mechanosensitive GPCR that directly participates in the effects of MET on

hearing through functional coupling to CNGA3.

Results

Expression of LPHN2 in cochlear hair cells

Previous single-cell RNA sequencing (scRNA-seq) data have indicated that LPHN2 is expressed primarily in cochlear hair cells with a dynamic pattern (Ranum et al., 2019). The LPHN2 expression frequency decreased from approximately 75% in embryonic cochlear hair cells to ~ 42% after birth (Figure S1A). Consistently, the mRNA level of *Lphn2* in the mouse cochlear epithelium showed an approximately 30% decline from E20 to P14 and then reached a steady-state level at P30 (Figure 1A). The protein level of LPHN2 in cochlear hair cells was approximately 387 fmol/mg as measured by western blotting (using the purified LPHN2 proteins as the reference), which is approximately 1.5-fold higher than that of VLGR1 at P10 (Figure S1B-1C).

Importantly, via separation of the stereocilia from the hair cell body using the twist-off method (Avenarius et al., 2017), we were able to detect the expression of LPHN2 in both stereocilia and cell bodies of cochlear hair cells using PCDH15 and Myosin7a as specific markers for stereocilia and the hair cell body, respectively (Figure 1B and Figure S1D-1E). In contrast, LPHN2 was detected only at the cell body but not in the stereocilia of utricular hair cells, which was consistent with the immunostaining results in our parallel study (Figure 1B and Figure S1D-1E). To further explore the expression pattern of LPHN2 in mouse cochlear hair cells, we employed whole-mount immunostaining using a commercial LPHN2 antibody, which showed specificity in both HEK293 cells overexpressing LPHN2 or cochleae derived from *Lphn2*$^{-/-}$ mice at E18 (Figure S1F-1G). Different optical sections from the stereocilia to the cochlear hair cell body were examined to monitor the subcellular localization of LPHN2 (Figure 1C). Notably, in cochlear outer hair cells (OHCs) of mice at P30, LPHN2 immunofluorescence puncta were observed along the length of stereocilia, with staining at the tips of several stereocilia (Figure 1D and Figure S1H). LPHN2 immunofluorescence was also observed at the apical surfaces of OHCs, which was further supported by coimmunostaining of LPHN2 with Spectrin, a specific marker of the cuticular plate, and by staining of sections of the cochlear sensory epithelium using Myosin7a as the hair cell marker (Figure 1E-1F). In contrast, LPHN2 expression was not detected at the basolateral membrane along the cell bodies of OHCs (Figure 1F). A similar expression pattern of LPHN2 was also observed in

cochlear inner hair cells (IHCs), showing a distribution along the stereocilia and at the cuticular plate (Figure 1D-1F and Figure S1H). The specific expression pattern of mechanosensitive LPHN2 in cochlear hair cells, especially its localization at the stereocilia, suggests that LPHN2 might have unique functions in the vestibular systems.

LPHN2 deficiency severely impairs hearing

To inspect the functional roles of LPHN2 in hearing, we initially employed heterozygous *Lphn2* gene knockout (*Lphn2$^{+/-}$*) mice since *Lphn2$^{-/-}$* mice show embryonic lethality (Anderson et al., 2017) (Figure S2A-2C). The *Lphn2$^{+/-}$* mice were viable and showed normal growth but displayed vestibular defects in our parallel study. Measurement of the auditory brainstem response (ABR) revealed that *Lphn2$^{+/-}$* mice at P30 had an approximately 2-fold (~30 dB) higher auditory threshold than their wild-type (WT) littermates in response to click stimuli (Figure 2A). Moreover, compared with the WT mice, the *Lphn2$^{+/-}$* mice showed significantly elevated ABR thresholds at different frequencies (4, 8, 12, 16, 24 and 32 kHz) (Figure 2A-2B). Analysis of ABR wave I tests further indicated a lower wave amplitude and increased wave latency of *Lphn2$^{+/-}$* mice, suggesting that the function of IHCs in *Lphn2$^{+/-}$* mice was impaired (Figure 2C). Furthermore, the distortion-product otoacoustic emission (DPOAE) in *Lphn2$^{+/-}$* mice was significantly elevated between 12 and 24 kHz, implying that OHC function was affected in *Lphn2$^{+/-}$* mice (Figure 2D). Since the deterioration of hearing functions might have been due to developmental defects of hair cells, we examined the morphology of hair cells by immunostaining and scanning electron microscopy. Importantly, the results from both analyses revealed no detectable impairment in the structure or morphology of OHCs and IHCs in the cochleae of *Lphn2$^{+/-}$* mice, suggesting that the heterozygous ablation of LPHN2 in mice did not, at least at the age of 4 weeks, cause developmental defects of cochlear hair cells (Figure S2D-2F).

To further assess the specific role of LPHN2 in cochlear hair cells, we crossed *Lphn2$^{fl/fl}$* mice with an inducible *Pou4f3-CreER$^{+/-}$* transgenic mouse line (Figure S2G). Previous scRNA-seq data (GSE114157) have indicated that POU4F3, which is a commonly used hair cell marker, is expressed in all LPHN2-positive cochlear hair cells; therefore, we deemed that the use of *Pou4f3-CreER$^{+/-}$Lphn2$^{fl/fl}$* mice should enable hair cell-specific ablation of LPHN2.

Consistently, specific ablation of LPHN2 in the cochleae and utricles of *Pou4f3-CreER*[+/-]

Lphn2[fl/fl] mice was indicated by both immunofluorescent and western blotting (Figure S2H-

2J). In contrast, LPHN2 expression in other tissues of *Pou4f3-CreER*[+/-]*Lphn2*[fl/fl] mice, such as

the brain, was not significantly affected (Figure S2H-2I). Notably, compared with their

control *Pou4f3-CreER*[+/-]*Lphn2*[+/+] littermates, the *Pou4f3-CreER*[+/-]*Lphn2*[fl/fl] mice displayed

severely impaired hearing in terms of the elevated ABR threshold at all testing frequencies

and deteriorated DPOAE between 8 and 32 kHz (Figure 2E-2H). Consistent with the

observation in *Lphn2*[+/-] mice, the *Pou4f3-CreER*[+/-]*Lphn2*[fl/fl] mice at P30 also showed normal

structure and morphology of cochlear hair cells (Figure 2I-2J and Figure S2K). Collectively,

these findings support the idea that LPHN2 expressed in cochlear hair cells is essential for

hearing and suggest that LPHN2 regulates hearing functions potentially through an

instantaneous physiological process.

163   Force sensation by LPHN2 results in increased cAMP and rapid Ca^{2+} signals

GPCRs act as direct sensors of itch or in the senses of vision, smell, and taste by converting

extracellular stimuli into alterations in the intracellular levels of second messengers, such as

cAMP, cGMP and Ca^{2+} (Filipek et al., 2003; Guo et al., 2023; Kaupp, 2010; Smith, 2023;

Yang et al., 2021; Yang et al., 2023). Our parallel study indicated that LPHN2 activates Gs

and Gi signalling in response to force stimulation (Figure S3A-3B). We therefore measured

cAMP levels after delivering force to LPHN2 in cochlear explants using paramagnetic beads

coated with an anti-LPHN2 antibody (referred to as LPHN2-M-beads) that recognize the N-

terminus (residues 551-600) of LPHN2 and can exert quantifiable force on LPHN2 in a

magnetic field. Importantly, application of 3 pN of force on the cochlear sensory epithelium

derived from *Pou4f3-CreER*[+/-]*Lphn2*[+/+] mice induced an approximately 1.6-2.2-fold increase

in the intracellular cAMP levels at different sections (apex turn: from 1.49 ± 0.11 mM to 2.41

± 0.13 mM; middle turn: from 1.58 ± 0.07 mM to 3.41 ± 0.18 mM; base turn: from 1.51 ±

0.08 mM to 2.48 ± 0.14 mM), which was abrogated in the cochlear explants pretreated with

the Gs-specific inhibitor NF449 or in cochlear explants from *Pou4f3-CreER*[+/-]*Lphn2*[fl/fl] mice at

P10 (Figure 3A). In contrast, the force applied by paramagnetic beads coated with polylysine

(referred to as Ctrl-beads) did not induce detectable cAMP responses in the cochlear explants

(Figure S3C). These results suggest that the Gs-cAMP pathway downstream of LPHN2 is

activated in response to force stimulation in the cochlea. Consistent with these findings, the

Gs protein was found to be the most abundantly expressed among all the G protein subtypes

in LPHN2-expressing cochlear hair cells, as revealed by both scRNA-seq data and the single-

cell qRT-PCR analysis of 20 isolated cochlear hair cells (Figure 3B and Figure S3D).

The mechanical stimulation-triggered Ca^{2+} signal in cochlear hair cells was hypothesized to

regulate the rate of neurotransmitter release, which is essential for the transmission of sound

information. To record the force-induced Ca^{2+} response at the single-cell level, we treated P3

mice with a modified AAV-ie-*Lphn2pr*-mCherry, which enabled specific fluorescent labelling

of the LPHN2-expressing hair cells, through round window membrane injection. The

cochleae were isolated at P10, and mCherry-labelled hair cells were selected for Ca^{2+} imaging

using a Fura-2 fluorescent probe. Notably, approximately 25% (15 out of 60) of the LPHN2-

expressing cochlear hair cells showed a rapid Ca^{2+} response to LPHN2-M-bead-mediated

force stimulation, which was recorded within 140 to approximately 340 ms after force

application (Figure 3C-3E and Figure S3E-3F). In contrast, none of the unlabelled hair cells

or the mCherry-labelled hair cells derived from *Pou4f3-CreER$^{+/-}$Lphn2$^{fl/fl}$* mice showed

detectable Ca^{2+} signals in response to force application, suggesting the specificity of the

mechanical force-mediated Ca^{2+} response through activation of LPHN2 (Figure 3C-3E and

Figure S3E-3F). The force-stimulated Ca^{2+} response in LPHN2-expressing cochlear hair cells

was concentration-dependently inhibited by pretreatment with the Gs inhibitor NF-449 with

an EC50 of 412 ± 19 nM, which was similar to the inhibitory EC50 of NF-449 (272 ± 44 nM)

in a heterologous system towards force-stimulated LPHN2-Gs-cAMP signalling (Figure 3F-

3G and Figure S3G-3K). In contrast, the force-stimulated Ca^{2+} response was not affected by

pretreatment with the Gq inhibitor YM-254890 (Figure 3F and Figure S3H). Moreover, the

force application via LPHN2-M-beads showed no detectable effects on the intracellular levels

of IP3, the critical downstream messenger of the Gq pathway (Figure S3L). These results

suggested that force-stimulated activation of LPHN2 in cochlear hair cells induces rapid Ca^{2+}

signals in a Gs-dependent and Gq-independent manner, potentially due to extracellular Ca^{2+}

influx rather than mobilization from the intracellular Ca^{2+} store of IP3R. Consistent with this

hypothesis, incubation of the cochlear explants in a Ca^{2+}-deficient bath completely abolished

the force-induced Ca^{2+} response in LPHN2-expressing hair cells (Figure 3F and Figure S3I).

We therefore speculate that an ion channel activated by Gs signalling may contribute to

extracellular Ca^{2+} influx in response to LPHN2 activation after force engagement in cochlear

hair cells.

Force sensation by LPHN2 elicits acute currents in cochlear hair cells

The opening of Ca^{2+}-permeable MET channels at the stereocilia bundles of hair cells is a

key step in auditory transduction, which induces inward current and depolarizes the hair cell

membrane (Douguet and Honore, 2019; Jeong et al., 2022; Jung and Muller, 2023; Wu et al.,

2017; Zheng and Holt, 2021). The force-induced rapid Ca^{2+} influx in cochlear hair cells

through activation of LPHN2 suggests the potential participation of LPHN2 in the MET

process. We therefore examined the excitability of cochlear hair cells in response to force

stimulation of LPHN2. The whole-cell patch-clamp recording revealed that approximately 23%

(15 out of 65, cluster I) of AAV-i.e.-*Lphn2pr*-mCherry-labelled cochlear hair cells derived

from *Pou4f3-CreER$^{+/-}$Lphn2$^{+/+}$* mice showed an approximately 50% increase in current

amplitude in response to 3 pN force stimulation applied by LPHN2-M-beads, whereas the

other 77% (50 out of 65, cluster II) showed no detectable changes (Figure 4A-4B and Figure

S4A-4B). In contrast, the AAV-ie-*Lphn2pr*-mCherry-labelled hair cells in the cochlear

explants derived from *Pou4f3-CreER$^{+/-}$Lphn2$^{fl/fl}$* mice showed no significant

electrophysiological responses to force stimulation with LPHN2-M-beads (Figure 4A-4B and

Figure S4A). Nevertheless, similar to those of their WT counterparts, the cochlear hair cells

of *Pou4f3-CreER$^{+/-}$Lphn2$^{fl/fl}$* mice could also be divided into cluster I cells (10 out of 40, 25%)

and cluster II cells (30 out of 40, 75%) according to their electrophysiological characteristics

(Figure 4A-4B). In particular, the cluster I cochlear cells derived from either *Pou4f3-CreER$^{+/-}$*

Lphn2$^{+/+}$ or *Pou4f3-CreER$^{+/-}$Lphn2$^{fl/fl}$* mice featured outwardly rectifying voltage-dependent

currents, showing a higher maximum amplitude than the cluster II cells at the same holding

potential (Figure 4A-4B).

To further analyse the excitability of cochlear hair cells downstream of LPHN2 in response

to force sensation, we recorded the resting membrane potential of cluster I cells using a

current clamp. Notably, when holding at 0 pA, 3 pN force stimulation with LPHN2-M-beads

induced a significant depolarization of the WT cluster I cells by elevating the resting

membrane potential from -80 mV to approximately -65 mV (Figure 4C). As the negative

controls, the cluster I hair cells showed no significant alteration of membrane potential in

response to force stimulation with Ctrl-beads (Figure 4C).

Notably, the kinetic analysis of the force-induced membrane potential alteration revealed

that the activation time, which was defined as the time between the force application and the

rise of the membrane potential, was 94.6 ± 16.4 ms (Figure 4C-4D). In addition, after force

removal, the hair cell membrane potential quickly returned to the basal level with a delay time

of 97.0 ± 15.0 ms and a recovery time constant (τ_{rec}) of 110.4 ± 18.5 ms (Figure 4C-4D). The

very rapid activation time indicates the presence of an acute current in cochlear hair cells in

response to force stimulation applied by LPHN2-M-beads, which is consistent with the force-

induced rapid Ca^{2+} response in LPHN2-expressing hair cells. Similar to the dependency of the

rapid Ca^{2+} response on the LPHN2-Gs pathway, the LPHN2-M-bead-mediated current change

was also dose-dependently inhibited by pretreatment with the Gs inhibitor NF449, with an

inhibitory EC50 of 515 ± 84 μM (Figure 4E and Figure S4C-4D). Moreover, the force-

stimulated membrane potential alteration of cochlear hair cells was also abrogated by

treatment with NF449 (Figure 4F). Notably, the LPHN2-mediated current response was much

faster than that obtained for the LPHN2-expressing utricular hair cells in our parallel study

(94.6 ± 16.4 ms vs. 704.7 ± 9.0 ms), suggesting that LPHN2 might be part of the MET

machinery in the cochlear hair cells that directly participates in auditory perception (Figure

4G). Consistently, FM1-43 uptake by *Pou4f3-CreER^{+/-}Lphn2^{+/+}* cochlear hair cells, which is

commonly utilized to evaluate MET channel activity, was significantly increased in response

to force stimulation with LPHN2-M beads within 5 seconds (Figure 4H and Figure S4E).

Notably, the force-induced stimulatory effect on FM1-43 uptake was abolished in *Pou4f3-

CreER^{+/-}Lphn2^{fl/fl}* cochlear hair cells or by pretreatment with 10 μM NF449 in *Pou4f3-

CreER^{+/-}Lphn2^{+/+}* cochlear explants (Figure 4H and Figure S4E). Collectively, these results

support the idea that force-stimulated LPHN2 activation in cochlear hair cells promotes

membrane depolarization by eliciting acute currents in a Gs-dependent manner (Figure 4G).

Functional coupling of mechanosensitive LPHN2 to CNGA3 in cochlear hair cells

The increased intracellular cAMP levels and the Gs-dependent rapid Ca^{2+} response and

membrane excitation downstream of the mechanosensitive LPHN2 in cochlear hair cells

recall the signalling mechanism for olfactory perception, which involves cyclic nucleotide-

gated (CNG) cation channels. Suspecting that a similar mechanism might exist in cochlear

hair cells, we examined the expression patterns of CNG channels and another family of cyclic

nucleotide-regulated cation channels, hyperpolarization-activated cyclic nucleotide–

modulated (HCN) channels, in LPHN2-expressing cochlear hair cells by analysing scRNA-

seq data (GSE114157). We found that 4 out of 6 CNG channels (CNGA1, CNGA2, CNGA3

and CNGB1) and all 4 HCN channels were expressed in LPHN2-positive cochlear hair cells

(Figure S5A). Notably, 5 of these channels, including CNGA3, CNGB1 and HCN1-3, showed

a high expression frequency (37~58%) in LPHN2-expressing hair cells (Figure S5A). Further

qRT☐PCR analysis of single LPHN2-expressing hair cells revealed that CNGA3 was

specifically expressed in cluster I hair cells (Figure 5A). HCN1 also showed elevated

expression in cluster I hair cells, whereas the other channels were similarly expressed in

cluster I and II cells (Figure 5A). Consistent with our findings in the utricular hair cells in the

parallel study, the LPHN2 precipitated from the cochlear hair cells by LPHN2-M-beads

showed specific interaction with CNGA3 but not the other channel proteins (Figure 5B and

Figure S5B). Moreover, LPHN2 coimmunostained with CNGA3 both at the stereocilia and

the cuticular plates of the cochlear hair cells derived from mice at P30, suggesting the

occurrence of a direct interaction between LPHN2 and CNGA3 in vivo (Figure 5C).

To explore the potential roles of CNGA3 in hearing, we exploited the CNG inhibitor L-cis-

diltiazem and examined its effects on the auditory function of mice. Notably, intraperitoneal

injection of L-cis-diltiazem for 3 consecutive days significantly elevated the ABR threshold at

click and different testing frequencies of mice, which resembled the hearing performance of

Lphn2-deficient mice (Figure S5C-5E). Moreover, L-cis-diltiazem administration induced

deteriorated DPOAE compared with that of the mice treated with the control vehicle,

resulting in elevated thresholds at testing frequencies between 12 and 24 kHz (Figure S5F).

To inspect the specific role of CNGA3 in cochlear hair cells, we generated a *Pou4f3-CreER$^{+/-}$*

Cnga3$^{fl/fl}$ mouse line in which *Cnga3* was specifically ablated in hair cells in an inducible

manner. Compared with their *Pou4f3-CreER$^{+/-}$Cnga3$^{+/+}$* littermates, the heterozygous *Pou4f3-*

CreER$^{+/-}$Cnga3$^{fl/+}$ mice showed significantly impaired auditory performance, including

elevated ABR thresholds at all testing frequencies, increased latency and reduced amplitudes

of ABR wave I, and deteriorated DPOAE between 12 and 24 kHz (Figure 5D-5E and Figure

S5G-5H). These results indicated a critical role of the CNGA3 channel in normal hearing.

To further investigate whether CNGA3 directly participates in the mechanosensitive

LPHN2-mediated rapid auditory response, we examined the effects of L-cis-diltiazem

application on the Ca^{2+} response and acute current in LPHN2-expressing hair cells in response

to force applied by LPHN2-M-beads. Our results indicated that both the force-stimulated

rapid Ca^{2+} response and acute currents were significantly diminished in cochlear hair cells

pretreated with L-cis-diltiazem in a dose-dependent manner, with EC50 values of 36 ± 2 μM

and 31 ± 5 μM, respectively (Figure 5F-5I and Figure S5I-5J). Moreover, administration of

100 μM L-cis-diltiazem abolished the force-stimulated membrane potential alteration of

cochlear hair cells and decreased the force-stimulated FM1-43 uptake in the cochlear hair

cells by approximately 80% (Figure 5J-5K and Figure S5K). To investigate the potential

nonspecificity of L-cis-diltiazem by inhibiting other ion channels, we examined the inhibitory

potency of L-cis-diltiazem on different CNG or HCN channels in the heterologous system.

The inhibitory EC50 of L-cis-diltiazem on CNGA3 in HEK293 cells was 38 ± 5 μM, which

was similar to that recorded for L-cis-diltiazem on the force-induced current changes in the

cochlear explants (31 ± 5 μM) (Figure S6A-6B). In contrast, L-cis-diltiazem showed

significantly lower potency in inhibiting CNGA1 in the heterologous system (EC50: 70 ± 10

μM), and it did not affect the HCN activities even at the high concentration of 500 μM

(Figure S6C-6H). Therefore, these results support the hypothesis that L-cis-diltiazem

regulates the mechanosensitive LPHN2-mediated effects primarily by inhibiting the CNGA3

channel in vivo. In addition, our parallel study indicated that force application to LPHN2 in

heterologous HEK293 recombination systems was able to activate CNGA3 channels in vitro

(Figure S6I-6J). Collectively, these results suggest the existence of a functional coupling

between mechanosensitive LPHN2 and CNGA3 in cochlear hair cells that regulates the

LPHN2-mediated rapid response and potentially contributes to auditory perception.

LPHN2 re-expression in cochlear hair cells of Lphn2-deficient mice reverses hearing loss

To further explore the functional roles of LPHN2 in the peripheral auditory system,

especially in cochlear hair cells, we next performed an in vivo rescue experiment by

delivering LPHN2-encoding AAV-ie into the cochleae of *Pou4f3-CreER*$^{+/-}$*Lphn2*$^{fl/fl}$ mice and

examined its effects on the auditory functions of these *Lphn2*-deficient mice. The LPHN2-

GAIN construct with the deletion of 523 N-terminal residues, which showed

mechanosensitive potential compared to that of full-length LPHN2, was packed into the AAV-

ie-*Lphn2pr*-mCherry vector (referred to as AAV-ie-LPHN2) and delivered into P3 *Pou4f3-*

CreER$^{+/-}$*Lphn2*$^{fl/fl}$ mice through round window membrane injection (Figure S7A). The specific

expression of LPHN2-GAIN in the cochleae and utricles, but not the brains, of *Pou4f3-*

CreER$^{+/-}$*Lphn2*$^{fl/fl}$ mice was observed using western blotting after the administration of AAV-

ie-LPHN2 (Figure S7B). In particular, LPHN2 was re-expressed at the stereocilia and

cuticular plates of both cochlear OHCs and IHCs of AAV-ie-LPHN2-treated *Pou4f3-CreER*$^{+/-}$

Lphn2$^{fl/fl}$ mice (Figure S7C). In contrast, no LPHN2-GAIN expression was detected in the

cochleae of *Pou4f3-CreER*$^{+/-}$*Lphn2*$^{fl/fl}$ mice treated with the control virus AAV-ie-*Lphn2pr*-

mCherry (Figure S7C).

We next assessed the auditory functions of mice receiving LPHN2-GAIN gene delivery.

Notably, the *Lphn2*-deficient mice treated with AAV-ie-LPHN2 showed significantly

improved performance in the ABR test. The ABR thresholds at all testing frequencies, as well

as the amplitude and latency of ABR wave I, of AAV-ie-LPHN2-treated *Pou4f3-CreER*$^{+/-}$

Lphn2$^{fl/fl}$ mice returned to levels comparable to those of *Pou4f3-CreER*$^{+/-}$*Lphn2*$^{+/+}$ mice

(Figure 6A-6C and Figure S7D). Moreover, the DPOAE thresholds of the *Lphn2*-deficient

mice were also restored to nearly normal levels after AAV-ie-LPHN2 treatment (Figure 6D).

As negative controls, *Lphn2*-deficient mice treated with AAV-ie-*Lphn2pr*-mCherry showed

no significant improvements in any of the above parameters (Figure 6A-6D and Figure S7D).

The mechanosensitive potentials of the LPHN2-expressing cochlear hair cells were also

evaluated, and our results indicated that approximately 22% of the mCherry-labelled cochlear

hair cells in AAV-ie-LPHN2-treated *Pou4f3-CreER*$^{+/-}$*Lphn2*$^{fl/fl}$ mice showed significantly

increased current amplitude in response to force applied by LPHN2-M-beads, which was

comparable to the proportion of force-sensitive cluster I hair cells in the WT mice (Figure 6E-

6F and Figure S7E). In contrast, none of the cochlear hair cells labelled by the control virus

AAV-ie-*Lphn2pr*-mCherry in *Pou4f3-CreER$^{+/-}$Lphn2$^{fl/fl}$* mice showed detectable force-induced

excitability (Figure 6E-6F and Figure S7E). Collectively, these findings suggest that LPHN2

in cochlear hair cells directly contributes to auditory perception through the sensation of

mechanical force.

Discussion

Cochlear hair cells convert sound stimuli into electrochemical signals through activation

of MET channels. Despite decades of intensive efforts, the composition of the MET channels

and the detailed mechanism underlying the MET process remain elusive. Because of the very

rapid activation and ultralow response latency of cochlear hair cells in the submillisecond

range, the MET process was conventionally believed to be directly regulated by ion channels

but not by GPCRs, which usually transduce signals although second messengers or enzymatic

pathways. In the present work and our parallel study, we showed that a mechanosensitive

GPCR, LPHN2, can elicit MET in both utricle and cochlear hair cells by activating Gs-cAMP

signalling, leading to positive modulation of membrane excitability after force sensation in

both the auditory and vestibular systems. Notably, sensation of force by LPHN2 elicits a

much faster Ca^{2+} response and membrane current change in cochlear hair cells than in

utricular hair cells. These different time scales of LPHN2-regulated MET in these two

systems may be related to the distinct expression patterns of LPHN2. Whereas LPHN2 is

expressed at the cuticular plates of both utricle and cochlear hair cells, it shows a unique

distribution along the stereocilia of the cochlear hair cells but not those of utricular hair cells.

We therefore speculate that the different microenvironments within the stereocilia and the

cuticular plate, such as the different cross-sectional areas (approximately 0.1 μm^2 and 19.6

μm^2 for stereocilia and the cuticular plate, respectively) and the potential

compartmentalization with other signalling molecules, might affect the diffusion dynamics of

cAMP downstream of LPHN2 activation, contributing to different time scales of LPHN2-

mediated MET in the auditory and vestibular systems. Previous studies have suggested that

the diffusion coefficients of cAMP in olfactory cilia, neonatal cardiac myocytes and neurons

range from 136 to approximately 780 μm^2/s (Bacskai et al., 1993; Chen et al., 1999; Neves

et al., 2008; Nikolaev et al., 2004; Saucerman et al., 2006). If we assume that the cAMP

coefficient is similar in hair cells, then the theoretical time required for the transduction of the

information of LPHN2-sensed force by cAMP is approximately 12.8 μs~73.5 μs in stereocilia

and approximately 25.1~144 ms in the cuticular plate, which is partially consistent with the

faster signals of LPHN2 in cochlear hair cells than in vestibular hair cells. Interestingly, we

have noticed that both *Lphn2*[+/-] mice and *Pou4f3-CreER*[+/-]*Lphn2*[fl/fl] mice show abnormal ABR

wave I latency and amplitude compared with their respective control littermates, which

happens within 2 ms after sound stimulation and is consistent with the theoretically calculated

time latency of LPHN2-induced MET. However, both this theoretically calculated MET

latency and the latency change caused by LPHN2 deficiency during the ABR tests are

significantly shorter than our experimentally observed MET time after LPHN2 activation (~

95ms). We therefore speculate several possibilities to reconcile these discrepancies: (1) the

time latency caused by the linkage of LPHN2 antibody coated beads and LPHN2 receptor, as

the force applied on the beads may take certain time to reach the receptor because of the

flexible linker; (2) We measured MET under room temperature whereas the ABR experiments

are performed under physiological temperature; (3) Sound sensing is a complicated process.

Sound stimulation may activate multiple receptors at the same time but our magnetic beads

only activate one receptor, which caused latency difference. Or LPHN2 might only be one of

several critical components of the sound sensing complex. Importantly, application of force

on the magnetic beads coated with specific antibody against protocadherin 15 or cadherin 23,

which are central components of tip-link (Kazmierczak et al., 2007; Siemens et al., 2004;

Sotomayor et al., 2012), induces a Ca^{2+} response in cochlear hair cells with a similar time

latency (100 ~ 200 ms) to that recorded for LPHN2-M-beads (unpublished observation). The

hair bundle deflection accompanied by the movement of tip-link has been hypothesized as the

main driving force for the MET occurred at the stereocilia tips. Therefore, the comparable

time latency recorded for the magnetic beads targeting LPHN2 and tip-link proteins suggests

that the longer time latency via magnetic beads than in vivo ABR differences caused by

LPHN2 deficiency is most likely due to the experimental conditions. These results also

support the functional importance of LPHN2 comparable to the key component of the tip link

during MET process. Clearly, further experimental condition optimizations, biophysics

studies using super-resolution fluorescence imaging with highly sensitive cAMP sensors or

detailed mathematical calculations might be helpful for elucidating the mechanism underlying

the dynamics of LPHN2-mediated MET.

Our work indicates that LPHN2 promotes MET in cochlear hair cells through Gs

engagement, which elevates intracellular cAMP levels and positively regulates hair cell

excitability. The LPHN2-mediated auditory MET is more similar to the signalling mechanism

for smell perception than to that for visual perception, which relies mainly on the Gt-mediated

inhibition of constitutive cGMP levels in photoreceptor cells. Consistent with our findings,

cAMP has been proposed to modulate the sensitivity of the MET in cochlear hair cells (Mecca

et al., 2022). Notably, a loss-of-function mutation in adenylyl cyclase 1 (AC1) caused by the

deletion of the C-terminal 82 amino acids is linked to hearing loss in humans (Santos-Cortez

et al., 2014). Moreover, AC1 was expressed at the stereocilia of cochlear hair cells, supporting

our speculation that localized Gs-cAMP signalling plays important roles in auditory

perception. The *Drosophila* LPHN homologue, Ca^{2+}-independent receptor of latrotoxin (Cirl),

has been reported to respond to mechanical stimuli in vivo through interaction with the Toll-

like receptor Tollo, which modulates the sensitivity of tactile and auditory stimuli in

Drosophila (Scholz et al., 2023; Scholz et al., 2015). However, in contrast to the function of

LPHN2 in cochlear hair cells, Cirl has been found to convert mechanical stimulation into a

decrease in intracellular cAMP level in the sensory neurons of *Drosophila* (Scholz et al.,

2017). However, the downstream ion channel of Cirl that putatively transmits the cAMP

alteration into electrical signals remains unknown. Intriguingly, our parallel study also

indicated that the other two LPHN subfamily members, LPHN1 and LPHN3, could activate

Gi signalling in response to force stimulation but the heterozygous deficiency of single-

receptor resulted in no obvious hearing loss phenotype (unpublished observation). Therefore,

according to previous publications and our experimental results, mechanosensitive LPHNs

and their homologues might represent one subfamily of evolutionarily conserved auditory

sensors that act through potentially different mechanisms in mammals and lower organisms.

These differences might be the result of evolutionary adaptation to sound stimuli in more

complex environments. In addition to LPHN2, we observed that VLGR1 was able to promote

an intracellular cAMP increase in response to force sensation. Many VLGR1 mutants cause

hearing loss, so we speculate that VLGR1 could be another candidate sound sensor in

cochlear hair cells.

Ion channels located at the tips of the stereocilia, such as TMC1 and TMC2, are proposed

to be key components of the MET complex. Recent studies have also revealed the expression

of the mechanosensitive protein Piezo2 at the cuticular plates of hair cells, where Piezo2

possibly functions as a subunit of the reverse-polarity channel (Wu et al., 2017). In the present

study, we identified a novel regulatory role of the cation channel CNGA3 in the MET process

and auditory perception. Our results suggest that CNGA3 is expressed at both the stereocilia

and cuticular plates of cochlear hair cells, consistent with the localization of CNGA3 in

saccular hair cells suggested by a previous study (Selvakumar et al., 2012). Importantly, the

force sensation by LPHN2 is converted into rapid electoral signals in cochlear hair cells

through coupling to CNGA3, and pharmacological inhibition of CNGA3 activity in mice

severely impairs auditory functions. These findings not only support an essential role of the

LPHN2-CNGA3 axis in hearing but also indicate that CNGA3 is a key component of the

MET machinery. Moreover, we observed a direct association between LPHN2 and CNGA3 in

vivo, which might shorten the diffusion time of cAMP, thus partially accounting for the fast

signals of LPHN2-mediated MET. However, the detailed mechanism underlying the LPHN2-

CNGA3 association and the potential contribution of this direct interaction to LPHN2-

mediated MET require further studies, such as mutational analysis and gene knock-in mouse

model studies. In addition, we found that LPHN2 was able to directly interact with TMC1

(Figure S5B). It will be of interest to further explore the potential contribution of the LPHN2-

TMC1 interaction to the MET exerted by hair bundle deflection and to auditory perception in

vivo. Collectively, our findings identify a force-sensitive aGPCR, LPHN2, that is essential for

normal auditory perception and can sense force in cochlear hair cells and convert the force

sensation into downstream CNGA3 activation and membrane depolarization. Both LPHN2

and CNGA3 are active components during MET processes in hair cells.

Author contributions

J.-P.S. and X.Y. initiated, designed and supervised the overall project. J.-P.S. conceived the

idea of searching for MET-regulating GPCRs in the cochlear system at 2011 and initiated the

screening process by collaborating with Z.-G.X.. X.Y. and C.W. designed and supervised all

electrophysiology experiments. Z.-G.X provided guidance and suggestions on the cochlear studies. Z.Y. and J.-P.S. developed the magnetic force stimulation assay to examine the activation of G proteins and second messengers downstream of GPCRs. Z.Y., M.-W.W., S.-H.Z. and Z.-C.S. performed the cell-based assay of LPHNs. Z.Y., Z.-C.S., M.-W.W., S.-H.Z. and J.-R.G. performed the electrophysiological experiments and Ca^{2+} imaging supervised by J.-P.S. and X.Y.. J.-P.S., Z.Y., M.-W.W. and S.-H.Z. participated in the design of gene knockout mice and modified AAV-ie. Z.Y., M.-W.W., Z.-C.S., Q.-Y.Z., W.-W.L. and L.X. performed cochlear behaviour studies. Z.Y., M.-W.W., and Z.-C.S. isolated mouse cochlea and performed immunofluorescence studies and ex vivo experiments. M.-W.W. and Z.-C.S. performed single cell qRT-PCR. M.-W.W. and Y. G. performed coimmunoprecipitation and western blotting analyses. Z.Y., M.-W.W., S.-H.Z., Z.-C.S., J.-R.G. and W.-W.L. performed the AAV-ie-related experiments. J.-P.S., X.Y., Z.-G.X, C.W., Z.Y., M.-W.W., S.-H.Z., Z.-C.S. and J.-R.G. participated in data analysis and interpretation. Z.-G.X., C.W., W.-W.L. and L.X. provided insightful ideas related to the auditory sensation. Z.Y., M.-W.W., S.-H.Z., Z.-C.S. and J.-R.G. prepared the figures. J.-P.S., X.Y., Z.Y., Z.-G.X and C.W. wrote the manuscript. All the authors have seen and commented on the manuscript.

Competing interests

The authors declare no competing interests.

Submission history

The manuscript was submitted on Sep 20[th],2023 and was sent for peer review on Sep 26[th] , 2023 by the journal editor.

Figure legends

Figure 1. Expression of LPHN2 in cochlear hair cells

(A) mRNA levels of *Lphn2* in the cochlear epithelium isolated from WT mice at E20, P1, P7, P14, P30, P60, P120 and P240 (n = 6). Data are normalized to the mRNA levels of *Lphn2* at E20. E, embryonic period; P, postnatal day.

(B) Representative blotting showing the protein levels of LPHN2, Myosin7a (specific marker

of hair cell body) and PCDH15 (specific marker of the stereocilia) in the stereocilia and cell

body of cochlear or utricle hair cells isolated from WT mice at P10. Representative blotting

from three independent experiments are shown (n = 3). LPHN2 antibody against the N

terminus (551 - 600 residues) was used, which revealed both the full-length (~130 kDa) and

the N-terminal fragment of LPHN2 (~120 kDa) in the cochlear or utricle hair cells. Data are

correlated to Figure S1D-1E.

(C) Schematic diagram showing the expression pattern of LPHN2 in cochlear and utricle hair

cells, respectively.

(D) Co-immunostaining of LPHN2 (green) and Phalloidin (red) in the stereocilia of OHC (left)

and IHC (right) in cochlear whole mounts derived from WT mice at P30 (n = 6 mice per

group). Scale bar, 5 μm. White arrows show the localization of LPHN2 at the tips of

stereocilia. Data are correlated to Figure S1H.

(E) Co-immunostaining of LPHN2 (green) and Spectrin (red) at the cuticular plate of OHC

(left) and IHC (right) in cochlear whole mounts derived from WT mice at P30 (n = 6 mice per

group). Scale bar, 5 μm.

(F) Co-immunostaining of LPHN2 (green) and Myosin7a (red) in cochlear sections derived

from WT mice at P30 (n = 6 mice per group). Scale bar, 20 μm. White arrows show the

specific distribution of LPHN2 at the cuticular plate.

(A) *P < 0.05; ***P < 0.001. WT mice at P1, P7, P14, P30, P60, P120 and P240 compared

with WT mice at E20. The bars indicate mean ± SEM values. All data were statistically

analyzed using one-way ANOVA with Dunnett's post hoc test.

Figure 2. LPHN2 deficiency severely impairs hearing

(A-D) ABR thresholds (A), ABR waveforms at click (B), ABR peak1 (P1) latency (left) and

amplitude (right) (C), and DPOAE thresholds (D) of WT and *Lphn2*$^{+/-}$ mice at P30 (n = 8

mice per group). ABR, auditory brainstem response; DPOAE, distortion product otoacoustic

emission.

(E-H) ABR thresholds (E), ABR waveforms at click (F), ABR peak1 (P1) latency (left) and

amplitude (right) (G), and DPOAE thresholds (H) of $Pou4f3\text{-}CreER^{+/-}Lphn2^{+/+}$ (referred to as

$Pc\text{-}Lphn2^{+/+}$) and $Pou4f3\text{-}CreER^{+/-}Lphn2^{fl/fl}$ (referred to as $Pc\text{-}Lphn2^{fl/fl}$) mice at P30 (n = 8

mice per group).

(I) Immunostaining of Myosin7a (red) in cochlear hair cells derived from $Pc\text{-}Lphn2^{+/+}$ and

$Pc\text{-}Lphn2^{fl/fl}$ mice at P30 (n = 6 mice per group). Scale bar, 20 μm. Data are correlated to

Figure S2K.

(J) Representative SEM images showing hair bundle morphology of the OHC (left) and IHC

(right) from $Pc\text{-}Lphn2^{+/+}$ and $Pc\text{-}Lphn2^{fl/fl}$ mice at P30 (n = 6 mice per group). Scale bar, 1 μm

(OHC) or 2 μm (IHC).

(A, C and D) **$P < 0.01$; ***$P < 0.001$; ns, no significant difference. $Lphn2^{+/-}$ mice compared

with WT mice. (E, G and H) **$P < 0.01$; ***$P < 0.001$. $Pc\text{-}Lphn2^{fl/fl}$ mice compared with $Pc\text{-}$

$Lphn2^{+/+}$ mice. The bars indicate mean ± SEM values. All data were statistically analyzed

using two-way ANOVA with Dunnett's post hoc test.

Figure 3. Force sensation by LPHN2 induces cAMP and Ca^{2+} signals

(A) Endogenous cAMP level changes in the cochlea (apex, middle or base) and utricle

explant derived from $Pc\text{-}Lphn2^{+/+}$ or $Pc\text{-}Lphn2^{fl/fl}$ mice at P10 in response to force (3 pN)

applied by LPHN2-M-beads (n = 6). NF449 (10 μM), a specific Gs inhibitor.

(B) mRNA levels of different G protein subtypes in LPHN2-expressing cochlear hair cells

derived from WT mice at P10 measured by single-cell qRT-PCR (n = 20 hair cells). The

mRNA levels of respective G protein subtypes in the brain were used as the reference

(normalized to 1).

(C) Heatmaps showing the Ca^{2+} responses in LPHN2-expressing cochlear hair cells derived

from P10 $Pc\text{-}Lphn2^{+/+}$ mice labelled by AAV-ie-$Lphn2pr$-mCherry (n = 60 hair cells). The

vertical yellow full line denotes the onset of force application (0 s). Each row represents the

temporal Ca^{2+} response of one hair cell. Data are correlated to Figure 3D-3E and Figure S3E-

3F.

(D) Representative Ca^{2+} traces recorded in individual cochlear hair cells derived from P10 $Pc\text{-}$

$Lphn2^{+/+}$ mice labelled by AAV-ie-$Lphn2pr$-mCherry. Data are correlated to Figure 3C and

Figure S3E-3F.

(E) Proportion of force-sensitive and force-insensitive cochlear hair cells derived from P10

Pc-Lphn2[+/+] mice labelled by AAV-ie-*Lphn2pr*-mCherry. Data are correlated to Figure 3C-3D

and Figure S3E-3F.

(F) Effects of Gs inhibitor NF449 (10 μM), Gq inhibitor YM-254890 (60 μM) and Ca^{2+}-free

buffer on the force-induced Ca^{2+} responses in cochlear hair cells labelled by AAV-ie-*Lphn2pr*-

mCherry (n = 10 hair cells per group). Data are correlated to Figure S3G-3I.

(G) Dose-dependent inhibitory effects of NF449 treatment on force-stimulated Ca^{2+} responses

in cochlear hair cells labelled by AAV-ie-*Lphn2pr*-mCherry (n = 18 hair cells). Data are

correlated to Figure S3K.

Figure 4. Force sensation by LPHN2 elicits acute currents in cochlear hair cells

(A-B) Representative whole-cell currents (upper panel) elicited by voltage steps between -100

mV and +100 mV and the corresponding I-V curves (lower panel) in AAV-ie-*Lphn2pr*-

mCherry-labeled cluster I (A) or cluster II (B) cochlear hair cells derived from *Pc-Lphn2*[+/+] or

Pc-Lphn2[fl/fl] mice in response to force (3 pN) stimulation. N = 15 for cluster I cells and n = 50

for cluster II cells in *Pc-Lphn2*[+/+] mice. N = 10 for cluster I cells and n = 30 for cluster II cells

in *Pc-Lphn2*[fl/fl] mice. Data are correlated to Figure S4A.

(C) Representative force-induced membrane potential alteration of cluster I cochlear hair cells

derived from *Pc-Lphn2*[+/+] mice when holding at absolute zero current (n = 10 hair cells). The

time latency between force application or force removal and the change of membrane

potential was defined as T_{pr} and T_{po}, respectively. The calculated time for recovery time

constant was denoted as τ_{rec}.

(D) The calculated time parameters in force-stimulated membrane potential alteration of *Pc-*

Lphn2[+/+] cluster I cochlear hair cells (n = 10 hair cells). Data are correlated to Figure 4C.

(E) Effect of NF449 treatment on the whole-cell current in *Pc-Lphn2*[+/+] cluster I hair cells in

response to 3 pN force stimulation (n = 6 hair cells). Data are correlated to Figure S4C-4D.

(F) Effect of NF449 treatment on the force-stimulated membrane potential alteration in *Pc-*

Lphn2[+/+] cluster I hair cells (n = 6 hair cells). Data are correlated to Figure S4C-4D.

(G) Schematic diagram showing the different latency time of the force-induced current

responses through activating LPHN2 in cochlear (upper panel, ~95 ms) or utricle (lower panel,

~705 ms) hair cells.

(H) Force (3 pN)-induced FM1-43 (green) uptake in cochlea explant derived from P10 *Pc-*

Lphn2$^{+/+}$ or *Pc-Lphn2$^{fl/fl}$* mice within 5 sec at the presence or absence of NF449 (n = 6 per

group). Data are correlated to Figure S4E.

Figure 5. LPHN2 regulates MET in cochlear hair cells through coupling to CNGA3 cells

(A) mRNA levels of CNG and HCN channels in WT cluster I and cluster II cochlear hair cells

examined by single-cell qRT-PCR (n = 10 cells per cluster). The mRNA levels of these ion

channels in the brain were used as the reference (normalized to 1).

(B) Representative blotting showing the co-immunoprecipitation of LPHN2 with CNGA3

channel in the lysates of mouse cochlea. Representative blotting form three independent

experiments are shown (n = 3). Data are correlated to Figure S5B.

(C) Co-immunostaining of LPHN2 (green), CNGA3 (magenta) and phalloidin (red) in

cochlea sections derived from WT mice at P30 (n = 6 mice per group). Scale bar: 5 μm. White

arrows show co-localization of LPHN2 and CNGA3 at the stereocilia and cuticular plate.

(D-E) ABR thresholds (D) and DPOAE thresholds (E) of *Pou4f3-CreER$^{+/-}$Cnga3$^{+/+}$* (referred

to as *Pc-Cnga3$^{+/+}$*) and *Pou4f3-CreER$^{+/-}$Cnga3$^{fl/+}$* (referred to *Pc-Cnga3$^{fl/+}$*) mice at P30 (n =

8 mice per group). Data are correlated to Figure S5G-5H.

(F-G) Representative traces (F) and the dose-response curve (G) showing the effect of CNG

inhibitor L-cis-diltiazem on the force (3 pN)-induced Ca^{2+} responses in individual cochlear

hair cell derived from P10 *Pc-Cnga3$^{+/+}$* cluster I hair cells (n = 18 hair cells). Data are

correlated to Figure S5I.

(H-I) Representative I-V curves (H) and the dose-response curve (I) showing the effect of L-

cis-diltiazem on the force (3 pN)-induced the whole-cell current in individual cochlear hair

cell derived from P10 *Pc-Cnga3$^{+/+}$* cluster I hair cells (n = 6 hair cells). Data are correlated to

Figure S5J.

(J-K) Representative fluorescence images (J) and quantification analysis (K) of the effect of

L-cis-diltiazem treatment on the force (3 pN)-induced FM1-43 uptake in cochlea explant

derived from P10 *Pc-Cnga3$^{+/+}$* mice within 5 sec (n = 6 mice per group).

(D-E) ***P < 0.001; ns, no significant difference. *Pc-Cnga3*$^{fl/+}$ mice compared with *Pc-*

Cnga3$^{+/+}$ mice at P30. (K) **P < 0.01; ***P < 0.001. Cochlea explants treated with force

compared with the control. $^{###}$P < 0.001. Cochlea explants treated with L-cis-diltiazem

compared with those treated with control vehicle. The bars indicate mean ± SEM values. All

data were statistically analyzed using two-way ANOVA with Dunnett's post hoc test.

Figure 6. LPHN2 re-expression in cochlear hair cells of *Lphn2*-deficient mice rescues

hearing loss

(A-D) ABR thresholds (A), ABR peak1 (P1) latency (B) and amplitude (C), and DPOAE

thresholds (D) of *Pc-Lphn2*$^{+/+}$, *Pc-Lphn2*$^{fl/fl}$, AAV-ie-*Lphn2pr*-LPHN2-mCherry-treated *Pc-*

Lphn2$^{fl/fl}$ mice (referred to as AAV-ie-LPHN2 mice) and AAV-ie-*Lphn2pr*-mCherry-treated

Pc-Lphn2$^{fl/fl}$ mice (referred to as AAV-ie-*Lphn2pr* mice) at P30 (n = 6 mice per group). Data

are correlated to Figure S7D.

(E) Representative whole-cell currents (left) elicited by voltage steps between -100 mV and

+100 mV and the corresponding I-V curves (right) in cluster I cochlear hair cells derived from

Pc-Lphn2$^{+/+}$, AAV-ie-LPHN2 and AAV-ie-*Lphn2pr* mice in response to force (3 pN)

stimulation (n = 5 ~ 6 hair cells per group). Data are correlated to Figure S7E.

(F) Quantification of force-sensitive and force-insensitive cochlear hair cells derived from *Pc-*

Lphn2$^{+/+}$, AAV-ie-LPHN2 and AAV-ie-*Lphn2pr* mice. Data are correlated to Figure 6E.

(A-D) **P < 0.01; ***P < 0.001; ns, no significant difference. *Pc-Lphn2*$^{fl/fl}$, AAV-ie-LPHN2

or AAV-ie-*Lphn2pr* mice compared with *Pc-Lphn2*$^{+/+}$ mice. The bars indicate mean ± SEM

values. All data were statistically analyzed using one-way ANOVA with Dunnett's post hoc

test.

References

Anderson, G.R., Maxeiner, S., Sando, R., Tsetsenis, T., Malenka, R.C., and Sudhof, T.C.

(2017). Postsynaptic adhesion GPCR latrophilin-2 mediates target recognition in entorhinal-

hippocampal synapse assembly. J Cell Biol *216*, 3831-3846.

Avenarius, M.R., Krey, J.F., Dumont, R.A., Morgan, C.P., Benson, C.B., Vijayakumar, S.,

Cunningham, C.L., Scheffer, D.I., Corey, D.P., Muller, U., *et al.* (2017). Heterodimeric capping

protein is required for stereocilia length and width regulation. J Cell Biol *216*, 3861-3881.

Bacskai, B.J., Hochner, B., Mahaut-Smith, M., Adams, S.R., Kaang, B.K., Kandel, E.R., and

Tsien, R.Y. (1993). Spatially resolved dynamics of cAMP and protein kinase A subunits in

Aplysia sensory neurons. Science *260*, 222-226.

Buck, L., and Axel, R. (1991). A novel multigene family may encode odorant receptors: a

molecular basis for odor recognition. Cell *65*, 175-187.

Chen, C., Nakamura, T., and Koutalos, Y. (1999). Cyclic AMP diffusion coefficient in frog

olfactory cilia. Biophys J *76*, 2861-2867.

Cunningham, C.L., Qiu, X., Wu, Z., Zhao, B., Peng, G., Kim, Y.H., Lauer, A., and Muller, U.

(2020). TMIE Defines Pore and Gating Properties of the Mechanotransduction Channel of

Mammalian Cochlear Hair Cells. Neuron *107*, 126-143 e128.

Douguet, D., and Honore, E. (2019). Mammalian Mechanoelectrical Transduction: Structure

and Function of Force-Gated Ion Channels. Cell *179*, 340-354.

Duncan, J.S., Stoller, M.L., Francl, A.F., Tissir, F., Devenport, D., and Deans, M.R. (2017).

Celsr1 coordinates the planar polarity of vestibular hair cells during inner ear development.

Dev Biol *423*, 126-137.

Filipek, S., Stenkamp, R.E., Teller, D.C., and Palczewski, K. (2003). G protein-coupled

receptor rhodopsin: a prospectus. Annu Rev Physiol *65*, 851-879.

Guan, Y., Du, H.B., Yang, Z., Wang, Y.Z., Ren, R., Liu, W.W., Zhang, C., Zhang, J.H., An,

W.T., Li, N.N., *et al.* (2023). Deafness-Associated ADGRV1 Mutation Impairs USH2A Stability

through Improper Phosphorylation of WHRN and WDSUB1 Recruitment. Adv Sci (Weinh) *10*,

e2205993.

Guo, L., Cheng, J., Lian, S., Liu, Q., Lu, Y., Zheng, Y., Zhu, K., Zhang, M., Kong, Y., Zhang,

C., *et al.* (2023). Structural basis of amine odorant perception by a mammal olfactory receptor.

Nature *618*, 193-200.

Jeong, H., Clark, S., Goehring, A., Dehghani-Ghahnaviyeh, S., Rasouli, A., Tajkhorshid, E.,

and Gouaux, E. (2022). Structures of the TMC-1 complex illuminate mechanosensory

transduction. Nature *610*, 796-803.

Jia, Y., Zhao, Y., Kusakizako, T., Wang, Y., Pan, C., Zhang, Y., Nureki, O., Hattori, M., and

Yan, Z. (2020). TMC1 and TMC2 Proteins Are Pore-Forming Subunits of Mechanosensitive

Ion Channels. Neuron *105*, 310-321 e313.

Jung, J., and Muller, U. (2023). Mechanoelectrical transduction-related genetic forms of

hearing loss. Curr Opin Physiol *32*.

Kaupp, U.B. (2010). Olfactory signalling in vertebrates and insects: differences and

commonalities. Nat Rev Neurosci *11*, 188-200.

Kazmierczak, P., Sakaguchi, H., Tokita, J., Wilson-Kubalek, E.M., Milligan, R.A., Muller, U.,

and Kachar, B. (2007). Cadherin 23 and protocadherin 15 interact to form tip-link filaments in

sensory hair cells. Nature *449*, 87-91.

Liang, X., Qiu, X., Dionne, G., Cunningham, C.L., Pucak, M.L., Peng, G., Kim, Y.H., Lauer, A.,

Shapiro, L., and Muller, U. (2021). CIB2 and CIB3 are auxiliary subunits of the

mechanotransduction channel of hair cells. Neuron *109*, 2131-2149 e2115.

Mecca, A.A., Caprara, G.A., and Peng, A.W. (2022). cAMP and voltage modulate rat auditory

mechanotransduction by decreasing the stiffness of gating springs. Proc Natl Acad Sci U S A

119, e2107567119.

Neves, S.R., Tsokas, P., Sarkar, A., Grace, E.A., Rangamani, P., Taubenfeld, S.M., Alberini,

C.M., Schaff, J.C., Blitzer, R.D., Moraru, II*, et al.* (2008). Cell shape and negative links in

regulatory motifs together control spatial information flow in signaling networks. Cell *133*, 666-

680.

Nikolaev, V.O., Bunemann, M., Hein, L., Hannawacker, A., and Lohse, M.J. (2004). Novel

single chain cAMP sensors for receptor-induced signal propagation. J Biol Chem *279*, 37215-

37218.

Pan, B., Akyuz, N., Liu, X.P., Asai, Y., Nist-Lund, C., Kurima, K., Derfler, B.H., Gyorgy, B.,

Limapichat, W., Walujkar, S.*, et al.* (2018). TMC1 Forms the Pore of Mechanosensory

Transduction Channels in Vertebrate Inner Ear Hair Cells. Neuron *99*, 736-753 e736.

Pan, B., Geleoc, G.S., Asai, Y., Horwitz, G.C., Kurima, K., Ishikawa, K., Kawashima, Y.,

Griffith, A.J., and Holt, J.R. (2013). TMC1 and TMC2 are components of the

mechanotransduction channel in hair cells of the mammalian inner ear. Neuron *79*, 504-515.

Ranum, P.T., Goodwin, A.T., Yoshimura, H., Kolbe, D.L., Walls, W.D., Koh, J.Y., He, D.Z.Z.,

and Smith, R.J.H. (2019). Insights into the Biology of Hearing and Deafness Revealed by

Single-Cell RNA Sequencing. Cell Rep *26*, 3160-3171 e3163.

Santos-Cortez, R.L., Lee, K., Giese, A.P., Ansar, M., Amin-Ud-Din, M., Rehn, K., Wang, X.,

Aziz, A., Chiu, I., Hussain Ali, R.*, et al.* (2014). Adenylate cyclase 1 (ADCY1) mutations cause

recessive hearing impairment in humans and defects in hair cell function and hearing in

zebrafish. Hum Mol Genet *23*, 3289-3298.

Saucerman, J.J., Zhang, J., Martin, J.C., Peng, L.X., Stenbit, A.E., Tsien, R.Y., and

McCulloch, A.D. (2006). Systems analysis of PKA-mediated phosphorylation gradients in live

cardiac myocytes. Proc Natl Acad Sci U S A *103*, 12923-12928.

Scholz, N., Dahse, A.K., Kemkemer, M., Bormann, A., Auger, G.M., Vieira Contreras, F.,

Ernst, L.F., Staake, H., Korner, M.B., Buhlan, M.*, et al.* (2023). Molecular sensing of

mechano- and ligand-dependent adhesion GPCR dissociation. Nature *615*, 945-953.

Scholz, N., Gehring, J., Guan, C., Ljaschenko, D., Fischer, R., Lakshmanan, V., Kittel, R.J.,

and Langenhan, T. (2015). The adhesion GPCR latrophilin/CIRL shapes mechanosensation.

Cell Rep *11*, 866-874.

Scholz, N., Guan, C., Nieberler, M., Grotemeyer, A., Maiellaro, I., Gao, S., Beck, S., Pawlak,

M., Sauer, M., Asan, E.*, et al.* (2017). Mechano-dependent signaling by Latrophilin/CIRL

quenches cAMP in proprioceptive neurons. Elife *6*.

Selvakumar, D., Drescher, M.J., Dowdall, J.R., Khan, K.M., Hatfield, J.S., Ramakrishnan,

N.A., and Drescher, D.G. (2012). CNGA3 is expressed in inner ear hair cells and binds to an

intracellular C-terminus domain of EMILIN1. Biochem J *443*, 463-476.

Shi, F., Hu, L., and Edge, A.S. (2013). Generation of hair cells in neonatal mice by beta-

catenin overexpression in Lgr5-positive cochlear progenitors. Proc Natl Acad Sci U S A *110*,

13851-13856.

Siemens, J., Lillo, C., Dumont, R.A., Reynolds, A., Williams, D.S., Gillespie, P.G., and Muller,

U. (2004). Cadherin 23 is a component of the tip link in hair-cell stereocilia. Nature *428*, 950-

955.

Smith, S.O. (2023). Mechanism of Activation of the Visual Receptor Rhodopsin. Annu Rev

Biophys *52*, 301-317.

Sotomayor, M., Weihofen, W.A., Gaudet, R., and Corey, D.P. (2012). Structure of a force-

conveying cadherin bond essential for inner-ear mechanotransduction. Nature *492*, 128-132.

Tan, F., Chu, C., Qi, J., Li, W., You, D., Li, K., Chen, X., Zhao, W., Cheng, C., Liu, X.*, et al.*

(2019). AAV-ie enables safe and efficient gene transfer to inner ear cells. Nat Commun *10*,

3733.

Tang, Y.Q., Lee, S.A., Rahman, M., Vanapalli, S.A., Lu, H., and Schafer, W.R. (2020).

Ankyrin Is An Intracellular Tether for TMC Mechanotransduction Channels. Neuron *107*, 112-

125 e110.

Wang, Y., Guo, N., and Nathans, J. (2006). The role of Frizzled3 and Frizzled6 in neural tube

closure and in the planar polarity of inner-ear sensory hair cells. J Neurosci *26*, 2147-2156.

Wu, Z., Grillet, N., Zhao, B., Cunningham, C., Harkins-Perry, S., Coste, B., Ranade, S.,

Zebarjadi, N., Beurg, M., Fettiplace, R.*, et al.* (2017). Mechanosensory hair cells express two

molecularly distinct mechanotransduction channels. Nat Neurosci *20*, 24-33.

Xiong, W., Grillet, N., Elledge, H.M., Wagner, T.F., Zhao, B., Johnson, K.R., Kazmierczak, P.,

and Muller, U. (2012). TMHS is an integral component of the mechanotransduction machinery

of cochlear hair cells. Cell *151*, 1283-1295.

Yang, F., Guo, L., Li, Y., Wang, G., Wang, J., Zhang, C., Fang, G.X., Chen, X., Liu, L., Yan,

X.*, et al.* (2021). Structure, function and pharmacology of human itch receptor complexes.

Nature *600*, 164-169.

Yang, Z., Cheng, J., Shang, P., Sun, J.P., and Yu, X. (2023). Emerging roles of olfactory

receptors in glucose metabolism. Trends Cell Biol *33*, 463-476.

Yau, K.W., and Hardie, R.C. (2009). Phototransduction motifs and variations. Cell *139*, 246-

264.

Zhang, Y., Hoon, M.A., Chandrashekar, J., Mueller, K.L., Cook, B., Wu, D., Zuker, C.S., and

Ryba, N.J. (2003). Coding of sweet, bitter, and umami tastes: different receptor cells sharing

similar signaling pathways. Cell *112*, 293-301.

Zheng, W., and Holt, J.R. (2021). The Mechanosensory Transduction Machinery in Inner Ear

Hair Cells. Annu Rev Biophys *50*, 31-51.

Figure 1

A Cochlea (n=6)
Lphn2 mRNA level — E20, P1, P7, P14, P30, P60, P120, P240 — Cochlear epithelium

B (P10 / Hair cells)
Cochlea / Utricle
LPHN2, Myosin7a, PCDH15
Stereocilia / Cell body

C Utricle / Cochlea
LPHN2 — Stereocilia, Cuticular plate, Cell body

D P30 / OHC (LPHN2 / Phalloidin); P30 / IHC (LPHN2 / Phalloidin)

E P30 (LPHN2 / Spectrin) — OHC, IHC

F P30 (LPHN2 / Myosin7a)

Figure 2

Figure 3

A P10 / cAMP level (mM / n=6)

Pc-Lphn2+/+ — Force, Ctrl
NF449 — Force Ctrl
Pc-Lphn2fl/fl — Force Ctrl

Utricle | Apex Middle Base
Cochlea

B P10 / mRNA level (n=20)

Gnas, Gnaolf, Gnai1, Gnai2, Gnai3, Gnao, Gnat1, Gnat2, Gnag, Gnaz, Gnaq, Gna11, Gna12, Gna13, Gna14, Gna15

C P10 / Pc-Lphn2+/+ (n=60)
Delay time 140 ~ 340 ms

Z-score

Hair cell number
Time (s)

F P10 / Pc-Lphn2+/+ (n=10)

Z-score 20
25s
Time (s)

Force

NF449
YM-254890
Ca2+-free buffer

D mCherry+/sensitive (n=15)
mCherry+/insensitive (n=45)

P10 / Pc-Lphn2+/+ (n=60)

Z-score
Force
Time (s)

E P10 / Pc-Lphn2+/+
Force sensitive
Force insensitive

25% 75%

mCherry+
Total = 60 cells

G

ΔZ-score
(n=6)
P10 / Pc-Lphn2+/+
$EC_{50} = 412 \pm 19$ nM
Log [NF449] (M)

Figure 4

Figure 5

A — mRNA level (n=10)

Cluster 1 Brain / Cluster 2

Cnga1 *Cnga2* *Cnga3* *Cnga4* *Cngb1* *Cngb3* *Hcn1* *Hcn2* *Hcn3* *Hcn4*

B

LPHN2 / CNGA3

Output / Heavy chain / Input

180 130 100 70 55 40

IgG LPHN2 / IgG LPHN2

IP with

C — P30 (LPHN2 / CNGA3 / Phalloidin)

Stereocilia / Cuticular plate

5 µm

LPHN2 CNGA3 Merge

D

ABR threshold (dB)

Pc-Cnga3$^{+/+}$
Pc-Cnga3$^{fl/+}$

P30 (n=8)

*** *** *** *** *** *** ***

Click 4 8 12 16 24 32

Frequency (kHz)

E

DPOAE threshold (dB)

Pc-Cnga3$^{+/+}$
Pc-Cnga3$^{fl/+}$

P30 (n=8)

ns *** *** *** ns

8 12 16 24 32

Frequency (kHz)

F

Force

Control

L-cis-diltiazem

Z-score

20 / 25s

Time (s)

G

$EC_{50} = 36 \pm 2$ µM

Δ Z-score

P10 (n=6)

Log [L-cis-diltiazem] (µM)

H

I/Imax

Control
Force
Force-1 µM
Force-3 µM
Force-30 µM
Force-300 µM

Dilt

Voltage (mV)

I

$EC_{50} = 31 \pm 5$ µM

Δ I/Imax

P10 (n=6)

Log [L-cis-diltiazem] (µM)

J — FM1-43 (P10, 5 s)

Control / L-cis-diltiazem (100 µM)

Control

Force

OHC IHC

10 µm

K

Relative fluorescence intensity of FM1-43

Control
Force

P10 (n=6)

*** ### **

Control L-cis-diltiazem

Figure 6