Force sensation by a Gs-coupled adhesion GPCR mediates equilibrioception by increasing vestibular hair cell membrane excitability and CNGA3 coupling

Zhao Yang[1,4#], Shu-Hua Zhou[1,2,3#], Qi-Yue Zhang[1#], Wen-Wen Liu[5#], Yu Sun[6#], Jie-Yu Qi[2#], Ying Guan[1], Ming-Wei Wang[4,7], Zhi-Chen Song[3], Yu-Nan Sun[8], Wei Yang[9], Xiao Yu[3*], Ren-Jie Chai[2,10*], Jin-Peng Sun[1,4,7,8*]

Affiliations:

[1]Key Laboratory Experimental Teratology of the Ministry of Education, Department of Biochemistry and Molecular Biology, School of Basic Medical Sciences, Cheeloo College of Medicine, Shandong University, Jinan 250012, China.

[2]State Key Laboratory of Digital Medical Engineering, Department of Otolaryngology Head and Neck Surgery, Zhongda Hospital, School of Life Sciences and Technology, Advanced Institute for Life and Health Jiangsu Province High-Tech Key Laboratory for Bio-Medical Research, Southeast University, Nanjing 210096, China.

[3]Key Laboratory of Experimental Teratology of the Ministry of Education, Department of Physiology, School of Basic Medical Sciences, Cheeloo College of Medicine, Shandong University, Jinan 250012, China.

[4]Advanced Medical Research Institute, Cheeloo College of Medicine, Shandong University, Jinan 250012, China.

[5]Department of Otolaryngology-Head and Neck Surgery, Shandong Provincial ENT Hospital, Cheeloo College of Medicine, Shandong University, Jinan, 250014, China.

[6]Department of Otorhinolaryngology, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China.

[7]Department of Clinical Laboratory, The Second Hospital, Cheeloo College of Medicine, Shandong University, Jinan 250012, China.

[8]Key Laboratory of Molecular Cardiovascular Science of the Ministry of Education, Department of Physiology and Pathophysiology, School of Basic Medical Sciences, Peking University, Beijing 100191, China.

[9]Department of Biophysics, and Department of Neurology of the Fourth Affiliated Hospital,

31   Zhejiang University School of Medicine, Hangzhou 310058, China.

32   [10]Co-Innovation Center of Neuroregeneration, Nantong University, Nantong 226001, China.

33   # These authors contributed equally to this work.

34   * Correspondence: Jin-Peng Sun, E-mail: sunjinpeng@sdu.edu.cn

35   Ren-Jie Chai, E-mail: renjiec@seu.edu.cn

36   Xiao Yu, E-mail: yuxiao@sdu.edu.cn

Abstract

Equilibrioception is essential for the perception and navigation of mammals in the three-dimensional world. The mechanoelectrical transduction (MET) process in vestibular hair cells is believed to play central roles in positional and motional sensation. However, the direct equilibrioception sensor at the molecular level is not clear, and the mechanism underlying the MET process in vestibular hair cells remains elusive. Here, we identified that the G protein-coupled receptor LPHN2, which is expressed in the cuticular plate of utricular hair cells, is required for normal balance maintenance. The sensation of force by LPHN2 in the utricular hair cells increased intracellular cAMP levels in a Gs-dependent manner. The direct association of LPHN2 with the ion channel CNGA3 and increased cAMP levels enabled the activation of CNGA3, which induced calcium influx and membrane depolarization. These cellular signals and balance-related functions were significantly weakened in $Lphn2^{+/-}$ mice, $Pou4f3\text{-}CreER^{+/-}$ $Lphn2^{fl/fl}$ mice and $Pou4f3\text{-}CreER^{+/-}Cnga3^{fl/+}$ mice and could be suppressed by Gs or CNGA3 inhibitors. Collectively, our data suggested that LPHN2 could be a direct equilibrioception sensor in utricular hair cells by converting the force sensation to electrical signals through Gs and CNGA3 coupling. Both GPCR members and the ion channel CNGA3 could be essential elements for equilibrioception.

Introduction

The sense of balance and motion enables our perception and navigation in the three-dimensional world and is thus critical to our interactions with the world. The positional or motional information is primarily perceived by the hair cells located in the sensory epithelium of the vestibular end organs, which include two perpendicularly arranged otolith organs, the

utricle and saccule, and three semicircular canals[1,2]. Vestibular hair cells transmit balance

information about head motion or tilt through a mechanoelectrical transduction (MET) process

that converts mechanical force into electrical signals. The MET process is thought to be mainly

regulated by mechanotransduction ion channels, with the mechanism remaining largely

elusive[3,4]. In addition to ion channels, recent studies have also uncovered a regulatory role of G

protein-coupled receptors (GPCRs) in the vestibular system[5,6]. For example, CELSR1 regulates

vestibular functions by coordinating the planar polarity of hair cells, and CELSR1 deficiency

in mice leads to balance dysfunction[7]. However, it is still unknown whether GPCRs directly

regulate the mechanotransduction in balance sensation.

In our parallel study described in a companion paper, which was started three years earlier

than the current manuscript, we identified 5 receptors, including GPR133, GPR126, LPHN2,

LPHN3 and VLGR1, as mechanosensitive GPCRs in the utricle by screening the

mechanosensitive potential of GPCRs in utricular hair cells using a combination of cellular

assays and an animal model of balance dysfunction. Importantly, we identified that GPR133

converts mechanical stimuli into intracellular cAMP signals in utricular hair cells through Gi

engagement, which suppress MET and inhibits plasma membrane excitability. This work

suggests a potential of GPCR-Gi signalling-modulated inhibition of membrane excitability in

equilibrioception, which appears similar to the events that occur during vision sensation[8].

However, whether a positive modulation of vestibular membrane excitability by GPCR-

mediated MET, a process similar to those in odour and taste sensation[9,10], is involved in the

activation of Gs signalling remains unclear (Figure 1A).

Notably, of the 5 mechanosensitive GPCRs in the utricle, LPHN2 and VLGR1 activate Gs

signalling in response to force stimulation. Therefore, in the present study, we further explored

the potential role of Gs signalling of mechanosensitive GPCRs in equilibrioception using

LPHN2 as the representative receptor since VLGR1 knockout did not result in significant

balance phenotypes. We found that LPHN2 specifically expressed in the cuticular plate of

utricular hair cells is indispensable for equilibrioception. Global knockout or hair cell-specific

knockout of *Lphn2* in mice impairs balance. Mechanistically, force sensation by LPHN2

promotes an increase in the cAMP levels of utricular hair cells, which activates CNGA3

channels and induces membrane depolarization. Consistently, inhibiting CNGA3 activity by

pharmacological or genetic approaches eliminates LPHN2-mediated MET, leading to balance

dysfunction. Therefore, our study identifies an equilibrioception-regulating GPCR that

participates in MET by activating Gs-cAMP signalling.

Results

Mechanosensitive LPHN2 is essential for normal vestibular functions

Quantitative RT–PCR (qPCR) analysis of LPHN expression from mouse utricles harvested

at different time points revealed that all three LPHN subfamily members (LPHN1-3) showed

constant and stable expression from late embryo (E15) to adulthood (P120) (Figure S1A). Both

single-cell RNA sequencing (scRNA-seq) data and the results from our single-hair-cell qRT–

PCR analysis indicated that all three LPHNs were expressed in the utricular hair cells of mice,

ranging from 57% to 95% for expression frequency[11] (Figures 1B and S1B). Compared with

that of LPHN1, the average expression levels of LPHN2 and LPHN3 were relatively higher,

approximately 6- and 3-fold, respectively (Figures 1C and S1C). We next examined the

mechanosensitive property of LPHNs using a magnetic force stimulation system in which

magnetic force was applied to paramagnetic beads coated with anti-Flag antibody (referred to

as Flag-M-beads) and interacted with N-terminal Flag-tagged LPHNs expressed in HEK293

cells (Figure S1D). The force-induced activation of LPHNs was determined by inspecting the

downstream dissociation of G protein heterotrimers as revealed by a bioluminescence

resonance energy transfer (BRET) assay. All three LPHNs showed significant Gi activity in

response to 3 pN force stimulation (Figure 1D). Notably, only LPHN2, but not LPHN1 or

LPHN3, displayed detectable force-stimulated Gs activity (Figures 1E and S1E). The higher

expression levels of LPHN2 and LPHN3 in the murine utricular hair cells compared with that

of LPHN1 and their mechanosensitive capability collectively prompted us to explore the

potential vestibular functions of these two receptors.

Initially, heterozygous *Lphn2* and *Lphn3* gene knockout KO mice were used for

equilibration-related phenotype analysis since constitutive KO of *Lphn2* or *Lphn3* caused

embryonic lethality and developmental defects, respectively[12,13]. The deficiency of one copy of

LPHN2 or LPHN3 in the heterozygous KO mice was verified by genotyping PCR and western

blotting analysis (Figures S2A-S2H). These mice were viable and had body weights

comparable to those of their WT littermates (Figure S2I). The vestibular behaviours of the mice at P40 were assessed through the rotarod test and forced swimming test. Despite their nearly normal performance in the rotarod test ($Lphn2^{+/-}$ 109.5 ± 6.43 s vs. WT 106.2 ± 5.36 s), the $Lphn2^{+/-}$ mice showed impaired performance in the open field test, displaying significantly elevated circulating and travelling activities in both 2 min and 10 min time frames (Figures 1F-1H). Moreover, compared with their WT littermates, the $Lphn2^{+/-}$ mice showed impaired swimming ability, with a swimming score of 0.88± 0.13 (the 0–3 scoring system was used, and WT mice had scores of 0) (Figure 1I). In contrast, compared with the WT mice, the $Lphn3^{+/-}$ mice showed no significant difference in the open field test, rotarod test or forced swimming tests (Figures 1F-1I).

To further investigate the vestibular functions of these heterozygous LPHN-deficient mice, we measured the vestibular-ocular reflex (VOR) responses of the mice to sinusoidal head rotations at P40. Notably, compared with the WT mice, the $Lphn2^{+/-}$ mice showed an approximately 25%-50% decrease in compensatory VOR gains in response to either earth-vertical or off-vertical axis rotation, suggesting a deficiency in both the semicircular canals and otolith system (Figures 1J-1M). In contrast, the $Lphn3^{+/-}$ mice showed normal eye movement responses in both VOR tests (Figures 1J-M). Collectively, these results indicated that LPHN2 but not LPHN3 plays an important role in balance maintenance.

Expression pattern of LPHN2 in utricular hair cells

We next examined the expression patterns of LPHN2 in the mouse utricle by whole-mount immunostaining, which showed specific expression in approximately 80% of myo7a-positive hair cells (Figures 2A and S3A). The specificity of the LPHN2 antibody was supported by both western blotting results in HEK293 cells expressing different LPHNs and immunostaining of utricles derived from $Lphn2^{-/-}$ mice at E18 ($Lphn2^{-/-}$ mice showed embryonic lethality) (Figures 2A and S3B). The expression pattern of LPHN2 was further supported by RNAscope in situ hybridization, which revealed a comparable LPHN2-expressing hair cell proportion (~80%) to that inspected by immunostaining (Figures 2B and 2C). To further determine the subcellular localization of LPHN2 in utricular hair cells, we examined the expression pattern of LPHN2 in different optical sections spanning from the stereocilia to the hair cell body. LPHN2 expression

was detected only at the apical surface of the hair cells, as revealed by its coimmunostaining with spectrin, which is a specific marker of the cuticular plate in hair cells[14,15] (Figures 2D and S3C). In contrast, LPHN2 expression was not detected within the stereocilia or along the basolateral membrane of hair cells (Figure 2D). The specific expression of LPHN2 at the cuticular plate was further supported by section staining of the utricle sensory epithelium using MYO7A and SOX2 as the hair cell and supporting cell markers, respectively (Figure 2E). The unique expression pattern of mechanosensitive LPHN2 in the utricular hair cells suggests that it might participate in the interactions between the hair cells and the otoconial matrix, which play important roles in force sensation.

LPHN2 in vestibular hair cells specifically regulates balance sensation

To assess the specific functional role of LPHN2 in vestibular hair cells, we crossed $Lphn2^{fl/fl}$ mice with inducible $Pou4f3\text{-}CreER^{+/-}$ transgenic mice (Figure 3A). Pou4f3 is the transcriptional target of ATOH1 and a commonly used marker of hair cells, which is expressed in all LPHN2-positive utricular hair cells, as revealed by scRNA-seq data[11,16] (Figures S3D and S3E). The specificity of LPHN2 ablation in the vestibule of $Pou4f3\text{-}CreER^{+/-}Lphn2^{fl/fl}$ mice was indicated by the loss of LPHN2 immunostaining in Pou4f3-expressing utricular hair cells but not in other tissues, such as the brainstem (Figures 3B and S3F). The specific elimination of Lphn2 expression in the vestibule of $Pou4f3\text{-}CreER^{+/-}Lphn2^{fl/fl}$ mice was further supported by western blotting analysis (Figure S3G).

A battery of behavioural tests was conducted to assess the effects of LPHN2 ablation in vestibular hair cells on balance maintenance. Notably, in addition to their significantly declined swimming performance, the $Pou4f3\text{-}CreER^{+/-}Lphn2^{fl/fl}$ mice showed approximately 50% shorter durations in the rotarod test than their $Pou4f3\text{-}CreER^{+/-}Lphn2^{+/+}$ littermates, which was not observed in the $Lphn2^{+/-}$ mice (Figures 3C and 3D). Moreover, compared with their $Pou4f3\text{-}CreER^{+/-}Lphn2^{+/+}$ littermates, the $Pou4f3\text{-}CreER^{+/-}Lphn2^{fl/fl}$ mice exhibited an approximately 2-fold increase in circling times in both the 2 min and 10 min testing time frames, accompanied by significantly increased travelling distances (Figures 3E,3F and S3H). Furthermore, compared with littermate control mice, $Pou4f3\text{-}CreER^{+/-}Lphn2^{fl/fl}$ mice showed an approximately 25%~50% decrease in the VOR response in either the earth-vertical or off-

vertical axis rotation test (Figures 3G and 3H). Collectively, these results suggest a balance-

regulatory role of LPHN2 in vestibular hair cells.

Force sensing by LPHN2 increases the cAMP levels of utricular hair cells

Downstream of GPCR activation, changes in the intracellular levels of second messengers,

such as cAMP and Ca^{2+}, play central roles in the sensation of odour, taste and vision[9,10,17-20]. For

instance, the sensation of odour by olfactory receptors triggers the cAMP signalling cascade

that finally results in an influx of Ca^{2+} and excitation of olfactory neurons[9,17]. Detection of the

sweet or bitter by taste receptors produced DAG and IP3, followed by the opening of the taste-

transduction channel TRPM5[10,21]. Since our G protein activity analysis indicated that the force

sensation by LPHN2 was able to activate Gs and Gi, two G protein subtypes modulating

intracellular cAMP levels, we measured the changes in cAMP levels downstream of LPHN2 in

the utricle explant using paramagnetic beads coated with anti-LPHN2 antibody (referred to as

LPHN2-M-beads), which recognizes the N-terminal residues 551-600 of LPHN2. In the

heterologous system, we showed that upon magnetic force application, the LPHN2-M-beads

induced dose-dependent cAMP accumulation in LPHN2-overexpressing HEK293 cells but not

in cells transfected with the empty vector pcDNA3.1 (Figures 4A and S4A). The EC50 for the

LPHN2-M-bead-induced cAMP level increase in vitro was 1.70 ± 0.34 pN. In contrast, the

polylysine-coated paramagnetic beads (Ctrl-beads) could not induce cAMP production in

LPHN2-expressing HEK293 cells in response to force application, thus supporting the

specificity of the LPHN2-M-bead in activating LPHN2 (Figure 4A). We subsequently applied

the LPHN2-M-bead to the murine utricle explant and revealed that the administration of

magnetic force to the beads dose-dependently increased the intracellular cAMP level from

approximately 4.4 mM to 6.4 mM in the *Pou4f3-CreER$^{+/-}$Lphn2$^{+/+}$* utricle explant (with an

EC50 of 1.64 ± 0.13 pN) but not in the *Pou4f3-CreER$^{+/-}$Lphn2$^{fl/fl}$* utricle explant, using the

adenylyl cyclase activator forskolin as the positive control (Figures 4B and S4B). In contrast,

the endogenous cAMP levels in the utricle explants were not significantly affected by force

application via Ctrl-beads (Figure 4B). These results indicated that Gs-cAMP signalling

downstream of LPHN2 was activated in utricular hair cells in response to force stimulation.

Consistently, the expression profiling of isolated utricular hair cells by single-cell qRT–PCR

revealed that Gs is the most abundantly expressed G protein subtype in 15 tested LPHN2-positive cells (Figures 4C and 4D). Moreover, the force stimulation activated the Gs pathway of LPHN2 in the heterologous system with an EC50 of 1.75 ± 0.02 pN, which is significantly stronger than the EC50 values of Gi subtype pathways (Gi1: 3.79 ± 0.05 pN, Gi2: 4.61 ± 0.09 pN, Gi3: 3.52 ± 0.09 pN) (Figures 4E-4G and S4C-S4D). In contrast, the mechanical force applied by LPHN2-M-beads did not activate the Go or G12 pathway downstream of LPHN2 (Figures S4E and S4F).

Force sensing by LPHN2 induces Gs-dependent Ca^{2+} influx in utricular hair cells

Intracellular Ca^{2+} is a key element for neurotransmitter release and may play a pivotal role in equilibration sensation[21-24]. We next examined the Ca^{2+} response downstream of LPHN2 in single hair cells after force stimulation. The AAV-ie-*Lphn2pr*-mCherry, in which the original CAG promoter in AAV-ie was replaced with the promoter region (800 bp) of *Lphn2*, was delivered to the P3 mice through round window membrane injection (Figures 4H, S4G and S4H). This modified AAV-ie-*Lphn2pr*-mCherry enabled mCherry expression specifically in LPHN2-expressing utricular hair cells, which was supported by immunofluorescence and single-cell RT–PCR (Figure S4I). The Ca^{2+} signal in LPHN2-expressing utricular hair cells in response to force stimulation was recorded by a Fura-2 fluorescent probe. Approximately 35% of the LPHN2-expressing hair cells derived from the *Pou4f3-CreER$^{+/-}$Lphn2$^{+/+}$* utricle showed an increase in Ca^{2+} levels in response to force stimulation (Figures 4I-4K). In contrast, the AAV-ie-*Lphn2pr*-mCherry-labelled utricular hair cells derived from *Pou4f3-CreER$^{+/-}$Lphn2$^{fl/fl}$* mice showed no detectable Ca^{2+} signals in response to force application via LPHN2-M-beads (Figures 4I-4K and S4J). Importantly, the force-stimulated Ca^{2+} response in LPHN2-expressing hair cells was completely abolished by pre-treatment with the Gs inhibitor NF-449 but was not affected by the PKA inhibitor PKI 14-22 or the Gq inhibitor YM-254890 (Figure 4J). These results suggested that the force-induced Ca^{2+} response was Gs-dependent and Gq-independent. Consistent with this hypothesis, force stimulation of LPHN2 did not induce detectable production of IP3, the messenger molecule downstream of the Gq pathway, in utricular hair cells (Figure 4M). Moreover, the force-induced Ca^{2+} response in LPHN2-expressing hair cells was abolished when the utricle explants were incubated in Ca^{2+}-free buffer (Figure 4N).

Collectively, these results suggested that the activation of LPHN2 by force stimulation induced

Gs-dependent Ca^{2+} influx in utricular hair cells, which was most likely regulated by an

unknown cation channel.

LPHN2 activation by force elicits membrane depolarization in utricular hair cells

The force-induced Ca^{2+} influx in utricular hair cells through activating LPHN2 suggested

that an unknown downstream ion channel localized in the hair cell was open and might cause

membrane potential changes. Therefore, we examined the excitability of the utricular hair cells

in response to force activation of LPHN2 via LPHN2-M-beads by whole-cell patch-clamp

recording. We found that approximately 32% of LPHN2-expressing hair cells (labelled with

AAV-ie-*Lphn2pr*-mCherry) in *Pou4f3-CreER$^{+/-}$Lphn2$^{+/+}$* utricle explants showed an

approximately 50% increase in current amplitude in response to 3 pN force stimulation (17 out

of 53, cluster I), whereas the other 68% were force insensitive (36 out of 53, cluster II) (Figures

5A-5C). In general, cluster I cells had a higher maximum current amplitude than cluster II cells

by approximately 1.7-fold at a holding potential of 100 mV (Figure S5A). The proportion of

force-sensitive utricular hair cells was in accordance with that recorded in the force-stimulated

calcium response assay (Figure 4K). As the negative control, AAV-ie-*Lphn2pr*-mCherry-

labelled hair cells in utricle explants derived from *Pou4f3-CreER$^{+/-}$Lphn2$^{fl/fl}$* mice showed no

detectable electrophysiological response to force stimulation via LPHN2-M-beads (Figures 5A-

5C and S5B). Notably, similar to those in their *Pou4f3-CreER$^{+/-}$Lphn2$^{+/+}$* counterparts, the

mCherry-labelled utricular hair cells in *Pou4f3-CreER$^{+/-}$Lphn2$^{fl/fl}$* mice could also be divided

into two clusters, with cluster I cells (13 out of 39, 33%) showing significantly higher current

amplitude than cluster II cells (26 out of 39, 67%) at the same holding potentials (Figures 5A-

5B). The recorded currents in cluster I cells were voltage-dependent and showed pronounced

outwards rectification, with a reversed potential of approximately -20 mV (Figures 5A-5B).

To further explore the functional consequences of force-stimulated LPHN2 in utricular hair

cells, we recorded the resting membrane potential of the force-sensitive cluster I cells using a

current clamp. When holding at 0 pA, force stimulation with LPHN2-M-beads induced dose-

dependent depolarization of the *Pou4f3-CreER$^{+/-}$Lphn2$^{+/+}$* cluster I cells with an EC50 of 1.83

$\pm$ 0.05 pN (Figures 5D, S5C and S5D), which is similar to that recorded in the Gs protein

activation assay in LPHN2-overexpressing HEK293 cells (Figure 4E). In contrast, the cluster I

hair cells from *Pou4f3-CreER*$^{+/-}$*Lphn2*$^{fl/fl}$ mice showed no significant membrane potential

changes in response to force stimulation via LPHN2-M-beads (Figure 5D). Notably, the

alteration of the hair cell membrane potential was recorded as 704.73 ± 8.99 ms after the force

application, which was recovered to the basal level after 3 s of force stimulation (with a time

latency of 365.13 ± 10.10 ms after force removal) (Figures 5D and 5E). The response kinetics

of the force-induced membrane potential change could be fitted with a monoexponential

equation, with a response time constant (τ_p) and a recovery time constant (τ_{rec}) of 253.70 ± 3.78

ms and 250.33 ± 10.15 ms, respectively (Figure 5E). Consistent with the current clamp data,

when holding at a potential of +100 mV or -70 mV, the current amplitudes of the cluster I hair

cells derived from the *Pou4f3-CreER*$^{+/-}$*Lphn2*$^{+/+}$ utricle explants, but not those from *Pou4f3-*

CreER$^{+/-}$*Lphn2*$^{fl/fl}$ utricle explants, were significantly increased in response to force stimulation

via LPHN2-M-beads (Figures 5F and S5E). Importantly, both the force-stimulated increase in

current and membrane potentials were completely blocked by pre-treatment with the Gs

inhibitor NF449 (Figures 5G,5H and S5F). Collectively, these results suggest that force-

stimulated LPHN2 activation in utricular hair cells promoted Ca^{2+} influx and membrane

depolarization, probably via an unknown cation channel.

LPHN2 regulates mechanoelectrical transduction through coupling to CNGA3

The dependency of LPHN2-mediated calcium influx and membrane depolarization on Gs-

cAMP signalling implies that the force sensation of LPHN2 might be converted to electrical

signals by activating cAMP-sensitive ion channel(s). Therefore, we analysed the expression

pattern of cyclic nucleotide–gated (CNG) channels and hyperpolarization-activated cyclic

nucleotide–modulated (HCN) channels, two principle subfamilies of cyclic nucleotide-

regulated cation channels that play essential roles in various neuronal activities[25,26], in utricular

hair cells. The scRNA-seq data indicated that at least 3 CNG channels (CNGA1, CNGA3 and

CNGB1) and 2 HCN channels (HCN1 and HCN4) were found to be expressed in LPHN2-

expressing utricular hair cells (Figure 6A). Further qRT–PCR analysis of single utricular hair

cells revealed that CNGA3 was specifically expressed in force-sensitive LPHN2-expressing

hair cells (cluster I cells), whereas the other channels were almost equally expressed in cluster

I and II cells (Figure 6B). Furthermore, only CNGA3, but not the other cyclic nucleotide-regulated channels expressed in LPHN2-expressing utricular hair cells, showed direct association with LPHN2, as revealed by an *in vivo* coimmunoprecipitation assay (Figure 6C). Consistently, a substantial portion of LPHN2 showed coimmunostaining with CNGA3 at the cuticular plate of utricular hair cells, supporting interactions between these two membrane proteins (Figure 6D).

Next, we tested whether force-induced activation of LPHN2 was directly coupled with CNGA3 current changes using an in vitro HEK293 overexpression system. Notably, co-expression of CNGA3, but not CNGA1 or HCN1/4 channels, enabled the membrane excitability of the LPHN2-expressing HEK293 cells in response to 3 pN force stimulation applied by LPHN2-M-beads, thus suggesting the importance of the physical association between LPHN2 and CNGA3 channels (Figures 6E and S6A-S6C). At the holding potential of +100 mV, the force induced a dose-dependent increase in current amplitude only in HEK293 cells co-expressing LPHN2 and CNGA3, with an EC50 of 1.71 ± 0.01 pN (Figures 6F and S6D). This EC50 value was close to those recorded for the force-induced membrane potential changes or cAMP accumulation via LPHN2-M-beads in utricle explants (Figure 6G). Collectively, these results suggest the direct interaction and functional coupling between CNGA3 and LPHN2 in response to force stimulation.

Inhibition of CNGA3 activity severely impairs balance

To further explore the functional importance of CNG channels in balance maintenance, we assessed the effects of the CNG inhibitor L-cis-diltiazem on the equilibration-related behaviours of mice. Notably, intraperitoneal injection of L-cis-diltiazem for 3 consecutive days significantly increased the circling behaviour and travelling distance of mice compared with the control group in both the 2 min time frame (9.75 ± 1.0 circles vs. 5.75 ± 0.63 circles) and 10 min time frame (30.00 ± 1.68 circles vs. 21.50 ± 1.85 circles) (Figures S7A-S7C). Moreover, L-cis-diltiazem administration significantly deteriorated mouse performance in both forced swimming (swimming scores of 1.50 ± 0.29 vs. 0) and rotarod tests (staying time of 66.70 ± 5.34 s vs. 103.54 ± 9.32 s) (Figures S7D and S7E). These data suggested a regulatory role of the CNG channel in vestibular functions.

To exclude the potential nonselectivity of L-cis-diltiazem by inhibiting multiple CNG

channel subtypes and to inspect the specific role of CNGA3 in vestibular hair cells, we

generated a hair cell-specific *Cnga3*-knockout mouse line by crossing *Cnga3^{fl/fl}* mice with

Pou4f3-CreER^{+/-} transgenic mice (Figures S7F and S7G). The specific knockdown

(approximately 50%) of CNGA3 expression in the vestibule of *Pou4f3-CreER^{+/-}Cnga3^{fl/+}* mice

was verified by both western blotting (Figures S7H and S7I). In contrast, the CNGA3

expression levels in other tissues, such as the brainstem and skin, were not significantly affected

(Figures S7H and S7I). Importantly, compared with their *Pou4f3-CreER^{+/-}Cnga3^{+/+}* littermates,

the *Pou4f3-CreER^{+/-}Cnga3^{fl/+}* mice showed significantly impaired balance behaviours,

including increased circling and travelling distance, reduced duration time on the rotarod and

deteriorated swimming performance (Figures 7A-7E). Moreover, the *Pou4f3-CreER^{+/-}Cnga3^{fl/+}*

mice showed a more than 80% reduction in VOR gain in off-vertical axis rotation (otolith

function test) and a 25%-40% reduction in VOR responses to earth-vertical axis rotation

(semicircular canal function test) (Figures 7F-7H). These results collectively indicated a critical

role of CNGA3 channels, especially those expressed in otolith organs, in normal balance

maintenance. These channels most likely act as a key signalling component downstream of

force-sensing LPHN2 and directly converts the chemical signals delivered by second

messengers into electrical signals. Consistent with this hypothesis, the application of 100 μM

L-cis-diltiazem in cluster I utricular hair cells completely abolished the force-stimulated

calcium signals or current amplitude change after the activation of LPHN2 by LPHN2-M-beads

(Figures 7I-7K). In other words, inhibition of CNGA3 activity eliminated the force sensitivity

of LPHN2-expressing utricular hair cells and turned all of the LPHN2-expressing cells into

cluster II cells (Figures S7J and S7K). Altogether, these results suggest that an LPHN2-CNGA3

functional axis modulates the membrane excitability of utricular hair cells in response to force

sensation, which contributes to equilibrioception.

Notably, CNGA3 expression is not confined to the cuticular plate of utricular hair cells where

it colocalizes with LPHN2 but is also observed at the lateral and basal membranes (Figure 6D).

Therefore, we speculated that CNGA3 might also regulate utricular hair cell functions through

other pathways or connect with other GPCRs, in addition to its crosstalk with LPHN2. This

notion might explain the aggravated phenotypes of *Cnga3*-deficient mice in balance behaviour

tests compared with those of *Lphn2*-deficient mice.

Re-expression of LPHN2 in vestibular hair cells rescued the equilibrioception of LPHN2-deficient mice

To further inspect the functional roles of LPHN2 specifically expressed in the vestibular

system, we reintroduced LPHN2 expression in utricular hair cells of *Pou4f3-CreER*$^{+/-}$*Lphn2*$^{fl/fl}$

mice via AAV delivery. Our previous structural and functional analyses revealed that the force

sensation by adhesion GPCRs was primarily mediated by the juxtamembrane GAIN domain[27,28].

Consistently, the Flag-tagged LPHN2-GAIN construct with the deletion of 523 N-terminal

residues retained mechanosensitive potential and showed dose-dependent Gs signalling in

response to force stimulation with Flag-M-beads, and the response was comparable to that of

full-length LPHN2 when expressed at similar levels (Figures S8A and S8B). Therefore, we

packaged the Flag-LPHN2-GAIN sequences into the AAV-ie-*Lphn2pr*-mCherry vector

(mCherry following an IRES element was fused to the C-terminus of LPHN2-GAIN, referred

to as AAV-ie-LPHN2) and delivered the virus into P3 *Pou4f3-CreER*$^{+/-}$*Lphn2*$^{fl/fl}$ mice through

round window membrane injection using AAV-ie-*Lphn2pr*-mCherry as a negative control

(Figure S8C). The specific expression of LPHN2-GAIN in the utricular hair cells, but not in

the brains, of *Pou4f3-CreER*$^{+/-}$*Lphn2*$^{fl/fl}$ mice was observed 14 days after the administration of

AAV-ie-LPHN2 using western blotting and immunofluorescence (Figures S8D and S8E). As a

negative control, *Pou4f3-CreER*$^{+/-}$*Lphn2*$^{fl/fl}$ mice infected with AAV-ie-*Lphn2pr*-mCherry

revealed only specific utricular hair cell labelling but no detectable LPHN2-GAIN expression

(Figures S8D and S8E).

We next assessed the equilibration-related behaviours of mice receiving LPHN2-GAIN gene

delivery. Notably, the LPHN2-deficient mice treated with AAV-ie-LPHN2 showed significantly

improved performance in all behaviour studies related to equilibrioception, including the forced

swimming, open field and rotarod tests, and the performances were comparable to those of the

WT mice (Figures 8A-8D). Moreover, the VOR gain values of AAV-ie-LPHN2-treated *Pou4f3-*

CreER$^{+/-}$*Lphn2*$^{fl/fl}$ mice at all tested frequencies in both the earth-vertical and off-vertical axis

rotation tests recovered to levels comparable to those of WT mice (Figures 8E and 8F). As a

negative control, LPHN2-deficient mice treated with AAV-ie-*Lphn2pr*-mCherry showed no

significant improvement in any of the above tests (Figures 8A-8F). Furthermore, compared

with the mCherry-labelled utricular hair cells in control virus-treated *Pou4f3-CreER*[+/-]*Lphn2*[fl/fl]

mice, which showed no Ca^{2+} response or current amplitude changes to Flag-M-bead-induced

force stimulation, approximately 31% and 30% of the labelled utricular hair cells in AAV-ie-

LPHN2-treated *Pou4f3-CreER*[+/-]*Lphn2*[fl/fl] mice regained force-induced excitability, resembling

that observed for WT utricle cluster I hair cells (Figures 8G, S8F and S8G). Therefore,

reintroduction of the mechanosensitive LPHN2-GAIN specifically in *Lphn2* promoter-labelled

utricular hair cells was able to rescue the equilibrioception of LPHN2-deficient mice. These

findings further supported that force-activated LPHN2 in the vestibular system directly

contributes to normal equilibrioception.

Discussion

Despite its importance for our daily living and motion in three-dimensional space, the

molecular mechanism underlying the sense of balance remains elusive. Previous studies have

revealed that ion channels, such as TMC1/2, are potential key components of the MET process,

which is essential for equilibrioception[29-31]. However, direct evidence for force sensation by

TMC1/2 is lacking, and the direct force sensors that participate in equilibrioception are not clear.

In addition to ion channels, another group of membrane receptors, GPCRs, are known as direct

sensors for vision, odour and touch[8,32,33]. Although recent studies have identified that the GPCR

member CELSR is essential for equilibrioception by modulating the planar polarity of hair

cells[7], whether GPCRs, especially those with mechanosensitive potentials, directly participate

in the MET process of equilibrioception remains unclear. In our parallel work, which we started

three years earlier than the present manuscript, we found that the mechanosensitive GPCR

GPR133 participates in MET and is indispensable for balance maintenance in mice. Here, we

identified that another mechanosensitive GPCR, LPHN2, is also able to sense force in

vestibular hair cells to mediate MET and is essential for equilibrioception. These two parallel

works support the likely existence of multiple GPCRs that function as force sensors and MET

mediators essential for equilibrioception, although further exploration is needed to confirm this

hypothesis.

In our study, heterozygous *Lphn2*-deficient mice showed reduced balance performance and

impaired VOR, which were further deteriorated in mice with hair cell-specific ablation of

LPHN2 in an inducible manner. Importantly, specific re-expression of LPHN2 in vestibular hair

cells rescued the equilibrioception of *Pou4f3-CreER^{+/-}Lphn2^{fl/fl}* mice. These data collectively

support the notion that mechanosensitive LPHN2 is at least a part of the balance-sensing

machinery in peripheral vestibular end organs. Intriguingly, LPHN2 is specifically expressed

at the cuticular plate of utricular hair cells, which is similar to the expression pattern of the

mechanotransduction channel PIEZO2[34]. Therefore, we speculated that the MET process is not

confined to hair cell stereocilia and that mechanosensitive GPCRs and/or channels at the apical

surface might also play important roles in the MET process through direct mechanical force

sensation or interaction with the extracellular or otoconial matrix.

Notably, GPCRs play important roles in the sensation of vision, olfaction and three taste

qualities (sweet, bitter and umami) by modulating the intracellular levels of second messengers.

For instance, the sensation of vision by rhodopsin relies on the Gt-mediated reduction of cGMP

levels in photoreceptor cells, and the perception of odour by odourant receptors depends on the

G_{olf}-regulated accumulation of cAMP in olfactory neurons[19,35]. In a parallel paper, we showed

that GPR133 decreased the basal cAMP levels of vestibular hair cells in a Gi-dependent manner

after force stimulation. We speculated that the inhibitory effect of the mechanosensitive

GPR133 on intracellular cAMP levels might resemble the regulatory processes that occur

during vision sensation. In contrast, our present work indicated that LPHN2 could promote

MET in utricular hair cells by activating Gs-cAMP signalling, demonstrating a positive

modulation of vestibular membrane excitability. This special GPCR-mediated positive MET

resembles the signalling mechanisms for olfaction and sweet/bitter taste perception. Therefore,

we speculated that the force sensation mediated by LPHN2 and GPR133, as well as potentially

other GPCRs, in vestibular hair cells might encode combinatory positional information. For a

simplified model, we hypothesized that a given force applied to hair cells with different

expression patterns of these two receptors (LPHN2^{+}GPR133^{+}, LPHN2^{+}GPR133^{-}, or LPHN2^{-}

GPR133^{+}) might result in distinct intracellular cAMP signalling patterns, which may cause

differences in membrane excitability or neurotransmitter release from specific spiral ganglion

neurons (SGNs). We further hypothesized that the potential involvement of other

mechanosensitive GPCRs may increase the temporal and spatial complexity of the information

input.

Our study also reveals that a previously uncharacterized vestibular MET-regulating ion

channel CNGA3, in addition to TMC1/2 and proton-conducting Otopetrin-1[36], is essential for

equilibrioception. Here, our results suggested that CNGA3 channel activity is coupled with the

activation of at least two mechanosensitive GPCRs, GPR133 and LPHN2, and likely in

response to changes in intracellular cAMP levels, modulates the membrane excitability of

vestibular hair cells during force sensation. Whereas LPHN2 activates the CNGA3 channel and

induces membrane depolarization, GPR133 inhibits CNGA3 channel activation and induces

membrane hyperpolarization. Importantly, inhibiting CNGA3 activity eliminates LPHN2-

mediated MET both *in vitro* and *in vivo*, leading to significant balance dysfunction in mice.

Kinetics analysis of LPHN2-CNGA3 coupling revealed that the delayed time between force

application and the electrophysiological response was approximately 704.73 ± 8.99 ms, which

is close to that recorded for GPR133-CNGA3 coupling (658.0 ± 11.14 ms). Therefore, we

speculated that equilibrioception regulated by certain GPCRs in the vestibular system might be

a relatively long process, which opposes the traditional views on hearing sensation and

equilibrioception[37,38]. Notably, CNG family members play essential roles in vision and olfaction

by converting the intracellular chemical signals carried by second messengers to electrical

signals[25]. In particular, whereas the rod photoreceptor cells mediating vision in dim light contain

CNGA1-CNGB heteromeric channels, the cone photoreceptor cells responsible for bright light

perception and colour vision have CNGA3-CNGB3 channels. Moreover, CNG channels in

olfactory neurons are composed of CNGA2-CNGA4-CNGB1b subunits. Our results indicated

an essential role of CNGA3 in equilibrioception in utricular hair cells and further supported the

functional importance of the GPCR-CNG axis in the regulation of basic senses. However,

whether CNGA3 forms homomeric channels in vestibular hair cells or interacts with other CNG

subunits to form heteromeric channels awaits further investigation. Due to its functional

importance in cone photoreceptor cells, CNGA3 mutation is one of the most common causes

of achromatopsia, an inherited eye disease associated with congenital absence of cone

photoreceptor function[39]. Although the vestibular functions of the affected individuals have

rarely been reported, they usually suffer from nystagmus, one of the typical syndromes of

vestibular diseases[40]. Therefore, it will be of interest to investigate the pathophysiological

effects of these CNGA3 mutations on the vestibular system, which will provide further insight into CNGA3 functions.

In conclusion, we identified a force-sensitive GPCR expressed in the cuticular plate of vestibular hair cells that is essential for equilibrioception and mediates MET by increasing intracellular cAMP levels and modulating CNGA3 activity. Our results indicated that GPCRs could be direct force sensors during equilibrioception.

Data availability

All data are included in the main text or the extended materials. Data are available when requested from the corresponding authors.

Acknowledgments

This work was supported by National Key R&D Program of China (2019YFA0904200 to J.-P.S., 2021YFA1101300 to R.-J.C.), National Science Fund for Distinguished Young Scholars Grant (81825022 to J.-P.S., 82225011 to X.Y.), National Natural Science Foundation of China (32130055 and 81773704 to J.-P.S., 92057121 to X.Y., 82030029 to R.-J.C., 82271176 to W.-W.L.), the Key Research Project of the Beijing Natural Science Foundation, China (Z200019 to J.-P.S.), Major Fundamental Research Program of Natural Science Foundation of Shandong Province, China (ZR2020ZD39 to J.-P.S. and W.-W.L, ZR2021ZD18 to X.Y.), Taishan Scholars Program of Shandong Province (tsqn201909189 to W.-W.L.).

Author contributions

J.-P.S., X.Y. and R.-J.C initiated, designed and supervised the overall project. J.-P.S., X.Y. and Z.Y. starting the screening of mechanosensitive GPCRs in vestibular and cochlear systems from 2019. R.-J.C provided guidance and suggestions on the vestibular and cochlear studies. Z.Y. and J.-P.S. developed the magnetic force stimulation assay to examine the activation of G proteins and second messengers downstream of GPCRs. X.Y. designed and supervised all electrophysiology experiments. Z.Y., S.-H.Z. and Q.-Y.Z. performed the cell-based assay of LPHNs. S.-H.Z., Z.Y., Q.-Y.Z. and Z.-C.S. performed the electrophysiological experiments and Ca^{2+} imaging supervised by J.-P.S. and X.Y.. J.-P.S., Z.Y., S.-H.Z. and W.-W.L. participated in

the design of gene knockout mice and modified AAV-ie. Z.Y., S.-H.Z., Q.-Y.Z. and W.-W.L. performed vestibular behaviour studies. Z.Y., S.-H.Z., Q.-Y.Z. and Y.G. isolated mouse utricle and performed immunofluorescence studies and ex vivo experiments. Z.Y., M.-W.W., S.-H.Z., and Q.-Y.Z. performed single cell qRT-PCR. Y.G., M.-W.W., S.-H.Z. and Y.-N.S. performed coimmunoprecipitation and western blotting analyses. Z.Y., S.-H.Z., Q.-Y.Z. and W.-W.L. performed the AAV-ie-related experiments under the supervision of J.-P.S. and R.-J.C.. J.-P.S., X.Y., R.-J.C., Z.Y., S.-H.Z., Q.-Y.Z., Y.S. and W.Y. participated in data analysis and interpretation. Y.S. provided insightful ideas related to the balance sensation based on clinical practice. Z.Y., S.-H.Z., and Q.-Y.Z. prepared the figures. J.-P.S., X.Y., Z.Y. and R.-J.C wrote the manuscript. All the authors have seen and commented on the manuscript.

Competing interests

The authors declare no competing interests.

Submission history

The manuscript was submitted on Sep 20, 2023 and was sent for peer review on Sep 26, 2023 by the journal editor.

Figure Legends

Figure 1. Deficiency of mechanosensitive LPHN2 but not LPHN3 impairs balance in mice

(A) Schematic representation of the potential roles of TMC1/2 and GPCRs in MET in vestibular hair cells.

(B) Expression frequency of LPHN subfamily members in mouse utricular hair cells. Data represent the average values of the LPHN expression frequency from a single-cell RNA-seq dataset (GSE71982) and the results of single-hair-cell qRT–PCR analysis in the current study.

(C) The mRNA levels of LPHN subfamily members in mouse utricular hair cells measured by single-cell qRT–PCR, using their respective expression levels in the brain as a reference (normalized to 1) (n=3). Data are correlated to Figure S1B.

(D-E) Gi3 (**D**) or Gs (**E**) activation in HEK293 cells overexpressing different LPHNs in

response to 3 pN force stimulation measured by G protein dissociation BRET assay (n=3).

HEK293 cells transfected with the empty vector pcDNA3.1 and G protein probes were used as

a negative control.

(F) Schematic view (left) and representative tracks (right) of WT, $Lphn2^{+/-}$, and $Lphn3^{+/-}$ mice

in open-field test during 2 min or 10 min tracking period.

(G-I) Quantification of the circling and travelling activity in open-field test (**G**), duration time

on the rotating rod (**H**), and swimming scores (**I**) of WT, $Lphn2^{+/-}$, and $Lphn3^{+/-}$ mice (n=9 mice

per group).

(J) Schematic view (left) and representative recording curves (right) of the VOR response of

WT, $Lphn2^{+/-}$, and $Lphn3^{+/-}$ mice to earth-vertical axis rotation (0.20-1.5 Hz, 40°/s peak velocity

sinusoidal, whole-body passive rotation).

(K) Quantification of the VOR gain response of WT, $Lphn2^{+/-}$, and $Lphn3^{+/-}$ mice to earth-

vertical axis rotation (n=9 mice per group).

(L) Schematic view (left) and representative recording curves (right) of the VOR response of

WT, $Lphn2^{+/-}$, and $Lphn3^{+/-}$ mice to off-vertical axis rotation (50°/s, whole-body passive

rotation).

(M) Quantification of the VOR gain response of WT, $Lphn2^{+/-}$, and $Lphn3^{+/-}$ mice to off-vertical

axis rotation (n=9 mice per group).

(D-E) $*P < 0.05$; $**P < 0.01$; ns, no significant difference. HEK293 cells transfected with

LPHNs compared with those transfected with pcDNA3.1. **(G-I, K, M)** $**P < 0.01$; $***P <$

0.001; ns, no significant difference. Gene knockout mice compared with WT mice. The bars

indicate mean ± SEM values. All data were statistically analysed using one-way ANOVA with

Dunnett's post hoc test.

Figure 2. LPHN2 is specifically expressed at the cuticular plate of utricular hair cells

(A) Coimmunostaining of LPHN2 (green) with Myosin7a (red) in utricle whole mounts derived

from WT mice at P40 or from $Lphn2^{-/-}$ mouse embryos at E18 (n = 3 mice per group). Scale

bar, 50 μm.

(B) Representative images of whole-mount RNAscope in situ hybridization of LPHN2 (white)

combined with immunostaining of Myosin7a (red) in the utricle of P40 mice (n = 3 mice).

Arrows indicate LPHN2 staining in myosin7a-expressing hair cells. Scale bars, 50 μm and 10

μm for low- and high-magnification views, respectively.

(C) Quantitative analysis of LPHN2 expression in myosin7a-positive utricular hair cells from

WT mice or *Lphn2$^{-/-}$* embryos. Data are correlated to Fig. 2a-b (n = 3 mice per group).

(D) Left panel: Diagram of utricular hair cells showing the selected optical planes (stereocilia,

cuticular plate or basolateral section) for imaging by confocal microscopy. Right panel:

Coimmunostaining of spectrin (red), LPHN2 (green) and phalloidin (magenta) at different

optical planes of hair cells in utricle whole mounts derived from WT mice (P40) or from *Lphn2^{-}*

$^{/-}$ embryos (E18). Arrows indicate coimmunostaining of LPHN2 with spectrin at the cuticular

plate. The line of polarity reversal (LPR) is depicted as a white dotted line. Scale bar, 10 μm.

(E) Left panel: Schematic view of LPHN2 (red) expression in utricular hair cells. Right panel:

Coimmunostaining of LPHN2 (green) with Myosin7a (red) or SOX2 (magenta) in utricle

sections derived from WT mice at P40 (n = 3 mice per group). Arrows indicate the distribution

of LPHN2 immunofluorescence puncta at the cuticular plate of utricular hair cells (depicted by

white dotted lines). Scale bars, 50 μm and 10 μm for low- and high-magnification views,

respectively.

Figure 3. Conditional knockout of LPHN2 in mouse hair cells impaired balance

(A) Schematic representation of the crossbreeding strategy to generate hair cell-specific *Lphn2*

knockout mice and vestibular function analysis.

(B) Immunostaining of LPHN2 (magenta) and POU4F3 (magenta) in utricle whole mounts

derived from *Pou4f3-CreER$^{+/-}$Lphn2$^{fl/fl}$* mice (referred to as *Pc-Lphn2$^{fl/fl}$*) or *Pou4f3-CreER$^{+/-}$*

Lphn2$^{+/+}$ mice (referred to as *Pc-Lphn2$^{+/+}$*). Enlarged images show the ablation of LPHN2 in

the utricular hair cells of *Pc-Lphn2$^{fl/fl}$* mice (n=3 mice per group). Scale bar: 50 μm and 20 μm

for low and high magnification views, respectively.

(C-F) Quantification of the swimming scores (**C**), time on the rotating rod (**D**), circling activity

(**E**) and travelling activity (**F**) in the open field test of *Pc-Lphn2$^{fl/fl}$* mice and *Pc-Lphn2$^{+/+}$* mice

(n=8 mice per group).

(G) Representative recording curves (top) and quantification of the VOR gain response (bottom)

of *Pc-Lphn2$^{fl/fl}$* mice and *Pc-Lphn2$^{+/+}$* mice to earth-vertical axis rotation (n=8 mice per group).

(H) Representative recording curves (left) and quantification of the VOR gain response (right) of *Pc-Lphn2*$^{fl/fl}$ mice and *Pc-Lphn2*$^{+/+}$ mice to off-vertical axis rotation (n=8 mice per group).

(C-H) *P < 0.05; **P < 0.01; ***P < 0.001; ns, no significant difference. *Pc-Lphn2*$^{fl/fl}$ mice compared with *Pc-Lphn2*$^{+/+}$ mice. The bars indicate mean ± SEM values. All data were statistically analysed using unpaired two-sided Student's *t* test.

Figure 4. Force sensation by LPHN2 induces Ca^{2+} influx in utricular hair cells by activating Gs-cAMP signalling

(A) Dose-dependent cAMP production in LPHN2-overexpressing HEK293 cells in response to force applied by LPHN2-M-beads. Polylysine-coated beads (Ctrl-beads) were used as a negative control (n=3).

(B) cAMP production in utricle explants derived from *Pc-Lphn2*$^{fl/fl}$ mice and *Pc-Lphn2*$^{+/+}$ mice in response to stimulation with 3 pN force or 10 μM forskolin (n=3).

(C) mRNA levels of different G protein subtypes in the utricle sensory epithelium using their respective expression levels in the brain as a reference (normalized to 1) (n=3).

(D) Heatmap showing the relative mRNA levels of G protein subtypes in 15 isolated LPHN2-expressing utricular hair cells measured by single-cell qRT–PCR. Data are normalized to the respective expression levels of these G protein subtypes in the brain.

(E-F) Dose-dependent Gs (**E**) or Gi3 (**F**) activation in LPHN2-overexpressing HEK293 cells in response to force applied by LPHN2-M-beads or Ctrl-beads (n=3).

(G) Summary of the potency of force-induced Gs or Gi subtype activation in LPHN2-overexpressing HEK293 cells (n=3). Data are correlated to Figure 4E-4F and Figure S4C-S4F.

(H) Schematic illustration of Ca^{2+} imaging in mouse utricular hair cells labelled with AAV-ie-*Lphn2pr*-mCherry in response to force stimulation.

(I) Heatmaps showing the Ca^{2+} responses in individual utricular hair cells derived from *Pc-Lphn2*$^{+/+}$ mice labelled with AAV-ie-*Lphn2pr*-mCherry (n = 84 cells from 11 mice). The vertical dashed line denotes the onset of force application (0 s). Each row represents the temporal Ca^{2+} response of one hair cell.

(J) Representative Ca^{2+} traces recorded in individual utricular hair cells derived from *Pc-Lphn2*$^{fl/fl}$ or *Pc-Lphn2*$^{+/+}$ mice labelled with AAV-ie-*Lphn2pr*-mCherry.

(K) Proportion of force-sensitive and force-insensitive utricular hair cells derived from *Pc-Lphn2*$^{+/+}$ mice labelled with AAV-ie-*Lphn2pr*-mCherry. Data are correlated to Fig. 4i.

(L) Effect of NF449, PKI 14-22 or YM-254890 treatment on the force-induced Ca^{2+} response in utricular hair cells labelled with AAV-ie-*Lphn2pr*-mCherry (n=10 cells per group).

(M) IP3 levels in the utricle explant of WT mice in response to stimulation with 3 pN force (n=3).

(N) Effect of Ca^{2+} ablation in the incubation buffer on the force-induced Ca^{2+} response in LPHN2-positive utricular hair cells (n=10).

(B, M) **$P < 0.01$; ns, no significant difference. Utricle explants treated with force or forskolin compared with control utricle explants. (L) ***$P < 0.001$; ns, no significant difference. Utricular hair cells treated with inhibitors compared with the control utricular hair cells. (N) ***$P < 0.001$. Utricular hair cells incubated with Ca^{2+}-free buffer compared with the control cells. The bars indicate mean ± SEM values. All data were statistically analysed using one-way ANOVA with Dunnett's post hoc test (B) or Student's *t* test (L-N).

Fig. 5 | Force-induced LPHN2 activation regulates the membrane excitability of utricular hair cells

(A-B) Representative whole-cell currents (upper panel) elicited by voltage steps between -100 mV and +100 mV and the corresponding I-V curves (lower panel) in AAV-ie-*Lphn2pr*-mCherry-labelled cluster I (A) or cluster II (B) utricular hair cells derived from *Pc-Lphn2*$^{fl/fl}$ or *Pc-Lphn2*$^{+/+}$ mice in response to force (3 pN) stimulation. n=17 for cluster I cells and n=36 for cluster II cells in *Pc-Lphn2*$^{+/+}$ mice. n=13 for cluster I cells and n=26 for cluster II cells in *Pc-Lphn2*$^{fl/fl}$ mice.

(C) Proportion of force-sensitive and force-insensitive utricular hair cells labelled with AAV-ie-*Lphn2pr*-mCherry derived from *Pc-Lphn2*$^{+/+}$ mice. Data are correlated to Fig. 5a-b.

(D) Representative force-induced membrane potential alteration of cluster I utricular hair cells derived from *Pc-Lphn2*$^{fl/fl}$ or *Pc-Lphn2*$^{+/+}$ mice when holding at absolute zero current (n=10).

The time latency between force application or force removal and the change in membrane

potential was defined as T_{pr} and T_{po}, respectively. The calculated time constant for the response

and the recovery time constant were denoted as τ_p and τ_{rec}, respectively.

(E) The calculated time parameters in force-stimulated membrane potential alteration of cluster

I utricular hair cells (n=10).

(F) Representative whole-cell currents in cluster I hair cells derived from *Pc-Lphn2*$^{+/+}$ mice in

response to 3 pN force stimulation at a holding potential of +100 mV or -70 mV (n=3).

(G) Effect of NF449 treatment on the whole-cell current in *Pc-Lphn2*$^{+/+}$ cluster I hair cells in

response to 3 pN force stimulation (n=11).

(H) Effect of NF449 treatment on the force-stimulated membrane potential alteration in *Pc-*

Lphn2$^{+/+}$ cluster I hair cells (n=13).

Figure 6. LPHN2 regulates MET in utricular hair cells through coupling to CNGA3

(A) Expression profiles of cyclic nucleotide–activated ion channels in LPHN2-expressing

utricular hair cells. A single-cell RNA-seq dataset (GSE71982) was used in this experiment.

(B) mRNA levels of CNG and HCN channels in WT cluster I and cluster II utricular hair cells

examined by single-cell qRT–PCR (n=10 cells per cluster). Data are normalized to the

respective expression levels of these ion channels in the brain.

(C) Coimmunoprecipitation of LPHN2 with CNG or HCN channels in mouse utricle cell

lysates. Representative blots from three independent experiments are shown (n=3).

(D) Left panel: Coimmunostaining of LPHN2 (green) with CNGA3 (red) in utricle sections

derived from WT mice (n = 3). Scale bar: 10 μm. Right panel: Pearson's correlation analysis of

LPHN2 and CNGA3 fluorescence intensities. The Pearson's correlation coefficient was 0.68.

(E) Representative whole-cell currents (upper panel) elicited by voltage steps between -100 mV

and +100 mV and the corresponding I-V curves (lower panel) in HEK293 cells overexpressing

LPHN2/CNGA3/CNGB1 (referred to as LPHN2$^+$) or CNGA3/CNGB1 (referred to as LPHN2$^-$)

in response to force (3 pN) stimulation (n=5).

(F) Representative whole-cell currents (left) and the corresponding dose–response curves (right)

in LPHN2$^+$ or LPHN2$^-$ HEK293 cells in response to force stimulation at a holding potential of

+100 mV (n=3). Data are correlated to Figure S4F.

(G) Summary of the potency of force-induced cAMP production or membrane potential/current

alteration in utricle explants from WT mice or in LPHN2-overexpressing HEK293 cells.

Figure 7. Heterozygous hair cell-specific *Cnga3* knockout mice showed impaired balance

(A) Representative tracks of *Pc-Cnga3$^{+/+}$* and *Pc-Cnga3$^{fl/+}$* mice in the open-field test during a

2 min or 10 min tracking period.

(B-E) Quantification of the circling activity **(B)** and travelling activity **(C)** in the open field test,

swimming scores **(D)** and duration time on the rotating rod **(E)** of *Pc-Cnga3$^{+/+}$* and *Pc-Cnga3$^{fl/+}$*

mice (n=10 for *Pc-Cnga3$^{+/+}$*mice and n=6 for *Pc-Cnga3$^{fl/+}$* mice).

(F) Representative recording curves (top) and quantification of the VOR gain response (bottom)

of *Pc-Cnga3$^{+/+}$* and *Pc-Cnga3$^{fl/+}$* mice to earth-vertical axis rotation (n=10 for *Pc-*

*Cnga3$^{+/+}$*mice and n=6 for *Pc-Cnga3$^{fl/+}$* mice).

(G) Representative recording curves (left) and quantification of the VOR gain response (right)

of *Pc-Cnga3$^{+/+}$* and *Pc-Cnga3$^{fl/+}$* mice to off-vertical axis rotation (n=10 for *Pc-Cnga3$^{+/+}$*mice

and n=6 for *Pc-Cnga3$^{fl/+}$* mice).

(H) Representative whole-cell currents (upper) elicited by voltage steps between -100 mV and

+100 mV and the corresponding I-V curves (lower) in AAV-ie-*Lphn2pr*-mCherry-labelled

cluster I utricular hair cells in response to force (3 pN) stimulation in the absence or presence

of 100 μM L-cis-diltiazem (n=10).

(I) Representative Ca^{2+} traces recorded in individual AAV-ie-*Lphn2pr*-mCherry-labelled

utricular hair cells in response to force (3 pN) stimulation in the absence or presence of 100 μM

L-cis-diltiazem.

(J) Quantification of the effect of L-cis-diltiazem treatment on the force-induced Ca^{2+} response

in utricular hair cells labelled with AAV-ie-*Lphn2pr*-mCherry (n=10 cells).

(B-G) ***P < 0.001, **P < 0.01; ns, no significant difference. *Pc-Cnga3$^{fl/+}$* mice compared with

Pc-Cnga3$^{+/+}$ mice. **(J)** ***P < 0.001. Utricular hair cells treated with L-cis-diltiazem compared

with control cells. The bars indicate mean ± SEM values. All data were statistically analysed

using unpaired **(B-G)** or paired **(J)** two-sided Student's *t* test.

Figure 8. Specific re-expression of LPHN2 in hair cells rescued the vestibular function of LPHN2-deficient mice

(A-D) Quantification of the circling (**A**) and travelling activity (**B**) in the open-field test during 10 min, swimming scores (**C**) and duration time on the rotating rod (**D**) of *Pc-Lphn2*$^{+/+}$, *Pc-Lphn2*$^{fl/fl}$, AAV-ie-LPHN2 mice and AAV-ie-*Lphn2pr* mice (n=6 mice per group).

(E-F) Representative recording curves (left) and quantification of the VOR gain responses (right) of *Pc-Lphn2*$^{+/+}$, *Pc-Lphn2*$^{fl/fl}$, AAV-ie-LPHN2 mice and AAV-ie-*Lphn2pr* mice to earth-vertical axis (**E**) or off-vertical axis (**F**) rotation (n=6 mice per group).

(G) Representative whole-cell currents (left) elicited by voltage steps between -100 mV and +100 mV and the corresponding I-V curves (right) in cluster I utricular hair cells derived from *Pc-Lphn2*$^{+/+}$, AAV-ie-LPHN2 mice and AAV-ie-*Lphn2pr* mice in response to force (3 pN) stimulation (n=5, 7, and 6 for *Pc-Lphn2*$^{+/+}$, AAV-ie-LPHN2 and AAV-ie-*Lphn2pr* group, respectively).

(A-F) *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$; ns, no significant difference. *Pc-Lphn2*$^{fl/fl}$ mice compared with *Pc-Lphn2*$^{+/+}$ mice. The bars indicate mean ± SEM values. All data were statistically analysed using one-way ANOVA with Dunnett's post hoc test.

References

1   Cullen, K. E. Vestibular processing during natural self-motion: implications for perception and action. *Nat Rev Neurosci* **20**, 346-363, doi:10.1038/s41583-019-0153-1 (2019).

2   Eatock, R. A. & Songer, J. E. Vestibular hair cells and afferents: two channels for head motion signals. *Annu Rev Neurosci* **34**, 501-534, doi:10.1146/annurev-neuro-061010-113710 (2011).

3   Zheng, W. & Holt, J. R. The Mechanosensory Transduction Machinery in Inner Ear Hair Cells. *Annu Rev Biophys* **50**, 31-51, doi:10.1146/annurev-biophys-062420-081842 (2021).

4　Cunningham, C. L. & Muller, U. Molecular Structure of the Hair Cell
Mechanoelectrical Transduction Complex. *Cold Spring Harb Perspect Med* **9**,
doi:10.1101/cshperspect.a033167 (2019).

5　Kindt, K. S. *et al.* EMX2-GPR156-Galphai reverses hair cell orientation in
mechanosensory epithelia. *Nat Commun* **12**, 2861, doi:10.1038/s41467-021-22997-1
(2021).

6　Zhang, Z. & Chai, R. Hear the sounds: the role of G protein-coupled receptors in the
cochlea. *Am J Physiol Cell Physiol* **323**, C1088-C1099,
doi:10.1152/ajpcell.00453.2021 (2022).

7　Duncan, J. S. *et al.* Celsr1 coordinates the planar polarity of vestibular hair cells during
inner ear development. *Dev Biol* **423**, 126-137, doi:10.1016/j.ydbio.2017.01.020
(2017).

8　Yau, K. W. & Hardie, R. C. Phototransduction motifs and variations. *Cell* **139**, 246-264,
doi:10.1016/j.cell.2009.09.029 (2009).

9　Chen, T. Y. & Yau, K. W. Direct modulation by Ca(2+)-calmodulin of cyclic nucleotide-
activated channel of rat olfactory receptor neurons. *Nature* **368**, 545-548,
doi:10.1038/368545a0 (1994).

10　Chandrashekar, J., Hoon, M. A., Ryba, N. J. & Zuker, C. S. The receptors and cells for
mammalian taste. *Nature* **444**, 288-294, doi:10.1038/nature05401 (2006).

11　Burns, J. C., Kelly, M. C., Hoa, M., Morell, R. J. & Kelley, M. W. Single-cell RNA-
Seq resolves cellular complexity in sensory organs from the neonatal inner ear. *Nat*
Commun **6**, 8557, doi:10.1038/ncomms9557 (2015).

12　Anderson, G. R. *et al.* Postsynaptic adhesion GPCR latrophilin-2 mediates target
recognition in entorhinal-hippocampal synapse assembly. *J Cell Biol* **216**, 3831-3846,
doi:10.1083/jcb.201703042 (2017).

13　Sando, R., Jiang, X. & Sudhof, T. C. Latrophilin GPCRs direct synapse specificity by
coincident binding of FLRTs and teneurins. *Science* **363**, doi:10.1126/science.aav7969
(2019).

14　Liu, Y. *et al.* Critical role of spectrin in hearing development and deafness. *Sci Adv* **5**,
eaav7803, doi:10.1126/sciadv.aav7803 (2019).

15   Cortese, M. *et al.* Spectrin betaV adaptive mutations and changes in subcellular

location correlate with emergence of hair cell electromotility in mammalians. *Proc Natl*

Acad Sci U S A **114**, 2054-2059, doi:10.1073/pnas.1618778114 (2017).

16   Vahava, O. *et al.* Mutation in transcription factor POU4F3 associated with inherited

progressive   hearing   loss   in   humans.   *Science*   **279**,   1950-1954,

doi:10.1126/science.279.5358.1950 (1998).

17   Nakamura, T. & Gold, G. H. A cyclic nucleotide-gated conductance in olfactory

receptor cilia. *Nature* **325**, 442-444, doi:10.1038/325442a0 (1987).

18   Zhang, Y. *et al.* Coding of sweet, bitter, and umami tastes: different receptor cells

sharing   similar   signaling   pathways.   *Cell*   **112**,   293-301,   doi:10.1016/s0092-

8674(03)00071-0 (2003).

19   Arshavsky, V. & Bownds, M. D. Regulation of deactivation of photoreceptor G protein

by its target enzyme and cGMP. *Nature* **357**, 416-417, doi:10.1038/357416a0 (1992).

20   Nakatani, K. & Yau, K. W. Calcium and light adaptation in retinal rods and cones.

Nature **334**, 69-71, doi:10.1038/334069a0 (1988).

21   Taruno, A. *et al.* CALHM1 ion channel mediates purinergic neurotransmission of sweet,

bitter and umami tastes. *Nature* **495**, 223-226, doi:10.1038/nature11906 (2013).

22   Street, V. A., McKee-Johnson, J. W., Fonseca, R. C., Tempel, B. L. & Noben-Trauth,

K. Mutations in a plasma membrane Ca2+-ATPase gene cause deafness in deafwaddler

mice. *Nat Genet* **19**, 390-394, doi:10.1038/1284 (1998).

23   Beutner, D., Voets, T., Neher, E. & Moser, T. Calcium dependence of exocytosis and

endocytosis at the cochlear inner hair cell afferent synapse. *Neuron* **29**, 681-690,

doi:10.1016/s0896-6273(01)00243-4 (2001).

24   Parsons, T. D., Lenzi, D., Almers, W. & Roberts, W. M. Calcium-triggered exocytosis

and endocytosis in an isolated presynaptic cell: capacitance measurements in saccular

hair cells. *Neuron* **13**, 875-883, doi:10.1016/0896-6273(94)90253-4 (1994).

25   Craven, K. B. & Zagotta, W. N. CNG and HCN channels: two peas, one pod. *Annu Rev*

Physiol **68**, 375-401, doi:10.1146/annurev.physiol.68.040104.134728 (2006).

26   Jiang, Z., Yue, W. W. S., Chen, L., Sheng, Y. & Yau, K. W. Cyclic-Nucleotide- and

HCN-Channel-Mediated Phototransduction in Intrinsically Photosensitive Retinal

Ganglion Cells. *Cell* **175**, 652-664 e612, doi:10.1016/j.cell.2018.08.055 (2018).

27   Ping, Y. Q. *et al.* Structural basis for the tethered peptide activation of adhesion GPCRs.

Nature **604**, 763-770, doi:10.1038/s41586-022-04619-y (2022).

28   Xiao, P. *et al.* Tethered peptide activation mechanism of the adhesion GPCRs ADGRG2

and ADGRG4. *Nature* **604**, 771-778, doi:10.1038/s41586-022-04590-8 (2022).

29   Jia, Y. *et al.* TMC1 and TMC2 Proteins Are Pore-Forming Subunits of

Mechanosensitive Ion Channels. *Neuron* **105**, 310-321 e313,

doi:10.1016/j.neuron.2019.10.017 (2020).

30   Pan, B. *et al.* TMC1 Forms the Pore of Mechanosensory Transduction Channels in

Vertebrate Inner Ear Hair Cells. *Neuron* **99**, 736-753 e736,

doi:10.1016/j.neuron.2018.07.033 (2018).

31   Pan, B. *et al.* TMC1 and TMC2 are components of the mechanotransduction channel

in hair cells of the mammalian inner ear. *Neuron* **79**, 504-515,

doi:10.1016/j.neuron.2013.06.019 (2013).

32   Buck, L. & Axel, R. A novel multigene family may encode odorant receptors: a

molecular basis for odor recognition. *Cell* **65**, 175-187, doi:10.1016/0092-

8674(91)90418-x (1991).

33   Yang, F. *et al.* Structure, function and pharmacology of human itch receptor complexes.

Nature **600**, 164-169, doi:10.1038/s41586-021-04077-y (2021).

34   Wu, Z. *et al.* Mechanosensory hair cells express two molecularly distinct

mechanotransduction channels. *Nat Neurosci* **20**, 24-33, doi:10.1038/nn.4449 (2017).

35   Firestein, S., Darrow, B. & Shepherd, G. M. Activation of the sensory current in

salamander olfactory receptor neurons depends on a G protein-mediated cAMP second

messenger system. *Neuron* **6**, 825-835, doi:10.1016/0896-6273(91)90178-3 (1991).

36   Hurle, B. *et al.* Non-syndromic vestibular disorder with otoconial agenesis in

tilted/mergulhador mice caused by mutations in otopetrin 1. *Hum Mol Genet* **12**, 777-

789, doi:10.1093/hmg/ddg087 (2003).

37   Fettiplace, R., Ricci, A. J. & Hackney, C. M. Clues to the cochlear amplifier from the

turtle ear. *Trends Neurosci* **24**, 169-175, doi:10.1016/s0166-2236(00)01740-9 (2001).

38   Fettiplace, R. & Kim, K. X. The physiology of mechanoelectrical transduction channels

in hearing. *Physiol Rev* **94**, 951-986, doi:10.1152/physrev.00038.2013 (2014).

39   Michalakis, S., Becirovic, E. & Biel, M. Retinal Cyclic Nucleotide-Gated Channels:

From Pathophysiology to Therapy. *Int J Mol Sci* **19**, doi:10.3390/ijms19030749 (2018).

40   Remmer, M. H., Rastogi, N., Ranka, M. P. & Ceisler, E. J. Achromatopsia: a review.

Curr Opin Ophthalmol **26**, 333-340, doi:10.1097/ICU.0000000000000189 (2015).

41   Tan, F. *et al.* AAV-ie enables safe and efficient gene transfer to inner ear cells. *Nat*

Commun **10**, 3733, doi:10.1038/s41467-019-11687-8 (2019).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8