Higher meiotic chromosome condensation: a potential function of kinetochore through polo-

like kinase

Authors: Deepika Trakroo, Prakhar Agarwal[#], Anushka Alekar[#], and Santanu Kumar Ghosh*

Affiliation: Department of Biosciences and Bioengineering, Indian Institute of Technology,

Bombay, Powai, Mumbai-400076, India.

# Equal contribution

*Corresponding author

Email: santanughosh@iitb.ac.in

Tel. +(91-22) 2576 7766

Key words: Meiosis, chromosome, kinetochore, condensin, budding yeast

Abstract

Chromosome condensation plays a pivotal role during faithful chromosome segregation, hence understanding the factors that drive condensation is crucial to get mechanistic insight into chromosome segregation. The results of our in vivo chromosome condensation assays in budding yeast reveal that chromosomes undergo a higher degree of condensation during meiosis than in mitosis. Earlier the non-essential kinetochore proteins were shown to have several significant meiotic functions. We conclude that these proteins might also have a role in achieving higher meiotic condensation since we observed meiotic-specific condensation defects in the absence of the non-essential kinetochore protein, Ctf19. While investigating the mechanism involved, we observed an accumulation of the polo-like kinase Cdc5 owing to its higher protein stability in *ctf19Δ* meiotic cells. High Cdc5 activity leads to hyper-phosphorylation of the condensin which shows reduced stability with concomitant decreased association with the chromatin. Overall, our findings highlight the role of Ctf19 in promoting meiotic chromosome condensation by influencing the activity of Cdc5 and thereby affecting the stability and association of condensin with the chromatin.

Introduction

Faithful chromosome segregation across eukaryotes requires that the chromosomes become condensed and attached properly to the microtubule spindle during cell division. The attachment is mediated by a conserved multiprotein complex called 'kinetochore' formed at the centromeres of the chromosomes (Schrader,1939). The proteins within a kinetochore are organized in a hierarchical fashion with respect to the centromeric DNA and the microtubules. The inner and the outer proteins associate with the centromeres and microtubules, respectively whereas the central

proteins act as linker (Meraldi et al., 2006, Dong et al., 2022, Dou et al., 2023). Over the last three decades studies on kinetochore from myriad organisms have revealed that this protein complex, besides establishing a connection with the microtubules, also controls other cellular activities that facilitate high-fidelity chromosome segregation. For instance, the promotion of pericentric enrichment of cohesin complex that keeps the sister chromatids together from the S phase till anaphase (Glynn et al., 2004, Eckert et al. 2007, Holt et al., 2005, Fernius et al., 2013), early replication of the centromeres that ensures correct deposition of the centromeric histone 3 variant (Raghuraman et al., 2001, Muller et al., 2012, Natsume et al., 2013), late assembly of the kinetochore in meiosis to facilitate co-orientation of the sister kinetochores during meiosis I (Meyer et al., 2015) and homolog pairing in meiosis by causing 'centromere coupling' (Tsubouchi et al., 2005, Kim et al., 2013).

In budding and fission yeasts, the kinetochore proteins, in particular the essential inner and central proteins, are also shown to promote chromosome condensation (Kruitwagen et al., 2018, Nakazawa et al., 2008). Subsequent studies revealed that this happens through kinetochore-mediated centromeric and pericentromeric recruitment of several proteins including cohesins (Guacci et al., 1997, Lavoie et al., 2004), shugoshin (Peplowska et al., 2014, Kruitwagen et al., 2018, Yahya et al., 2020), 14-3-3 protein, Bmh1 (Jain et al., 2021) and enzymes such as aurora B kinase, Ipl1 (Giet et al., 2001, Petersen et al., 2003, Lavoie et al., 2004,), polo-like kinase, Cdc5 (St. Pierre et al., 2009, Bazile et al., 2010, Leonard et al., 2015, Lamothe et al., 2020,) and a histone deacetylase, Hst2 (Wilkins et al., 2014, Kruitwagen et al., 2015, 2018, Jain et al., 2021). However, the non-essential proteins of the Ctf19 sub-complex (Ctf19c) of the kinetochore (Mehta et al., 2022) were not shown to hinder chromosome condensation (Kruitwagen et al., 2018) as expected from their dispensability in mitotic growth. However, we have shown earlier that in absence of these non-essential proteins, meiosis is severely affected as several meiosis specific

events are perturbed (Ghosh et al., 2004, Mehta et al., 2014, Agarwal et al., 2015). This happens

as there appears to be a differential organization of the kinetochore assembly between mitosis and

meiosis (Mehta et al., 2014, Borek et al., 2021) for which the non-essential proteins perhaps

facilitate certain protein-protein and/or protein-DNA interactions in meiosis but not in mitosis.

Strikingly, earlier we observed that Ctf19 has a role to play in meiotic chromosome

condensation since the cells in the absence of Ctf19 harbor defects in rDNA condensation and

show reduced association of condensin with the chromatin in meiosis but not in mitosis (Mehta et

al., 2014). While why chromatin-condensin association is compromised in *ctf19Δ* meiotic cells is

not known, we hypothesize that to facilitate prophase I events, meiosis may require extra

chromosome condensation than mitosis and the nonessential kinetochore proteins might contribute

to regulating that. This notion is also fueled by the distinct thread-like appearance of the paired

homologous chromosomes in pachytene of meiosis I which is never visible in mitosis and unlikely

to happen in meiosis solely due to pairing.

In this report, to test the above hypothesis we sought to compare the extent of chromosome

condensation between mitosis and meiosis and to find a molecular explanation of condensation

defect in the non-essential kinetochore mutants. Interestingly, we observed using three cell

biological in vivo assays that the chromosomes are indeed more condensed in meiosis than in

mitosis. We further demonstrate that lack of Ctf19 causes misregulation of Cdc5 specifically in

meiosis, which leads to hyper-phosphorylation of condensin resulting its dissociation from the

chromatin and defect in chromosome condensation. Additionally, using wild type cells

overexpressing Cdc5, we show that the misregulation of Cdc5 is indeed the reason for condensin

failing to associate properly with the chromatin. This study highlights yet another meiosis-specific

essential contribution of the mitotically dispensable kinetochore proteins.

Results

Chromosomes are more condensed in meiosis as compared to mitosis

We hypothesize that the level and determinants of chromosome condensation might differ between mitosis and meiosis because earlier using Ctf19c mutant we observed that the kinetochore has a role in rDNA condensation in meiosis but not in mitosis (Mehta et al., 2014). However, whether such a role is confined to a specific condensed locus like rDNA or it is observed at other chromosomal locales remains unknown. To investigate this, we first sought out to compare the extent of chromosomal condensation of the kinetochore proximal pericentromeric regions between mitosis and meiosis. Diploid strains were constructed where one copy of chromosome V was marked with $[TetO]_{224}$ –[TetR-GFP] system at a locus 1.4 kb away from the *CEN V* (henceforth called *CEN V*-GFP). The fluorescence intensity of *CEN V*-GFP was used as a surrogate marker to decipher the in vivo condensation level as described before (Kruitwagen et al., 2015). In this microscopy-based assay, the fluorescence intensity of TetR-GFP bound to the $[TetO]_n$ array inserted at a decondensed locus will be more compared to when the array is inserted at a condensed locus. This is because, due to condensation, a linear shrinkage of the operators happens which brings the TetR-GFP molecules closer to each other causing quenching of the fluorescence and hence reduction in the intensity (Fig. 1A). The *CEN V*-GFP marked strains also harbored SPBs (Spc42) marked with mCherry to judge different cell cycle stages. The GFP signal intensities were measured in the cells residing at pre-S phase (interphase) stage or at metaphase/metaphase I, which is presumed to contain decondensed and condensed chromatin, respectively. An unbudded cell with a single SPB was taken as a pre-S phase cell in mitosis or meiosis; a large budded cell with two separated SPBs in the mother compartment was taken as a metaphase cell (Fig. 1B) whereas

same in an unbudded meiotic cell was considered as a metaphase I cell (Fig. 1C). The timepoint

to harvest meiotic pre-S phase cells was determined based on Ime1 (an inducer of meiosis in yeast)

expression and FACS analysis. The Ime1 expression is induced by an acetate-based rich medium

(YPA), however, it starts accumulating in the nucleus after the depletion of G1 cyclins (Colomina

et al., 1999, 2003). In our experimental condition also, Ime1 expression started in YPA followed

by its nuclear localization around 30 mins in SPM with no nuclear localization either in YPA or

hardly any in SPM at 10 mins (Fig. S1A, B). FACS analysis showed that the DNA synthesis started

before the cells were at least 90 mins in SPM (Fig. S1C). Based on these observations, 1 hr into

SPM was taken as the timepoint to harvest meiotic pre-S phase cells. For metaphase I cells, the

sample was harvested at 6 hrs in SPM, and cells with SPB-SPB distance more than 1 but less than

3 μm were analyzed as described before (Tsuchiya et al., 2014, Yu and Koshland, 2005). The

mitotic pre-S phase was determined based on Cln2 expression (yeast G1/S-CDK) as its expression

marks the beginning of mitosis (Hadwiger et al., 1989) (Fig. S1D).

The strategy followed for accurately measuring *CEN V*-GFP signal intensity normalized

with internal control (Spc42-mCherry) from four different types of samples – mitotic interphase,

metaphase, meiotic interphase, and metaphase I cells is described in materials and methods. As

expected, both in mitosis and meiosis, the *CEN V*-GFP fluorescence intensity decreases in the

metaphase and metaphase I cells, respectively than in the corresponding interphase cells

reconfirming that condensation occurs as the cells enter into the cell cycle (Lavoie et al., 2004;

Vas et al., 2007). Interestingly, while in mitosis, the extent of decrease was 30.9% (1.13 vs 0.78

A.U., Fig. 1B), the same was 47% (1.33 vs 0.7 A.U., Fig. 1C) in meiosis. The ~1.5-fold decrease

in signal intensity in meiosis over mitosis argues for an increased level of chromosome

condensation in meiosis than in mitosis. Since *CEN V*-GFP fluorescence intensity was measured

from different mitotic and meiotic cells, an error in measuring the intensity across the samples may

lead to the observed difference in fluorescence intensity between mitosis and meiosis. To rule out

this possibility we mixed the mitotic and meiotic cells and measured the fluorescence intensity

from the metaphase or metaphase I cells residing in the same field of view. The metaphase cells

with two SPBs and a bud were distinguishable from the metaphase I cells harboring 2 SPBs but

no bud. Consistent with the previous observation, we found ~1.7-fold decreases in signal intensity

in meiotic metaphase I cells compared to mitotic metaphase cells (0.825 vs. 0.485 A.U., Fig. 1D)

indicating chromosomes are indeed more condensed in meiosis than in mitosis.

The non-essential kinetochore protein Ctf19 influences chromosome condensation in

meiosis but not in mitosis

Earlier we and others have reported that the mitotically non-essential kinetochore proteins have

essential roles in meiosis (Mehta et al., 2014, Agarwal et al., 2015, Borek et al., 2021). In support

of this, we observed that such a protein Ctf19 promotes rDNA condensation in meiosis but not in

mitosis (Mehta et al., 2014). Since we revealed that the extent of chromosome condensation, at

least at the pericentromeres, is more in meiosis than in mitosis, we wished to investigate the

contribution of Ctf19 to this difference. We measured the *CEN V*-GFP signal intensity in the

diploid *ctf19Δ* mutant cells from the pre-S phase and metaphase/metaphase I in both mitosis and

meiosis. Any meiosis-specific function of Ctf19 on condensation is expected to ameliorate the

difference in condensation (or *CEN V*-GFP signal intensity) between the pre-S phase and M phase

in meiosis but not in mitosis. We observed a significant (39%) decrease in signal intensity in

metaphase cells compared to the pre-S phase cells in the *ctf19Δ* mutant (1.67 vs 1.01 A.U., Fig.

2A) like the wild type (Fig. 1B) suggesting that condensation is not compromised in the mutant

compared to the wild type in mitosis. In contrast, in meiosis, unlike the wild type which showed a

47% decrease in the intensity in meiosis (Fig. 1C), only a ~20% decrease (3.13 vs 2.4 A.U., Fig.

2B) was found in the mutant which was not statistically significant indicating that a meiosis-

specific chromosome condensation defect occurs in absence of Ctf19. Further, we performed a

comparative analysis of condensation status between wild-type and *ctf19Δ* cells across both

mitotic and meiotic metaphase stages. We found that the mean intensity value is significantly

higher in the mutant in meiosis but not in mitosis, which provides additional validation of our

claim that the condensation defect occurs in the mutant, specifically in meiosis (Figure S2) .

To examine if the observed condensation defect in the *ctf19Δ* mutant in meiosis is specific

to the Ctf19c proteins of the kinetochore, we monitored the *CEN V*-GFP signal in the pre-S phase

and metaphase I stage in the cells lacking Slk19, another non-essential protein residing at the

kinetochore but not as a component of the Ctf19c (Zeng et al., 1999). We could not observe any

chromosome condensation defect in the *slk19Δ* cells as the GFP signal intensity decreased in

metaphase I compared to the interphase stage (~50%, 1.12 vs 0.55 A.U., Fig. 2C) as found in the

wild type (Fig. 1C). Since we used identical conditions during mitosis and meiosis, the possibility

of other factors viz., accessibility of TetR-GFP to the operators or a lower concentration of TetR-

GFP responsible for reducing the fluorescence intensity, can be void. Moreover, the control mutant

slk19Δ behaved as expected.

It is possible that the quenching of the fluorophores can happen due to local compaction of

the chromatin through nucleosome-nucleosome interaction independent of condensin-mediated

condensation function (Kruitwagen et al., 2015, 2018) or the observed difference in fluorescence

intensity can be due to differential binding behaviour of TetR-GFP to the [TetO$_n$] array among the

strains and the cell cycle stages. Therefore, to correlate the observed condensation defect in the

meiotic cells of the kinetochore mutant with condensin-mediated condensation, we used an

alternative in-vivo assay to further validate the above observed condensation defect. We evaluated

the condensation status in meiosis and mitosis for both the wild type and *ctf19Δ* by using Second

Harmonic Generation (SHG) microscopy which relies on generating a second harmonic by the chromatin fibers upon excitation by two-photon microscopy (Yamin et al., 2019, 2022). In this assay, the chromosomes were stained with DAPI and a higher ratio of integrated density to area obtained through the SHG technique corresponds to higher condensation. As expected, we indeed observed an increased condensation in both wild type and *ctf19Δ* in mitotic metaphase as compared to pre-S phase (Fig. S3 A, B). However, when the meiotic cells of the same strains were analyzed, the wild type cells showed significant increase in the ratio of integrated density to area at metaphase I as compared to pre-S phase, but the *ctf19Δ* cells did not show any significant increase in the ratio in meiotic metaphase I (Fig. S3 C, D). These findings further confirm that the *ctf19Δ* cells are indeed compromised in meiotic condensation with no significant effect on mitotic condensation.

For further validation, we compared the extent of rDNA condensation using Net1-GFP as the marker as described before (Torres-Rosell et al., 2004, Machín et al, 2005, Mehta et al., 2014) among the wild type, *ctf19Δ*, and *slk19Δ* strains (Fig. 2D). The Net1-GFP signal pattern was visualized in these strains when the cells were arrested at the metaphase I stage (*pCLB2-CDC20*). While around 85% of the wild-type cells showed the 'tight knit' and 'crescent' signal which represent a condensed state of rDNA (Mehta et al., 2014; Kerr et al., 2011), *ctf19Δ* cells showed a reduction (50%) in the frequency and a corresponding increase (50%) in the occurrence of the 'puff' signal which corroborates a decondensed state. On the other hand, *slk19Δ* cells showed similar results to the wild type. In mitosis, however, there was no difference in condensation amongst the wild type, *ctf19Δ,* and *slk19Δ* cells (Fig. S4) where the 'line', 'loop', and 'puff' patterns designate condensed, intermediate, and decondensed forms of the rDNA locus, respectively (D'Ambrosio et al., 2008; Dauban et al., 2019). Altogether, from these results, we infer that Ctf19 promotes chromosome condensation in meiosis but not in mitosis and this function

209   appears to be specific for the Ctf19 protein or perhaps for all the non-essential proteins of the

210   Ctf19c of the kinetochore.

Loss of Ctf19 affects axial condensation of the chromosomal arm

The loss of pericentric and rDNA condensation in the *ctf19Δ* mutant prompted us to examine the

mutant for linear (axial) condensation of the chromosomal arms which relies on condensin

(Kruitwagen et al., 2015). To investigate this, we first wished to test the condensation status of a

locus proximal to the telomere V (*TEL V*) as compared to *CEN V*-GFP. For this, we marked

chromosomes at two different loci 400 kb away from each other, one was near the *CEN V* (1.4 kb

away), as before, and the other one was 21 kb away from *TEL V* using $[TetO]_{224}$ –[TetR-GFP] and

$[TetO]_{448}$–[TetR-GFP] systems, respectively. Since the number of operators used to visualize *TEL*

V-GFP was double that of *CEN V*-GFP, the two loci could be visually distinguishable by the

brightness of the GFP dots where *TEL* V-GFP appeared much brighter than the *CEN V*-GFP (Fig.

3A). The GFP fluorescence intensities were measured from *CEN V* and *TEL V* dots both in the

wild type and *ctf19Δ* mutant cells from pre-S phase and metaphase I stages (Fig. 3B-3C). Both

CEN V-GFP and *TEL V*-GFP dots showed equivalent (*CEN V*= 33.3% decrease; *TEL V*= 25%

decrease) and a statistically significant decrease in GFP intensity in metaphase I cells compared to

pre-S phase cells in the wild type. The difference in *CEN V*-GFP intensity between here to what

has been described in the previous section (33.3% vs 47%) is possibly because of the difference in

the intensity of Spc42-mCherry which was used as an internal control for normalization purposes.

Nevertheless, in contrast to the wild type, in the *ctf19Δ* mutant the decreases in *CEN V* and *TEL V*

intensities in metaphase I cells compared to pre-S phase cells were 15.6% and 12.5%, respectively

which were not statistically significant. These results confirm that Ctf19 is important for the

condensation of the region proximal to centromeres or telomeres.

Next, we wished to examine the axial condensation of the chromosome V arm by

measuring the distance between the *CEN V*-GFP (light) and *TEL V*-GFP (bright) dots.   In the wild

type, we observed a 37% decrease in the mean distance in the metaphase I stage (0.5 µm) compared

to the pre-S phase stage (0.8 µm) owing to the condensation (Fig. 3D, left panel). However, in the

ctf19Δ mutant, there was no significant decrease in the mean distances between the two stages (0.9

vs 1.08 µm) suggesting a defect in the condensation (Fig. 3D, right panel).   While looking at the

axial condensation in mitosis, there was a statistically significant decrease in the distance in

metaphase compared to the pre-S phase both in the wild type and the mutant (21% and 24 %,

respectively) (Fig. S5). Notably, since the mean distance in the mutant (1.08 µm) was found higher

than the wild type (0.8 µm) in the meiotic pre-S phase that precedes condensin function, it is

possible that at this phase the compaction of the chromatin is compromised in the mutant due to

relaxed nucleosome-nucleosome interaction. Altogether, these results suggest that the contribution

of Ctf19 in promoting chromatin condensation in meiosis is not only restricted to the

pericentromeric regions but also extends axially along the chromosome arms at the distal loci.

Ctf19 is required for maintenance of chromatin-condensin association in meiosis.

Earlier we reported that in absence of Ctf19, the chromatin-condensin association is perturbed in

meiosis (Mehta et al., 2014) which supports our findings from the condensation assays described

here. We wished to re-verify our earlier observation with further investigation to find whether the

effect on chromatin-condensin association is specific to Ctf19c proteins and whether the effect is

on the establishment and/or maintenance of the association. To address these, we first compared

the localization of the condensin protein Brn1 on the chromatin between the wild type and *ctf19Δ*

mutant along with the *slk19Δ* mutant which, as control, showed no effect on condensation (Fig.

2C and 2D). Brn1 fused to 6XHA at the C-terminal (Verzijlbergen et al., 2014) was detected on

the chromatin in the meiotic cells arrested at metaphase I by Cdc20 depletion. While a similar

number (~ 13 to 15) of bright Brn1-6HA foci were visible in the wild type and *slk19Δ* mutant, the

number of foci was much lower (~7) and the foci were less bright in the *ctf19Δ* mutant suggesting

a defect in chromatin-condensin association in the latter (Fig. S6 A). On the other hand, as expected

from the condensation assay results, no difference in Brn1-6HA localization was observed between

the wild type and *ctf19Δ* spreads prepared from the cells arrested at mitotic metaphase by Cdc20

destruction as mentioned in materials and methods (Fig. S6 B, C). To test if the observed

condensation defect is specific to Ctf19 or also occurs in other non-essential proteins of Ctf19c,

we analyzed the association of Brn1-6HA with chromatin in cells lacking another protein of

Ctf19c, Ctf3. We found that in *ctf3Δ* cells also, the number of Brn1-6HA foci was lesser (~4) than

the wild type (~9) (Fig. S7). The lesser number of foci here in the wild type compared to the

previous chromatin spread of Brn1-6HA in the wild type (Fig. S6) can be because of technical

difference in antibody staining, as these experiments were performed at different times

To further validate our chromatin spread data from meiosis, we performed a ChIP assay to

quantify the association of Brn1 with different regions of chromosome - centromere,

pericentromere, and arm (Fig. 4A). The cells were harvested after 8 and 10 hrs into SPM to obtain

metaphase I arrested wild type and *ctf19Δ* mutant, respectively by Cdc20 depletion. We observed

a significant reduction in the level of Brn1 at all the regions in the mutant compared to the wild

type; however, the reduction was more at the centromeric region compared to other regions (Fig.

4B). These results, re-confirming our earlier chromatin spread data (Mehta et al., 2014), further

indicate that the meiotic defect in chromatin-condensin association is specific to the lack of Ctf19c

proteins.

Next, we wished to address whether Ctf19 is required for the establishment or maintenance

of chromatin-condensin association. It has been previously shown that during early meiosis,

condensin first associates with rDNA as a cluster and then at later time points associates with the

rest of the chromatin leaving a minuscule amount at the rDNA (Markowitz et al., 2020; Li et al.,

2014). To compare the dynamics of chromatin-condensin association between wild type and

ctf19Δ we monitored the association of Brn1-6HA with chromatin using chromatin spreads at

different time points in the cells released into synchronized meiosis (Fig. 4C). Types 1 – 3 are

categorized as localization of Brn1-6HA first at rDNA (type 1) and then gradually spreading to

rest of the chromatin (types 2 – 3). Type 4 implies a gross reduction in chromatin-condensin

association. While the initial rDNA localization of Brn1-6HA (0 hr, type 1) in the mutant was not

found lesser than the wild type, its gross chromatin localization (type 3) in the mutant was merely

below 10% of the spreads compared to the wild type (40%) at 4 hrs. This implies that the

establishment of condensin association with the gross chromatin, but not with the rDNA, is

perturbed in the absence of Ctf19. Consequently, at 8 hrs, while the mutant showed a major

population of spreads (70%) with reduced condensin on the chromatin (type 4), only a minor

population (<20%) of wild type spreads demonstrated this phenotype. Moreover, as the type1

population decreased to 75% in the mutant compared to 50% in the wild type at 6 hrs, this

observation may be interpreted as a defect in the maintenance of the rDNA association of

condensin in the mutant.

The condensation defect in *ctf19Δ* may occur partially through Sgo1 mislocalization but

independent of DDK- and Ipl1-mediated pathways

Next, we wished to address the mechanism through which Ctf19 may regulate condensation in

meiosis. Besides protecting the pericentromeric cohesin in meiosis I, Sgo1 (Shugoshin) has a role

in mitotic chromosome condensation (Kruitwagen et al., 2018) through condensin loading at

pericentromeric region (Peplowska et al., 2014). It has also been reported that in the absence of

Ctf19c proteins, Iml3 and Chl4, the Sgo1 localization becomes compromised at the chromatin both

in mitosis (Deng et al., 2018) and meiosis (Kiburz et al., 2005). To investigate if Ctf19, unlike the

peripheral Ctf19c components Iml3 and Chl4, differs in recruiting Sgo1 on the chromatin between

mitosis and meiosis which may cause the differential effect on condensation, we analyzed the

localization of Sgo1-EGFP in mitosis (metaphase) and meiosis (metaphase I) in presence or

absence of Ctf19. We observed that the absence of Ctf19 affected Sgo1 localization both in mitosis

(Fig. 5A) and meiosis (Fig. 5B), however, the extent of the defect was more in meiosis where 79%

of the cells showed Sgo1-EGFP mislocalization (no signal) compared to only 53% in mitosis.

Whereas the fraction of cells showing no signal of Sgo1-EGP in the wild type was found similar

in mitosis and meiosis (Fig. 5A, B). To further validate our cell biological data, we performed a

ChIP assay to quantify the association of Sgo1-9Myc with the centromeres. We observed a

significant reduction in the level of Sgo1-9Myc at the centromeres both in mitosis (Fig. 5C) and

meiosis (Fig. 5D) in *ctf19Δ* mutant compared to the wild type. However, the reduction was more

(~80% for both *CEN III* and *CEN IV*) in meiosis than in mitosis (~ 45% for *CEN III* and ~ 40%

for *CEN IV*) (Fig. 5C, D; see materials and methods). Based on these observations, we conclude

that a higher reduction in Sgo1's association with the chromatin in meiosis over mitosis in *ctf19Δ*

cells may contribute partially, if at all, to the meiosis-specific chromatin-condensin association

defect in these cells.

It has been reported earlier that till metaphase the rDNA condensation in mitosis is

dependent on cohesin (Lavoie *et al*., 2002). As the cohesin loading at the pericentromeres is

dependent on Ctf19 through recruitment of DDK (Dbf4-dependent kinase) both in mitosis and

meiosis (Hinshaw et al., 2015, 2017), therefore, the reduction of cohesin level at the centromeres

in *ctf19Δ* mutant is unlikely the cause for observed meiosis-specific condensation defect. However,

recently DDK and Cdc5 (budding yeast polo-like kinase) have been reported to have roles in the

meiotic prophase-like pathway (cleavage-independent cohesin removal from the chromatin) which

has been shown important for chromosome compaction in meiosis probably by making space on

chromatin for condensin loading (Challa et al., 2019). Based on this, we hypothesized that perhaps

DDK kinase localization at the centromeres is perturbed in the absence of Ctf19 which in turn

results in an impaired prophase-like pathway hence leading to chromosome condensation defect.

If our hypothesis is correct, then a DDK mutant unable to be localized at the centromeres should

show a condensation defect in meiosis. It has been reported that the fusion of epitopes at the C-

terminal of the Dbf4 protein renders it inactive at the centromeres however its localization at the

origins of replications remains unaffected (Hinshaw et al., 2017). We constructed a strain

expressing Dbf4-9myc with heterozygous tagging of *CEN V* and *TEL V* with GFP. We confirmed

that the *DBF4-9MYC* allele is acting as a mutant as we observed a reduced frequency of sister

chromatid cohesion (Fig. S8A) as reported earlier (Hinshaw et al., 2015). However, we failed to

observe any defect either in axial contraction by measuring the distance between *CEN V* and *TEL*

V (Fig. S8B) or in condensation of centromeric and telomeric regions measured by fluorescence-

based quenching assay (Fig. S8C). Based on these observations we conclude that the absence of

DDK at the centromeres in *ctf19Δ* cells is not responsible for the observed condensation defect in

these cells in meiosis.

Phosphorylation of histone H3 at serine 10 by Aurora kinase B (Ipl1 in budding yeast), has

been known to play a crucial role in chromosome compaction both in yeast (Kruitwagen et al.,

2015) and mammals (Mora-Bermúdez et al., 2007). Additionally, an in vitro assay has

demonstrated an interaction between Ctf19 and Ipl1, which is essential for stabilizing the

chromosomal passenger complex (CPC) at the centromeres (Fischböck-Halwachs et al., 2019).

However, it is uncertain whether Ctf19 interacts with Ipl1 in vivo and regulates Ipl1's centromeric

localization and possibly thereby promotes chromosome condensation in meiosis. To address this

question, we performed a ChIP assay to quantify the association of Ipl1 at the centromeres in

metaphase I arrested wild-type and *ctf19Δ* mutant cells, harvested at 8 and 10 hours into SPM,

respectively. However, our analysis did not reveal any significant reduction in the centromeric

enrichment of Ipl1 in the *ctf19Δ* mutant compared to the wild type (Fig. S9). Based on these

findings, we conclude that Ctf19 does not regulate Ipl1 localization at the centromeres, at least

during meiosis and hence Ctf19's involvement in chromosome condensation is independent of the

Ipl1-mediated pathway.

Polo-like kinase (Cdc5) level increases in the cells in the absence of Ctf19 in meiosis

Having shown that DDK and Ipl1 kinases are not involved, we wished to identify other factor(s)

that may promote meiotic condensation via Ctf19. A kinase switch model for chromosome

condensation says that different kinases take turns to regulate the condensation at different stages

of the cell cycle (Bazile et al., 2010). For instance, casein kinase II, cyclin-dependent kinase, and

polo-like kinase regulate condensation at the interphase, metaphase, and anaphase, respectively.

Recently, in budding yeast, the polo-like kinase Cdc5 has been shown to be important for both

condensation establishment and maintenance in mitosis (Lamothe et al., 2020). Furthermore,

earlier we reported that in meiosis, Ctf19c protein mutants show SPB misduplication and

supernumerary SPBs (Agarwal et al., 2015), a phenotype also noticed in meiotic cells

overexpressing Cdc5 (Shirk et al., 2011). Therefore, we wished to examine if there is any meiosis-

specific misregulation of Cdc5 in the *ctf19Δ* cells which might cause the condensation defect.

Surprisingly, we observed a higher level of Cdc5 in the *ctf19Δ* cells in meiosis but not in mitosis

when the cells were arrested at metaphase I and metaphase, respectively (Fig. 6A and Fig. S10A).

To investigate if the high level is due to increased stability of Cdc5 and/or high expression of the

corresponding gene, we first detected the protein level of Cdc5-6HA over time in meiosis after

blocking the protein synthesis using cycloheximide (CHX) which was added at 8 hrs and 10 hrs

timepoints when the wild type and mutant cells were into SPM medium, respectively. We found

that Cdc5 is more stable in the mutant compared to the wild type (Fig. 6B). The difference between

the slopes of the lines was significant as calculated using GraphPad Prism 8.0.1 ($p = 0.009$).

However, we failed to observe any increased stability of Cdc5 in the mutant in mitosis (Fig. S10B).

Moreover, there was no increased expression of the *CDC5* gene observed in the mutant (Fig.

S10C). From these results, we conclude that perhaps increased stability and a consequent higher

level of Cdc5 in *ctf19Δ* cells in meiosis cause the condensation defect.

Despite the high level of Cdc5, its association with chromatin was reduced in *ctf19Δ* mutant

compared to the wild type as observed by chromatin spread in meiosis (Fig. 6C). Interestingly,

more spreads with reduced Cdc5 (type II in 6C) were found in meiosis compared to mitosis in the

mutant (Fig. 6D). The reduction in Cdc5-chromatin association can be possible because of reduced

cohesin association with chromatin in the absence of Ctf19 (Mehta et al., 2014, Hinshaw et al.,

2017). In the absence of cohesin, the chromatin localization of Cdc5 is compromised and its

activity becomes somehow misregulated causing hyper-phosphorylation of condensin subunits

leading to condensation defect (Lamothe et al., 2020). Therefore, we hypothesized that the

increased level of chromatin unbound Cdc5 in *ctf19Δ* meiotic cells may misregulate the condensin

phosphorylation status resulting in a reduced association of condensin with the chromatin with

consequent condensation defect (Fig. 4). To test this hypothesis, we monitored the phosphorylation

status of the condensin protein Ycg1, as it is an essential subunit for condensin complex

recruitment to chromatin (Piazza et al., 2014) and is known to be phosphorylated at multiple sites

(St-Pierre et al., 2009). We monitored Ycg1-6HA expression pattern over different time points in

398 cells released synchronously into meiosis till metaphase I arrest (Fig. 7A). At later time points (8

399 hrs and 10 hrs) in SPM, which corresponded to the metaphase I arrested stage, two bands of higher

400 (B1) and lower (B2) mobilities were visible while at early time points (0 and 6 hrs) only one band

401 was visible both in wild type and mutant indicating post-translational modification of Ycg1-6HA

402 as reported earlier (St-Pierre et al., 2009). Since the B1 bands at 6 hrs are of higher intensity

403 compared to the bands at 0 hr, we speculated that 0 and 6 hrs bands harbor unmodified and

404 (unmodified + modified) forms of Ycg1-6HA, respectively. However, it was interesting to note

405 that in the mutant the top band B2 was reproducibly of higher intensity compared to the wild type.

406 We speculated that the B2 band depicts a higher-level modification of Ycg1-6HA and was found

407 more in the mutant compared to the wild type. Treatment of the protein extract from metaphase I

408 arrested cells with lambda phosphatase showed that the modification is phosphorylation (Fig. 7B).

409 A better resolution of the protein bands using Phos-tag gel revealed that B1 band indeed contained

410 unphosphorylated and phosphorylated forms of Ycg1-6HA whereas B2 band harbored hyper-

411 phosphorylated Ycg1-6HA (Fig. 7B). These results suggest that hyper-phosphorylation of

412 condensin protein in the mutant possibly occurs due to excess Cdc5 level in the mutant as observed

413 earlier.

414 It is possible that the hyper-phosphorylated form of Ycg1 in the mutant is less stable which

415 impedes the chromatin-condensin association in the mutant, as observed for Brn1 of the condensin

416 complex (Fig. 4). To verify this, we monitored the Ycg1 level in the CHX chase experiment where

417 the protein synthesis was blocked by adding the drug at 8 hrs and 10 hrs in SPM for the wild type

418 and the mutant, respectively; the samples were harvested at the indicated time points (Fig. 7C).

419 We found that Ycg1 is indeed less stable in the mutant compared to the wild type. The difference

420 between the slopes of the lines depicting the Ycg1 level was also significant as calculated using

421 GraphPad Prism 8.0.1 ($p = 0.03$). We also tested whether the Ycg1 stability was also impaired in

the *ctf19Δ* cells in mitosis. For that, we first arrested both the wild type and *ctf19Δ* cells in

metaphase by nocodazole (Fig. S11A) and then added CHX in the media to monitor Ycg1 stability.

We found no difference in the Ycg1 stability in the wild type and *ctf19Δ* cells (Fig. S11B, C),

indicating that Ctf19 influences the stability specifically in meiosis. From our results, we conclude

that higher stability of Cdc5 causes accumulation of the protein in *ctf19Δ* meiotic cells, which

results in hyper-phosphorylation of condensin. This hyper-phosphorylated condensin is less stable

and possibly why its association with the chromatin decreases (Fig. 4, S12).

429   Misregulation of Cdc5 influences chromatin-condensin association

If excess Cdc5 is responsible for reduced chromatin-condensin association in *ctf19Δ* meiotic cells,

then overexpression of Cdc5 in the wild type cells should emulate that, whereas its depletion

should rescue the defect in the mutant. The overexpression of Cdc5 in the wild type was achieved

by shuffling the endogenous promoter of *CDC5* with the *CUP1* promoter. We verified the

overexpression as a phenotype of metaphase I delay as reported earlier (Shirk et al., 2011, Li et

al., 2014) when the cells were treated with 50 μM CuSO$_4$ in SPM against a normal meiotic

progression at 10 μm CuSO$_4$ (Fig S13A, B). The cells were released into SPM either in 10 or in

50 μm CuSO$_4$ and were harvested at 7 hr (Fig. 8A), since this time point produced maximum

number of metaphase I cells while treated with 50 μm CuSO$_4$ (Fig. S13B). In the harvested cells,

when we visualized the level of chromatin-condensin association using chromatin spreads, the

normal level of Cdc5 (10 μm CuSO$_4$) produced 11 to 13 bright Brn1-6HA foci, while with excess

Cdc5 (50 μm CuSO$_4$), the number was much lower (~7) (Fig. 8B, C, S13 C) indicating a defect in

the association which is consistent to an earlier report (Li et al., 2014).

Next, to examine if the defect in chromatin-condensin association in the *ctf19Δ* meiotic

cells can be rescued by depleting Cdc5, we shuffled the *CDC5* promoter with *CUP1* promoter in

those cells and followed the above strategy except used 0 μm CuSO$_4$ instead of 50 μm (Fig. 8D).

We verified that at 7-8 hrs of SPM release, maximum number of metaphase I cells were present

(Fig. S13D, E). With normal level of Cdc5 (10 μm CuSO$_4$), we observed around 6 Brn1-6HA foci,

while with no Cdc5 expression (0 μm CuSO$_4$), we could see a significantly higher number of Brn1-

6HA foci (~10) (Fig. 8E, F, S13 F) indicating a clear amelioration of the defect. Taken together,

the above results suggest that the observed meiosis specific condensation defect in absence of

Ctf19 is indeed due to accumulation of Cdc5 in the cells.

Discussion

For the last few decades, the kinetochore-centromere has emerged as a controlling hub for various

events of chromosome segregation viz., kinetochore-microtubule attachment, early replication of

centromeres, pericentromeric cohesin enrichment, homolog pairing, and chromosome

condensation. In budding yeast, the essential kinetochore proteins have been shown to initiate

chromosome condensation in *cis* which is believed to extend distal to the centromeres through

Sgo1 and a histone deacetylase, Hst2 (Kruitwagen et al., 2018). On the other hand, through a series

of studies we and others have demonstrated that the organizational assembly of the kinetochore

complex differs between mitosis and meiosis and possibly due to this difference it was found that

the mitotically non-essential kinetochore proteins play significant roles in meiosis; consequently,

in their absence, meiosis is very poor but not mitosis (Mehta et al., 2014, Aggarwal et al., 2015,

Borek et al., 2021). In this work, we surprisingly found that the level of chromosome condensation

is more in meiosis than in mitosis and the non-essential kinetochore proteins play a role in the

regulation of the meiotic condensation through modulation of polo-like kinase Cdc5.

Chromosome condensation: mitosis vs meiosis

Regardless of the mode of the cell cycle, chromosomes need to be condensed as they embark into

partitioning. However, there are fundamental differences in how the chromosomes behave as they

enter into mitosis or meiosis (meiosis I, in particular). One of the major differences is the pairing

of the homolog in meiosis which is believed to start at the centromeres by 'centromere coupling'

(Tsubouchi et al., 2005) and proceeds towards the telomeres in a zipper-like fashion. It is obvious

that finding the right partners and the completion of lengthwise pairing will be more efficient if

the units undergoing pairing are shorter and more condensed. Moreover, the post-pairing events

including chiasmata formation and resolution of the chiasmata by 'terminalization' of the cross-

over point are also presumably favored by more condensed chromosomes. Notably, in pachytene

during meiosis I, the paired homologs are individually visible as thread-like structures by whole

chromatin staining using DAPI even in small yeast nuclear spreads. However, although the

homologs never pair in mitosis, they never appear as thread-like structures even with half the DAPI

intensity of the pachytene spreads. By using two different methods of in vivo condensation assays,

we observed that in budding yeast meiotic chromosomes are indeed more condensed than the

equivalent mitotic stage chromosomes (Fig. 1, 3D, S5A). However, since we observed a greater

decrease in intensity of GFP foci and distance between them, as readout for chromatin's local

compaction and global contraction, respectively, in meiosis, both nucleosome-nucleosome

interaction and condensin function may culminate into higher meiosis-specific condensation.

Evidence of more chromosome condensation in meiosis is available in nature. For instance, in Rye

(*Secale cereale*) it is reported that chromosomes are compacted in meiosis by a factor of 1.7 in

length and 1.3 in diameter more compared to mitosis (Zoller et al., 2004). Notably, we have also

found 1.7 times more condensation in meiosis compared to mitosis by in vivo assay in budding

yeast (Fig. 1D) indicating a similar mechanism across eukaryotes might exist to achieve more

chromosome condensation in meiosis.

Kinetochore proteins controlling chromosome condensation

Chromosome condensation can be divided into two stages viz., establishment and maintenance, which are regulated by kinases. As the cells enter into the cell cycle, the condensin becomes active and condensation is established through CDKs (Kimura et al., 1998, Sutani et al., 1999). After the metaphase to anaphase transition, as the CDK level starts declining, aurora B kinase and polo-like kinases phosphorylate condensin subunits and thus maintain the condensation (Lavoie et al., 2004, Takemoto et al., 2007, St. Pierre et al., 2009, Walters et al., 2014, Kruitwagen et al., 2018). In budding yeast mitosis, the essential inner kinetochore proteins promote condensation at the centromeres/pericentromeres by recruiting Ipl1 (aurora B) and Cdc5 (polo-like) kinases at the centromeres (Lavoie et al., 2004, St. Pierre et al., 2009). In contrast, the outer essential and the inner non-essential kinetochore proteins were found to be dispensable for condensation (Nakazawa et al., 2008, Kruitwagen et al., 2018) possibly because they do not regulate chromatin localization of Ipl1 and Cdc5. However, given our earlier study reporting meiosis specific reduction in chromatin-condensin association in absence of the non-essential kinetochore protein Ctf19 (Mehta et al., 2014), in this study we show that the effect is specific to the lack of Ctf19c proteins and it is the establishment of the association which is compromised.

Since our and other studies showed reduced pericentric cohesin enrichment in *ctf19Δ* cells (Mehta et al., 2014, Fernius and Marston 2009), it can be argued that the observed reduction in condensin-chromatin association in *ctf19Δ* meiotic cells is merely due to pericentric cohesin enrichment defect. This can happen provided i) compared to mitosis, in meiosis, the arm cohesin is sufficiently low due to the meiosis-specific prophase-like pathway of cohesin removal (Challa et al., 2019), which becomes aggravated due to simultaneous loss of cohesin from the preicentromeres and ii) the presence of cohesin must promote condensin binding. But we

categorically negate the above possibilities because i) half of the arm cohesin still remains after

cohesin removal by a prophase-like pathway (Challa et al., 2019) although earlier works failed to

detect any significant difference in arm cohesin between mitosis and meiosis (Glynn et al., 2004,

Fernius and Marston 2009) and ii) it was shown that cohesin does not influence condensin binding

using biochemical and cell biological assays (Lamothe et al., 2020, Lavoie et al., 2002).

Condensation distal from the centromeres can be influenced by non-essential kinetochore

proteins through Bub1 and DDK kinases mediated recruitment of Sgo1 (Fernius et al., 2007,

Peplowska et al., 2014) and cohesin (Lavoie et al., 2004, Natsume et al., 2013, Hinshaw et al.,

2015), respectively, at the pericentromeres which is then believed to extend towards the telomeres

through functions of Sgo1 and Hst2 (Kruitwagen et al., 2018). Therefore, we tested whether Sgo1's

centromeric localization and DDK activity are further down in meiosis in *ctf19Δ* cells to explain

meiosis-specific condensation defects in these cells. We failed to see any effect on condensation

in DDK-compromised cells (Fig. S8A, B and C). However, a higher reduction of Sgo1 level at the

pericentromeres in meiosis than in mitosis in *ctf19Δ* cells (Fig. 5) argued this as a possible cause

of meiosis-specific condensation defect in these cells. Additionally, we observed that the

centromeric association of Ipl1 kinase remains unaffected in *ctf19Δ* cells, indicating that Ctf19

does not play a regulatory role in condensation through the Ipl1-mediated pathway (Fig. S9).

Further studies surprisingly revealed that it is the higher accumulation of Cdc5 in the *ctf19Δ*

meiotic cells that perturbs chromatin-condensin association as discussed below.

Interaction of kinetochore with polo-like kinase

In budding yeast, the polo-like kinase Cdc5 is targeted to the centromeres through interaction with

the kinetochore proteins in mitosis (Snead et al., 2007, Mishra et al., 2019). It is also shown that

pericentromeric cohesin enriches Cdc5 at the centromeres which in turn promotes cohesin removal

(Mishra et al., 2016). Therefore, as *ctf19Δ* cells show reduced pericentric cohesin enrichment (Fernius et al., 2009, Mehta et al., 2014), it is expected that the association of Cdc5 with the chromatin would be less and that is what we observed (Fig. 6C). But more reduction of Cdc5-chromatin association in meiosis than in mitosis (Fig. 6D) may account for a greater effect of pericentric cohesin loss in meiosis in *ctf19Δ* cells. Earlier it was reported that Cdc5 dislodged from the chromatin, due to global cohesin inactivation, hyper-phosphorylates condensin causing their misfolding but the misfolded and non-functional condensin remains associated with the chromatin (Lamothe et al., 2020). We also noticed hyper-phosphorylation of condensin (Fig. 7A, B), but in contrast observed a huge drop in chromatin-condensin-association in the spreads made from *ctf19Δ* cells (Fig. 4, S6A, S12), possibly because the hyper-phosphorylated condensin generated in our conditions is less stable (Fig. 7C). Notably, while Lamothe et al., 2020, could not find the difference in total Cdc5 level between wild type and cohesin-depleted cells, we observed a substantial increase in Cdc5 level in *ctf19Δ* cells compared to the wild type. That might have caused a different type of hyper-phosphorylation of condensin in our cells compared to their conditions, leading to the dissociation of condensin from the chromatin. Therefore, we conclude that in *ctf19Δ* cells, the meiosis-specific altered behavior of Cdc5 is not only due to pericentric cohesin reduction but also due to its higher stability (Fig. 6B). Currently, it is not known that how the loss of Ctf19 leads to a higher stability of Cdc5 in meiosis. Cdc5 has meiosis-specific roles including degradation of Pds1 in meiosis I (Attner et al., 2013), mono-orientation of sister chromatids by regulating the association of Mam1 protein with kinetochores, step-wise cohesin removal (Lee and Amon, 2003) and dissolution of homolog linkage (Yu and Koshland, 2005). Moreover, unlike mitosis, in meiosis, Cdc5 is degraded by APC associated with meiosis-specific co-factor Ama1 (Okaz et al., 2012). Therefore, a meiosis-specific protein-protein interaction involving Cdc5 is expected. In this context, given a difference in the assembly of kinetochore

exists between mitosis and meiosis (Mehta et al., 2014, Borek et al., 2021), the loss of Ctf19 and

possibly other non-essential proteins of Ctf19c might create a condition that disfavors normal

degradation of Cdc5. One possibility is that in the *ctf19Δ* mutant cells the activity of Mnd2, an

antagonist to Ama1 (Oelschlaegel et al., 2005), increases as Mnd2 is believed to physically interact

with Ctf19 (Wong et al., 2007). It will be interesting to investigate the status of Mnd2 and Ama1

in the *ctf19Δ* mutant.

Overall, in this work, we reveal that the chromosomes are more condensed during meiosis

than in mitosis which has implications for understanding the complex meiotic chromosome

behavior in mammals where the gametogenesis is largely asymmetric and mechanistically varies

profoundly between the sexes. In animal cells, there is no evidence of kinetochore inactivation

affecting chromosome condensation in meiosis. Since we observed that the non-essential proteins

affect condensation in meiosis but not in mitosis, it would be intriguing to examine the

condensation status in animal meiotic cells harboring mutations in kinetochore. Finally, the

interaction between kinetochore integrity and the functionality of polo-like kinase (Plk) can be

explored in animal cells as improper polo-like kinase activity has been linked to aneuploidy and

different disease states in humans viz., elevated Plk level has been reported in several types of

cancers and neurological disorders (Takai et al., 2005, Ackermann et al., 2011, Korns et al., 2022).

Material and methods

Yeast strains

All the strains utilized in this work were derived from the SK1 background and are mentioned in

Table S1. Chromosome V was fluorescently marked near *CEN V* and *TEL V* using [TetO]$_n$ –[TetR-

GFP] system in strains expressing TetR-GFP. An array of 224 copies of the TetO sequence was

inserted at locus 1.4 kb away from *CEN V* through homologous recombination (Michaelis et al.,

1997, Tanaka et al., 2000). Similarly, [TetO]$_{448}$ array was inserted near *TEL V* between *BMH1* and *PDA1* genes (Kumar et al., 2021). To perform gene deletion, promoter shuffling, and C-terminal tagging of proteins, a PCR-based method was used (Wach, 1996, Longtine et al.,1998, Janke et al., 2004). Relevant plasmid-borne cassettes procured from Euroscarf were employed for these techniques. Yeast transformation was performed as described previously (Guldener et al., 1996).

The cells were synchronized in the G1 phase in the YPA medium and then released in YPD and SPM for mitotic and meiotic sample collection, respectively. For the mitotic pre-S phase, the cells were harvested at 45 mins post-release in YPD which corresponds with the start of Cln2 expression. The cells were identified microscopically by unbudded cells with one SPB focus. For metaphase, the cells were harvested after 1.5 hrs post-release in YPD and distinguished microscopically by large-budded cells with two SPB foci in the mother bud. For meiosis, the cells were harvested at 1 hr and 6 hrs post-release in SPM for pre-S phase and metaphase I cell, respectively.

Meiotic synchronization

Meiotic induction was carried out as described elsewhere (Cha et al., 2000). Typically, the diploid strain of interest was streaked onto a YPD plate (1% yeast extract, 2% peptone, 2% dextrose, 2% agar) for a single colony followed by 48 hrs of incubation. A single colony was then inoculated into 5 ml of YPD broth and left to incubate overnight at 30°C for ~15 hrs. Subsequently, cells from the overnight grown culture were transferred to a 50 ml YPA medium (1% yeast extract, 2% peptone, and 1% potassium acetate) at 0.2 OD$_{600}$ and grown till ~1.4-1.6 OD$_{600}$ which took approximately 13-15 hrs. 1 ml of YPA culture was taken, the cells were diluted appropriately and subjected to sonication at 50% amplitude for 3 secs and observed under microscope for G1 arrest.

Once over 85% of G1 arrest was achieved, the cells were washed with sterile distilled water and

resuspended in 50 ml of sporulation media (SPM; 0.3% potassium acetate and 0.02% raffinose).

Meiotic synchronization was performed at 30°C with continuous shaking at 200 RPM. Samples

were harvested from the SPM at specified time intervals for further experimentation. For

metaphase I arrest, the cells harboring *CDC20* promoter shuffled with mitotic *CLB2* promoter (Lee

and Amon, 2003), were harvested at 8 hrs and 10 hrs for wild type and *ctf19Δ* mutant, respectively.

The arrest of yeast strains at mitotic metaphase

Using the auxin-induced degron system

Mitotic metaphase arrest was achieved by degrading Cdc20 using the AID degron (Auxin-

Inducible Degron) system (Nishimura et al., 2009). In cells expressing *OsTIR1* from the *ADH1*

promoter, the AID coding sequence was C-terminally fused to *CDC20*. Cells were grown in YPD

from 0.2 to 0.5 OD_{600} before auxin (3-Indoleacetic acid, Sigma) was added at 1.5 mM

concentration and the culture was incubated for a further 2 hrs at 30⁰C. Metaphase arrest was

confirmed by DAPI staining (more than 85% large budded cells with undivided DAPI).

Using nocodazole, a microtubule depolymerizing drug

The cells were inoculated in YPD at 0.2 OD_{600} and grown till 0.5-0.6 OD_{600} before nocodazole (20

µg/ml, Sigma) was added to the culture which was incubated for a further 2 hrs under shaking

conditions at 30⁰C. Metaphase arrest was confirmed by DAPI staining.

Fluorescence microscopy

The cells were grown in YPD or SPM medium for mitotic or meiotic culture, respectively. The

cells were harvested and fixed with 4% formaldehyde for 5 mins and then washed twice with 0.1

M phosphate buffer before imaging. All fluorescence images were acquired with a Zeiss Axio

Observer.Z1 (63X objective, NA = 1.40; Carl Zeiss Micro-Imaging Inc.) equipped with Axiocam camera. The images were processed and analyzed using Zeiss Zen 3.1 software. Image J and Imaris (8.0.2) software were used for intensity and distance measurements, respectively. For all the box plots for the intensity and distance measurements, mean values were compared across the samples.

Intensity measurement

For intensity calculation of a fluorescence signal, Z-stacks with maximum intensity were merged. A Region Of Interest (ROI) was drawn around the signal of interest, and the integrated density was determined to get the 'signal intensity'. ROI of the same size was put elsewhere in the nucleus for background signal and integrated density was determined to get the 'background intensity'. Each signal intensity value was normalized with the background and Spc42-mCherry intensity values to get the 'normalized signal intensity' which was used to plot the graphs. For *CEN V*-GFP intensity comparison between pre-S phase cells and metaphase I cells in meiosis, the normalized signal intensity obtained from metaphase I cells was divided by 2 to obtain intensity per *CEN V*-GFP as at this stage two sister centromeres are glued together by cohesin and two sister *CEN V*-GFP dots appeared as a single dot (Fig S14).

Second Harmonic Generation (SHG) microscopy and analysis

The sample preparation for SHG microscopy was followed as mentioned earlier (Yamin et al., 2019; 2022). Briefly, the cells were harvested and fixed in 4% paraformaldehyde for 20 mins at RT; they were washed with 0.1 M phosphate buffer and stained with DAPI (Invitrogen, #D1306) for 20 mins in dark. The cells were subsequently placed on a polylysine coated slide and allowed to adhere to the slide. After washing the immobilized cells with PBS, mounting solution was added and the slide was covered with a coverslip. Cells were then analyzed by a Zeiss LSM780 inverted Confocal Microscope, with multi-photon Chameleon vision II Laser (laser power > 3 W, pulse

width and repetition rate at default settings, tuning range 690‑1040 nm); Lambda−PMT−

identifying all the wavelength spectrums. The slides were excited by 768 nm and emission was

collected at 414-628 nm. The quantification was carried out using ImageJ software as mentioned

earlier (Yamin et al., 2019; 2022).

Chromatin spread and the analysis

Chromatin spread was performed by following the protocol mentioned elsewhere (Mehta et al.,

2014) with certain modifications. Cells were harvested from the mitotic culture of 1.0 OD_{600} and

at the required time point for meiotic culture. The cells were washed once with spheroplasting

solution (1.2 M sorbitol in 0.1 M phosphate buffer) and harvested by centrifugation at 3000 RPM

for 3 mins. The cells were resuspended in 0.5 ml of spheroplasting solution supplemented with 5

µl of β-mercaptoethanol and 12.5 µl of zymolyase 20T (10 mg/ml, MP Biomedicals) and incubated

at 30ºC for 1.5 hrs. Once 80-90% of the cells were spheroplasted (by observing under the

microscope), the spheroplasting was stopped by adding 1 ml of ice-cold stop solution (0.1 M MES

- pH 6.4, 1 mM EDTA, 0.5 mM $MgCl_2$, 1 M Sorbitol). The spheroplasts were centrifuged at 2000

RPM for 3 mins at 4 ºC and resuspended gently in 125 µl of ice-cold stop solution. 60 µl of

spheroplasted cells were put on acid-washed glass slides followed by the addition of 40 µl of

fixative solution (4% paraformaldehyde, 3.4% sucrose, 10 N NaOH) and 80 µl of 1% lipsol (LIP

Equipment and services) over it. After gentle mixing 80 µl of fixative solution was again added to

the slides. The content on the slides was mixed and spread gently using a pipette tip and kept at

room temperature overnight for drying. The slides were then washed with 2 ml of 0.4% photo

flow-200 (Kodak) and then submerged in 1X PBS for 15 mins. The excess liquid was removed

and 100 µl of blocking solution (10 mg/ml BSA in 1X PBS + 5% Skim milk) was immediately

added over the fixed nuclear spreads which were then covered with coverslips and incubated at

room temperature in a humid chamber for 20 mins. The coverslips were removed and 100 µl of

primary antibody (1:200) was added to each slide and incubated at room temperature for 60 mins.

The primary antibody was rinsed off by submerging the slides 3 times in 1X PBS for 5 mins each

and 100 µl of secondary antibody (1:200) was added followed by incubating at room temperature

for 60 mins. The slides were again washed in the same way followed by incubation with 100 µl of

DAPI (1 µg/ml, Invitrogen) for 30 mins. After PBS wash for 5 mins, 100 µl of mounting solution

(90% glycerol supplemented with 1 mg/ml p-phenylenediamine) was added over the slides

followed by a clean coverslip which was sealed with transparent nail paint. The primary antibodies

used: rat anti-HA (3F10, Roche), mouse anti-HA (12CA5, Roche) and rat anti-Tubulin (MCA78G,

Serotec). The secondary antibodies used: Rhodamine (TRITC)-conjugated goat anti-Rat (Jackson)

and Alexa Fluor 488 conjugated goat anti-Mouse IgG (Jackson).

For calculation of the number of fluorescence foci associated with each chromatin spread,

all the Z-stacks with the signals were first merged and then the file was exported to Imaris (8.0.2)

software. The images were visualized using '3-D view' tool of the software. Subsequently, the

'Spot' tool in the algorithm setting of the software was used to mark the different foci present in

the chromatin spreads. The background fluorescence was subtracted using an automated

'subtraction' tool, and the number of foci visible for each spread was scored.

Indirect immunofluorescence

Immunofluorescence was performed, as mentioned previously (Mittal et al., 2020; Shah et al.,

2023). Briefly, cells from meiotic culture were harvested at indicated time points and fixed with

4% formaldehyde at room temperature (RT) for 15 mins. The fixed cells were washed once with

PBS and spheroplasted in spheroplasting solution (1.2 M sorbitol, 0.1 M phosphate buffer pH 7.5),

in the presence of Zymolyase 20T (MP Biomedicals, #32092) (10 mg/ml) and 25 mM β-mercaptoethanol or 1 hr at 30°C. The spheroplasts were mounted on a polylysine coated slide. The cells were then flattened by immersing the slide in -20°C methanol for 5 mins and in acetone for 30 s. The slides were dried for 1-2 mins before blocking using 5% skim milk solution prepared in dilution buffer (10 mg/ml BSA in PBS). The cells were incubated with appropriate primary and secondary antibodies for 1 hr at RT. Following several washes with PBS, samples were incubated with DAPI (Invitrogen, #D1306) (1 μg/ml) solution for 20 mins in dark. Finally, the glass slides were sealed with coverslips for imaging after spreading the mounting solution (1 mg/ml phenylenediamine in 90% glycerol). The antibodies and their dilutions (in dilution buffer) were mentioned as follows. Primary antibodies rat anti-tubulin (Serotec, MCA78G) at 1:5000. Secondary antibodies: (TRITC)-labeled goat anti-rat (Jackson, #115485166).

ChIP (Chromatin immunoprecipitation)

Chromatin immunoprecipitation was performed as described previously (Makrantoni et al., 2019) with certain modifications. Chromatin was cross-linked with 1% formaldehyde (30 mins for Brn1-6HA, 1 hr for Sgo1-9Myc, 2 hrs for Ipl1-6HA) at 25°C with shaking of the culture at 100 RPM, followed by the addition of glycine at a final concentration of 125 mM to quench the crosslinking. Cells were harvested and washed twice with 10 ml ice-cold TBS buffer and once with 10 ml 1X FA lysis buffer (50 mM Hepes–KOH pH 7.5, 150 mM NaCl, 1 mM EDTA, 1% Triton X-100, 0.1% Na-deoxycholate) supplemented with 0.1 % SDS. The cell pellet was resuspended in 500 μl of ice-cold 1X FA lysis buffer supplemented with 0.5% SDS, 1 mM PMSF and 1X PIC (Protease Inhibitor Cocktail, Roche). An equal volume of glass beads was added and cells were lysed in a mini bead beater (BIOSPEC, India) for 9 cycles (1 min ON, 1 min OFF on ice). The lysate was collected in a pre-chilled 1.5 ml eppendorf tube and centrifuged at 16,000 x g for 15 min at 4°C.

The supernatant was removed and the pellet was resuspended in 1 ml ice-cold 1X FA lysis buffer

supplemented with 0.1% SDS, 1 mM PMSF and PIC. The tubes were again centrifuged at 16,000

x g for 15 mins. The supernatant was discarded and the chromatin pellet was resuspended in 300

µl ice-cold 1X FA lysis buffer supplemented with 0.1% SDS, 1 mM PMSF and PIC. The chromatin

was then sheared to 300-500 bp using a water bath sonicator (Diagenode SA, Picoruptor, BC 100,

LAUDA Germany) for 30 cycles of 30 s ON / 30 s OFF at 4°C. The sonicated sample was then

centrifuged at 16,000 x g for 15 mins at 4°C. The supernatant was collected to which 1X FA lysis

buffer was again added and re-centrifuged at the same settings. 10 µl of the supernatant was stored

as input at -20°C (Whole cell extract, WCE) and the rest was divided into 'plus antibody' (+Ab)

and 'minus antibody' (-Ab), 600 µl each. Aliquots (+Ab) were incubated with 5 µg antibody [anti-

HA (3F10, Roche) or anti-MYC (AB9106, Abcam)] overnight in rotating conditions followed by

incubation with 50 µl of 50% Protein A Sepharose-beads (GE Healthcare,17-0780-01) for 2 hrs at

4°C in rotating condition. The beads were washed at room temperature twice with 1 ml of IP wash

buffer I (1X FA lysis buffer, 0.1%SDS, 275 mM NaCl), twice with 1 ml of IP wash buffer II (1X

FA lysis buffer, 0.1%SDS, 500 mM NaCl), once with 1 ml of IP wash buffer III (10 mM Tris pH

8.0, 250 mM LiCl, 0.5% NP-40, 0.5% sodium deoxycholate, 1 mM EDTA) and once with 1 ml of

1X TE (Tris EDTA). The beads were resuspended in elution buffer (50 mM Tris, pH 8.0, 10 mM

EDTA, 1% SDS) and incubated overnight at 65°C for de-crosslinking. De-crosslinked samples

were treated with Proteinase K (SRL, India) at 45 °C for 2 hrs followed by P:C:I

(Phenol:Chloroform:Isoamyl, Himedia) purification. Purified samples were further used for qPCR

analysis using CFX96 Real-Time system (BioRad) and SYBR green dye (BioRad) for detection.

Enrichment/Input was calculated by the following equation: $\% \, Enrichment/input =$

$E^\wedge(-\Delta Ct)$; where, E is the primer efficiency and was calculated by plotting Ct values of serially

diluted input with respect to Log dilution. $E = (-1)/slope$ and $\Delta Ct = Ct \, _{(Test \, sample)} - (Ct \, _{(Input)} - Log$

E (Input dilution factor). Primers used for PCR amplification are given in Table S2. For calculation

of percentage decrease of Sgo1-9Myc enrichment at the centromeres in the ChIP assays (Fig 5C

and D) following formulae was used: % decrease $= \frac{WT - MT}{WT} x\ 10$; (WT and MT are average

enrichment/input values from the wild type and the mutant, respectively).

Protein isolation and western blotting

Samples for western blotting were prepared by resuspending the cells in 200 µl of 20% TCA.

Equal volume of glass beads was added, and bead beating was done using a mini bead beater

(BIOSPEC, India) for 5 cycles (1 min ON, 1 min OFF on ice). Supernatant was collected by

centrifugating at 2800 RPM for 2 mins. 400 µl of 5% TCA was again added to the beads and

supernatant was again collected after centrifugation. The total supernatant obtained (600 µl) was

centrifuged at maximum speed (13,500 RPM) for 5 mins to precipitate the protein. The supernatant

was removed, and the protein pellet was resuspended in Laemmli buffer and loaded on SDS-PAGE

after boiling at 95°C for 5 mins for further analysis. For phosphatase assay, protocol mentioned

elsewhere (Ontoso et al., 2013) was followed with certain modifications. TCA extract in Laemmli

buffer was diluted 10-fold in 1X PMP buffer (NEB, Protein Metallo Phosphatases) supplemented

with 1 mM Mncl$_2$. The diluted sample was treated with 2 µl Lambda protein phosphatase (NEB)

and incubated in water bath at 30 °C for 4 hrs. As control, a similar diluted extract but without

adding phosphatase was incubated at similar conditions. The protein sample was TCA precipitated

and resuspended in Laemmli buffer followed by heating at 95°C for 5 mins. Western blotting was

performed using standard procedure. The blots were developed using following antibodies: rat

anti-tubulin antibody (MCA78G, Serotec,), mouse anti-HA (12CA5 and 3F10, Roche), HRP-

conjugated goat anti-mouse IgG (Jackson), HRP-conjugated goat anti-rat (Jackson). For CHX-

chase experiments, the percentage of protein remaining at each time point was calculated by taking

the protein level at 0 hr as 100%.

RNA isolation

Total RNA was extracted from meiotic cell culture of 1.5-2 OD_{600} (~2.5 x10^7 cells) using TRIzol®

reagent and PureLink® RNA mini kit (Ambion Life Technologies™). The cell pellet was re-

suspended in TRIzol® and cell lysis was done using DEPC treated sterile 0.5 mm glass beads. The

rest of the protocol was followed as instructed by the supplier (PureLink® RNA mini kit). RNA

was eluted in DEPC (Sigma Aldrich Chemicals Pvt. Ltd.) treated sterile water. DNase treatment

was done at 37°C for 10 mins using RNAse-free DNase I (Thermo Fisher Scientific) in the buffer

supplied with the enzyme. The DNase and salts in the buffer were removed by purification using

trizol-based layer separation and pure RNA was recovered from the aqueous layer. To ensure the

absence of DNA impurity, end-point PCR was performed.

Biological replicates and statistical methods

The quantitative data shown in all the figures were obtained from two or three biological replicates.

The N values in the figure legends are the total number of cells analyzed from combined replicates

of individual assay. Box plots were generated using OriginPro software (OriginLab). Statistical

significance was estimated using two-tailed student's t-test for mean with $p < .05$ considered to be

significant.

Competing Interest Statement

The authors declare no competing interests

Acknowledgement

We acknowledge the central instrumentation facility of IIT Bombay. The SKG laboratory is

supported by the Science and Engineering Research Board (SERB), Govt. of India

(CRG/2020/000444). DT and PA are supported by CSIR fellowships (09/087(0886)/2017- EMR-

I), and (09/087(0972)/2019- EMR-I), respectively. AA is supported by JRF fellowship from

Department of Biotechnology (DBT), Govt of India (BT/PR43050/BRB/10/1992/2021).

Author contributions: DT, PA and AA performed experiments. DT and SKG designed

experiments, analyzed and interpreted the data. DT and SKG wrote the manuscript.

800   References

Ackermann, S., Goeser, F., Schulte, J.H., Schramm, A., Ehemann, V., Hero, B., Eggert, A.,

Berthold, F. and Fischer, M., 2011. Polo-like kinase 1 is a therapeutic target in high-risk

neuroblastoma. *Clinical Cancer Research*, *17*(4), pp.731-741.

Agarwal, M., Mehta, G. and Ghosh, S.K., 2015. Role of Ctf3 and COMA subcomplexes in meiosis:

Implication in maintaining Cse4 at the centromere and numeric spindle poles. *Biochimica et*

Biophysica Acta (BBA)-Molecular Cell Research, *1853*(3), pp.671-684.

Attner, M.A., Miller, M.P., Ee, L.S., Elkin, S.K. and Amon, A., 2013. Polo kinase Cdc5 is a central

regulator of meiosis I. *Proceedings of the National Academy of Sciences*, *110*(35), pp.14278-

14283.

Bazile, F., St-Pierre, J. and D'Amours, D., 2010. Three-step model for condensin activation during

mitotic chromosome condensation. *Cell Cycle*, *9*(16), pp.3263-3275.

Borek, W.E., Vincenten, N., Duro, E., Makrantoni, V., Spanos, C., Sarangapani, K.K., de Lima

Alves, F., Kelly, D.A., Asbury, C.L., Rappsilber, J. and Marston, A.L., 2021. The proteomic

landscape of centromeric chromatin reveals an essential role for the Ctf19CCAN complex in

meiotic kinetochore assembly. *Current Biology*, *31*(2), pp.283-296.

Cha, R.S., Weiner, B.M., Keeney, S., Dekker, J. and Kleckner, N., 2000. Progression of meiotic

DNA replication is modulated by interchromosomal interaction proteins, negatively by Spo11p

and positively by Rec8p. *Genes & development*, *14*(4), pp.493-503.

Challa, K., Shinohara, M. and Shinohara, A., 2019. Meiotic prophase-like pathway for cleavage-independent removal of cohesin for chromosome morphogenesis. *Current genetics*, *65*, pp.817-827.

Colomina, N., Garí, E., Gallego, C., Herrero, E. and Aldea, M., 1999. G1 cyclins block the Ime1 pathway to make mitosis and meiosis incompatible in budding yeast. *The EMBO journal*, *18*(2), pp.320-329.

Colomina, N., Liu, Y., Aldea, M. and Garí, E., 2003. TOR regulates the subcellular localization of Ime1, a transcriptional activator of meiotic development in budding yeast. *Molecular and cellular biology*, *23*(20), pp.7415-7424.

D'Ambrosio, C., Schmidt, C.K., Katou, Y., Kelly, G., Itoh, T., Shirahige, K. and Uhlmann, F., 2008. Identification of cis-acting sites for condensin loading onto budding yeast chromosomes. *Genes & development*, *22*(16), p.2215.

Dauban, L., Kamgoué, A., Wang, R., Léger-Silvestre, I., Beckouët, F., Cantaloube, S. and Gadal, O., 2019. Quantification of the dynamic behaviour of ribosomal DNA genes and nucleolus during yeast Saccharomyces cerevisiae cell cycle. *Journal of structural biology*, *208*(2), pp.152-164.

Deng, X. and Kuo, M.H., 2018. Tripartite chromatin localization of budding yeast shugoshin involves higher-ordered architecture of mitotic chromosomes. *G3: Genes, Genomes, Genetics*, *8*(9), pp.2901-2911.

Dong, Q. and Li, F., 2022. Cell cycle control of kinetochore assembly. *Nucleus*, *13*(1), pp.210-222.

Dou, Z., Liu, R., Zang, J., Yao, X. and Liu, X., 2023. The function, structure and dynamic organization of centromeres and kinetochores. *JUSTC*, *53*, pp.1-9.

Eckert, C.A., Gravdahl, D.J. and Megee, P.C., 2007. The enhancement of pericentromeric cohesin association by conserved kinetochore components promotes high-fidelity chromosome segregation and is sensitive to microtubule-based tension. *Genes & development*, *21*(3), pp.278-291.

Fernius, J. and Hardwick, K.G., 2007. Bub1 kinase targets Sgo1 to ensure efficient chromosome biorientation in budding yeast mitosis. *PLoS genetics*, *3*(11), p.e213.

Fernius, J. and Marston, A.L., 2009. Establishment of cohesion at the pericentromere by the Ctf19
kinetochore subcomplex and the replication fork-associated factor, Csm3. *PLoS genetics*, *5*(9),
p.e1000629.

Fernius, J., Nerusheva, O.O., Galander, S., de Lima Alves, F., Rappsilber, J. and Marston, A.L.,
2013. Cohesin-dependent association of scc2/4 with the centromere initiates pericentromeric
cohesion establishment. *Current Biology*, *23*(7), pp.599-606.

Fischböck-Halwachs, J., Singh, S., Potocnjak, M., Hagemann, G., Solis-Mezarino, V., Woike, S.,
Ghodgaonkar-Steger, M., Weissmann, F., Gallego, L.D., Rojas, J. and Andreani, J., 2019. The
COMA complex interacts with Cse4 and positions Sli15/Ipl1 at the budding yeast inner
kinetochore. *Elife*, *8*, p.e42879.

Ghosh, S.K., Sau, S., Lahiri, S., Lohia, A. and Sinha, P., 2004. The Iml3 protein of the budding
yeast is required for the prevention of precocious sister chromatid separation in meiosis I and for
sister chromatid disjunction in meiosis II. *Current genetics*, *46*, pp.82-91.

Giet, R. and Glover, D.M., 2001. Drosophila aurora B kinase is required for histone H3
phosphorylation and condensin recruitment during chromosome condensation and to organize the
central spindle during cytokinesis. *The Journal of cell biology*, *152*(4), pp.669-682.

Glynn, E.F., Megee, P.C., Yu, H.G., Mistrot, C., Unal, E., Koshland, D.E., DeRisi, J.L. and Gerton,
J.L., 2004. Genome-wide mapping of the cohesin complex in the yeast Saccharomyces
cerevisiae. *PLoS biology*, *2*(9), p.e259.

Guacci, V., Koshland, D. and Strunnikov, A., 1997. A direct link between sister chromatid
cohesion and chromosome condensation revealed through the analysis of MCD1 in S.
cerevisiae. *Cell*, *91*(1), pp.47-57.

Güldener, U., Heck, S., Fiedler, T., Beinhauer, J. and Hegemann, J.H., 1996. A new efficient gene
disruption cassette for repeated use in budding yeast. *Nucleic acids research*, *24*(13), pp.2519-
2524.

Hinshaw, S.M., Makrantoni, V., Harrison, S.C. and Marston, A.L., 2017. The kinetochore receptor
for the cohesin loading complex. *Cell*, *171*(1), pp.72-84.

Hinshaw, S.M., Makrantoni, V., Kerr, A., Marston, A.L. and Harrison, S.C., 2015. Structural evidence for Scc4-dependent localization of cohesin loading. *Elife*, *4*, p.e06057.

Holt, S.V., Vergnolle, M.A., Hussein, D., Wozniak, M.J., Allan, V.J. and Taylor, S.S., 2005. Silencing Cenp-F weakens centromeric cohesion, prevents chromosome alignment and activates the spindle checkpoint. *Journal of cell science*, *118*(20), pp.4889-4900.

Jain, N., Janning, P. and Neumann, H., 2021. 14-3-3 Protein Bmh1 triggers short-range compaction of mitotic chromosomes by recruiting sirtuin deacetylase Hst2. *Journal of Biological Chemistry*, *296*.

Janke, C., Magiera, M.M., Rathfelder, N., Taxis, C., Reber, S., Maekawa, H., Moreno-Borchart, A., Doenges, G., Schwob, E., Schiebel, E. and Knop, M., 2004. A versatile toolbox for PCR-based tagging of yeast genes: new fluorescent proteins, more markers and promoter substitution cassettes. *Yeast*, *21*(11), pp.947-962.

Kerr, G.W., Sarkar, S., Tibbles, K.L., Petronczki, M., Millar, J.B. and Arumugam, P., 2011. Meiotic nuclear divisions in budding yeast require PP2ACdc55-mediated antagonism of Net1 phosphorylation by Cdk. *Journal of Cell Biology*, *193*(7), pp.1157-1166.

Kiburz, B.M., Reynolds, D.B., Megee, P.C., Marston, A.L., Lee, B.H., Lee, T.I., Levine, S.S., Young, R.A. and Amon, A., 2005. The core centromere and Sgo1 establish a 50-kb cohesin-protected domain around centromeres during meiosis I. *Genes & development*, *19*(24), pp.3017-3030.

Kim, S., Meyer, R., Chuong, H. and Dawson, D.S., 2013. Dual mechanisms prevent premature chromosome segregation during meiosis. *Genes & development*, *27*(19), pp.2139-2146.

Kimura, K., Hirano, M., Kobayashi, R. and Hirano, T., 1998. Phosphorylation and activation of 13 S condensin by Cdc2 in vitro. *Science*, *282*(5388), pp.487-490.

Korns, J., Liu, X. and Takiar, V., 2022. A review of Plks: Thinking outside the (polo) box. *Molecular Carcinogenesis*, *61*(2), pp.254-263.

Kruitwagen, T., Chymkowitch, P., Denoth-Lippuner, A., Enserink, J. and Barral, Y., 2018. Centromeres license the mitotic condensation of yeast chromosome arms. *Cell*, *175*(3), pp.780-795.

Kruitwagen, T., Denoth-Lippuner, A., Wilkins, B.J., Neumann, H. and Barral, Y., 2015. Axial
contraction and short-range compaction of chromatin synergistically promote mitotic chromosome
condensation. *Elife*, *4*, p.e10396.

Kumar, D., Prajapati, H.K., Mahilkar, A., Ma, C.H., Mittal, P., Jayaram, M. and Ghosh, S.K.,
2021. The selfish yeast plasmid utilizes the condensin complex and condensed chromatin for
faithful partitioning. *PLoS Genetics*, *17*(7), p.e1009660.

Lamothe, R., Costantino, L. and Koshland, D.E., 2020. The spatial regulation of condensin activity
in chromosome condensation. *Genes & Development*, *34*(11-12), pp.819-831.

Lavoie, B.D., Hogan, E. and Koshland, D., 2002. In vivo dissection of the chromosome
condensation machinery: reversibility of condensation distinguishes contributions of condensin
and cohesin. *The Journal of cell biology*, *156*(5), pp.805-815.

Lavoie, B.D., Hogan, E. and Koshland, D., 2004. In vivo requirements for rDNA chromosome
condensation reveal two cell-cycle-regulated pathways for mitotic chromosome folding. *Genes &
development*, *18*(1), pp.76-87.

Lee, B.H. and Amon, A., 2003. Role of Polo-like kinase CDC5 in programming meiosis I
chromosome segregation. *Science*, *300*(5618), pp.482-486.

Leonard, J., Sen, N., Torres, R., Sutani, T., Jarmuz, A., Shirahige, K. and Aragon, L., 2015.
Condensin relocalization from centromeres to chromosome arms promotes Top2 recruitment
during anaphase. *Cell Reports*, *13*(11), pp.2336-2344.

Li, P., Jin, H. and Yu, H.G., 2014. Condensin suppresses recombination and regulates double-
strand break processing at the repetitive ribosomal DNA array to ensure proper chromosome
segregation during meiosis in budding yeast. *Molecular biology of the cell*, *25*(19), pp.2934-2947.

Longtine, M.S., Mckenzie III, A., Demarini, D.J., Shah, N.G., Wach, A., Brachat, A., Philippsen,
P. and Pringle, J.R., 1998. Additional modules for versatile and economical PCR-based gene
deletion and modification in Saccharomyces cerevisiae. *Yeast*, *14*(10), pp.953-961.

Makrantoni, V., Robertson, D. and Marston, A.L., 2019. Analysis of the chromosomal localization
of yeast SMC complexes by chromatin immunoprecipitation. *SMC Complexes: Methods and
Protocols*, pp.119-138.

Markowitz, T.E., Heldrich, J. and Hochwagen, A., 2020. DNA break formation induces
Scc2/cohesin-dependent recruitment of condensin to meiotic chromosomes. *bioRxiv*, pp.2020-07.

Mehta, G., Sanyal, K., Abhishek, S., Rajakumara, E. and Ghosh, S.K., 2022. Minichromosome
maintenance proteins in eukaryotic chromosome segregation. *BioEssays*, *44*(1), p.2100218.

Mehta, G.D., Agarwal, M. and Ghosh, S.K., 2014. Functional characterization of kinetochore
protein, Ctf19 in meiosis I: an implication of differential impact of Ctf19 on the assembly of
mitotic and meiotic kinetochores in Saccharomyces cerevisiae. *Molecular microbiology*, *91*(6),
pp.1179-1199.

Meraldi, P., McAinsh, A.D., Rheinbay, E. and Sorger, P.K., 2006. Phylogenetic and structural
analysis of centromeric DNA and kinetochore proteins. *Genome biology*, *7*, pp.1-21.

Meyer, R.E., Chuong, H.H., Hild, M., Hansen, C.L., Kinter, M. and Dawson, D.S., 2015.
Ipl1/Aurora-B is necessary for kinetochore restructuring in meiosis I in Saccharomyces
cerevisiae. *Molecular biology of the cell*, *26*(17), pp.2986-3000.

Michaelis, C., Ciosk, R. and Nasmyth, K., 1997. Cohesins: chromosomal proteins that prevent
premature separation of sister chromatids. *Cell*, *91*(1), pp.35-45.

Mishra, P.K., Ciftci-Yilmaz, S., Reynolds, D., Au, W.C., Boeckmann, L., Dittman, L.E., Jowhar,
Z., Pachpor, T., Yeh, E., Baker, R.E. and Hoyt, M.A., 2016. Polo kinase Cdc5 associates with
centromeres to facilitate the removal of centromeric cohesin during mitosis. *Molecular Biology of*
the Cell, *27*(14), pp.2286-2300.

Mishra, P.K., Olafsson, G., Boeckmann, L., Westlake, T.J., Jowhar, Z.M., Dittman, L.E., Baker,
R.E., D'Amours, D., Thorpe, P.H. and Basrai, M.A., 2019. Cell cycle–dependent association of
polo kinase Cdc5 with CENP-A contributes to faithful chromosome segregation in budding
yeast. *Molecular Biology of the Cell*, *30*(8), pp.1020-1036.

Mora-Bermúdez, F., Gerlich, D. and Ellenberg, J., 2007. Maximal chromosome compaction occurs
by axial shortening in anaphase and depends on Aurora kinase. *Nature cell biology*, *9*(7), pp.822-
831.

Müller, C.A. and Nieduszynski, C.A., 2012. Conservation of replication timing reveals global and
local regulation of replication origin activity. *Genome research*, *22*(10), pp.1953-1962.

Nakazawa, N., Nakamura, T., Kokubu, A., Ebe, M., Nagao, K. and Yanagida, M., 2008. Dissection of the essential steps for condensin accumulation at kinetochores and rDNAs during fission yeast mitosis. *The Journal of cell biology*, *180*(6), pp.1115-1131.

Natsume, T., Müller, C.A., Katou, Y., Retkute, R., Gierliński, M., Araki, H., Blow, J.J., Shirahige, K., Nieduszynski, C.A. and Tanaka, T.U., 2013. Kinetochores coordinate pericentromeric cohesion and early DNA replication by Cdc7-Dbf4 kinase recruitment. *Molecular cell*, *50*(5), pp.661-674.

Nishimura, K., Fukagawa, T., Takisawa, H., Kakimoto, T. and Kanemaki, M., 2009. An auxin-based degron system for the rapid depletion of proteins in nonplant cells. *Nature methods*, *6*(12), pp.917-922.

Oelschlaegel, T., Schwickart, M., Matos, J., Bogdanova, A., Camasses, A., Havlis, J., Shevchenko, A. and Zachariae, W., 2005. The yeast APC/C subunit Mnd2 prevents premature sister chromatid separation triggered by the meiosis-specific APC/C-Ama1. *Cell*, *120*(6), pp.773-788.

Okaz, E., Argüello-Miranda, O., Bogdanova, A., Vinod, P.K., Lipp, J.J., Markova, Z., Zagoriy, I., Novak, B. and Zachariae, W., 2012. Meiotic prophase requires proteolysis of M phase regulators mediated by the meiosis-specific APC/CAma1. *Cell*, *151*(3), pp.603-618.

Ontoso, D., Acosta, I., van Leeuwen, F., Freire, R. and San-Segundo, P.A., 2013. Dot1-dependent histone H3K79 methylation promotes activation of the Mek1 meiotic checkpoint effector kinase by regulating the Hop1 adaptor. *PLoS genetics*, *9*(1), p.e1003262.

Peplowska, K., Wallek, A.U. and Storchova, Z., 2014. Sgo1 regulates both condensin and Ipl1/Aurora B to promote chromosome biorientation. *PLoS Genetics*, *10*(6), p.e1004411.

Petersen, J. and Hagan, I.M., 2003. S. pombe aurora kinase/survivin is required for chromosome condensation and the spindle checkpoint attachment response. *Current biology*, *13*(7), pp.590-597.

Piazza, I., Rutkowska, A., Ori, A., Walczak, M., Metz, J., Pelechano, V., Beck, M. and Haering, C.H., 2014. Association of condensin with chromosomes depends on DNA binding by its HEAT-repeat subunits. *Nature structural & molecular biology*, *21*(6), pp.560-568.

Raghuraman, M.K., Winzeler, E.A., Collingwood, D., Hunt, S., Wodicka, L., Conway, A.,
Lockhart, D.J., Davis, R.W., Brewer, B.J. and Fangman, W.L., 2001. Replication dynamics of the
yeast genome. *science*, *294*(5540), pp.115-121.

Schrader, F., 1939. Kinetic regions in chromosomes. *Nature*, *143*(3612), pp.122-122.

Shirk, K., Jin, H., Giddings Jr, T.H., Winey, M. and Yu, H.G., 2011. The Aurora kinase Ipl1 is
necessary for spindle pole body cohesion during budding yeast meiosis. *Journal of cell*
science, *124*(17), pp.2891-2896.

Snead, J.L., Sullivan, M., Lowery, D.M., Cohen, M.S., Zhang, C., Randle, D.H., Taunton, J.,
Yaffe, M.B., Morgan, D.O. and Shokat, K.M., 2007. A coupled chemical-genetic and
bioinformatic approach to Polo-like kinase pathway exploration. *Chemistry & biology*, *14*(11),
pp.1261-1272.

St-Pierre, J., Douziech, M., Bazile, F., Pascariu, M., Bonneil, É., Sauvé, V., Ratsima, H. and
D'Amours, D., 2009. Polo kinase regulates mitotic chromosome condensation by hyperactivation
of condensin DNA supercoiling activity. *Molecular cell*, *34*(4), pp.416-426.

Sutani, T., Yuasa, T., Tomonaga, T., Dohmae, N., Takio, K. and Yanagida, M., 1999. Fission yeast
condensin complex: essential roles of non-SMC subunits for condensation and Cdc2
phosphorylation of Cut3/SMC4. *Genes & development*, *13*(17), pp.2271-2283.

Takai, N., Hamanaka, R., Yoshimatsu, J. and Miyakawa, I., 2005. Polo-like kinases (Plks) and
cancer. *Oncogene*, *24*(2), pp.287-291.

Takemoto, A., Murayama, A., Katano, M., Urano, T., Furukawa, K., Yokoyama, S., Yanagisawa,
J., Hanaoka, F. and Kimura, K., 2007. Analysis of the role of Aurora B on the chromosomal
targeting of condensin I. *Nucleic acids research*, *35*(7), pp.2403-2412.

Tanaka, T., Fuchs, J., Loidl, J. and Nasmyth, K., 2000. Cohesin ensures bipolar attachment of
microtubules to sister centromeres and resists their precocious separation. *Nature cell*
biology, *2*(8), pp.492-499.

Tsubouchi, T. and Roeder, G.S., 2005. A synaptonemal complex protein promotes homology-
independent centromere coupling. *Science*, *308*(5723), pp.870-873.

Vas, A.C., Andrews, C.A., Kirkland Matesky, K. and Clarke, D.J., 2007. In vivo analysis of chromosome condensation in Saccharomyces cerevisiae. *Molecular biology of the cell*, *18*(2), pp.557-568.

Verzijlbergen, K.F., Nerusheva, O.O., Kelly, D., Kerr, A., Clift, D., de Lima Alves, F., Rappsilber, J. and Marston, A.L., 2014. Shugoshin biases chromosomes for biorientation through condensin recruitment to the pericentromere. *Elife*, *3*, p.e01374.

Visintin, C., Tomson, B.N., Rahal, R., Paulson, J., Cohen, M., Taunton, J., Amon, A. and Visintin, R., 2008. APC/C-Cdh1-mediated degradation of the Polo kinase Cdc5 promotes the return of Cdc14 into the nucleolus. *Genes & development*, *22*(1), pp.79-90.

Wach, A., 1996. PCR-synthesis of marker cassettes with long flanking homology regions for gene disruptions in S. cerevisiae. *Yeast*, *12*(3), pp.259-265.

Walters, A.D., May, C.K., Dauster, E.S., Cinquin, B.P., Smith, E.A., Robellet, X., D'Amours, D., Larabell, C.A. and Cohen-Fix, O., 2014. The yeast polo kinase Cdc5 regulates the shape of the mitotic nucleus. *Current Biology*, *24*(23), pp.2861-2867.

Wilkins, B.J., Rall, N.A., Ostwal, Y., Kruitwagen, T., Hiragami-Hamada, K., Winkler, M., Barral, Y., Fischle, W. and Neumann, H., 2014. A cascade of histone modifications induces chromatin condensation in mitosis. *Science*, *343*(6166), pp.77-80.

Wong, J., Nakajima, Y., Westermann, S., Shang, C., Kang, J.S., Goodner, C., Houshmand, P., Fields, S., Chan, C.S., Drubin, D. and Barnes, G., 2007. A protein interaction map of the mitotic spindle. *Molecular biology of the cell*, *18*(10), pp.3800-3809.

Yahya, G., Wu, Y., Peplowska, K., Röhrl, J., Soh, Y.M., Bürmann, F., Gruber, S. and Storchova, Z., 2020. Phospho-regulation of the Shugoshin-Condensin interaction at the centromere in budding yeast. *PLoS genetics*, *16*(8), p.e1008569.

Yamin, K., Assa, M., Matityahu, A., and Onn, I., 2020. Analyzing chromosome condensation in yeast by second-harmonic generation microscopy. *Current Genetics, 66*(2), 437–443. doi: 10.1007/s00294-019-01034-1

Yamin, K., Bijlani, S., Berman, J., Soni, A., Shlomai, J., Buragohain, B. M., and Onn, I., 2022.
Fold-change of chromatin condensation in yeast is a conserved property. *Scientific Reports, 12*(1),
17393. doi: 10.1038/s41598-022-22340-8

Yu, H.G. and Koshland, D., 2005. Chromosome morphogenesis: condensin-dependent cohesin
removal during meiosis. *Cell*, *123*(3), pp.397-407.

Zeng, X., Kahana, J.A., Silver, P.A., Morphew, M.K., McIntosh, J.R., Fitch, I.T., Carbon, J. and
Saunders, W.S., 1999. Slk19p is a centromere protein that functions to stabilize mitotic
spindles. *The Journal of cell biology*, *146*(2), pp.415-425.

Zoller, J.F., Herrmann, R.G. and Wanner, G., 2004. Chromosome condensation in mitosis and
meiosis of rye (Secale cereale L.). *Cytogenetic and Genome Research*, *105*(1), pp.134-144.

1048

1049

1050

1051

1052

1053

1054

1055

1056

1057

1058

1059

1060

1061

Figure 1

1062

Figure 1. Chromosomes are more condensed in meiosis than in mitosis. (A) Schematic diagrams showing the strategy adopted for in vivo measurement of chromosome condensation. One copy of chromosome V was tagged using [TetO]$_{224}$-[TetR-GFP] system near the centromere V. As the [TetO]$_{224}$ repeats come closer due to axial condensation in the metaphase stage, the GFP signal intensity decreases because of fluorescence quenching. The fluorescence intensity of *CEN V*-GFP measured in (B) mitotic and

(C) meiotic cells (see materials methods for detail) from the indicated cell cycle stages are graphically

represented on the right; the corresponding representative images are shown on the left. N $\geq$ 30-45 from

three independent experiments, scale bar = 2 µm. (D) The cells used in (B) and (C) were mixed and

fluorescence intensity was measured from mitotic (budded), and meiotic (unbudded) cells present in the

same field of view as represented on the left; the fluorescence intensities are presented graphically on right.

N $\geq$ 30 from two independent experiments, scale bar = 5 µm. The intensity values were normalized with

background and Spc42-mCherry (internal control) intensities. For statistical significance, p values were

estimated by the two-tailed student's t-test for the mean.

Figure 2

Figure 2. Ctf19 is involved in chromosome condensation in meiosis but not in mitosis. In vivo, chromosome condensation in *ctf19Δ* cells was measured as in Fig. 1 at the indicated stages in mitosis (A) and in meiosis (B). (C) Similar measurements were done in *slk19Δ* meiotic cells. The intensity values are graphically represented on the right; the corresponding representative images are shown on the left. N = 25-30 from two independent experiments, scale bar = 2 µm. (D) rDNA condensation in meiosis was analyzed by visualizing the rDNA-binding protein Net1 tagged with GFP in the wild type, *ctf19Δ*, and *slk19Δ* cells arrested in the metaphase I stage. N > 200, scale bar = 2 µm. Error bars represent the standard deviation from the mean values obtained from three independent experiments. For statistical significance, *p* values were estimated by the two-tailed student's t-test for the mean.

Figure 3

Figure 3. Ctf19 contributes to chromosome condensation along the chromosomal arm (A) One copy of chromosome V was marked with GFP at two different chromosomal loci ~400 kb away from each other,

each proximal to the centromere or telomere was marked with [TetO]$_{224}$–[TetR-GFP] as *CEN V*-GFP or

with [TetO]$_{448}$ –[TetR-GFP] as *TEL V*-GFP, respectively. The *TEL V*-GFP dot appeared brighter than the

CEN V-GFP dot (arrow). (B) The representative images are shown for the indicated strains and the cell

cycle stages. Scale bar = 2 μm (C) The *CEN V*-GFP (*left*) and *TEL V*-GFP (*right*) intensities measured as

described in materials and methods are shown for the indicated cells harvested from the indicated cell cycle

stages. N = 25-30 from two independent experiments. (D) The 3D distances between the *CEN V*-GFP and

TEL V-GFP dots measured as described in the materials and methods are shown for the wild type (left) and

ctf19Δ (right) cells harvested from the indicated cell cycle stages. N = 45-50 from two independent

experiments. The statistical significance *p* value was estimated by the two-tailed student's t-test for the

mean.

1099

Figure 4

1100

Figure 4. The loss of Ctf19 causes reduced association of condensin with chromatin by impairing the establishment of the association in meiosis (A) Schematic diagram showing the locations of PCR amplicons (horizontal bars) on chromosomes III and IV used for the qPCR analysis of the ChIP samples. The yellow oval shapes represent the centromeres (*CEN III* and *CEN IV*), pink and blue regions indicate the pericentromeric (P) and arm (A) regions, respectively. (B) ChIP assay using anti-HA antibodies (3F10) for quantifying the association of Brn1-6HA with the indicated chromosomal loci in the indicated cells arrested at metaphase I. The graph represents % enrichment per input obtained by qPCR analysis. Error bars represent the standard deviation from the mean values obtained from three independent experiments. The statistical significance *p* value was estimated by the two-tailed student's t-test for the mean. (C) The loss of Ctf19 impairs the establishment of the association of condensin with chromatin in meiosis. The left panel shows the representative images of the chromatin spreads showing different types of association (described in the text) of the condensin subunit (Brn1-6HA) with chromatin at different time points in SPM in wild type and *ctf19Δ* cells harboring homozygous *BRN1-6HA* allele. Anti-HA antibodies (3F10) were used to detect Brn1-6HA. The right panel represents the percentage of different types of spreads at the indicated time points. with respect to the Brn1-6HA signal. Arrowhead shows rDNA region. N >100 from two independent experiments, scale bar = 2 μm.

Figure 5

1117

Figure 5. Pericentric localization of Sgo1 is perturbed more in meiosis than in mitosis in the absence of Ctf19. The localization of Sgo1-GFP with respect to a kinetochore protein, Ndc80-mCherry in the wild type and *ctf19Δ* cells arrested at metaphase upon treatment with nocodazole (A) and harvested at metaphase I following their release into SPM for 8 and 10 hrs (B), respectively. The left panel shows the representative images of indicated categories and the right panel depicts the percentage of cells in each category. Error bars represent the standard deviation from the mean values obtained from two independent experiments. N = 90-120, scale bar = 2 μm. ChIP assays using anti-Myc antibodies (ab9106) for quantifying association of Sgo1-9Myc with the indicated chromosomal loci in the wild type and *ctf19Δ* cells arrested at metaphase (C) and metaphase I (D). The graph represents % enrichment per input obtained by qPCR analysis. Error bars represent the standard deviation from the mean values obtained from three independent experiments. The statistical significance *p* value was estimated by the two-tailed student's t-test for the mean.

Figure 6

Figure 6. Increase in the level of Cdc5 and decrease in its chromatin association in *ctf19Δ* mutant in meiosis. (A) The wild type and *ctf19Δ* cells harboring *CDC5-6HA pCLB2-CDC20* were arrested at the metaphase I following 8 and 10 hrs of meiotic induction, respectively. Left, the proteins extracted from the cells were run on a gel in dilutions and were detected by western blotting using anti-HA (3F10) and anti-tubulin (YOL1/34) antibodies. Right, the intensity of the bands was calculated using ImageJ software and normalized to the tubulin band intensity used as loading control. *p* value was estimated by the two-tailed student's t-test for the mean (B) Cdc5-6HA stability in wild type and *ctf19Δ* cells. Following release into synchronized meiosis for 8 and 10 hrs for the wild type and the mutant, respectively, the cells were treated with CHX (t = 0). The cells were harvested at indicated time points. Cdc5-6HA and tubulin levels were analyzed by western blotting (*left*) as in (A) and quantified (*right*) at indicated time points after CHX treatment. Error bars represent the standard deviation from the mean values obtained from three independent experiments. (C) Left, the representative images of the chromatin spreads show the association of Cdc5-6HA with chromatin in metaphase I arrested wild type and *ctf19Δ* cells harboring homozygous *CDC5-6HA* alleles. Right, quantitative analysis of Cdc5-6HA signal (*right*) on the chromatin spreads as type I (bright signal) and type II (low signal) categories. The metaphase I stage of the spreads was determined by SPB localization revealed by tubulin immunostaining. (D) The percentage of chromatin spreads with type II signal, as mentioned in (C), from the cells arrested in metaphase (mitosis) or metaphase I (meiosis). Error bars represent the standard deviation from the mean values obtained from two independent experiments (N = 83, scale bar = 2 µm).

Figure 7

Figure 7. Evidence of hyper-phosphorylation and reduced stability of Ycg1 in *ctf19* mutant in meiosis. (A) The wild type and *ctf19Δ* cells harboring *YCG1-6HA pCLB2-CDC20* were released into synchronized meiosis and were harvested at the indicated time points; proteins extracted by the TCA method (materials and methods) from the cells were resolved by running 6% SDS-PAGE gel and were detected by western blotting using anti-HA antibodies (12CA5). Notably, the B2 band is more prominent than the B1 band in the mutant. (B) The wild type and *ctf19Δ* cells harboring *YCG1-6HA pCLB2-CDC20* were released into SPM and the samples were harvested at 8 hrs and 10 hrs, respectively. Protein extracts treated with (+) or without (-) lambda phosphatase (λ P) were run on 7.5% SDS gel made with Phos-tag acrylamide (Wako) to detect Ycg1-6HA by western blotting using anti-HA antibodies (12CA5). A non-specific band is shown as a loading control. (C) Ycg1-6HA stability in wild type and *ctf19Δ* cells. The cells were treated with CHX (t = 0) following 8 and 10 hrs into meiotic induction for the wild type and *ctf19Δ*, respectively. Left, the protein extracts were resolved on 10% SDS-PAGE gel and the protein level was analyzed by western blotting using anti-HA (12CA5) and anti-tubulin (YOL1/34) antibodies in the cells harvested at indicated time points. Right, quantification of Ycg1-6HA bands normalized with the tubulin bands using ImageJ software at indicated time points after CHX treatment. Error bars represent the standard deviation from the mean values obtained from two independent experiments.

Figure 8

A

pCUP1-CDC5 BRN1-6HA
(YPA + 10 μM Cu) ——— 16 hr ——— SPM
(10 μM Cu)

7 hr → Sample collection

50 μM Cu

4 hr | 3 hr

B

| Brn1-6HA | DAPI | Merged |

10 μM Cu (normal expression)

50 μM Cu (over expression)

C

Number of Brn1-6HA foci / spread

p = 7.4 x 10⁻⁷

Cu concentration (μM)
10 μM | 50 μM

D

ctf19Δ
pCUP1-CDC5 BRN1-6HA
(YPA + 10 μM Cu) ——— 16 hr ——— SPM ——— 7-8 hr ——— Sample collection

0 μM Cu

10 μM Cu

E

| Brn1-6HA | DAPI | Merged |

0 μM Cu (no expression)

10 μM Cu (normal expression)

F

Number of Brn1-6HA foci / spread

p = 1.1 x 10⁻⁵

Cu concentration (μM)
0 μM | 10 μM

Figure 8. Misregulation of Cdc5 influences chromatin-condensin association.   (A)

Schematic representation of the experimental strategy. Wild type cells with homozygous *pCUP1-CDC5*

and *BRN1-6HA* alleles were grown in YPA supplemented with 10 µM copper used for normal expression

of *CDC5*. The cells were released into SPM supplemented with either 10 µM or 50 µM (used for over

expression of *CDC5*) copper, respectively. The samples were harvested at the indicated time point for

analysis by chromatin spread assay. (B) Representative images of the chromatin spreads showing

chromatin-Brn1-6HA association under normal or overexpression of *CDC5* in the wild type cells. Anti-HA

antibody (3F10) was used to detect Brn1-6HA. (C) Box plots showing the number of Brn1-6HA foci per

nuclear spread of the wild type cells. N = 60 from two independent experiments. (D, E, F) Same as in A, B

and C, respectively except *ctf19Δ* cells and 0 µM copper was used instead of wild type cells and 50 µM

copper, respectively. For F, N = 80 from two independent experiments. Scale bar = 2 µM. The statistical

significance *p* value was estimated by the two-tailed student's t-test for the mean.

1181

1182