Meta-analysis on the effects of chemical stressors on freshwater ecosystem functions

Alexander Feckler[1,2,*], Ralf Schulz[1,2], Ralf B. Schäfer[1]

[1]iES, Institute for Environmental Sciences, University of Kaiserslautern-Landau (RPTU),

Fortstraße 7, D-76829 Landau, Germany

[2]Eusserthal Ecosystem Research Station, University of Kaiserslautern-Landau (RPTU),

Birkenthalstraße 13, D-76857 Eusserthal, Germany

*Contact details for the corresponding author:

Name: Alexander Feckler

Address: iES, Institute for Environmental Sciences, University of Kaiserslautern-Landau

(RPTU), Fortstraße 7, 76829 Landau in der Pfalz, Germany

Telephone: +49 6341 28031312

Email: alexander.feckler@rptu.de

Abstract

Despite considerable progress in our predictive capacity for the response of different trophic

levels to chemical stressors, generalizable relations between chemical stressors and

ecosystem functions are lacking. We addressed this knowledge gap by conducting a meta-

analysis (153 studies; 350 observations) on the responses of freshwater ecosystem functions

(community respiration, organic matter decomposition, nutrient cycling, photosynthesis, and

primary production) under controlled conditions (laboratory or outdoor mesocosms) to

pesticides, pharmaceuticals, and metals. We identified monotonic dose-response relationships

between standardized chemical concentrations, in terms of toxic units, for selected chemical

use groups and organic matter decomposition by decomposer-detritivore-systems as well as

photosynthesis by algae and macrophytes. By contrast, consistent relationships were not

found for other ecosystem functions, such as organic matter decomposition by microbial

decomposers alone and primary production under the conditions studied. Importantly, the

shape and direction of the relationships identified here match those reported in field-based

studies, indicating a decrease in functioning as chemical stress increases, strengthening the

ecological relevance of our findings. Finally, we found a disconnect between regulatory

ecological quality targets and ecological outcomes, highlighting a need to re-evaluate risk

assessment approaches if they are supposed to be ecologically meaningful and protective of

ecosystem functions.

Keywords

chemical stress-ecosystem function, contaminants, ecological integrity, freshwater

ecosystems, pollution, sublethal effects, toxicants

1. Introduction

Freshwater habitats (streams, rivers, lakes, and ponds) host at least 9.5% of the currently described animal species biodiversity despite comprising just around 0.01% of the global water resources and covering 2.3% of the global surface area (Shiklomanov, 1993; Strayer and Dudgeon, 2010; Reid et al., 2019). Furthermore, they play a key role with respect to ecosystem functions. Approximately 20% of all land-based nitrogen sources are removed by denitrification in freshwaters (Seitzinger et al. 2006) and the transport and partial processing of carbon in streams and rivers is equivalent to about 25% of the estimated terrestrial net ecosystem production globally (Battin et al., 2023). Despite their paramount importance, freshwater habitats experience more pronounced declines in biodiversity compared to terrestrial ecosystems, leading to what has been labelled as the "freshwater crisis" (Harrison et al., 2018). The freshwater crisis is the result of an interplay of anthropogenic disturbances: over-exploitation, chemical pollution, alterations in water flow patterns, habitat destruction or degradation, and the introduction of non-native species (Reid et al., 2019). However, chemical pollution, except for nutrients, has often been marginalized in freshwater ecosystem studies, even though it can play a substantial role in deteriorating these ecosystems (Bernhardt et al., 2017). Extensive meta-analyses on continental and global scales consistently reveal that a substantial proportion of water bodies are affected by chemicals. In Europe, nearly half of the evaluated water bodies were identified as being at risk due to chronic chemical effects (Malaj et al., 2014). Similarly, on a global scale, insecticide contamination was observed to surpass thresholds indicating ecological impacts in nearly 70% of stream sites within agricultural landscapes (Stehle and Schulz, 2015). Recent investigations underscore the substantial level of chemical risk faced by freshwater organisms, both in Europe and the United States (Schulz et al., 2021; Wolfram et al., 2021).

Numerous studies showed that toxic chemicals can influence the organisms providing

pivotal ecosystem functions or the functions themselves, but their results remain scattered

and generalizations on relationships remain open. Meta-analyses and reviews (e.g., Crane et

al., 2006; Kümmerer, 2009; Schäfer et al., 2012b) showed that both microorganisms and

invertebrates, which are crucial for ecosystem functions like nutrient cycling, organic matter

decomposition, and primary production, are impacted by pharmaceuticals and pesticides at

environmentally relevant concentrations. Additionally, several meta-analyses and reviews

have drawn connections between effects of chemical stressors on specific groups of

organisms and changes in aquatic ecosystem functions these organism groups contribute to,

such as organic matter decomposition (Beaumelle et al., 2020; Feckler and Bundschuh, 2020;

Ferreira et al., 2016; Peters et al., 2013; Schäfer et al., 2012b), photosynthesis (Feckler et al.,

2015; Kovalakova et al., 2020), primary production (Feckler et al., 2015; Guo et al., 2015;

Kovalakova et al., 2020; Peters et al., 2013; Schäfer et al. 2012a), nutrient cycling (Peters et

al., 2013), and ecosystem respiration (Schäfer et al. 2012a). Recent studies also demonstrated

that chemicals from the same chemical type and with the same mode of toxic action (MoA)

exhibit qualitatively similar effects in communities (Halstead et al., 2014; Rumschlag et al.,

2019; Rumschlag et al., 2020). Thus, it can be hypothesized that the response patterns of

ecosystem functions towards chemicals from the same chemical type and MoA display a

generalizable relationship between the intensity of chemical stress and the rate of ecosystem

function. An earlier review by Peters et al. (2013) failed to establish consistent dose-response

relationships between various groups of toxic chemicals and several ecosystem functions,

such as nutrient cycling, organic matter decomposition and primary production. However,

these authors rather aimed at identifying effect thresholds and only considered studies that

exceeded a defined negative effect size instead of the full spectrum of responses of ecosystem

functions. Indeed, the responses of ecosystem functions to toxic chemicals can be negative

(e.g., Beaumelle et al., 2020; Robson et al., 2020; Rosi-Marshall and Royer, 2012), positive

(e.g., Corcoll et al., 2015; Richmond et al., 2019) or non-distinguishable from a control

treatment or a reference (e.g., Jepsen et al., 2019; Schäfer et al., 2012a). Thus, it remains

open whether an analysis of a broader number of studies, including novel studies that have

been conducted in the last decade, enables the derivation of generalizable relationships

between chemical stressors and ecosystem functions.

In the present study, we aimed to derive such generalizable relationships to help

predict chemical effects on ecosystems. Therefore, we conducted a meta-analysis and

extracted 350 observations (effect sizes and respective chemical concentrations) from 153

published studies conducted under controlled conditions (laboratory or outdoor mesocosms)

that cover a wide range of chemical use groups (metals, pesticides, and pharmaceuticals) and

five ecosystem functions (community respiration, organic matter decomposition, nutrient

cycling, photosynthesis, and primary production). In addition, we contextualized the observed

effects within an environmental risk assessment framework by calculating risk quotients

(RQs), relating the observed effects to regulatory acceptable concentrations (RACs) provided

by the German Federal Environment Agency . We hypothesized that where chemicals target

similar organism groups, generalizable relationships between the chemical stress intensity

(standardized by their toxic potency) and rate of ecosystem function can be established (H1).

Furthermore, we hypothesized that the relationship between chemical stress intensity and the

ecosystem function organic matter decomposition by microbial decomposers would be less

pronounced than for other functions studied (H2). This is because toxicity data for microbial

decomposers (bacteria and fungi) are largely lacking, hampering a reliable characterisation of

toxic potency towards them.

2. Materials and methods

2.1 Literature collection and inclusion criteria

We searched across Web of Science and Scopus to compile literature concerning the effects of chemical use groups (pesticides, pharmaceuticals, and metals) on five ecosystem functions (community respiration, nutrient cycling, organic matter decomposition, photosynthesis, and primary production). The search was limited to literature published between January 2000 and December 2024 to primarily focus on current-use pesticides and pharmaceuticals as well as metals. These chemical use groups were pre-selected because an initial screening showed that data would be too scarce for a meta-analysis of other chemical types, such as biocides. Our search string was constructed using a combination of freshwater habitat-related terms (freshwater* OR stream* OR river* OR pond* OR lake*), chemical stressor-related terms (chemical stress* OR contaminant* OR pollutant* OR toxicant* OR pesticide* OR heavy metal* OR metal* OR fungicide* OR herbicide* OR insecticide* OR pharmaceut*), and ecosystem function-related terms (ecosystem function* OR photosynth* OR primary product* OR nutrient cycl* OR organic matter decomp* OR leaf breakdown OR decomposition* OR respirat*). The asterisks (*) were employed as wildcards, capturing all terms sharing the same root. Moreover, we conducted footnote chasing by screening the reference lists of relevant articles for further literature within the considered time period. The selection process for studies encompassed three stages. Initially, titles were screened to identify suitable studies. Subsequently, the abstracts were evaluated, leading to the exclusion of non-matching studies. Lastly, the full texts of selected studies were assessed.

Studies were included based on a set of criteria: (1) laboratory-based studies and outdoor mesocosm studies; we deliberately excluded field studies from further analyses, as these represented a minor share of our dataset ($n = 60$ across all ecosystem functions), of which a major fraction (>90%) did not meet one or multiple of the following inclusion

criteria, (2) focus on species, populations, or communities dwelling within freshwater habitats (excluding groundwater, marshlands, coastal waters, and marine systems), (3) experimental exposure to at least one of the pre-defined chemical types (fungicides, herbicides, insecticides, pharmaceuticals, and metals) from the list of chemical use groups above (excluding eutrophication, salinization, and acidification), (4) presence of a control treatment (absence of chemical stress), and (5) measurement of the designated ecosystem functions as well as uncertainty around the central tendency of the measurement. Furthermore, we only included studies that addressed direct effects on the organisms under scrutiny, thereby excluding indirect effects (e.g. shifts in the palatability of leaf material for detritivores due to direct effects on leaf-associated microorganisms). We excluded studies on teleost fish, as they do not contribute directly to the ecosystem functions examined here. An overview of the selected studies and moderator variables (see below) will be provided on Zenodo upon article acceptance.

2.2 Data extraction and effect size calculation

We extracted central tendencies, uncertainty estimators, sample sizes, chemical concentrations, and moderator variables (see below) from the text, tables, and figures of the selected studies. For cases where the data were presented in figures, we utilized the "digitize" package (Poisot, 2011) within the open-source software R (version 4.4.0 for Mac OS X; R Core Team, R Core Team, 2024).

As our response variable, we calculated the ratio of the ecosystem function observed in the stressor treatment to that in the control treatment. From each study, we extracted only the results for the concentration with the highest effect size observed per functional endpoint for the same species, population, or community. This approach was adopted to derive unequivocal cause-effect relationships and prevent pseudoreplication. Due to this method and

the diverse test designs employed in individual studies, the timing of functional

measurements varied.

For studies investigating community respiration, the functional indicators included

measurements of O_2-consumption (e.g., mg O_2 g^{-1} dry weight h^{-1}) and CO_2-production (e.g.,

µg CO_2 g^{-1} dry weight h^{-1}). In the context of organic matter decomposition, the indicators

consisted of breakdown rates (k (d^{-1})) and mass loss (e.g., mg d^{-1}) of leaf material, mediated

by microbial decomposers or a combination of microbial decomposers and detritivorous

macroinvertebrates (decomposer-detritivore-systems) as endpoints. As the literature search

on nutrient cycling only resulted in data for nitrogen cycling, the indicators encompassed

anaerobic NH_4^+ oxidation rates (g-N g-volatile suspended solids^{-1} d^{-1}), NO_2- and NO_3-

turnover rates (e.g., mg NO_2-N mg^{-1} d^{-1}), and N_2-production (e.g., N_2 g^{-1} h^{-1}). In the case of

photosynthesis, we determined effect sizes based on maximum and effective photosystem II

quantum yields (F_v/F_m and ΦPSII, respectively) and O_2-production (e.g., mg O_2 g^{-1} dry

weight h^{-1}). Finally, for primary production, the effect sizes related to fixed carbon quantities

(mg ^{14}C g^{-1} C h^{-1}) and changes in the standing biomass (e.g., g C h^{-1}). Studies that exclusively

reported dissolved O_2 as a response variable were excluded, as net dissolved O_2 could arise

from multiple sources simultaneously, such as photosynthesis and the ambient atmosphere.

2.3 Calculation of toxic units and risk quotients

The toxic effects caused by different chemical stressors vary for a given test

concentration, due to differing ecotoxicological potentials, making direct comparisons

challenging. Ideally, we would have used the HC_{50} – the hazardous concentration of a

stressor at which 50% of the species are exposed to a concentration above their EC_{50} (the

concentration at which an effect is observed in 50% of exposed organisms) derived from a

species sensitivity distribution (SSD) – as a benchmark for the toxicity of individual stressors.

However, the calculation of SSDs is not reliable if less than 10 data points are available (Hiki

and Iwasaki, 2020), which was the case for >90% (especially for macrophytes) of substances.

Therefore, we normalized the test concentrations of individual chemical stressors to their

acute effects by calculating toxic units (TUs). Ideally, TUs would be computed for each

combination of chemical stressor and organism included in the selected literature. However,

given substantial data gaps, we relied on surrogate species results to facilitate comparisons

between different stressors. Unlike previous studies that sought to establish generalized

chemical stress–ecosystem function relationships (e.g., Peters et al., 2013), we avoided

confining the surrogate species selection to conventional test species as stipulated by

chemical regulations (i.e., *Daphnia magna*, *Lemna* sp., and *Raphidocelis subcapitata*).

Instead, we estimated stressor toxicity based on the EC_{50} values for the most sensitive tested

freshwater algal, plant, and invertebrate species (not necessarily the conventional test species

used for risk assessment) for the respective chemical stressor, because this approach was

superior to using a surrogate species in a previous comparison (Schäfer et al., 2013).

Notwithstanding, this approach may incur some variance given its dependency on the testing

effort for each chemical and the selected test species available in the literature. We employed

these EC_{50} values to compute toxic units (TU) defined here as the logarithmic ratio between

experimental exposure concentrations and a defined effective concentration:

$$TU = \log\left(\frac{c_i}{EC50_i}\right)$$

206 where *c* represents the concentration of a toxicant *i* and $EC50_i$ is the median effective

207 concentration of the respective toxicant for the most sensitive species. In case of chemical

208 mixtures (23% of cases), we employed the EC_{50} values to compute sum toxic units (sumTU)

209 defined as the logarithmic sum of the ratios between exposure concentrations and a defined

210 effective concentration:

$$sumTU = \log\left(\sum_{i=1}^{n} \frac{c_i}{EC50_i}\right)$$

where n gives the number of toxicants that caused an effect in the respective ecosystem

function. Although the assumption of concentration addition underlying the sumTU concept

cannot be guaranteed – given that mixture components often have distinct MoAs – previous

research suggests that the sumTU can reliably represent mixture toxicity, even when MoAs

differ (cf. Escher et al., 2020). EC_{50} values were extracted from various sources including

Standartox (Scharmüller et al., 2020), the Pesticide Property Database (Lewis et al., 2016),

the Veterinary Substance Database (Lewis et al., 2011), the Norman Database

(https://www.norman-network.com/nds/), and WikiPharma (Molander et al., 2009), with

most of the EC_{50} being related to mortality (invertebrates) or growth (algae, macrophytes).

We primarily extracted EC_{50} values corresponding to a 48-hour exposure duration or, when

not available, the closest exposure time (algae: 15-120 h, invertebrates: 48-96 h,

macrophytes: 15-120 h).

In addition, we extracted RACs for pesticides – used as benchmarks in environmental

risk assessment – from the Information System Ecotoxicology and Environmental Quality

Targets (ETOX), provided by the German Federal Environment Agency. We selected

German data for its comprehensiveness and to avoid mixing data from different agencies,

which may derive RACs with slightly different approaches, thereby incurring variability.

RQs were calculated as the ratio between exposure concentrations and the respective RAC:

$$RQ = \frac{c_i}{RAC_i}$$

where c represents the concentration of a toxicant i and RAC_i is the corresponding regulatory

acceptable concentration. In case of toxicant mixtures, we employed the combined risk

quotient that sums up the RQs of individual pesticides, assuming that the effects of mixtures

are additive. As such benchmarks are lacking for pharmaceuticals and metals, RQ

calculations could not be performed for these chemical use groups. A RQ > 1 means that the

exposure exceeded the RAC. All EC_{50} values, the respective species, RACs, measured

endpoints and their sources will be provided on Zenodo upon article acceptance.

2.4 Moderator variables

Besides central tendencies, uncertainty estimators, sample sizes, and chemical

concentrations, we extracted seven moderator variables: (1) chemical type with levels: (a)

fungicide, (b) herbicide, (c) insecticide, (d) pesticide mixture, (e) pharmaceutical, (f) metal,

(g) metal mixture, and (h) other (such as biocides); (2) classification of the MoAs extracted

from Lewis et al. (2016) or Molander et al. (2009); (3) organism group with levels: (a) algae

(including cyanobacteria and phytoplankton), (b) macrophytes, (c) microbial decomposers

(bacteria and fungi), (d) joint systems of microbial decomposers and invertebrate

decomposers, hereafter termed decomposer-detritivore-systems; (4) exposure time as a

continuous variable in days; (5) pH as a continuous variable; (6) test temperature as a

continuous variable in °C; (7) study system with levels: (a) laboratory and (b) mesocosm.

2.5 Data analysis

To identify the moderator variables affecting the assessed ecosystem functions and

establish relationships between chemical stressors and ecosystem functions, we performed

separate data analyses for each of the included ecosystem functions. We used the calculated

effect sizes (along with the respective standard errors of the effects) as the response variables,

while including the sumTUs and moderator variables as predictors. Initially, we checked for

multi-collinearity by examining correlations ($r \geq 0.8$) among our predictors, ensuring that the

meta-regression coefficient estimates are robust. Subsequently, we identified the best-fitting

models using the "multimodel.inference" function in the R package "dmetar" (Harrer et al.,

2019) based on corrected Akaike Information Criteria. Using the results of the multimodel

inference, we build models to evaluate the individual impact of the identified predictors and

their interactions. We achieved this by fitting meta-regression models using the R package

"metafor" (Viechtbauer, 2010). We employed the "maximum likelihood" estimator to assess

the amount of residual heterogeneity (τ^2) and applied the Knapp-Hartung method for random-

effects meta-analyses (van Aert and Jackson, 2019). We conducted a permutation test running

1,000 iterations before reporting *p*-values, as recommended for meta-regression models, to

control for type I error rates of frequentist meta-regressions with small sample sizes (Higgins

and Thompson, 2004).

Afterwards, we fitted (quasi-)binomial Generalized Linear Models (GLMs) to explore

potential dose-response relationships between ecosystem functions and sumTUs as well as

RQs. Models were fitted for all stressors combined or separately for chemical types

(fungicides, herbicides, insecticides, pharmaceuticals, and metals) or MoAs (only if the

number of observations n ≥ 8, given the statistical limitations of small sample sizes; Jenkins

and Quintana-Ascencio, 2020). To exclude the influence of extreme data points (such as very

low or very high responses of ecosystem functions), we employed bootstrapping (1,000

times). Models were then run on the individual bootstrapped data sets, and we calculated the

model medians to represent the central tendency of the fitted models. For statistical analyses

and graphics, we used the open-source software R (R Core Team, 2024), along with the

necessary add-on packages. Data and code will be freely available on Zenodo upon

acceptance.

3. Results

3.1 Overview of included studies

Through our database query and footnote chasing, we identified a total of 7,661

publications, of which 7,508 publications were excluded due to violations of our predefined

inclusion criteria, absence of reported effect sizes and associated uncertainties, or the inability

to calculate sumTUs due to a lack of toxicity data. Ultimately, we focused on 153 studies that

met our criteria, from which we extracted 350 effect sizes (# of observations), uncertainty

estimators, sumTUs, and moderator variables (see Fig. A1 for details). Among the selected

studies, 37% of observations each focused on organic matter decomposition and

photosynthesis, while 21% addressed primary production. Due to limited representation

(5%), community respiration and nutrient cycling were excluded from further analysis.

Pesticides were the most frequently studied chemical stressors (66%, n=232), comprising

fungicides (n=67), herbicides (n=86), insecticides (n=43), and pesticide mixtures (n=36).

Metals and their mixtures accounted for 26% (n=91), and pharmaceuticals for 8% (n=27; Fig.

1). SumTU values ranged from –5.13 to 2.68 (metals), –7.60 to 3.99 (pesticides), and –5.68

to 5.59 (pharmaceuticals), with 0 indicating toxicity equivalent to the EC50 of the most

sensitive species (Table 1).

Table 1: Descriptive statistics on sumTU values for the individual chemical types ($n = 350$

effects sizes or # of observations).

Chemical type	n	Mean	Min	1. Quartile	2. Quartile (Median)	3. Quartile	4. Quartile
Fungicides	67	-0.01	-7.60	-0.82	0.11	0.59	2.31
Herbicides	86	1.26	-3.03	0.87	1.39	1.57	3.99
Insecticides	43	0.34	-4.93	-0.90	0.75	2	3.60
Pesticide mixtures	36	0.09	-2.31	-1.00	0.08	0.94	2.78
Pharmaceuticals	27	0.08	-5.68	-1.05	0.26	0.87	5.59

Metals	81	0.33	-5.13	-0.49	0.56	1.66	2.68
Metal mixtures	10	0.86	-1.26	0.55	0.99	1.43	1.77

3.2 Influence of moderator variables and relationships between sumTU and effects on

ecosystem functions

3.2.1 Organic matter decomposition

For the ecosystem function organic matter decomposition, our best-fit model included

two significant predictors: "organism group" (p = 0.001), which categorized microbial

decomposers alone and decomposer-detritivore-systems, and "pH" (p = 0.007). Additionally,

"organism group" and "sumTU" exhibited a significant interaction (p = 0.039). This model

accounted for a substantial portion of the variance (meta-regression: R^2 = 58.93%, τ^2 =

0.0434; Table 2). We observed a general dose-response relationship for organic matter

decomposition across all groups of chemicals for studies examining decomposer-detritivore

systems (as indicated by the GLM for sumTU and the respective ecosystem function; p =

0.022; n = 93, Fig. 2a). However, when focusing on microbial decomposers alone, we did not

find a consistent relationship between sumTU and organic matter decomposition (p = 0.801;

n = 82; Fig. 2b), nor did we find such a relationship for any individual chemical type or MoA

($p \geq$ 0.288; n = 8-22).

3.2.2 Photosynthesis

For the ecosystem function photosynthesis, our best-fit model included two

significant predictors: "sumTU" (p = 0.001) and "temperature" (p = 0.023), collectively

explaining a portion of the variance (R^2 = 18.22%, τ^2 = 0.0707; Table 2). In contrast to

organic matter decomposition, we did not observe an influence of the parameter "organism

group" (distinguishing between algae and macrophytes; Fig. 2c&d). When we examined the

effects of individual chemical types or MoA on specific organism groups, no clear dose-

response relationships were evident ($p \geq$ 0.210; n = 12-50).

3.2.3 Primary production

For the ecosystem function primary production, our best-fit model included "pH" ($p =$

0.008) as significant predictor, explaining a portion of the variance (R^2 = 16.06%, $\tau^2 =$

0.0437; Table 2), but not "sumTU". When we examined the effects of individual chemical

types or MoA on algae (lack of data on macrophytes), no clear dose-response relationships

were   evident   (p $\geq$ 0.285; n $=$ 12-24).

Table 2: Outcome of meta-regression models assessing the influence of moderator variables on ecosystem functions.

Function	Predictor	Estimate	T-value	df[1]	p-value	Lower CI[2]	Upper CI
Organic matter decomposition	Organism group (decomposer vs. decomposer-detritivore)	0.2549	6.2973	171	0.001	0.1750	0.3348
	sumTU	0.0111	0.4160	171	0.723	-0.0415	0.0637
	pH	-0.2455	-2.7898	171	0.007	-0.4205	-0.0705
	Organism group $\times$ sumTU	0.0714	2.4485	171	0.039	0.0138	0.1290
Photosynthesis	sumTU	0.0877	4.4305	99	0.001	0.0484	0.1270
	Temperature	-0.0221	-2.8695	99	0.019	-0.0374	-0.0068
Primary production	pH	0.2932	2.8056	42	0.008	0.0823	0.5041

[1]df = Degrees of freedom, [2]CI = confidence interval

3.3 Relationships between risk quotients for regulatory thresholds and effects on ecosystem

functions

We did not find a general relationship between RQs and ecosystem functioning

related to organic matter decomposition across all chemical groups, neither when focusing on

decomposer–detritivore systems (GLM of $\log_{10}$(RQ) and the respective ecosystem

function; p = 0.933; n = 71) nor when focusing on decomposers only (p = 0.335; n = 41). For

photosynthesis, a general relationship was found for algae (p = 0.008; n = 35), whereas no

such relationship was observed for studies examining primary production (p = 0.989; n = 15;

Fig. 3).

4. Discussion

4.1 Influence of moderator variables and relationships between sumTU and effects on

ecosystem functions

To the best of our knowledge, our study is among the first in establishing

relationships between chemical stressors and ecosystem functions. This represents a crucial

stride, as such relationships can ultimately aid in predicting the toxicity of chemicals that

have not yet been thoroughly studied, although some degree of uncertainty remains due to the

broad range of MoAs. Nevertheless, the transferability and predictive capacity of these

relationships could potentially reduce costs and efforts, diminish the reliance on animal

testing, and mitigate the suffering of living organisms – an issue that has received

considerable attention in the last decade (Burden et al., 2015).

In accordance with our first hypothesis (H1), we found a few generalizable

relationships between the toxic potency of chemicals that target similar organism groups and

the rate of ecosystem functions. These relationships exhibited an inverted S-shaped

monotonicity, signifying that the measured parameters declined as the toxic potency of the

chemical stressors increased. Our findings therefore align with the dose-response patterns frequently observed in the field of ecotoxicology (Newman, 2009). The S-shape observed in these relationships suggests that organisms carry specific defense or repair mechanisms that provide protection against a certain amount of a chemical stressor, up to a critical threshold level. Beyond this threshold, organisms become incapable of defending or repairing themselves effectively. Consequently, the biological response to chemical stressors showed a graded response due to the evolutionary adaptive capabilities of organisms to react to a range of low doses. The absence of relationships between toxic potency and ecosystem functioning apart from those identified for organic matter decomposition and photosynthesis may be explained by three factors. First, the toxic pressure from certain chemical groups may have been below thresholds triggering adverse effects. Second, bias related to our toxicity estimation particularly for microbial decomposers may have obscured relationships (H2). Third, given considerable variation in the obtained regression relationships, incurred by pooling studies with very different conditions and chemicals with differing MoAs, may have hidden a potential relationship.

For the ecosystem function organic matter decomposition by decomposer-detritivore-systems, we observed a 50% reduction occurring at a sumTU of approximately 0. This finding aligns with a field study that observed a substantial decrease in organic matter decomposition by decomposer-detritivore-systems of approximately 50% at a an even lower $\log(TU_{D. magna})$ of 0.01 (Schäfer et al., 2012a). Consequently, there is a good overlap of field data and the relationships between chemical exposures and their effect on organic matter decomposition observed in our study, despite the inherent challenges of comparing data sets – for instance studies mainly focused on individual chemicals in laboratory or mesocosm settings in comparison to complex mixtures in the field – highlighting the value of our results.

Despite this good overlap of field data and the chemical stress-ecosystem function

relationship derived here, it is essential to acknowledge that the heavy emphasis on specific

chemical use groups in research often hampers the broader applicability of laboratory

findings to real-world conditions. Over the past 25 years, a predominant focus has been on

metals, pesticides (particularly herbicides), and pharmaceuticals. However, these chemical

use groups constitute only a small fraction of the vast array of chemicals present in the

environment, given that approximately 350,000 chemicals are registered for production and

use (Wang et al., 2020). Studies employing bioassays have revealed that target chemicals can

only account for a small fraction of the overall response to chemicals (as low as < 0.1 %;

Escher et al., 2013), suggesting that numerous ecotoxicologically important compounds exist

within freshwater ecosystems (Escher et al., 2013; Neale et al., 2020). In addition to the

emphasis on pesticides, metals, and pharmaceuticals, only around one quarter of the included

studies have considered that chemicals often co-occur as mixtures. Typically, the effect of

chemicals in mixtures is additive, but in a relevant proportion of cases, the simultaneous

presence of multiple chemicals can result in stronger (synergistic) or weaker (antagonistic)

effects compared to the assumption of additive effects (cf. Warne and Hawker, 1995). A

recent review, encompassing various chemical groups, indicated an average of 20% of cases

with synergism and 15% of cases with antagonism (Martin et al., 2021). Another review

reported synergistic effects in 3% to 26% of cases involving binary mixtures, with the

prevalence varying by chemical group (Cedergreen, 2014). Furthermore, freshwater

ecosystems are typically subject to multiple stressors concurrently (Birk et al., 2020), which

can lead to interactions between chemical and non-chemical stressors (Liess et al., 2016). To

provide a more comprehensive understanding of chemical effects in freshwater ecosystems

and derive additional chemical stress-ecosystem function relationships, it is essential to

broaden the scope. This entails identifying ecotoxicologically relevant chemical groups,

assessing interactions within chemical mixtures, and considering the context of multiple

(non-chemical) stressors in the ecosystem.

Furthermore, finding chemical stressor-ecosystem function relationships can be

confounded by abiotic factors, such as pH (Ragnarsdottir, 2000; Wang et al., 2016) and

temperature (Soares et al., 2020), which influence the fate and toxicity of metals and organic

chemicals in aquatic environments. This phenomenon is often referred to as "context

dependency" (Catford et al., 2022). While pH and temperature had direct effects on

ecosystem functions, which are discussed below, no interactions between pH or temperature

and other moderator variables such as the toxic potency were observed. However, such

interactions have been observed earlier. For instance, a meta-analysis revealed a substantial

influence of pH on metal toxicity (Wang et al., 2016). This can be explained by the pH-

dependency of metal speciation, which in turn affects metal bioavailability. Furthermore,

pesticides can be ionizable compounds (Ragnarsdottir, 2000), and their degradation and

subsequent toxicity are influenced by pH as well as molecular structural properties (Fenner et

al., 2013). Temperature also influences the effects of chemical stressors through two key

concepts. The "climate-induced toxicant sensitivity" (CITS) concept describes how climate-

related environmental factors alter chemical toxicity, while the "toxicant-induced climate

change sensitivity" (TICS) concept explains how pesticide exposure increases susceptibility

to climate change (Verheyen et al., 2022). These concepts are interconnected, as global

warming may enhance pesticide toxicity (CITS), which in turn could further reduce heat

tolerance (TICS; Verheyen et al., 2022). Moreover, aside from their role in modifying

chemical bioavailability, speciation, and concentration, abiotic conditions have been shown

to increase species sensitivity by up to 30-fold in certain experiments (Liess et al., 2001;

Liess and Beketov, 2011).

431 Additionally, biotic factors can impede the derivation of chemical stressor-ecosystem

432 function relationships. However, the relevance of these factors largely depends on the

433 ecosystem function under investigation and the design of the experimental approach.

434 Laboratory experiments typically involve the use of species that are maintained in culture

435 over extended periods, often under conditions that differ from their natural habitats (e.g.,

436 variations in light, temperature, and pH) and that may lay outside their niche width and

437 optima. Thus, these conditions affect the survival, growth, reproduction, and functioning of

438 species, and thereby the related ecosystem functions (Blackman, 1905). Indeed, pH and

439 temperature significantly influenced organic matter decomposition, photosynthesis, and

440 primary production. These findings suggest a mismatch between species-specific

441 requirements and the laboratory conditions under which the respective experiments were

442 conducted. In studies that make use of complex natural communities rather than single

443 species tests (such as microbially-mediated organic matter decomposition), additional biotic

444 factors can introduce uncertainties into the derivation of chemical stressor-ecosystem

445 function relationships. These studies typically involve communities with different

446 compositions, and these structural differences have been demonstrated to influence the

447 sensitivity of communities to chemical stress (Beketov et al., 2008; Feckler et al., 2018), for

448 instance due to functional similarity among species that has not been considered in the

449 present study (Louca et al., 2018). Consequently, these variations can lead to differing effects

450 on ecosystem functions due to differences in the capacity of communities to compensate for

451 species loss caused by chemical stressors (Cadotte et al., 2011).

453 *4.2 Study bias and confounding factors in the present meta-analysis*

454 We specifically chose biologically-mediated ecosystem functions for our analysis,

455 focusing on those we deemed most ecologically relevant within freshwater ecosystems.

Among the five selected ecosystem functions, primary production and photosynthesis by

algae and macrophytes were notably prevalent. This prevalence can be attributed to several

factors. First, these functions are likely to be highly responsive to the most studied group of

pesticides, namely herbicides (Hedlund et al., 2020), as well as pharmaceuticals.

Cyanobacteria, as photosynthetic prokaryotes, are considered sensitive to antibiotics due to

their close relationship with pathogenic bacteria (González-Pleiter et al., 2013), and green

algae and macrophytes can also be affected by antibiotics because of the prokaryotic origin of

their semi-autonomous cell organelles (Nie et al., 2009). Second, maintaining algal cultures

and macrophytes in laboratory settings is relatively straightforward, and measurement

techniques for primary production and photosynthesis enable rapid quantitative assessment.

Despite their importance for freshwater ecosystem functioning, their potential of

being used as indicators of streams' ecological status and in the assessment of streams'

restoration (Klocker et al., 2009; Udy et al., 2006; Young et al., 2008), only a minor share of

studies addressed chemical stressor effects on microbial community respiration and nutrient

cycling. Unlike for algae, standardized protocols for culturing heterotrophic microbial

communities (bacteria and fungi) in the laboratory are largely lacking, hindering their use in

research and complicating comparisons between studies. Moreover, studies assessing

ecosystem functions through microbial decomposers (approximately 24 ± 2 days; mean ±

standard error) required considerably longer exposure times compared to those focusing on

primary production and photosynthesis by algae (approximately 5 ± 0.5 days), increasing the

labor-intensity. Nevertheless, protocols to quantify various processes related to community

respiration and nutrient cycling have been developed and are frequently applied in field

studies to assess stream ecosystem health (von Schiller et al., 2017). While these protocols

are similarly complex and costly as other methods for quantifying ecosystem functions, they

could be partially automated for use in laboratory studies (von Schiller et al., 2017). Thus, we

recommend their application in future laboratory studies to broaden our mechanistic

understanding of how chemical stressors impact ecosystem functions.

Lastly, the taxonomic mismatch between the most sensitive tested species used for

calculating TUs and the species assessed for the chemical stress–ecosystem function

relationship likely introduced some uncertainty. However, we deliberately used the most

sensitive reported species for each chemical stressor to provide a conservative estimate of

their effects on ecosystem functions. Given that these sensitive species are present in

freshwater communities and are affected by chemical exposure, the observed effects are

expected to be comparable to those identified in this meta-analysis (cf. Schäfer et al., 2012b).

4.3 Relationships between risk quotients for regulatory thresholds and effects on ecosystem

functions

Only one of four relationships between RQs of RACs and effects on ecosystem

functions was significant. Ideally, ecosystem functions would be at control levels for RQs < 1

and only decline when this threshold is exceeded, equaling the exceedance of RACs. The fact

that ecosystem functions were frequently much lower than control levels for RQs < 1 and that

only one consistent relationship could be established questions the protectiveness of RACs

for ecosystem functions and their accuracy in reflecting real-world ecological dynamics.

Several factors likely explain this shortcoming of RACs. First, RACs are derived from

toxicity data that relate to single-species laboratory or multispecies mesocosm studies. The

relationship between single-species experiments and ecosystem functions can vary strongly

because of variability in the contribution of the single species to the function under scrutiny

and complex interspecies interactions not accounted for (Kraft et al., 2015). Likewise, the

responses to chemicals can vary strongly between mesocosm studies due to experimental

conditions (e.g., Beketov and Lies, 2008) and the most sensitive response variable selected

for RAC derivation may vary regarding its relationship with ecosystem functioning. Second,

the thresholds derived from current risk assessments for single species or populations may

underestimate functional impairments originating from sublethal effects. Consequently, key

ecosystem processes may be disrupted even when stressor concentrations remain below

levels deemed "acceptable" for non-functional responses. Third, the relationship between risk

quotients for RAC and ecosystem functioning may be nonlinear and moderated by non-

additive mixture effects that are not adequately captured by current risk assessment

approaches.

5. Conclusions

Along with the ever-increasing production values and usage of new chemicals, it

becomes insurmountable to experimentally study the effects of all chemicals alone and in

combination with other chemicals and stressors in a wide range of ecosystems, making

generalizations invaluable. In this context, our analyses revealed the potential of deriving

chemical stress-ecosystem function relationships using currently available data that show

adverse effects for certain ecosystem functions. Nevertheless, we identified several important

research gaps – partially beyond the scope of the present study – that should be considered in

the future to support deriving more specific (e.g., in terms of MoA) chemical stress-

ecosystem function relationships and ultimately facilitate the prediction of chemical effects.

As we showed in our analysis, chemicals of certain groups as well as specific ecosystem

functions were underrepresented in the extracted literature published between 2000 and 2024.

Therefore, data likely remains insufficient for important but understudied chemical groups

and ecosystem functions, hampering a more complete evaluation of chemical stress.

Furthermore, we showed that the context-dependency of chemical stress effects on ecosystem

functions is largely ignored. Therefore, future laboratory research should proactively consider

a series of abiotic and biotic factors as well as the composition of stressors (single stressors

vs. diverse chemical mixtures; combination of chemical and non-chemical stressors) to

ensure the transferability of the data to various field situations. Ideally, such studies should

simulate different climatic and biogeographical regions as well as chemical exposure

histories as these factors were shown to determine the stress response (Zubrod et al., 2019;

Sinclair et al., 2024). In addition, laboratory studies should be appended by semi-field and

field studies as verification and validation benchmarks to assess the accuracy and reliability

of laboratory data for field situations. Finally, there is a disconnect between regulatory

ecological quality targets and ecological outcomes, highlighting a need to re-evaluate risk

assessment approaches if they are supposed to be ecologically meaningful and protective of

ecosystem functions.

Funding statement

This work was funded by the EU grant HORIZON-HLTH-2021-ENVHLTH-03-01 -

European partnership for the assessment of risks from chemicals (PARC) Grant agreement

ID: 101057014. Alexander Feckler was funded by the Strategy Fund of the University of

Kaiserslautern-Landau.

Data accessibility statement

All data used for the analyses of the present study were extracted from previously published

articles. All data and R code used in this paper will be uploaded to Zenodo upon article

acceptance.

References

References

Balian, E.V., Segers, H., Martens, K., Lévéque, C., 2008. The Freshwater Animal Diversity
Assessment: an overview of the results, in: Balian, E.V., Lévêque, C., Segers, H.,
Martens, K. (Eds.), Freshwater Animal Diversity Assessment, Developments in
Hydrobiology. Springer Netherlands, Dordrecht, pp. 627–637.
https://doi.org/10.1007/978-1-4020-8259-7_61

Battin, T.J., Lauerwald, R., Bernhardt, E.S., Bertuzzo, E., Gener, L.G., Hall, R.O., Hotchkiss,
E.R., Maavara, T., Pavelsky, T.M., Ran, L., Raymond, P., Rosentreter, J.A., Regnier,
P., 2023. River ecosystem metabolism and carbon biogeochemistry in a changing
world. Nature 613, 449–459. https://doi.org/10.1038/s41586-022-05500-8

Beaumelle, L., De Laender, F., Eisenhauer, N., 2020. Biodiversity mediates the effects of
stressors but not nutrients on litter decomposition. eLife 9, e55659.
https://doi.org/10.7554/eLife.55659

Beketov, M.A., Liess, M., 2008. Variability of pesticide exposure in a stream mesocosm
system: Macrophyte-dominated vs. non-vegetated sections. Environmental Pollution
156, 1364–1367. https://doi.org/10.1016/j.envpol.2008.08.014

Beketov, M.A., Schäfer, R.B., Marwitz, A., Paschke, A., Liess, M., 2008. Long-term stream
invertebrate community alterations induced by the insecticide thiacloprid: Effect
concentrations and recovery dynamics. Sci. Total Environ. 405, 96–108.
https://doi.org/10.1016/j.scitotenv.2008.07.001

Bernhardt, E.S., Rosi, E.J., Gessner, M.O., 2017. Synthetic chemicals as agents of global
change. Front. Ecol. Environ. 15, 84–90. https://doi.org/10.1002/fee.1450

Birk, S., Chapman, D., Carvalho, L., Spears, B.M., Andersen, H.E., Argillier, C., Auer, S.,
Baattrup-Pedersen, A., Banin, L., Beklioğlu, M., Bondar-Kunze, E., Borja, A.,
Branco, P., Bucak, T., Buijse, A.D., Cardoso, A.C., Couture, R.-M., Cremona, F., de
Zwart, D., Feld, C.K., Ferreira, M.T., Feuchtmayr, H., Gessner, M.O., Gieswein, A.,
Globevnik, L., Graeber, D., Graf, W., Gutiérrez-Cánovas, C., Hanganu, J., Işkın, U.,
Järvinen, M., Jeppesen, E., Kotamäki, N., Kuijper, M., Lemm, J.U., Lu, S., Solheim,
A.L., Mischke, U., Moe, S.J., Nõges, P., Nõges, T., Ormerod, S.J., Panagopoulos, Y.,
Phillips, G., Posthuma, L., Pouso, S., Prudhomme, C., Rankinen, K., Rasmussen, J.J.,
Richardson, J., Sagouis, A., Santos, J.M., Schäfer, R.B., Schinegger, R., Schmutz, S.,
Schneider, S.C., Schülting, L., Segurado, P., Stefanidis, K., Sures, B., Thackeray, S.J.,
Turunen, J., Uyarra, M.C., Venohr, M., von der Ohe, P.C., Willby, N., Hering, D.,

2020. Impacts of multiple stressors on freshwater biota across spatial scales and
ecosystems. Nat. Ecol. Evol. 4, 1060–1068. https://doi.org/10.1038/s41559-020-1216-
4

Blackman, F.F., 1905. Optima and limiting factors. Ann. Bot. 19, 281–295.

Burden, N., Sewell, F., Chapman, K., 2015. Testing chemical safety: what is needed to ensure
the widespread application of non-animal approaches? PLOS Biology 13, e1002156.
https://doi.org/10.1371/journal.pbio.1002156

Cadotte, M.W., Carscadden, K., Mirotchnick, N., 2011. Beyond species: functional diversity
and the maintenance of ecological processes and services. J. Appl. Ecol. 48, 1079–
1087. https://doi.org/10.1111/j.1365-2664.2011.02048.x

Cardinale, B.J., Duffy, J.E., Gonzalez, A., Hooper, D.U., Perrings, C., Venail, P., Narwani,
A., Mace, G.M., Tilman, D., Wardle, D.A., Kinzig, A.P., Daily, G.C., Loreau, M.,
Grace, J.B., Larigauderie, A., Srivastava, D.S., Naeem, S., 2012. Biodiversity loss and
its impact on humanity. Nature 486, 59–67. https://doi.org/10.1038/nature11148

Catford, J.A., Wilson, J.R.U., Pyšek, P., Hulme, P.E., Duncan, R.P., 2022. Addressing
context dependence in ecology. Trends Ecol. Evol. 37, 158–170.
https://doi.org/10.1016/j.tree.2021.09.007

Cedergreen, N., 2014. Quantifying synergy: a systematic review of mixture toxicity studies
within environmental toxicology. PLOS ONE 9, e96580.
https://doi.org/10.1371/journal.pone.0096580

Corcoll, N., Casellas, M., Huerta, B., Guasch, H., Acuña, V., Rodríguez-Mozaz, S., Serra-
Compte, A., Barceló, D., Sabater, S., 2015. Effects of flow intermittency and
pharmaceutical exposure on the structure and metabolism of stream biofilms.
Sci.Total Environ. 503–504, 159–170. https://doi.org/10.1016/j.scitotenv.2014.06.093

Crane, M., Watts, C., Boucard, T., 2006. Chronic aquatic environmental risks from exposure
to human pharmaceuticals. Sci. Total Environ. 367, 23–41.
https://doi.org/10.1016/j.scitotenv.2006.04.010

Dirzo, R., Young, H.S., Galetti, M., Ceballos, G., Isaak, N.J.B., Collen, B., 2014.
Defaunation in the Anthropocene. Science 345, 401-406.
https://10.1126/science.1251817

Escher, B.I., van Daele, C., Dutt, M., Tang, J.Y.M., Altenburger, R., 2013. Most oxidative
stress response in water samples comes from unknown chemicals: the need for effect-
based water quality trigger values. Environ. Sci. Technol. 47, 7002–7011.
https://doi.org/10.1021/es304793h

Escher, B., Braun, G., Zarfl, C., 2020. Exploring the Concepts of Concentration Addition and
621   Independent Action Using a Linear Low-Effect Mixture Model. Environmental
622   Toxicology and Chemistry 39, 2552–2559. https://doi.org/10.1002/etc.4868

624   Feckler, A., Bundschuh, M., 2020. Decoupled structure and function of leaf-associated
625   microorganisms under anthropogenic pressure: Potential hurdles for environmental
626   monitoring. Freshw. Sci. 39, 652–664. https://doi.org/10.1086/709726

627   Feckler, A., Goedkoop, W., Konschak, M., Bundschuh, R., Kenngott, K.G.J., Schulz, R.,
628   Zubrod, J.P., Bundschuh, M., 2018. History matters: heterotrophic microbial
629   community structure and function adapt to multiple stressors. Glob. Change Biol. 24,
630   e402–e415. https://doi.org/10.1111/gcb.13859

631   Feckler, A., Kahlert, M., Bundschuh, M., 2015. Impacts of contaminants on the ecological
632   role of lotic biofilms. Bull. Environ. Contam. Toxicol. 95, 421–427.
633   https://doi.org/10.1007/s00128-015-1642-1

634   Fenner, K., Canonica, S., Wackett, L.P., Elsner, M., 2013. Evaluating pesticide degradation
635   in the environment: blind spots and emerging opportunities. Science 341, 752–758.
636   https://doi.org/10.1126/science.1236281

637   Ferreira, V., Koricheva, J., Duarte, S., Niyogi, D.K., Guérold, F., 2016. Effects of
638   anthropogenic heavy metal contamination on litter decomposition in streams – A
639   meta-analysis. Environ. Pollut. 210, 261–270.
640   https://doi.org/10.1016/j.envpol.2015.12.060

641   González-Pleiter, M., Gonzalo, S., Rodea-Palomares, I., Leganés, F., Rosal, R., Boltes, K.,
642   Marco, E., Fernández-Piñas, F., 2013. Toxicity of five antibiotics and their mixtures
643   towards photosynthetic aquatic organisms: Implications for environmental risk
644   assessment. Water Res. 47, 2050–2064. https://doi.org/10.1016/j.watres.2013.01.020

645   Guo, J., Boxall, A., Selby, K., 2015. Do pharmaceuticals pose a threat to primary producers?
646   Crit. Rev. Environ. Sci. Technol. 45, 2565–2610.
647   https://doi.org/10.1080/10643389.2015.1061873

648   Halstead, N.T., McMahon, T.A., Johnson, S.A., Raffel, T.R., Romansic, J.M., Crumrine,
649   P.W., Rohr, J.R., 2014. Community ecology theory predicts the effects of
650   agrochemical mixtures on aquatic biodiversity and ecosystem properties. Ecology
651   Letters 17, 932–941. https://doi.org/10.1111/ele.12295

652   Harrer, M., Cuijpers, P., Furukawa, T., Ebert, D.D., 2019. dmetar: Companion R Package For
653   The Guide "Doing Meta-Analysis in R."

Harrison, I., Abell, R., Darwall, W., Thieme, M.L., Tickner, D., Timboe, I., 2018. The

freshwater biodiversity crisis. Science 362, 1369–1369.

https://doi.org/10.1126/science.aav9242

Harrison, P.A., Berry, P.M., Simpson, G., Haslett, J.R., Blicharska, M., Bucur, M., Dunford,

R., Egoh, B., Garcia-Llorente, M., Geamănă, N., Geertsema, W., Lommelen, E.,

Meiresonne, L., Turkelboom, F., 2014. Linkages between biodiversity attributes and

ecosystem services: a systematic review. Ecosyst. Serv. 9, 191–203.

https://doi.org/10.1016/j.ecoser.2014.05.006

Hedlund, J., Longo, S.B., York, R., 2020. Agriculture, pesticide use, and economic

development: a global examination (1990–2014). Rural Sociol. 85, 519–544.

https://doi.org/10.1111/ruso.12303

Hiki, K., Iwasaki, Y., 2020. Can We Reasonably Predict Chronic Species Sensitivity

Distributions from Acute Species Sensitivity Distributions? Environ. Sci. Technol. 54,

13131–13136. https://doi.org/10.1021/acs.est.0c03108

Higgins, J.P.T., Thompson, S.G., 2004. Controlling the risk of spurious findings from meta-

regression. Stat. Med. 23, 1663–1682. https://doi.org/10.1002/sim.1752

Jenkins, D.G., Quintana-Ascencio, P.F., 2020. A solution to minimum sample size for

regressions. PLOS ONE 15, e0229345. https://doi.org/10.1371/journal.pone.0229345

Jepsen, R., He, K., Blaney, L., Swan, C., 2019. Effects of antimicrobial exposure on detrital

biofilm metabolism in urban and rural stream environments. Sci. Total Environ. 666,

1151–1160. https://doi.org/10.1016/j.scitotenv.2019.02.254

Klocker, C.A., Kaushal, S.S., Groffman, P.M., Mayer, P.M., Morgan, R.P., 2009. Nitrogen

uptake and denitrification in restored and unrestored streams in urban Maryland,

USA. Aquat. Sci. 71, 411–424. https://doi.org/10.1007/s00027-009-0118-y

Kovalakova, P., Cizmas, L., McDonald, T.J., Marsalek, B., Feng, M., Sharma, V.K., 2020.

Occurrence and toxicity of antibiotics in the aquatic environment: a review.

Chemosphere 251, 126351. https://doi.org/10.1016/j.chemosphere.2020.126351

Kraft, N.J.B., Adler, P.B., Godoy, O., James, E.C., Fuller, S., Levine, J.M., 2015. Community

assembly, coexistence and the environmental filtering metaphor. Functional Ecology

29, 592–599. https://doi.org/10.1111/1365-2435.12345

Kümmerer, K., 2009. Antibiotics in the aquatic environment – a review – Part I.

Chemosphere 75, 417–434. https://doi.org/10.1016/j.chemosphere.2008.11.086

Lewis, K., Tzilivakis, J., Green, A., Warner, D.J., 2011. Veterinary Substances Database

(VSDB).

Lewis, K.A., Tzilivakis, J., Warner, D.J., Green, A., 2016. An international database for
689   pesticide risk assessments and management. Hum. Ecol. Risk Assess. 22, 1050–1064.
690   https://doi.org/10.1080/10807039.2015.1133242

691   Liess, M., Beketov, M., 2011. Traits and stress: keys to identify community effects of low
692   levels of toxicants in test systems. Ecotoxicology 20, 1328–1340.
693   https://doi.org/10.1007/s10646-011-0689-y

694   Liess, M., Champeau, O., Riddle, M., Schulz, R., Duquesne, S., 2001. Combined effects of
695   ultraviolet-B radiation and food shortage on the sensitivity of the Antarctic amphipod
696   *Paramoera walkeri* to copper. Environ. Toxicol. Chem. 20, 2088–2092.
697   https://doi.org/10.1002/etc.5620200931

698   Liess, M., Foit, K., Knillmann, S., Schäfer, R.B., Liess, H.-D., 2016. Predicting the synergy
699   of multiple stress effects. Sci. Rep. 6, 32965. https://doi.org/10.1038/srep32965

700   Liess, M., Liebmann, L., Vormeier, P., Weisner, O., Altenburger, R., Borchardt, D., Brack,
701   W., Chatzinotas, A., Escher, B., Foit, K., Gunold, R., Henz, S., Hitzfeld, K.L.,
702   Schmitt-Jansen, M., Kamjunke, N., Kaske, O., Knillmann, S., Krauss, M., Küster, E.,
703   Link, M., Lück, M., Möder, M., Müller, A., Paschke, A., Schäfer, R.B., Schneeweiss,
704   A., Schreiner, V.C., Schulze, T., Schüürmann, G., von Tümpling, W., Weitere, M.,
705   Wogram, J., Reemtsma, T., 2021. Pesticides are the dominant stressors for vulnerable
706   insects in lowland streams. Water Res. 201, 117262.
707   https://doi.org/10.1016/j.watres.2021.117262

708   Louca, S., Polz, M.F., Mazel, F., Albright, M.B.N., Huber, J.A., O'Connor, M.I., Ackerman,
709   M., Hahn, A.S., Srivastava, D.S., Crowe, S.A., Doebeli, M., Parfrey, L.W., 2018.
710   Function and functional redundancy in microbial systems. Nature Ecol. Evol. 2, 936-
711   943. https://doi.org/10.1038/s41559-018-0519-1

712   Malaj, E., Ohe, P.C. von der, Grote, M., Kühne, R., Mondy, C.P., Usseglio-Polatera, P.,
713   Brack, W., Schäfer, R.B., 2014. Organic chemicals jeopardize the health of freshwater
714   ecosystems on the continental scale. PNAS 111, 9549–9554.
715   https://doi.org/10.1073/pnas.1321082111

716   Martin, O., Scholze, M., Ermler, S., McPhie, J., Bopp, S.K., Kienzler, A., Parissis, N.,
717   Kortenkamp, A., 2021. Ten years of research on synergisms and antagonisms in
718   chemical mixtures: a systematic review and quantitative reappraisal of mixture
719   studies. Environ. Int. 146, 106206. https://doi.org/10.1016/j.envint.2020.106206

720   Molander, L., Ågerstrand, M., Rudén, C., 2009. WikiPharma – A freely available, easily
721   accessible, interactive and comprehensive database for environmental effect data for

pharmaceuticals. Regul. Toxicol. Pharm. 55, 367–371.
https://doi.org/10.1016/j.yrtph.2009.08.009

Naeem, S., Wright, J.P., 2003. Disentangling biodiversity effects on ecosystem functioning:
deriving solutions to a seemingly insurmountable problem. Ecol. Lett. 6, 567–579.
https://doi.org/10.1046/j.1461-0248.2003.00471.x

Neale, P.A., Braun, G., Brack, W., Carmona, E., Gunold, R., König, M., Krauss, M.,
Liebmann, L., Liess, M., Link, M., Schäfer, R.B., Schlichting, R., Schreiner, V.C.,
Schulze, T., Vormeier, P., Weisner, O., Escher, B.I., 2020. Assessing the mixture
effects in in vitro bioassays of chemicals occurring in small agricultural streams
during rain events. Environ. Sci. Technol. 54, 8280–8290.
https://doi.org/10.1021/acs.est.0c02235

Newman, M.C., 2009. Fundamentals of Ecotoxicology. CRC Press.

Nie, X., Gu, J., Lu, J., Pan, W., Yang, Y., 2009. Effects of norfloxacin and butylated
hydroxyanisole on the freshwater microalga Scenedesmus obliquus. Ecotoxicology
18, 677–684. https://doi.org/10.1007/s10646-009-0334-1

Peters, K., Bundschuh, M., Schäfer, R.B., 2013. Review on the effects of toxicants on
freshwater ecosystem functions. Environ. Pollut. 180, 324–329.
https://doi.org/10.1016/j.envpol.2013.05.025

Poisot, T., 2011. The digitize package: extracting numerical data from scatterplots. The R
Journal 3, 25–26.

R Core Team, 2024. R: a language and environment for statistical computing.

Ragnarsdottir, K.V., 2000. Environmental fate and toxicology of organophosphate pesticides.
J. Geol. Soc. 157, 859–876. https://doi.org/10.1144/jgs.157.4.859

Reid, A.J., Carlson, A.K., Creed, I.F., Eliason, E.J., Gell, P.A., Johnson, P.T.J., Kidd, K.A.,
MacCormack, T.J., Olden, J.D., Ormerod, S.J., Smol, J.P., Taylor, W.W., Tockner,
K., Vermaire, J.C., Dudgeon, D., Cooke, S.J., 2019. Emerging threats and persistent
conservation challenges for freshwater biodiversity. Biol. Rev. 94, 849–873.
https://doi.org/10.1111/brv.12480

Richmond, E.K., Rosi, E.J., Reisinger, A.J., Hanrahan, B.R., Thompson, R.M., Grace, M.R.,
2019. Influences of the antidepressant fluoxetine on stream ecosystem function and
aquatic insect emergence at environmentally realistic concentrations. J. Freshw. Ecol.
34, 513–531. https://doi.org/10.1080/02705060.2019.1629546

Robson, S.V., Rosi, E.J., Richmond, E.K., Grace, M.R., 2020. Environmental concentrations of pharmaceuticals alter metabolism, denitrification, and diatom assemblages in artificial streams. Freshw. Sci. 39, 256-267. https://doi.org/10.1086/708893

Rosi-Marshall, E.J., Royer, T.V., 2012. Pharmaceutical compounds and ecosystem function: an emerging research challenge for aquatic ecologists. Ecosystems 15, 867–880. https://doi.org/10.1007/s10021-012-9553-z

Rumschlag, S.L., Halstead, N.T., Hoverman, J.T., Raffel, T.R., Carrick, H.J., Hudson, P.J., Rohr, J.R., 2019. Effects of pesticides on exposure and susceptibility to parasites can be generalised to pesticide class and type in aquatic communities. Ecology Letters 22, 962–972. https://doi.org/10.1111/ele.13253

Rumschlag, S.L., Mahon, M.B., Hoverman, J.T., Raffel, T.R., Carrick, H.J., Hudson, P.J., Rohr, J.R., 2020. Consistent effects of pesticides on community structure and ecosystem function in freshwater systems. Nature Communications 11, 6333. https://doi.org/10.1038/s41467-020-20192-2

Schäfer, R.B., Bundschuh, M., Rouch, D.A., Szöcs, E., von der Ohe, P.C., Pettigrove, V., Schulz, R., Nugegoda, D., Kefford, B.J., 2012a. Effects of pesticide toxicity, salinity and other environmental variables on selected ecosystem functions in streams and the relevance for ecosystem services. Sci. Total Environ. 415, 69–78. https://doi.org/10.1016/j.scitotenv.2011.05.063

Schäfer, R.B., von der Ohe, P.C., Rasmussen, J., Kefford, B.J., Beketov, M.A., Schulz, R., Liess, M., 2012b. Thresholds for the effects of pesticides on invertebrate communities and leaf breakdown in stream ecosystems. Environ. Sci. Technol. 46, 5134–5142. https://doi.org/10.1021/es2039882

Schäfer, R.B., Gerner, N., Kefford, B.J., Rasmussen, J.J., Beketov, M.A., de Zwart, D., Liess, M., von der Ohe, P.C., 2013. How to Characterize Chemical Exposure to Predict Ecologic Effects on Aquatic Communities? Environ. Sci. Technol. 47, 7996–8004. https://doi.org/10.1021/es4014954

Schäfer, R.B., Baikova, D., Bayat, H., Beermann, A., Berger, S.A., Boenigk, J., Bruans, M., Burfeid-Castallanos, A., Cardinale, B., David, G., Feckler, A., Feld, C.K., Fink, P., Gessner, M.O., Hadziomerovic, U., Hering, D., le, T.T.Y., Macaulay, S., Madariaga, G.M., Mayombo, S., Pimentel, I.M., Orr, J., Oskapolor, S., Schlenker, A., Sures, B., Vermiert, A.-M., Weitere, M., Vos, M., Schürings, C., in preparation. Multiple stressors increase the effects of taxon loss on ecosystem functioning.

Scharmüller, A., Schreiner, V.C., Schäfer, R.B., 2020. Standartox: standardizing toxicity
data. Data 5, 46. https://doi.org/10.3390/data5020046

Schulz, R., Bub, S., Petschick, L.L., Stehle, S., Wolfram, J., 2021. Applied pesticide toxicity
shifts toward plants and invertebrates, even in GM crops. Science 372, 81–84.
https://doi.org/10.1126/science.abe1148

Shiklomanov, I.A., 1993. World freshwater resource. Water in crisis: a guide to the world's
freshwater resources. Oxford University Press, New York.

Sinclair, T., Craig, P., Maltby, L.L., 2024. Climate warming shifts riverine macroinvertebrate
communities to be more sensitive to chemical pollutants. Global Change Biology 30,
e17254. https://doi.org/10.1111/gcb.17254

Soares, M.P., Jesus, F., Almeida, A.R., Domingues, I., Hayd, L., Soares, A.M.V.M., 2020.
Effects of pH and nitrites on the toxicity of a cypermetrin-based pesticide to shrimps.
Chemosphere 241, 125089. https://doi.org/10.1016/j.chemosphere.2019.125089

Stehle, S., Schulz, R., 2015. Agricultural insecticides threaten surface waters at the global
scale. PNAS 112, 5750–5755. https://doi.org/10.1073/pnas.1500232112

Strayer, D.L., Dudgeon, D., 2010. Freshwater biodiversity conservation: recent progress and
future challenges. J. N. Am. Benthol. Soc. 29, 344–358. https://doi.org/10.1899/08-
171.1

Tilman, D., Isbell, F., Cowles, J.M., 2014. Biodiversity and ecosystem functioning. Annu.
Rev. Ecol. Evol. Syst. 45, 471–493. https://doi.org/10.1146/annurev-ecolsys-120213-
091917

Udy, J.W., Fellows, C.S., Bartkow, M.E., Bunn, S.E., Clapcott, J.E., Harch, B.D., 2006.
Measures of nutrient processes as indicators of stream ecosystem health.
Hydrobiologia 572, 89–102. https://doi.org/10.1007/s10750-005-9006-1

van Aert, R.C.M., Jackson, D., 2019. A new justification of the Hartung-Knapp method for
random-effects meta-analysis based on weighted least squares regression. Res. Synth.
Meth. 10, 515–527. https://doi.org/10.1002/jrsm.1356

Verheyen, J., Delnat, V., Theys, C., 2022. Daily temperature fluctuation can magnify the
toxicity of pestides. Curr. Opin. Insect Sci. 51, 100919.
https://doi.org/10.1016/j.cois.2022.100919

Viechtbauer, W., 2010. Conducting meta-analyses in R with the metafor package. J. Stat.
Softw. 36, 1–48. https://doi.org/10.18637/jss.v036.i03

von Schiller, D., Acuña, V., Aristi, I., Arroita, M., Basaguren, A., Bellin, A., Boyero, L.,
Butturini, A., Ginebreda, A., Kalogianni, E., Larrañaga, A., Majone, B., Martínez, A.,

Monroy, S., Muñoz, I., Paunović, M., Pereda, O., Petrovic, M., Pozo, J., Rodríguez-
Mozaz, S., Rivas, D., Sabater, S., Sabater, F., Skoulikidis, N., Solagaistua, L.,
Vardakas, L., Elosegi, A., 2017. River ecosystem processes: a synthesis of
approaches, criteria of use and sensitivity to environmental stressors. Sci. Total
Environ. 596–597, 465–480. https://doi.org/10.1016/j.scitotenv.2017.04.081

Wang, Z., Meador, J.P., Leung, K.M.Y., 2016. Metal toxicity to freshwater organisms as a
function of pH: a meta-analysis. Chemosphere 144, 1544–1552.
https://doi.org/10.1016/j.chemosphere.2015.10.032

Wang, Z., Walker, G.W., Muir, D.C.G., Nagatani-Yoshida, K., 2020. Toward a global
understanding of chemical pollution: a first comprehensive analysis of national and
regional chemical inventories. Environ. Sci. Technol. 54, 2575–2584.
https://doi.org/10.1021/acs.est.9b06379

Warne, M.S.J., Hawker, D.W., 1995. The number of components in a mixture determines
whether synergistic and antagonistic or additive toxicity predominate - the funnel
hypothesis. Ecotoxicol. Environ. Safe. 31, 23–28.
https://doi.org/10.1006/eesa.1995.1039

Wolfram, J., Stehle, S., Bub, S., Petschick, L.L., Schulz, R., 2021. Water quality and
ecological risks in European surface waters – Monitoring improves while water
quality decreases. Environ. Int. 152, 106479.
https://doi.org/10.1016/j.envint.2021.106479

Young, R.G., Matthaei, C.D., Townsend, C.R., 2008. Organic matter breakdown and
ecosystem metabolism: functional indicators for assessing river ecosystem health. J.
N. Am. Benthol. Soc. 27, 605–625. https://doi.org/10.1899/07-121.1

Zubrod, J.P., Bundschuh, M., Arts, G., Brühl, C.A., Imfeld, G., Knäbel, A., Payraudeau, S.,
Rasmussen, J.J., Rohr, J., Scharmüller, A., Smalling, K., Stehle, S., Schulz, R.,
Schäfer, R.B., 2019. Fungicides: an overlooked pesticide class? Environ. Sci.
Technol. 53, 3347–3365. https://doi.org/10.1021/acs.est.8b04392

Figure captions

Figure 1: Alluvial plot displaying the extracted effect data. The size of each box is

proportional to the number of extracted observations per organism group (left) or ecosystem

function (right). The link width is proportionate to the number of observations per chemical

type (indicated by color).

Figure 2: Relationships between sumTU and (a) organic matter decomposition by

decomposer-detritivore-systems, (b) organic matter decomposition by microbial

decomposers, (c) photosynthesis by algae, and (d) photosynthesis by macrophytes, shown as

the proportion of the functioning in the stressor-treatment compared to the stressor-free

control. Colors code chemical types (see Fig. 1), and shapes represent the study system,

namely laboratory (circles) and micro-/mesocosms (squares). The shaded area represents the

95% confidence interval based on the bootstrapped data, while the black line represents the

median model. Please note the differences in the x-axis scale.

Figure 3: Relationships between $\log_{10}(RQ)$ and (a) organic matter decomposition by

decomposer-detritivore-systems, (b) organic matter decomposition by microbial

decomposers, (c) photosynthesis by algae, and (d) photosynthesis by macrophytes, shown as

the proportion of the functioning in the stressor-treatment compared to the stressor-free

control. Colors code chemical types (see Fig. 1), and shapes represent the study system,

namely laboratory (circles) and micro-/mesocosms (squares). The shaded area represents the

95% confidence interval based on the bootstrapped data, while the black line represents the

median model. Please note the differences in the x-axis scale.