Meta-analysis on the effects of chemical stressors on five ecosystem functions: are there

generalizable relationships?

Alexander Feckler[1,2,*], Ralf Schulz[1,2], Ralf B. Schäfer[1]

[1]iES, Institute for Environmental Sciences, RPTU Kaiserslautern-Landau, Fortstraße 7, D-

76829 Landau, Germany

[2]Eusserthal Ecosystem Research Station, RPTU Kaiserslautern-Landau, Birkenthalstraße 13,

D-76857 Eusserthal, Germany

* Contact details for the corresponding author:

Name: Alexander Feckler

Address: iES, Institute for Environmental Sciences, RPTU Kaiserslautern-Landau, Fortstraße

7, 76829 Landau in der Pfalz, Germany

Telephone: +49 6341 28031312

Email: alexander.feckler@rptu.de

Abstract

Despite considerable progress in our predictive capacity for the response of different trophic

levels to chemicals, generalizable relations between chemicals and ecosystem functions are

lacking. We addressed this knowledge gap by conducting the largest meta-analysis to date (136

studies; ~350 observations) on the responses of ecosystem functions to chemicals. Based on

this analysis, for selected classes of chemicals (pesticides, pharmaceuticals, and metals), we

are the first to establish relations between chemical concentrations and the ecosystem functions

organic matter decomposition and photosynthesis, showing monotonic dose-response

relationships. These results concur with existing field data on chemical effects, which

underpins the value of our findings. Taking into account the outlined research gaps and

shortcomings, our findings and recommendations can be seen as a promising starting point to

advance predictions of responses of ecosystem functions to selected chemical classes.

Keywords

Freshwater ecosystems, stress-ecosystem function relationship, context-dependency, effect

modeling, chemical risk

Introduction

Freshwater habitats, despite comprising just around 0.01% of the global water resources and covering only 2% of the global surface area (Shiklomanov, 1993), hold a remarkable species biodiversity (Strayer and Dudgeon, 2010). Hosting over 140,000 species (Reid et al., 2019), with approximately 80% being invertebrate species and 15% vertebrate species (Balian et al., 2008), these ecosystems are pivotal for species biodiversity. However, a stark contrast emerges as freshwater habitats experience more pronounced declines in biodiversity compared to terrestrial ecosystems, leading to what has been labelled as the "freshwater crisis" (Harrison et al., 2018). The freshwater crisis is the result of an intricate interplay of anthropogenic disturbances: over-exploitation, toxic chemical pollution, alterations in water flow patterns, habitat destruction or degradation, and the introduction of non-native species (as comprehensively reviewed in Reid et al., 2019). Despite their paramount importance, toxic chemicals have often been marginalized in freshwater ecosystem studies, even though they play a substantial role in deteriorating these ecosystems (Bernhardt et al., 2017). Extensive meta-analyses on continental and global scales consistently reveal that a substantial proportion of water bodies are affected by toxic chemicals. In Europe, nearly half of the evaluated water bodies were identified as being at risk due to chronic chemical effects (Malaj et al., 2014). Similarly, on a global scale, insecticide contamination was observed to surpass thresholds indicating ecological impacts in nearly 70% of stream sites within agricultural landscapes (Stehle and Schulz, 2015). Recent investigations underscore the escalating chemical risk faced by freshwater organisms, both in Europe and the United States (Schulz et al., 2021; Wolfram et al., 2021), pointing toward a concerning thread of increasing contamination.

The decline in freshwater biodiversity raises considerable concerns regarding its repercussions on the functioning of aquatic ecosystems (Cardinale et al., 2012; Hooper et al., 2005; Loreau et al., 2001). Over the past few decades, substantial research efforts have been

dedicated to unraveling the connections between biodiversity and ecosystem functioning

(Naeem and Wright, 2003). Numerous empirical studies have unveiled a connection between

these variables, demonstrating that the loss of species generally diminishes ecosystem

functioning (Harrison et al., 2014; Tilman et al., 2014). Nevertheless, the interplay between

biodiversity and ecosystem functioning under the influence of chemical stress remains less

elucidated. A handful of meta-analyses and reviews (Crane et al., 2006; Kümmerer, 2009;

Schäfer et al., 2012b) showed that specific organisms, which are crucial for functions like

nutrient cycling, organic matter decomposition, and primary production, are impacted by

chemical stress at environmentally relevant concentrations. Nevertheless, the broader

implications on the functional level remain a subject of ongoing investigation. Additionally,

several meta-analyses and reviews have drawn connections between changes in ecosystem

functions and the effects of chemical stressors on specific groups of organisms (Beaumelle et

al. 2020; DeLorenzo et al., 2001; Feckler et al., 2015; Feckler and Bundschuh, 2020; Ferreira

et al., 2016; Guo et al., 2015; Kovalakova et al., 2020; Peters et al., 2013; Schäfer et al., 2012b).

Despite compelling evidence highlighting the impact of chemical stressors on ecosystem

functions, we still lack a generalizable relationship between chemical stress and ecosystem

functioning. Notably, a meta-analysis failed to establish a consistent dose-response relationship

between various groups of chemical stressors and several ecosystem functions, such as nutrient

cycling, organic matter decomposition and primary production (Peters et al., 2013). However,

Peters et al. (2013) rather aimed at identifying effect thresholds and only considered studies

that exceeded a defined negative effect size, whereas the responses of ecosystem functions to

chemical stress can be negative (e.g., Beaumelle et al., 2020; Robson et al., 2020; Rosi-

Marshall et al., 2013), positive (e.g., Corcoll et al., 2015; Richmond et al., 2019) or non-

distinguishable from a control treatment or a reference (e.g., Jepsen et al., 2019; Schäfer et al.,

2012a). Thus, it remains open whether an analysis of all studies, including novel studies that

have been conducted in the last decade, enables the derivation of generalizable relationships

between chemical stressors and their effects on ecosystem functions.

We aimed to derive such generalizable relationships to help predict chemical effects on

ecosystems. Therefore, we conducted a literature meta-analysis and extracted 347 observations

(effect sizes and respective chemical concentration) from 136 published laboratory studies that

cover a wide range of chemical groups (metals, pesticides, and pharmaceuticals) and five

ecosystem functions (community respiration, organic matter decomposition, nutrient cycling,

photosynthesis, and primary production). We hypothesized to find a negative association

between chemical stressors standardized by their toxic potency and levels of ecosystem

functions. Furthermore, we hypothesized that the effects of particular chemical groups would

be most pronounced for ecosystem functions performed by organism groups targeted by these

chemical stressors. For instance, we anticipated the most substantial effects of fungicides and

herbicides on microbially-mediated organic matter decomposition and photosynthesis,

respectively. Our results indicate generalizable relationships for the ecosystem functions

organic matter decomposition and photosynthesis, showing a decrease in the function as the

chemical pressure increased. The match of the results found here with data from field studies

suggests that the relationships are relevant for ecosystems. Finally, we distilled research gaps

that require further scrutiny in the future.

Materials and methods

Literature collection and inclusion criteria

We conducted an extensive search across Web of Science and Scopus to compile literature concerning the effects of chemical stressors (pesticides, pharmaceuticals, and metals) on five ecosystem functions (community respiration, nutrient cycling, organic matter decomposition, photosynthesis, and primary production). The search was limited to literature published between January 2000 and December 2020 to focus on currently used chemical stressors. Our search string was constructed using a combination of freshwater habitat-related terms (freshwater* OR stream* OR river* OR pond* OR lake*), stressor-related terms (chemical stress* OR contaminant* OR pollutant* OR toxicant* OR pesticide* OR heavy metal* OR metal* OR fungicide* OR herbicide* OR insecticide* OR pharmaceut*), and ecosystem function-related terms (ecosystem function* OR photosynth* OR primary product* OR nutrient cycl* OR organic matter decomp* OR leaf breakdown OR decomposition* OR respirat*). The asterisks (*) were employed as wildcards, capturing all terms sharing the same root. Moreover, we conducted footnote chasing by screening the reference lists of relevant articles for further literature. The selection process for studies encompassed several stages. Initially, titles were screened to identify suitable studies. Subsequently, the abstracts were evaluated, leading to the exclusion of non-matching studies. Lastly, the full texts of selected studies were assessed.

Studies were included based on a set of criteria: (1) laboratory-based investigations, as other study types (semi-field or field) yielded insufficient data points for meta-analysis (Figure A1 in the Supporting Information), (2) focus on species, populations, or communities dwelling within freshwater habitats (excluding groundwater, marshlands, coastal waters, and marine systems), (3) exposure to at least one of the pre-defined chemical stressors from the list above (excluding eutrophication, salinization, and acidification), (4) presence of a control treatment

(absence of chemical stress), and (5) measurement of the designated ecosystem functions as

well as uncertainty around the central tendency of the measurement. An overview of the

selected studies and moderator variables (see below) will be provided in a data repository upon

acceptance of the manuscript.

Data extraction and effect size calculation

We extracted central tendencies, uncertainty estimators, sample sizes, and chemical

concentrations from the text, tables, and figures of the selected studies. For cases where the

data were presented in figures, we utilized the "digitize" package (Poisot, 2011) within the

open-source software R (version 4.2.3 for Mac OS X; R Core Team, 2023).

As our response variable, we computed the ratio of the ecosystem function observed in the

stressor treatment to that in the control treatment. From each study, we extracted only the

highest effect observed per functional endpoint for the same species, population, or community.

This approach was adopted to derive unequivocal cause-effect relationships and prevent

pseudoreplication. Due to this method and the diverse test designs employed in individual

studies, the timing of functional measurements varied.

For studies investigating community respiration, the functional indicators included

measurements of O_2 consumption (e.g., mg O_2 g^{-1} dry weight h^{-1}) and CO_2 production (e.g., µg

CO_2 g^{-1} dry weight h^{-1}). In the context of organic matter decomposition, the indicators consisted

of breakdown rates (k (d^{-1})) and mass loss (e.g., mg d^{-1}) of leaf material, mediated by microbial

decomposers or a combination of microbial decomposers and detritivorous macroinvertebrates

(decomposer-detritivore-systems) as endpoints. Concerning nutrient cycling (primarily

nitrogen), the indicators encompassed anaerobic NH_4^+ oxidation rates (g-N g-volatile

suspended solids^{-1} d^{-1}), NO_2 and NO_3 turnover rates (e.g., mg NO_2-N mg^{-1} d^{-1}), and N_2

production (e.g., N_2 g^{-1} h^{-1}). In the case of photosynthesis, we determined effect sizes based on

maximum and effective photosystem II quantum yields (F_v/F_m and $\Phi PSII$, respectively) and

O_2 production (e.g., mg O_2 g^{-1} dry weight h^{-1}). Finally, for primary production, the effect sizes

related to fixed carbon quantities (mg ^{14}C g^{-1} C h^{-1}) and changes in the standing biomass (net

primary production). Studies that exclusively reported dissolved oxygen as a response variable

were excluded, as net dissolved oxygen could arise from multiple sources simultaneously, such

as photosynthesis and the ambient atmosphere.

Calculation of toxic units

The toxic effects caused by different chemical stressors vary for a given test

concentration, due to differing ecotoxicological potentials, making direct comparisons

challenging. Therefore, we normalized the test concentrations of individual chemical stressors

to their acute effects by calculating toxic units (TUs). Ideally, TUs would be computed for each

combination of stressor and organism included in the selected literature. However, given

substantial data gaps, we relied on surrogate species results to facilitate comparisons between

different stressors. Unlike previous meta-analyses that sought to establish a generalized

chemical stress–ecosystem function relationship (e.g., Peters et al., 2013), we avoided

confining the surrogate species selection to conventional test species as stipulated by chemical

regulations (i.e., *Daphnia magna*, *Lemna* sp., and *Raphidocelis subcapitata*). Instead, we

estimated stressor toxicity based on the EC_{50} values – the concentration at which an effect is

observed in 50% of exposed organisms – for the most sensitive freshwater algal, plant, and

invertebrate species for the respective chemical stressor. We employed these EC_{50} values to

compute the logarithmic sum of toxic units (sumTU) as:

$$sumTU = log\left(\sum_{i=1}^{n} \frac{c_i}{EC50_i}\right)$$

where c represents the concentration of each toxicant i, $EC50_i$ is the median effect

concentration of the respective toxicant for the most sensitive species i, and n gives the number

of toxicants that caused an effect in the respective ecosystem function. EC_{50} values were

extracted from various sources including Standartox (Scharmüller et al., 2020), the Pesticide

Property Database (Lewis et al., 2016), the Veterinary Substance Database (Lewis et al., 2011),

the Norman Database (https://www.norman-network.com/nds/), and WikiPharma (Molander

et al., 2009). We chose EC_{50} values corresponding to a 48-hour exposure duration or, when not

available, the closest exposure time. It is important to note that this chosen exposure time serves

merely as a reference for benchmarking the relative toxicity of compounds and does not need

to align with the experimental exposure time. See Supplementary Data (Table A1) for an

overview of all EC_{50} values and their sources.

Moderator variables

Besides the sumTU, we extracted seven moderator variables: (1) ecosystem function

with levels: (a) community respiration, (b) organic matter decomposition, (c) nutrient cycling,

(d) photosynthesis, and (e) primary production; (2) detailed chemical group with levels: (a)

fungicide, (b) herbicide, (c) insecticide, (d) pesticide mixture, (e) pharmaceutical, (f) metal, (g)

metal mixture, and (h) other (such as biocides); (3) classification of the mode of toxic action

extracted from Lewis et al. (2016) or Molander et al. (2009); (4) organism group with levels:

(a) algae (including cyanobacteria), (b) aquatic plants (phytoplankton, macrophytes, etc.), (c)

microbial   decomposers   (bacteria   and   fungi),   (d)   decomposer-detritivore-systems

(microorganisms and macroinvertebrates); (5) exposure time as a continuous variable in days;

(6) pH as a continuous variable; (7) test temperature as a continuous variable in °C.

Data analysis

To identify the moderator variables affecting the assessed ecosystem functions and

establish relationships between chemical stressors and ecosystem functions, we performed

separate data analyses for each of the included ecosystem functions. We used the calculated

effect size (along with the standard error of the effect size for each study) as the response

variable, while including the sumTU and moderator variables as predictors. Initially, we

checked for multi-collinearity, ensuring that the meta-regression coefficient estimates are

robust by examining correlations ($r \geq 0.8$) among our predictors. Subsequently, we identified

the best-fitting models using the "multimodel.inference" function in the R package "dmetar"

(Harrer et al., 2019) based on corrected Akaike Information Criteria. Using the results of the

multimodel inference, we build models to evaluate the individual impact of the identified

predictors and their interactions. We achieved this by fitting meta-regression models using the

R package "metafor" (Viechtbauer, 2010). We employed the "maximum likelihood" estimator

to assess the amount of residual heterogeneity (τ^2) and applied the Knapp-Hartung method for

random-effects meta-analysis (van Aert and Jackson, 2019). We conducted a permutation test

before reporting *p*-values, as recommended for meta-regression models (Higgins and

Thompson, 2004).

In a subsequent step, we fitted (quasi-)binomial Generalized Linear Models (GLMs) to

explore potential dose-response relationships between ecosystem functions and sumTUs of all

stressors or separately for chemical groups (fungicides, herbicides, insecticides,

pharmaceuticals, and metals) or modes of toxic action (only if $n \geq 8$; Jenkins and Quintana-

Ascencio, 2020). For statistical analyses and graphics, we used the open-source software R (R

Core Team, 2023), along with the necessary add-on packages. Upon manuscript acceptance,

we will make all code and data available in a data repository.

Results

Overview of included studies

Through our database query and footnote chasing, we identified a total of 7,599

publications, of which 7,463 publications were excluded due to various reasons including

violations of our predefined inclusion criteria, absence of reported effect sizes and associated

uncertainties, or the inability to calculate sumTUs due to a lack of toxicity data. Ultimately, we

focused on 136 laboratory studies that met our criteria, from which we extracted 347 effect

sizes, uncertainty estimators, sumTUs, and moderator variables (see Fig. A1 for details). We

deliberately excluded semi-field and field studies from further analyses. This decision was

driven by the fact that these types of studies only represented a minor share of our dataset after

initial screening ($n = 38$ and 60 for semi-field and field studies, respectively), while a major

fraction of these 98 studies did not meet our predefined inclusion criteria and would have been

excluded (see overview of studies).

Among the selected studies, we found that most observations (37% each) were related

to organic matter decomposition and photosynthesis. Additionally, there were 75 observations

(22%) related to primary production, while community respiration and nutrient cycling studies

constituted 9 observations (4%; Fig. 1a). Regarding the types of chemical stressors examined,

pesticides were the most extensively studied group, representing 63% of the observations (218

observations in total), with fungicides accounting for 56 observations, herbicides for 102,

insecticides for 34, and pesticide mixtures for 26. Metals and metal mixtures constituted 30%

of the observations (104 observations in total), while pharmaceuticals made up 7% (26

observations; Figs. 1a & 1b). The sumTUs ranged from -5.13 to 2.68 for metals and metal

mixtures, -4.54 to 3.99 for pesticides and pesticide mixtures, and -5.68 to 5.58 for

pharmaceuticals (Table 1). A value of 0 indicated that the toxicity corresponded to the EC_{50} of

the most sensitive species for which toxicity data were available. When comparing the tested

concentrations to the range of detected environmental concentrations for each stressor group,

which we obtained from field measurements through meta-analyses (Deo, 2014; Wolfram et

al., 2021), we observed that these two datasets only partly overlap (Fig. A2). However, this

(substantial) non-overlap should be interpreted cautiously due to our specific selection

procedure for effect sizes (utilizing the highest effect from each study) and potential differences

between the tested and detected substances.

Influence of moderator variables and relationships between sumTU and effects on ecosystem

functions

Organic matter decomposition

For the ecosystem function organic matter decomposition, our best-fit model included

two predictors: "organism group", which categorized microbial decomposers alone and

decomposer-detritivore-systems, and "pH". Additionally, there was a significant interaction

between "organism group" and "sumTU". This model accounted for a substantial portion of

the variance ($R^2 = 58.93\%$, $\tau^2 = 0.0434$; Table 2). We observed a general dose-response

relationship for organic matter decomposition across all groups of chemicals for studies

examining decomposer-detritivore systems (as indicated by the GLM for sumTU and the

respective ecosystem function; $p = 0.022$; $n = 62$, Fig. 2). However, when focusing on microbial

decomposers alone, we did not identify a consistent relationship between sumTU and organic

matter decomposition ($p = 0.801$; $n = 60$; Fig. 2). When we examined the effects of individual

chemical groups or modes of toxic action on specific organism groups, no clear dose-response

relationships were evident ($p \geq 0.288$; $n = 8\text{-}22$).

 Photosynthesis

For the ecosystem function photosynthesis, our best-fit model included two predictors:

"sumTU" and "temperature", collectively explaining a portion of the variance ($R^2 = 18.22\%$,

$\tau^2 = 0.0707$; Table 2). In contrast to organic matter decomposition, we did not observe an

influence of the parameter "organism group" (distinguishing between algae and aquatic plants;

Fig. 3). When we examined the effects of individual chemical groups or modes of toxic action

on specific organism groups, no clear dose-response relationships were evident ($p \geq 0.210$; $n =$

12-50).

Primary production

For the ecosystem function primary production, our best-fit model included only "pH"

as predictor, explaining a portion of the variance ($R^2 = 16.06\%$, $\tau^2 = 0.0437$; Fig. 4; Table 2),

but not sumTU. When we examined the effects of individual chemical groups or modes of toxic

action on algae (lack of data on aquatic plants), no clear dose-response relationships were

evident ($p \geq 0.285$; $n = 12$-24).

Discussion

Chemical stressor-ecosystem function relationships

Currently, over 350,000 chemicals are registered for global production and use (Wang

et al., 2020), often occurring as mixtures in surface waters. Furthermore, the Chemical

Abstracts Service has records for over 100 million substances, with approximately 15,000 new

substances being registered daily (Rohr et al., 2016). Considering this ever-expanding array of

chemicals, undertaking comprehensive studies on all currently in use or future developmental

chemicals appears insurmountable (Rohr et al., 2006). Consequently, there is a need to establish

generalizable relationships between chemical exposures and their effects on the environment.

To the best of our knowledge, our study is pioneering in establishing relationships between

chemical stressors and ecosystem functions (Figs. 2 and 3). This represents a crucial initial

stride, as such relationships can ultimately aid in predicting the toxicity of chemicals that have

not yet been thoroughly studied, especially if these chemicals share similar modes of toxic

action. The transferability and predictive capacity of these relationships could potentially

reduce costs and efforts, diminish the reliance on animal testing, and mitigate the suffering of

living organisms – an issue that has received considerable attention in the last decade (Burden

et al., 2015).

All established relationships between chemical exposures and their effects on

ecosystem functioning exhibited an inverted S-shaped monotonicity (Figs. 2 and 3), signifying

that the measured parameters declined as the dose of the chemical stressor increased. Our

findings therefore align with the dose-response patterns frequently observed in the field of

ecotoxicology (Newman, 2009). The S-shape observed in these relationships suggests that

organisms carry specific defense and/or repair mechanisms that provide protection against a

certain amount of a chemical stressor, up to a critical threshold level. Beyond this threshold,

organisms become incapable of defending or repairing themselves effectively. Consequently,

the biological response to chemical stressors showed a graded response due to the evolutionary

adaptive capabilities of organisms to react to a range of low doses. The absence of additional

relationships between chemical exposures and their effects on ecosystem functioning, apart

from those identified for organic matter decomposition and photosynthesis across all chemical

groups, may indicate that the toxic pressure from certain chemical groups did not surpass the

threshold required to overcome the defense and/or repair mechanisms of enough species to

trigger adverse effects.

For the ecosystem function organic matter decomposition by decomposer-detritivore-

systems, for example, we observed a 50% reduction occurring at a sumTU of approximately -

0.5 (Fig. 2). This finding aligns with the sensitivity of macroinvertebrates in the field, as

demonstrated in a large-scale field study involving 101 streams (Liess et al., 2021). In that

study, a sumTU of -0.5 had a pronounced impact on the SPEcies At Risk index, which detects

pesticide effects on the macroinvertebrate community composition in the field (SPEAR$_{pesticides}$).

This indicator pointed to a poor or deteriorated environmental quality that led to the elimination

of key species crucial for ecosystem functioning. These observations align well with another

field study (e.g., Schäfer et al., 2012b), which similarly demonstrated a sharp decline in

SPEAR$_{pesticides}$ at a sumTU of -0.5. This decline resulted in a substantial decrease of

approximately 70% in organic matter decomposition by decomposer-detritivore-systems.

Consequently, there is a good overlap of field data and the relationships between chemical

exposures and their effect on organic matter decomposition observed in our study, despite the

inherent challenges of comparing data sets (e.g., studies mainly focused on individual

chemicals vs. complex mixtures in the field), highlighting the value of our results.

Despite this fairly good overlap of field data and the chemical stress-ecosystem function

relationship derived here, it is essential to acknowledge that the heavy emphasis on specific

chemical groups in research often hampers the broader applicability of laboratory findings to

real-world conditions. Over the past two decades, a predominant focus has been on metals,

pesticides (particularly herbicides), and pharmaceuticals (Figs. 1a and 1b). These chemical

groups closely link to prominent anthropogenic activities such as agriculture, mining, and

wastewater discharge (Bommarco et al., 2013; Nriagu, 1996; Rodriguez-Mozaz et al., 2020).

Modern intensive agriculture heavily relies on pesticide application to mitigate crop losses.

However, following their application, pesticides can be transported into adjacent non-target

ecosystems like freshwater bodies, leading to diffuse pollution (Schulz, 2004). Mining and

wastewater discharge, on the other hand, represent point sources of pollution, resulting in more

localized spatial impacts primarily by metals and pharmaceuticals, though wastewater

treatment plants are highly abundant in densely populated regions of industrialized countries

(Sonter et al., 2018).

The predominant research focus on pesticides, metals, and pharmaceuticals can hence

be justified by their widespread presence and severe impacts on the environment. However, it

is important to recognize that these chemical groups constitute only a small fraction of the vast

array of chemicals present in the environment, given that approximately 350,000 chemicals are

registered for production and use (Wang et al., 2020). Studies employing bioassays have

revealed that target chemicals can only account for a small fraction (as low as < 0.1 %) of the

overall response to chemicals, suggesting that numerous ecotoxicologically important

compounds exist within freshwater ecosystems (Escher et al., 2013; Neale et al., 2020). In

addition to the emphasis on pesticides, metals, and pharmaceuticals, relatively few studies have

considered that chemicals often co-occur as mixtures (Fig. 1b). Typically, the effect of

chemicals in mixtures is additive, but in a relevant portion of cases, the simultaneous presence

of multiple chemicals can result in stronger (synergistic) or weaker (antagonistic) effects

compared to the assumption of additive effects. A recent review, encompassing various

chemical groups, indicated an average of 20% of cases with synergism and 15% of cases with

antagonism (Martin et al., 2021). Another review reported synergistic effects in 3% to 26% of

cases involving binary mixtures, with the prevalence varying by chemical group (Cedergreen,

2014). Furthermore, freshwater ecosystems are typically subject to multiple stressors

concurrently (Birk et al., 2020), which can lead to interactions between chemical and non-

chemical stressors (Liess et al., 2016). To provide a more comprehensive understanding of the

effects of chemicals in freshwater ecosystems and derive additional chemical stress-ecosystem

function relationships, it is essential to broaden the scope. This entails identifying

ecotoxicologically relevant chemical groups, assessing interactions within chemical mixtures,

and considering the context of multiple (non-chemical) stressors in the ecosystem.

Furthermore, finding chemical stressor-ecosystem function relationships can be

confounded by abiotic factors, such as pH and temperature, which influence the fate and

toxicity of metals and organic chemicals in aquatic environments (Ragnarsdottir, 2000; Soares

et al., 2020; Wang et al., 2016). This phenomenon is often referred to as "context dependency"

(Catford et al., 2022). While our findings did not reveal any significant interactions between

pH or temperature as moderators of the toxicity of the chemical stressors on ecosystem

functioning, such interactions have been observed earlier. For instance, a meta-analysis (Wang

et al., 2016) revealed a substantial influence of pH changes on metal toxicity. This can be

explained by the pH-dependency of metal speciation, which in turn affects metal

bioavailability. Furthermore, many pesticides are ionizable compounds (Ragnarsdottir, 2000),

and their degradation and subsequent toxicity are influenced by pH as well as molecular

structural properties (Fenner et al., 2013). Temperature also interacts with the effect of

chemical stressors, a concept known as climate-induced toxicant sensitivity (Noyes et al.,

2009). This interaction can result in lower effects due to increased biotransformation of

chemical stressors at higher temperatures, as well as reduced activity (and consequently, lower

exposure) of organisms at lower temperatures. Moreover, aside from their role in modifying

chemical bioavailability, speciation, and concentration, abiotic conditions have been shown to

increase species sensitivity by up to 30-fold in certain experiments (Liess et al., 2001; Liess

and Beketov, 2011).

Additionally, biotic factors can impede the derivation of chemical stressor-ecosystem

function relationships. However, the relevance of these factors largely depends on the specific

ecosystem function under investigation and the design of the experimental approach.

Laboratory experiments typically involve the use of species that are maintained in culture over

extended periods, often under conditions that differ from their natural habitats (e.g., variations

in light, temperature, and pH) and that may lay outside their niche width and optima. Thus,

these conditions affect the survival, growth, reproduction and functioning of species, and thereby the related ecosystem functions (Blackman, 1905). Indeed, pH and temperature influenced organic matter decomposition and photosynthesis, respectively (Table 2). These findings suggest a mismatch between species-specific requirements and the laboratory conditions under which the respective experiments were conducted. In addition, the acclimation to laboratory conditions can have a notable impact on the sensitivity of these species to chemicals, as highlighted in a recent review (Lavoie et al., 2017). Furthermore, the presence of cryptic species complexes and demographic variability, encompassing both demographic stochasticity (random variations in birth and death rates) and demographic heterogeneity (variations in vital rates among individuals within the same population), is frequently overlooked or inadequately considered in laboratory studies. Nevertheless, these factors have the potential to confound the results of laboratory investigations (Feckler et al., 2013; Melbourne and Hastings, 2008).

In studies that make use of complex natural communities rather than single species tests, additional biotic factors can introduce uncertainties into the derivation of chemical stressor-ecosystem function relationships. These studies typically involve communities with different compositions, and these structural differences have been demonstrated to influence the sensitivity of communities to chemical stress (Beketov et al., 2008; Feckler et al., 2018). Consequently, these variations can lead to differing effects on ecosystem functions due to differences in the capacity of communities to compensate for species loss caused by chemical stressors (Cadotte et al., 2011). Therefore, integrating theoretical advances from community ecology into the toolkit of chemical stressor effect research holds promise (Clements and Rohr, 2009). This approach can help address questions such as "Which types or structures of communities are most sensitive to pollution?", "Which species are most vulnerable to pollution?", and "On which [...] of the chemicals should we focus our attention?". Community

ecology theory, particularly when employing methods like community viability analysis and

network theory, can generate testable predictions by considering factors that influence

community stability and contribute to the propagation of indirect effects (for further reading

see Rohr et al., 2006). Such an integration could enhance our understanding of the intricate

relationships between chemical stressors, community structure, and ecosystem functions.

Finally, chemical stressors have the potential to influence the shapes of biodiversity-

ecosystem function relationships, an effect mediated by species interactions (Baert et al., 2016).

In this context, ecological theory distinguishes between interspecific interactions, which can

change species' contributions to ecosystem functions through competitive replacement, and

intraspecific interactions, which can alter species' contributions to ecosystem functions but

without leading to competitive replacement (Fox, 2005). As the impact of chemical stress on

ecosystem functions is mediated through its effects on species interactions, these interactions

play a pivotal role not only in theories related to biodiversity-ecosystem function relationships

but also in the field of stress ecology (Baert et al., 2016). Consequently, understanding how

and to what extent chemical stress alters species interactions is crucial for developing theories

about stress-induced changes in biodiversity effects. However, the mechanisms behind stress-

induced alterations in the shapes of biodiversity-ecosystem function relationships remain

largely unexplored (Baert et al., 2018, 2016). Therefore, ecotoxicology can benefit from the

application of theoretical ecological approaches utilized in biodiversity-ecosystem function

research. By establishing connections between these research fields, we can work towards

bridging this knowledge gap and gain a mechanistic understanding of how chemical stressors

can modify the shapes of biodiversity-ecosystem function relationships.

Study bias in the selection of ecosystem functions?

We specifically chose biologically mediated functions for our analysis, focusing on

those we deemed most ecologically relevant within freshwater ecosystems. Among the five

selected ecosystem functions, primary production and photosynthesis by algae and aquatic

plants were notably prevalent (Fig. 1a). This prevalence can be attributed to several factors.

Firstly, these functions are likely to be highly responsive to the most studied group of

pesticides, namely herbicides (Hedlund et al., 2020), as well as pharmaceuticals.

Cyanobacteria, as photosynthetic prokaryotes, are considered sensitive to antibiotics due to

their close relationship with pathogenic bacteria (González-Pleiter et al., 2013), and green algae

and plants can also be affected by antibiotics because of the prokaryotic origin of their semi-

autonomous cell organelles (Nie et al., 2009). Secondly, maintaining algal cultures and plants

in laboratory settings is relatively straightforward, and measurement techniques for primary

production and photosynthesis enable rapid quantitative assessment.

Despite their importance for freshwater ecosystem functioning, their potential of being

used as indicators of streams' ecological status and in the assessment of streams' restoration

(Klocker et al., 2009; Udy et al., 2006; Young et al., 2008), only a minor share of studies

addressed chemical stressor effects on microbial community respiration and nutrient cycling

(Fig. 1a). Unlike for algae, standardized protocols for culturing heterotrophic microbial

communities (bacteria and fungi) in the laboratory are largely lacking, hindering their use in

research and complicating comparisons between studies. Moreover, studies assessing

ecosystem functions through microbial decomposers (approximately 24 ± 2 days; mean $\pm$

standard error) typically required considerably longer exposure times compared to those

focusing on primary production and photosynthesis by algae (approximately 5 ± 0.5 days),

which increases the labor-intensity of these studies. Nevertheless, protocols to quantify various

processes related to community respiration and nutrient cycling have been developed and are

frequently applied in field studies to assess stream ecosystem health (von Schiller et al., 2017).

While these protocols are similarly complex and costly as other methods for quantifying

ecosystem functions, they can potentially be partially automated for use in laboratory studies

(von Schiller et al., 2017). Thus, we recommend their application in future laboratory studies

to broaden our mechanistic understanding of how chemical stressors impact ecosystem

functions.

Research gaps and recommendations

Along with the ever-increasing production values and usage of new chemicals, it

becomes insurmountable to experimentally study the effects of all chemicals alone and in

combination with other chemicals and stressors in a wide range of ecosystems, making

generalizations invaluable. In this context, our analyses revealed the potential of deriving

chemical stress-ecosystem function relationships using currently available data that reveal

adverse effects for certain ecosystem functions. Nevertheless, we identified several important

research gaps that should be considered in the future to support deriving more specific (e.g., in

terms of mode of action) chemical stress-ecosystem function relationships and ultimately

facilitate the prediction of chemical effects.

(1) Stronger focus on underrepresented chemical stressors and ecosystem functions

As we showed in our analysis, chemicals of certain groups as well as specific ecosystem

functions were underrepresented in the extracted literature. Therefore, data remains insufficient

for important but understudied chemical groups and ecosystem functions, hampering a more

complete evaluation of chemical stress.

(2) Consideration of environmental variation and complexity of field conditions

Above we showed that the context-dependency of chemical stress effects on ecosystem

functions is largely ignored. Therefore, future laboratory research should proactively consider

a series of abiotic and biotic factors as well as the composition of stressors (single stressors vs.

diverse chemical mixtures; combination of chemical and non-chemical stressors) to ensure the

transferability of the data to various field situations. Ideally, such studies should simulate

different climatic and geological regions as well as chemical exposure histories as both factors

were shown to determine the stress response (cf. Zubrod et al., 2019). In addition, laboratory

studies should be appended by semi-field and field studies as verification and validation

benchmarks to assess the accuracy and reliability of laboratory data for field situations.

(3) Increased effect modeling

Given the high number of used chemicals and the predicted increase in registered chemicals in

the future as well as the context-dependency of chemical effects, empirical testing will be

insurmountable. In this context, modeling approaches offer useful tools to identify general

findings at different levels of biological organization (e.g., microbes and invertebrates). In

addition, effect modeling can help to fill data gaps.

Acknowledgements

The authors thank Jakob Wolfram for making data on field concentrations of pesticides

available.

Funding statement

Alexander Feckler was funded by the Strategy Fund of the RPTU Kaiserslautern-Landau.

Conflict of interest

The authors have declared no conflict of interest.

Data accessibility statement

All data used for the analyses of the present study were extracted from previously published

articles. All data and R code used in this paper will be uploaded to GitHub upon article

acceptance.

Statement of authorship

Alexander Feckler: Conceptualization (equal); Methodology (equal); Investigation (lead);

Formal analysis (lead); Visualization (lead); Data curation (lead); Writing – Original Draft,

Review & Editing (lead). **Ralf Schulz**: Conceptualization (equal); Writing – Original Draft,

Review & Editing (support); **Ralf Schäfer**: Conceptualization (equal); Methodology (equal);

Investigation (support); Formal analysis (support); Writing – Original Draft, Review & Editing

(support); Funding acquisition (lead).

References

Baert JM, Eisenhauer N, Janssen CR, De Laender F. 2018. Biodiversity effects on ecosystem

functioning respond unimodally to environmental stress. *Ecology Letters* **21**:1191–

1199. doi:10.1111/ele.13088

Baert JM, Janssen CR, Sabbe K, Laender FD. 2016. Per capita interactions and stress tolerance

drive stress-induced changes in biodiversity effects on ecosystem functions. *Nature*

Communications **7**:1–8. doi:10.1038/ncomms12486

Balian EV, Segers H, Martens K, Lévéque C. 2008. The Freshwater Animal Diversity

Assessment: an overview of the results In: Balian EV, Lévêque C, Segers H, Martens

K, editors. Freshwater Animal Diversity Assessment, Developments in Hydrobiology.

Dordrecht: Springer Netherlands. pp. 627–637. doi:10.1007/978-1-4020-8259-7_61

Beaumelle L, De Laender F, Eisenhauer N. 2020. Biodiversity mediates the effects of stressors

but not nutrients on litter decomposition. *eLife* **9**:e55659. doi:10.7554/eLife.55659

Beketov MA, Schäfer RB, Marwitz A, Paschke A, Liess M. 2008. Long-term stream

invertebrate community alterations induced by the insecticide thiacloprid: Effect

concentrations and recovery dynamics. *Science of The Total Environment* **405**:96–108.

doi:10.1016/j.scitotenv.2008.07.001

Bernhardt ES, Rosi EJ, Gessner MO. 2017. Synthetic chemicals as agents of global change.

Frontiers in Ecology and the Environment **15**:84–90.

doi:https://doi.org/10.1002/fee.1450

Birk S, Chapman D, Carvalho L, Spears BM, Andersen HE, Argillier C, Auer S, Baattrup-

Pedersen A, Banin L, Beklioğlu M, Bondar-Kunze E, Borja A, Branco P, Bucak T,

Buijse AD, Cardoso AC, Couture R-M, Cremona F, de Zwart D, Feld CK, Ferreira MT,

Feuchtmayr H, Gessner MO, Gieswein A, Globevnik L, Graeber D, Graf W, Gutiérrez-

Cánovas C, Hanganu J, Işkın U, Järvinen M, Jeppesen E, Kotamäki N, Kuijper M,

Lemm JU, Lu S, Solheim AL, Mischke U, Moe SJ, Nõges P, Nõges T, Ormerod SJ,

Panagopoulos Y, Phillips G, Posthuma L, Pouso S, Prudhomme C, Rankinen K,

Rasmussen JJ, Richardson J, Sagouis A, Santos JM, Schäfer RB, Schinegger R,

Schmutz S, Schneider SC, Schülting L, Segurado P, Stefanidis K, Sures B, Thackeray

SJ, Turunen J, Uyarra MC, Venohr M, von der Ohe PC, Willby N, Hering D. 2020.

Impacts of multiple stressors on freshwater biota across spatial scales and ecosystems.

Nature Ecology and Evolution **4**:1060–1068. doi:10.1038/s41559-020-1216-4

Blackman FF. 1905. Optima and limiting factors. Annals of Botany 19:281–295.

Bommarco R, Kleijn D, Potts SG. 2013. Ecological intensification: harnessing ecosystem

services for food security. *Trends in Ecology & Evolution* **28**:230–238.

doi:10.1016/j.tree.2012.10.012

Burden N, Sewell F, Chapman K. 2015. Testing Chemical Safety: What Is Needed to Ensure

the Widespread Application of Non-animal Approaches? *PLOS Biology* **13**:e1002156.

doi:10.1371/journal.pbio.1002156

Cadotte MW, Carscadden K, Mirotchnick N. 2011. Beyond species: functional diversity and

the maintenance of ecological processes and services. *Journal of Applied Ecology*

48:1079–1087. doi:10.1111/j.1365-2664.2011.02048.x

Cardinale BJ, Duffy JE, Gonzalez A, Hooper DU, Perrings C, Venail P, Narwani A, Mace GM,

Tilman D, Wardle DA, Kinzig AP, Daily GC, Loreau M, Grace JB, Larigauderie A,

Srivastava DS, Naeem S. 2012. Biodiversity loss and its impact on humanity. *Nature*

486:59–67. doi:10.1038/nature11148

Catford JA, Wilson JRU, Pyšek P, Hulme PE, Duncan RP. 2022. Addressing context

dependence in ecology. *Trends in Ecology & Evolution* **37**:158–170.

doi:10.1016/j.tree.2021.09.007

Cedergreen N. 2014. Quantifying Synergy: A Systematic Review of Mixture Toxicity Studies

within Environmental Toxicology. *PLOS ONE* **9**:e96580.

doi:10.1371/journal.pone.0096580

Clements WH, Rohr JR. 2009. Community responses to contaminants: Using basic ecological

principles to predict ecotoxicological effects. *Environmental Toxicology and Chemistry*

28:1789–1800. doi:10.1897/09-140.1

Corcoll N, Casellas M, Huerta B, Guasch H, Acuña V, Rodríguez-Mozaz S, Serra-Compte A,

Barceló D, Sabater S. 2015. Effects of flow intermittency and pharmaceutical exposure

on the structure and metabolism of stream biofilms. *Science of The Total Environment*

503–504:159–170. doi:10.1016/j.scitotenv.2014.06.093

Crane M, Watts C, Boucard T. 2006. Chronic aquatic environmental risks from exposure to

human pharmaceuticals. *Science of The Total Environment* **367**:23–41.

doi:10.1016/j.scitotenv.2006.04.010

DeLorenzo ME, Scott GI, Ross PE. 2001. Toxicity of pesticides to aquatic microorganisms: A

review. *Environmental Toxicology and Chemistry* **20**:84–98.

doi:10.1002/etc.5620200108

Deo RP. 2014. Pharmaceuticals in the Surface Water of the USA: A Review. *Current*

Environmental Health Reports **1**:113–122. doi:10.1007/s40572-014-0015-y

Escher BI, van Daele C, Dutt M, Tang JYM, Altenburger R. 2013. Most oxidative stress

response in water samples comes from unknown chemicals: the need for effect-based

water quality trigger values. *Environmental Science and Technology* **47**:7002–7011.

doi:10.1021/es304793h

Feckler A, Bundschuh M. 2020. Decoupled structure and function of leaf-associated

microorganisms under anthropogenic pressure: potential hurdles for environmental

monitoring. *Freshwater Science* **39**:652–664. doi:10.1086/709726

Feckler A, Goedkoop W, Konschak M, Bundschuh R, Kenngott KGJ, Schulz R, Zubrod JP,

Bundschuh M. 2018. History matters: Heterotrophic microbial community structure

and function adapt to multiple stressors. *Global Change Biology* **24**:e402–e415.

doi:10.1111/gcb.13859

Feckler A, Kahlert M, Bundschuh M. 2015. Impacts of Contaminants on the Ecological Role

of Lotic Biofilms. *Bulletin of Environmental Contamination and Toxicology* **95**:421–

427. doi:10.1007/s00128-015-1642-1

Feckler A, Schulz R, Bundschuh M. 2013. Cryptic lineages—same but different? *Integrated*

Environmental   Assessment   and   Management   **9**:172–173.

doi:https://doi.org/10.1002/ieam.1370

Fenner K, Canonica S, Wackett LP, Elsner M. 2013. Evaluating Pesticide Degradation in the

Environment: Blind Spots and Emerging Opportunities. *Science* **341**:752–758.

doi:10.1126/science.1236281

Ferreira V, Koricheva J, Duarte S, Niyogi DK, Guérold F. 2016. Effects of anthropogenic

heavy metal contamination on litter decomposition in streams – A meta-analysis.

Environmental Pollution **210**:261–270. doi:10.1016/j.envpol.2015.12.060

Fox JW. 2005. Interpreting the 'selection effect' of biodiversity on ecosystem function.

Ecology Letters **8**:846–856. doi:10.1111/j.1461-0248.2005.00795.x

Gessner MO, Chauvet E. 2002. A case for using litter breakdown to assess functional stream

integrity. *Ecological Applications* **12**:498–510. doi:10.2307/3060958

González-Pleiter M, Gonzalo S, Rodea-Palomares I, Leganés F, Rosal R, Boltes K, Marco E,

Fernández-Piñas F. 2013. Toxicity of five antibiotics and their mixtures towards

photosynthetic aquatic organisms: Implications for environmental risk assessment.

Water Research **47**:2050–2064. doi:10.1016/j.watres.2013.01.020

Guo J, Boxall A, Selby K. 2015. Do Pharmaceuticals Pose a Threat to Primary Producers?

Critical Reviews in Environmental Science and Technology **45**:2565–2610.

doi:10.1080/10643389.2015.1061873

Harrer M, Cuijpers P, Furukawa T, Ebert DD. 2019. dmetar: Companion R Package For The

Guide "Doing Meta-Analysis in R."

Harrison I, Abell R, Darwall W, Thieme ML, Tickner D, Timboe I. 2018. The freshwater

biodiversity crisis. *Science* **362**:1369–1369. doi:10.1126/science.aav9242

Harrison PA, Berry PM, Simpson G, Haslett JR, Blicharska M, Bucur M, Dunford R, Egoh B,

Garcia-Llorente M, Geamănă N, Geertsema W, Lommelen E, Meiresonne L,

Turkelboom F. 2014. Linkages between biodiversity attributes and ecosystem services:

A systematic review. *Ecosystem Services* **9**:191–203. doi:10.1016/j.ecoser.2014.05.006

Hedlund J, Longo SB, York R. 2020. Agriculture, Pesticide Use, and Economic Development:

A Global Examination (1990–2014). *Rural Sociology* **85**:519–544.

doi:10.1111/ruso.12303

Higgins JPT, Thompson SG. 2004. Controlling the risk of spurious findings from meta-

regression. *Statistics in Medicine* **23**:1663–1682. doi:10.1002/sim.1752

Hooper DU, Chapin FS, Ewel JJ, Hector A, Inchausti P, Lavorel S, Lawton JH, Lodge DM,

Loreau M, Naeem S, Schmid B, Setälä H, Symstad AJ, Vandermeer J, Wardle DA.

2005. Effects of Biodiversity on Ecosystem Functioning: A Consensus of Current

Knowledge. *Ecological Monographs* **75**:3–35. doi:https://doi.org/10.1890/04-0922

Jenkins DG, Quintana-Ascencio PF. 2020. A solution to minimum sample size for regressions.

PLOS ONE **15**:e0229345. doi:10.1371/journal.pone.0229345

Jepsen R, He K, Blaney L, Swan C. 2019. Effects of antimicrobial exposure on detrital biofilm

metabolism in urban and rural stream environments. *Science of The Total Environment*

666:1151–1160. doi:10.1016/j.scitotenv.2019.02.254

Klocker CA, Kaushal SS, Groffman PM, Mayer PM, Morgan RP. 2009. Nitrogen uptake and

denitrification in restored and unrestored streams in urban Maryland, USA. *Aquatic*

Science **71**:411–424. doi:10.1007/s00027-009-0118-y

Kovalakova P, Cizmas L, McDonald TJ, Marsalek B, Feng M, Sharma VK. 2020. Occurrence

and toxicity of antibiotics in the aquatic environment: A review. *Chemosphere*

251:126351. doi:10.1016/j.chemosphere.2020.126351

Kümmerer K. 2009. Antibiotics in the aquatic environment – A review – Part I. *Chemosphere*

75:417–434. doi:10.1016/j.chemosphere.2008.11.086

Lavoie I, Hamilton PB, Morin S, Kim Tiam S, Kahlert M, Gonçalves S, Falasco E, Fortin C,

Gontero B, Heudre D, Kojadinovic-Sirinelli M, Manoylov K, Pandey LK, Taylor JC.

2017. Diatom teratologies as biomarkers of contamination: Are all deformities

ecologically meaningful? *Ecological Indicators* **82**:539–550.

doi:10.1016/j.ecolind.2017.06.048

Lewis K, Tzilivakis J, Green A, Warner DJ. 2011. Veterinary Substances Database (VSDB).

Lewis KA, Tzilivakis J, Warner DJ, Green A. 2016. An international database for pesticide

risk assessments and management. *Human and Ecological Risk Assessment: An*

International Journal **22**:1050–1064. doi:10.1080/10807039.2015.1133242

Liess M, Beketov M. 2011. Traits and stress: keys to identify community effects of low levels

of toxicants in test systems. *Ecotoxicology* **20**:1328–1340. doi:10.1007/s10646-011-

0689-y

Liess M, Champeau O, Riddle M, Schulz R, Duquesne S. 2001. Combined effects of

ultraviolet-B radiation and food shortage on the sensitivity of the Antarctic amphipod

Paramoera walkeri to copper. *Environmental Toxicology and Chemistry* **20**:2088–2092.

doi:10.1002/etc.5620200931

Liess M, Foit K, Knillmann S, Schäfer RB, Liess H-D. 2016. Predicting the synergy of multiple

stress effects. *Scientific Reports* **6**:32965. doi:10.1038/srep32965

Liess M, Liebmann L, Vormeier P, Weisner O, Altenburger R, Borchardt D, Brack W,

Chatzinotas A, Escher B, Foit K, Gunold R, Henz S, Hitzfeld KL, Schmitt-Jansen M,

Kamjunke N, Kaske O, Knillmann S, Krauss M, Küster E, Link M, Lück M, Möder M,

Müller A, Paschke A, Schäfer RB, Schneeweiss A, Schreiner VC, Schulze T,

Schüürmann G, von Tümpling W, Weitere M, Wogram J, Reemtsma T. 2021.

Pesticides are the dominant stressors for vulnerable insects in lowland streams. *Water*

Research **201**:117262. doi:10.1016/j.watres.2021.117262

Loreau M, Naeem S, Inchausti P, Bengtsson J, Grime JP, Hector A, Hooper DU, Huston MA,

Raffaelli D, Schmid B, Tilman D, Wardle DA. 2001. Biodiversity and Ecosystem

Functioning: Current Knowledge and Future Challenges. *Science* **294**:804–808.

doi:10.1126/science.1064088

Malaj E, Ohe PC von der, Grote M, Kühne R, Mondy CP, Usseglio-Polatera P, Brack W,

Schäfer RB. 2014. Organic chemicals jeopardize the health of freshwater ecosystems

on the continental scale. *Proceedings of the National Academy of Sciences* **111**:9549–

9554. doi:10.1073/pnas.1321082111

Martin O, Scholze M, Ermler S, McPhie J, Bopp SK, Kienzler A, Parissis N, Kortenkamp A.

2021. Ten years of research on synergisms and antagonisms in chemical mixtures: A

systematic review and quantitative reappraisal of mixture studies. *Environment*

International **146**:106206. doi:10.1016/j.envint.2020.106206

Melbourne BA, Hastings A. 2008. Extinction risk depends strongly on factors contributing to

stochasticity. *Nature* **454**:100–103. doi:10.1038/nature06922

Molander L, Ågerstrand M, Rudén C. 2009. WikiPharma – A freely available, easily

accessible, interactive and comprehensive database for environmental effect data for

pharmaceuticals. *Regulatory Toxicology and Pharmacology* **55**:367–371.

doi:10.1016/j.yrtph.2009.08.009

Naeem S, Wright JP. 2003. Disentangling biodiversity effects on ecosystem functioning:

deriving solutions to a seemingly insurmountable problem. *Ecology Letters* **6**:567–579.

doi:10.1046/j.1461-0248.2003.00471.x

Neale PA, Braun G, Brack W, Carmona E, Gunold R, König M, Krauss M, Liebmann L, Liess

M, Link M, Schäfer RB, Schlichting R, Schreiner VC, Schulze T, Vormeier P, Weisner

O, Escher BI. 2020. Assessing the Mixture Effects in In Vitro Bioassays of Chemicals

Occurring in Small Agricultural Streams during Rain Events. *Environmental Science*

and Technology **54**:8280–8290. doi:10.1021/acs.est.0c02235

Newman MC. 2009. Fundamentals of Ecotoxicology. CRC Press.

Nie X, Gu J, Lu J, Pan W, Yang Y. 2009. Effects of norfloxacin and butylated hydroxyanisole

on the freshwater microalga Scenedesmus obliquus. *Ecotoxicology* **18**:677–684.

doi:10.1007/s10646-009-0334-1

Noyes PD, McElwee MK, Miller HD, Clark BW, Van Tiem LA, Walcott KC, Erwin KN, Levin

ED. 2009. The toxicology of climate change: Environmental contaminants in a

warming world. *Environment International* **35**:971–986.

doi:10.1016/j.envint.2009.02.006

Nriagu JO. 1996. A History of Global Metal Pollution. *Science* **272**:223–223.

doi:10.1126/science.272.5259.223

Peters K, Bundschuh M, Schäfer RB. 2013. Review on the effects of toxicants on freshwater

ecosystem functions. *Environmental Pollution* **180**:324–329.

doi:10.1016/j.envpol.2013.05.025

Poisot T. 2011. The digitize package: extracting numerical data from scatterplots. *The R*

Journal **3**:25–26.

R Core Team. 2023. R: a language and environment for statistical computing.

Ragnarsdottir KV. 2000. Environmental fate and toxicology of organophosphate pesticides.

Journal of the Geological Society **157**:859–876. doi:10.1144/jgs.157.4.859

Reid AJ, Carlson AK, Creed IF, Eliason EJ, Gell PA, Johnson PTJ, Kidd KA, MacCormack

TJ, Olden JD, Ormerod SJ, Smol JP, Taylor WW, Tockner K, Vermaire JC, Dudgeon

D, Cooke SJ. 2019. Emerging threats and persistent conservation challenges for

freshwater biodiversity. *Biological Reviews* **94**:849–873. doi:10.1111/brv.12480

Richmond EK, Rosi EJ, Reisinger AJ, Hanrahan BR, Thompson RM, Grace MR. 2019.

Influences of the antidepressant fluoxetine on stream ecosystem function and aquatic

insect emergence at environmentally realistic concentrations. *Journal of Freshwater*

Ecology **34**:513–531. doi:10.1080/02705060.2019.1629546

Robson SV, Rosi EJ, Richmond EK, Grace MR. 2020. Environmental concentrations of

pharmaceuticals alter metabolism, denitrification, and diatom assemblages in artificial

streams. *Freshwater Science* 000–000. doi:10.1086/708893

Rodriguez-Mozaz S, Vaz-Moreira I, Varela Della Giustina S, Llorca M, Barceló D, Schubert

S, Berendonk TU, Michael-Kordatou I, Fatta-Kassinos D, Martinez JL, Elpers C,

Henriques I, Jaeger T, Schwartz T, Paulshus E, O'Sullivan K, Pärnänen KMM, Virta

M, Do TT, Walsh F, Manaia CM. 2020. Antibiotic residues in final effluents of

European wastewater treatment plants and their impact on the aquatic environment.

Environment International **140**:105733. doi:10.1016/j.envint.2020.105733

Rohr JR, Kerby JL, Sih A. 2006. Community ecology as a framework for predicting

contaminant effects. *Trends in Ecology & Evolution* **21**:606–613.

doi:10.1016/j.tree.2006.07.002

Rohr JR, Salice CJ, Nisbet RM. 2016. The pros and cons of ecological risk assessment based

on data from different levels of biological organization. *Critical Reviews in Toxicology*

46:756–784. doi:10.1080/10408444.2016.1190685

Rosi-Marshall EJ, Kincaid DW, Bechtold HA, Royer TV, Rojas M, Kelly JJ. 2013.

Pharmaceuticals suppress algal growth and microbial respiration and alter bacterial

communities in stream biofilms. *Ecological Applications* **23**:583–593. doi:10.1890/12-

0491.1

Schäfer RB, Bundschuh M, Rouch DA, Szöcs E, von der Ohe PC, Pettigrove V, Schulz R,

Nugegoda D, Kefford BJ. 2012a. Effects of pesticide toxicity, salinity and other

environmental variables on selected ecosystem functions in streams and the relevance

for ecosystem services. *Science of The Total Environment* **415**:69–78.

doi:10.1016/j.scitotenv.2011.05.063

Schäfer RB, von der Ohe PC, Rasmussen J, Kefford BJ, Beketov MA, Schulz R, Liess M.

2012b. Thresholds for the Effects of Pesticides on Invertebrate Communities and Leaf

Breakdown in Stream Ecosystems. *Environmental Science and Technology* **46**:5134–

5142. doi:10.1021/es2039882

Scharmüller A, Schreiner VC, Schäfer RB. 2020. Standartox: Standardizing Toxicity Data.

Data **5**:46. doi:10.3390/data5020046

Schulz R. 2004. Field Studies on Exposure, Effects, and Risk Mitigation of Aquatic Nonpoint-

Source Insecticide Pollution: A Review. *Journal of Environmental Quality* **33**:419–

448. doi:https://doi.org/10.2134/jeq2004.4190

Schulz R, Bub S, Petschick LL, Stehle S, Wolfram J. 2021. Applied pesticide toxicity shifts

toward plants and invertebrates, even in GM crops. *Science* **372**:81–84.

doi:10.1126/science.abe1148

Shiklomanov IA. 1993. World freshwater resource. Water in crisis: a guide to the qorld's

779   freshwater resources. New York: Oxford University Press.

780   Soares MP, Jesus F, Almeida AR, Domingues I, Hayd L, Soares AMVM. 2020. Effects of pH

781   and nitrites on the toxicity of a cypermetrin-based pesticide to shrimps. *Chemosphere*

782   **241**:125089. doi:10.1016/j.chemosphere.2019.125089

783   Sonter LJ, Ali SH, Watson JEM. 2018. Mining and biodiversity: key issues and research needs

784   in conservation science. *Proceedings of the Royal Society B: Biological Sciences*

785   **285**:20181926. doi:10.1098/rspb.2018.1926

786   Stehle S, Schulz R. 2015. Agricultural insecticides threaten surface waters at the global scale.

787   *Proceedings of the National Academy of Sciences* **112**:5750–5755.

788   doi:10.1073/pnas.1500232112

789   Strayer DL, Dudgeon D. 2010. Freshwater biodiversity conservation: recent progress and

790   future challenges. *Journal of the North American Benthological Society* **29**:344–358.

791   doi:10.1899/08-171.1

792   Tilman D, Isbell F, Cowles JM. 2014. Biodiversity and Ecosystem Functioning. *Annual Review*

793   *of Ecology, Evolution, and Systematics* **45**:471–493. doi:10.1146/annurev-ecolsys-

794   120213-091917

795   Udy JW, Fellows CS, Bartkow ME, Bunn SE, Clapcott JE, Harch BD. 2006. Measures of

796   Nutrient Processes as Indicators of Stream Ecosystem Health. *Hydrobiologia* **572**:89–

797   102. doi:10.1007/s10750-005-9006-1

798   van Aert RCM, Jackson D. 2019. A new justification of the Hartung-Knapp method for

799   random-effects meta-analysis based on weighted least squares regression. *Research*

800   *Synthesis Methods* **10**:515–527. doi:10.1002/jrsm.1356

801   Viechtbauer W. 2010. Conducting Meta-Analyses in R with the metafor Package. *Journal of*

802   *Statistical Software* **36**:1–48. doi:10.18637/jss.v036.i03

von Schiller D, Acuña V, Aristi I, Arroita M, Basaguren A, Bellin A, Boyero L, Butturini A,

Ginebreda A, Kalogianni E, Larrañaga A, Majone B, Martínez A, Monroy S, Muñoz I,

Paunović M, Pereda O, Petrovic M, Pozo J, Rodríguez-Mozaz S, Rivas D, Sabater S,

Sabater F, Skoulikidis N, Solagaistua L, Vardakas L, Elosegi A. 2017. River ecosystem

processes: A synthesis of approaches, criteria of use and sensitivity to environmental

stressors. *Science of The Total Environment* **596–597**:465–480.

doi:10.1016/j.scitotenv.2017.04.081

Wang Z, Meador JP, Leung KMY. 2016. Metal toxicity to freshwater organisms as a function

of pH: A meta-analysis. *Chemosphere* **144**:1544–1552.

doi:10.1016/j.chemosphere.2015.10.032

Wang Z, Walker GW, Muir DCG, Nagatani-Yoshida K. 2020. Toward a Global Understanding

of Chemical Pollution: A First Comprehensive Analysis of National and Regional

Chemical Inventories. *Environmental Science and Technology* **54**:2575–2584.

doi:10.1021/acs.est.9b06379

Wolfram J, Stehle S, Bub S, Petschick LL, Schulz R. 2021. Water quality and ecological risks

in European surface waters – Monitoring improves while water quality decreases.

Environment International **152**:106479. doi:10.1016/j.envint.2021.106479

Young RG, Matthaei CD, Townsend CR. 2008. Organic matter breakdown and ecosystem

metabolism: functional indicators for assessing river ecosystem health. *Journal of the*

North American Benthological Society **27**:605–625. doi:10.1899/07-121.1

Zubrod JP, Bundschuh M, Arts G, Brühl CA, Imfeld G, Knäbel A, Payraudeau S, Rasmussen

JJ, Rohr J, Scharmüller A, Smalling K, Stehle S, Schulz R, Schäfer RB. 2019.

Fungicides: An Overlooked Pesticide Class? *Environmental Science and Technology*

53:3347–3365. doi:10.1021/acs.est.8b04392

Figure captions

Figure 1: (a) Alluvial plot displaying the extracted effect data. The size of each box is

proportional to the number of extracted observations per organism group (left) or ecosystem

function (right). The link width is proportionate to the number of observations per chemical

stressor type (indicated by color). (b) Cumulative number of studies on each type of stressor

studied through time.

Figure 2: Dose-response relationships with 95% confidence bands (grey) between sumTU and

organic matter decomposition by decomposers or decomposer-detritivore-systems, shown as

the proportion of the functioning in the stressor-treatment compared to the stressor-free control.

Colors indicate the chemical stressor groups (see Fig. 1). Please note the slight differences in

the x-axis scale.

Figure 3: Dose-response relationships with 95% confidence bands (grey) between sumTU and

photosynthesis by algae or aquatic plants, shown as the proportion of the functioning in the

stressor-treatment compared to the stressor-free control. Colors indicate the chemical stressor

groups (see Fig. 1). Please note the slight differences in the x-axis scale.

Figure 4: Dose-response relationships with 95% confidence bands (grey) between sumTU and

primary production by algae, shown as the proportion of the functioning in the stressor-

treatment compared to the stressor-free control. Colors indicate the chemical stressor groups

(see Fig. 1). Please note the differences in the x-axis scale.

Table 1: Descriptive statistics on sumTU values for the individual chemical stressor groups.

Chemical stressor group	n	Mean	Min	1. Quartile	2. Quartile (Median)	3. Quartile	4. Quartile
Fungicides	56	0.20	-4.54	-0.82	0.32	1.67	3.49
Herbicides	102	1.53	-2.88	0.87	1.39	2.20	3.99
Insecticides	34	0.74	-4.25	0.43	0.86	1.86	3.60
Pesticide mixtures	26	0.10	-2.31	-1.03	-0.10	0.94	3.27
Pharmaceuticals	26	-0.18	-5.68	-1.52	0.28	0.77	5.56
Metals	94	0.41	-5.13	-0.43	0.69	1.65	2.68
Metal mixtures	10	0.86	-1.26	0.55	0.99	1.43	1.77

855 Table 2: Outcome of meta-regression models assessing the influence of moderator variables on ecosystem functions.

Function	Predictor	Estimate	T-value	df[1]	p-value	Lower CI[2]	Upper CI
Organic matter decomposition	Organism group	0.3793	7.1855	85	0.001	0.2744	0.4843
	sumTU	-0.0326	-0.9749	85	0.485	-0.0992	0.0339
	pH	-0.2455	-2.7898	85	0.007	-0.4205	-0.0705
	Organism group × sumTU	0.1399	3.7094	85	0.020	0.0649	0.2149
Photosynthesis	sumTU	0.0877	4.4305	99	0.001	0.0484	0.1270
	Temperature	-0.0221	-2.8695	99	0.023	-0.0374	-0.0068
Primary production	pH	0.2932	2.8056	42	0.088	0.0823	0.5041

856 [1]df = Degrees of freedom, [2]CI = confidence interval

(a)