A single bout of vigorous exercise decreases subsequent non-exercise physical activity and body temperature

Daisuke Funabashi[1, 2], Shohei Dobashi[1], Kazuki Sameshima[1], Hiroyuki Sagayama[3],

Takeshi Nishijima[4], Takashi Matsui[1, 2*]

[1]Exercise Biochemistry & Sport Neurobiology Division, Institute of Health and Sport Sciences, University of Tsukuba, Tsukuba, Japan.

[2] Advanced Research Initiative for Human High Performance, University of Tsukuba, Tsukuba, Japan

[3]Laboratory of Exercise Nutrition, Institute of Health and Sport Sciences, University of Tsukuba, Ibaraki, Japan

[4]Department of Health Promotion Sciences, Graduate School of Human Health Sciences, Tokyo Metropolitan University, Tokyo, Japan.

*Correspondence

Takashi Matsui, Ph.D.

Assistant Professor

Exercise Biochemistry & Sport Neurobiology Division,

Institute of Health and Sport Sciences, University of Tsukuba

Tel: +81-29-853-2750 E-mail: matsui.takashi.ga@u.tsukuba.ac.jp

Abstract

Exercise benefits the body and mind, but its weight loss effect is less than generally expected.

Although this phenomenon is likely due to a decrease in non-exercise physical activity (NEPA)

and thermogenesis following exercise, the underlying mechanisms and effects of exercise

intensity remain unknown. Here we show the potential role of corticosterone dynamics in the

decline in NEPA and heat production following vigorous exercise. Adult male C57BL/6J mice

were categorized into three groups: sedentary, moderate exercise, and vigorous exercise, with

exercise groups undergoing a 30 min treadmill session. Using an intraperitoneally implanted

activity monitor, NEPA and body temperature (BT) were monitored for two days before and three

days after exercise. The daily synchrony between the NEPA and BT was evaluated using a cross-

correlation function. Plasma corticosterone was also detected at 6 and 24 h after exercise. Notably,

only the vigorous exercise group exhibited a decline in both NEPA and BT, and increased body

weight compared to the sedentary group, despite no observed changes in food intake. Furthermore,

a distinct negative time lag was identified between the daily dynamics of NEPA and BT following

vigorous exercise. A positive correlation was observed between plasma corticosterone levels and

changes in NEPA before and after exercise across all exercise groups. Our findings provide

evidence for vigorous exercise-specific reduction in NEPA, BT, and their synchrony, likely due

to the disturbed circadian rhythm of corticosterone. Declined BT could imply lowered

thermogenesis, providing novel insights into the role of exercise intensity in influencing the

weight loss effects of exercise.

Introduction

Exercise has been widely recommended as a strategy for improving physiological and mental health (1–3). Given the growing prevalence of obesity, a leading cause of many health problems, optimal exercise prescriptions are required to promote a better energy balance and prevent obesity. However, it has been reported that the effect of exercise on body weight loss is often less than the theoretically predicted reduction based on exercise-induced energy expenditure (4–6). This might be because exercise-induced increases in energy expenditure reduce the energy spent on other physiological activities (7, 8). Therefore, we should better understand the physiological and behavioral changes in response to exercise to develop an effective exercise prescription for preventing obesity and promoting a healthy lifestyle.

Non-exercise physical activity (NEPA) is a crucial component of total energy expenditure and is closely associated with fat gain and the risk of obesity (9). Several studies have shown that a reduction in NEPA is likely a critical contributor to insufficient weight loss by exercise (10, 11). However, evidence of the effects of exercise on NEPA remains inconclusive in human studies (12, 13). In rodents, it has been reported that the introduction of a running wheel consistently decreases NEPA in mice in several studies (14–16), suggesting that voluntary exercise negatively affects NEPA. Furthermore, it was recently reported that a single bout of treadmill running decreased NEPA following exercise in mice (17). As described above, exercise certainly decreases NEPA, and this compensatory response may contribute to maintaining the total energy expenditure within a narrow range and insufficient exercise-induced weight loss. However, the effects of exercise type, intensity, and volume on compensatory reduction in NEPA induced by exercise are poorly understood and compensatory changes in other physiological activities after exercise have not been sufficiently investigated.

Although moderate-to vigorous-intensity physical activity is generally recommended for promoting overall health, the benefits of exercise vary depending on exercise intensity (18, 19). There is evidence that vigorous exercise increases stress-related physiological responses compared to moderate exercise intensity (20). Considering that vigorous exercise increases energy demands and contributes to the development of fatigue, it can be reasonably assumed that a substantial compensatory response is observed after vigorous exercise compared with moderate exercise.

However, it is unknown whether a compensatory reduction in NEPA and other physiological activities after exercise occurs in an exercise intensity-dependent manner. Body temperature is a physiological index that is increased by psychiatric stress and affected by metabolic rate (21, 22), whereas a reduction in body temperature is an adaptive mechanism to reduce energy expenditure when food availability is limited (23). Therefore, for a better

understanding of compensatory responses after exercise, it is important to investigate whether
compensatory changes in body temperature also depend on exercise intensity.

Vigorous exercise activates the hypothalamic-pituitary-adrenal axis and increases
corticosterone release depending on exercise intensity (24). Corticosterone exhibits circadian
rhythms that typically rise in the late afternoon to early evening, just prior to the onset of a dark
period in rodents (25, 26), which is synchronized with that of physical activity. In addition,
corticosterone plays a key role in the regulation of stress responses, energy metabolism, and
physical activity in rodents. For example, previous studies have reported that chronic
corticosterone exposure causes obesity and decreases physical activity in mice (27, 28).
Considering these findings showing a close link between corticosterone and physiological
activities related to energy balance, exercise-induced changes in corticosterone are likely to be
associated with compensatory responses in NEPA after exercise. However, whether corticosterone
is associated with a compensatory reduction in NEPA after exercise remains unclear.

Here, we aimed to examine whether NEPA and body temperature decrease following
exercise in an exercise intensity-dependent manner, and whether corticosterone is associated with
decreased NEPA in mice. Furthermore, we attempted to investigate the relationship between daily
changes in NEPA and body temperature following exercise using cross-correlation analysis to
better understand the compensatory responses in NEPA and body temperature after exercise.

100
101

Materials and Methods

Animals
Male C57BL/6 mice (11 weeks old; n = 27) (SLC, Shizuoka, Japan) were used in these
experiments. Three mice were excluded due to an accident during the experiment and the resulting
change in group size. The mice were housed in standard laboratory cages (225 × 338 × 140 mm)
under controlled temperature (22 ± 2 °C) and light (12:12-h light and dark cycle, light at 7:00–
19:00). Food and water were provided ad libitum. All experimental protocols were conducted in
accordance with the guidelines of the University of Tsukuba Animal Experiment Committee.

Experimental designs
A schematic of the experimental procedure is shown in Figure 1A. After 1-week habituation to
the housing environment, mice underwent surgery to implant a device intraperitoneally to
measure NEPA and body temperature, as described previously (29). After the 1-week of recovery,
all mice were habituated to running on a treadmill (Natsume, Tokyo, Japan) for five sessions over
6 days for 30 min/day with incremental increases in speed (from 0 to 25 m/min). After the

habituation period for all mice, we randomly divided them into three groups: sedentary (SE),
moderate (ME), and vigorous (VE) exercise, and exercised on a treadmill for 30 min at each
intensity. Based on a previous study that determined the lactate threshold (LT) during treadmill
running in mice (18), treadmill running was performed at SE (0 m/min), ME (sub-LT, 15 m/min),
and VE (supra-LT, 25 m/min). The NEPA and body temperature were continuously measured in
the home cage (Fig. 1B) from 2 days before to 3 days after the exercise intervention using an
intraperitoneally implanted device. Three days after the first exercise intervention, the mice
performed the same treadmill running again, and blood was sampled at 6 h and 24 h after exercise
to measure plasma corticosterone.

Measuring NEPA and body temperature
NEPA and body temperature were measured using nanotag® (Kissei Comtec Ltd, Japan), a
recently developed body-implantable device (29). Mice aged ten weeks were implanted with
nanotag® intraperitoneally under anesthesia via the intraperitoneal injection of an anesthetic
cocktail (hydrochloric acid medetomidine, 0.3 mg/kg; midazolam, 4 mg/kg; butorphanol tartrate,
5 mg/kg). After surgery, the mice were intraperitoneally administered atipamezole hydrochloride
(an antagonist of medetomidine hydrochloride, 0.3 mg/kg) to recover from anesthesia, thereby
preventing a sustained decline in body temperature. The mice underwent a 1-week recovery
period after surgery because the negative effect on physical activity was confirmed to be nearly
negligible after this period (29). Prior to nanotag® implantation, we set the device to start and
stop recording automatically, according to the experimental procedure (Fig. 1A). The number of
counts was recorded in the nanotag® at 5-minute intervals. After the experiment was completed,
we removed the nanotag® from deeply anesthetized mice, and the data stored in the nanotag®
were retrieved into a computer using the FeliCa® communication system.

Synchrony analysis
Nanotag® allows us to continuously measure NEPA and body temperature simultaneously in mice.
These data were used to calculate the cross-correlation function (CCF) between daily changes in
NEPA and body temperature on day 1 post-exercise, which can detect the strength of correlations
and time lags among the temporal data. The time lag representing the peak CCF was obtained for
each participant to investigate the causality between daily changes in NEPA and body temperature.
A negative time lag indicates that the circadian rhythm of body temperature precedes that of NEPA.

Corticosterone
Corticosterone concentrations in mouse plasma samples were measured using the Corticosterone
ELISA Kit (ARB, #K014-H1) according to the manufacturer's recommended protocols. The

absorbance was determined using a fluorescence microplate reader (Varioskan LUX multimode

microplate reader, Thermo Fisher Scientific, MA, USA).

Statistical analysis

Daily changes in NEPA and body temperature were analyzed using a two-way (time × exercise-

intervention) repeated-measures analysis of variance (ANOVA). Percent changes in NEPA were

also analyzed using two-way (time × group) repeated-measures ANOVA. If a significant

interaction or main effect was observed, Tukey's or Dunnett's multiple comparison tests were

performed. CCF analysis was performed by an original program by Python 11. Body weight

changes, plasma corticosterone levels, CCF, and lag were analyzed using ordinary one-way

ANOVA. Pearson's correlation analysis (plasma corticosterone levels vs. percentage changes in

physical activity) was also performed. All data are presented as mean ± standard error (SEM).

The threshold for statistical significance was set at $P < 0.05$.

Results

Vigorous exercise decreases subsequent non-exercise physical activity and increases body

weight without altering the food intake.

We first compared the daily changes in NEPA during pre-exercise with day 1 post-exercise within

each group (Fig. 1C–E). Although there was no time × exercise-intervention interaction [F (19,

95) = 0.925, $P = 0.554$] in the SE group, the ME and VE groups showed time × exercise-

intervention interaction [F (19, 152) = 1.779, $P < 0.05$; F (19, 152) = 4.470, $P < 0.0001$] (Fig. 1C).

In ME, we observed a significant difference in NEPA at only one time point between pre-exercise

and day 1 post-exercise, while VE exhibited significant differences at some time points (Fig. 1D,

E).

In the analysis of percentage changes in NEPA from pre-exercise to days 1–3 post-

exercise (Fig. 1F), two-way repeated-measures ANOVA revealed a time × group interaction [F (6,

63) = 4.87, $P < 0.001$]. In the post hoc analysis, a significant reduction from pre-exercise to day

1 post-exercise was observed only in the VE group ($P < 0.01$). Furthermore, the percent change

on day 1 post-exercise from pre-exercise in the VE group was significantly lower than in the SE

and ME groups ($P < 0.01$).

Body weight and food intake were measured weekly during the experiment. Changes in

body weight after 24 h of exercise were greater in the VE group than in the SE group ($P < 0.01$,

Fig. 1G). The average food intake per cage divided by the number of mice is shown in Figure 1H.

Vigorous exercise leads to a subsequent reduction in body temperature during the active
dark period.
Daily changes in body temperature were compared between pre-exercise and day 1 post-exercise
within each group (Fig. 2A–C). In the SE group, daily changes in body temperature did not differ
between pre- and post-exercise [$F(19, 95) = 1.39$, $P = 0.152$]. In contrast, ME and VE showed a
time × exercise-intervention interaction [$F(19, 152) = 4.14$, $P < 0.0001$; $F(19, 152) = 5.34$, $P <$
0.0001]. Post hoc analysis revealed a significant decline in body temperature during day 1 post-
exercise compared to pre-exercise at some time points during the dark period in the ME and VE
groups (Fig. 2B, C).
In the pre-exercise period, the average body temperature during the light and dark
periods did not differ between the groups (Fig. 2D). However, on day 1 post-exercise, the average
body temperature during the light period was not different among the groups, while a significant
difference was observed during the dark period ($P < 0.01$). Post hoc analysis revealed that the
body temperature in the VE group was significantly lower than that in the SE group ($P < 0.01$,
Fig. 2E).
204
Vigorous exercise induces a distinct negative time lag between the daily dynamics of non-
exercise physical activity and body temperature.
We showed typical data representing the daily changes in NEPA and body temperature during day
1 post-exercise in the ME and VE, respectively (CCF = 0.683, Lag = 10 min, Fig. 3A; CCF =
0.641, Lag = -10 min, Fig. 3B). There was no significant difference in the CCF values among the
groups ($P = 0.391$, Fig. 3C). On the other hand, the lag time in VE was significantly shifted toward
the negative direction compared to SE and ME ($P < 0.01$, $P < 0.05$, Fig. 3D).
212
Plasma corticosterone levels are positively correlated with changes in NEPA following
exercise in all exercise groups.
No difference was found in plasma corticosterone levels 6 h after exercise among the groups (P
$= 0.160$, Fig. 4A). There were also no differences in plasma corticosterone levels 24 h after
exercise among the groups ($P = 0.138$). On the other hand, plasma corticosterone levels 6 h after
exercise were significantly correlated with the percent changes in NEPA from pre-exercise to day
1 post-exercise in the ME ($r = 0.919$, $P < 0.05$) and VE ($r = 0.832$, $P < 0.05$), respectively (Fig.
4B, C).
221
222
223
224

Discussion

We examined whether a reduction in NEPA and body temperature following exercise depends on

exercise intensity, and investigated the underlying mechanisms whereby NEPA decreases with

exercise. Our study revealed that vigorous exercise specifically causes a substantial compensatory

response, including a reduction in NEPA and body temperature, supporting the insufficient weight

loss effect induced by exercise. Furthermore, delayed circadian rhythms of NEPA from that of

body temperature induced by vigorous exercise and the lack of a rise in corticosterone before

awakening may be associated with a compensatory reduction in NEPA following exercise.

Compensatory reduction in NEPA and body temperature following exercise

So far, there is growing evidence that the effect of exercise on body weight loss is less than

expected. Pontzer et al. demonstrated that increasing physical activity does not simply add total

energy expenditure in a dose-response manner (30). In addition, an animal study reported that the

total energy expenditure does not increase despite the fact that the daily running distance

considerably increases in mice (16). These findings suggest the possibility of exercise-induced

compensatory responses related to the regulation of the energy balance. Although several studies

have reported a compensatory reduction in NEPA after exercise (14, 15), whether the volume or

intensity of exercise decreases NEPA is insufficiently understood. The present study demonstrated

for the first time that NEPA decreases following vigorous exercise, but not moderate exercise (Fig.

1F), suggesting that a compensatory reduction in NEPA following exercise was specifically

caused by vigorous exercise.

In addition, body weight gain was higher only in the vigorous exercise group than in the

sedentary group, despite the fact that energy expenditure during exercise was the highest in the

vigorous exercise group (Fig. 1G). This phenomenon might be attributed to a reduction in the

NEPA. These findings indicate the possibility that vigorous exercise makes mice physically

inactive following exercise, and thus, moderate exercise might be an optimal exercise program

for promoting a physically active lifestyle and preventing obesity.

Body temperature is affected by whole-body metabolic activity, immune activity, and

physical activity levels. Therefore, if exercise causes compensatory changes in physiological

activity, the body temperature is likely to change. In addition, the circadian rhythm of resting

energy expenditure parallels the changes in core body temperature (31). Body temperature

represents the basal metabolic rate when mice are in a physically inactive phase. Our study

demonstrated that moderate exercise decreased body temperature during the late dark period (Fig.

2B), whereas vigorous exercise decreased body temperature subsequent to both the early and late

dark periods (Fig. 2C). During the early dark period, both NEPA and body temperature decreased

following vigorous exercise, and the reduction in body temperature during this period was

considered to result from decreased NEPA.

However, the exercise-induced reduction in body temperature during the late dark
period is likely to result from a decreased basal metabolic rate because NEPA did not decrease
during this period. Taken together, exercise was likely to cause a compensatory reduction in the
basal metabolic rate in mice, independent of exercise intensity. Furthermore, considering that a
compensatory response was observed even following moderate exercise, this result may suggest
that a reduction in the basal metabolic rate preferentially occurs as a compensatory response
following exercise.

In this study, because there was no body weight reduction or increased food intake
following exercise, a reduction in NEPA and body temperature following exercise certainly
contributed to energy compensation. Considering that body weight gain was not increased in the
moderate exercise group, decreased NEPA might largely contribute to body weight gain compared
to the decreased basal metabolic rate. Altogether, we consider that exercise might cause a
compensatory reduction in basal metabolic rate, and thereafter, a compensatory reduction in
NEPA in response to increased exercise intensity, which may partly explain the background of
Pontzer's constrained total energy expenditure model (30). However, we did not assess energy
expenditure. For a better understanding of compensatory responses after exercise, further studies
examining the effect of exercise on the metabolic rate assessed by respiratory analysis are required.
279

Investigation of synchrony analysis in circadian rhythms between NEPA and body
temperature
NEPA and body temperature exhibit circadian rhythms that are highly synchronized; for example,
an increase in these levels is observed during the early dark period. We attempted to investigate
the synchrony in circadian rhythms between NEPA and body temperature as a possible underlying
mechanism of the compensatory reduction in NEPA following exercise. Although the CCF
between daily changes in NEPA and body temperature on day 1 post-exercise did not differ among
groups (Fig. 3C), the time lag significantly shifted toward a negative direction in VE compared
with SE and ME (Fig. 3D). This result indicates that the circadian rhythm of NEPA is different
from that of body temperature following vigorous exercise.

If increased body temperature is partly responsible for the initiation of NEPA, there is a
possibility that a reduction in the sensitivity of NEPA to a rise in body temperature contributes to
compensatory reduction in NEPA following vigorous exercise. For a detailed investigation, NEPA
and body temperature should be measured simultaneously at shorter intervals than those measured
in this study. The synchrony between daily changes in NEPA and body temperature can potentially
be used to investigate the underlying mechanism of the decline in NEPA following exercise.
296

The possible underlying molecular mechanism of compensatory reduction in NEPA following exercise

We first focused on corticosterone, a hormone that represents a circadian rhythm similar to that of physical activity and responds to a bout of exercise depending on its intensity (24, 26). In this study, plasma corticosterone was measured 6 h and 24 h after exercise, at which time points represent the late light period and the early light period, respectively. Plasma corticosterone commonly shows a higher value in the late light to early dark period, namely, before awakening (26). Consistent with previous studies, the average plasma corticosterone levels were higher at 6 h than at 24 h after exercise. Although plasma corticosterone levels were not different among the groups at either time point, plasma corticosterone levels at 6 h after exercise positively correlated with percent changes in NEPA in both the ME and VE groups (Fig. 4B, C). This result indicates that an increase in corticosterone levels before awakening is important to prevent decreased NEPA following exercise. Daily corticosterone release is regulated by many chronobiological mechanisms such as clock-related genes, neuronal activity in the suprachiasmatic nucleus, and hypocretin neurons in the paraventricular nucleus (32, 33). Therefore, daily changes in corticosterone represent a state of biological circadian rhythm. Compensatory changes in physiological activities, including NEPA, following exercise are also associated with impaired biological rhythms of the circadian clock.

In the present study, exercise intensity in the VE group (25 m/min) was probably higher than LT in mice (18). Considering that compensatory reduction in NEPA is specifically caused by vigorous exercise, compensatory reduction in NEPA following exercise might be associated with increased blood lactate levels. It was recently reported that the activation of lactate receptors, known as hydroxycarboxylic acid receptor 1, decreases spontaneous locomotor activity in mice (34). Therefore, it is worth investigating whether lactate signaling is associated with a reduction in NEPA following vigorous exercise. As described above, further studies featuring corticosterone, lactate, or other molecules are required to better understand the underlying mechanism of compensatory reduction in NEPA following exercise.

Conclusion

Our findings provide evidence for vigorous exercise-specific decreases in NEPA, body temperature, and their synchrony. These compensatory responses must be induced by the disturbed circadian rhythm of corticosterone. Adjusting exercise intensity not only dictates the effectiveness of the exercise protocol but also presents potential implications for post-exercise NEPA and thermogenesis. This novel insight paves the way for a comprehensive exercise strategy to address physical and mental health issues, and mitigate the challenges of overweight and obesity.

Acknowledgments

This study was supported by Grant-in-Aid for Scientific Research (B) (22H03478: Rep. T.M.),

Grant-in-Aid for Scientific Research (C) (22K11528: Rep. T.N.), Grant-in-Aid for Research

Activity Start-up (22K21199: Rep. D.F.) by JSPS KAKENHI, and by Japan Science and

Technology Agency (JST) (JPMJFR205M: Rep. T.M.).

Author contributions

D.F. and T.M. conceived and designed the study. D.F., S.D., and K.S. performed the experiments.

D.F. and T.M. analyzed the data. D.F., S.D., K.S., H.S., T.N., and T.M. interpreted the results of

the experiments. D.F. and T.M. prepared the figures. D.F. and T.M. drafted the manuscript. All

authors edited and revised the manuscript, and read and approved the final version of this

manuscript.

Declaration of interest

The authors have no conflicts of interest to declare.

Data availability

The data that support the findings of this study are available from the corresponding author upon

reasonable request.

References

1. Levine JA, Lanningham-Foster LM, McCrady SK, Krizan AC, Olson LR, Kane PH, Jensen

MD, Clark MM. Interindividual variation in posture allocation: possible role in human obesity.

Science. 2005. 307: 584–6. doi: 10.1126/science.1106561.

2. Stillman CM, Esteban-Cornejo I, Brown B, Bender CM, Erickson KI. Effects of Exercise on

Brain and Cognition Across Age Groups and Health States. Trends Neurosci. 2020. 43: 533–

543. doi: 10.1016/j.tins.2020.04.010.

3. Taylor JA, Greenhaff PL, Bartlett DB, Jackson TA, Duggal NA, Lord JM. Multisystem

physiological perspective of human frailty and its modulation by physical activity. Physiol

Rev. 2023. 103: 1137–1191. doi: 10.1152/physrev.00037.2021.

4. Broskey NT, Martin CK, Burton JH, Church TS, Ravussin E, Redman LM. Effect of Aerobic

Exercise-induced Weight Loss on the Components of Daily Energy Expenditure. Med Sci

Sports Exerc. 2021. 53: 2164–2172. doi: 10.1249/MSS.0000000000002689.

5. Melanson EL. The effect of exercise on non-exercise physical activity and sedentary behavior

in adults. Obes Rev. 2017. 1: 40–49. doi: 10.1111/obr.12507.

6. Myers CA, Johnson WD, Earnest CP, Rood JC, Tudor-Locke C, Johannsen NM, Cocreham
S, Harris M, Church TS, Martin CK. Examination of mechanisms (E-MECHANIC) of
exercise-induced weight compensation: study protocol for a randomized controlled trial.
Trials. 2014. 15: 212. doi: 10.1186/1745-6215-15-212.

7. Mansfeldt JM, Magkos F. Compensatory Responses to Exercise Training As Barriers to
Weight Loss: Changes in Energy Intake and Non-exercise Physical Activity. Curr Nutr Rep.
2023. 12: 327–337. doi: 10.1007/s13668-023-00467-y.

8. Riou MÈ, Jomphe-Tremblay S, Lamothe G, Finlayson GS, Blundell JE, Décarie-Spain L,
Gagnon JC, Doucet É. Energy Compensation Following a Supervised Exercise Intervention
in Women Living With Overweight/Obesity Is Accompanied by an Early and Sustained
Decrease in Non-structured Physical Activity. Front Physiol. 2019. 10: 1048. doi:
10.3389/fphys.2019.01048.

9. Levine JA, Eberhardt NL, Jensen MD. Role of nonexercise activity thermogenesis in
resistance to fat gain in humans. Science. 1999. 283: 212–4. doi:
10.1126/science.283.5399.212.

10. Drenowatz C, Grieve GL, DeMello MM. Change in energy expenditure and physical activity
in response to aerobic and resistance exercise programs. Springerplus. 2015. 4: 798. doi:
10.1186/s40064-015-1594-2.

11. Liu XM, Wang K, Zhu Z, Cao ZB. Compensatory effects of different exercise durations on
non-exercise physical activity, appetite, and energy intake in normal weight and overweight
adults. Front Physiol. 2022. 13: 932846. doi: 10.3389/fphys.2022.932846.

12. Drenowatz C. Reciprocal Compensation to Changes in Dietary Intake and Energy
Expenditure within the Concept of Energy Balance. Adv Nutr. 2015. 6: 592–9. doi:
10.3945/an.115.008615.

13. Silva AM, Júdice PB, Carraça EV, King N, Teixeira PJ, Sardinha LB. What is the effect of
diet and/or exercise interventions on behavioral compensation in non-exercise physical
activity and related energy expenditure of free-living adults? A systematic review. Br J Nutr.
2018. 119: 1327–1345. doi: 10.1017/S000711451800096X.

14. de Carvalho FP, Benfato ID, Moretto TL, Barthichoto M, de Oliveira CA. Voluntary running
decreases nonexercise activity in lean and diet-induced obese mice. Physiol Behav. 2016. 165:
249–56. doi: 10.1016/j.physbeh.2016.08.003.

15. Lark DS, Kwan JR, McClatchey PM, James MN, James FD, Lighton JRB, Lantier L,
Wasserman DH. Reduced Nonexercise Activity Attenuates Negative Energy Balance in Mice
Engaged in Voluntary Exercise. Diabetes. 2018. 67: 831–840. doi: 10.2337/db17-1293.

16. O'Neal TJ, Friend DM, Guo J, Hall KD, Kravitz AV. Increases in Physical Activity Result in
Diminishing Increments in Daily Energy Expenditure in Mice. Curr Biol. 2017. 27: 423–430.

404   doi: 10.1016/j.cub.2016.12.009.

405 17. Quintanilha ACS, Benfato ID, Santos RLO, Antunes HKM, de Oliveira CAM. Effects of
406   acute exercise on spontaneous physical activity in mice at different ages. BMC Sports Sci
407   Med Rehabil. 2021. 13: 78. doi: 10.1186/s13102-021-00311-2.

408 18. Morikawa R, Kubota N, Amemiya S, Nishijima T, Kita I. Interaction between intensity and
409   duration of acute exercise on neuronal activity associated with depression-related behavior in
410   rats. J Physiol Sci. 2021. 71: 1. doi: 10.1186/s12576-020-00788-5.

411 19. Okamoto M, Yamamura Y, Liu YF, Min-Chul L, Matsui T, Shima T, Soya M, Takahashi K,
412   Soya S, McEwen BS, Soya H. Hormetic effects by exercise on hippocampal neurogenesis
413   with glucocorticoid signaling. Brain Plast. 2015. 1: 149–158. doi: 10.3233/BPL-150012.

414 20. Otsuka T, Nishii A, Amemiya S, Kubota N, Nishijima T, Kita I. Effects of acute treadmill
415   running at different intensities on activities of serotonin and corticotropin-releasing factor
416   neurons, and anxiety- and depressive-like behaviors in rats. Behav Brain Res. 2016. 298: 44–
417   51. doi: 10.1016/j.bbr.2015.10.055.

418 21. Škop V, Guo J, Liu N, Xiao C, Hall KD, Gavrilova O, Reitman ML. The metabolic cost of
419   physical activity in mice using a physiology-based model of energy expenditure. Mol Metab.
420   2023. 71: 101699. doi: 10.1016/j.molmet.2023.101699.

421 22. Houshyar H, Cooper ZD, Woods JH. Paradoxical effects of chronic morphine treatment on
422   the temperature and pituitary-adrenal responses to acute restraint stress: a chronic stress
423   paradigm. J Neuroendocrinol. 2001. 13: 862–74. doi: 10.1046/j.1365-2826.2001.00713.x.

424 23. Severinsen T, Munch IC. Body core temperature during food restriction in rats. Acta Physiol
425   Scand. 1999. 165: 299–305. doi: 10.1046/j.1365-201x.1999.00488.x.

426 24. Soya H, Nakamura T, Deocaris CC, Kimpara A, Iimura M, Fujikawa T, Chang H, McEwen
427   BS, Nishijima T. BDNF induction with mild exercise in the rat hippocampus. Biochem
428   Biophys Res Commun. 2007. 358: 961–7. doi: 10.1016/j.bbrc.2007.04.173.

429 25. Atkinson HC, Waddell BJ. Circadian variation in basal plasma corticosterone and
430   adrenocorticotropin in the rat: sexual dimorphism and changes across the estrous cycle.
431   Endocrinology. 1997. 138: 3842–8. doi: 10.1210/endo.138.9.5395.

432 26. Dalm S, Enthoven L, Meijer OC, van der Mark MH, Karssen AM, de Kloet ER, Oitzl MS.
433   Age-related changes in hypothalamic-pituitary-adrenal axis activity of male C57BL/6J mice.
434   Neuroendocrinology. 2005. 81: 372–80. doi: 10.1159/000089555.

435 27. Burke SJ, Batdorf HM, Huang TY, Jackson JW, Jones KA, Martin TM, Rohli KE, Karlstad
436   MD, Sparer TE, Burk DH, Campagna SR, Noland RC, Soto PL, Collier JJ. One week of
437   continuous corticosterone exposure impairs hepatic metabolic flexibility, promotes islet β-
438   cell proliferation, and reduces physical activity in male C57BL/6 J mice. J Steroid Biochem
439   Mol Biol. 2019. 195: 105468. doi: 10.1016/j.jsbmb.2019.105468.

28. Bowles NP, Karatsoreos IN, Li X, Vemuri VK, Wood JA, Li Z, Tamashiro KL, Schwartz GJ,
Makriyannis AM, Kunos G, Hillard CJ, McEwen BS, Hill MN. A peripheral endocannabinoid
mechanism contributes to glucocorticoid-mediated metabolic syndrome. PNAS. 2015. 112:
285–90. doi: 10.1073/pnas.1421420112.

29. Funabashi D, Wakiyama Y, Muto N, Kita I, Nishijima T. Social isolation is a direct
determinant of decreased home-cage activity in mice: A within-subjects study using a body-
implantable actimeter. Exp Physiol. 2022. 107: 133–146. doi: 10.1113/EP090132.

30. Pontzer H, Durazo-Arvizu R, Dugas LR, Plange-Rhule J, Bovet P, Forrester TE, Lambert EV,
Cooper RS, Schoeller DA, Luke A. Constrained Total Energy Expenditure and Metabolic
Adaptation to Physical Activity in Adult Humans. Curr Biol. 2016. 26: 410–7. doi:
10.1016/j.cub.2015.12.046.

31. Zitting KM, Vujovic N, Yuan RK, Isherwood CM, Medina JE, Wang W, Buxton OM,
Williams JS, Czeisler CA, Duffy JF. Human Resting Energy Expenditure Varies with
Circadian Phase. Curr Biol. 2018. 28: 3685–3690.e3. doi: 10.1016/j.cub.2018.10.005.

32. Bonnavion P, Jackson AC, Carter ME, de Lecea L. Antagonistic interplay between hypocretin
and leptin in the lateral hypothalamus regulates stress responses. Nat Commun. 2015. 6: 6266.
doi: 10.1038/ncomms7266.

33. Jones JR, Chaturvedi S, Granados-Fuentes D, Herzog ED. Circadian neurons in the
paraventricular nucleus entrain and sustain daily rhythms in glucocorticoids. Nat Commun.
2021. 12: 5763. doi: 10.1038/s41467-021-25959-9.

34. Li J, Xia Y, Xu H, Xiong R, Zhao Y, Li P, Yang T, Huang Q, Shan F. Activation of brain lactate
receptor GPR81 aggravates exercise-induced central fatigue. Am J Physiol Regul Integr
Comp Physiol. 2022. 323: R822–R831. doi: 10.1152/ajpregu.00094.2022.

Figures

Figure 1 Vigorous exercise decreases subsequent non-exercise physical activity and increases body weight without altering food intake.

Schematic of experimental procedure (A). Exercise was performed on the day of exercise as an EX. Images of the home cage of mice (B). Daily changes in NEPA pre-exercise and on day 1 post-exercise in SE (C), ME (D), and VE (E). The pre-exercise was averaged for two days of NEPA before exercise. *P < 0.05, **P < 0.01 vs. pre-exercise. Percent changes in NEPA from pre-exercise to days 1–3 post-exercise (F). Body weight change (G) and food intake (H) after 24 h of exercise. **P < 0.01 vs. SE, ##P < 0.01 vs. ME, $$P < 0.01 vs. pre-exercise. Values represent mean ± SEM (n = 6–9 per group).

Figure 2 Vigorous exercise leads to a subsequent reduction in body temperature during the active dark period.

Daily changes in core body temperature pre-exercise and on day 1 post-exercise in SE (A), ME (B), and VE (C). The pre-exercise was averaged for two days of body temperature before exercise. *P < 0.05, **P < 0.01 vs. Pre-exercise. Average core body temperature during light and dark periods pre-exercise (D) and on day 1 post-exercise (E). **P < 0.01 vs. SE. Values represent mean ± SEM (n = 6–9 per group).

Figure 3 Vigorous exercise induces a distinct negative time lag between the daily dynamics

of non-exercise physical activity and body temperature.

Typical data of daily changes in NEPA and body temperature during day 1 post-exercise in the

ME (A) and VE (B). CCF (C) and lag time (D) of daily changes on day 1 post-exercise. A negative

time lag means that circadian rhythms of body temperature precede that of NEPA. **P < 0.01 vs.

SE, ##P < 0.01 vs. ME. Values represent mean ± SEM (n = 6–9 per group).

564 **Figure 4 Plasma corticosterone levels positively correlated with changes in NEPA following**

565 **exercise in all exercise groups.**

566 Plasma corticosterone levels 6 h after exercise (A). Correlation between plasma corticosterone 6

567 h after exercise and percent changes in NEPA from pre-exercise to day 1 post-exercise in ME (B)

568 and VE (C). Values represent mean ± SEM (n = 3–6 per group).