Title

Elevated epigenetic age in wild compared to lab-raised house mice

Authors

Eveliina Hanski[1,2], Susan Joseph[3], Aura Raulo[1,4], Klara M Wanelik[1,5], Áine O'Toole[6], Sarah C L Knowles[1*], Tom J Little[6*]

[1]University of Oxford

[2]University of Helsinki

[3]King's College London

[4]University of Turku

[5]University of Surrey

[6]University of Edinburgh

*Authors contributed equally; corresponding authors (sarah.knowles@biology.ox.ac.uk and tom.little@ed.ac.uk)

Competing interests

Authors declare no competing interests.

Abstract

Age is a key parameter in population ecology, with myriad biological processes changing with age

as organisms develop early in life and later senesce. As age is often hard to accurately measure

with non-lethal methods, epigenetic methods of age estimation (epigenetic clocks) have become a

popular tool in animal ecology, and are often developed or calibrated using captive animals of

known age. However, few studies have directly compared epigenetic age estimates between wild

and captive or lab-reared animals of the same species, even though age-related epigenetic changes

can be influenced by environmental conditions. Here, we built an epigenetic clock from standard

laboratory house mice (C57BL/6, *Mus musculus)* and then used it to estimate age in a population

of wild mice (*Mus musculus domesticus*) of unknown age. We show that this clock accurately

predicts adult wild mice to be older than juveniles and that wild mice typically increase in

epigenetic age over time, but with wide variation in epigenetic ageing rate among wild individuals.

We further show that, for a given body mass, wild mice are epigenetically older than lab mice, and

that this difference is not explained by accelerated ageing post-capture but is observed even among

the smallest juvenile mice. This suggests different epigenetic age profiles in mice with contrasting

genetic and environmental backgrounds arise very early in life and may be driven by peri- and

postnatal effects on offspring DNA methylation.

Keywords

Epigenetic clock, DNA methylation, ageing, *Mus musculus*

Introduction

Age is a key characteristic for any organism, with numerous biological processes from immune

maturation to reproduction being age-related. Yet, measuring age in wild individuals can be

challenging as date of birth is often unknown. Classic methods for assessing age in wild animals

are often imprecise (e.g., body size) or destructive (e.g., eye lens weight), raising ethical concerns

and preventing longitudinal studies. An alternative approach for measuring the age of wild

individuals relies on the measurement of epigenetic marks in particular genomic regions.

Specifically, at some CpG sites (those where a cytosine is followed by a guanine; Moore et al.,

2013) the proportion of methylated cytosines appears to change linearly with age, and together

these sites can be used to derive an 'epigenetic clock'. An epigenetic clock trained using samples

from individuals of known age can then be used to predict age in individuals of unknown age.

Such epigenetic clocks can provide a more accurate estimate of chronological age among wild

animals than visible characteristics (Mayne et al., 2022; Larison et al., 2021). Epigenetic clocks

have now been developed for a wide range of animal species including baboons, chimpanzees,

humpback whales, wolves, green turtles, and zebras (Jarman et al., 2015; Anderson et al., 2021;

Pinho et al., 2022; De Paoli-Iseppi et al., 2017; Polanowski et al., 2014; Thompson et al., 2017;

Wright et al., 2018; Mayne et al., 2022; Larison et al., 2021; Bors et al., 2021; Sullivan et al., 2022;

Tangili et al., 2023; Ito et al., 2018; Fairfield et al., 2021; Wilkinson et al., 2021), as well as plants

(Gardner et al., 2023).

Alongside measuring chronological age, epigenetic clocks also appear to capture signals of

biological age, typically considered to reflect the accumulated damage and functional decline in

cells, tissues, and organs (Yousefzadeh et al., 2021). Accelerated epigenetic ageing has been linked

to various communicable and non-communicable diseases in both humans and laboratory mice

(Joyce et al., 2021; Morales Berstein et al., 2022; Ambatipudi et al., 2017; Harvanek et al., 2021;

Cao et al., 2022; Peng et al., 2019). Insights have also come from the wild: high social rank is

associated with accelerated epigenetic ageing in wild baboons (Anderson et al., 2021), and

hibernation slows down ageing in marmots and bats (Pinho et al., 2022; Sullivan et al., 2022).

Thus, the use of epigenetic clocks may provide a means of estimating chronological age among

wild animals while simultaneously providing insight into biological ageing in natural settings.

Previous studies using epigenetic clocks have focused on humans (Joyce et al., 2021; Morales

Berstein et al., 2022; Ambatipudi et al., 2017; Harvanek et al., 2021; Cao et al., 2022; Peng et al.,

2019), laboratory (Han et al., 2018; Kerepesi et al., 2022) or wild animals (Prado et al., 2021;

Polanowski et al., 2014; Anderson et al., 2021; De Paoli-Iseppi et al., 2018; Lemaître et al., 2022).

A number of studies have also included both wild and captive (e.g., from a zoo) individuals (Mayne

et al., 2022; Robeck et al., 2021; Ito et al., 2018; Fairfield et al., 2021; Wilkinson et al., 2021).

However, a comparison of wild and laboratory individuals has not been previously conducted.

Comparing epigenetic age and ageing between lab and wild individuals of the same species could

help us understand drivers of biological ageing and its variability in individuals from contrasting

genetic and environmental backgrounds.

Here, we build an epigenetic clock using samples from laboratory mice (*Mus musculus*) and use

this lab-based clock to predict age in house mice (*Mus musculus domesticus*) from a wild

population. Our aims were threefold: (1) to see whether we could develop an epigenetic clock from

lab-based animals capable of accurately capturing differences in chronological age within a wild

population (2) to compare estimates of epigenetic age in lab compared to wild mice of a given size,

to gain insight into biological ageing in lab vs wild settings and (3) to assess the extent of variability

in biological (epigenetic) ageing rate among wild mice. We used faecal samples as a source of

DNA, to develop a non-invasive method that allows longitudinal sampling without ethical or

logistical limitations on sampling frequency, and allows applications of epigenetic age estimates

in contexts where animal capture or handling are not possible. To our knowledge, this is the first

epigenetic clock built with faecal samples. Our results show the potential of such an approach, but

also indicate substantial differences in the estimated epigenetic age of an inbred laboratory

population and an outbred wild population, even from very early in life.

Materials and Methods

Sample collection

A total of 137 faecal samples were collected from 65 individual *Mus musculus* C57BL/6 laboratory

mice (30 females, 35 males) from two animal facilities. The samples were collected in May–

November 2021 at the Biomedical Services Building, Oxford, UK (Animal facility A), and King's

College, London, UK (Animal facility B). The chronological age of the mice varied from 7 to 339

days. The mice were kept in standard housing and were not subject to any interventions before or

during sampling. During sample collection, body mass was recorded for mice from Animal facility

B but not for mice from Animal facility A. However, body mass is tightly correlated with age

among juvenile house mice (Spangenberg et al., 2014; Jax, 2022a; Jax, 2022b) allowing accurate

estimation of mass from age. As such, we estimated body mass for 25 mice under seven weeks of

age from Animal Facility A (for older mice from this facility (*n*=8) body mass was not estimated

and consequently samples from those mice were not included in analyses including body mass).

Body mass estimation was done based on Spangenberg et al. (2014) for 7–20-day old pups and

The Jackson Laboratory C57BL/6 body mass references for 3–7-week-old pups (Jax, 2022a; Jax,

2022b). For the latter age group, estimation was done separately for females and males using the

Jackson Laboratory sex-specific data (Jax, 2022a; Jax, 2022b). To collect faecal samples, mice

were briefly placed on a sterile surface until defecation. Faecal pellets were collected in a sterile

manner, immediately preserved in DNA/RNA Shield, and stored frozen at -80°C until further

processing.

Wild house mouse (*Mus musculus domesticus*) sampling was conducted in April–May 2019, July

2019, September–October 2019, August–September 2020, and April–May 2021 on Skokholm

Island, Wales, UK. Mice were trapped overnight using small Sherman live traps baited with

peanuts and non-absorbent cotton wool for bedding, and with a spray of sesame oil outside the

trap as a lure. Across each of two broad sampling areas (one near the coast and one in the island

interior, named "Quarry" and "Observatory" respectively), on each trapping night 150 traps were

set at dusk and checked at dawn. To prevent cross-contamination, any traps showing signs of

mouse presence were washed and sterilised before being reset using bleach solution (including a

≥60 min soak in 20% bleach) to destroy bacterial cells and DNA.

All newly captured mice were permanently identified by subcutaneous injection of a passive

integrated transponder (PIT) tag. Upon each capture, each mouse was either tagged or identified

(if a recapture), aged, sexed, and measured before being released within 3m of its trapping point.

Age category was assigned based on size and pelage characteristics: small (typically <15g of mass

and <80mm of length) mice were classified as juveniles and full-sized mice (typically >20g and

>80mm of length) were classified as adults. Mice falling between these two size categories were

ranked as sub-adults. Sex was determined using anogenital distance and reproductive state.

Reproductive state was recorded as either non-perforate, perforate, suspected pregnant or lactating

for females, and testes abdominal, small or large for males. At each capture, body mass and body

condition were recorded. Body condition was scored from 1 to 4 by palpating the lower spine and

hips to estimate the amount of subcutaneous fat.

Faecal samples were collected from traps in a sterile manner (shortly after mice were collected

from traps the following day), preserved in DNA/RNA Shield and stored in a -20°C freezer until

the end of fieldwork (maximum 6 weeks after sample collection). At this point they were returned

to the laboratory frozen and stored at -80°C until DNA extraction. A total of 215 samples were

selected from all collected samples (>900) for further processing, by first selecting a longitudinal

dataset (mice sampled ≥2 times over time; total of 54 individuals) with as much variation as

possible in morphometric variables (age, body mass, sex, and reproductive status), as well as

environmental variables (sampling season and sampling area) and then supplementing this with

additional (equally variable) cross-sectional samples (one sample per animal) to increase number

of individuals for cross-sectional analyses up to 130. Variation in variables was achieved by

randomly selecting approximately equal numbers of samples across categories e.g., across

juvenile, sub-adult and adult mice.

DNA extraction, bisulfite conversion and PCR amplification

DNA was extracted from faecal samples using the ZymoBIOMICS DNA MiniPrep Kit according

to the manufacturer's protocol (Zymo Research, Irvine, California, USA). DNA was then bisulfite-

converted using the Zymo EZ DNA Methylation-Gold Kit to convert unmethylated cytosines to

uracil and then thymine (Zymo Research, Irvine, California, USA). PCR amplification was

conducted for five genes previously reported to correlate with chronological age in *Mus musculus*;

Prima1, Hsf4, Kcns1, Gm9312, and *Gm7325* (Han et al., 2018). Amplification was conducted

using the PyroMark PCR Kit according to manufacturer's instructions and primers for the five

genes (QIAGEN, Hilden, Germany; Table S2; Han et al., 2018). PCR conditions were as follows:

15 min initial denaturation at 95℃, 50 cycles of 30 sec denaturation at 58℃, 30 sec primer

annealing at 58℃ and 30 sec extension at 72℃, followed by a 10 minute final extension at 72℃.

Amplification success was confirmed using gel electrophoresis. PCR was repeated with the same

conditions for any reaction that did not produce a band on the gel. The five amplicons (PCR

products) were pooled for each sample into 5-gene libraries by combining all PCR products per

sample. DNA was quantified with Qubit Fluorometer High Sensitivity dsDNA kit and normalised

to 6.25 ng/μl (Thermo Fisher Scientific, Waltham, Massachusetts, USA).

Sequencing, basecalling, and demultiplexing

The Oxford Nanopore Technology (ONT) platform was used for library sequencing, and all ONT

procedures were conducted according to manufacturer's instructions and ONT protocol

NBA_9102_v109_revl_09Jul2020 (Oxford Nanopore Technologies, Oxford, UK). We first used

the ONT Ligation Sequencing Kit (SQK-LSK109) to repair and dA-tail the DNA ends, followed

by ligation of sequencing adaptors to the prepared ends. We then barcoded libraries using the ONT

Native Barcoding Expansion kit (EXP-NBD104 or EXP-NBD196). Approximately 15 ng of the

prepared library was loaded onto a prepared ONT MinION Mk1B R9.4.1 flow cell and sequenced

using the ONT MinKNOW software v21.10.4. Libraries were sequenced across a total of six runs

resulting in a mean of 49,969 reads per sample. A negative control (where DNAse free H_2O was

used instead of pooled amplicons at the start of Nanopore pipeline) was included in three

sequencing runs, and these generated a mean of 200 (range 17–519) reads. One flow cell was used

twice, and washed between runs with the ONT Flow Cell Wash kit (EXP-WSH003). Different

barcodes were used for negative controls across the two sequencing runs where the same flow cell

was used to enable testing for carry-over of reads (only 17 potential carry-over reads were

detected). Raw sequencing data was basecalled and demultiplexed using High Accuracy

basecalling on the ONT Guppy software v5.0.11. The basecalled FASTQ files were then run

through the Apollo pipeline v0.1 (https://github.com/WildANimalClocks/apollo, DOI:

10.5281/zenodo.8426692) to acquire methylation rates for each CpG site within the five genes.

Using the alignment with the reference genes, target sites (cytosines within CpGs) were identified

in each read and determined as either methylated (cytosine) or unmethylated (uracil). The process

was continued for each read, resulting in a proportion of methylated cytosines at each CpG site.

Analyses

The data was analysed and visualised in R v4.1.2 (R Core Team, 2023). The epigenetic clock was

built using the cross-sectional data of 50 samples from C57BL/6 mice housed in two facilities.

Using CpG site specific methylation rates, we used the package glmnet v4.1–3 (Friedman et al.,

2010) to perform a cross-validated elastic net regulatisation using the cv.glmnet function using a

LASSO model (mixing parameter alpha=1) and a leave-one-out cross validation (nfolds=nrow).

We then fitted a final glmnet model using optimal lambda value determined by the cross-

validation. Epigenetic age was then predicted based on the glmnet model using the predict()

function. This clock was validated on an additional 30 samples from C57BL/6 mice (*n*=15 mice,

two samples per animal). We assessed clock performance for estimating chronological age of lab

mice using mean absolute error (MAE; Tangili et al., 2023). We then built another epigenetic clock

as described above using only those CpG sites that (1) showed parallel trends in their methylation

rates against body mass in lab and wild mice (i.e., ones that increased in methylation with body

mass in both systems; Fig. S2–5) and (2) had methylation rates for the majority of wild mouse

samples (we failed to acquire sufficient read counts for methylation rate measurement for all 11

CpG sites from the gene *Gm7325* in 9% of all wild mouse samples). Following these principles,

we chose in total 53 CpG sites. This revised clock was similarly validated with the independent

lab dataset and then used to estimate epigenetic age for all wild house mice for which methylation

rates at the CpG sites included in the clock were successfully measured (*n*=201; 93% of all wild

mouse samples) using linear modeling.

To test for the effect of covariates on predicted epigenetic age in the lab mouse validation dataset,

we fit a linear model with epigenetic age as the dependent variable and chronological age, sex,

cage, and sequencing run ID as predictor variables. A Wilcoxon rank sum test with 1,000

permutations was used to test whether the predicted epigenetic age of wild mice ranked as adults

was older than that of wild mice ranked as juveniles. The ability of the clock to detect an increase

in age among wild mice sampled on two consecutive occasions was tested using a binomial test.

This test used a null hypothesis $p=0.5$ that predicted age was higher at the later time-point than the

earlier time point (i.e. mice are equally likely to increase or decrease in epigenetic age over time).

We also used a linear model to test whether time between sampling points predicted absolute

change in epigenetic age. Age category at the first time-point and sex were also included as

predictors. Mice with less than 25 days between time-points were excluded from this longitudinal

analysis as the mean absolute error (MAE) of the clock in the validation dataset was 25 days (Fig.

1B).

To explore whether lab and wild mice might differ in epigenetic age for a given chronological age,

we made use of the fact that body mass is a good predictor of age when mice are young, in both

lab and wild settings (Spangenberg et al., 2014; Jax, 2022a; Jax, 2022b; Gerber et al., 2021; Gray

et al., 2015). We used a linear model to test whether source (lab/wild) predicted epigenetic age

when controlling for the effects of body mass (a proxy of age). Epigenetic age was the dependent

variable and source, body mass, sex and an interaction between body mass and sex (as the

relationship between mass and age is somewhat sex-dependent, Jax, 2022a; Jax, 2022b; Gerber et

al., 2021; Gray et al., 2015) were fitted as predictor variables. Female wild mice with signs of

ongoing or recent pregnancy (those recorded as suspected pregnant) were excluded, as body mass

will be a less accurate age proxy in these individuals, who are significantly heavier than other

females (one sample per Animal ID; non-pregnant n=48, suspected pregnant n=9; linear model,

$F_{1,54}$= 29.28, p<0.001). Lastly, to explore whether the *rate* of epigenetic ageing differed between

lab and wild mice we used two methods. First, we used a linear model which tested whether the

increase in epigenetic age observed between two consecutive time-points in repeat-sampled mice

varied as a function of the time between samples, source population (lab/wild) and their

interaction. Second, we used Levene's test to ask whether variation in the rate of epigenetic ageing

differed according to source. Here, we used the ratio between change in epigenetic age and change

in time as the response variable, and source (lab/wild) as the explanatory variable.

Ethical statement

Wild mouse work was conducted under Home Office license PPL PB0178858 held at the

University of Oxford, and with research permits from the Islands Conservation Advisory

Committee (ICAC), and Natural Resources Wales.

Results

Construction of a non-invasive epigenetic clock

We used faecal samples from C57BL/6 laboratory mice (*Mus musculus*, n=50, one sample per

animal) from two animal facilities to generate a DNA methylation-based epigenetic clock. We first

built an epigenetic clock using 73 targeted CpG sites from five genes that were previously

associated with age in laboratory mice (*Hsf4*, *Gm9312*, *Kcns1, Gm7325,* and *Prima1;* Table. S1;

Han et al., 2018). Elastic net regression identified 22 CpG sites from the five targeted genes; three

from *Hsf4*, six from *Gm9312*, six from *Kcns1,* one from *Gm7325* and one from *Prima1* (Table

S2). The epigenetic clock built using methylation patterns across these sites had a mean absolute

error (MAE) of 7 days (Pearson's r=0.996, p<0.001; Fig. 1A). We validated the clock by applying

it to an independent set of C57BL/6 mice that were not used in training the clock (n=15, one sample

per animal). Among these lab mice, predicted (epigenetic) age was also strongly correlated with

chronological age (Pearson's r=0.935, p<0.001, MAE=23 days; Fig. 1B). Neither sex, cage nor

sequencing run had a significant effect on epigenetic age (linear model: chronological age

$F_{1,8}$=63.797, p<0.001; sex $F_{1,8}$=0.328, p=0.583; cage $F_{3,8}$=0.040, p=0.989; sequencing run

$F_{1,8}$=0.057, p=0.817). This demonstrates that non-invasive faecal samples can be used to generate

an epigenetic clock in laboratory mice with equivalent or higher accuracy in estimating

chronological age compared to a clock previously derived using blood samples (Han et al., 2018;

MAE=35–41 days in two validation datasets).

As our aim was to develop a lab-based epigenetic clock that could be used to predict age in wild

mice, we next inspected methylation rates of targeted CpG sites (n=73) against body mass (a proxy

for age) in lab and wild mice. The majority of CpG sites showed similar trends (i.e., increase in

methylation rate with body mass; Fig. S2–5). However, nine CpG sites from genes *Prima1* and

Gm9312 showed contradicting patterns, where methylation rate increased in wild but decreased in

lab mice with body mass (all CpG sites from *Prima1*, CpG sites at positions 10, 33, 116, 218, and

224 from *Gm9312*; Fig. S2, S3), and as such we excluded these CpG sites. Further, we failed to

acquire sufficient read counts and thus methylation rates for the 11 CpG sites from the gene

Gm7325 in 20 wild mouse samples (9% of all wild mouse samples), and consequently decided to

remove this gene from the clock model. We then used the above described approach to build an

epigenetic clock with the remaining 53 CpG sites from four genes (Table S3), whose methylation

rates showed a positive linear relationship with body mass (Fig. S2–4). For this clock, elastic net

regression identified 11 CpG sites from genes *Hsp4* and *Kcns1* (Table S3). Here, the slope deviated

more from 1 (where 1 would indicate perfect positive linear relationship between chronological

and epigenetic age) than did the first clock (slope estimate 0.81 ± 0.02 standard error vs $0.98 \pm$

0.01 standard error in training set; Fig 1A, 1C). However, the clock still had a high accuracy with

a MAE of 19 days in the training set (Pearson's $r=0.979$, $p<0.001$; Fig. 1C) and 24 days in the

validation set (Pearson's $r=0.939$, $p<0.001$; Fig. 1D). This second clock was then used for further

analyses in the present study.

Figure 1. The relationship between DNA methylation-based (epigenetic) age and chronological age in C57BL/6 mice (**A**, **C**) used to train the epigenetic clock model or (**B**, **D**) an independent set of mice (a validation set) not used in building the epigenetic clock model. Two epigenetic clocks were built: one with 73 input CpG sites from five genes (**A**, **B**), and one with 53 input CpG sites from four genes (**C**, **D**). Circles represent individual mice, circle colour in A indicates animal facility (*grey*= facility A, *black* = facility B). Solid lines are linear regression lines, while dashed lines are a reference line of *y=x* (the hypothetical relationship if chronological age and epigenetic age estimates were exactly equivalent).

Chronological age prediction in wild mice

We next applied the lab-mouse derived epigenetic clock (Fig. 1C, 1D) to 205 faecal samples from

119 wild house mice to test if it can be used to estimate chronological age in wild individuals of

unknown age. Mice of all available body sizes were included with the aim of capturing as much

age variation as possible (body mass range 5.9–43.0g, mean 19.0, median 19.8). The epigenetic

age of wild mice varied from -26 to 460 days (mean 227, median 207; 2 out of 205 samples (1%)

had a negative epigenetic age).

Mice assigned as adults in the field using external characteristics were estimated by the epigenetic

clock model to be significantly older than those assigned as juveniles (permutational Wilcoxon

rank sum test, $p<0.001$, $n=166$; Fig. 2A). Moreover, among 35 wild mice sampled twice between

30 and 340 days apart (mean 129, median 81), a great majority (31; 89%) were epigenetically older

at the latter timepoint (binomial test for H_0 p=0.5, $p<0.001$). In general, the number of days

between sampling time-points positively predicted change in epigenetic age (linear model,

$F_{1,17}=8.516$, $p=0.010$; Pearson's $r=0.406$, $p=0.01$, $n=35$), but there was wide variation in the

slope observed among individuals (Fig. 2B). Rate of epigenetic ageing

$(\frac{absolute\ change\ in\ epigenetic\ age}{days\ elapsed})$ was not predicted by any investigated variables (sex $F_{1,18}=0.366$,

$p=0.552$; reproductive status $F_{11,18}=0.653$, $p=0.763$, body mass $F_{1,18}=0.046$, $p=0.834$, season

$F_{2,18}=0.370$, $p=0.696$, and trapping area $F_{1,18}=1.986$, $p=0.176$ at first sampling point). Together

these results indicate that an epigenetic clock trained with samples from inbred lab mice can be

used to provide an estimate of chronological age in outbred wild mice, though not one that is highly

precise.

Figure 2. (**A**) Epigenetic age of wild mice phenotypically characterised as adult (*n*=109) and

juvenile (*n*=57) predicted with an epigenetic clock built using C57BL/6 lab mice. Age category

was assigned in the field using body size and appearance (see Methods). Median epigenetic age

was 316 days in adults and 110 days in juveniles. Circles are individual samples. (**B**) Change in

epigenetic age between two timepoints in wild mice sampled twice between 30 and 340 days apart

(*n*=35). Lines represent individual mice. Epigenetic age increased with time for 31 out of 35 (89%)

mice.

Wild mice are multiple times older epigenetically than laboratory mice

We next assessed whether epigenetic age for a given chronological age differed between wild and

laboratory mice. In the absence of known chronological age for wild mice, we used body mass to

provide an upper limit age estimate for individuals classed as juveniles. Others have reported that

12–13-day-old wild house mice from mainland Europe weigh around 7g (range 3.6–10.5g, mean

6.8; Gerber et al., 2021) and another study showed that 14-day-old wild-derived but captive house

mice from Gough Island (home to the largest wild house mice recorded) weigh around 8.5g (range

~7–10.5g, raw data not available; Gray et al., 2015; Fig. S1). Thus, irrespective of context, house

mice between 12–14 days typically are expected to weigh 7–10.5g (Gerber et al., 2021; Gray et

al., 2015). We therefore examined epigenetic age from a cross-sectional set of juvenile wild

Skokholm Island mice that fall within this body mass range and thus which we expect to be no

more than 25 days old (n=19, body mass 6.1–10.4g with a mean of 8.4g). Among these individuals,

the mean epigenetic age estimate was 106 days (range -26–173, median 110; 1 sample had a

negative epigenetic age), which is more than four times their expected chronological age (Fig. S1).

To explore whether wild mice were epigenetically older than lab mice beyond early life, we

explored the relationship between epigenetic age and body mass across mice of all sizes. While

the reliability of body mass as an indicator of age declines after initial growth during first few

weeks of life, it continues to increase with chronological age in both C57BL/6 lab and wild mice

beyond this time and thus can be used as a rough estimate of age in adults as well (Jax, 2022a; Jax,

2022b; Gerber et al., 2021; Gray et al., 2015). In a cross-sectional subset of 57 lab and 99 wild

mice (excluding wild mice with visual indication of ongoing/recent pregnancy, see Methods, and

those with unknown sex), source (lab/wild) predicted epigenetic age (linear model,

$F_{1,135}$=93.0143, p<0.001; controlling for an interaction term between sex and body mass,

$F_{1,135}$=0.8519, p=0.357), with wild mice having a higher mean epigenetic age than that of lab mice

consistently across investigated range of body mass (Fig. 3A). In line with this finding, wild mice

also had higher rates of DNA methylation in the targeted genes (Fig. S2–4), explaining the higher

epigenetic age profile.

To further examine whether older epigenetic age profiles among wild mice might be due to

accelerated ageing through exposure to environmental stressors, such as food shortage or climatic

variation, we studied the rate of epigenetic ageing across lab and wild mice for which two

timepoints were available. In lab mice (n=15), the rate of epigenetic ageing was similar to that

observed when originally validating the clock using the cross-sectional data (longitudinal data:

slope estimate = 0.77 ± 0.15 standard error; cross-sectional data: slope estimate = 0.80 ± 0.08

standard error; Fig. 1B). This rate was slightly shallower in wild mice (n=35; slope estimate = 0.49

± 0.19 standard error), but not significantly so (linear model: $F_{1,48}$=0.085, p=0.772, Fig. 3B). Lab

mice included were all adults, while the longitudinal data of wild mice included juvenile mice;

however, exclusion of juvenile mice (<15g of body mass) did not appear to have a strong effect

on these trends (Fig. S6). Further, although the rate of epigenetic ageing appeared to be more

variable among wild than lab mice, the variance in epigenetic ageing rate did not differ statistically

across settings (Levene's test, $F_{1,48}$=3.140, p=0.083; correlation between change in epigenetic age

and days elapsed: lab mice, Pearson's r=0.81, p<0.001; wild mice, Pearson's r=0.48, p=0.003; Fig.

3B).

Figure 3. Wild mice have elevated epigenetic age compared to lab mice for a given body mass,

but do not show a different rate of epigenetic ageing in adulthood (**A**) Epigenetic age of wild

(*n*=99) and laboratory mice (*n*=57) in relation to body mass. Empty circles are wild mice and filled

circles are lab mice; black circles are lab mice for which body mass was recorded during sample

collection, and grey circles are lab mice for which body mass was estimated post hoc (see

Methods). (**B**) Change in epigenetic age in relation to time elapsed in laboratory (*n*=15) and wild

mice (*n*=35) sampled twice a minimum 25 days apart. Body mass varies 19.2–26.3g for lab mice

and 7.0–37.7g for wild mice. Empty circles are wild mice and filled circles are lab mice. Lines are

linear regression lines (*dashed* = wild mice, *solid* = lab mice) and shading indicates 95%

confidence intervals.

Discussion

Here, we tested an approach for estimating age in wild house mice, by building an epigenetic clock

using samples from inbred C57BL/6 laboratory mice and using it to estimate age in outbred wild

mice of unknown chronological age. The clock effectively distinguished wild juveniles from

adults, and typically predicted increases in age over time among repeat-captured individuals at a

similar success rate (~90% individuals predicted older at later time point) to the wild baboon study

of Anderson et al (2021)).

However, while the clock accurately predicted age (with error of $\pm$ 25 days) in an independent set

of laboratory mice, we observed high variation among wild mice in how their epigenetic age

changed over chronological time, suggesting our clock had reduced accuracy in this different

ecological context. Others have had better success in applying clocks built with captive individuals

to wild individuals (e.g. Mayne et al., 2022 in green turtles; Robeck et al., 2021 in cetaceans;

correlation between chronological and epigenetic age in these studies 0.67–0.98 *vs* 0.41 between

change in time and change in epigenetic age in our study). However, these studies have built

epigenetic clocks using samples from captive individuals where there is genetic and environmental

variation, such as zoos or outdoor enclosures. Our clock was built with samples from inbred lab

mice housed under very stable environmental conditions, but applied to wild mice that are outbred

and exposed to a highly variable temperate climate. Studies of different lab strains have also

confirmed that epigenetic clocks may behave differently in different genetic backgrounds. For

instance, Han et al demonstrated DBA/2 mice to be up to twice as old epigenetically as C57BL/6

mice (Han et al., 2018).

Moreover, DNA methylation may be influenced by environmental factors (Zocher et al., 2021;

Parrott et al., 2014; Viitaniemi et al., 2019), and wild animals are generally exposed to more

variable environments. As such, the contrasting genetic and environmental backgrounds in our

mouse systems may partly explain why age estimates in wild mice based on a clock from lab mice

had low accuracy. Finally, while all samples from lab mice were preserved immediately after

defecation, the time between defecation and sample preservation varied in wild mice (where

samples were collected from traps which animals had been in overnight, up to 13 hours). It is

possible some DNA degradation by nuclease and gut microbes occurred before the samples were

preserved (in DNA/RNA Shield, which inactivates microbes and preserves DNA integrity),

affecting the methylation profiles.

The epigenetic age of wild mice from Skokholm Island varied from -26 to 460 days (mean 227,

median 207; 2 (1% of all 205) samples had negative epigenetic ages, -26 and -21 days). The

presence of negative predicted ages in wild individuals may be due to measurement error, but it

may also be due to the inherent biological differences between wild and lab populations. DNA

methylation-based age predictions may reflect the distinct age trajectories and relative age of wild

individuals compared to their lab counterparts, highlighting the importance of considering the

natural variation in epigenetic ageing processes across different populations.

Despite our epigenetic clock having a very close relationship with chronological age in laboratory

mice, several lines of evidence suggest these epigenetic age estimates for wild mice are

overestimates of their chronological age. First, when investigating juveniles, in which body mass

is an accurate predictor of age across both wild and lab mice (Jax, 2022a; Jax, 2022b; Gerber et

al., 2021; Gray et al., 2015), we found that wild juveniles were predicted to be several times older

than their expected chronological age from body mass (Gerber et al., 2021; Gray et al., 2015).

Second, by comparing the epigenetic age estimates to body mass in mice of all sizes, we found

that the estimated epigenetic age of wild mice was on average several times higher at any given

body mass than that of lab mice, with similar patterns seen in CpG site methylation rates. While

accelerated weight gain in *ad libitum*-fed lab mice may contribute to a lower epigenetic age among

adult lab mice compared to wild mice, these findings suggested that the more challenging

environment experienced by wild mice may accelerate epigenetic clocks. Our comparison of the

epigenetic age of wild vs lab mice is specific to a comparison with the C57BL/6 strain; however,

the difference between the epigenetic age of e.g., C57BL/6 and DBA/2 mice is substantially

smaller than the observed difference between C57BL/6 and wild mice (DBA/2 mice are only twice

as old epigenetically than are C57BL/6; Han et al., 2018). As such, while the choice of strain may

have influenced our findings, it is unlikely to entirely explain the difference between lab and wild

mice.

To test whether the older epigenetic age profile of wild mice could be explained by accelerated

ageing post-weaning (i.e., from when they are trappable) we investigated the rate of epigenetic

ageing using individuals captured and sampled twice over time. If anything, the rate of epigenetic

age appeared slightly slower and more variable in wild mice, though this observation relied on a

relatively small sample size and was not statistically significant. Moreover, as we observed

heightened epigenetic age in wild compared to lab mice even during the first ~2 weeks of life, we

suggest that peri- and early postnatal effects on offspring DNA methylation may vary between

laboratory and wild mice and contribute to their different epigenetic age profiles. Various human,

mouse, and other animal studies have demonstrated the association between prenatal maternal

experience (such as food shortage, diet, infection, substance exposure, and stress) and offspring

DNA methylation patterns, with differences from the prenatal (foetal) phase still detectable in later

life (Tobi et al., 2009; Heijmans et al., 2008; Lan et al., 2013; Richetto et al., 2017; Camerota et

al., 2021; Joubert et al., 2016; Kertes et al., 2016; Vangeel et al., 2017).

While our approach of training an epigenetic clock with lab individuals and using it to estimate

age in wild individuals did not allow precise estimation of chronological age, our results

demonstrate such an approach can still be effective in distinguishing between juvenile and adult

individuals. Such information may be useful in contexts where a faecal deposit is found but the

individual is not observed, such as in field-based projects of animals that are hard or impossible to

capture. At the same time, this method can provide interesting insights into biological ageing when

applied to wild animals of known chronological age or to individuals sampled longitudinally such

that changes in epigenetic age can be estimated (De Paoli-Iseppi et al., 2017; Powell & Proulx,

2003; Brivio et al., 2015). Considering the much greater variability in epigenetic ageing rates we

observed in wild compared to laboratory animals, our results suggest wild systems may provide

an informative environment in which to study drivers of epigenetic age acceleration. In the present

study we were not able to identify drivers of epigenetic ageing rates in the wild due to a small

sample size, but further work with larger samples sizes could address this important question.

We used faeces as a source of host DNA and demonstrated that an epigenetic clock can be built

with faecal samples at a similar or even improved accuracy to a previously published blood-based

mouse epigenetic clock (Han et al., 2018). As faecal samples can be collected non-invasively, a

given individual can be sampled over time without limitations to sampling frequency. Further, it

may be possible to collect faecal samples without capturing the animal, making this

methodological approach particularly useful when estimating age in wild species that are

endangered, hard to capture or even detect.

In summary, our data indicate the potential to use a non-invasive, DNA methylation-based

epigenetic clock built with samples from laboratory mice to estimate the age of entirely wild mice.

While this approach did not provide highly precise estimates of chronological age, it can be used

to measure variation in biological ageing in future longitudinal studies, making it a promising tool

for studies of ontogeny and senescence in wild settings.

Funding

494 This work was funded by The Osk. Huttunen Foundation studentship and the National Geographic

495 Society (Early Career grant reference No. EC-58520R-19) to EH, the European Research Council

496 (ERC) under the European Union's Horizon 2020 research and innovation programme (Grant

497 agreement No. 851550) and a NERC fellowship (NE/L011867/1) to SCLK, and the British

498 Ecological Society (BES) to TJL.

Acknowledgements

501 We thank Giselle Eagle and Richard Brown, the wardens of Skokholm Island, the Friends of

502 Skokholm and Skomer, the Wildlife Trust of South and West Wales and field assistants for their

503 help in enabling the wild mouse data collection.

Author information

Contributions

EH and SCLK set up the wild mouse study system. EH, SJ, AR and KW collected the samples.

AO and TJL developed the software Apollo. EH conducted the laboratory work and analysed the

data with support from TJL, SCLK and AR. EH wrote the manuscript with contributions from all

authors.

Data accessibility

Data Accessibility

Data will be made publicly available upon publication.

References

Ambatipudi S, Horvath S, Perrier F, et al. DNA methylome analysis identifies accelerated
epigenetic ageing associated with postmenopausal breast cancer susceptibility. *Eur J Cancer*. 2017;75:299-307. doi:10.1016/j.ejca.2017.01.014

Anderson JA, Johnston RA, Lea AJ, et al. High social status males experience accelerated
epigenetic ageing in wild baboons. *Elife*. 2021;10:e66128. Published 2021 Apr 6. doi:10.7554/eLife.66128

Berry RJ, Jakobson ME. Life and death in an island population of the house mouse. *Exp Gerontol*. 1971;6: 187–197.

Bors EK, Baker CS, Wade PR, et al. An epigenetic clock to estimate the age of living beluga

whales. *Evol Appl*. 2021;14(5):1263-1273. Published 2021 Feb 3. doi:10.1111/eva.13195

Brivio F, Grignolio S, Sica N, Cerise S, Bassano B. Assessing the Impact of Capture on Wild

Animals: The Case Study of Chemical Immobilisation on Alpine Ibex. *PLoS One*.

2015;10(6):e0130957. Published 2015 Jun 25. doi:10.1371/journal.pone.0130957

Camerota M, Graw S, Everson TM, et al. Prenatal risk factors and neonatal DNA methylation in

very preterm infants. *Clin Epigenetics*. 2021;13(1):171. Published 2021 Sep 10.

doi:10.1186/s13148-021-01164-9

Cao X, Li W, Wang T, et al. Accelerated biological ageing in COVID-19 patients. *Nat Commun*.

2022;13(1):2135. Published 2022 Apr 19. doi:10.1038/s41467-022-29801-8

De Paoli-Iseppi R, Deagle BE, McMahon CR, Hindell MA, Dickinson JL, Jarman SN. Measuring

Animal Age with DNA Methylation: From Humans to Wild Animals. *Front Genet*.

2017;8:106. Published 2017 Aug 17. doi:10.3389/fgene.2017.00106

Fairfield EA, Richardson DS, Daniels CL, Butler CL, Bell E, Taylor MI. Ageing European lobsters

(*Homarus gammarus*) using DNA methylation of evolutionarily conserved ribosomal

DNA. *Evol Appl*. 2021;14(9):2305-2318. Published 2021 Sep 23. doi:10.1111/eva.13296

Ferrari, Manuela; Lindholm, Anna K; König, Barbara (2015). The risk of exploitation during

communal nursing in house mice, *Mus musculus domesticus.* Animal Behaviour, 110:133-

143.

Flurkey, Currer, and Harrison, 2007. 'The mouse in biomedical research.' in James G. Fox (ed.),

American College of Laboratory Animal Medicine series (Elsevier, AP: Amsterdam;

Boston).

Friedman J, Hastie T, Tibshirani R. Regularization Paths for Generalized Linear Models via
Coordinate Descent. *J Stat Softw*. 2010;33(1):1-22.

Gardner ST, Bertucci EM, Sutton R, Horcher A, Aubrey D, Parrott BB. Development of DNA
methylation-based epigenetic age predictors in loblolly pine (Pinus taeda). *Mol Ecol*
Resour. 2023;23(1):131-144. doi:10.1111/1755-0998.13698

Gerber N, Auclair Y, König B, Lindholm AK. Population density and temperature influence the
return on maternal investment in wild house mice. *Front Ecol Evol.* 2021;8:602359.
https://doi.org/10.3389/fevo.2020.602359

Gray MM, Parmenter MD, Hogan CA, et al. Genetics of Rapid and Extreme Size Evolution in
Island Mice. *Genetics*. 2015;201(1):213-228. doi:10.1534/genetics.115.177790

Han Y, Eipel M, Franzen J, et al. Epigenetic age-predictor for mice based on three CpG sites. *Elife*.
2018;7:e37462. Published 2018 Aug 24. doi:10.7554/eLife.37462

Harvanek ZM, Fogelman N, Xu K, Sinha R. Psychological and biological resilience modulates the
effects of stress on epigenetic ageing. *Transl Psychiatry*. 2021;11(1):601. Published 2021
Nov 27. doi:10.1038/s41398-021-01735-7

Heijmans BT, Tobi EW, Stein AD, et al. Persistent epigenetic differences associated with prenatal
exposure to famine in humans. *Proc Natl Acad Sci U S A*. 2008;105(44):17046-17049.
doi:10.1073/pnas.0806560105

Ito H, Udono T, Hirata S, Inoue-Murayama M. Estimation of chimpanzee age based on DNA
methylation. *Sci Rep*. 2018;8(1):9998. Published 2018 Jul 3. doi:10.1038/s41598-018-
28318-9

Jarman SN, Polanowski AM, Faux CE, et al. Molecular biomarkers for chronological age in animal
ecology. *Mol Ecol*. 2015;24(19):4826-4847. doi:10.1111/mec.13357

JAX®. Body weight information for aged C57BL/6J. Available at: https://www.jax.org/jax-mice-
and-services/strain-data-sheet-pages/body-weight-chart-aged-b6 (Accessed: 30.11.2022)

JAX®. Body weight information for C57BL/6J. Available at: https://www.jax.org/jax-mice-and-
services/strain-data-sheet-pages/body-weight-chart-000664 (Accessed: 30.11.2022)

Joubert BR, Felix JF, Yousefi P, et al. DNA Methylation in Newborns and Maternal Smoking in
Pregnancy: Genome-wide Consortium Meta-analysis. *Am J Hum Genet*. 2016;98(4):680-
696. doi:10.1016/j.ajhg.2016.02.019

Joyce BT, Gao T, Zheng Y, et al. Epigenetic Age Acceleration Reflects Long-Term Cardiovascular
Health. *Circ Res*. 2021;129(8):770-781. doi:10.1161/CIRCRESAHA.121.318965

Kerepesi C, Meer MV, Ablaeva J, et al. Epigenetic ageing of the demographically non-ageing
naked mole-rat. *Nat Commun*. 2022;13(1):355. Published 2022 Jan 17.
doi:10.1038/s41467-022-27959-9

Kertes DA, Kamin HS, Hughes DA, Rodney NC, Bhatt S, Mulligan CJ. Prenatal Maternal Stress
Predicts Methylation of Genes Regulating the Hypothalamic-Pituitary-Adrenocortical
System in Mothers and Newborns in the Democratic Republic of Congo. *Child Dev*.
2016;87(1):61-72. doi:10.1111/cdev.12487

Lan X, Cretney EC, Kropp J, et al. Maternal Diet during Pregnancy Induces Gene Expression and
DNA Methylation Changes in Fetal Tissues in Sheep. *Front Genet*. 2013;4:49. Published
2013 Apr 5. doi:10.3389/fgene.2013.00049

Larison B, Pinho GM, Haghani A, et al. Epigenetic models developed for plains zebras predict age
in domestic horses and endangered equids. *Commun Biol*. 2021;4(1):1412. Published 2021
Dec 17. doi:10.1038/s42003-021-02935-z

Lemaître JF, Rey B, Gaillard JM, et al. DNA methylation as a tool to explore ageing in wild roe

deer populations. *Mol Ecol Resour*. 2022;22(3):1002-1015. doi:10.1111/1755-0998.13533

Mayne B, Mustin W, Baboolal V, et al. Age prediction of green turtles with an epigenetic clock.

Mol Ecol Resour. 2022;22(6):2275-2284. doi:10.1111/1755-0998.13621

Moore LD, Le T, Fan G. DNA methylation and its basic function. *Neuropsychopharmacology*.

2013;38(1):23-38. doi:10.1038/npp.2012.112

Morales Berstein F, McCartney DL, Lu AT, et al. Assessing the causal role of epigenetic clocks

in the development of multiple cancers: a Mendelian randomization study. *Elife*.

2022;11:e75374. Published 2022 Mar 29. doi:10.7554/eLife.75374

Parrott BB, Bowden JA, Kohno S, et al. Influence of tissue, age, and environmental quality on

DNA methylation in Alligator mississippiensis. *Reproduction*. 2014;147(4):503-513.

Published 2014 Mar 2. doi:10.1530/REP-13-0498

Peng C, Cardenas A, Rifas-Shiman SL, et al. Epigenetic age acceleration is associated with allergy

and asthma in children in Project Viva. *J Allergy Clin Immunol*. 2019;143(6):2263-

2270.e14. doi:10.1016/j.jaci.2019.01.034

Petkovich DA, Podolskiy DI, Lobanov AV, Lee SG, Miller RA, Gladyshev VN. Using DNA

Methylation Profiling to Evaluate Biological Age and Longevity Interventions. *Cell Metab*.

2017;25(4):954-960.e6. doi:10.1016/j.cmet.2017.03.016

Pinho GM, Martin JGA, Farrell C, et al. Hibernation slows epigenetic ageing in yellow-bellied

marmots. *Nat Ecol Evol*. 2022;6(4):418-426. doi:10.1038/s41559-022-01679-1

Polanowski AM, Robbins J, Chandler D, Jarman SN. Epigenetic estimation of age in humpback

whales. *Mol Ecol Resour*. 2014;14(5):976-987. doi:10.1111/1755-0998.12247

Powell RA, Proulx G. Trapping and marking terrestrial mammals for research: integrating ethics,

performance criteria, techniques, and common sense. *ILAR J.* 2003;44(4):259-276.

doi:10.1093/ilar.44.4.259

Prado NA, Brown JL, Zoller JA, et al. Epigenetic clock and methylation studies in elephants.

ageing Cell. 2021;20(7):e13414. doi:10.1111/acel.13414

R Core Team (2023). _R: A Language and Environment for Statistical Computing_. R Foundation

for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.

Richetto J, Massart R, Weber-Stadlbauer U, Szyf M, Riva MA, Meyer U. Genome-wide DNA

Methylation Changes in a Mouse Model of Infection-Mediated Neurodevelopmental

Disorders. *Biol Psychiatry*. 2017;81(3):265-276. doi:10.1016/j.biopsych.2016.08.010

Robeck TR, Fei Z, Lu AT, et al. Multi-species and multi-tissue methylation clocks for age

estimation in toothed whales and dolphins. *Commun Biol*. 2021;4(1):642. Published 2021

May 31. doi:10.1038/s42003-021-02179-x

Spangenberg E, Wallenbeck A, Eklöf AC, Carlstedt-Duke J, Tjäder S. Housing breeding mice in

three different IVC systems: maternal performance and pup development. *Lab Anim*.

2014;48(3):193-206. doi:10.1177/0023677214531569

Stubbs TM, Bonder MJ, Stark AK, et al. Multi-tissue DNA methylation age predictor in mouse.

Genome Biol. 2017;18(1):68. Published 2017 Apr 11. doi:10.1186/s13059-017-1203-5

Sullivan IR, Adams DM, Greville LJS, Faure PA, Wilkinson GS. Big brown bats experience

slower epigenetic ageing during hibernation. *Proc Biol Sci*. 2022;289(1980):20220635.

doi:10.1098/rspb.2022.0635

Tangili M, Slettenhaar AJ, Sudyka J, et al. DNA methylation markers of age(ing) in non-model

animals. *Mol Ecol*. 2023;32(17):4725-4741. doi:10.1111/mec.17065

Thompson MJ, vonHoldt B, Horvath S, Pellegrini M. An epigenetic ageing clock for dogs and

wolves. *ageing (Albany NY)*. 2017;9(3):1055-1068. doi:10.18632/ageing.101211

Tobi EW, Lumey LH, Talens RP, et al. DNA methylation differences after exposure to prenatal

famine are common and timing- and sex-specific. *Hum Mol Genet*. 2009;18(21):4046-

4053. doi:10.1093/hmg/ddp353

Vangeel EB, Pishva E, Hompes T, et al. Newborn genome-wide DNA methylation in association

with pregnancy anxiety reveals a potential role for *GABBR1*. *Clin Epigenetics*. 2017;9:107.

Published 2017 Oct 3. doi:10.1186/s13148-017-0408-5

Viitaniemi HM, Verhagen I, Visser ME, Honkela A, van Oers K, Husby A. Seasonal Variation in

Genome-Wide DNA Methylation Patterns and the Onset of Seasonal Timing of

Reproduction in Great Tits. *Genome Biol Evol*. 2019;11(3):970-983.

doi:10.1093/gbe/evz044

Wilkinson GS, Adams DM, Haghani A, et al. DNA methylation predicts age and provides insight

into exceptional longevity of bats [published correction appears in Nat Commun. 2021

May 5;12(1):2652] [published correction appears in Nat Commun. 2022 Sep

7;13(1):5266]. *Nat Commun*. 2021;12(1):1615. Published 2021 Mar 12.

doi:10.1038/s41467-021-21900-2

Wright PGR, Mathews F, Schofield H, et al. Application of a novel molecular method to age free-

living wild Bechstein's bats. *Mol Ecol Resour*. 2018;18(6):1374-1380. doi:10.1111/1755-

0998.12925

Yousefzadeh, M., Henpita, C., Vyas, R., Soto-Palma, C., Robbins, P., & Niedernhofer, L. (2021).

DNA   damage-how   and   why   we   age?.   *eLife*,   *10*,   e62852.

https://doi.org/10.7554/eLife.62852

Zocher S, Overall RW, Lesche M, Dahl A, Kempermann G. Environmental enrichment preserves

a young DNA methylation landscape in the aged mouse hippocampus. *Nat Commun*.

2021;12(1):3892. Published 2021 Jun 23. doi:10.1038/s41467-021-23993-1