Coupled ribosome allocation and nutrient sensing lead to incoherent feedback and oscillatory response in bacterial growth.

Rossana Droghetti,[1] Valerio Firmano,[2] Giorgio Tallarico,[1] Ludovico Calabrese,[1] Jacopo Grilli,[3] and Marco Cosentino Lagomarsino[1,2,4,*]

[1]*IFOM Foundation, FIRC Institute for Molecular Oncology, via Adamello 16, Milan, Italy*

[2]*Dipartimento di Fisica, Università degli Studi di Milano, via Celoria 16, Milan, Italy*

[3]*ICTP - Strada Costiera, 11 - 34151 Trieste Italy*

[4]*INFN sezione di Milano, via Celoria 16, Milan, Italy*

(Dated: October 25, 2023)

Abstract

Current theories describing bacterial growth physiology across environments have demonstrated an impressive predictive power, but they are typically phenomenological. Incorporating mechanistic details into these frameworks remains an open challenge that would greatly improve our ability to predict and control bacterial growth in varying environmental conditions. For example, the "Flux Controlled Regulation" (FCR) model is a reference out-of-equilibrium framework that links ribosome allocation to translation efficiency by means of a steady-state assumption. By making use of this assumption, this model does not account for ppGpp-mediated nutrient sensing and transcriptional regulation of ribosomal operons. In this study, we propose a simple model that integrates the FCR framework with a mechanistic description of three key components: (i) the amino-acid pool, (ii) ppGpp sensing of translation elongation rate, and (iii) transcriptional programming of protein allocation strategy by ppGpp-sensitive promoters. Our framework is fully coherent with observed steady-state growth laws and makes testable predictions for unobserved quantities. Furthermore, our theory predicts that under environmental changes the incoherent feedback between sensing and regulation leads to oscillatory relaxation towards new equilibria, a feature observed experimentally but not captured by previous phenomenological models.

SIGNIFICANCE STATEMENT

Existing theories that explain cellular growth by considering resource allocation do not provide mechanistic details connecting the nutrient precursors pools, the sensing process responsible for allocation and the allocation process itself. Instead, these theories view growth as an outcome of either flux balance or resource optimization, neglecting finer mechanistic aspects. An outstanding challenge is to describe mechanistically how resource allocation at the proteome level is implemented by regulatory circuits that can sense the environment and regulate protein synthesis accordingly. This study focuses on *E. coli* bacteria and proposes a theoretical framework based on contemporary formalisms that includes the description of the sensing and the regulation of protein synthesis at the mRNA level. Intriguingly, the interplay between sensing and regulatory architecture gives rise to oscillatory adaptation to environmental changes, which emerges in experiments but remains not well characterized.

---

* marco.cosentino-lagomarsino@ifom.eu

I. INTRODUCTION

The regulation of growth is critical for all living cells [1–4]. On general grounds, it can be seen on different levels as an optimization of catabolic fluxes and biosynthesis [5], a global resource allocation problem whereby enzymes and proteins that optimize the fluxes are produced [5], and the coordination of sensing of nutrients and other environmental cues leading to the regulatory circuits [6, 7]. The resource allocation problem determines the optimal levels of protein expression that maximize the growth rate (or more in general the fitness) in a given and possibly fluctuating environment [5, 8, 9]. For example, under carbon-limited growth, there is a trade-off between the expression of ribosomes, which carry out protein biosynthesis, and the metabolic enzymes providing the necessary amino acids and other precursors [5, 10–12]. The theories that originate from this observation work are conceptually powerful and quantitatively predictive both during exponential growth [10] and out-of-steady-state scenarios [13].

On the other hand, the current frameworks miss an explicit description of the regulatory circuits that coordinate the cellular response to perturbations by reading the environmental signals.

In bacteria, the circuit that implements this control is based on the small signaling molecule (p)ppGpp (guanosine tetraphosphate or pentaphosphate) [6, 7]. This small molecule can "read" the external environment by sensing changes in the uncharged tRNA caused by changes in amino-acid abundances or other essential nutrients. Under scarcity of these components, uncharged tRNA molecules accumulate in the ribosome, leading to the activation of the ribosome-associated RelA protein, which synthesizes ppGpp by transferring a pyrophosphate group from ATP to GTP or GDP, resulting in the production of ppGpp and AMP or ADP [14, 15]. The less-well-characterized SpoT protein can both degrade and catalyze the synthesis of ppGpp, which provides a mechanism for fine-tuning the cellular response to stress. Recent quantitative measurements of ppGpp levels during nutrient shifts lead to the hypothesis that ppGpp amounts may sense translation elongation speed through the concerted action of RelA and SpoT [16]. The sensing of amino acid levels by ppGpp results in the regulation of ribosomal biosynthesis through its function as a signaling molecule that modulates the activity of RNA polymerase on ribosomal and growth-related promoters. This process involves the DksA protein and a GC-rich "discriminator region" present in the promoter sequences and affects the relative amount of transcripts whose promoter has or does not have the discriminator region [7]. As ppGpp levels increase, the production of ribosomal transcripts decreases, enabling bacteria to adjust to nutrient and stress conditions by redirecting resources towards survival and growth [6, 17, 18].

Being able to predict how cells will respond to new perturbations is crucial, and to this end

a mechanistic understanding of growth control is essential. It is important to note that the response to a change is specific to the environment or perturbation under investigation [12], while the regulatory mechanism remains the same irrespective of the environment. However, obtaining a detailed description of the key circuits controlling resource allocation and growth is challenging as we still lack a comprehensive understanding of all the molecular players. A crucial aspect for gaining insight into the underlying sensing and regulation of growth is that during steady-state (i.e., balanced exponential) growth all the relevant molecular players are balanced [4, 19]. In such conditions, even if resource allocation is the result of the action of sensing and regulatory circuits, the mechanistic principles, and causal chains governing these links remain hidden. Thus, to understand the regulatory aspects, it is necessary to study the dynamic cellular response to perturbations and focus on the out-of-steady-state behavior. From a modeling perspective, non-steady conditions offer the opportunity to describe jointly growth laws, limiting components, and the role of the external nutrients and cues on cellular growth [9, 13, 20, 21]. Here we focus on perturbations performed by changing the external nutrient source [13, 22]. A number of modeling studies have focused on non-steady conditions, and they can be divided into models that incorporate the growth law theory by using a top-down approach [13, 23, 24], and more detailed descriptions of the regulatory mechanisms [9, 17, 20]. Recently, Wu and coworkers [16] have included the elongation-rate sensing mechanism in a top-down framework. However, this model does not include a description of the connection between the environment and the translational speed, which is mediated by the pool of amino acids available for protein synthesis.

In this study, we propose an intermediate approach between a top-down framework and a specific model of the circuits by introducing a comprehensive model that incorporates (I) an explicit description of amino acid sensing, (II) a detailed account of the mechanistic regulation of transcription via ppGpp, and (III) a framework for growth laws and global resource allocation. Notably, our model manifests an emergent property whereby the system's response to external perturbations exhibits oscillatory behavior, which arises due to the incoherent mutual feedback loop that emerges between amino acid pools and ribosome levels.

FIG. 1: **Mechanistic model for bacteria response to external perturbations.** Global scheme of the regulatory model. The out-of-steady-state dynamics is governed by four main ingredients: (1) the allocation functions χ_is, which set the target size for the proteome sectors ϕ_is (2) the ppGpp-mediated transcript dynamics (total transcript T and ribosomal transcripts T_R), which determine the values of the allocation functions by setting the composition of the transcript pool (3) the ppGpp dynamics, which reads the translation elongation rate ϵ and regulates the transcript production (4) the production and consumption of amino acids, ψ_A, which control ppGpp production by setting the translation speed. The first module on the dynamics of the sector size is derived from ref. [13].

II. RESULTS

A hybrid mechanistic-phenomenological framework for understanding bacterial response to nutrient changes

In order to describe the out-of-steady-state dynamics of cell growth, we designed a model framework that, starting from the framework defined in ref. [13] takes into account all the major mechanistic players involved in the response to the internal amino acids concentration, translation, nutrient sensing by ppGpp concentration, mRNA dynamics, and protein production (see Methods and SI Appendix for details). Fig. 1 shows the scheme of the proposed framework, which we describe in the following paragraphs.

In the model, ribosomes are responsible for the synthesis of all proteins, divided into "sectors" representing groups of co-regulated proteins (which usually have similar functions) [13]. Our

model focuses on three sectors: ribosomal (ϕ_R), catabolic (ϕ_P, assumed to be connected with the flux of amino acids [10]), and housekeeping (ϕ_Q, which is assumed to be kept homeostatically constant [10]). Out of steady state, the target size of each sector is regulated by "regulatory functions", denoted as χ_R, χ_P, and χ_Q, representing the fraction of ribosomes that actively translate proteins of the different sectors. The requirement that ribosomes make all proteins leads to the following equations [13]

$$\frac{\mathrm{d}\phi_i}{\mathrm{d}t} = \lambda[\chi_i - \phi_i]; i \in [R, P, Q] \ , \tag{1}$$

where λ is the growth rate (Fig. 1).

In the absence of post-transcriptional control [6, 25], we assume that ribosomes randomly attach to transcripts, initiate protein synthesis, and produce the proteins. Therefore, the composition of the mRNA pool determines the redistribution of the flux among the three sectors. We hence define χ_R as the ratio of ribosomal transcripts (T_R) to the total number of transcripts (T), and similarly for χ_P and χ_Q.

The ribosomal mRNA pool composition is regulated by the alarmone ppGpp, which controls the partitioning of the RNA polymerase (RNAP) [7]. Consequently, the number of ribosomal transcripts, T_R, depends on the concentration of ppGpp, denoted as G, via the partition of the RNA polymerases, denoted by ω_R. We can derive an equation for χ_R after an environmental nutrient change by combining our definition of χ_R with transcript dynamics. The resulting equation is given by

$$\frac{\mathrm{d}\chi_R}{\mathrm{d}t} = \frac{1}{\tau_\chi}[\omega_R(G) - \chi_R] \ , \tag{2}$$

where τ_χ is a time scale that includes ppGpp level adjustment to the new condition and transcript pool adjustment, which we estimate it to be around 10 min (see Fig. 1, Methods and SI Appendix). In our model the partitioning function of the RNA polymerases ω_R depends solely on the ppGpp pool, and its functional form derives from a fit of the steady state data (see SI material and SI Fig. 1). This assumption is based on the recent study by Balakrishnan and coworkers [25], who found that the mRNA pool composition across conditions is mostly determined by the specific gene on-rates and only depends weakly on other factors such as gene dosage or mRNA degradation. This is especially true for the proteins belonging to the ribosomal sector, whose genes on-rates are governed by ppGpp. An extension of our theory including other dependencies is straightforward.

Amino acid levels and translation rate are determined in our model by the interplay of nutrient uptake, biosynthesis, and volume growth (Fig. 1). Specifically, the model describes the abundance of one compound amino acid species by the equation

$$\frac{\mathrm{d}\psi_{\mathrm{A}}}{\mathrm{d}t} = \nu\phi_{\mathrm{P}} - \epsilon\phi_{\mathrm{R}} - \lambda\psi_{\mathrm{A}} \ . \tag{3}$$

Eq. (3) introduces the normalized amino acid mass ψ_{A}, which is the ratio of amino acid mass (A) to total protein mass (M_{tot}). The catabolic flux linking nutrients to amino acids is represented by the term $\nu\phi_{\mathrm{P}}$, where ν denotes the nutrient quality (an average catabolic flux per employed catabolic sector protein), and the biosynthesis flux is represented by $\epsilon\phi_{\mathrm{R}}$. The last term accounts for the dilution effect caused by volume growth, which we chose to not neglect, as this term has an impact on the time scales of the relaxation dynamics (see SI Fig. S3).

The framework describes the full connection between amino-acid levels, ppGpp pool, and transcriptional reallocation

A key aspect of our model is the explicit representation of the amino acid pool (which represents a significant departure from ref. [13]). This ingredient plays a vital role in the sensing mechanism of ppGpp, which governs the cellular response to perturbations. Indeed, following a shift in nutrient availability, the first change observed by the cell is in the metabolic flux $\nu\phi_{\mathrm{P}}$, due to alterations in nutrient quality (ν). Consequently, these changes in nutrient availability lead to variations in the levels of amino acids, which have a direct influence on tRNA charging and the translation rate. In turn, the altered translation rate induces changes in the level of ppGpp (G), thereby triggering a transcriptional reconfiguration of the cell's allocation strategy.

Let us further explain the relationship between the translation elongation rate ϵ and ψ_{A}: based on experimental findings (ref. [26]), we express the translation elongation rate (ϵ) as the following function of the concentration of charged tRNAs:

$$\epsilon = \tilde{\epsilon}\frac{[tRNA^{\mathrm{C}}]}{[tRNA^{\mathrm{C}}] + K_{\mathrm{C}}} \ , \tag{4}$$

Where ($\tilde{\epsilon}$) is the theoretical maximum value of the elongation rate. The concentration of charged tRNAs is, in turn, influenced by the pool of available amino acids, which affects the dynamics of tRNA charging. We assume a simple relationship between uncharged tRNAs and cognate amino acids: $[tRNA^{\mathrm{C}}] \propto \psi_{\mathrm{A}}$, i.e. that the fast time-scale changes of precursors are instantaneously mirrored by tRNA charging. For a detailed motivation of this assumption please refer to the SI Appendix.

The (simplified but) explicit description of charged tRNA sensing in our model allows us to make quantitative predictions regarding the size of the amino acid pool and its relationships with other variables (Fig. 2). We found that, regardless of the growth conditions, the model predicts that in any biologically accessible regime, the system reaches a steady state of balanced

growth (the fixed point solution is described in the next section and in SI Appendix) where $\psi_A \simeq k_A$, thereby regulating itself to avoid limiting amino acids and an excessively high fraction of loaded tRNA, which therefore are buffered in a self-organized way. The allocation of resources within the proteome regulates the influx of nutrients and efflux of amino acids resulting from biosynthesis. This regulation is influenced by the dynamic interplay of the ppGpp-mediated transcriptome. In order to connect ppGpp levels to the amino-acid pool, we use the model proposed by Wu and coworkers [16], who established that ppGpp level (G) is related to charged tRNA levels in a way that ppGpp is effectively a function of translation elongation rate (ϵ) through the equation

$$G = \frac{\tilde{\epsilon}}{\epsilon} - 1 \; . \tag{5}$$

It is important to note that so far ϵ solely represents the translation elongation rate. In the presence of stalled ribosomes (e.g., under chloramphenicol treatment [26, 27]) this is distinct from the average ribosomal activity (called σ in ref [13]). The relationship between G and tRNA charging is explained in terms of the typical times elongating ribosomes spend in elongation (τ_{trans}) and waiting for a charged tRNA (τ_{dwell}), both quantities that are captured by the translation elongation rate ϵ, see Fig. 1 and ref. [16]. Note that the model by Wu and coworkers is in contrast with previous models that assumed that ppGpp concentration depends on both ribosomal sector size and amino acid levels [9, 16, 17, 20].

The model reproduces the available steady-state data between growth speed and resource allocation

Two aspects of our model are noteworthy. Firstly, the ability to reproduce the known steady-state relationships between measured quantities is a crucial requirement, and our model satisfies this criterion. This is partly due to the original FCR framework from which our model is derived. However, our model incorporates additional observables, such as ppGpp and amino acid levels. ppGpp levels have been measured and are in agreement with our model. Furthermore, the model's ability to predict the amino acid pool levels is an outcome that can be tested by new experiments, making it a testable prediction.

Fig. 2 shows the steady state results of the simulations of the model, compared with the ppGpp pool experimental data presented in ref [16]. The simulations in Fig. 2 start from an arbitrary initial condition and collect the steady-state values of the main state variables once the system has reached equilibrium. Panels B and D of the figure show the dependency between the growth rate, the amino-acid pool ψ_A, ppGpp concentration G, while panel C shows the

FIG. 2: **The proposed framework predicts steady-state relationships between the amino acids pool, ppGpp levels and ribosomal allocation, and reproduces the available experimental data. A:** Model prediction of the dependence of the ribosomal sector fractional size (y-axis) on the exponential growth rate (x-axis). Simulations (red circles) are compared with experimental data from ref. [16] (blue reverse triangles) and ref. [26] (light blue crosses). In our framework such "first growth law" is not used in the definition of the regulatory circuit, and comes as a genuine prediction of the model. **B:** The plot shows the simulated model prediction (red circles) for the relationship between the steady-state exponential growth rate (x axis) and the size (mass fraction ψ_A) of the amino-acid pool (y axis). This quantity is proportional to the concentration of charged tRNA ([tRNA^C]), as explained in the text and in SI. **C:** Model prediction (red circles) for the relationship between the size of the amino-acid pool (x axis) and the relative translation rate ($\epsilon/\tilde{\epsilon}$, y axis) in steady-state growth. **D:** The model prediction (red circles) for the relationship between the steady-state exponential growth rate (x axis) and the ppGpp concentration (y axis) is in line with the experimental data (blue reverse triangles, data from ref. [16] out of the slow-growth regime, see SI Fig. S2). **E:** The model prediction (red circles) for the relationship between the ppGpp concentration (x axis) and the size of the ribosomal sector (y axis) agrees with the experimental data (blue reverse triangles, data from ref. [16]). For this figure, we restricted ourselves to the fast-growth regime ($\lambda > 0.5 h^{-1}$). For the slow growth regime, where additional phenomena such as degradation and inactive ribosomes also impact physiology, please refer to SI Fig. S2.

dependence of the translation elongation rate ϵ on the amino-acid pool and panel D shows the relationship between ribosomal allocation ϕ_R and ppGpp levels. Additionally, as we anticipated above Fig. 2 illustrates that the model can accurately reproduce available steady-state data (from ref. [16]) for ppGpp. We note that, unlike in other frameworks (including the FCR model), the ribosomal growth law is not an input of our model, therefore the trend shown in Fig. 2A is a genuine prediction.

In order to avoid unnecessary complications, we restricted this analysis to a "fast-growth" regime, which we defined by the condition $\lambda > 0.5h^{-1}$. Indeed, it is well known that in slow-growth conditions, other phenomena such as protein degradation and inactive ribosomes play a significant role in the growth physiology [26, 27]. The model discussed in the main text of this work does not account for these phenomena and therefore refers to the fast-growth regime. We have also studied an extended version of our framework that includes the essential ingredients to describe the slow-growth behavior. For a description of this version of the model and its steady-state predictions please refer to SI Appendix and SI Fig. S2.

Finally, a well-defined steady-state behavior requires the fixed point to be stable, which we investigated by analyzing the eigenvalues of the linearized dynamics. Specifically, we have studied in detail two versions of the fast-growth model, with instantaneous transcription and with a transcriptional time scale set by the mRNA degradation rate (see methods and, SI Fig. S6 and SI Appendix). In both cases, all eigenvalues have a negative real part in the physiologically relevant parameter region, showing that the steady-growth fixed points are always stable. More in general, we also show that in the absence of transcriptional delays, a generic model for ribosome translation regulation will display stable fixed points, as long as the regulatory function is monotonically decreasing in ϕ_R and increasing in ψ_A (details and demonstration in SI Appendix).

The relaxation dynamics towards the new steady state shows damped oscillations that do not depend on the transcriptional time scales

We next investigated the model behavior during a nutritional upshift. To realize such an upshift, we varied suddenly (in a stepwise fashion) the nutrient quality ν, which is the parameter that characterizes the external environment. Fig. 3 shows the predicted relaxation behavior, which presents damped oscillations.

We asked whether the presence of oscillations was linked to the transcriptional delay τ_χ, which sets the time scale for the adaptation of the mRNA pool composition after the ppGpp has adjusted to the new post-shift level. Fig. 4 shows that both the instantaneous-transcription ($\tau_\chi =$

FIG. 3: **Damped oscillations characterize the relaxation dynamics towards the new steady state.** The figure presents simulation predictions of the post-upshift dynamics of various state variables. Panels **A**, **B**, **C**, and **D** display the time-dependent behavior of the relative translation elongation rate $\epsilon/\tilde{\epsilon}$, the mass fraction of the ribosomal sector ϕ_R, the growth rate λ, and the ppGpp concentration G, respectively. As $\lambda = \epsilon\phi_R$, its dynamics are more intricate than those of the sector size and elongation speed. Panel **E** shows the post-shift dynamics of the ribosomal sector ϕ_R in relation to the amino acids pool ψ_A and the relative translation elongation rate $\epsilon/\tilde{\epsilon}$, with circles representing simulation results color-coded by time from the shift. The X-shaped symbols denote the pre- and post-shift steady-state values and the dashed line represents the steady-state relationship between the two plotted variables. Similarly, panel **F** shows the post-shift dynamics of the ppGpp concentration G versus ψ_A and $\tilde{\epsilon}/\epsilon - 1$, a factor which is found proportional to ppGpp [16]. In this panel, we also show shift data from [16] (blue reverse triangles). These plots further emphasize the presence of oscillatory relaxation dynamics.

0) model and the full model ($\tau_\chi > 0$) present oscillations during the shift, highlighted Fig. 4AB.

The analysis of the eigenvalues of the two systems (SI Fig. S6) confirms the presence of

damped oscillations in both the instantaneous and non-instantaneous transcriptional response

model variants. Indeed in both cases, the eigenvalues become complex with negative real part if

the nutrient quality ν is larger than a certain threshold ν^* (see SI Appendix for details on ν^*). By

studying the parametric dependence of ν^* we also find that the onset of this damped oscillatory

dynamics is not just determined by the resource allocation strategy, but the dynamics of the

amino acid pool also plays a role (see SI Appendix, sec. 9). More in detail, our analysis of the

eigenvalues of the system reveals three distinct regimes that depend on the parameters and the

growth rate (SI Fig.s S6-S8). The first regime is characterized by a shift without oscillations,

occurring at slow growth rates. The second regime features a shift with damped oscillations,

which occurs in the fast-growth regime. Finally, the third regime is a shift with sustained

FIG. 4: **The incoherent feedback between amino acid pool and ribosome production gives rise to oscillatory response regardless of delays due to ribosome transcription.** Simulations were performed with two different models: with instantaneous transcription (**A-C**) and with gradual mRNA and rRNA pool production (**B-D**). In the instantaneous transcription model, the regulatory functions χ_i follow ppGpp levels without delay (as in refs. [13, 16]), while they are given by Eq. 2 in the gradual transcription model. In all panels, circles indicate simulation results color-coded by the time from the shift, and X-shaped symbols represent the pre- and post-shift steady states. Panels **A** and **B** show the three phases of post-shift adaptation. The x axis displays the ribosomal sector mass fraction, and the y axis shows the growth rate. The dashed line in both panels connects $(0,0)$ and $(\phi_{\mathrm{R}}^{\mathrm{final}}, \lambda^{\mathrm{final}})$, and highlights the second phase (as in Fig. 9 of ref. [13]). Panel **C** and **D** show the ribosome allocation strategy through the shift. The x axis displays the relative translation rate, and the y axis shows the ribosomal sector regulatory function χ_{R}. The plots show that oscillations occur even when the regulatory function instantaneously follows the translation elongation rate along the steady-state relation. For both models, the relaxation is composed of the three highlighted phases. Panels E-F-G compare simulations of three different models proposed to describe nutrient shift dynamics: this work (red circles), the FCR model (ref. [13] (brown plus signs), and the Flux-Parity Model (FPM) (ref. [24] (pink crosses). Panel **E** shows the dynamics of the instantaneous growth rate for the three models, with our model being the only one that predicts an overshoot of the growth rate. Panel **F** shows the size of the amino acid pool ψ_{A}, which is not predicted by the FCR model, and with our model and the FPM model making different predictions, even during steady state. Panel **G** shows the ppGpp concentration G across the shift. This can be predicted with all three models, combining the FCR framework with Eq. 5, from ref. [16] (see SI Appendix for the upgraded FCR model definition). The discrepancies between the steady-state predictions are due to the different values of the parameters used. Panel **H** provides a sketch of the incoherent feedback loop between the amino acid pool ψ_{A} and the ribosomal fraction ϕ_{R}, which is explicitly described in this study and is responsible for the observed oscillatory behavior.

oscillations. The theoretical possibility of such "oscillatory growth" has been predicted by a previous generic growth model [28]. The threshold between these regimes varies depending on the parameters of the system (SI Appendix and SI Fig. S6-S8). However, we find that for experimentally derived parameters, damped oscillations should be observable in a wide range of growth rates, indeed SI Fig. S6 shows that the eigenvalues of the system present a non-zero imaginary part for every growth rate larger than approximately $0.5h^{-1}$.

In order to interpret the different relaxation time scales involved in a nutrient shift within our model, Fig. 4AB shows that the process of reaching a new steady state after a change in nutrient quality can be divided into three phases. In the first phase, there is a sudden increase in the translation elongation rate due to the increase in the amino acid pool size ψ_A. This change in the growth rate is detected by the sensory mechanism of the ppGpp, but the synthesis of mRNA and new proteins has not yet changed. Therefore, the first phase is characterized by a change in the growth rate without a change in the ribosomal sector size. In the second phase, the synthesis of new proteins starts to adjust to the new protein allocation strategy given by $\chi_R(G)$. This second phase is slower than the first one because the synthesis of new proteins is not immediate. During this phase, the translation elongation rate remains almost constant while the sector sizes change, and the relaxation proceeds along a straight line in the (ϕ_R, λ) plane. In the third and last phase, all the relevant variables, including translational activity, regulatory functions, and protein sectors, oscillate while relaxing around the new steady state. Our study provides a detailed analysis of these three phases and their underlying mechanisms. The study by Erickson and coworkers that introduced the FCR model (ref. [13]) already presented some of the different relaxation phases we identified in our study. However, their model only included the first and second phases and did not account for the oscillatory dynamics observed in our study (see ref. [13]).

Damped oscillations are not observed in models not accounting for the connection between amino-acid pool sensing and ribosome allocation.

To shed more light on the nature of the oscillatory behavior in our model, we further compared our framework with two other models, the original FCR model and the Flux Parity Model (FPM, ref. [24]), which were designed to describe nutritional upshifts (see SI Appendix and references for details). The FCR model is based on a phenomenological assumption that relates directly to the translation elongation rate and proteome sectors regulatory functions. This is similar to our model variant without transcriptional delay. However, Fig. 4CD shows that oscillations occur in our model even when the regulatory function instantaneously follows the translation elongation

rate, while they are not present in this version of the FCR model. Note that here we have integrated the FCR model with the description of the ppGpp circuit, in order to compare the prediction on ppGpp dynamics of the two models (see SI material and SI Fig. S5). The crucial difference is that our model directly connects the translation elongation rate with the amino acids level, and describes the dynamics of the amino acids pool and its interaction with the ribosomal proteins explicitly. The FPM model includes the sensing of charged tRNAs However, its predictions are distinct from both the FCR and our own framework, in that it assumes an optimization principle for defining the control of biosynthesis. Specifically, the resource allocation is designed to always maximize the growth rate. The authors argue that this strategy is compatible with the ppGpp regulation circuit, at least qualitatively, but they do not provide an explicit model for this part, breaking the feedback loop that in our model leads to oscillatory behavior. Consequently, Fig. 4E-F shows that the three models have different predictions, and only our model predicts the oscillatory relaxation toward the new steady state, because is the only one that explicitly describes the relation between amino acids pool and ribosomal allocation. To back up this point, in the SI we also show that a version of our model with a different ppGpp regulation (incorporating solely the RelA production term) also shows damped oscillation (see SI section 5 and SI Fig. S4).

Shift behavior compatible with damped oscillations has been reported experimentally for various quantities, albeit with different time scales

So far, we have shown that the oscillatory relaxation behavior arises naturally from a theoretical framework that incorporates biological knowledge on nutrient sensing and regulatory response. It is natural to ask whether this behavior has any experimental counterpart. While data from the experimental literature appear to support the emergence of damped oscillations during nutritional upshifts, the time scales of these oscillations vary significantly across different studies (SI Fig. 12). We observe clear oscillations in data from a nutritional upshift performed in a microfluidic device, from ref. [22] (Fig. 5). These damped oscillations are apparent in both the growth rate λ and the ribosomal sector ϕ_{R}. Note that to obtain data on the ribosomal sector, the authors of ref. [22] have used the signal of a GFP attached to a ribosomal promoter. Before using this signal as a proxy, we conducted several tests on the signal to see if it was compatible with the sector dynamics behavior (see SI material and SI Fig. 9-10 for details). However, the time scale of the experimentally observed oscillations differs from the one naturally predicted by our model. In order to fit the data in Fig. 5, we had to adjust the value of k_{A} (the Michaelis-Menten constant of $\epsilon(\psi_{\mathrm{A}})$, Eq. 9) to around 2, which appears quite large. This

Shifts in microfluidic device [Panlilio et al. 2021]

FIG. 5: **Oscillations in nutrient shift experiments. A-B:** growth rate λ and ribosomal sector size ϕ_R dynamics following an upshift in a microfluidic device (reanalyzed data from ref. [22]). The circles are sliding averages placed every 4min, with a window of 75min (see Methods and/or SI Appendix for details on the averaging procedures and the definition of ϕ_R). The red line represents the results of the simulation, where we set the value of k_A to 5 in order to fit the data. The data show oscillatory relaxation to the new steady state, both for the growth rate and the ribosomal sector size, and that the model can account for this behavior.

may be due to experimental features of such microfluidic nutrient upshifts. Indeed, a minute time-scale growth-rate overshoot is observed in such an experiment (SI Fig. 11). The same fast overshoot was also previously reported in similar single-cell shifts providing amino acids [29], and connected by direct observation of cell volume and mass to a dilution of the cell. Hence, cells entering an upshift effectively feel a hypo-osmotic shock. It is possible that relevant short time-scale biophysical processes, and consequent physiological adaptation not described by our model bias the effective fit parameters. Additionally, our *a priori* estimation of the parameter k_A is based on a simple consideration of the amino acid pool and may not be accurate enough to reproduce experimental data.

Therefore, we did not attempt to provide a precise fit or parameter estimation with our model from experimental data, and our main claims remain limited to the statement that a realistic architecture linking nutrient sensing to ribosome allocation would give rise to oscillations. This may be connected to experimental observations, but it is also possible that missing elements in the architecture that are not described by our model could change the prediction, and further systematic experiments are needed to clarify the situation.

III. DISCUSSION AND CONCLUSION

This work presents a dynamical modeling framework that describes the out-of-steady-state growth of *E. coli* cells through a global resource allocation dynamics. The model combines the framework of ref. [13] with a mechanistic description of nutrient sensing and global gene regulation by the ppGpp circuit. This combination of ingredients leads to predict an oscillatory

relaxation after a shift in the nutrient conditions, for all shifts that lead to a moderate-to-fast growth rate. While other models predict oscillatory behavior in this context [17, 20], recent work on growth laws in this area did not incorporate this aspect [13, 16]. Crucially, we have shown that removing any transcriptional delay in our model does not ablate the incoherent feedback loop, nor the oscillations. Hence, we conclude that the transcriptional waiting time *per se* should not be regarded as a cause for the oscillations. Our framework indeed connects the presence of oscillations to the feedback interplay between global flux balance and resource allocation and the mechanistic circuits that implement growth control. In other words, we show that the oscillatory behavior is caused by the incoherent feedback between the size of the amino acid pool and the expression of the ribosomal sector.

Importantly, we have shown that damped oscillations of growth rate, ribosome allocation and ribosome proteome fraction have been observed across studies of experimental nutrient shifts in *E. coli*. Interestingly, similar oscillations in response to perturbations have also been observed in yeast [30], where the mechanistic architecture of nutrient sensing and growth regulation is very different [2, 31–33]. This suggests that the global feedback described by our model may be a general feature, or strategy, for growth regulation based on nutrient sensing. However, there are still many unanswered questions regarding these experiments and their relationship with our model. While the mechanism provided by our model provides a possible explanation, it is possible that the oscillations observed experimentally are also influenced by other molecular players that are not considered in our study. Indeed, the experimental oscillations observed during the experiments exhibit a wide range of time scales, ranging from a few minutes to several hours, which in our model would result in an unrealistically wide range of fitting parameters across studies. To quantitatively match the experimental data, in future studies it will be necessary to conduct additional experiments under controlled conditions, measure all relevant quantities carefully, and adjust the model according to the specific characteristics of each experiment.

In our model, the oscillations arise because the system works as a thermostat, with a built-in feedback system involving a sensor and an effector. The origin of oscillations in our system could be attributed to architectural constraints, evolution, or a combination of both. In order to clarify this point, it will be crucial to study this system from a control-theory perspective, and to discover the optimization goals that it follows over evolution. A previous study [34] taking this approach, has concluded that oscillatory behavior may be due to optimal control towards the adjustment of ribosome synthesis in a switch-like manner. The authors demonstrate that a precursor-only control strategy, which alternates investment between gene expression and metabolism producing an oscillatory time profile of precursor concentration performs much

better than a nutrient-only strategy in a dynamical upshift scenario, by avoiding the inefficient transient accumulation of precursors.

A growth regulation system can be seen as a decision-making process that detects the environment. As nutrients in a natural environment tend to fluctuate over different time scales, we can expect that the nutrient quality can vary by fluctuations and net trends. Our model shows that sensing as a low-pass filter in response to time-varying input, preventing the system from reacting to environmental nutrient quality changes above a certain frequency, but also carries a characteristic oscillatory frequency that varies with the parameters. We speculate that these features could facilitate growth control, allowing a cell to spare energy and resources by avoiding reactions to changes that are too short-lived, but also "resonate" with specific frequencies. In our framework, the model variant without delay represents the fastest-reacting version of the network. Biologically, this fast-reacting architecture may be embodied by ribosome sequestration [35], at the cost of the extra proteome sector occupied by inactive ribosomes [26, 27]. We speculate that by studying the long-term fitness trade-offs of a system that can bypass the transcriptional delay and comparing it to one that does not one could address the question of whether *E. coli* hedges its bets on both architectures through regulated ribosome sequestration, in order to set its threshold frequency in a plastic fashion, and optimize its fitness flexibly.

IV. MATERIALS AND METHODS

Model description

We report here the equations needed for the definition and simulation of our mechanistic model. In these notes, we will distinguish the state variables from the parameters by expliciting the time dependence of the first ones. We start with the differential equations that define the dynamical system:

$$\frac{\mathrm{d}\psi_\mathrm{A}(t)}{\mathrm{d}t} = \nu\phi_\mathrm{P}(t) - \epsilon(t)\phi_\mathrm{R}(t) - \lambda(t)\psi_\mathrm{A}(t), \tag{6}$$

$$\frac{\mathrm{d}\chi_\mathrm{R}(t)}{\mathrm{d}t} = \frac{1}{\tau_\chi}[\chi_\mathrm{R}(t) - \omega_\mathrm{R}(t)], \tag{7}$$

$$\frac{\mathrm{d}\phi_\mathrm{R}(t)}{\mathrm{d}t} = \lambda(t)[\chi_\mathrm{R}(t) - \phi_\mathrm{R}(t)]. \tag{8}$$

Eq. 6 describes the dynamics of the amino-acid pool, given by the balance of the upcoming nutrient flux ($\nu\phi_\mathrm{P}$), the outgoing biosynthesis flux ($\epsilon\phi_\mathrm{R}$), and a dilution term given by volume growth ($\lambda\psi_\mathrm{A}$). Eq. 7 is derived from the transcript dynamics, which is explained in detail in the SI Appendix. Eq. 8 is derived from the sector definition as in ref. [13].

Next, we show the other definitions needed to close the system. We need to define the

translation elongation rate, which depends on the amount of charged tRNAs, and therefore on the amino acids level. Eq. 4 connects these quantities:

$$\epsilon(t) = \tilde{\epsilon}\frac{\psi_{\mathrm{A}}(t)}{\psi_{\mathrm{A}}(t) + k_{\mathrm{A}}} \ . \tag{9}$$

The ppGpp concentration is given by the following empirical relation from ref. [16]

$$G(t) = \mathrm{C}G^{\mathrm{ref}}\left(\frac{\tilde{\epsilon}}{\epsilon(t)} - 1\right) \ . \tag{10}$$

The equation presented in ref. [16] gives the fold change of the ppGpp with respect to a reference condition, therefore, in order to obtain the ppGpp concentration we add the parameter G^{ref} to the original equation for the fold change. The value for G^{ref} is $101.5\mu M$ and is given by ref. [17], see SI Appendix for details.

The RNAP allocation on ribosomal genes was assumed to follow the following relationship

$$\omega_R(t) = \frac{K_{\mathrm{G}}}{K_{\mathrm{G}} + G(t)} \ , \tag{11}$$

where K_{G} is given from a fit of the data presented in ref.s [16, 17], and its value is $14.5\mu M$ (see SI Appendix for details and a variant of this model).

The growth rate in our model corresponds to the biosynthesis rate:

$$\lambda(t) = \epsilon(t)\phi_{\mathrm{R}}(t). \tag{12}$$

Lastly, the constitutive sector is defined as the remaining part of the proteome, which does not belong to the ribosomal nor to the housekeeping one,

$$\phi_{\mathrm{P}}(t) = 1 - \phi_{\mathrm{Q}} - \phi_{\mathrm{R}}(t) = \phi_{\mathrm{R,max}} - \phi_{\mathrm{R}}(t) \ . \tag{13}$$

Data Analysis

All the information on the analysis of the data from ref [22], which are shown in Fig. 5, are reported in the SI material, section 11 and SI Fig. S9-S11. Other experimental data that show oscillations are reported in SI section 12 and SI Fig. S12.

Data availability

The code utilized in this work can be found in the repository at DOI: 10.17632/w294vd3pgh.1

CONFLICT OF INTEREST STATEMENT

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

AUTHOR CONTRIBUTIONS

MCL conceived the study. RD performed all the simulations with assistance from LC. RD, JG, and MCL designed the models. RD took care of the model simulations. RD took care of the data analysis with assistance from GT, VF. MCL, JG, and RD wrote the paper.

FUNDING

This work was supported by Associazione Italiana per la Ricerca sul Cancro AIRC IG grant no. 23258 (MCL and RD). RD was also supported by an AIRC fellowship (Italy Pre-Doc 2022 - ID 28176).

ACKNOWLEDGMENTS

We are very grateful to all the members of the Cosentino Lagomarsino lab for the useful discussions. We would like to thank our experimental collaborators Pietro Cicuta and Aske Petersen for the feedback on the data. We are grateful to Gregory Bokinsky, Frank Bruggeman, Luca Ciandrini, Ulrich Gerland, Gabriele Micali, Sander Tans, and Terence Hwa for the feedback on this work. We also thank Severin J. Schink for the useful feedback and for having shared with us the data of ref. [13].

 SI APPENDIX

435 This SI Appendix explains in more detail the model introduced in the main text, illustrates

436 new versions of this framework and the differences in terms of predictions. It also includes

437 the analytical calculations of the fixed points and the eigenvalues of the system, and lastly, it

438 describes how we made use of the data reported in ref. [22] to compare with our analytical

439 findings.

V. BRIEF DESCRIPTION OF THE MODEL.

To ensure clarity, we begin by providing a synthetic overview of all the components of the

model. The following presents a detailed description of the model equations. Our framework

starts from the existing "flux-control regulation" (FCR) model [13] and integrates it with an

explicit description of three different regulatory modules related to ppGpp dynamics. Fig. 1 of

the main text shows the four ingredients of the proposed framework, which we describe in the

following paragraphs. Throughout this SI Appendix, we distinguish the system state variables

from the system parameters by making explicit the time dependence of the formers.

a. Global allocation functions set the proteome sectors sizes. As in the FCR model, in our

model some "global allocation" functions determine the sizes of the proteome sectors. These

allocation functions, denoted as $\chi_i(t)$, set the fraction of the total biosynthesis flux that is

redirected toward a specific sector, thereby determining the dynamics of the sector sizes ($\phi_i(t)$).

Our model focuses on three sectors: ribosomal ($\phi_R(t)$), constitutive ($\phi_P(t)$), and housekeeping

($\phi_Q(t)$), with $\chi_R(t)$, $\chi_P(t)$, and $\chi_Q(t)$ determining the number of ribosomes making ribosomes,

constitutive proteins, and housekeeping proteins, respectively. The time course of the sector size

is described by the same equation as in the FCR model [13]:

$$\frac{\mathrm{d}\phi_i(t)}{\mathrm{d}t} = \lambda(t)[\chi_i(t) - \phi_i(t)]; i \in [\mathrm{R}, \mathrm{P}, \mathrm{Q}]. \tag{S1}$$

Here, $\lambda(t)$ represents the growth rate (i.e. the biosynthesis rate), and the term $\chi_i(t) - \phi_i(t)$

specifies the rate of change of the sector size.

b. ppGpp-mediated transcripts dynamics determine the proteome allocation functions. Our

model proposes that the allocation of the total biosynthesis flux is determined by the composition

of the mRNA pool, which is controlled, for the ribosomal sector, by ppGpp-mediated transcript

dynamics. Without post-transcriptional control [6, 25], ribosomes randomly attach to transcripts

and initiate protein synthesis. Therefore, the mRNA pool composition determines the partition

of the flux towards the three proteome sectors. We define χ_R as the ratio of ribosomal to

total transcripts ($\frac{T_R}{T}$), and χ_P and χ_Q in a similar way. The dynamics of ribosomal transcripts (T_R) is governed by the ppGpp circuit that interacts with RNA polymerase (RNAP) and this dependence is encoded in the regulatory function

$$\omega_R(G(t)) = \frac{K_G}{K_G + G(t)} \ , \tag{S2}$$

where ω_R represents the fraction of RNAP that transcribes ribosomal genes. The function for ω_R is a fit of experimental data presented in ref.s [16, 17] (SI Fig. S1). The dynamics of ribosomal transcripts is given by

$$\frac{dT_R(t)}{dt} = \beta \omega_R(G(t))P - \delta T_R(t) \ , \tag{S3}$$

where β and δ are the mRNA production and degradation rates (representing the average behavior of many genes), respectively, and P is the number of available RNAPs. Analogously, total transcripts follow the equation

$$\frac{dT(t)}{dt} = \beta P - \delta T(t) \ , \tag{S4}$$

The differential equation that governs χ_R derives from its definition and Eqs. (S3) and (S4)

$$\frac{d\chi_R(t)}{dt} = \frac{1}{\tau_\chi}[\omega_R(G(t)) - \chi_R(t)] \ , \tag{S5}$$

where $\tau_\chi \simeq 1/\delta$ is the time scale separating the (fast) adjustment of the ppGpp level to the new condition and the consequent adjustment of the transcript pool (on the order of minutes, we typically set $\tau_\chi = 10\text{min}$) [36].

    c.   Translation elongation speed determines ppGpp production and degradation. A recent study by Wu et al. [16] found a clear relationship between ppGpp level and translation elongation rate (ϵ),

$$G(t) \propto \frac{\tilde{\epsilon}}{\epsilon(t)} - 1, \tag{S6}$$

where $\tilde{\epsilon}$ is the theoretical maximum translation elongation rate. This relationship can be explained by a model where the rate of ppGpp synthesis depends on the time ribosomes spend elongating (τ_{trans}), while the rate of degradation is proportional to the time spent waiting for a charged tRNA (τ_{dwell}) [16]. This finding invalidates previous models [9, 17, 20] that only described ppGpp synthesis by RelA based on loading of uncharged tRNAs, and suggests ppGpp degradation by SpoT plays a more complex role than previously assumed.

Because of the rapid production and degradation rates of ppGpp (on the order of seconds) [20], we assume that ppGpp concentration is always at a quasi-steady state. Moreover, the data from Wu et al. [16] showed that this relationship between ϵ and G holds even during out-of-steady-state scenarios. The translation elongation speed ϵ is related to the fraction of charged tRNA, thus the last module that closes the circle is the dynamics of the free amino acids denoted as ψ_A.

   d. Nutrient influx, biosynthesis outflux, and growth govern the amino acid levels and the

translation rate.   The mass of amino acids A in the cell is determined by the equilibrium between nutrient influx and outflux due to biosynthesis [5]

$$\frac{\mathrm{d}A(t)}{\mathrm{d}t} = \nu M_\mathrm{P}(t) - \epsilon(t)M_\mathrm{R}(t) \ , \tag{S7}$$

where ν is a "nutrient quality" parameter representing a mass-specific catabolic flux [10, 13], and M_P and M_R are the masses of the constitutive and ribosomal proteins, respectively. To obtain an equation with the proteome sectors, we define $\psi_\mathrm{A} = A/M_\mathrm{tot}$, where M_tot is the mass of total proteins. If we assume a constant dry-mass density [13, 19], we obtain the following equation for ψ_A,

$$\frac{\mathrm{d}\psi_\mathrm{A}(t)}{\mathrm{d}t} = \nu\phi_\mathrm{P}(t) - \epsilon(t)\phi_\mathrm{R}(t) - \lambda(t)\psi_\mathrm{A}(t) \ , \tag{S8}$$

where the additional term $\lambda\psi_\mathrm{A}$ accounts for dilution due to cellular growth. Here, ϕ_P and ϕ_R denote the fractions of the proteome allocated to the constitutive and ribosomal proteins, respectively. The influx of nutrients and the outflux of amino acids due to biosynthesis are determined by the proteome allocation, which in turn is governed by the dynamics of the ppGpp-mediated transcriptome, as described earlier. The translation elongation rate, ϵ, is also influenced by the amino-acid levels via the availability of charged tRNAs, as discussed earlier.

VI.   MODEL DESCRIPTION - IN DEPTH.

This section presents all the equations required for defining and simulating our model. These equations are also presented in the main text; here we provide a comprehensive description and discuss their validity ranges.

A.   Dynamics of amino acid pools are set by catabolic flux, protein production and dilution due to volume growth

We start with the equation that describes the dynamics of the amino acids pool valid outside of balanced-growth conditions,

$$\frac{\mathrm{d}\psi_\mathrm{A}(t)}{\mathrm{d}t} = \nu\phi_\mathrm{P}(t) - \epsilon(t)\phi_\mathrm{R}(t) - \lambda(t)\psi_\mathrm{A}(t), \tag{S9}$$

where $\psi_\mathrm{A} = A/M_{tot}$ is the rescaled amino acids mass as defined in the main text. Although there are twenty different amino acid species, for simplicity we have considered one quantity to describe the amino acid pool. We interpret this as describing the limiting species. The terms on the right-hand side of Eq. (S9) above represent (I) the inflow of amino acids resulting

from the catabolic flux $J_A(t) = \nu\phi_P(t)$, (II) the biosynthesis flux $J_b(t) = \epsilon(t)\phi_R(t)$, and (III) a dilution term due to volume growth $-\lambda(t)\psi_A(t)$. The inflow of amino acids is a result of various processes such as nutrient uptake, the conversion of nutrients into central precursors, and the synthesis of specific amino acids. The nature of these processes varies based on the specific nutrients provided to the cells, for instance, whether they consist of solely carbon sources or a nutrient-rich medium containing amino acids. In our model, we have chosen to provide a high-level description of these processes, combining them into a single flux. The nutrient uptake and amino-acid flux is mediated by proteins that do not belong to the ribosomal sector ϕ_R [10] nor to the housekeeping sector ϕ_Q, justifying the assumption that the flux depends on the constitutive/catabolic sector ϕ_P.

Previous studies have examined the growth-dependent expression of these proteins, particularly those involved in carbon source uptake [11, 37]. This regulation has been incorporated into the FCR framework describing shifts between different carbon sources [13]. While our model derives from the FCR framework, we have chosen for simplicity a less detailed and more generic version of this regulation. As uptake proteome sectors are part of the unregulated constitutive sector ϕ_P, our model includes only this information. While our model may sacrifice some details of the regulation described in references [11, 37], it is not limited to describing carbon sources alone. Consequently, it avoids certain constraints, such as the maximum growth rate for growth on carbon-only nutrients $\lambda_C = 1.17h^{-1}$ set in ref. [13]. Additionally, to our knowledge, there is no quantitative analysis of substrate uptake in rich media containing amino acids. In any case, this modeling choice is not restrictive, and it is possible to adopt a more detailed description without affecting our main results.

The other main factor setting the amino-acid flux is the nutrient quality ν, which represents the sum of the nutrient qualities of every individual substrate present in the growth medium. Once again, for simplicity, we chose not to describe a specific uptake sector for each substrate component in our model as in ref. [13], but this extension is simple to explore. We can still group all nutrient qualities into a single global parameter interpreting it as follows. Each nutrient quality ν_i can be expressed as $\nu_i = \nu_i' f_i$, where ν_i' represents the uptake rate per protein mass [13], and f_i is the fraction of the P sector occupied by the uptake proteins of the i-th substrate, such that $\phi_P f_i = \phi_{up,i}$. Using these definitions, we can express the uptake flux as follows,

$$J_A(t) = \sum_i \nu_i \phi_i(t) = \sum_i \nu_i' \phi_{up,i}(t) \ . \tag{S10}$$

It should be noted that this simplified description of the catabolic flux has certain implications. With this approach, the only relevant change for a nutrient shift is a change in the value of ν. For instance, these restrictions do not make it possible to describe shifts between carbon

sources with very similar nutrient qualities, such as from glucose to lactose. Additionally, when we model an upshift by a stepwise increase in ν, and the catabolic flux $\nu\phi_P$ initially changes due to the change in ν, we implicitly assume that all the proteins responsible for the uptake of the new substrate are already expressed in the pre-shift condition. However, this assumption is not always valid [38]. For a more precise description of the uptake process, we refer to the FCR model [13], which is specifically designed to describe shift data between different carbon sources. Our model can be endowed with the same ingredients if necessary.

Turning to the biosynthesis flux, we choose to use a description similar to the one reported in ref [13]: $J_b(t) = \epsilon(t)\phi_R(t)$, but in our case $\epsilon(t)$ is the actual translation elongation rate instead of the ribosomal activity (which represents an *average* translation rate per ribosome). Note that, in general, the ribosomal activity can be different from the elongation rate, not only because the ribosome can have different speeds but also because the ribosomal activity takes into account ribosomes that can be inactive or sequestered [13, 16, 26]. Fig. S2 shows that at fast growth the contributions of the sequestered ribosomes (and degradation) can be neglected, therefore in the fast-growth regimes, the ribosomal activity and the elongation rate are approximately the same. We chose to model the translation elongation rate instead of the ribosomal activity because in our model the former is directly connected to the amino-acid availability, as we describe in the next paragraphs. In a variant of the model applicable to slow-growth data (see below), we have considered an additional factor f_{bg} representing the fraction of active ribosomes as well as protein degradation [27]. This version of the model is described in one of the subsequent sections of these notes.

Finally, the last term of Eq. (S9), the dilution term given by growth, appears in the equation because of the definition of the rescaled mass $\psi_A = M_A/M_{prot}$. Since the amino acid pool is very small in terms of concentration, previous studies have neglected this last term, making the assumption of perfect balancing between the metabolic and biosynthesis [5]. However, neglecting this term leads to inconsistencies in the equations for the concentration of amino acids that do not appear acceptable. Indeed, if we neglect the dilution term in Eq.S9, reduces to

$$\frac{\mathrm{d}\psi_A}{\mathrm{dt}} = \nu\phi_P - \epsilon\phi_R \qquad (S11)$$

Therefore the steady-state condition requires $\nu\phi_P = \epsilon\phi_R$. Inserting this condition into Eq. S7 contradicts the exponential accumulation of total amino acids necessary to sustain exponential growth. If we consider the dilution term, we obtain the correct equation for the concentration of amino acids, which includes the dilution term associated with cellular growth. Therefore, even if the amino-acid pool is small, we kept the dilution term in the definition of our model and in our numerical simulations, which also plays a role in how we describe tRNA charging

and elongation [17]. Note that this is not just a mathematical requirement, as neglecting this term has an impact on the relaxation dynamics after a perturbation, which is one of the main phenomena studied in this work, as also pointed out in ref. [24]. Fig. S3 shows that if we neglect the dilution term the system needs a much longer time to reach the new equilibrium.

B. The elongation rate senses the amino acids pool through charged tRNAs

It is widely recognized that the translation elongation rate is closely tied to the availability of amino acids and other precursors, quantities that play a crucial role in the bacteria's stringent response [13, 20, 26].

In our framework, we account for this precursor dependence by relating the translation elongation rate to the concentration of charged tRNAs, similar to what is presented in ref. [26]. Specifically, we define the elongation rate as follows,

$$\epsilon(t) = \tilde{\epsilon} \frac{[tRNA^{\mathrm{C}}]}{[tRNA^{\mathrm{C}}] + K_{\mathrm{C}}} \ . \tag{S12}$$

In this equation, $\tilde{\epsilon}$ represents a theoretical maximum translation elongation rate, which we estimate empirically as

$$\tilde{\epsilon} = \gamma \frac{\phi_{\mathrm{R}}(\tilde{\epsilon}) - \phi_{\mathrm{R}}^{\mathrm{min}}}{\phi_{\mathrm{R}}(\tilde{\epsilon})} \ , \tag{S13}$$

an empirical expression obtained from the condition $\lambda_{\mathrm{max}} = \tilde{\epsilon}\phi_{\mathrm{R}}(\tilde{\epsilon})$ and the first growth law $\phi_{\mathrm{R}} = \phi_{\mathrm{R}}^{\mathrm{min}} + \lambda/\gamma$. For this derivation, we use $\phi_{\mathrm{R}}(\tilde{\epsilon}) = 1$ in order to get the maximum theoretical elongation rate. This is a requirement of our framework. Indeed, by analyzing the expression of the fixed points of the system (see section 8, Eq. S60) one notices that ϵ can reach $\tilde{\epsilon}$ if and only if ϕ_{R} reaches unity. Note that this is just a theoretical requirement, as experimentally the ribosomal sector cannot reach unity because of the presence of the housekeeping sector ϕ_{Q}.

More specifically, we propose a model based on the following key results from ref. [26]: (I) in steady conditions, the ratio of the concentration of tRNA and rRNA is constant across growth rates, leading to approximately 7 tRNAs per ribosome, and maintained by gene regulation (II) the ratio of elongations factors to ribosomes is constant across growth rates (there are approximately 6 EF per ribosome) and lastly, (III) the percentage of charged tRNA is approximately the 70% of all tRNAs, and it is constant across growth conditions. These results suggest that the main factor that limits translation elongation speed is the diffusion of tRNA (charged or uncharged) toward the ribosomes. Indeed, this hypothesis would explain why the elongation speed increases as the growth rate increases, even if the ratio of tRNA to ribosomes remains constant. In a diffusion-limited scenario, the translation-elongation rate is given by the overall

concentration of tRNA, which increases with the growth rate, rather than the amount of tRNA per ribosome, leading to Eq. S12.

In a dynamic situation, for example, a nutrient shift, tRNA charging will change faster than the total amount of tRNA as the amount of available amino acids has increased. Once the cell reads this signal, it can up-regulate the transcription/translation tRNAs and ribosomes. Therefore, in order to make the above equation S12 compatible with the shift dynamics without describing explicitly tRNA regulation, we use the ansatz that the amount of charged tRNAs is proportional to the amino acids concentration: $[tRNA^C] \propto \psi_A$. By this ansatz obtain the relation

$$\epsilon(t) = \tilde{\epsilon} \frac{\psi_A}{\psi_A + k_A} \ . \tag{S14}$$

This equation captures the fast dependency of the translation elongation rate on the availability of amino acids and provides a link between the concentration of charged tRNAs and the translation activity. The constant k_A has not been measured to our knowledge. We set its value to 5×10^{-3}, as it is related to the order of magnitude of the amino acids mass fraction [5].

Finally, we observe that the available data [16] lead to the following empirical relation

$$\epsilon(t) = \tilde{\epsilon} \frac{\phi_R}{\phi_R + \phi_R^0} \ , \tag{S15}$$

suggesting a tight mirror relationship between precursor pools, charged tRNA concentration, and ribosomal sector size in a steady condition.

C. ppGpp senses the translation elongation rate through uncharged tRNAs

In order to model the ppGpp sensing and dynamics, we used the recent results obtained Wu and coworkers [16]. In brief, the authors find that the level of ppGpp is related to the translation elongation rate ϵ via a simple empirical relation

$$G(t) \propto C \left(\frac{\tilde{\epsilon}}{\epsilon(t)} - 1 \right), \tag{S16}$$

where C is a parameter fitted in ref. [16]. We write that G is proportional to the right-hand term of the above equation because Wu and coauthors measure the fold change of ppGpp with respect to a reference condition, which they chose as the exponential growth on glucose. In order to get the ppGpp concentration one needs to multiply the right-hand term of Eq. S16 with the value of G found in glucose, estimated to be $G^{ref} = 41.8858791333739 \mu M$. This estimate is derived from a simulation of our model, based on the parameters derived by Marr in ref. [17] (see SI section 5 for the details). Differently from the study of Wu and coworkers [16], this study

considers the concentration of ppGpp instead of the fold change and allows us to estimate the ppGpp concentration at various growth rates.

Importantly, Wu and coworkers [16]) demonstrate that the relationship between G and ϵ holds during dynamic regimes, beyond steady states. This suggests that the rapid synthesis and degradation dynamics of ppGpp maintain its pool in a steady-state relationship with the translation elongation rate. Consequently, we consider the regime in which the level of ppGpp G remains at a steady-state with ϵ, and hence it is described by an algebraic relationship rather than an ordinary differential equation (ODE).

It is important to stress that G depends on ϵ through the ratio $\psi_A/(\psi_A + k_A)$. In other words, ppGpp concentration is strictly related to the amount of charged tRNAs, which determines the ribosome waiting time τ_{dwell}. In the presence of an elongation-inhibiting drug, the ribosomes split into two populations, those that are stuck by the drug do not elongate and are not linked to ppGpp production or degradation, and the others feel an increased pool of charged tRNA, making the elongation rate (conditionally to where elongation is possible) higher [16].

D. Regulatory functions emerge from transcripts dynamics

In the FCR framework, regulatory functions are crucial in determining the target size for proteome sector production, ultimately leading to a steady-state growth condition, where $\phi_i(t) = \chi_i(t) \forall t$. Mathematically, these functions represent the fraction of the total biosynthetic flux directed towards the i-th sector. However, this definition does not provide insights into the microscopic processes that give rise to the emergence of χ_i. Intuitively, this flux fraction must be set by the fraction of ribosomes that are translating proteins of a given sector.

In the FCR model [13], the authors bypass the problem by using a quasi-steady-state approximation that sets the values of χ_i based on $\chi_i(t) = \phi_i^*(\sigma(t))$, where ϕ_i^* represents the steady-state relationship between ϕ_i and translational activity σ. This approach ensures that even during kinetic regimes following perturbations, the regulatory functions adjust to changes in ribosomal activity in a quasi-steady-state manner (note this formalism defines the translational activity σ as λ/ϕ_R, rather than solely the translation elongation rate ϵ [13], hence σ includes a factor from the fraction of active ribosomes). This assumption is justified by the fact that the allocation of ribosomes should reproduce steady-growth states. However, this top-down approach has a drawback in that it has a limited ability to provide insights into the biological mechanisms governing the regulation of these functions and their underlying definition. Our framework aims at overcoming this limitation by distinguishing translational activity from a translation rate linked to the charged tRNA pool and providing a more mechanistic description of the origin of

 regulatory functions.

 To distribute the biosynthetic flux among different proteome sectors, ribosomes must produce
 proteins in accordance with the ratios determined by the regulatory functions. In our model,
 ribosomes randomly attach to mRNA transcripts and initiate the synthesis of the correspond-
 ing proteins. Therefore, neglecting post-transcriptional regulation [25], the distribution of the
 biosynthesis flux is solely determined by the composition of the transcript pool.

 With this perspective, we can define the regulatory functions as the ratio of the number of
 transcripts coding for proteins belonging to the i-th sector, denoted as $T_i(t)$, to the total number
 of transcripts, denoted as $T(t)$. This is represented mathematically as

$$\chi_i(t) = \frac{T_i(t)}{T(t)}. \tag{S17}$$

 It is important to note that $\Sigma_i T_i(t) = T(t)$, as is the case for the proteome sectors.

 To study the dynamics of χ_i, we need to consider the dynamics of transcripts. Transcripts
 are rapidly degraded and produced by RNA polymerases, denoted as P and assumed to be
 constant. Thus, the dynamics of transcripts can be described by the following equation,

$$\frac{\mathrm{d}T(t)}{\mathrm{d}t} = \beta P - \delta T(t) , \tag{S18}$$

 where β and δ are constants representing the mRNA synthesis and degradation rates, respec-
 tively. This is analogous to the approach presented in ref. [25], where the authors break down
 the dynamics of the mRNA pool by studying the production and degradation flux. Eq. S18 rep-
 resents the dynamics of the total number of transcripts. To study the dynamics of transcripts
 belonging to the i-th sector, we can replace the total RNAP P with the fraction of RNAP
 producing transcripts for the i-th sector, denoted as $\omega_i(t)P$. This leads to the following equation

$$\frac{\mathrm{d}T_i(t)}{\mathrm{d}t} = \beta \omega_i(t)P - \delta T_i(t) . \tag{S19}$$

 By combining Eq. S18 and S19, along with the definition of the regulatory function in Eq. S17,
 we obtain

$$\frac{\mathrm{d}\chi_i(t)}{\mathrm{d}t} = \frac{1}{\tau_\chi}[\omega_i(t) - \chi_i(t)]. \tag{S20}$$

 Here, τ_χ is a constant defined as $\beta P/\delta$, which represents the typical timescale for adjusting the
 composition of the transcripts pool. Given that transcript degradation is much faster than the
 growth rate, we set $\tau_\chi = 10$ minutes [36].

 We chose to focus for simplicity on three sectors, namely ϕ_R, ϕ_P, and ϕ_Q, and the con-
 straint $P + R + Q = 1$ applies to both ϕ and ω, we only need to specify the ppGpp dependence
 for the RNAP partition function of ribosomal transcripts, ω_R.

A further aspect of the model we need to address is the allocation of RNA polymerases (RNAPs), specifically in relation to ribosomal transcripts. In *E. coli*, it is known that the stringent response, mediated by ppGpp, regulates the production of ribosomal transcripts by directly influencing the allocation of RNAPs (see ref.s[6, 7] for details). We have left implicit the role of gene dosage in this regulation. Ribosomal protein genes and ribosomal RNAs are found near the replication origin, which affects their copy number across growth conditions [39]. However the recent findings by Balakrishnan and coworkers [25] suggest that the mRNA pool composition is mainly determined by the regulation of the promoter on-rates rather than other factors such as gene dosage and mRNA degradation rates. Therefore, while it is possible to study the role of these other factors by incorporating them in model variants, we have assumed for simplicity that the functions ω_i just depend on promoter on-rates, which in our theory are just set by ppGpp in the case of the genes belonging to the ribosomal sectors (although one could also add catabolic regulators in a straightforward way).

We describe the allocation of RNAPs to ribosomal genes by the following equation,

$$\omega_R(t) = \frac{K_{\mathrm{G}}}{K_{\mathrm{G}} + G(t)}, \tag{S21}$$

where K_{G} is determined by fitting the data presented in ref.s [16, 17] and replotted in Fig. S1, with a fitted value of $14.5\mu M$.

We highlight the fact that alternative choices are possible for Eq. S21. For example, one can define a model variant where the constraint on ppGpp transcriptional regulation given by a constant Q sector, $\phi_{\mathrm{R}} < \phi_{\mathrm{R}}^{\mathrm{max}}$, is made explicit, by defining

$$\omega_{\mathrm{R}} = \phi_{\mathrm{R}}^{\mathrm{max}} \frac{K_{\mathrm{G}}}{K_{\mathrm{G}} + G(t)}. \tag{S22}$$

In this model variant, the equations at steady states become

$$\epsilon = \tilde{\epsilon} \frac{\psi_{\mathrm{A}}}{\psi_{\mathrm{A}} + k_{\mathrm{A}}} \, , \tag{S23}$$

$$G = G^{\mathrm{ref}} \left(\frac{\tilde{\epsilon}}{\epsilon} - 1 \right) \, , \tag{S24}$$

$$\lambda = \epsilon \phi_R \, , \tag{S25}$$

$$\phi_R = \phi_{max} \frac{K_{\mathrm{G}}}{K_{\mathrm{G}} + G(t)} \, . \tag{S26}$$

Hence, G affects the ribosomal and RNAP sectors in a more symmetric way. Since we fit the empirical relation $\phi_{\mathrm{R}}(G)$ in order to define ω_{R}, our model appears to be sufficiently good for our purposes. Indeed, we verified that none of the main results of this study are affected by this variant. However, the model variant with bounded ω_{R} leads to more consistent calculations and good agreement with data, and may be preferred in future studies.

728 The definition we have used for χ_i relies on a set of simplifying assumptions that may not

729 hold in some conditions. Specifically, the following requirements need to be fulfilled:

- For an accurate representation of ppGpp levels in transcript dynamics, both T and T_R
730

731 need to be sufficiently large. Otherwise, the ratio $\frac{T_R}{T}$ becomes an approximation of the

732 true allocation strategy.

- The number of ribosomes to be larger than the number of transcripts. This is required
733

734 because $\frac{T_R}{T}$ is the probability of a ribosome hitting a ribosomal transcript, and for the

735 law of large numbers, this probability is accurately achieved just for a sufficient amount

736 of ribosomes.

- Conversely, the number of ribosomes should not be excessively large, as our model does
737

not account for ribosomal traffic.

In summary, we have $\chi_R = \frac{T_R}{T}$ only when the following conditions hold: $LT \gg R \gg T$, where L

represents the maximum number of ribosomes that can be allocated to a transcript, and T

must be sufficiently large to accurately capture ppGpp changes. In other regimes, more general

definitions χ_R can be used [40], keeping the rest of the model ingredients similar.

Finally, it is important to emphasize that our framework exclusively focuses on ribosomal

proteins, leaving ribosomal RNA implicit. Given the tightly coordinated production of ribosomal

proteins and rRNA [41–43], this approximation appears reasonable. However, it is crucial to

highlight that in our framework, ω_R (representing the fraction of RNAP involved in transcribing

ribosomal protein transcripts) is computed by considering only the RNA polymerases engaged in

the transcription of protein transcripts, excluding those involved in rRNA synthesis. We chose

this approach due to the analogy between RNAP sectors and protein sectors, which enables

us to equate ω_i with ϕ_i during steady state. Nonetheless, it is worth noting that other studies

also incorporate rRNA explicitly [44–46], hence it is in principle feasible to study this model

extension, which would make sense in order to study conditions or perturbations where the

balance assumed here is or can be lost. Similar considerations hold for representing tRNA

biosynthesis in the model [46].

E. The dynamics of protein sectors are determined by their regulatory function and
755

756 **the growth rate**

757 The derivation of sector dynamics in our model is straightforward and follows from their

758 definition, precisely as for the FCR model [13]. Here are the main steps of the derivation. Each

sector is characterized by the mass ratio of the proteins belonging to the sector to the total protein mass, $\phi_i = M_i/M_{tot}$. The increment of the total mass is governed by biosynthesis

$$\frac{dM_{tot}(t)}{dt} = \epsilon(t)M_R(t) \ , \tag{S27}$$

while the increment of each sector mass fraction corresponds to a fraction of the total increment, as follows,

$$\frac{dM_i(t)}{dt} = \chi_i(t)\epsilon(t)M_R(t). \tag{S28}$$

Combining these equations leads to the differential equation describing the dynamics of each sector

$$\frac{d\phi_i(t)}{dt} = \lambda(t)[\chi_i(t) - \phi_i(t)] \ , \tag{S29}$$

where λ, by definition, is given by $\frac{1}{M_{tot}}\frac{dM_{tot}}{dt} = \epsilon(t)\phi_R(t)$.

In this study, we focus on three sectors: the ribosomal sector ϕ_R, the constitutive sector ϕ_P, and the housekeeping sector ϕ_Q. As ϕ_Q is assumed to be constant, the other two sectors must satisfy the constraint $\phi_P(t) = 1 - \phi_Q - \phi_R(t) = \phi_{R,max} - \phi_R(t)$ at each time t. This constraint also applies to the regulatory functions χ_R and χ_P, as well as the RNAP partition functions ω_R and ω_P. This constraint holds for all these quantities simply because of mass conservation. Thus, we can avoid including ω_P and χ_P in the simulation and use the relation between ϕ_R and ϕ_P to obtain the dynamics of the latter.

VII. MODEL VERSION WITH INACTIVE RIBOSOMES AND DEGRADATION.

As mentioned in the main text, the model described so far is not adapted to describe slow-growth behavior due to the significance of protein degradation and inactive ribosomes in cellular physiology [26, 27]. Therefore, to describe this regime, we need to incorporate these two ingredients into the model. To achieve this, we follow ref. [27] and introduce two key elements: the fraction of ribosomes contributing to growth, denoted as f_{bg}, and the degradation rate, denoted as η. Both quantities depend on the steady-state growth rate or, equivalently, on the nutrient quality ν.

It is known that the fraction of active ribosomes decreases at slow growth rates and that when the growth rate is zero, f_{bg} must also be zero by definition. We first define the fraction of ribosomes contributing to growth as

$$f_{bg}(\nu) = \frac{\nu}{\nu + \nu_{bg}}. \tag{S30}$$

By this choice, following ref. [27], we propose a Michaelis-Menten dependence on nutrient quality, with ν_{bg} as a parameter fitted to reproduce experimental slow-growth data on ϵ and ϕ_R. It is

important to note that currently there is no available data on f_{bg}, and the above equation represents an empirical approximation for this quantity.

Concerning the protein degradation rate, we have utilized the available data in *E. coli* [27, 47] to fit a hyperbolic function as follows,

$$\eta(\nu) = \frac{\eta_0 + \eta_\infty s\nu}{1 + s\nu} \ , \tag{S31}$$

where η_0 and η_∞ represent the protein degradation rate values for $\lambda = 0$ and $\lambda = \infty$, respectively, and s controls the steepness of the hyperbolic function.

Once these two mechanisms are defined, we need to integrate them into the system model equations which now become:

$$\frac{\mathrm{d}\psi_{\mathrm{A}}(t)}{\mathrm{d}t} = \nu\phi_{\mathrm{P}}(t) - \epsilon(t)\phi_{\mathrm{R}}(t)f_{\mathrm{bg}}(\nu)(1 + \psi_{\mathrm{A}}(t)) + \eta(\nu)\psi_{\mathrm{A}}(t), \tag{S32}$$

$$\frac{\mathrm{d}\chi_{\mathrm{R}}(t)}{\mathrm{d}t} = \frac{1}{\tau_\chi}[\omega_{\mathrm{R}}(t) - \chi_{\mathrm{R}}(t)], \tag{S33}$$

$$\frac{\mathrm{d}\phi_{\mathrm{R}}(t)}{\mathrm{d}t} = \epsilon(t)\phi_{\mathrm{R}}(t)f_{\mathrm{bg}}(\nu)[\phi_{\mathrm{R}}(t) - \chi_{\mathrm{R}}(t)]. \tag{S34}$$

$$\lambda(t) = \epsilon(t)\phi_{\mathrm{R}}(t)f_{\mathrm{bg}}(\nu) - \eta(\nu). \tag{S35}$$

It should be noted that the structure of the equations remains the same, but in Eq. S32, an additional positive term representing protein degradation is included. Furthermore, both Eq. S32 and S34 multiply the ribosome sector by f_{bg}, which is a consequence of the revised definition of the growth rate presented in Eq. S35, incorporating both active ribosomes and the degradation term. The other equations in the system remain unchanged. Note that the definitions provided for f_{bg} and η are suitable just for studying steady-state behavior. During a shift, one would expect these two quantities, along with other relevant variables, to exhibit oscillatory behavior. However, our current definition predicts an immediate shift to the new post-shift equilibrium value. This is a consequence of the dependence on nutrient quality ν, which changes in a stepwise fashion.

Figure S2 shows the predictions of this modified model. Specifically, the plots demonstrate that by incorporating the mechanisms of inactive ribosomes and degradation, the model can accurately reproduce the data even during the slow growth regime. A previous study [27] has already pointed out that degradation is necessary in order to explain the phenomenology of slow growth because the degradation (and the subsequent maintenance synthesis) is required to explain why at zero growth rate one still has measure a non-zero elongation rate [16, 26]. At the light of the current model, we can add that the maintenance activity of the ribosomes is also required to explain the observed ppGpp pool. Indeed, if we assume that the production

and degradation of ppGpp is related to ribosomal activity, as claimed in ref. [16], we need to acknowledge that at zero growth rate the fraction of active ribosomes cannot go to zero, which means that the ribosomes cannot be all sequestered. If this were the case, the translation elongation rate ϵ would go to zero in our model and the ppGpp level, according to Eq. S16, would diverge, which clearly appears as a non-physical solution.

VIII. MODEL VERSION WITH INSTANTANEOUS TRANSCRIPTION.

This section presents a variant of the model that neglects the transcriptional delay due to ppGpp-induced changes in ribosome transcription, which we used to verify if the predicted oscillatory behavior arises solely due to the delay introduced by the rearrangement of the transcripts.

In the absence of transcription delays, the regulatory function χ_R directly depends on the ppGpp levels, resulting in the following equation:

$$\chi_R(t) = \omega_R(t) = \frac{K_G}{K_G + G(t)} \ . \tag{S36}$$

Under this assumption, the system is reduced to just two ordinary differential equations

$$\frac{d\psi_A(t)}{dt} = \nu\phi_P(t) - \epsilon(t)\phi_R(t) - \lambda(t)\psi_A(t), \tag{S37}$$

$$\frac{d\phi_R(t)}{dt} = \lambda(t)[\omega_R(t) - \phi_R(t)]. \tag{S38}$$

All the other equations in the model remain unchanged.

IX. MODEL VERSION WITH PPGPP SYNTHESIS ONLY FROM RELA.

Before the direct measurement of the relationship between ppGpp levels and the translation elongation rate ϵ (Eq. S16) in ref. [16], the standard approach to describing ppGpp dynamics relied on the elongation-coupled ppGpp synthesis by the enzyme RelA, for which there is solid experimental evidence [18]. Specifically, ppGpp is synthesized by RelA using ATP and GTP as substrates in the presence of uncharged tRNAs. When uncharged tRNAs are loaded onto transcribing ribosomes, the ribosome becomes stalled on the transcript and RelA detects this and attaches to the ribosome, producing (p)ppGpp before detaching. This mechanism links ppGpp production to the state of central precursors, in particular, RelA can sense amino acid starvation (which influences the level of uncharged tRNAs) [6]. In contrast, less is known about how SpoT synthesizes ppGpp and how it senses stressful conditions. It appears that SpoT can be involved in the accumulation of ppGpp under various stressful conditions and nutrient

limitations, not necessarily related to amino acid starvation [6, 39]. Therefore, this pathway is typically modeled simply as a degradation term [9, 20].

This information about RelA and the assumption that SpoT works at a constant rate leads to the following equation for ppGpp [17, 20]

$$\frac{\mathrm{d}G}{\mathrm{dt}} = k_{\mathrm{RelA}} f_{\mathrm{c}} \phi_{\mathrm{R}}(1 - P_{\mathrm{C}}) - k_{\mathrm{spoT}} G, \tag{S39}$$

where the first term on the right-hand side describes the synthesis by RelA and the second term represents the degradation by SpoT. We observe that the RelA term depends on both the amount of ribosomes ϕ_{R} and the probability of having a charged tRNA $P_{\mathrm{C}} = \frac{\psi_{\mathrm{A}}}{\psi_{\mathrm{A}} + k_{\mathrm{A}}}$, while k_{RelA} and f_{c} are the synthesis constant and the conversion factor between the number of ribosomes and the sector size, respectively (their values can be found in Table S1). Assuming a quasi-steady-state description of ppGpp, the equation reduces to

$$G = \frac{k_{\mathrm{RelA}}}{k_{\mathrm{spoT}}} f_{\mathrm{c}} \phi_{\mathrm{R}}(1 - P_{\mathrm{C}}) \, , \tag{S40}$$

which differs significantly from the equation used here and by Wu and colleagues [16]. This fundamental difference falsifies this model, as shown in Fig.S4D, and suggests that the role of SpoT cannot be reduced to a simple degradation term (although the exact mechanism of SpoT is not yet known). Moreover, in this model, the relation between the ppGpp concentration and ribosome allocation is given by

$$\chi_{\mathrm{R}} = \frac{\tilde{K}_{\mathrm{G}}^2}{\tilde{K}_{\mathrm{G}}^2 + G^2} \tag{S41}$$

following the definition used in ref. [20] (see table 1 for the value of $\tilde{K}_{\mathrm{G}}$).

While the specific quantitative details may vary, it is worth mentioning that even when incorporating Marr's proposed model into our framework, the oscillatory behavior still emerges. Hence our main results appear to be more generic than these specific aspects of the model. As discussed in the main text, as long as there is a feedback network between amino acids and ribosomes, an oscillatory regime is expected, as shown in Fig. S4E-H.

X. VARIANT OF THE FCR MODEL INCORPORATING A DESCRIPTION OF PPGPP.

In the main text, we presented the predicted shift behavior based on the FCR model [13]. Since the original version of the FCR model does not incorporate the dynamics of ppGpp, in order to get a prediction for the ppGpp levels within this framework we defined an extended version of the FCR model by integrating the findings from Wu and coauthors [16] to account for the ppGpp description.

Before introducing the equations that define the extended model, it is important to highlight a few key points. In the FCR model, the parameter σ represents the ribosomal activity, defined as λ/ϕ_R, which is different from the translation elongation rate ϵ. This distinction becomes significant here because the model for ppGpp, as presented in ref. [16], also introduces sequestered ribosomes, denoted as $\phi_\text{R,seq}$, which are ribosomes that are synthesized but do not contribute to growth. Given that the peculiarity of this model is to define the sectors regulatory functions χ_i as function of σ, it becomes important to introduce both σ and ϵ, which is not strictly necessary in our framework.

Let us proceed by presenting the main equations used in our extended FCR model with a brief explanation, given that most of them are already presented in ref. [13]. We start with the growth rate, which is determined by the nutrient uptake fluxes:

$$\lambda(t) = \mu_1 \phi_\text{cat1}(t) + \mu_2 \phi_\text{cat2}(t) \ , \tag{S42}$$

where ϕ_cat 1 and 2 are the uptake sectors for the first and the second carbon source present in the substrate, and μ 1 and 2 are their nutrient qualities. From the growth rate we define the ribosomal activity

$$\sigma(t) = \frac{\lambda(t)}{\phi_\text{R}(t)} \ . \tag{S43}$$

Next, we want to define the translation elongation rate. To do so we introduce the fraction of sequestered ribosomes

$$f_\text{seq}(t) = k_\text{conv} \left(\frac{1 - \frac{\sigma(t)}{\gamma}}{\phi_\text{R}^\text{min}} \right)^2 \ . \tag{S44}$$

With this definition, we define the amount of sequestered ribosomes

$$\phi_\text{R,seq}(t) = \phi_\text{R}(t) f_\text{seq}(t) \ , \tag{S45}$$

and the translation elongation rate, following the definition in ref. [16]

$$\epsilon(t) = \frac{\lambda(t)}{\phi_\text{R}(t) - \phi_\text{R,seq}(t)}. \tag{S46}$$

The ppGpp dynamics follows the rules defined by Wu and coauthors, as in our model,

$$G(t) = C G^\text{ref} \left(\frac{\tilde{\epsilon}}{\epsilon(t)} - 1 \right). \tag{S47}$$

The two regulatory functions are given by steady-state relations [13]

$$\chi_\text{R}(t) = \chi_\text{R}^*(\sigma(t)), \tag{S48}$$

$$\chi_\text{cat}(t) = \chi_\text{cat}^*(\sigma(t)), \tag{S49}$$

and, as usual, the dynamics of the sectors are determined by the regulatory functions

$$\frac{d\phi_R(t)}{dt} = \lambda(t)[\chi_R(t) - \phi_R(t)], \tag{S50}$$

$$\frac{d\phi_{cat}(t)}{dt} = \lambda(t)[\chi_{cat}(t) - \phi_{cat}(t)]. \tag{S51}$$

Most of the equations presented here are directly taken from the FCR [13], with the addition of the ppGpp equation from Wu et al. [16]. However, we have introduced two new equations that describe the behavior of sequestered ribosomes during a shift, namely Eq.s S44 and S45. We have defined this extension in order to make testable predictions for ppGpp during shifts. While the authors of ref. [16] do not explicitly describe how sequestered ribosomes behave during a shift and only state that the sequestering process is independent of protein synthesis, we propose an ansatz to maintain the steady-state relationship between ribosomal activity σ and translation elongation rate ϵ as described in ref. [16],

$$\sigma(t) = \epsilon(t)\frac{\phi_R(t) - \phi_{R,seq}(t)}{\phi_R(t)}. \tag{S52}$$

During a shift, σ changes faster than ϕ_R, and in order to maintain the correspondence between σ and ϵ, it is necessary for the sequestered ribosomes to change as fast as σ does. To account for this, we model f_{seq} as a dynamic function of σ. Let us explain how we describe the sequestered ribosomes. According to ref. [16], the sequestered ribosomes constitute a fraction of the total number of ribosomes at steady-state. Using the equations $\phi_R = \left(\frac{A\eta_{R/P,\phi}}{\eta_{N,R/P}}\right)/G$ and $\phi_{R,seq} = \frac{B\eta_{R/P,\phi}}{\eta_{N,R/P}}G$, where G is the ppGpp level the rest are constants fitted from the growth law (see ref. [16] for details), we can write:

$$\phi_{R,seq} = \frac{k_{conv}}{\phi_R}, \tag{S53}$$

where $k_{conv} = \left(\frac{\eta_{R/P,\phi}}{\eta_{N,R/P}}\right)^2 AB$. We want to express the sequestered ribosomes as $\phi_{R,seq} = \phi_R f_{seq}$. By substituting the above equation, we obtain

$$f_{seq}(\sigma) = \frac{k_{conv}}{\phi_R^2(\sigma)} = k_{conv}\left(\frac{1 - \frac{\sigma}{\gamma}}{\phi_R^{min}}\right)^2, \tag{S54}$$

where we have explicitly included the dependence on ribosomal activity σ by using the first growth law [13].

We have defined this updated version of the FCR framework in order to compare its prediction with the ones of our model. The prediction and the dynamics of the updated FCR model are presented in Fig. S5, which shows that this model reproduces the observed dynamics, for both up and downshifts, as well as making predictions for additional observables. However, due to the steady-state assumptions intrinsic in the FCR model, it does not produce any damped oscillation in response to a nutrient change.

XI. COMPARISON WITH OTHER MODELS.

This section compares our framework with two other models describing nutritional upshifts available in the literature, the Flux-Controlled Regulation (FCR) model [13], from which our framework derives, and the Flux Parity Model (FPM) introduced by Chure and coauthors in ref. [24].

As anticipated in the main text, one important aspect by which our framework deviates from the FCR model is that the FCR model assumes that the steady-growth relationship between the ribosomal activity σ and the regulatory functions χ_i is also valid in non-steady conditions and during shifts. This assumption allows the authors to connect these two quantities directly, avoiding the description of the ppGpp mechanistic circuits that implement the synthesis control. By contrast, our model includes explicitly the relation between the translation elongation rate and the ppGpp concentration G, and the transcriptional regulation leading to χ_i, which we use to make predictions about the ppGpp concentration and transcriptional reprogramming during the shift.

Another difference between the FCR model and our model is the description of the amino-acid pool. In the FCR model, the main quantities that control the shift dynamics are the catabolic and biosynthesis fluxes, which are related to the accumulation of amino acids. However, the FCR framework does not explicitly account for the amino acids in the description, and the external condition is described by the fluxes only. In our model, the translation elongation rate is directly connected with the amino acid level, which is justified by the tRNA sensing model.

In contrast, the FPM model follows an optimization principle for defining the control of biosynthesis, always maximizing the growth rate. This model presents the sensing of the charged tRNAs, a quantity that can be related to the amino acid pool in our model. An important aspect of our model is that by explicitly accounting for the sensing and regulatory modules, it does not assume any optimization principle [48].

Fig. 4 of the main text compares the predictions of the three models for a nutrient upshift, showing that, due to their different assumptions, they differ. The FCR model does not predict the amino acid pool and the FPM needs to be integrated with our hypothesis on the charged tRNA pool to do so. Furthermore, the FCR and FPM models do not predict the oscillatory relaxation toward the new steady state found by our model, as they do not explicitly describe the dynamics of the amino acid pool and its feedback interaction with the ribosomal proteins.

TABLE S1: Parameters table

Parameter	Value	Units	Reference
ϕ_R^{max}	0.55	dimensionless [mass fraction]	ref. [10]
ϕ_R^{min}	0.049	dimensionless [mass fraction]	ref. [13]
γ	11.02	rate [h^{-1}]	ref. [13]
$\tilde{\epsilon}$	10.48	rate [h^{-1}]	this work, from Eq. S13
k_A	0.005	dimensionless [mass fraction]	ref. [5]
τ_X	0.16666	time [h]	assumption based on the short mRNA lifespan [36]
K_G	5.9882	concentration [μM]	fit of ref. [17] data (reported here in Fig. S1)
G_{ref}	41.885	concentration [μM]	fit of ref. [17] data (reported here in Fig. S1)
A	1.85E-05	concentration [mass fraction*μM]	ref. [16]
B	2.11E-06	inverse of concentration [mass fraction/μM]	ref. [16]
$\eta_{N,R/P}$	6.8E-05	dimensionless [mass fraction]	ref. [16]
$\eta_{R/P,\phi}$	0.46	dimensionless	ref. [13]
η_0	0.05	rate [h^{-1}]	fit of data reported in ref. [27] (original data from ref. [47])
η_∞	0.009	rate [h^{-1}]	fit of data reported in ref. [27] (original data from ref. [47])
s	5	h	fit of data reported in ref. [27] (original data from ref. [47])
ν_{bg}	0.5	rate [h^{-1}]	this work, fit of data reported here in Fig S2
k_1	1	rate [s^{-1}]	ref. [20]
k_2	0.035	rate [s^{-1}]	ref. [20]
f_c	80	μM	ref. [5] (box 1)
$\tilde{K}_G$	40	concentration [μM]	ref. [20]

 XII.   CALCULATION OF FIXED POINTS.

In order to gain a deeper mathematical understanding of the system, it is desirable to obtain

expressions for the fixed points of the dynamical system as functions of the parameters. It is

particularly informative to examine these expressions in terms of the nutrient quality parameter,

denoted as ν, as it encapsulates the state of the environment and governs nutritional shifts when

altered.

Since our system consists of three coupled differential equations, obtaining a closed analytical

expression for the steady state becomes requires solving a cubic equation. Unfortunately, solving

this cubic equation analytically yields lengthy and non-transparent solutions involving various

combinations of the parameters.

Hence, obtaining a simple expression of the fixed points as solely a function of the parameters

is a complicated task. An easier option is to compute the dependence on the growth rate λ,

which can still be useful. At steady-state our system reads:

$$\epsilon = \tilde{\epsilon}\frac{\psi_{\mathrm{A}}}{\psi_{\mathrm{A}} + k_{\mathrm{A}}}, \tag{S55}$$

$$g = G_{\mathrm{ref}}C\left(\frac{\tilde{\epsilon}}{\epsilon} - 1\right), \tag{S56}$$

$$\lambda = \epsilon\phi_{\mathrm{R}}, \tag{S57}$$

$$\phi_{\mathrm{R}} = \frac{K_{\mathrm{G}}}{K_{\mathrm{G}} + G}. \tag{S58}$$

964   We can solve this system for ϕ_{R}:

$$\phi_{\mathrm{R}}(\lambda) = \frac{\lambda}{\tilde{\epsilon}}\frac{1 - K_{\mathrm{G}}/(G_{\mathrm{ref}}C)}{2}\left(1 + \sqrt{1 + \frac{4K_{\mathrm{G}}/(G_{\mathrm{ref}}C)}{(1 - K_{\mathrm{G}}/(G_{\mathrm{ref}}C))^2}\frac{\tilde{\epsilon}}{\lambda}}\right), \tag{S59}$$

965   and use this expression to obtain the other fixed points as a function of λ:

$$\epsilon(\lambda) = \tilde{\epsilon}\frac{\phi_{\mathrm{R}}}{\phi_{\mathrm{R}} + \frac{K_{\mathrm{G}}}{G_{\mathrm{ref}}C}(1 - \phi_{\mathrm{R}})} = \frac{\lambda}{\phi_{\mathrm{R}}} = \frac{2}{1 - K_{\mathrm{G}}/(G_{\mathrm{ref}}C)}\frac{\tilde{\epsilon}}{1 + \sqrt{1 + \frac{4K_{\mathrm{G}}/(G_{\mathrm{ref}}C)}{(1 - K_{\mathrm{G}}/(G_{\mathrm{ref}}C))^2}\frac{\tilde{\epsilon}}{\lambda}}}, \tag{S60}$$

$$G(\lambda) = G_{\mathrm{ref}}C\left(\frac{\tilde{\epsilon}}{\epsilon} - 1\right) = G_{\mathrm{ref}}C\left(\phi_R\frac{\tilde{\epsilon}}{\lambda} - 1\right) = \frac{G_{\mathrm{ref}}C - K_{\mathrm{G}}}{2}\left(1 + \sqrt{1 + \frac{4K_{\mathrm{G}}/(G_{\mathrm{ref}}C)}{(1 - K_{\mathrm{G}}/(G_{\mathrm{ref}}C))^2}\frac{\tilde{\epsilon}}{\lambda}}\right) - G_{\mathrm{ref}}C, \tag{S61}$$

$$\psi_{\mathrm{A}} = k_{\mathrm{A}}\frac{1}{\frac{\tilde{\epsilon}}{\epsilon} - 1} = k_{\mathrm{A}}\frac{1}{\frac{1 - K_{\mathrm{G}}/(G_{\mathrm{ref}}C)}{2}\left(1 + \sqrt{1 + \frac{4K_{\mathrm{G}}/(G_{\mathrm{ref}}C)}{(1 - K_{\mathrm{G}}/(G_{\mathrm{ref}}C))^2}\frac{\tilde{\epsilon}}{\lambda}}\right) - 1}. \tag{S62}$$

966   Let us comment briefly on these results. The expression for $\phi_{\mathrm{R}}(\lambda)$, Eq. S59, is the so-called

967   "first growth law", and is usually modeled as a linear relation of the growth rate λ. Our solution

highlights that the dependence, in our model, is not exactly linear, but there is a correction term given by the features of the ppGpp regulatory circuit.

The expressions of the translation elongation rate ϵ and the one of the amino acids pool ψ_A are also interesting. Eq. S60 shows that ϵ cannot reach the value $\tilde{\epsilon}$ unless the size of the ribosomal sector ϕ_R reaches 1. This is empirically impossible given that the value of the maximum extension of ϕ_R is about 0.55 (see Table S1 and ref.s [10, 13]). Therefore, in this model, the translational speed is always lower than $\tilde{\epsilon}$, which may seem counterintuitive especially if one just looks at Eq. S14. This fact has consequences on the amino-acid pool, visible in Eq. S62. By having $\epsilon < \tilde{\epsilon}$ regardless of the nutrient quality, this equation shows that the amino-acid pool size never diverges, even in those environments characterized by very rich nutrients. This implies that (within our model) the rate of cellular growth is always limited by the amount of available amino acids, and not just by the allocation strategy of cellular resources.

XIII.   MATHEMATICAL STUDY OF THE APPEARANCE OF THE OSCILLATORY RELAXATION RESPONSE.

As described in the main text, in order to create a biological oscillator, a negative feedback loop with a substantial delay is a necessary requirement [49]. In our system, such a feedback loop exists between the amino acids pool and the ribosomal mass fraction. However, the feedback requirement per se is not sufficient, and oscillatory behavior is not always present in our model. As shown in Fig. S6C-D, there is a certain threshold nutrient quality, referred to as ν^*, beyond which the system exhibits complex eigenvalues, i.e. oscillatory dynamics. The value of ν^* depends on the parameter values.

In order to better understand the emergence of oscillations and their dependence on the parameters, we can perform stability analysis around the fixed points and compute the eigenvalues of the dynamical system, which encode information on how the system relaxes upon weak perturbations of the steady state. Relaxation with damped oscillations corresponds on general grounds to the complex conjugate eigenvalues [49]. In addition, for systems described by just two coupled ODEs (2×2 systems), the determinant and the trace of the Jacobian matrix correspond to the product and the sum of the eigenvalues, hence can be used to fully determine if the relaxation is oscillatory without computing directly the eigenvalues. Indeed in 2×2 systems (i.e. the system with instantaneous transcription described earlier), the determinant and trace of the Jacobian read

$$\det(J) = \frac{\partial \dot{\phi}_R}{\partial \phi_R} \frac{\partial \dot{\psi}_A}{\partial \psi_A} - \frac{\partial \dot{\phi}_R}{\partial \psi_A} \frac{\partial \dot{\psi}_A}{\partial \phi_R} \tag{S63}$$

$$\mathrm{Tr}(J) = \frac{\partial \dot{\phi}_\mathrm{R}}{\partial \phi_\mathrm{R}} + \frac{\partial \dot{\psi}_\mathrm{A}}{\partial \psi_\mathrm{A}} \tag{S64}$$

where the dot represents the time derivative of the quantity. Specifically, 2×2 system presents complex eigenvalues if and only if

$$4 \det(J) > \mathrm{Tr}(J)^2, \tag{S65}$$

if, in addition to this, the following relationship holds

$$\mathrm{Tr}(J) < 0 \wedge \det(J) > 0, \tag{S66}$$

then the fixed points are stable, therefore the oscillations are damped toward the new equilibrium point.

Stability conditions of a generic regulation. Before studying under which conditions Eq.s S65 and S66 hold for our system, we can study whether a system with an unspecified regulatory circuit presents stable fixed points. We can define a 2×2 system for translation regulation with a generic regulatory circuit as follows

$$\frac{\mathrm{d}\psi_\mathrm{A}}{\mathrm{d}t} = \nu \phi_\mathrm{R}^{\max} - \tilde{\epsilon} \frac{\psi_\mathrm{A}}{k_\mathrm{A} + \psi_\mathrm{A}} \phi_\mathrm{R}(1 + \psi_\mathrm{A}), \tag{S67}$$

$$\frac{\mathrm{d}\phi_\mathrm{R}}{\mathrm{d}t} = \tilde{\epsilon} \frac{\psi_\mathrm{A}}{k_\mathrm{A} + \psi_\mathrm{A}} \phi_\mathrm{R} \left(f(\psi_\mathrm{A}, \phi_\mathrm{R}) - \phi_\mathrm{R} \right) \tag{S68}$$

where the first equation corresponds to Eq. S9, because it is not directly tied to the regulation, while the second equation contains a generic function f, which can in principle depend on both ϕ_R and ψ_A, which sets the target size of the ribosomal sector. The partial derivatives of this generic system are

$$\frac{\partial \dot{\psi}_\mathrm{A}}{\partial \psi_\mathrm{A}} = -\phi_\mathrm{R} \tilde{\epsilon} \frac{\psi_\mathrm{A}}{k_\mathrm{A} + \psi_\mathrm{A}} \left(1 + \frac{k_\mathrm{A}}{\psi_\mathrm{A}} \frac{1 + \psi_\mathrm{A}}{k_\mathrm{A} + \psi_\mathrm{A}} \right), \tag{S69}$$

$$\frac{\partial \dot{\psi}_\mathrm{A}}{\partial \phi_\mathrm{R}} = -\nu - \tilde{\epsilon} \frac{\psi_\mathrm{A}}{k_\mathrm{A} + \psi_\mathrm{A}} (1 + \psi_\mathrm{A}), \tag{S70}$$

$$\frac{\partial \dot{\phi}_\mathrm{R}}{\partial \phi_\mathrm{R}} = \phi_\mathrm{R} \tilde{\epsilon} \frac{\psi_\mathrm{A}}{k_\mathrm{A} + \psi_\mathrm{A}} \left(\frac{\partial f}{\partial \phi_\mathrm{R}} - 1 \right), \tag{S71}$$

$$\frac{\partial \dot{\phi}_\mathrm{R}}{\partial \psi_\mathrm{A}} = \phi_\mathrm{R} \tilde{\epsilon} \frac{\psi_\mathrm{A}}{k_\mathrm{A} + \psi_\mathrm{A}} \frac{\partial f}{\partial \psi_\mathrm{A}}. \tag{S72}$$

Note that, given that the above expressions need to be evaluated in a fixed point, some of them are already simplified (for example the terms $f - \phi_\mathrm{R}$ are set to zero). We will now study the positivity of the determinant and the negativity of the trace, starting by noticing that both $\frac{\partial \dot{\psi}_\mathrm{A}}{\partial \psi_\mathrm{A}}$ and $\frac{\partial \dot{\psi}_\mathrm{A}}{\partial \phi_\mathrm{R}}$ are negative regardless of the nature of f.

The condition on the trace implies

$$\frac{\partial \dot{\phi}_{\mathrm{R}}}{\partial \phi_{\mathrm{R}}} < -\frac{\partial \dot{\psi}_{\mathrm{A}}}{\partial \psi_{\mathrm{A}}} \tag{S73}$$

which is the most permissive condition. This condition is surely satisfied if $\frac{\partial f}{\partial \phi_{\mathrm{R}}} < 0$, which implies a regulation that is monotonically decreasing in ϕ_{R}. Moreover, in this case, there is also an homeostatic mechanism towards the target size. Indeed, if ϕ_{R} is slightly lower than the target, the regulation will push for ϕ_{R} to increase, and vice versa in the opposite case.

The condition on the determinant is more complicated because it involves more factors. We can rearrange the terms of the determinant and obtain

$$\frac{\partial \dot{\psi}_{\mathrm{A}}/\partial \psi_{\mathrm{A}}}{\partial \dot{\psi}_{\mathrm{A}}/\partial \phi_{\mathrm{R}}} \left(\frac{\partial f}{\partial \phi_{\mathrm{R}}} - 1 \right) < \frac{\partial f}{\partial \psi_{\mathrm{A}}}, \tag{S74}$$

which defines the least restrictive condition on f in order to have a positive determinant. If we assume that $\frac{\partial f}{\partial \phi_{\mathrm{R}}} < 0$ (given by the condition on the trace), then the left-hand term of Eq. S74 becomes negative, therefore by requiring $\frac{\partial f}{\partial \psi_{\mathrm{A}}} > 0$ the above condition is satisfied. Also this latter condition has a physical interpretation, indeed, in this case, if the amino acids build up then the circuit will increase the target size of ϕ_{R}, and the increased biosynthesis will push back the amino acids pool size. We will not study the presence of oscillations in this generic system, as this requires many assumptions on f, making this study less generic that is intended to be.

Stability conditions in the case with instantaneous transcription. It is possible to gain more specific insight when applying the above requirements to our system, in particular, the model variant with instantaneous transcription, which is a 2×2 system. We start by pointing out that in this model the fixed points are always stable, as the partial derivatives of the regulatory function fulfill the conditions found above for the generic system. In addition, we can insert the definition of the trace and the determinant in Eq. S65, and we find the following requirement for the presence of oscillations,

$$\left(\frac{\partial \dot{\psi}_{\mathrm{A}}}{\partial \psi_{\mathrm{A}}} - \frac{\partial \dot{\phi}_{\mathrm{R}}}{\partial \phi_{\mathrm{R}}} \right)^2 < 4 \frac{\partial \dot{\psi}_{\mathrm{A}}}{\partial \phi_{\mathrm{R}}} \frac{\partial \dot{\phi}_{\mathrm{R}}}{\partial \psi_{\mathrm{A}}}. \tag{S75}$$

The left-hand side term of Eq. (S75) contains two characteristic time scales: $\frac{\partial \dot{\psi}_{\mathrm{A}}}{\partial \psi_{\mathrm{A}}}$ and $\frac{\partial \dot{\phi}_{\mathrm{R}}}{\partial \phi_{\mathrm{R}}}$, which are, respectively, the time scale of the reaction of ψ_{A} in response to changes in ϕ_{R} and the reaction time scale of ϕ_{R} in response to changes in ψ_{A} [49]. Similarly, the right-hand side term of Eq. (S75) can be interpreted as an interaction strength between the two quantities. For a system where the time scales of the feedback interactions match, the left-hand term will always be zero and therefore the system will always show damped oscillations during the relaxation to the new steady state.

The specific values of the system parameters determine whether or not inequality S75 holds,

setting a boundary between damped oscillations and exponential relaxation in the phase diagram of the system. We can calculate the partial derivatives of the system:

$$\frac{\partial \dot{\psi}_A}{\partial \psi_A} = -\phi_R \tilde{\epsilon} \frac{\psi_A}{k_A + \psi_A} \left(1 + \frac{k_A}{\psi_A} \frac{1 + \psi_A}{k_A + \psi_A}\right), \tag{S76}$$

$$\frac{\partial \dot{\psi}_A}{\partial \phi_R} = -\nu - \tilde{\epsilon} \frac{\psi_A}{k_A + \psi_A}(1 + \psi_A), \tag{S77}$$

$$\frac{\partial \dot{\phi}_R}{\partial \phi_R} = -\phi_R \tilde{\epsilon} \frac{\psi_A}{k_A + \psi_A}, \tag{S78}$$

$$\frac{\partial \dot{\phi}_R}{\partial \psi_A} = \phi_R \tilde{\epsilon} \frac{\psi_A}{k_A + \psi_A} \frac{\frac{K_G/(CG^{\text{ref}})}{k_A}}{\left(\frac{K_G/(CG^{\text{ref}})}{k_A}\psi_A + 1\right)^2}. \tag{S79}$$

Note that $\frac{\partial \dot{\phi}_R}{\partial \psi_A}$ depends solely on a single parameter, namely $\frac{K_G/(CG^{\text{ref}})}{k_A}$, which represents the ratio of two scales. The first, denoted as k_A, represents the concentration of amino acids at which the elongation rate ϵ attains half of its maximum value. The second, $K_G/(CG^{\text{ref}})$, corresponds to the ratio of the Michaelis-Menten constant of $\chi_R(G)$, K_G, governing the extent of ppGpp required for altering resource allocation, and the sensitivity of ppGpp concerning variations in the translation elongation rate ϵ: CG^{ref}. We refer to this factor as $\tilde{G} = K_G/(CG^{\text{ref}})$. This observation implies that the ribosomal resource allocation strategy is solely contingent on the relationship between these two scales. However, as we will demonstrate in this section, the overall system dynamics are more intricate.

By substituting the partial derivatives into Eq. S75, we derive the subsequent condition for ν to initiate oscillations,

$$\frac{\tilde{\epsilon} k_A \tilde{G}}{4 \phi_R^{\text{max}}} < \nu \frac{(\psi_A(\nu) + k_A)^3}{\psi_A(\nu)(1 + \psi_A(\nu))^2}. \tag{S80}$$

Since Eq. S80 still incorporates a state variable (the amino acids pool ψ_A), it is not possible to derive a phase diagram solely based on a single dimensionless parameter combination. Instead, we adopt the steady-state function fit $\psi_A^*(\nu)$ to obtain the trends shown by Fig. S7.

To acquire the function $\psi_A^*(\nu)$, we chose to simulate the system's evolution for varying nutrient qualities (ν), permitting the system to attain equilibrium. Subsequently, we collected the essential data, such as the equilibrium point coordinates ($\nu, \psi_A, \phi_R, \lambda$), or the relationship between ψ_A and ϕ_R concerning ν. This data was then subjected to a polynomial fitting process, simplifying analysis and interpretation

Dimensional analysis helps to characterize the phase diagram of damped oscillations. Although we could only analyze Eq. S80 numerically, we can isolate several key trends by dimensional analysis, leveraging Eq. S80 to better characterize the phase diagram. We initiated this by plotting the right-hand side of Eq. S80 divided by ν, as a function of ν itself. Figure S7A

indicates that the RHS term can be approximated as linear in ν, implying that it can be expressed as $\frac{(\psi_A(\nu)+k_A)^3}{\psi_A(\nu)(1+\psi_A(\nu))^2} = f_{\text{const}}(\tilde{G}, \tilde{\epsilon}, k_A)\nu$, where f_{const} represents a constant function of the system parameters.

To further break down the function f_{const}, we analyzed the dimensions of the terms in

$$\frac{\tilde{\epsilon} k_A \tilde{G}}{4\phi_R^{\max}} < \nu^2 f_{\text{const}}(\tilde{G}, \tilde{\epsilon}, k_A), \tag{S81}$$

which is Eq.S80 with the given RHS ansatz. We note that the LHS dimension is h^{-1} due to $\tilde{\epsilon}$, while the RHS dimension is h^{-2} because of ν^2. This implies that the RHS function must have the dimension of h, specifically $f_{\text{const}}(\tilde{G}, \tilde{\epsilon}, k_A) = h_{\text{const}}(\tilde{G}, k_A)/\tilde{\epsilon}$.

Substituting this relation into the previous equation, we derive

$$\left(\frac{k_A \tilde{G}}{4\phi_R^{\max}}\right)/h_{\text{const}}(\tilde{G}, k_A) < \frac{\nu^2}{\tilde{\epsilon}^2}, \tag{S82}$$

signifying that $\nu^*/\tilde{\epsilon}$ must be a constant. This prediction is verified by panel B of Fig. S7, illustrating that $\nu^*/\tilde{\epsilon}$ remains well approximated by a constant while altering the value of $\tilde{\epsilon}$.

Unfortunately, breaking down the dependence on the other parameters is not as straightforward. Nonetheless, we can use numerical simulations to show how $\nu^*/\tilde{\epsilon}$ changes with alterations in the parameter values (Fig. S7C-D-E). In particular, we can analyze how h_{const} changes when the other parameters involved in Eq. S82 change, by extracting and plotting the function

$$h_{\text{const}}(\tilde{G}, k_A) = \left(\frac{k_A \tilde{G}}{4\phi_R^{\max}}\right) \frac{\tilde{\epsilon}^2}{(\nu^*)^2}, \tag{S83}$$

as a function of the parameters.

Amino acid dynamics (and not only ppGpp regulation) contribute to the damped oscillations threshold. Lastly, we aim to ascertain whether the effective parameter discovered earlier appears in $h_{\text{const}}(\tilde{G}, k_A)$. To this aim, we examined the trend of $\nu^*/\tilde{\epsilon}$ in a scenario where K_G and C change while maintaining $\tilde{G} = K_G/(CG^{\text{ref}})$ constant. Fig. S7 panel C demonstrates that the parameter $\tilde{G}$ indeed functions as an effective parameter, as $\nu^*/\tilde{\epsilon}$ remains constant even if $\tilde{G}$ changes.

In contrast, panel D illustrates that even though the parameter combination $\tilde{G}/k_A$ appears in the ribosomal regulatory function χ_R, it is insufficient to classify it as an effective parameter for the entire system. Indeed in our system, k_A also appears in the equation governing amino acid dynamics, but not in combination with $\tilde{G}$. Consequently, amino acid dynamics play a pivotal role in the emergence of oscillatory behavior, as this characteristic is not solely reliant on biosynthesis regulation.

 Different results for the model including transcriptional delays. This SI section has so far
focused on the study of the reduced 2×2 dynamical system, because of the presence of simple
criteria that ensure the stability of the fixed points and the nature of the relaxation. It is
important to highlight that some of the considerations above do not apply to the complete model
(a 3×3 dynamical system), which includes the transcriptional timescale for mRNA production,
and hence should be more realistic.

For this complete model, there are no simple criteria that describe stability and relaxation,
therefore, as mentioned in the main text, we performed a direct study of the eigenvalues of the
system, shown in Fig. S6, together with the eigenvalues of the 2×2 system with instantaneous
transcription. This analysis shows that our model exhibits a variety of behaviors depending
on the parameters describing the nutritional upshift. From the analysis of the eigenvalues
of the system, we can identify three regimes: (I) a shift without oscillations at slow growth
rates, (II) a shift with damped oscillations in the fast growth regime, and (III) a shift with
sustained oscillations (Hopf bifurcation) at growth rates far above the maximum achievable one
experimentally [28]. The presence of sustained oscillations occurs when the real part of the
complex eigenvalues becomes positive. These oscillations are shown in Fig. S8. Note that all
these three growth regimes are compatible with a system that can achieve balanced growth
during steady-state [28]. As for the 2×2 system, the exact threshold between these regimes
depends on the parameters of the system.

XIV. THE PROPOSED MODEL IS ANALOGOUS TO A NEGATIVE AUTOREGULATION.

In this section, we argue that the emergent feedback from the global regulation in our model
can be paralleled to a negative autoregulation. A signature of the negative autoregulation is
that the production rate of a quantity x depends on the quantity itself, and it decreases as x
increases.

We will argue that this holds for ϕ_R, by focusing on the feedback loop between the amino
acid pool ψ_A and the ribosomal sector size ϕ_R, described by Eq. S9 and S29. The steady-state
expression of ψ_A reads

$$\psi_A^* = \frac{\nu}{\lambda}(\phi_R^{\max} - \phi_R) - \frac{\epsilon}{\lambda}\phi_R \ , \tag{S84}$$

which is a monotonic decreasing function of ϕ_R. We can insert this expression in the equation
for χ_R, which is the production term of ϕ_R (Eq. S29). We obtain the following expression

$$\chi_R = K_G \left(K_G + G^{\mathrm{ref}} C \left(\frac{\psi_A + k_A}{\psi_A} - 1 \right) \right)^{-1} \ , \tag{S85}$$

which shows that the regulatory function χ_R is monotonically increasing in the amino acid concentration ψ_A. This follows from the fact that as the amino acids builds up, ppGpp production decreases, and the target for ribosome transcription increases. In turn, ψ_A^* decreases with increasing ϕ_R, and this makes the regulatory function a decreasing function of ϕ_R. For small circuits, negative autoregulation feedback loops were argued to provide quicker response to perturbation, and speculated to be selected based on this trait [50].

XV.   ANALYSIS OF EXPERIMENTAL DATA FROM PANLILIO ET AL. 2021.

In this work, we utilize the nutritional shift data reported in ref. [22]. The authors of that study investigate the dynamics of the shift at the single-cell level using a mother-machine microfluidic device. We leverage their data to compare the model predictions with experimental results. Figure 5 of the main text presents the outcomes of this comparative analysis.

A.   Experimental apparatus and processed data

The microfluidic chip consists of parallel channels where the cells grow, with nutrients flowing from both the top and bottom of each channel. The data acquisition process used a microscope to capture images of approximately 10 channels per field of view simultaneously, with a time interval of 5 minutes between each capture. The acquired images were then processed using tracking software, which analyzed the cell growth on a frame-by-frame basis. The software recognized the cell's area and width and computed its volume using a sphero-cylindrical approximation. Panlilio and colleagues employed *E. coli* strains with fluorescent tags applied to the ribosomal (P1 and P1long) or constitutive promoters (P5). These tags were inserted near the replication origin ("ori" strains) or near the terminus ("ter" strains) of the genetic material. The constitutive P5 promoter is derived from the bacteriophage T5 [51]. On the other hand, the growth rate-dependent *E. coli* ribosomal promoter rrnB P1 is one of the two promoters responsible for initiating the transcription of the rRNA operon [52, 53]. The native P1 promoter contains several distinctive features, including a GC-rich discriminator region that renders it sensitive to the alarmone ppGpp [54, 55]. It also possesses an upstream sequence (UP element) that significantly enhances promoter activity [56]. Furthermore, the P1 promoter contains three Fis-binding sites located even further upstream, which further activate the promoter under nutrient-rich conditions [57–59]. P1long refers to the full-length rrnB P1 promoter, while P1 denotes the truncated version lacking the Fis-binding sites but still retaining the discriminator and UP element. It is important to note that the P5 promoter lacks the discriminator region,

UP element, and Fis-binding sites. For a more detailed description of the promoters refer to ref.s [22, 60]. By using these fluorescently labeled strains, the authors obtained information on the fluorescence intensity, which is indicative of protein expression levels, throughout the entire experiment.

The performed shift was an upshift from minimal medium (M9) + glucose to M9 + glucose + casaminoacids [22]. The relevant processed data for our analysis consists of the time series data for each cell, including volume and fluorescence intensity were downloaded from the Mendeley Data repository associated to the study (DOI: 10.17632/5d4yhyjn8j.1). These data report on volume, cell-division events and fluorescence intensity of the tracked cell lineages.

B. Computation of population data, growth rate, and sector size

We focused exclusively on the population-average time series of the growth rate and sector size (derived from the fluorescent reporters) across the shift. This section describes how we derived these average quantities from the single-cell data.

a. Population averages. We transformed the individual cell time series into population-average time series by calculating the average signal across all cells present at each experimental time interval using a sliding window average. The step size was 4 minutes, and the window size was 75 minutes, resulting in the coarse-grained population time series shown in Fig. 5 of the main text. In contrast, Fig. S9 displays a less coarse-grained population time series, with a step size of 4 minutes and a window size of 20 minutes.

b. Growth rate. To compute the growth rate, we used the single-cell time series of the volume. We applied a Savitzky-Golay filter to the data and then calculated the growth rate as the central derivative of the volume with respect to time, divided by the volume

$$\lambda(t_\mathrm{i}) = \frac{1}{V(t)}\frac{V(t_{\mathrm{i}+1}) - V(t_{\mathrm{i}-1})}{t_{\mathrm{i}+1} - t_{\mathrm{i}-1}}. \tag{S86}$$

This was done for each cell, and the resulting growth rates were averaged across all cells as described above to obtain the population time series.

c. Proteome sector size. To extract information on the proteome sector size from the fluorescence signal, we first normalized the fluorescence intensity by the cell volume. This ratio is expected to be proportional to the protein concentration. To determine the conversion constant between the fluorescence signal and the proteome sector size, we used different procedures for the ribosomal and constitutive protein signals.

For the ribosomal signal, we used the growth law at steady state, along with the data on the growth rate, to find the conversion factor between the GFP signal normalized by the volume

and the ribosomal sector size. This conversion factor was denoted as $k_{\text{conv,R}}$ and was defined such that it satisfies

$$k_{\text{conv,R}} \left(\frac{F_{\text{p1}}}{V} \right)_{\text{s1}} = \phi \text{R}^{\min} + \frac{\lambda_{\text{s1}}}{\gamma}, \tag{S87}$$

where the subscript $s1$ denotes the pre-shift steady state. The ribosomal time series was then obtained as the product of the conversion factor and the fluorescence signal normalized by the cell volume, $\phi_{\text{R}}(t) = k_{\text{conv,R}} \frac{F_{\text{p1}}(t)}{V(t)}$.

For the constitutive protein signal, we utilized the constraint that the sum of the ribosomal and constitutive protein sector sizes is constant $\phi_{\text{R}} + \phi_{\text{P}} = 0.55$. We defined the conversion factor $k_{\text{conv,P}}$ such that it satisfies

$$\phi_{\text{R,s1}} + k_{\text{conv,P}} \left(\frac{F_{\text{p5}}}{V} \right)_{\text{s1}} = 0.55. \tag{S88}$$

The constitutive protein time series was then obtained as the product of the conversion factor and the fluorescence signal normalized by the cell volume, $\phi \text{P}(t) = k_{\text{conv,P}} \frac{F_{\text{p5}}(t)}{V(t)}$.

C. Consistency checks on the fluorescence proxies

The approach used to compute the proteome sector size in this study is unique in the literature, as most previous studies have relied on macromolecular quantification or proteomic assays to assess the proteome [10, 12, 13, 37]. To ensure that using the fluorescence signal as a proxy for the proteome is valid, we conducted several consistency checks on the data.

Firstly, we examined whether the conversion factor $k_{\text{conv,R}}$ computed using the pre-shift steady-state data remained consistent with the post-shift data. If different conversion factors were obtained for the two steady states within a time series, it would indicate a potential issue with either the steady states or the experiment itself. Panel A of Fig. S10 shows that the conversion factor computed for the first steady state in the pair $(\lambda^{\text{p5-ter}}, \phi_{\text{R}}^{\text{p1long-ori}})$ correctly places the point of the second steady state on the trend of the first growth law from literature data, as expected. However, the other pairs, where $\phi_{\text{R}}^{\text{p1long-ori}}$ is coupled with the growth rate from other strains (p1-ori, p1long-ori, and p1long-ter), did not meet this requirement. It is important to note that we excluded the fluorescence signal of the p1-ori and p1long-ter strains from this test for two main reasons: (I) the promoter P1long is the true ribosomal promoter as it conserves all the discriminatory regions and the "ori" position ensures that phenomena like gene dosage and supercoiling are taken into account, (II) they do not exhibit a post-shift steady state (shown in Fig. S9B).

The second check we performed focused on the constitutive sector. We examined whether the constraint $\phi_{\text{R}} + \phi_{\text{P}} = 0.55$ held throughout the entire shift, despite the fact that the two

sector fractions were derived from independent fluorescent reporters in different experiments. Panel B of Fig. S10 demonstrates that this constraint holds during the entire shift, even if it is enforced on just the steady-state pre-shift data.

D.   A cell-density imbalance alters the shift dynamics

The high temporal resolution of the data collected by Panlilio and colleagues exhibits a recently studied biophysical phenomenon occurring during amino acid upshifts, which goes beyond our model. Specifically, in the first minutes following the shift, the volume and mass of a cell become temporarily uncoupled. This feature appears in similar shifts that were studied recently monitoring synchronously both the volume and the refractive index of individual cells [29]. In the Panlilio et al. data, upon close examination of the growth rate data presented in Fig. S9 and Fig. S11, it is apparent that all the population-average time series exhibit a rapid over-shoot in volume growth rate that is not matched by the protein dynamics. This discrepancy can be attributed to the osmotic imbalance that occurs immediately after the shift, as noted by Oldewurtel and colleagues [29].

This volume-mass growth imbalance has the potential to impact the shift dynamics by introducing a perturbation in the protein density. It is important to note that our model, as well as most shift models available in the literature, cannot account for this phenomenon, as a crucial assumption of most available models is that the protein density remains constant and that the sector sizes reflect the intracellular concentrations of the different proteins.

XVI.   OTHER SHIFT EXPERIMENTS OBSERVED THE OSCILLATORY DYNAMICS.

The observed oscillatory behavior does not seem to be limited to the data presented by Panlilio et al. We have come across two additional experiments that suggest the possibility of oscillations. The first experiment, described in ref. [61] and depicted in Fig. S12A, demonstrates rapid oscillations of the ribosomal regulatory function χ_R following an upshift in amino acids. The second set of experiments, reported in a more recent paper, ref. [13], involves various upshift experiments by introducing a new carbon source in addition to the one already present in the medium. The corresponding data is shown in Fig.S12B-C. Although the putative oscillations in this case may be less pronounced, this may be attributed to the lower temporal resolution of these data. We observe that the three instances of oscillatory behavior (from ref.S [13, 22, 61]) exhibit distinct time scales, ranging from minutes to hours, which are not captured by our current

model. It is important to emphasize that further studies, both theoretical and experimental, are necessary to gain a better understanding of this phenomenon.

 SUPPLEMENTARY FIGURES

FIG. S1: **Fit of the dependency RNAP regulatory function** ω_R **from ppGpp concentration from experimental data.**. The plot shows steady-state data from ref. [16, 17], the y axis represents the ribosomal sector size, while the x axis reports the ppGpp concentration. To obtain the plotted data, we have taken the data of ppGpp concentration versus the growth rate reported in ref. [17] and transformed the growth rate in the ribosomal sector size using the first growth law [13]. These data are shown as blue squares. Additionally, we have taken the data in ref [16], which reports the fold change of ppGpp levels versus the ribosomal sector size, and converted the fold change into absolute concentration using the data from ref. [17]. For the conversion, we used the value of ppGpp corresponding to a growth rate of approximately 0.9, which should be close to the growth on glucose that has been taken as a reference in ref. [16]: $G^{\mathrm{ref}} = 101.46 \mu M$.

FIG. S2: **Adding degradation and inactive ribosomes is sufficient to quantitatively describe the slow growth regime.** The figure compares two versions of the proposed framework: the fast-growth approximation (red circles) proposed in the main text and a full model that considers degradation and inactive ribosomes (orange circles). **A:** results for both models, along with experimental data from [16]. The x-axis indicates the steady-state exponential growth rate (shared with panels B-C), while the y-axis represents the size of the ribosomal sector. **B:** growth rate shown on the x-axis, with the relative translation rate shown on the y-axis. **C:** degradation rate vs growth rate, for the full model (orange rounds) and for the data reported in [27, 47]. This panel shows an input of the full model, whereby degradation depends on growth rate in a non-linear fashion. The results obtained from the fast-growth model predict that $\phi_R \to 0$ and $\epsilon \to 0$ as $\lambda \to 0$, which is in contrast with experimental evidence. However, both ϕ_R and ϵ show a non-zero offset at zero growth rate when the full model is considered. This is explained by the fact that at slow growth rates ($\lambda < 0.5h^{-1}$), the phenomena of protein degradation and inactive ribosomes become relevant for growth physiology, as described in [27].

Model with amino acids dilution   Model with(out) amino acids dilution

$$\frac{d\psi_A}{dt} = \nu\phi_P - \epsilon\phi_R - \lambda\psi_A \qquad \frac{d\psi_A}{dt} = \nu\phi_P - \epsilon\phi_R$$

A metabolic flux $J_a = \nu\phi_P$

B biosynthesis flux $J_b = \epsilon\phi_R$

C amino acids pool changes $\frac{d\psi_A}{dt}$

time [h]   time [h]

FIG. S3:   **Neglecting the dilution of the amino acid pool affects the predicted time scales of adaptation.** The plots compare the dynamics of a model that considers amino acid dilution due to cell growth (main text, right panels) and a model that neglects it (left panels). In all panels, the x axis represents the time from the shift (note that the span of the axis is different in the right and left panels). **A:** metabolic flux dynamics following an upshift. The metabolic flux, which supplies the amino acids pool, is given by the nutrient quality ν times the size of the constitutive sector ϕ_P. We assume ν changes instantaneously over the shift, and so does the metabolic flux. **B:** biosynthesis flux dynamics following an upshift. This flux represents the protein production and is given by the translation elongation rate ϵ times the ribosomal sector size ϕ_R. **C:** changes in the amino acids pool following an upshift. This plot shows the dynamics of the derivative of the size of the amino acids pool, which is also the difference between the two fluxes and, when considered, the dilution term (see equations at the top of the panels). The figure shows that both models reach equilibrium after being perturbed, contrary to what is shown in ref [24]. This happens because in our model the feedback between amino acid pool and ribosomes ensures that the disparity between the fluxes is adjusted by tuning the proteome composition, and this is true also for the model without dilution. However, we note that if the dilution is neglected the system needs more time to reach equilibrium after the perturbation.

FIG. S4: **The model version that includes just the action of RelA still presents the oscillatory dynamics.** Panels **A-C** show the steady state prediction of this version of the model and experimental data from ref.s [26] (cyan crosses) and [16] (blue triangles), in particular, is plotted the ribosomal sector size (**A**) and the relative translation elongation rate (**B**) versus the growth rate, and the ppGpp concentration versus the inverse of the relative translation rate (**C**). These panels show that overall this model is performing well in the fast growth regime, except for the last prediction on ppGpp, which is the one that falsifies it. Panels **D-G** show the model predictions following an upshift (**D:** relative translation rate, **E:** ribosomal sector, **F:** growth rate, **G:** ppGpp concentration). These plots show that even if the relation between ppGpp and ϵ has changed the model still predicts an oscillatory relaxation towards the new steady state.

FIG. S5: **The FCR model variant can still reproduce the overall shift dynamics.** The plot shows the dynamics of the principal quantities involved during nutrient shifts predicted by the FCR model equipped with the description of the ppGpp dynamics introduced in this work, and it compares with shift data collected in ref. [13]. The first two rows show the upshift dynamics, while the two bottom ones are the downshift one. All plots display the time from the shift on the x-axis. The first column (**A1-4**) shows the comparison between the predicted growth rate (red line) and the one observed in ref. [13] (blue reverse triangles). The second column (**B1-4**) shows the dynamics of the relative ribosomal activity $\sigma/\sigma_{\max}$, which rapidly increases during upshifts and decreases and recovers during downshifts. The third column (**C1-4**) shows the dynamics of the ribosomal sector and its regulatory function χ_{R} (grey fine dashed line). In particular, we show the dynamics of the total ribosomes (red solid line) and of the sequestered ones (red dashed line), which are necessary in order to include the ppGpp dynamics, as described in these notes. The last column (**D1-4**) shows the dynamics of the catabolic sector ϕ_{cat} (solid lines) and of its regulatory function χ_{cat} (grey fine dashed line). This model describes a different uptake sector for each substrate present in the medium, therefore in this case we included two different sectors, one of which gets activated or diluted after an up or downshift, respectively.

FIG. S6: **The eigenvalues from linear-stability analysis predict the presence of stable fixed points and oscillatory relaxation towards the new steady state.** This figure shows results for two versions of the model, the one where transcriptional reprogramming from ppGpp has a timescale given by the degradation of the transcripts and a variant where transcription is instantaneous. **A:** real part of the eigenvalues plotted as a function of the nutrient quality ν. The real part of the eigenvalues is always negative in the relevant range of parameters, indicating stable fixed points. **B:** the real part of the eigenvalues of the model variant with instantaneous transcription remains negative. Panels **C** and **D** show the imaginary parts of the eigenvalues of the instantaneous and gradual transcription models as a function of nutrient quality ν. Above a threshold value of ν, the eigenvalues become complex, leading to oscillatory relaxation. In all panels, the grey areas represent the slow-growth regime, defined as $\lambda < 0.5h^{-1}$, the dashed grey line is an empirical maximum growth rate $\lambda \simeq 2h^{-1}$.

FIG. S7: **The onset of oscillatory behavior depends on key parameters such as maximum elongation speed and sensitivity of ppGpp to elongation rate.** These panels show the study of the threshold nutrient quality ν^* value triggering damped oscillations (a value that depends on the model parameters). **A:** the x-axis represents the nutrient quality ν, while the y-axis shows the right-hand side of eq S80 (the definition id near the vertical axis). The red round points are the result of the numerical simulation of the RHS, while the blue dashed line is a linear fit of these points (for $\nu > 2$). This plot shows that for large ν the RHS can be approximated as a linear function of ν. In all the subsequent panels we have studied the parameter dependence of ν^* in order to understand better the phase diagram of the system. The green points correspond to $\nu^*/\tilde{\epsilon}$ given the parameter values, the green shaded area indicates the parameter region where the model predicts oscillatory behavior, while the white area corresponds to overdamped relaxation. The explored range of all parameters spans from 0.5 to 2 times the value of the parameter estimated for the wild type, reported in table 1. **B:** trend of $\nu^*/\tilde{\epsilon}$ versus $\tilde{\epsilon}$. The panel shows that $\nu^*/\tilde{\epsilon}$ is almost constant in $\tilde{\epsilon}$, as predicted by the dimensionality study. **C:** $\nu^*/\tilde{\epsilon}$ is constant if K_G and C are changed in such a way that $K_G/(CG^{\mathrm{ref}}$ remains constant. This demonstrates that three parameters K_G, C and G^{ref} appears always in combination in Eq. S80. **D:** trend of h_{const} (Eq. S82) versus k_A, the parameter that controls the sensitivity of the translation elongation rate to amino acid levels (Eq. S14). **E:** trend of h_{const} (Eq. S82) versus $\tilde{G}$.

FIG. S8: **Presence of Hopf bifurcation in non-biological regimes.** All the plots are obtained by simulating an upshift in a regime where growth exceeds the maximum achievable growth rate. **A:** dynamics of the relative translation elongation rate $\epsilon/\tilde{\epsilon}$ across the shift. **B:** dynamics of the ribosomal sector size ϕ_R across the shift. **C:** dynamics of the instantaneous growth rate λ across the shift. The plots show that in this regime the oscillations are not damped, and the system shows sustained oscillations around the new equilibrium. Mathematically, this happens because the real part of the complex eigenvalues becomes positive if the nutrient quality exceeds a certain threshold, as can be seen in Fig. S6A. This phenomenon is called Hopf bifurcation in the theory of dynamical systems.

FIG. S9:   **Time series of the shift dynamics for the different strains utilized in Panlilio et al. 2021 [22]. A-C:** data from all the five studied strains, averaged across experimental replicates. **D-F:** time series used in this work. The first row shows the shift time series of the growth rate (**A**), the fluorescence signal associated with the ribosomal promoter (p1 and p1long) divided by the volume (**B**) and the fluorescence over volume for the constitutive promoter p5 (**C**). The second row shows the same data, but just for the selected strains. Panels **E** and **F** also show the standard deviation of the population average. The chosen growth rate is from p5-ter because we find it consistent with the first growth law. The fluorescent signal of the ribosomal promoter is p1long-ori because is the only one that displays a post-shift steady state. The fluorescent signal of the constitutive promoter is p5-ter because is consistent with the constraint $\phi_P + \phi_R = 0.55$.

FIG. S10:   **The fluorescent reporters from Panlilio et al. consistently obey the proteome sector relationships. A:** steady states for the pair of time series $(\lambda^{\mathrm{p5-ter}}, \phi_R^{\mathrm{p1long-ori}})$. Once fixed the conversion factor $k_{\mathrm{conv,R}}$ using the data from the pre-shift steady state (circle) the post-shift steady state (plus sign) automatically obeys the first growth law (black line [13]), as expected. The inset shows the performance of the other pairs, obtained by pairing up $\phi_R^{\mathrm{p1long-ori}}$ with growth rate from the other remaining strains The symbols are blue for $(\lambda^{\mathrm{p5-ter}}, \phi_R^{\mathrm{p1long-ori}})$, and other colors correspond to the other pairs. **B:** once fixed the conversion factor for ϕ_P using the pre-shift conditions, the constraint $\phi_R + \phi_P = 0.55$ is satisfied throughout the shift. The results shown in these two panels suggest that the fluorescent reporters can be a robust proxy for the ribosomal and constitutive sector size. Original data from ref. [22].

FIG. S11: **The rapid growth rate overshoot in the data from Panlilio et al. may be linked with a cell-density unbalance. A:** shift dynamics of the volume growth rate of the strain p5-ter [22], The grey box highlights the fast rapid overshoot of this quantity. **B:** Replotted data from ref. [29] form a similar amino-acid upshift (from minimal medium+mannose to minimal medium+glucose+casaminoacids) on agar pads, monitoring accumulation rates of single-cell volume and mass following a shift. Panel B shows that following a shift, the volume and the mass display very different dynamics and the cells become diluted. Indeed, cell volume increases 15 % faster than mass in the first 50 min [29]. This dilution, possibly of osmotic origin, could be the reason why the rapid overshoot in the volume growth rate is not matched by an overshoot in protein production, as monitored by the fluorescent reporters.

FIG. S12: **Oscillations in nutrient shift experiments occur on varying time scales.** Panel **A:** shows data for the ribosomal allocation χ_R dynamics following an upshift (data from ref. [61]). The plot shows two different experimental replicates where the relative rate of ribosome production was measured by leucine incorporation. **B-C:** growth rate λ dynamics following an upshift in bulk cultures (data from ref. [13]). In both panels, the experimental data appear to present an overshoot around 1h after the shift.

[1] J. Zhu and C. B. Thompson, Metabolic regulation of cell growth and proliferation, Nature Reviews Molecular Cell Biology **20**, 436 (2019).

[2] L. Galdieri, S. Mehrotra, S. Yu, and A. Vancura, Transcriptional regulation in yeast during diauxic shift and stationary phase, OMICS: A Journal of Integrative Biology **14**, 629 (2010).

[3] P. C. Marijuán, J. Navarro, and R. del Moral, On prokaryotic intelligence: Strategies for sensing the environment, Biosystems **99**, 94 (2010).

[4] M. Scott and T. Hwa, Bacterial growth laws and their applications, Current Opinion in Biotechnology **22**, 559 (2011).

[5] M. Scott, S. Klumpp, E. M. Mateescu, and T. Hwa, Emergence of robust growth laws from optimal regulation of ribosome synthesis, Molecular Systems Biology **10**, 747 (2014).

[6] L. U. Magnusson, A. Farewell, and T. Nyström, ppGpp: a global regulator in escherichia coli, Trends in Microbiology **13**, 236 (2005).

[7] K. Potrykus, H. Murphy, N. Philippe, and M. Cashel, ppGpp is the major source of growth rate control in e. coli, Environmental Microbiology **13**, 563 (2010).

[8] B. D. Towbin, Y. Korem, A. Bren, S. Doron, R. Sorek, and U. Alon, Optimality and sub-optimality in a bacterial growth law, Nature Communications **8**, 10.1038/ncomms14123 (2017).

[9] E. Bosdriesz, D. Molenaar, B. Teusink, and F. J. Bruggeman, How fast-growing bacteria robustly tune their ribosome concentration to approximate growth-rate maximization, The FEBS Journal **282**, 2029 (2015).

[10] M. Scott, C. W. Gunderson, E. M. Mateescu, Z. Zhang, and T. Hwa, Interdependence of cell growth and gene expression: Origins and consequences, Science **330**, 1099 (2010).

[11] R. Hermsen, H. Okano, C. You, N. Werner, and T. Hwa, A growth-rate composition formula for the growth of e. coli on co-utilized carbon substrates, Molecular Systems Biology **11**, 801 (2015).

[12] S. Hui, J. M. Silverman, S. S. Chen, D. W. Erickson, M. Basan, J. Wang, T. Hwa, and J. R. Williamson, Quantitative proteomic analysis reveals a simple strategy of global resource allocation in bacteria, Molecular Systems Biology **11**, 784 (2015).

[13] D. W. Erickson, S. J. Schink, V. Patsalo, J. R. Williamson, U. Gerland, and T. Hwa, A global resource allocation strategy governs growth transition kinetics of escherichia coli, Nature **551**, 119 (2017).

[14] M. V. Rojiani, H. Jakubowski, and E. Goldman, Relationship between protein synthesis and concentrations of charged and uncharged tRNATrp in escherichia coli., Proceedings of the National

Academy of Sciences **87**, 1511 (1990).

[15] E. Goldman and H. Jakubowski, Uncharged tRNA, protein synthesis, and the bacterial stringent response, Molecular Microbiology **4**, 2035 (1990).

[16] C. Wu, R. Balakrishnan, N. Braniff, M. Mori, G. Manzanarez, Z. Zhang, and T. Hwa, Cellular perception of growth rate and the mechanistic origin of bacterial growth law, Proceedings of the National Academy of Sciences **119**, 10.1073/pnas.2201585119 (2022).

[17] A. G. Marr, Growth rate of escherichia coli, Microbiological Reviews **55**, 316 (1991).

[18] T. M. Wendrich, G. Blaha, D. N. Wilson, M. A. Marahiel, and K. H. Nierhaus, Dissection of the mechanism for the stringent factor RelA, Molecular Cell **10**, 779 (2002).

[19] H. Bremer and P. P. Dennis, Modulation of chemical composition and other parameters of the cell at different exponential growth rates, EcoSal Plus **3**, 10.1128/ecosal.5.2.3 (2008).

[20] I. Shachrai, A. Zaslaver, U. Alon, and E. Dekel, Cost of unneeded proteins in e. coli is reduced after several generations in exponential growth, Molecular Cell **38**, 758 (2010).

[21] A. Bren, Y. Hart, E. Dekel, D. Koster, and U. Alon, The last generation of bacterial growth in limiting nutrient, BMC Systems Biology **7**, 10.1186/1752-0509-7-27 (2013).

[22] M. Panlilio, J. Grilli, G. Tallarico, I. Iuliani, B. Sclavi, P. Cicuta, and M. C. Lagomarsino, Threshold accumulation of a constitutive protein explains e. coli cell-division behavior in nutrient upshifts, Proceedings of the National Academy of Sciences **118**, 10.1073/pnas.2016391118 (2021).

[23] M. Mori, S. Schink, D. W. Erickson, U. Gerland, and T. Hwa, Quantifying the benefit of a proteome reserve in fluctuating environments, Nature Communications **8**, 10.1038/s41467-017-01242-8 (2017).

[24] G. Chure and J. Cremer, An optimal regulation of fluxes dictates microbial growth in and out of steady-state, eLife **12**, 10.7554/elife.84878 (2023).

[25] R. Balakrishnan, M. Mori, I. Segota, Z. Zhang, R. Aebersold, C. Ludwig, and T. Hwa, Principles of gene regulation quantitatively connect DNA to RNA and proteins in bacteria, Science **378**, 10.1126/science.abk2066 (2022).

[26] X. Dai, M. Zhu, M. Warren, R. Balakrishnan, V. Patsalo, H. Okano, J. R. Williamson, K. Fredrick, Y.-P. Wang, and T. Hwa, Reduction of translating ribosomes enables escherichia coli to maintain elongation rates during slow growth, Nature Microbiology **2**, 10.1038/nmicrobiol.2016.231 (2016).

[27] L. Calabrese, J. Grilli, M. Osella, C. P. Kempes, M. C. Lagomarsino, and L. Ciandrini, Protein degradation sets the fraction of active ribosomes at vanishing growth, PLOS Computational Biology **18**, e1010059 (2022).

[28] W.-H. Lin, E. Kussell, L.-S. Young, and C. Jacobs-Wagner, Origin of exponential growth in nonlinear reaction networks, Proceedings of the National Academy of Sciences **117**, 27795 (2020).

[29] E. R. Oldewurtel, Y. Kitahara, and S. van Teeffelen, Robust surface-to-mass coupling and turgor-dependent cell width determine bacterial dry-mass density, Proceedings of the National Academy of Sciences **118**, 10.1073/pnas.2021416118 (2021).

[30] A. P. Gasch, P. T. Spellman, C. M. Kao, O. Carmel-Harel, M. B. Eisen, G. Storz, D. Botstein, and P. O. Brown, Genomic expression programs in the response of yeast cells to environmental changes, Molecular Biology of the Cell **11**, 4241 (2000).

[31] T. Schmelzle and M. N. Hall, TOR, a central controller of cell growth, Cell **103**, 253 (2000).

[32] A. G. Hinnebusch, Translational regulation of gcn4 and the general amino acid control of yeast, Annual Review of Microbiology **59**, 407 (2005).

[33] Y. J. Joo, J.-H. Kim, U.-B. Kang, M.-H. Yu, and J. Kim, Gcn4p-mediated transcriptional repression of ribosomal protein genes under amino-acid starvation, The EMBO Journal **30**, 859 (2010).

[34] N. Giordano, F. Mairet, J.-L. Gouzé, J. Geiselmann, and H. de Jong, Dynamical allocation of cellular resources as an optimal control problem: Novel insights into microbial growth strategies, PLOS Computational Biology **12**, e1004802 (2016).

[35] X. Dai and M. Zhu, Coupling of ribosome synthesis and translational capacity with cell growth, Trends in Biochemical Sciences **45**, 681 (2020).

[36] J. A. Bernstein, P.-H. Lin, S. N. Cohen, and S. Lin-Chao, Global analysis of escherichia coli RNA degradosome function using DNA microarrays, Proceedings of the National Academy of Sciences **101**, 2758 (2004).

[37] C. You, H. Okano, S. Hui, Z. Zhang, M. Kim, C. W. Gunderson, Y.-P. Wang, P. Lenz, D. Yan, and T. Hwa, Coordination of bacterial proteome with metabolism by cyclic AMP signalling, Nature **500**, 301 (2013).

[38] N. M. Belliveau, G. Chure, C. L. Hueschen, H. G. Garcia, J. Kondev, D. S. Fisher, J. A. Theriot, and R. Phillips, Fundamental limits on the rate of bacterial growth and their influence on proteomic composition, Cell Systems **12**, 924 (2021).

[39] H. Bremer and P. Dennis, Feedback control of ribosome function in escherichia coli, Biochimie **90**, 493 (2008).

[40] L. Calabrese, L. Ciandrini, and M. C. Lagomarsino, How total mRNA influences cell growth, biorxiv 10.1101/2023.03.17.533181 (2023).

[41] M. Nomura, J. L. Yates, D. Dean, and L. E. Post, Feedback regulation of ribosomal protein gene expression in escherichia coli: structural homology of ribosomal RNA and ribosomal protein MRNA., Proceedings of the National Academy of Sciences **77**, 7084 (1980).

[42] B. J. Paul, W. Ross, T. Gaal, and R. L. Gourse, rRNA transcription in escherichia coli, Annual Review of Genetics **38**, 749 (2004).

[43] J. J. Lemke, P. Sanchez-Vazquez, H. L. Burgos, G. Hedberg, W. Ross, and R. L. Gourse, Direct regulation of escherichia coli ribosomal protein promoters by the transcription factors ppGpp and DksA, Proceedings of the National Academy of Sciences **108**, 5712 (2011).

[44] P. P. Dennis, M. Ehrenberg, and H. Bremer, Control of rRNA synthesis in escherichia coli: a systems biology approach, Microbiology and Molecular Biology Reviews **68**, 639 (2004).

[45] S. Kostinski and S. Reuveni, Ribosome composition maximizes cellular growth rates in e. coli, Physical Review Letters **125**, 028103 (2020).

[46] A. Roy, D. Goberman, and R. Pugatch, A unifying autocatalytic network-based framework for bacterial growth laws, Proceedings of the National Academy of Sciences **118**, 10.1073/pnas.2107829118 (2021).

[47] M. J. Pine, Regulation of intracellular proteolysis in escherichia coli, Journal of Bacteriology **115**, 107 (1973).

[48] M. Scott and T. Hwa, Shaping bacterial gene expression by physiological and proteome allocation constraints, Nature Reviews Microbiology **21**, 327 (2022).

[49] U. Alon, *An introduction to systems biology : design principles of biological circuits* (CRC Press, 2019).

[50] N. Rosenfeld, M. B. Elowitz, and U. Alon, Negative autoregulation speeds the response times of transcription networks, Journal of Molecular Biology **323**, 785 (2002).

[51] J. Wang, Y. Jiang, M. Vincent, Y. Sun, H. Yu, J. Wang, Q. Bao, H. Kong, and S. Hu, Complete genome sequence of bacteriophage t5, Virology **332**, 45 (2005).

[52] M. Maeda, T. Shimada, and A. Ishihama, Strength and regulation of seven rRNA promoters in escherichia coli, PLOS ONE **10**, e0144697 (2015).

[53] X. Zhang and H. Bremer, Control of the escherichia coli rrnB p1 promoter strength by ppGpp, Journal of Biological Chemistry **270**, 11181 (1995).

[54] B. Gummesson, M. Lovmar, and T. Nyström, A proximal promoter element required for positive transcriptional control by guanosine tetraphosphate and DksA protein during the stringent response, Journal of Biological Chemistry **288**, 21055 (2013).

[55] Q. Zhang, E. Brambilla, R. Li, H. Shi, M. C. Lagomarsino, and B. Sclavi, A decrease in transcription capacity limits growth rate upon translation inhibition, mSystems **5**, 10.1128/msystems.00575-20 (2020).

[56] L. Rao, W. Ross, J. Appleman, T. Gaal, S. Leirmo, P. J. Schlax, M. Record, and R. L. Gourse, Factor independent activation of rrnB p1, Journal of Molecular Biology **235**, 1421 (1994).

[57] P. Hengen, Information analysis of fis binding sites, Nucleic Acids Research **25**, 4994 (1997).

[58] W. Ross, J. F. Thompson, J. T. Newlands, and R. L. Gourse, E.coli fis protein activates ribosomal RNA transcription in vitro and in vivo., The EMBO Journal **9**, 3733 (1990).

[59] C. A. Ball, R. Osuna, K. C. Ferguson, and R. C. Johnson, Dramatic changes in fis levels upon nutrient upshift in escherichia coli, Journal of Bacteriology **174**, 8043 (1992).

[60] M. M. C. A. Panlilio, *Non-equilibrium growth laws: growth and gene expression dynamics in Escherichia coli under a nutritional upshift*, Ph.D. thesis, Department of Physics, University of Cambridge (September 2020).

[61] K. Gausing, *Regulation of ribosomal synthesys in E.coli. From the book "Ribosomes : structure, function and genetics. Proceedings of the 9th Steenbock symposium held at the university of Wisconsin-Madison, July 5-8, 1979"*, edited by G. Chambliss (Baltimora : University park press, 1980).