Dysregulated neuroimmune interactions and sustained type I interferon signaling after human immunodeficiency virus type 1 infection of human iPSC derived microglia and cerebral organoids

Andrew J. Boreland[1,2], Alessandro C. Stillitano[1], Hsin-Ching Lin[1], Yara Abbo[1], Ronald P. Hart[3], Peng Jiang[3], Zhiping P. Pang[1,2*], Arnold B. Rabson[1,4*]

[1] Child Health Institute of New Jersey, Rutgers Robert Wood Johnson Medical School, New Brunswick, NJ

[2] Department of Neuroscience, Rutgers Robert Wood Johnson Medical School, New Brunswick, NJ

[3] Department of Cell Biology and Neuroscience, Rutgers University, Piscataway, NJ

[4] Departments of Pharmacology, Pathology & Laboratory Medicine, and Pediatrics, Rutgers Robert Wood Johnson Medical School, New Brunswick, NJ

*Correspondence: zhiping.pang@rutgers.edu and rabsonab@rwjms.rutgers.edu

Highlights of the work:

• HIV-1 productively infects iPSC-derived microglia and triggers inflammatory activation.

• HIV-1 infection of microglia results in sustained type I interferon signaling.

• Microglia infected by HIV-1 incorporate into sliced neocortical organoids with persistent

type I interferon signaling and disease risk gene expression.

ABSTRACT

Human immunodeficiency virus type-1 (HIV-1) associated neurocognitive disorder (HAND)

affects up to half of HIV-1 positive patients with long term neurological consequences, including

dementia. There are no effective therapeutics for HAND because the pathophysiology of HIV-1

induced glial and neuronal functional deficits in humans remains enigmatic. To bridge this

knowledge gap, we established a model simulating HIV-1 infection in the central nervous system

using human induced pluripotent stem cell (iPSC) derived microglia combined with sliced

neocortical organoids. Upon incubation with two replication-competent macrophage-tropic HIV-1

strains (JRFL and YU2), we observed that microglia not only became productively infected but

also exhibited inflammatory activation. RNA sequencing revealed a significant and sustained

activation of type I interferon signaling pathways. Incorporating microglia into sliced neocortical

organoids extended the effects of aberrant type I interferon signaling in a human neural context.

Collectively, our results illuminate the role of persistent type I interferon signaling in HIV-1 infected

microglial in a human neural model, suggesting its potential significance in the pathogenesis of

HAND.

INTRODUCTION

The global burden of human immunodeficiency virus-1 (HIV-1) infection remains profound, with neurocognitive complications manifesting in a significant proportion of infected individuals despite the remarkable success of combined antiretroviral therapy (cART) in reducing morbidity and mortality. Up to 55% of HIV-1 positive people experience neurological complications that may develop into HIV associated neurocognitive disorder (HAND) (Simioni et al. 2010; Sacktor et al. 2016; Eggers et al. 2017; Sacktor 2018). This debilitating disorder manifests across a spectrum of cognitive, motor, affective, and behavioral impairments, which can severely impact the quality of life for affected individuals, emphasizing an urgent need for a better understanding of its underlying mechanisms (Lew et al. 2018; Israel et al. 2019). The etiology and neuropathogenesis of HAND remain enigmatic, as there has been no evidence of neuronal HIV-1 infection. Decades of research suggest that HAND development is caused by a combination of both direct and indirect effects involving viral proteins, inflammatory cytokines and chemokines, and dysregulated neuroimmune interactions (Koenig et al. 1986; Gendelman et al. 1994; Rappaport et al. 1999; González-Scarano and Martín-García 2005; Kraft-Terry et al. 2009; Lindl et al. 2010; Burdo, Lackner, and Williams 2013; Alvarez-Carbonell et al. 2020). Interestingly, clinical symptoms of HAND correlate more closely with cytokine production, neuronal pathology, decreased synaptic density, and increased microglia than with central nervous system (CNS) viral load (Kaul, Garden, and Lipton 2001; Kaul et al. 2005). Currently there is a lack of mechanistic understanding due to a need for better experimental models that can address these questions, especially regarding neuroimmune interactions.

A pivotal component of neuroimmune interactions is the innate immune response, including the production and signaling of interferons, which have recently been implicated in the neuropathology of various neurological diseases [Reviewed in (Hofer and Campbell 2013; McGlasson et al. 2015)]. Interferons and their downstream signaling through interferon stimulated genes (ISGs) are primarily an innate immune response to viral infection [reviewed in (Platanias

2005)]. However, recent literature suggests a role for dysregulated type I interferon signaling in

neurological diseases such as Alzheimer's disease (Roy et al. 2020; Roy et al. 2022), COVID-19

related brain fog (Lee and Shin 2020; Phetsouphanh et al. 2022), aging associated dementia

(Baruch et al. 2014), and HIV associated dementia (Rho et al. 1995; Krivine et al. 1999; Perrella

et al. 2001; Anderson et al. 2017; Solomon et al. 2020). Clinically, type I interferons have been

shown to be a driver of cognitive impairment both in diseased states and when administered

exogenously (Merimsky et al. 1990). Indeed, many studies using interferon treatment for various

severe cancers and chronic viral hepatitis C have reported that patients receiving interferon-alpha

displayed cognitive, mood, and behavioral deficits that arose and worsened with prolonged,

multiweek treatment regimens (Christina, Randall, and Arthur 1991; Pavol et al. 1995; Capuron,

Ravaud, and Dantzer 2001; Capuron et al. 2002; Wichers et al. 2005; Raison et al. 2005). Taken

together, these studies indicate a role for aberrant type I interferon signaling in compromising

neuronal and glial function, leading to cognitive deficits and degeneration.

Microglia, as the resident immune cells of the CNS and the primary CNS target of HIV-1

infection, contribute significantly to the pathophysiology of HAND, largely through their secretory

and functional dysregulation. Under homeostatic conditions, microglia play important roles in

synaptic pruning, phagocytic functions, immune surveillance, and extracellular matrix remodeling

(Paolicelli et al. 2011; Casano and Peri 2015). When challenged, microglial activation can be

either neurotoxic, producing proinflammatory cytokines and chemokines, or neurotrophic,

producing anti-inflammatory cytokines (Durafourt et al. 2012). Microglia exhibit a heterogeneous

spectrum of activation and disease-associated states with unique properties and consequences

on neuronal functions (Gosselin et al. 2017; Song and Colonna 2018; Hammond et al. 2019).

Since HIV-1 specifically infects humans, the ability of traditional animal model organisms

to replicate the complex disease processes of HAND is inherently limited. HIV-1-Simian

immunodeficiency virus chimeras (SHIVs) have been studied in monkey models. However, these

models are limited in their availability and utility for molecular analyses [reviewed in (Byrnes et al.

2022; Moretti et al. 2021)]. Recent advances in stem cell technologies have opened up new

possibilities for modeling and studying HAND, particularly the development of human iPSC

derived neurons, astrocytes, and microglia capable of mimicking complex in vivo neuroimmune

interactions when grown in three dimensional cerebral organoid cultures (Abud et al. 2017; Ormel

et al. 2018; Xu et al. 2021). Cerebral organoids offer a powerful and physiologically relevant

platform to study complex neuroimmune interactions, but traditional organoid differentiation

protocols do not produce microglia. Recent studies have demonstrated the feasibility of infecting

iPSC-derived macrophages (Vaughan-Jackson et al. 2021) and microglia (Rai et al. 2020) with

HIV-1. Moreover, these infected microglia have been effectively cocultured in various model

systems, including human stem cell derived neural tricultures (Alvarez-Carbonell et al. 2020; Ryan

et al. 2020), primary neuron progenitor cell derived organoids (dos Reis et al. 2020), iPSC-derived

cerebral organoids (Gumbs et al. 2022), and a humanized mouse model (Min et al. 2023).

In this study, we hypothesized that by infecting iPSC-derived microglia with replication

competent HIV-1 and integrating them into sliced neocortical organoids, we could recapitulate

neuroimmune dysregulation related to HAND pathology. We demonstrate that HIV-1 infection of

iPSC-derived microglia leads to persistent activation of type I interferon signaling and microglial

activation in both microglial monocultures and in the context of cerebral organoids.

RESULTS

Human microglia are efficiently generated from iPSCs

To induce microglial differentiation, we used three iPSC lines and employed previous protocols

(Haenseler et al. 2017; Xu et al. 2021). To mimic in vivo microglial development from the yolk sac,

iPSC embryoid bodies were patterned toward a mesodermal yolk sac lineage that generates

primitive macrophage precursors (PMPs) after 3 weeks of culture (**Figure 1A**). The identity of the

PMPs was confirmed by immunocytochemistry against CD235, a marker for yolk sac (Claes, Van

den Daele, and Verfaillie 2018), and CD43, a marker for hematopoietic progenitor like cells

(Vodyanik, Thomson, and Slukvin 2006) (**Figure 1B**). Upon further maturation under the influence

of a colony stimulating factor 1 receptor (CSF1R) antagonist, interleukin 34 (IL-34), and

granulocyte maturation factor (GM-CSF), PMPs exhibit a more pronounced microglial phenotype.

This was evidenced by a significant increase in mRNA expression of the microglial markers

ionized calcium binding molecule 1 (IBA1) and purinergic receptor P2Y12 across all three cell

lines (**Figure 1C,** biological replicates of each cell line shown in **Figure S1**). Supporting these

gene expression results, we noted robust immunocytochemical staining of the microglial markers

IBA1, P2Y12, CX3CR1, transmembrane protein 119 (TMEM119), and transcription factor PU.1 in

maturing microglia from these same lines (**Figure 1D**). Upon coculture of microglia with iPSC-

derived neurons (Xu et al. 2021), we found that microglia display a more elaborate morphology

(**Figure 1E**). Furthermore, costaining with the neuronal dendritic marker microtubule-associated

protein 2 (MAP2) and the presynaptic marker synapsin 1 (SYN1) revealed that microglial

processes interact with dendrites and colocalize with synaptic puncta, indicating putative

microglia-synapse interactions (**Figure 1F**).

Given the tropism of HIV-1 for CD4 receptor expressing cells and the requirement of CCR5

coreceptor expression for infection of monocyte/macrophage related cell lineages, we assessed

the gene expression of *CD4* and *CCR5* in iPSC-derived microglia. Results showed significantly

increased mRNA expression of *CD4* and *CCR5* in microglia differentiated from all three iPSC

lines (**Figure 1G,** biological replicates of each cell line shown in **Figure S1**). These results

demonstrate that microglia can be successfully and reproducibly generated from iPSCs. Crucially,

these cells express appropriate markers for microglial identity and infection by HIV-1.

iPSC derived microglia are productively infected by HIV-1 and exhibit inflammatory

activation

Having confirmed the generation and maturation of iPSC-derived microglia, we next determined

whether these cells could sustain a productive infection by replication competent, macrophage

tropic HIV-1. To this end, we incubated male 03SF iPSC-derived microglia with JRFL HIV (2

ng/ml), an R5 tropic HIV-1 strain isolated from the cerebral spinal fluid (CSF) of a patient with

severe HAND (Koyanagi et al. 1987; Merrill et al. 1992). Compared to uninfected controls, HIV-

1-infected microglia robustly expressed HIV-1 *gag/pol* RNA within 6-9 days after infection (Yu et

al. 1996) (**Figure 2A**). As a key indicator of productive HIV-1 infection, the production of secreted

p24 was measured (Chargelegue and O'Toole 1992; Marodon et al. 1999). Using ELISA to detect

secreted p24 in the culture supernatant, we observed increased protein production starting 1

week post infection, peaking around day 16, and then maintaining consistent levels for more than

2 weeks (**Figure 2B**). Finally, we found that after just 4 days post infection, circular polymerase

chain reaction (PCR) products from both long terminal repeats (LTR) were detected in HIV-1

incubated cultures indicating the presence of active viral replication (Butler, Hansen, and

Bushman 2001) (**Figure S2**).

After establishing the infectability of iPSC microglia with HIV-1 JRFL, we then asked

whether HIV-1 infection elicits differential expression of inflammatory and canonical microglial

mRNAs using reverse transcription–quantitative PCR (RT-qPCR). In this second experiment

found we found HIV-1 infection augmented the expression of the inflammation-associated genes

interleukin 1ß (IL1ß, on day 16) and tumor necrosis factor alpha (TNFα), while the expression of

interleukin 6 (IL6) displayed no change (**Figure 2C**). We then examined the mRNA expression of

the chemokines CCL2 (monocyte chemoattractant protein 1, MCP-1), CCL3 (macrophage

inflammatory protein 1 alpha, MIP-1), and CCL5 (RANTES). These three chemokines play

important roles in immune cell recruitment and migration to sites of inflammation and infection

and have been found to be elevated in cognitively impaired individuals with HIV-1 (Letendre,

Lanier, and McCutchan 1999). A significant elevation in the mRNA expression of CCL2 and CCL3

was observed by day 16 post-infection, whereas the expression of CCL5 remained unchanged

(**Figure 2D**). To assess whether microglial identity was altered by HIV-1 infection, we assessed

the mRNA expression of the canonical microglial genes IBA1 and P2Y12. While IBA1 remained

stable, P2Y12 exhibited significantly increased expression by day 23 post infection (**Figure 2E**).

Next, we asked how HIV-1 infection might alter microglial chemokine and cytokine production.

Although the levels of most cytokines were below the detection threshold (**Table S1**), we noted a

significant surge in the secretion of chemokines M-CSF and interferon inducible protein 10 (IP-

10/CXCL10) following HIV-1 infection in the same supernatant used in figure 2B (**Figure 2F**).

Finally, to visualize HIV-1 infected cells, we detected HIV-1 p24 protein using a biotin-streptavidin

amplification protocol (Eugenin and Berman 2016). On day 13 post infection with JRFL, p24

positive microglia were found (**Figure 2G**). Collectively, our findings suggest that iPSC-derived

microglia can sustain an R5 tropic HIV-1 infection and exhibit an inflammatory activation profile

consistent with previous studies.

To confirm that HIV-1 JRFL infection of human iPSC-derived microglia resulted in

productive infection and inflammatory activation, we chose a female iPSC line, CD06, and

infected it with the JRFL strain as well as a second R5 tropic HIV-1 strain, YU2 (Li et al. 1991),

isolated from the CSF of a different patient with severe HAND. We confirmed infection by RT–

qPCR of viral *gag-pol* RNA, which exhibited significantly increased expression for both strains by

day 16 post-infection, with a subsequent decrease (**Figure 3A**). Using ELISA to detect secreted

p24 revealed productive infection for both viral strains (**Figure 3B**). We then assessed the

expression of key inflammatory markers and microglial core mRNAs by qPCR. Both JRFL and

YU2 infections led to a significant increase in IL1β expression compared to uninfected microglia

(**Figure 3C**). Similarly, TNFα was significantly upregulated in the first two weeks of infection but

decreased as the productive infection waned (**Figure 3C**). We also observed increased

expression of the chemokines CCL2 and CCL3 by days 16 and 23 post infection with both strains

(**Figure 3D**). This upregulation then decreased by day 27, correlating with the reduction of

productive infection as indicated by the p24 ELISA (**Figure 3A**). In comparison to the unaltered

IBA1 expression we found previously, HIV-1 led to a significant reduction in IBA1 expression in

CD06 infected microglia (**Figure 3E**). Similar to 03SF infected microglia, CD06 had significant

increased expression of P2RY12 (**Figure 3E**). Together, these results demonstrate infectability

of human iPSC-derived microglia with multiple, macrophage tropic HIV-1 strains and their ability

to elicit an inflammatory response.

Type I interferon stimulated gene expression and sustained activation following HIV-1

infection

To investigate the transcriptional landscape of HIV-1 infected microglia, we performed bulk RNA

sequencing on male 03SF iPSC-derived microglia infected by HIV-1 JRFL day 16 post infection.

Principal component analysis (PCA) revealed clear separation of the infected vs mock infected

samples (**Figure S2**). Differential gene analysis revealed differentially expressed genes (DEGs)

including several classic ISGs, such as IFIT3, MX2, ISG15, RSAD2 and others (**Figure 4A**) (**Table**

S2). Gene set enrichment analysis (GSEA) confirmed a significant enrichment of genes related

to the interferon alpha response (**Table S3**) (**Figure 4B**). Gene ontology analysis identified

activation of innate immune responses and several related terms in the biological process

component (**Figure 4C, D** and **Figure S3**). The upregulated DEG GO set includes pathways

falling into the overall signaling clusters related to interferon signaling and positive regulation of

antiviral biotic, interferon alpha/beta, defense against virus, cytokine mediated and interferon

mediated pathways, and negative regulation of viral processes. Among the downregulated DEG

GO clusters are pathways related to positive regulation of antigen processing, extracellular

organization, granulocyte/leukocyte migration, chemokine-mediated signaling, and spindle

assembly checkpoints. Importantly, HIV-1 infection was found to lead to significant induction of

many ISGs, including ISG15, IRF7, MX1, OAS1, OAS2, OAS3, STAT1, CXCL10, and IFITM3

(**Figure 4A, E**). **Figure 4F** shows the expression of various genes encoding SIGLEC proteins that

are responsible for regulating microglial cell–cell interactions [Reviewed in (Crocker, Paulson, and

Varki 2007)]. Consistent with previous studies, we found that HIV-1 infection causes significant

upregulation of SIGLEC1, which has been implicated in the direct cell–cell transmission of HIV

(Izquierdo-Useros et al. 2012; Zou et al. 2011). However, we found that SIGLEC11 and

SIGLEC16 were significantly downregulated after infection.

We then validated the expression of a subset of the ISGs identified in the sequencing

dataset by RT–qPCR of JRFL infected 03SF iPSC-derived microglia. We found significant

upregulation of CXCL10, RSAD2, ISG15, and IDO1 (**Figure 4G**). Elevated expression of these

genes persisted over an additional week of infection on day 23 (**Figure 4G**). Additionally, we

performed RT–qPCR on CD06 iPSC-derived microglia infected with either JRFL or YU2 HIV

strains. We also found increased expression of CXCL10 and RSAD2 in these HIV-infected cells;

however, only the day 16 YU2 condition reached significance (**Figure 4H**). Comparison of Figure

4G and 4H indicates potential differences with infecting different cell lines. Together, these data

support the conclusion that HIV-1 infection induces sustained type I interferon signaling in iPSC-

derived microglia and alters the gene regulation of many microglial biological processes including

interferon signaling.

HIV-1 infected microglia incorporate into cerebral organoids

After characterizing the response of isolated human iPSC-derived microglia to HIV-1 infection we

then investigated whether HIV-1 infected microglia could incorporate into iPSC-derived cerebral

organoids and whether interferon signaling was similarly upregulated in microglia and potentially

other neural cell types. Cerebral organoids were grown using modified, published protocols

(Lancaster and Knoblich 2014; Paşca et al. 2015; Sloan et al. 2018; Yoon et al. 2019; Qian et al.

2020; Xu et al. 2021). To create microglia-containing organoids, we cocultured human microglia

with sliced neocortical organoids (SNO), prepared as described previously (Qian et al. 2020).

SNOs exhibit enhanced neurogenesis and laminar expansion (Qian et al. 2020). We found that

microglia exhibit the ability to incorporate with organoid slices to form human microglia containing

sliced neocortical organoids (mSNOs). **Figure 5A** shows mSNO containing neurons (MAP2$^+$),

astrocytes (GFAP$^+$), and microglia (IBA1$^+$). These neuroimmune organoids provide a model

system for studying the impact of HIV-1 infection on the physiology, localization, and interactions

of these three cell types.

To create an HIV-1 infected human cerebral organoid, we infected 03SF iPSC-derived

microglia with JRFL HIV. Following 2 weeks in culture, infected or noninfected microglia were

added to 400 μM organoid slices and cultured for 2 additional weeks. On day 14 post coculture

with microglia, mSNOs were fixed for immunohistochemistry or total RNA was extracted for RT–

qPCR analysis. HIV-1-infected microglia (p24$^+$, IBA1$^+$) were detected in a mSNO (**Figure 5B&C**).

Interestingly, neurons (MAP2$^+$) appeared to be sparser in regions of HIV-1-infected microglia

(**Figure 5B** "arrows"). These regions had scattered MAP2$^+$ processes (**Figure 5B1**) compared

with the denser MAP2 staining throughout the bulk of the organoid. HIV-1 *gag/pol* RNA exhibited

significant expression in mSNOs containing HIV-1 infected microglia (**Figure 5D**). To provide

evidence of productive infection in mSNOs, we identified secreted p24 only in the media of

organoid cultures that had received HIV-1 infected microglia (**Figure 5E**).

Because we observed significant and sustained interferon signaling in microglial

monocultures, we evaluated mRNA expression of a panel of ISGs (ISG15, RSAD2, IFITM3, MX1,

IFI44, IFI27). We found that ISG15 mRNA was significantly upregulated and RSAD2 also showed

an increasing trend of expression in mSNOs containing HIV-1 infected microglia (**Figure 5F**).

IFITM3 has been touted as a marker for interferon-responsive microglia and has been shown to

play a newly discovered role in amyloid precursor protein processing to amyloid beta (Hur et al.

2020). Interestingly, we found significant upregulation of IFITM3 mRNA in mSNOs containing

HIV-1 infected microglia (**Figure 5F).** We also evaluated the expression of MX1, IFI27, and IFI44,

which, along with ISG15 and RSAD2, were among the top 10 persistently upregulated genes in

a recent study on two cohorts of 22 HIV-1 seroconverts from Uganda and 22 from South Africa

(Mackelprang et al. 2023). Expression of MX1, IFI27, and IFI44 appeared to be increased but

failed to reach significance. Especially considering the small fraction of HIV-1 infected microglia

in the cultured mSNO (**Figure 5B**), these increases suggest that HIV-1 infected microglia are

associated with increased interferon signaling in the organoids and may have indirect

consequences for neuronal viability and inflammatory responses in other cell types.

DISCUSSION:

In this study, we used iPSC-derived human microglia and microglia containing cortical organoids

to model HIV-1 related neuroimmune dysregulation and pathology. In this model, iPSC-derived

microglia were shown to be infected with two strains of replication competent HIV-1, JRFL and

YU2, and exhibited inflammatory activation, as shown by gene expression changes and

cytokine/chemokine secretion (**Figures 2 & 3**). Importantly, we found persistent activation of type

I interferon response pathways and downstream interferon stimulated genes (ISGs), which may

have important implications for understanding the pathogenesis of HAND.

In comparison to the recently developed primary human neural progenitor cell-derived

organoid HIV-1 model (dos Reis et al. 2020) and the 2D neuron, astrocyte, and microglia

"triculture" HIV-1 model (Ryan et al. 2020), an iPSC-derived human microglia containing sliced

neocortical organoid model has distinct advantages. Organoids recapitulate 3D cell-cell and cell-

matrix interactions more faithfully than 2D cultures and develop complex tissue architecture

(Lancaster et al. 2013; Paşca et al. 2015; Birey et al. 2017; Quadrato et al. 2017; Velasco et al.

2019); thus, organoids are well suited to studying aspects of altered brain anatomy and physiology

seen in individuals with *HAND* (Di Sclafani et al. 1997; Clifford et al. 2017; Israel et al. 2019).

While others have also generated microglia containing organoids using a protocol in which

microglia innately develop within organoids, this system is associated with a high degree of

heterogeneous cell types (Gumbs et al. 2022), which could make dissection of the relative

contributions and effects on specific cell types more complex. In our method, we have greater

freedom to manipulate the timing, proportion, and infection stage of microglia addition in sliced

neocortical organoids. While some previous studies used pseudotyped virus (dos Reis et al.

2020), our study examined infection using two macrophage tropic, replication competent strains

of HIV-1. Our findings, showing induction of inflammatory gene expression associated with active

viral expression align with recent studies on brain samples from HIV-1[+] patients from the National

NeuroAIDS Tissue Consortium (NNTC). In that study, HIV-1 RNA load was found to correlate with

increased interferon signaling and inflammatory gene sets while inversely correlating with

neuronal and synaptic gene sets, suggesting neuronal pathology (Sanna et al. 2021).

Microglia secrete diverse cytokines and chemokines implicated in HIV-1 neuropathology

(Hanisch 2002; Ramesh, MacLean, and Philipp 2013; Chen et al. 2017). Cytokine/chemokine

analysis of the supernatant of HIV-1 infected microglia identified the secretion of M-CSF and IP-

10/CXCL10 (**Figure 2F**). M-CSF overexpression has been linked to microglial proliferation,

increased phagocytosis, cytokine secretion, and inflammatory responses in both Alzheimer's

disease (AD) and HAND, as well as increased microglial susceptibility to HIV-1 infection and

increased virus replication (Gallo et al. 1990; Kalter et al. 1991; Kutza et al. 2000; Mitrasinovic et

al. 2001; Kutza et al. 2002; Smith et al. 2013). IP-10/CXCL10 is a chemotactic chemokine that

recruits leukocytes, T cells, and monocytes while also being neurotoxic and strongly correlated

with HIV-1 related disease progression (van Marle et al. 2004; Jiao et al. 2012; Lei et al. 2019).

Clinical studies have shown elevated IP-10/CXCL10 levels (~200-1000 pg/ml) in the cerebral

spinal fluid of HIV-1[+] individuals (Kolb et al. 1999; Peluso et al. 2017); these levels are strikingly

similar to our observations (**Figure 2F**). CXCL10/IP-10 has been shown to be induced by HIV-1

proteins gp120, Nef, and Tat (Kutsch et al. 2000; Asensio et al. 2001; van Marle et al. 2004; Lei

et al. 2019). Importantly, IP-10/CXCL10 and RANTES (CCL5) were the strongest predictors of

neurocognitive impairment in a screen of 139 HIV-1[+] individuals (Ruhanya et al. 2021), similar to

findings from earlier studies (Valverde-Villegas et al. 2018; Burlacu et al. 2020). HIV-1 brain

infection leads to an abnormal inflammatory state and dysregulated cytokine/chemokine

production (Kaul, Garden, and Lipton 2001; González-Scarano and Martín-García 2005; Burdo,

Lackner, and Williams 2013). Despite this knowledge, there remains a large gap in our

understanding of how CXCL10 and other cytokines/chemokines associated with CNS HIV-1

infection alter aspects of neuronal excitability, synaptic transmission, and inward and outward

currents vital to healthy neuronal physiology.

A striking finding in our study was the persistent activation of interferon response genes.

Induction of these pathways is known to be a hallmark of HIV-1 infection in monocytes and

macrophages, much more prominently than in T cells (Berg et al. 2014; Aso et al. 2019; Elsner et

al. 2020; Jeremiah et al. 2023). Thus, it is not surprising to have seen this strong response in

microglia, which share many physiologic properties with macrophages [reviewed in (Li and Barres

2018)]. Most studies suggest that the interferon response is relatively transient; thus, the

persistent expression of at least a subset of ISGs may be a relatively unique feature of microglial

HIV-1 infection that may have important pathophysiologic consequences. Increasing evidence

suggests a role for type I interferons in cognitive-related deficits seen in patients with various

neurodegenerative diseases. Using single-cell sequencing, researchers have dissected a unique

sub-population of interferon-responsive microglia that may potentiate dysregulated immune

interactions in the form of aberrant pruning of synapses, altered phagocytic function, and

dysregulated cytokine/chemokine production (Jin et al. 2022). Constitutive overexpression of type

I interferons seen in the "interferonopathies" can be associated with profound CNS disease

[reviewed in (Goldmann, Blank, and Prinz 2016), and (Blank and Prinz 2017)](Wiley, Steinman,

and Wang 2023). Patients receiving type I interferons therapeutically may experience cognitive

impairment and neurological symptoms (Christina, Randall, and Arthur 1991; Pavol et al. 1995;

Capuron, Ravaud, and Dantzer 2001; Capuron et al. 2002; Wichers et al. 2005; Raison et al.

2005).

Whether interferon induction inhibits or enhances HIV infection and latency in microglia

remains to be understood. Recent studies have highlighted that interferon can inhibit HIV infection

of primary microglia but less so of iPSC-derived microglia, as well as macrophages (Nasr et al.

2017; Woodburn et al. 2022). Interestingly, interferon may promote latency in macrophages

(Dickey et al. 2022). An important feature of HIV infection of microglia identified in previous in vitro

studies has been the rapid induction of viral latency (Alvarez-Carbonell et al. 2019; Sreeram et

al. 2022). Our cerebral organoid model will provide an interesting system to examine the induction

of HIV latency, its reactivation, and its effects on both adjacent and more distant neural cells.

An important future direction, facilitated by the development of this model system, will be

experiments aimed at understanding the functional consequences that HIV infection,

dysregulated interferon signaling, inflammatory activity, and changes in other important genes

expressed in the CNS, may have on neuronal synaptic transmission. A number of ISGs have

predicted downstream effects that could trigger or enhance neurodegeneration. For example,

indoleamine 2,3-dioxygenase 1 (IDO) is a key enzyme in tryptophan catabolism, leading to the

formation of kynurenic acid, an antagonist of NMDA receptors (Schwarcz and Stone 2017). This

pathway and its products, such as the neurotoxin and NMDAR antagonist quinolinic acid, have

been previously implicated in neurodegenerative diseases, including Huntington's, Alzheimer's

and Parkinson's disease (Malpass 2011; Maddison and Giorgini 2015).

Overall, the HIV-1 mSNO model presents an effective in vitro platform for investigating the

underlying molecular and cellular mechanisms associated with HIV-1 and human neural cell

interactions. **Our results predict that interferon stimulated genes likely play a role in altering**

neuronal function and contributing to HAND.

ACKNOWLEDGEMENTS

This work was supported in part by grants from the NIH (R21NS120806), a Rutgers Busch

Biomedical Grant, a Rutgers Brain Health Institute Pilot Grant, and the Robert Wood Johnson

Foundation (Grant #74260). AJB was supported by NIH NIGMS T32GM8339 and NIH NCATS

TL1TR003019.

AUTHOR CONTRIBUTIONS

AJB, ZPP, PJ, RPH, and ABR conceived the study. AJB designed the experiments and

interpreted the data. AJB performed most of the experiments with technical assistance from ACS,

HL, and YA. ACS and AJB performed qPCR. HL and AJB performed ELISA. AJB performed

immunocytochemistry, immunohistochemistry, and confocal imaging. AJB and HS performed viral

infections. RPH and AJB performed computational analysis of RNA-seq data. AJB prepared the

figures and wrote the manuscript with input from all coauthors.

COMPETING FINANCIAL INTERESTS

The authors declare no competing financial interests.

METHODS

Human induced pluripotent stem cell maintenance

Three human iPSC lines were used in this study (male 03SF iPSC line; female CD06 iPSC line;

male CD52 iPSC line). Human iPSCs were cultured in stem cell medium consisting of MACs iPS-

Brew XF complete media. The culture medium was changed daily. Human iPSCs were passaged

one to two times per week onto new plates coated with Matrigel. Human iPSCs were detached

with Accutase treatment for 5 min and centrifuged at 120 rcf for 5 min. The pellet was resuspended

in stem cell medium plus Y-27632 (10 µM). The iPSCs used in this study were below passage

50. All studies were performed with protocols approved by Rutgers University.

Generation of microglia

Primitive microglia precursors (PMPs) were generated from 3 iPSC cell lines using a previously

established protocol (Haenseler et al. 2017). iPSCs were detached with Accutase, plated at

20,000 cells per well in a 96-well low-adherence plate in yolk sac induction medium plus the rock

inhibitor Y-27632 (10 µM), and centrifuged at 1000 rpm for 3 min to aggregate the cells for

embryoid body formation. Yolk-sac induction medium consisted of MAC iPS-Brew XF complete

media plus bone morphogenetic protein 4 (BMP4, 50 ng/ml), vascular endothelial growth factor

(VEGF, 50 ng/ml), stem cell factor (SCF, 20 ng/ml), and ß-mercaptoethanol. Yolk-sac induction

media was given for the first 5 days of culture. On day 6, YS-EBs were plated in 100 mm cell

culture plates in factory media consisting of XVIVO15 supplemented with 0.5% NEAA, ß-

mercaptoethanol, macrophage colony stimulating factor (M-CSF, 100 ng/ml) and interleukin-3 (25

ng/ml). At 2-4 weeks after plating, human PMPs egress from the yolk sac lumen into the

surrounding media and are continuously produced for more than 3 months.

To mature PMPs into microglia, PMPs were transitioned to maturation media composed

of XVIVO15 supplemented with N2, Glutamax, 0.5% NEAA, interleukin-34 (IL-34, 100 ng/ml),

granulocyte macrophage colony stimulating factor (GM-CSF, 10 ng/ml, and M-CSF (25 ng/ml) for

1-2 weeks. The medium was changed every 4 days. Cells were then detached with 20 min

Accutase treatment and used for experiments.

Generation of iPSC-derived Cerebral Organoids

Cerebral organoids were generated from 1 iPSC line using adapted previously published

protocols (Qian et al. 2020). In this study, 9,000 iPSCs were cultured in low-adherence 96-well

plates to form uniform embryoid bodies. On day 1, the medium was changed to neural induction

medium composed of DMEM/F12 supplemented with 1X N2, 2 µM A83-01, and 2 µM

dorsomorphin. On days 5 and 6, half of the medium was replaced with DMEM/F12 supplemented

with 1X N2, 10 µM SB431542, and 1 µM CHIR99021. On day 7, EBs were embedded in a Growth-

Factor-Reduced Matrigel 'cookie' as previously described (Qian et al. 2020). Half of the medium

was changed every two days until day 14. On day 14, organoids were dissociated from the

Matrigel cookie and placed in low-adherence 6-well plates on an orbital shaker at 85 rpm. On day

16, organoids were fed 1:1 DMEF12: Neurobasal, 1X N2, 1X B27-RA, 0.5% NEAA, and 200 µM

L-AA. Media was replenished every 2-3 days until day 30. On day 30, the medium was changed

to 1:1 DMEF12: Neurobasal, 1X N2, 1X B27, 0.5% NEAA, 200 µM L-AA, and.25% GFR-Matrigel.

Half of the media was changed every 3-4 days.

Organoid Slicing

To create sliced neocortical organoid cultures, cerebral organoids were sectioned with a

vibratome and transferred back into in vitro cell culture. Briefly, a fresh stock of 3% low-melting

temperature agarose in PBS was prepared and kept at 39°C to remain liquid. Artificial cerebral

spinal fluid (ACSF) for sectioning contained (in mM): NaCl 125, KCl 2.5, $NaH_2PO_4 \cdot H_2O$ 1.25,

$NaHCO_3$ 25, $MgCl_2$ 1.2, $CaCl_2$ 2.5, glucose ($C_6H_{12}O_6$) 2.5, and sucrose ($C_{12}H_{22}O_{11}$) 22.5. The

ACSF was then bubbled with 5% CO_2/95% O_2 for 10 minutes before bivalent cations (in mM):

$MgCl_2$ 1.2, $CaCl_2$ 0.625. The solution was then filtered through a 0.2-micron PES filter and returned

to bubbling on ice. Cerebral organoids were then transferred to 37°C DMEF12, 200 μM L-AA, and

55 μM beta-mercaptoethanol before being embedded in 3% low-melting temperature agarose.

The agarose block containing organoids was then sectioned at 400 μM thickness. Slices were

then subjected to two washing processes in Primocin-treated media to reduce the chance of

contamination.

HIV-1 Stock Preparation

HIV-1 JRFL virus (ARP-395) and HIV-1 YU2 proviral DNA (ARP-1350) were obtained through

the NIH HIV Reagent Program, Division of AIDS, NIAID. HIV-1 JRFL was propagated in

phytohemagglutinin (PHA, 5 μg/ml)-stimulated human peripheral blood mononuclear cells.

Viruses were harvested on days 7-10 post infection. HIV-1 YU2 stocks were generated by

transfection of 293T cells with HIV-1 YU2 proviral DNA. The transfected cells were cocultivated

with PHA-PBMCs 48 h after transfection, and cell supernatants were harvested 7 days after

coculture. The YU2 HIV stocks were further propagated in PHA-PBMCs as described. The

amount of viral p24 was determined using the HIV-1 Gag p24 DuoSet ELISA (R &D System,

DY7360-05).

HIV-1 infection

PMPs were matured for 1-2 weeks in microglia maturation medium. Matured PMPs were then

incubated with 2 ng/ml p24 of either JRFL strain or YU2 strain HIV-1 (NIH AIDS Repository) for

24 hours at 300,000 cells per 100 μl of media. After 24 hours, DMEF12 was added, and the cells

were centrifuged for 3 min at 120 rcf. The cells were then resuspended in microglia maturation

media minus M-CSF and plated. Half of the media was changed every 3-6 days.

HIV-1 2-LTR circles PCR

For detection of HIV 2LTR circles, total DNA (including small nongenomic DNA) was isolated

with a DNeasy Blood and Tissue kit (Qiagen 69504). End point PCR analysis (Taq 2X Master

Mix, NEB) was performed using 300 ng DNA template and 2-LTR circle primers: forward,

MH535: 5′-AACTAGGGAACCCACTGCTTAAG-3′; reverse, MH536: 5′-

TCCACAGATCAAGGATATCTTGTC-3′ (Butler, Hansen, and Bushman 2001).

Tissue preparation

Organoids were fixed in 4% PFA overnight at 4°C. Organoids were then transferred to 15%

sucrose and 0.05% sodium azide in PBS for 24 hours, followed by another 24 hours in 30%

sucrose and 0.05% sodium azide in PBS. Organoids were then embedded in OCT freezing

medium and snap frozen on an ethanol dry ice slurry. Frozen embedded organoids were given at

least 3 hours to warm to -20°C before cryosectioning on a Leica cryostat at thicknesses between

10-30 μM. Sections were mounted on charged microscope slides (VWR) and placed on a slide

warmer at 37°C for 30 min. Slides were then stored in a slide box at -20°C until used for staining.

Immunofluorescence

For immunocytochemistry, half of the medium was left in each well with an equal amount of 4%

PFA and then added at a final concentration of 2% PFA overnight at 4°C. The next day, PFA was

removed, and the cells were washed with PBS plus 0.3 M glycine for 30 min. The cells were then

permeabilized with 0.2% Triton X-100 in PBS for 7 min. The cells were then blocked for 1 hour in

blocking buffer consisting of 5% normal goat serum (NGS), 4% bovine serum albumin (BSA),

0.05% Triton X-100, 0.05% sodium azide, and PBS.

For immunohistochemistry, cryosectioned organoid slides were warmed to 37°C for 5 min.

A hydrophobic barrier was then drawn around the specimens and allowed to dry for 5 min. PBS

plus 0.3 M glycine was then added in a liquid dome over the specimens to rehydrate for 10 min.

Sections were permeabilized with 0.5% Triton X-100, 0.3 M glycine, and PBS for 1 hr. After

permeabilization, the sections were blocked with blocking solution consisting of 5% normal goat

serum, 4% BSA, 0.05% Triton X-100, 0.05% sodium azide, and PBS for 1 hr. Primary antibodies

were diluted in blocking solution and applied to the sections overnight at 4°C. The primary

antibodies and their dilutions are summarized in the Key Resources Table. After washing with

0.05% Triton X-100 and PBS (PBST) 5 times, secondary antibodies diluted in blocking buffer

were added for 1 hr at RT. Finally, the sections were washed five times with PBST before a final

wash in deionized water and mounting with DAPI Fluoroshield (Fisher). Secondary antibodies

were Alexa Fluor 488-, 546-, 633- or 647-conjugated goat antibodies (Invitrogen) used at a 1:500

dilution. Images were taken on a Zeiss LSM700 confocal microscope.

RNA isolation

Total RNA was prepared with an RNeasy Plus kit (Qiagen) with preprocessing through

Qiashredder (Qiagen) and frozen at -80°C.

RT–qPCR

cDNA was generated using SuperScript VILO Master mix. RNA expression was measured using

SYBR green on a QuantStudio 3 (Agilent). Probes were generated using NCBI primer blast and

are listed in Table X. Two technical replicates were performed per sample. When a high standard

deviation between replicates occurred, replicates were rerun in triplicate. All expression levels

were normalized to GAPDH expression and represented as $2^{-(\Delta\Delta CT)}$. For null result values where

a ct was not achieved, a max ct value of 40 was used for statistical analysis.

Bulk RNA sequencing and analysis

We performed bulk RNA-sequencing analysis of iPSC-derived microglia. Total RNA was prepared

using an RNeasy kit (QIAGEN). Library construction and sequencing were performed by

Novogene. The libraries were subjected to 75 bp paired read sequencing using a NextSeq500

Illumina sequencer. Approximately 30–36 million paired reads were generated for each sample.

Bc12Fastq software, version 1.8.4, was used to generate Fastq files. The genome sequence was

then indexed using the rsem-prepare-reference command.

Each fastq file was trimmed and checked for quality with fastp (v. 0.12.2) and then aligned

to the UCSC hg38 human genome using HISAT2 (v.2.1.0)[71,72]. Transcript counts were extracted

using the featureCounts function of the Rsubread package.

Statistics and reproducibility. All data represent the mean ± s.e.m. When only two independent

groups were compared, significance was determined by a two-tailed unpaired t test with Welch's

correction. A p value <0.05 was considered significant. The analyses were performed in

GraphPad Prism v.9. All experiments were independently performed at least three times with

similar results.

References:

Abud, Edsel M., Ricardo N. Ramirez, Eric S. Martinez, Luke M. Healy, Cecilia H. H. Nguyen, Sean A.
Newman, Andriy V. Yeromin, Vanessa M. Scarfone, Samuel E. Marsh, Cristhian Fimbres, Chad A.
Caraway, Gianna M. Fote, Abdullah M. Madany, Anshu Agrawal, Rakez Kayed, Karen H. Gylys,
Michael D. Cahalan, Brian J. Cummings, Jack P. Antel, Ali Mortazavi, Monica J. Carson, Wayne
W. Poon, and Mathew Blurton-Jones. 2017. 'iPSC-Derived Human Microglia-like Cells to Study
Neurological Diseases', *Neuron*, 94: 278-93.e9.
Alvarez-Carbonell, D., F. Ye, N. Ramanath, Y. Garcia-Mesa, P. E. Knapp, K. F. Hauser, and J. Karn. 2019.
'Cross-talk between microglia and neurons regulates HIV latency', *PLoS Pathog*, 15: e1008249.
Alvarez-Carbonell, David, Fengchun Ye, Nirmala Ramanath, Yoelvis Garcia-Mesa, Pamela E. Knapp, Kurt
F. Hauser, and Jonathan Karn. 2020. 'Cross-talk between microglia and neurons regulates HIV
latency', *PLOS Pathogens*, 15: e1008249.
Anderson, Albert M., Jeffrey L. Lennox, Mark M. Mulligan, David W. Loring, Henrik Zetterberg, Kaj
Blennow, Cari Kessing, Rajeth Koneru, Kirk Easley, and William R. Tyor. 2017. 'Cerebrospinal
fluid interferon alpha levels correlate with neurocognitive impairment in ambulatory HIV-
Infected individuals', *Journal of NeuroVirology*, 23: 106-12.
Asensio, V. C., J. Maier, R. Milner, K. Boztug, C. Kincaid, M. Moulard, C. Phillipson, K. Lindsley, T.
Krucker, H. S. Fox, and I. L. Campbell. 2001. 'Interferon-independent, human immunodeficiency
virus type 1 gp120-mediated induction of CXCL10/IP-10 gene expression by astrocytes in vivo
and in vitro', *Journal of Virology*, 75: 7067-77.
Aso, H., J. Ito, Y. Koyanagi, and K. Sato. 2019. 'Comparative Description of the Expression Profile of
Interferon-Stimulated Genes in Multiple Cell Lineages Targeted by HIV-1 Infection', *Front
Microbiol*, 10: 429.
Baruch, Kuti, Aleksandra Deczkowska, Eyal David, Joseph M. Castellano, Omer Miller, Alexander
Kertser, Tamara Berkutzki, Zohar Barnett-Itzhaki, Dana Bezalel, Tony Wyss-Coray, Ido Amit, and
Michal Schwartz. 2014. 'Aging-induced type I interferon response at the choroid plexus
negatively affects brain function', *Science*, 346: 89-93.
Berg, R. K., S. H. Rahbek, E. Kofod-Olsen, C. K. Holm, J. Melchjorsen, D. G. Jensen, A. L. Hansen, L. B.
Jørgensen, L. Ostergaard, M. Tolstrup, C. S. Larsen, S. R. Paludan, M. R. Jakobsen, and T. H.
Mogensen. 2014. 'T cells detect intracellular DNA but fail to induce type I IFN responses:
implications for restriction of HIV replication', *PLoS One*, 9: e84513.
Birey, Fikri, Jimena Andersen, Christopher D. Makinson, Saiful Islam, Wu Wei, Nina Huber, H. Christina
Fan, Kimberly R. Cordes Metzler, Georgia Panagiotakos, Nicholas Thom, Nancy A. O'Rourke, Lars
M. Steinmetz, Jonathan A. Bernstein, Joachim Hallmayer, John R. Huguenard, and Sergiu P.
Paşca. 2017. 'Assembly of functionally integrated human forebrain spheroids', *Nature*, 545: 54-
59.
Blank, T., and M. Prinz. 2017. 'Type I interferon pathway in CNS homeostasis and neurological
disorders', *Glia*, 65: 1397-406.
Burdo, Tricia H., Andrew Lackner, and Kenneth C. Williams. 2013. 'Monocyte/macrophages and their
role in HIV neuropathogenesis', *Immunological Reviews*, 254: 102-13.
Burlacu, R., A. Umlauf, T. D. Marcotte, B. Soontornniyomkij, C. C. Diaconu, A. Bulacu-Talnariu, A.
Temereanca, S. M. Ruta, S. Letendre, L. Ene, and C. L. Achim. 2020. 'Plasma CXCL10 correlates
with HAND in HIV-infected women', *Journal of NeuroVirology*, 26: 23-31.

Butler, S. L., M. S. Hansen, and F. D. Bushman. 2001. 'A quantitative assay for HIV DNA integration in vivo', *Nat Med*, 7: 631-4.

Byrnes, S. J., T. A. Angelovich, K. Busman-Sahay, C. R. Cochrane, M. Roche, J. D. Estes, and M. J. Churchill. 2022. 'Non-Human Primate Models of HIV Brain Infection and Cognitive Disorders', *Viruses*, 14.

Capuron, Lucile, Jane F. Gumnick, Dominique L. Musselman, David H. Lawson, Andrea Reemsnyder, Charles B. Nemeroff, and Andrew H. Miller. 2002. 'Neurobehavioral Effects of Interferon-α in Cancer Patients: Phenomenology and Paroxetine Responsiveness of Symptom Dimensions', *Neuropsychopharmacology*, 26: 643-52.

Capuron, Lucile, Alain Ravaud, and Robert Dantzer. 2001. 'Timing and Specificity of the Cognitive Changes Induced by Interleukin-2 and Interferon-α Treatments in Cancer Patients', *Psychosomatic Medicine*, 63.

Casano, Alessandra Maria, and Francesca Peri. 2015. 'Microglia: Multitasking Specialists of the Brain', *Developmental Cell*, 32: 469-77.

Chargelegue, Daniel, and Carol M. O'Toole. 1992. 'Development of a sensitive ELISA for HIV-1 p24 antigen using a fluorogenic substrate for monitoring HIV-1 replication in vitro', *Journal of Virological Methods*, 38: 323-32.

Chen, Natalie C., Andrea T. Partridge, Christian Sell, Claudio Torres, and Julio Martín-García. 2017. 'Fate of microglia during HIV-1 infection: From activation to senescence?', *Glia*, 65: 431-46.

Christina, A. Meyers, S. Scheibel Randall, and D. Forman Arthur. 1991. 'Persistent neurotoxicity of systemically administered interferon-alpha', *Neurology*, 41: 672.

Claes, Christel, Johanna Van den Daele, and Catherine M. Verfaillie. 2018. 'Generating tissue-resident macrophages from pluripotent stem cells: Lessons learned from microglia', *Cellular Immunology*, 330: 60-67.

Clifford, Katherine M., Vishal Samboju, Yann Cobigo, Benedetta Milanini, Gabriel A. Marx, Joanna M. Hellmuth, Howard J. Rosen, Joel H. Kramer, Isabel E. Allen, and Victor G. Valcour. 2017. 'Progressive Brain Atrophy Despite Persistent Viral Suppression in HIV Patients Older Than 60 Years', *Journal of acquired immune deficiency syndromes (1999)*, 76: 289-97.

Crocker, Paul R., James C. Paulson, and Ajit Varki. 2007. 'Siglecs and their roles in the immune system', *Nature Reviews Immunology*, 7: 255-66.

Di Sclafani, V., R. D. Mackay, D. J. Meyerhoff, D. Norman, M. W. Weiner, and G. Fein. 1997. 'Brain atrophy in HIV infection is more strongly associated with CDC clinical stage than with cognitive impairment', *Journal of the International Neuropsychological Society : JINS*, 3: 276-87.

Dickey, L. L., L. J. Martins, V. Planelles, and T. M. Hanley. 2022. 'HIV-1-induced type I IFNs promote viral latency in macrophages', *J Leukoc Biol*, 112: 1343-56.

dos Reis, Roberta S., Shilpa Sant, Hannah Keeney, Marc C. E. Wagner, and Velpandi Ayyavoo. 2020. 'Modeling HIV-1 neuropathogenesis using three-dimensional human brain organoids (hBORGs) with HIV-1 infected microglia', *Scientific Reports*, 10: 15209.

Durafourt, B. A., C. S. Moore, D. A. Zammit, T. A. Johnson, F. Zaguia, M. C. Guiot, A. Bar-Or, and J. P. Antel. 2012. 'Comparison of polarization properties of human adult microglia and blood-derived macrophages', *Glia*, 60: 717-27.

Eggers, C., G. Arendt, K. Hahn, I. W. Husstedt, M. Maschke, E. Neuen-Jacob, M. Obermann, T. Rosenkranz, E. Schielke, and E. Straube. 2017. 'HIV-1-associated neurocognitive disorder: epidemiology, pathogenesis, diagnosis, and treatment', *J Neurol*, 264: 1715-27.

Elsner, C., A. Ponnurangam, J. Kazmierski, T. Zillinger, J. Jansen, D. Todt, K. Döhner, S. Xu, A. Ducroux, N. Kriedemann, A. Malassa, P. K. Larsen, G. Hartmann, W. Barchet, E. Steinmann, U. Kalinke, B. Sodeik, and C. Goffinet. 2020. 'Absence of cGAS-mediated type I IFN responses in HIV-1-infected T cells', *Proc Natl Acad Sci U S A*, 117: 19475-86.

Eugenin, E. A., and J. W. Berman. 2016. 'Improved Methods to Detect Low Levels of HIV Using Antibody-Based Technologies', *Methods Mol Biol*, 1354: 265-79.

Gallo, Paolo, Silvana Pagni, Bruno Giometto, Maria G. Piccinno, Felice Bozza, Vincenza Argentiero, and Bruno Tavolato. 1990. 'Macrophage-colony stimulating factor (M-CSF) in the cerebrospinal fluid', *Journal of Neuroimmunology*, 29: 105-12.

Gendelman, Howard E., Stuart A. Lipton, Marc Tardieu, Michael I. Bukrinsky, and Hans S. L. M. Nottet. 1994. 'The neuropathogenesis of HIV-1 infection', *Journal of Leukocyte Biology*, 56: 389-98.

Goldmann, T., T. Blank, and M. Prinz. 2016. 'Fine-tuning of type I IFN-signaling in microglia--implications for homeostasis, CNS autoimmunity and interferonopathies', *Curr Opin Neurobiol*, 36: 38-42.

González-Scarano, Francisco, and Julio Martín-García. 2005. 'The neuropathogenesis of AIDS', *Nature Reviews Immunology*, 5: 69-81.

Gosselin, David, Dylan Skola, Nicole G. Coufal, Inge R. Holtman, Johannes C. M. Schlachetzki, Eniko Sajti, Baptiste N. Jaeger, Carolyn O'Connor, Conor Fitzpatrick, Martina P. Pasillas, Monique Pena, Amy Adair, David D. Gonda, Michael L. Levy, Richard M. Ransohoff, Fred H. Gage, and Christopher K. Glass. 2017. 'An environment-dependent transcriptional network specifies human microglia identity', *Science*, 356: eaal3222.

Gumbs, Stephanie B. H., Amber Berdenis van Berlekom, Raphael Kübler, Pauline J. Schipper, Lavina Gharu, Marco P. Boks, Paul R. Ormel, Annemarie M. J. Wensing, Lot D. de Witte, and Monique Nijhuis. 2022. "Characterization of HIV-1 Infection in Microglia-Containing Human Cerebral Organoids." In *Viruses*.

Haenseler, W., S. N. Sansom, J. Buchrieser, S. E. Newey, C. S. Moore, F. J. Nicholls, S. Chintawar, C. Schnell, J. P. Antel, N. D. Allen, M. Z. Cader, R. Wade-Martins, W. S. James, and S. A. Cowley. 2017. 'A Highly Efficient Human Pluripotent Stem Cell Microglia Model Displays a Neuronal-Co-culture-Specific Expression Profile and Inflammatory Response', *Stem Cell Reports*, 8: 1727-42.

Hammond, Timothy R., Connor Dufort, Lasse Dissing-Olesen, Stefanie Giera, Adam Young, Alec Wysoker, Alec J. Walker, Frederick Gergits, Michael Segel, James Nemesh, Samuel E. Marsh, Arpiar Saunders, Evan Macosko, Florent Ginhoux, Jinmiao Chen, Robin J. M. Franklin, Xianhua Piao, Steven A. McCarroll, and Beth Stevens. 2019. 'Single-Cell RNA Sequencing of Microglia throughout the Mouse Lifespan and in the Injured Brain Reveals Complex Cell-State Changes', *Immunity*, 50: 253-71.e6.

Hanisch, U. K. 2002. 'Microglia as a source and target of cytokines', *Glia*, 40: 140-55.

Hofer, Markus J., and Iain L. Campbell. 2013. 'Type I interferon in neurological disease—The devil from within', *Cytokine & Growth Factor Reviews*, 24: 257-67.

Hur, J. Y., G. R. Frost, X. Wu, C. Crump, S. J. Pan, E. Wong, M. Barros, T. Li, P. Nie, Y. Zhai, J. C. Wang, J. Tcw, L. Guo, A. McKenzie, C. Ming, X. Zhou, M. Wang, Y. Sagi, A. E. Renton, B. T. Esposito, Y. Kim, K. R. Sadleir, I. Trinh, R. A. Rissman, R. Vassar, B. Zhang, D. S. Johnson, E. Masliah, P. Greengard, A. Goate, and Y. M. Li. 2020. 'The innate immunity protein IFITM3 modulates γ-secretase in Alzheimer's disease', *Nature*, 586: 735-40.

Israel, Sarah M., Shiva Hassanzadeh-Behbahani, Peter E. Turkeltaub, David J. Moore, Ronald J. Ellis, and
Xiong Jiang. 2019. 'Different roles of frontal versus striatal atrophy in HIV-associated
neurocognitive disorders', *Human Brain Mapping*, 40: 3010-26.
Izquierdo-Useros, Nuria, Maier Lorizate, Maria C. Puertas, Maria T. Rodriguez-Plata, Nadine Zangger,
Elina Erikson, Maria Pino, Itziar Erkizia, Bärbel Glass, Bonaventura Clotet, Oliver T. Keppler,
Amalio Telenti, Hans-Georg Kräusslich, and Javier Martinez-Picado. 2012. 'Siglec-1 Is a Novel
Dendritic Cell Receptor That Mediates HIV-1 Trans-Infection Through Recognition of Viral
Membrane Gangliosides', *PLOS Biology*, 10: e1001448.
Jeremiah, N., H. Ferran, K. Antoniadou, K. De Azevedo, J. Nikolic, M. Maurin, P. Benaroch, and N.
Manel. 2023. 'RELA tunes innate-like interferon I/III responses in human T cells', *J Exp Med*, 220.
Jiao, Yanmei, Tong Zhang, Rui Wang, Hongwei Zhang, Xiaojie Huang, Jiming Yin, Liguo Zhang, Xiaoning
Xu, and Hao Wu. 2012. 'Plasma IP-10 Is Associated with Rapid Disease Progression in Early HIV-
1 Infection', *Viral Immunology*, 25: 333-37.
Jin, M., R. Xu, L. Wang, M. M. Alam, Z. Ma, S. Zhu, A. C. Martini, A. Jadali, M. Bernabucci, P. Xie, K. Y.
Kwan, Z. P. Pang, E. Head, Y. Liu, R. P. Hart, and P. Jiang. 2022. 'Type-I-interferon signaling drives
microglial dysfunction and senescence in human iPSC models of Down syndrome and
Alzheimer's disease', *Cell Stem Cell*, 29: 1135-53.e8.
Kalter, D. C., M. Nakamura, J. A. Turpin, L. M. Baca, D. L. Hoover, C. Dieffenbach, P. Ralph, H. E.
Gendelman, and M. S. Meltzer. 1991. 'Enhanced HIV replication in macrophage colony-
stimulating factor-treated monocytes', *J Immunol*, 146: 298-306.
Kaul, M., J. Zheng, S. Okamoto, H. E. Gendelman, and S. A. Lipton. 2005. 'HIV-1 infection and AIDS:
consequences for the central nervous system', *Cell Death Differ*, 12 Suppl 1: 878-92.
Kaul, Marcus, Gwenn A. Garden, and Stuart A. Lipton. 2001. 'Pathways to neuronal injury and
apoptosis in HIV-associated dementia', *Nature*, 410: 988-94.
Koenig, S., H. E. Gendelman, J. M. Orenstein, M. C. Dal Canto, G. H. Pezeshkpour, M. Yungbluth, F.
Janotta, A. Aksamit, M. A. Martin, and A. S. Fauci. 1986. 'Detection of AIDS virus in
macrophages in brain tissue from AIDS patients with encephalopathy', *Science*, 233: 1089-93.
Kolb, Stefan A., Bernd Sporer, Fritz Lahrtz, Uwe Koedel, Hans-Walter Pfister, and Adriano Fontana.
1999. 'Identification of a T cell chemotactic factor in the cerebrospinal fluid of HIV-1-infected
individuals as interferon-γ inducible protein 10', *Journal of Neuroimmunology*, 93: 172-81.
Koyanagi, Y., S. Miles, R. T. Mitsuyasu, J. E. Merrill, H. V. Vinters, and I. S. Chen. 1987. 'Dual infection of
the central nervous system by AIDS viruses with distinct cellular tropisms', *Science*, 236: 819-22.
Kraft-Terry, Stephanie D., Shilpa J. Buch, Howard S. Fox, and Howard E. Gendelman. 2009. 'A Coat of
Many Colors: Neuroimmune Crosstalk in Human Immunodeficiency Virus Infection', *Neuron*, 64:
133-45.
Krivine, A., G. Force, J. Servan, A. Cabée, F. Rozenberg, L. Dighiero, F. Marguet, and P. Lebon. 1999.
'Measuring HIV-1 RNA and interferon-alpha in the cerebrospinal fluid of AIDS patients: insights
into the pathogenesis of AIDS Dementia Complex', *J Neurovirol*, 5: 500-6.
Kutsch, O., J. Oh, A. Nath, and E. N. Benveniste. 2000. 'Induction of the chemokines interleukin-8 and
IP-10 by human immunodeficiency virus type 1 tat in astrocytes', *J Virol*, 74: 9214-21.
Kutza, J., K. Fields, T. A. Grimm, and K. A. Clouse. 2002. 'Inhibition of HIV replication and macrophage
colony-stimulating factor production in human macrophages by antiretroviral agents', *AIDS Res
Hum Retroviruses*, 18: 619-25.

Kutza, Joseph, Lynne Crim, Steven Feldman, Mark P. Hayes, Marion Gruber, Judy Beeler, and Kathleen
A. Clouse. 2000. 'Macrophage Colony-Stimulating Factor Antagonists Inhibit Replication of HIV-
1 in Human Macrophages', *The Journal of Immunology*, 164: 4955.
Lancaster, Madeline A., and Juergen A. Knoblich. 2014. 'Generation of cerebral organoids from human
pluripotent stem cells', *Nature Protocols*, 9: 2329-40.
Lancaster, Madeline A., Magdalena Renner, Carol-Anne Martin, Daniel Wenzel, Louise S. Bicknell,
Matthew E. Hurles, Tessa Homfray, Josef M. Penninger, Andrew P. Jackson, and Juergen A.
Knoblich. 2013. 'Cerebral organoids model human brain development and microcephaly',
Nature, 501: 373-79.
Lee, Jeong Seok, and Eui-Cheol Shin. 2020. 'The type I interferon response in COVID-19: implications for
treatment', *Nature Reviews Immunology*, 20: 585-86.
Lei, Jie, Xiaowan Yin, Hong Shang, and Yongjun Jiang. 2019. 'IP-10 is highly involved in HIV infection',
Cytokine, 115: 97-103.
Letendre, S. L., E. R. Lanier, and J. A. McCutchan. 1999. 'Cerebrospinal fluid beta chemokine
concentrations in neurocognitively impaired individuals infected with human immunodeficiency
virus type 1', *J Infect Dis*, 180: 310-9.
Lew, Brandon J., Timothy J. McDermott, Alex I. Wiesman, Jennifer Neill, Mackenzie S. Mills, Kevin R.
Robertson, Howard S. Fox, Susan Swindells, and Tony W. Wilson. 2018. 'Neural dynamics of
selective attention deficits in HIV-associated neurocognitive disorder', *Neurology*, 91: e1860.
Li, Qingyun, and Ben A. Barres. 2018. 'Microglia and macrophages in brain homeostasis and disease',
Nature Reviews Immunology, 18: 225-42.
Li, Y., J. C. Kappes, J. A. Conway, R. W. Price, G. M. Shaw, and B. H. Hahn. 1991. 'Molecular
characterization of human immunodeficiency virus type 1 cloned directly from uncultured
human brain tissue: identification of replication-competent and -defective viral genomes',
Journal of Virology, 65: 3973-85.
Lindl, Kathryn A., David R. Marks, Dennis L. Kolson, and Kelly L. Jordan-Sciutto. 2010. 'HIV-Associated
Neurocognitive Disorder: Pathogenesis and Therapeutic Opportunities', *Journal of
Neuroimmune Pharmacology*, 5: 294-309.
Mackelprang, R. D., A. Filali-Mouhim, B. Richardson, F. Lefebvre, E. Katabira, A. Ronald, G. Gray, K. W.
Cohen, N. R. Klatt, T. Pecor, C. Celum, M. J. McElrath, S. M. Hughes, F. Hladik, M. J. Cameron,
and J. R. Lingappa. 2023. 'Upregulation of IFN-stimulated genes persists beyond the transitory
broad immunologic changes of acute HIV-1 infection', *iScience*, 26: 106454.
Maddison, D. C., and F. Giorgini. 2015. 'The kynurenine pathway and neurodegenerative disease',
Semin Cell Dev Biol, 40: 134-41.
Malpass, K. 2011. 'Neurodegenerative disease: the kynurenine pathway--promising new targets and
therapies for neurodegenerative disease', *Nat Rev Neurol*, 7: 417.
Marodon, Gilles, David Warren, Michael C. Filomio, and David N. Posnett. 1999. 'Productive infection of
double-negative T cells with HIV in vivo', *Proceedings of the National
Academy of Sciences*, 96: 11958.
McGlasson, Sarah, Alexa Jury, Andrew Jackson, and David Hunt. 2015. 'Type I interferon dysregulation
and neurological disease', *Nature Reviews Neurology*, 11: 515-23.
Merimsky, Ofer, Irith Reider-Groswasser, Moshe Inbar, and Samario Chaitchik. 1990. 'Interferon-
related mental deterioration and behavioral changes in patients with renal cell carcinoma',
European Journal of Cancer and Clinical Oncology, 26: 596-600.

Merrill, J. E., Y. Koyanagi, J. Zack, L. Thomas, F. Martin, and I. S. Chen. 1992. 'Induction of interleukin-1 and tumor necrosis factor alpha in brain cultures by human immunodeficiency virus type 1', *Journal of Virology*, 66: 2217.

Min, A. K., B. Javidfar, R. Missall, D. Doanman, M. Durens, S. S. Vil, Z. Masih, M. Graziani, A. Mordelt, S. Marro, L. de Witte, B. K. Chen, T. H. Swartz, and S. Akbarian. 2023. 'HIV-1 infection of genetically engineered iPSC-derived central nervous system-engrafted microglia in a humanized mouse model', *bioRxiv*.

Mitrasinovic, Olivera M., Grace V. Perez, FeiFei Zhao, Yuen Ling Lee, Clara Poon, and Greer M. Murphy. 2001. 'Overexpression of Macrophage Colony-stimulating Factor Receptor on Microglial Cells Induces an Inflammatory Response*', *Journal of Biological Chemistry*, 276: 30142-49.

Moretti, S., S. Virtuoso, L. Sernicola, S. Farcomeni, M. T. Maggiorella, and A. Borsetti. 2021. 'Advances in SIV/SHIV Non-Human Primate Models of NeuroAIDS', *Pathogens*, 10.

Nasr, N., A. A. Alshehri, T. K. Wright, M. Shahid, B. M. Heiner, A. N. Harman, R. A. Botting, K. J. Helbig, M. R. Beard, K. Suzuki, A. D. Kelleher, P. Hertzog, and A. L. Cunningham. 2017. 'Mechanism of Interferon-Stimulated Gene Induction in HIV-1-Infected Macrophages', *J Virol*, 91.

Ormel, Paul R., Renata Vieira de Sá, Emma J. van Bodegraven, Henk Karst, Oliver Harschnitz, Marjolein A. M. Sneeboer, Lill Eva Johansen, Roland E. van Dijk, Nicky Scheefhals, Amber Berdenis van Berlekom, Eduardo Ribes Martínez, Sandra Kling, Harold D. MacGillavry, Leonard H. van den Berg, René S. Kahn, Elly M. Hol, Lot D. de Witte, and R. Jeroen Pasterkamp. 2018. 'Microglia innately develop within cerebral organoids', *Nature Communications*, 9: 4167.

Paolicelli, Rosa C., Giulia Bolasco, Francesca Pagani, Laura Maggi, Maria Scianni, Patrizia Panzanelli, Maurizio Giustetto, Tiago Alves Ferreira, Eva Guiducci, Laura Dumas, Davide Ragozzino, and Cornelius T. Gross. 2011. 'Synaptic Pruning by Microglia Is Necessary for Normal Brain Development', *Science*, 333: 1456.

Paşca, Anca M., Steven A. Sloan, Laura E. Clarke, Yuan Tian, Christopher D. Makinson, Nina Huber, Chul Hoon Kim, Jin-Young Park, Nancy A. O'Rourke, Khoa D. Nguyen, Stephen J. Smith, John R. Huguenard, Daniel H. Geschwind, Ben A. Barres, and Sergiu P. Paşca. 2015. 'Functional cortical neurons and astrocytes from human pluripotent stem cells in 3D culture', *Nature Methods*, 12: 671-78.

Pavol, M. A., C. A. Meyers, J. L. Rexer, A. D. Valentine, P. J. Mattis, and M. Talpaz. 1995. 'Pattern of neurobehavioral deficits associated with interferon alfa therapy for leukemia', *Neurology*, 45: 947-50.

Peluso, Michael J., Victor Valcour, Nittaya Phanuphak, Jintanat Ananworanich, James L. K. Fletcher, Thep Chalermchai, Shelly J. Krebs, Merlin L. Robb, Joanna Hellmuth, Magnus Gisslén, Henrik Zetterberg, Serena Spudich, Rv Search Rv254Search, and Search Study Teams. 2017. 'Immediate initiation of cART is associated with lower levels of cerebrospinal fluid YKL-40, a marker of microglial activation, in HIV-1 infection', *AIDS (London, England)*, 31: 247-52.

Perrella, O., P. B. Carreiri, A. Perrella, C. Sbreglia, F. Gorga, D. Guarnaccia, and G. Tarantino. 2001. 'Transforming growth factor beta-1 and interferon-alpha in the AIDS dementia complex (ADC): possible relationship with cerebral viral load?', *Eur Cytokine Netw*, 12: 51-5.

Phetsouphanh, Chansavath, David R. Darley, Daniel B. Wilson, Annett Howe, C. Mee Ling Munier, Sheila K. Patel, Jennifer A. Juno, Louise M. Burrell, Stephen J. Kent, Gregory J. Dore, Anthony D. Kelleher, and Gail V. Matthews. 2022. 'Immunological dysfunction persists for 8 months following initial mild-to-moderate SARS-CoV-2 infection', *Nature Immunology*, 23: 210-16.

Platanias, Leonidas C. 2005. 'Mechanisms of type-I- and type-II-interferon-mediated signalling', *Nature*
Reviews Immunology, 5: 375-86.
Qian, Xuyu, Yijing Su, Christopher D. Adam, Andre U. Deutschmann, Sarshan R. Pather, Ethan M.
Goldberg, Kenong Su, Shiying Li, Lu Lu, Fadi Jacob, Phuong T. T. Nguyen, Sooyoung Huh, Ahmet
Hoke, Sarah E. Swinford-Jackson, Zhexing Wen, Xiaosong Gu, R. Christopher Pierce, Hao Wu,
Lisa A. Briand, H. Isaac Chen, John A. Wolf, Hongjun Song, and Guo-li Ming. 2020. 'Sliced Human
Cortical Organoids for Modeling Distinct Cortical Layer Formation', *Cell Stem Cell*, 26: 766-81.e9.
Quadrato, Giorgia, Tuan Nguyen, Evan Z. Macosko, John L. Sherwood, Sung Min Yang, Daniel R. Berger,
Natalie Maria, Jorg Scholvin, Melissa Goldman, Justin P. Kinney, Edward S. Boyden, Jeff W.
Lichtman, Ziv M. Williams, Steven A. McCarroll, and Paola Arlotta. 2017. 'Cell diversity and
network dynamics in photosensitive human brain organoids', *Nature*, 545: 48-53.
Rai, Mohammad A., Jason Hammonds, Mario Pujato, Christopher Mayhew, Krishna Roskin, and Paul
Spearman. 2020. 'Comparative analysis of human microglial models for studies of HIV
replication and pathogenesis', *Retrovirology*, 17: 35.
Raison, Charles L., Marina Demetrashvili, Lucile Capuron, and Andrew H. Miller. 2005.
'Neuropsychiatric Adverse Effects of Interferon-α', *CNS Drugs*, 19: 105-23.
Ramesh, Geeta, Andrew G. MacLean, and Mario T. Philipp. 2013. 'Cytokines and Chemokines at the
Crossroads of Neuroinflammation, Neurodegeneration, and Neuropathic Pain', *Mediators of*
Inflammation, 2013: 480739.
Rappaport, Jay, Jeymohan Joseph, Sidney Croul, Guillermo Alexander, Luis Del Valle, Shohreh Amini,
and Kamel Khalili. 1999. 'Molecular pathway involved in HIV-1-induced CNS pathology: role of
viral regulatory protein, Tat', *Journal of Leukocyte Biology*, 65: 458-65.
Rho, M. B., S. Wesselingh, J. D. Glass, J. C. McArthur, S. Choi, J. Griffin, and W. R. Tyor. 1995. 'A
Potential Role for Interferon-α in the Pathogenesis of HIV-Associated Dementia', *Brain,*
Behavior, and Immunity, 9: 366-77.
Roy, Ethan R., Gabriel Chiu, Sanming Li, Nicholas E. Propson, Rupa Kanchi, Baiping Wang, Cristian
Coarfa, Hui Zheng, and Wei Cao. 2022. 'Concerted type I interferon signaling in microglia and
neural cells promotes memory impairment associated with amyloid β plaques', *Immunity*, 55:
879-94.e6.
Roy, Ethan R., Baiping Wang, Ying-wooi Wan, Gabriel Chiu, Allysa Cole, Zhuoran Yin, Nicholas E.
Propson, Yin Xu, Joanna L. Jankowsky, Zhandong Liu, Virginia M. Y. Lee, John Q. Trojanowski,
Stephen D. Ginsberg, Oleg Butovsky, Hui Zheng, and Wei Cao. 2020. 'Type I interferon response
drives neuroinflammation and synapse loss in Alzheimer disease', *The Journal of Clinical*
Investigation, 130: 1912-30.
Ruhanya, Vurayai, Graeme B. Jacobs, Shalena Naidoo, Robert H. Paul, John A. Joska, Soraya Seedat,
George Nyandoro, Susan Engelbrecht, and Richard H. Glashoff. 2021. 'Impact of Plasma IP-
10/CXCL10 and RANTES/CCL5 Levels on Neurocognitive Function in HIV Treatment-Naive
Patients', *AIDS Research and Human Retroviruses*.
Ryan, Sean K., Michael V. Gonzalez, James P. Garifallou, Frederick C. Bennett, Kimberly S. Williams,
Nathaniel P. Sotuyo, Eugene Mironets, Kieona Cook, Hakon Hakonarson, Stewart A. Anderson,
and Kelly L. Jordan-Sciutto. 2020. 'Neuroinflammation and EIF2 Signaling Persist despite
Antiretroviral Treatment in an hiPSC Tri-culture Model of HIV Infection', *Stem Cell Reports*, 14:
703-16.

Sacktor, N., R. L. Skolasky, E. Seaberg, C. Munro, J. T. Becker, E. Martin, A. Ragin, A. Levine, and E.
306   Miller. 2016. 'Prevalence of HIV-associated neurocognitive disorders in the Multicenter AIDS
307   Cohort Study', *Neurology*, 86: 334-40.
308   Sacktor, Ned. 2018. 'Changing clinical phenotypes of HIV-associated neurocognitive disorders', *Journal*
309   *of NeuroVirology*, 24: 141-45.
310   Sanna, Pietro Paolo, Yu Fu, Eliezer Masliah, Celine Lefebvre, and Vez Repunte-Canonigo. 2021. 'Central
311   nervous system (CNS) transcriptomic correlates of human immunodeficiency virus (HIV) brain
312   RNA load in HIV-infected individuals', *Scientific Reports*, 11: 12176.
313   Schwarcz, R., and T. W. Stone. 2017. 'The kynurenine pathway and the brain: Challenges, controversies
314   and promises', *Neuropharmacology*, 112: 237-47.
315   Simioni, S., M. Cavassini, J. M. Annoni, A. Rimbault Abraham, I. Bourquin, V. Schiffer, A. Calmy, J. P.
316   Chave, E. Giacobini, B. Hirschel, and R. A. Du Pasquier. 2010. 'Cognitive dysfunction in HIV
317   patients despite long-standing suppression of viremia', *Aids*, 24: 1243-50.
318   Sloan, Steven A., Jimena Andersen, Anca M. Paşca, Fikri Birey, and Sergiu P. Paşca. 2018. 'Generation
319   and assembly of human brain region–specific three-dimensional cultures', *Nature Protocols*, 13:
320   2062-85.
321   Smith, Amy M., Hannah M. Gibbons, Robyn L. Oldfield, Peter M. Bergin, Edward W. Mee, Maurice A.
322   Curtis, Richard L. M. Faull, and Mike Dragunow. 2013. 'M-CSF increases proliferation and
323   phagocytosis while modulating receptor and transcription factor expression in adult human
324   microglia', *Journal of Neuroinflammation*, 10: 859.
325   Solomon, Isaac H., Sukrutha Chettimada, Vikas Misra, David R. Lorenz, Robert J. Gorelick, Benjamin B.
326   Gelman, Susan Morgello, and Dana Gabuzda. 2020. 'White Matter Abnormalities Linked to
327   Interferon, Stress Response, and Energy Metabolism Gene Expression Changes in Older HIV-
328   Positive Patients on Antiretroviral Therapy', *Molecular Neurobiology*, 57: 1115-30.
329   Song, Wilbur M., and Marco Colonna. 2018. 'The identity and function of microglia in
330   neurodegeneration', *Nature Immunology*, 19: 1048-58.
331   Sreeram, S., F. Ye, Y. Garcia-Mesa, K. Nguyen, A. El Sayed, K. Leskov, and J. Karn. 2022. 'The potential
332   role of HIV-1 latency in promoting neuroinflammation and HIV-1-associated neurocognitive
333   disorder', *Trends Immunol*, 43: 630-39.
334   Valverde-Villegas, J. M., R. M. de Medeiros, J. H. Ellwanger, B. R. Santos, M. G. Melo, S. E. M. Almeida,
335   and J. A. B. Chies. 2018. 'High CXCL10/IP-10 levels are a hallmark in the clinical evolution of the
336   HIV infection', *Infect Genet Evol*, 57: 51-58.
337   van Marle, G., S. Henry, T. Todoruk, A. Sullivan, C. Silva, S. B. Rourke, J. Holden, J. C. McArthur, M. J.
338   Gill, and C. Power. 2004. 'Human immunodeficiency virus type 1 Nef protein mediates neural
339   cell death: a neurotoxic role for IP-10', *Virology*, 329: 302-18.
340   Vaughan-Jackson, Alun, Szymon Stodolak, Kourosh H. Ebrahimi, Cathy Browne, Paul K. Reardon,
341   Elisabete Pires, Javier Gilbert-Jaramillo, Sally A. Cowley, and William S. James. 2021.
342   'Differentiation of human induced pluripotent stem cells to authentic macrophages using a
343   defined, serum-free, open-source medium', *Stem Cell Reports*, 16: 1735-48.
344   Velasco, Silvia, Amanda J. Kedaigle, Sean K. Simmons, Allison Nash, Marina Rocha, Giorgia Quadrato,
345   Bruna Paulsen, Lan Nguyen, Xian Adiconis, Aviv Regev, Joshua Z. Levin, and Paola Arlotta. 2019.
346   'Individual brain organoids reproducibly form cell diversity of the human cerebral cortex',
347   *Nature*, 570: 523-27.

Vodyanik, Maxim A., James A. Thomson, and Igor I. Slukvin. 2006. 'Leukosialin (CD43) defines
hematopoietic progenitors in human embryonic stem cell differentiation cultures', *Blood*, 108:
2095-105.
Wichers, M. C., G. H. Koek, G. Robaeys, A. J. Praamstra, and M. Maes. 2005. 'Early increase in
vegetative symptoms predicts IFN-α-induced cognitive-depressive changes', *Psychological*
Medicine, 35: 433-41.
Wiley, C. A., R. A. Steinman, and Q. Wang. 2023. 'Innate immune activation without immune cell
infiltration in brains of murine models of Aicardi-Goutières Syndrome', *Brain Pathol*, 33:
e13118.
Woodburn, B. M., K. Kanchi, S. Zhou, N. Colaianni, S. B. Joseph, and R. Swanstrom. 2022.
'Characterization of Macrophage-Tropic HIV-1 Infection of Central Nervous System Cells and the
Influence of Inflammation', *J Virol*, 96: e0095722.
Xu, R., A. J. Boreland, X. Li, C. Erickson, M. Jin, C. Atkins, Z. P. Pang, B. P. Daniels, and P. Jiang. 2021.
'Developing human pluripotent stem cell-based cerebral organoids with a controllable microglia
ratio for modeling brain development and pathology', *Stem Cell Reports*, 16: 1923-37.
Yoon, Se-Jin, Lubayna S. Elahi, Anca M. Pașca, Rebecca M. Marton, Aaron Gordon, Omer Revah, Yuki
Miura, Elisabeth M. Walczak, Gwendolyn M. Holdgate, H. Christina Fan, John R. Huguenard,
Daniel H. Geschwind, and Sergiu P. Pașca. 2019. 'Reliability of human cortical organoid
generation', *Nature Methods*, 16: 75-78.
Yu, H., A. B. Rabson, M. Kaul, Y. Ron, and J. P. Dougherty. 1996. 'Inducible human immunodeficiency
virus type 1 packaging cell lines', *Journal of Virology*, 70: 4530-37.
Zou, Zhongcheng, Ashley Chastain, Susan Moir, Jennifer Ford, Kathryn Trandem, Elena Martinelli,
Claudia Cicala, Paul Crocker, James Arthos, and Peter D. Sun. 2011. 'Siglecs Facilitate HIV-1
Infection of Macrophages through Adhesion with Viral Sialic Acids', *PLoS One*, 6: e24559.
372
373

FIGURE LEGENDS

Figure 1: Human microglia are efficiently generated from induced pluripotent stem cells

(A) Representative brightfield images showing the induction of iPSCs to mesodermal yolk sac embryoid

bodies (EB) and then to primitive microglia progenitor cells (PMPs) and microglia.

(B) Representative confocal images of PMPs expressing the yolk sac marker CD235 and the

hematopoietic progenitor cell marker CD43 in the 03SF cell line.

(C) RT–qPCR analysis of *IBA1* and *P2RY12* mRNA expression in iPSCs and iPSC-derived microglia;

n=3 cell lines (03SF, CD06, CD52); t test; data are presented as the mean ± SEM.

(D) Representative confocal images of microglia stained for the microglial markers IBA1, P2Y12,

CX3CR1, TMEM119, and PU1 and costained with DAPI; n = 3 cell lines (03SF, CD06, CD52).

(E) Representative confocal image of a microglial cell displaying ramified morphology after 4 weeks of

coculture with neurons.

(F) Representative confocal images of a 03SF cell line iPSC-derived microglial neuronal coculture

stained for neurons (MAP2), synapses (SYN1), and microglia (IBA1). (F1 and F2) Orthogonal view of

microglial (IBA1+) processes colocalized with neuronal dendrites (MAP2) and synapses (SYN1).

(G) RT–qPCR analysis of *CD4* and *CCR5* mRNA from iPSCs and microglia; n=3 cell lines (03SF,

CD06, CD52); t test; data are presented as the mean ± SEM.

Figure 2: HIV-1-infected microglia exhibit productive infection, activation, and dysregulated

cytokine/chemokine production

(A) RT–qPCR analysis of viral *gag/pol* mRNA after infection with HIV-1 JRFL in microglia monocultures;

n=3 independent infections of 03SF line iPSC-derived microglia.

(B) ELISA analysis of secreted p24 in the culture supernatant of microglia infected with HIV-1 JR-FL

reveals productive infection; n=3 independent infections of 03SF iPSC-derived microglia.

(C) RT–qPCR analysis of *IL1β, TNFα*, and *IL-6* mRNA at days 16 and 23 post infection with HIV-1 JRFL; n=3 independent infections of 03SF iPSC-derived microglia, t test; data are presented as the mean ± SEM.

(D) RT–qPCR analysis of *CCL2, CCL3*, and *CCL5* mRNA at days 16 and 23 post infection with HIV-1 JRFL; n=3 independent infections of 03SF iPSC-derived microglia.

(E) RT–qPCR analysis of *IBA1* and *P2RY12* mRNA at days 16 and 23 post infection with HIV-1 JRFL; n=3 independent infections of 03SF iPSC-derived microglia.

(F) Luminex cytokine/chemokine analysis of secreted M-CSF and IP-10/CXCL10 culture supernatant from HIV-1 JRFL-infected microglia; n=3 independent infections of 03SF microglia, two-way ANOVA; data are presented as the mean ± SEM.

(G) Representative confocal images of nucleocapsid protein p24 staining in HIV-1 JRFL-infected 03SF-derived microglia (IBA1+).

Figure 3: HIV-1-infected microglia exhibit productive infection, activation, and dysregulated cytokine/chemokine production

(A) RT–qPCR analysis of viral *gag/pol* mRNA at days 9, 16, 23, and 30 post infection with HIV-1 JRFL and HIV-1 YU2; n=3 independent infections of CD06 iPSC-derived microglia. t test

(B) ELISA analysis of secreted p24 in the culture supernatant of microglia infected with HIV-1 JRFL and HIV-1 YU2 at days 9, 15, 21, and 27 reveals productive infection; n=3 independent infections of CD06 iPSC-derived microglia.

(C) RT–qPCR analysis of *IL1β* and *TNFα* mRNA at days 9, 16, 23, and 30 post infection with HIV-1 JRFL and HIV-1 YU2; n=3 independent infections of CD06 iPSC-derived microglia.

(D) RT–qPCR analysis of *CCL2* and *CCL3* mRNA at days 9, 16, 23, and 30 post infection with HIV-1 JRFL and HIV-1 YU2; n=3 independent infections of CD06 iPSC-derived microglia.

(E) RT–qPCR analysis of *IBA1* and *P2RY12* mRNA at days 9, 16, 23, and 30 post infection with HIV-1 JRFL and HIV-1 YU2; n=3 independent infections of CD06 iPSC-derived microglia.

Figure 4: HIV-1 causes enrichment of type I interferon signaling pathways and sustained interferon activation

(A) A volcano plot illustrating differentially expressed genes (DEGs) in HIV-1 JRFL-infected vs mock-infected microglia on day 16 post infection; n=3 independent infections of 03SF line iPSC-derived microglia.

(B) GSEA plot showing enrichment of IFNα-responsive genes in HIV-1 JRFL-infected vs mock-infected microglia on day 16 post infection (NES, normalized enrichment score; FDR, false discovery rate).

(C) Gene Ontology (GO) analysis of upregulated DEGs in HIV-1 JRFL vs mock-infected microglia on day 16 post infection; n=3 independent infections of 03SF microglia.

(D) Gene Ontology (GO) analysis of downregulated DEGs in HIV-1 JRFL vs mock-infected microglia on day 16 post infection; n=3 independent infections of 03SF microglia.

(E) A heatmap illustrating significant upregulation of interferon-stimulated genes in HIV-1-infected vs mock-infected microglia on day 16 post infection; n=3 independent infections of 03SF microglia.

(F) A heatmap illustrating SIGLEC gene expression in HIV-1-infected vs mock-infected microglia on day 16 post infection; n=3 independent infections of 03SF microglia.

(G) RT–qPCR analysis of *CXCL10*, *RSAD2*, *ISG15*, and *IDO1* mRNA from HIV-1 JRFL-infected vs mock-infected microglia at days 16 and 23 post infection; n = 3 independent infections of 03SF iPSC-derived microglia, t test; *p < 0.05, **p < 0.01, ***p < 0.001; data are presented as the mean ± SEM.

(H) RT–qPCR analysis of *CXCL10* and *RSAD2* mRNA from HIV-1 JRFL- and HIV-1 YU2-infected and mock-infected microglia at days 16 and 23 post infection; n = 4 independent infections of CD06 iPSC-derived microglia, t test; *p < 0.05, **p < 0.01, ***p < 0.001; data are presented as the mean ± SEM.

Figure 5: HIV-1-infected microglia incorporate into sliced cerebral organoids

(A) Representative confocal images of a day 144 organoid with neurons (MAP2$^+$), astrocytes (GFAP$^+$), and microglia (IBA1$^+$).

(B) Representative confocal images of day 144 organoids 14 days after the addition of HIV-1 JRFL-

infected or mock-infected microglia; (B1) shows a high-magnification image of HIV-1-infected microglia

from (B).

(C) High-magnification image of a single HIV-1-infected microglial cell within a sliced organoid.

(D) RT–qPCR of *gag/pol* mRNA from HIV-1 JRFL-infected or mock-infected microglia organoids on day

14; n=3 independent infections of 03SF microglia; one-tailed t test; ****$P<0.001$; error bars represent

SEM.

(E) ELISA analysis of secreted p24 in culture supernatant from HIV-1 JRFL-infected or mock-infected

microglia organoid day 14; n=3 independent infections of 03SF microglia; one-tailed t test; **$p<0.01$;

data are presented as the mean ± SEM.

(F) RT–qPCR analysis of *ISG15*, *RSAD2*, *IFITM3*, *MX1*, *IFI44*, and *IFI27* mRNA from HIV-1 JRFL-

infected or mock-infected microglia organoids on day 14; n=3 independent infections of 03SF microglia;

one-tailed t test; *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$; data are presented as the mean ± SEM.

Figure S1: Microglial RT-qPCR analysis

(A) RT–qPCR analysis of *IBA1* and *P2RY12* mRNA expression in iPSCs and iPSC-derived microglia;

n=3 cell lines (03SF, CD06, CD52) and 3 biological replicates/line; t test; data are presented as the

mean ± SEM.

(B) RT–qPCR analysis of *CD4* and *CCR5* mRNA expression in iPSCs and iPSC-derived microglia; n=3

cell lines (03SF, CD06, CD52) and 3 biological replicates/line; t test; data are presented as the mean ±

SEM.

Figure S2: PCR analysis of 2-LTR circles

(A) A schematic representation of HIV-1 structure.

(B) A schematic representation of an HIV-1 2-LTR circle.

(C) PCR analysis of 2-LTR circles on day 4 post HIV-1 JRFL infection and mock infection.

Figure S3: RNA-seq analysis of HIV-1 JRFL- vs mock-infected 03SF microglia

(A) A PCA plot of RNA-seq data collected from HIV-1 JRFL-infected microglia vs mock-infected

microglia on day 16; n=3 independent infections of 03SF microglia.

(B) A REVIGO treemap indicating major dysregulated pathways from gene ontology (GO) analysis of

HIV-1 JRFL-infected microglia vs mock-infected microglia on day 16; n=3 independent infections of

03SF microglia.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Supplemental Figure 1

Supplemental Figure 2

Supplemental Figure 3