NRC immune receptor networks show diversified hierarchical genetic architecture across plant lineages

Foong-Jing Goh[1,2,3], Ching-Yi Huang[1], Lida Derevnina[4], and Chih-Hang Wu[1,2,5*]

[1]Institute of Plant and Microbial Biology, Academia Sinica, Taipei, Taiwan

[2]Molecular and Biological Agricultural Sciences Program, Taiwan International Graduate Program, National Chung-Hsing University and Academia Sinica, Taipei, Taiwan

[3]Graduate Institute of Biotechnology, National Chung-Hsing University, Taichung, Taiwan

[4]Crop Science Centre, Department of Plant Science, University of Cambridge, Cambridge, United Kingdom

[5]Biotechnology Center, National Chung-Hsing University, Taichung, Taiwan

*Correspondence: wuchh@gate.sinica.edu.tw

Short title: Diversity of NRC networks in asterids

One-sentence summary: The NRC networks show degrees of complexity across asterids, encompassing largely conserved NRC0 networks and diversified family-specific NRC networks

Abstract

Plants developed sophisticated immune systems with nucleotide-binding domain and leucine-rich repeat-containing (NLR) proteins to repel invading pathogens. The NRC (NLR required for cell death) family includes helper NLRs that form a complex genetic network with multiple sensor NLRs to provide resistance against pathogens of solanaceous plants. However, the evolution and function of NRC networks outside solanaceous plants is currently unknown. We conducted phylogenomic and macroevolutionary analyses comparing NLRs identified from different asterids lineages and found that NRC networks expanded significantly in most lamiids but not in Ericales and campanulids. Using transient expression assays in *Nicotiana benthamiana*, we show that NRC networks are simple in Ericales and campanulids, but are with high complexity in lamiids. Phylogenetic analyses grouped the NRC helper NLRs into three NRC0 subclades that are conserved, and several family-specific NRC subclades of lamiids that show signatures of diversifying selection. Functional analyses revealed that members of the NRC0 subclades are partially interchangeable, whereas family-specific NRC members in lamiids lack

interchangeability. Our findings highlight the distinctive evolutionary patterns of the NRC
networks in asterids and provide potential insights into transferring disease resistance across plant
lineages.

Keywords: plant immunity, disease resistance genes, nucleotide-binding domain and leucine-rich
repeat-containing protein (NLR), NLR-required for cell death (NRC), immune receptor network,
evolution, asterids

Introduction

Plants have evolved intricate immune systems to protect themselves from pathogen invasion.
Intracellular nucleotide-binding domain and leucine-rich repeat (NLR) immune receptors play
major roles in plant immunity by detecting effector proteins delivered from pathogens (Dodds and
Rathjen 2010; Jones et al. 2016; Ngou et al. 2022). NLR activation often results in a form of
programmed cell death known as the hypersensitive response, leading to the restriction of pathogen
growth. NLRs exhibit a conserved tripartite structure comprised of an N-terminal domain involved
in cell death initiation, a central nucleotide-binding domain (NBD) involved in activation, and a
C-terminal leucine-rich repeat (LRR) domain involved in ligand binding and NLR self-regulation
(Duxbury et al. 2021). NLRs are classified based on their N-terminal domains into TNLs that
contain Toll/interleukin-1 receptor/R protein (TIR) domains, RNLs that contain Resistance to
Powdery mildew 8 (RPW8)-like coiled-coil (CC) domains, and CNLs that contain G10-type
coiled-coil domains (CC_{G10}-NLR) or Rx-type coiled-coil domains (CC_{Rx}-NLR) (Shao et al. 2016;
Duxbury et al. 2021; Kourelis et al. 2021; Lee et al. 2021).

Molecular and genetic studies have categorised NLRs into three groups based on their mode of
action: singletons, pairs, and networks (Adachi et al. 2019b; Contreras et al. 2023a). Singleton
NLRs can directly or indirectly detect pathogen effectors and initiate downstream immune
responses without the assistance of additional NLRs (Contreras et al. 2023s). The CNLs ZAR1
(HopZ-Activated Resistance 1) from *Arabidopsis thaliana* and Sr35 from wheat are two well-
studied examples of singleton NLRs (Wang et al. 2015, 2019; Förderer et al. 2022). ZAR1
indirectly recognizes effectors via its RLCK (Receptor-like cytoplasmic kinase) partners and form
pentameric membrane-associated resistosome complexes that function as calcium-permeable
channels to activate immune responses (Wang et al. 2019; Bi et al. 2021). Sr35 directly binds the
stem rust effector AvrS35 through its LRR domain, forming a pentameric resistosome complex
similar to ZAR1 (Förderer et al. 2022). NLRs can also function in pairs, in which a sensor NLR,
specialised to recognise pathogen effectors, is coupled with a helper NLR that is involved in
immune signalling. Paired NLRs often physically interact and exist as linked gene pairs or clusters
on the chromosomes, suggesting that they are co-regulated and may have co-evolved exclusively

with each other throughout their evolutionary trajectory (Xi et al. 2022). Classical examples of
paired NLRs include RRS1/RPS4 of *A. thaliana*, and RGA4/RGA5 and Pik-1/Pik-2 of rice
(Ashikawa et al. 2008; Narusaka et al. 2009; Césari et al. 2014; Sohn et al. 2014; Xi et al. 2022).
In these pairs, the sensor NLRs often contain integrated domains (ID) that play critical roles in
sensing pathogen effectors; for example, the WRKY domain in RRS1 and HMA (Heavy-Metal-
Associated) domain in RGA5 and Pik-1 (Cesari et al. 2013; Maqbool et al. 2015; Sarris et al. 2015;
Marchal et al. 2022).

NLRs can also function in networks, in which multiple sensor NLRs that detect different pathogen
effectors signal through a set of helper NLRs to mediate immunity (Wu et al. 2017, 2018). NRG1
(N-Required Gene 1) and ADR1 (Activated Disease Resistance 1), two RNL-type helper NLRs,
are required by multiple sensor NLRs for immune signalling (Castel et al. 2019; Saile et al. 2020).
In Arabidopsis, the TNLs RPS4/RRS1, RPP2, and RPS6, preferentially signal through NRG1,
whereas the CNLs RPS2 and RPP4 preferentially signal through ADR1, supporting the idea that
the two groups of helper NLRs show complex genetic redundancy (Saile et al. 2020). Similar to
NRG1 and ADR1, NRCs (NLR-required for cell death) are required by multiple Solanaceae sensor
NLRs. Within the NRC network, the NRC-dependent sensor NLRs detect different pathogen
effectors and require partially redundant helper NRCs for immune response. For example, sensor
NLRs Rpi-blb2, Mi-1.2, and R1 transmit signals via NRC4 to trigger cell death. The sensor NLR
Prf and Rpi-amr1 operates with either NRC2 or NRC3, while Sw5b, R8, Rx, Bs2, and Rpi-amr3
can signal redundantly through NRC2, NRC3, or NRC4 (Wu et al. 2016, 2017; Chen et al. 2021;
Witek et al. 2021; Wu and Kamoun 2021; Lin et al. 2022; Zhang et al. 2024).

Phylogenetically, NRCs and NRC-dependent sensor NLRs form the NRC superclade, which is a
subgroup of CC_{Rx}-NLR (Wu et al. 2017; Kourelis et al. 2021; Contreras et al. 2023a). The NRC
superclade is present across the asterids and in some Caryophyllales, but is not present in monocots
or rosids. This suggests that the ancestral sequences of the NRC superclade arose before the
diversification of Caryophyllales and asterids (Wu et al. 2017). A recent report defined NRC0, an
NRC helper NLR, as the only conserved family member found in various asterids (Sakai et al.
2023). NRC0 often exists in a gene cluster with NRC0-dependent sensor NLRs, strengthening the
hypothesis that the NRC superclade originated from an ancient helper-sensor NLR gene cluster
(Wu et al. 2017; Sakai et al. 2023). Another intriguing evolutionary aspect of the NRC superclade
is the presence of noncanonical extended N-terminal domains (exNT), which exist before the CC
domain in some clades of sensor NLRs (Seong et al. 2020). Although the detailed evolutionary
history of the sensor NLRs is not clear, the exNT were also found in sugar beet, suggesting that
the exNT of NRC-dependent sensor NLRs emerged between the common ancestor of asterids and
Amaranthaceae (Caryophyllales) (Seong et al. 2020).

Upon sensor NLR activation, NRCs form higher-molecular-weight complexes that localize to the
plasma membrane and likely act as calcium-permeable channels similar to ZAR1 (Duggan et al.
2021; Ahn et al. 2023; Contreras et al. 2023b). The first alpha helixes of most NRCs possess a

conserved α1 helix domain known as the MADA motif that plays a major role in executing cell death (Adachi et al. 2019a). The MADA motif is conserved in ZAR1 and many other singleton NLRs but is absent in the NRC-dependent sensor NLRs, suggesting that degeneration of the MADA motif may be an evolutionary feature associated with the functional diversification of the NRC superclade members (Adachi et al. 2019a). Interestingly, members of the NRC family have also evolved functions beyond triggering cell death induced by sensor NLRs. The cell surface receptor Cf-4 induces NRC3-dependent hypersensitive responses upon detection of the plant pathogen effector AVR4, indicating that the NRC superclade not only functions in intracellular receptor defence but also contributes to cell surface receptor-mediated defences (Kourelis et al. 2022). Furthermore, NRCx, an unusual NRC family member lacking a functional MADA motif, modulates NRC2/NRC3-mediated cell death in *Nicotiana benthamiana* plants. NRCx functions as a negative regulator of cell death execution, thus, playing a key role in maintaining homeostasis within the NRC network (Adachi et al. 2023a).

NLRs represent one of the most diverse protein families in angiosperms, with many species encoding large and diverse repertoires of NLR genes in the genome (Barragan and Weigel 2021). As they play key roles in the survival of a plant species under emerging pathogen pressures, *NLR* genes are known to show distinguished signs of rapid evolution, even within a single species (Kuang et al. 2004; Jacob et al. 2013; Van De Weyer et al. 2019). Given that NLRs exhibit a high turnover rate, the birth-and-death model has been proposed to describe the evolutionary process of *NLR* genes (Michelmore and Meyers 1998). In this model, the emergence of new NLRs occurs through repetitive cycles of gene duplication. Some genes are maintained in the genome and develop novel capabilities to detect pathogens, while others are either lost or nonfunctionalized due to the accumulation of deleterious mutations (Michelmore and Meyers 1998; Barragan and Weigel 2021). These dynamic evolutionary patterns allow the plant immune system to effectively adapt to the rapidly evolving effector repertoires of pathogenic microbes. Indeed, some NLRs have been lost in certain plant lineages but highly expanded in other plant lineages (Shao et al. 2016; Martin et al. 2023). TNLs and NRG1, along with their associated signalling partner SAG101, have been lost in monocots, some dicots, and some Magnoliids (Tarr and Alexander 2009; Collier et al. 2011; Shao et al. 2016; Liu et al. 2021; Wu et al. 2021). However, TNL and RNL are more highly expanded and diversified in Gymnosperms and Rosids (Terefe-Ayana et al. 2012; Shao et al. 2016; Van De Weyer et al. 2019; Van Ghelder et al. 2019; Woudstra et al. 2023). CNLs are in general more abundant in the genomes of most angiosperms, whereas TNLs have been lost frequently in dicots (Liu et al. 2021). Studies of wild tomato and Arabidopsis demonstrated a wide range of NLR polymorphisms within single-species populations, suggesting a correlation between plant adaptation to pathogens and NLR diversity (Stam et al. 2019; Van De Weyer et al. 2019). These NLR polymorphisms may contribute to species-specific signalling pathways in NLR-mediated resistance. For example, CaRpi-blb2, a pepper homolog of wild tomato Rpi-blb2, initiates cell death via NRC8 and NRC9 which were identified only in pepper but not other solanaceous species, highlighting the contribution of lineage-specific NLR clades that emerged recently in resistance against pathogens (Oh et al. 2023).

The availability and improvement of genome databases have enabled the analyses of NLR genes from many model and non-model species, leading to a more comprehensive understanding of NLR evolution and diversity (Thind et al. 2018; Van De Weyer et al. 2019; Barragan and Weigel 2021; Kourelis et al. 2021; Ence et al. 2022). Plant NLRs likely originated from the common ancestor of green algae before rapidly diversifying and evolving in land plants (Gao et al. 2018; Ortiz and Dodds 2018; Shao et al. 2019). The discovery of the MAEPL motif and the MADA motif, which are both crucial in executing cell death, provides evidence of the shared function of the CC domain between non-flowering and flowering plants, strengthening the concept of the origin of NLR from common green algae (Adachi et al. 2019a; Chia et al. 2022). While a high number of NLR genes could offer potential survival advantages, several factors have been proposed to influence the number of NLR genes in a particular plant species. Adaptations to aquatic, parasitic, and carnivorous lifestyles correlated to low NLR numbers among closely related species (Baggs et al. 2020; Liu et al. 2021). For example, the aquatic plant duckweed (*Lemna minor*) has only 11 NLR genes, and the carnivorous aquatic plant *Utricularia gibba* has completely lost all NLR genes (Baggs et al. 2020; Liu et al. 2021). To date, close to 500 NLRs from 31 genera belonging to 11 orders of flowering plants have been experimentally validated (Kourelis et al. 2021). These NLRs emerged in different plant species and likely diversified to function against various pathogens. The diversity of NLRs found in plants may reflect the outcome of their long-standing coevolutionary arms race with pathogens (Upson et al. 2018).

Asterids constitute a highly diverse group of angiosperms and encompass numerous economically important crops. The NRC superclade exists in several asterids species, suggesting that NRC networks may play roles in disease resistance in these plants (Wu et al. 2017; Sakai et al. 2023). Understanding the diversity of NRC networks across asterids may provide useful information on lineage- or species-specific sensor-helper NLR connections, which could be useful for disease resistance breeding and help facilitate interspecies resistance gene transfer. However, our knowledge of the evolutionary history and diversity of NRC networks beyond Solanaceae is limited. To address this, we performed macroevolutionary analyses of NRC networks across asterids and experimentally validated the sensor-helper dependency using heterologous expression in *N. benthamiana*. We found that the NRC superclade displays distinct duplication and expansion patterns in the three asterids lineages (Ericales, campanulids, and lamiids). Combined with the results from transient expression assays in *N. benthamiana*, we revealed that the NRC networks in Ericales, campanulids, and lamiids show different complexity and hierarchical structures. The NRC helper NLR family can be further grouped into three NRC0 subclades that are conserved, and several family-specific NRC subclades of lamiids that show signatures of diversifying selection. Further inter-species comparisons revealed that members of the NRC0 subclades were partially interchangeable, as many of them can function with NRC0-dependent sensor NLRs from different lineages. In contrast, members of the family-specific NRC subclades of lamiids lack interchangeability, with only some of the NRCs showing compatibility with sensor NLRs across plant families. Our study sheds light on the unique evolutionary patterns of the NRC networks

within the asterids and offers valuable insights into the potential transfer of disease resistance
mechanisms across different plants.

Results

The NRC superclade is expanded differentially in distinct plant lineages

To understand the evolutionary diversity of the NRC network in plants, we performed comparative
phylogenomic analyses of 46 angiosperms, including basal angiosperms, monocots, rosids,
Caryophyllales, and asterids (comprising Ericales, campanulids and lamiids) (Supplemental Data
Set 1). We used MAST - MEME Suite and 20 previously defined motifs to predict potential NLR
encoding sequences from the genome of all these angiosperms (Jupe et al. 2012). After generating
the NLR phylogenetic trees of each species using the conserved NB-ARC domain, we grouped
these NLRs into four categories including TNL, RNL, CC_{G10}-NLR, and the CC_{Rx}-NLR. To
identify the NRC superclades in these species, we performed additional phylogenetic analyses of
the CC_{Rx}-NLR using the solanaceous NLRs (NRC, NRC-dependent and NRC-independent NLRs)
as references. These second phylogenetic trees classified the CC_{Rx}-NLR into the NRC superclade
and non-NRC superclade CNLs. Consistent with the previous finding, the NRC superclade was
found only in species of some Caryophyllales and most asterids (Supplemental Data Set 2) (Wu et
al. 2017).

Next, we calculated the percentage of NRC superclade members relative to the total number of
NLRs in these genomes (Fig. 1A). Interestingly, the percentages of NRC superclade members
relative to the total number of NLRs show striking variations across different plant lineages. In
Caryophyllales, two out of the seven species we analysed contained the NRC superclade (Fig. 1A).
Sugar beet (*Beta vulgaris*) and carnation (*Dianthus caryophyllus*) have approximately 4% and 15%
of their total NLRs belonging to the NRC superclade, respectively. In Ericales, all three species
we analysed contain members of the NRC superclade (Fig. 1A). The percentages of NRC
superclade members out of total NLR in kiwifruit (*Actinidia chinensis*), tea (*Camelia sinensis*) and
miracle fruit (*Synsepalum dulcificum*) are 7%, 3% and 1%, respectively. In campanulids, the
percentage of NLRs belonging to the NRC superclade is variable, with an average of 8% out of
total NLRs (Fig. 1A). Among the campanulids analysed, ginseng (*Panax ginseng*) has the lowest
percentage of NRC superclade members, with 3% (4 out of 131) of its NLRs belonging to the NRC
superclade; candyleaf (*Stevia rebaudiana*) has the highest percentage, with 14% (44 out of 298)

of its NLRs belonging to the NRC superclade. In lamiids, the proportions of NRC superclade
members among total NLRs are highly variable (Fig. 1A). Overall, the number of NRC superclade
members in most lamiids is higher than that of Caryophyllales, Ericales, and campanulids. In most
lamiids species, over 40% of the total NLRs belong to the NRC superclade. In solanaceous plants
(such as tomato *Solanum lycopersicum* and *N. benthamiana*), where the NRC superclade and NRC

network were originally described (Wu et al., 2017), around 50% of total NLRs belong to the NRC superclade. In contrast, over 75% of total NLRs from the three related *Ipomoea* species (sweet potato, *Ipomoea batatas*; threefork morning glory, *Ipomoea trifida*; and littlebell, *Ipomoea triloba*) belong to the NRC superclade (Fig. 1A). Remarkably, Asiatic witchweed (*Striga asiatica*), a parasitic plant, has the highest percentage (89%) of NRC superclade members among all analysed plant species, while dodder (*Cuscuta australis*), another parasitic plant, has the lowest percentage (14%) of NRC superclade members among lamiids, with only one out of seven NLRs belonging to the superclade. Hardy rubber tree (*Eucommia ulmoides*) was the only lamiids species where the NRC superclade was not found (Fig. 1A). These results suggest that the NRC superclade shows very different evolutionary trajectories among plant lineages.

To understand the degree to which the expansion of the NRC superclade and other NLRs correlates with genome expansion, we compared the size of the genome or the number of protein-coding genes with the number of NLRs or the size of the NRC superclade of each species. The data revealed that there was no correlation between genome size with the amounts of NLRs or the size of the NRC superclade (Supplemental Fig. S1A and S1B), but there was a weak correlation between NLRs and the numbers of total protein-coding genes (R=0.36, p=0.052) (Supplemental Fig. S1C). Additionally, the size of the NRC superclade was very weakly correlated with total protein-coding genes (R=0.28, p=0.13) (Fig. 1B), suggesting that the overall evolutionary pattern of total NLRs is not well aligned with that of the NRC superclade in these species. To further compare the features of the NRC superclade and NLR expansion in different plant lineages, we assessed the number of total NLR/total protein-coding genes and the size of the NRC superclade/total protein-coding genes. Most lamiids had higher ratios of total NRCs to total protein-coding genes compared to other plant lineages (Supplemental Fig. S1D). Based on these analyses, we concluded that the expansion of the NRC superclade is a unique characteristic of lamiids.

The NRC superclade encompasses two types of NLRs, namely the NRC-dependent sensor NLRs (NRC-S) and the NRC helper NLRs (NRC-H). Since the NRC superclade originated from a linked NLR pair or cluster, we expect the ancestral sensor-helper ratio of the NRC superclade to be around 1:1 or 2:1. To gain further insights into whether the sensor and helper NLRs within the NRC superclade have expanded differentially during evolution, we calculated the NRC sensor-helper ratio for all NRC superclade containing plant species (Fig. 1C). Most of the species have NRC sensor-helper ratios greater than 2:1, suggesting that the NRC-S experienced more frequent duplication events in comparison to NRC-H. In Ericales and campanulids, where the NRC superclade constitutes only a small portion of total NLRs, the NRC sensor-helper ratio varied. In Ericales, the NRC sensor-helper ratio ranged from 1:1 to 4:1, while in campanulids this ratio ranged from 3:1 to 14:1. In lamiids, where the NRC superclade is often extensively expanded, most species displayed sensor-helper ratios higher than the ancestral ratio (Fig. 1C). Together, these findings indicate that sensor NLRs are more prone to duplication during evolution compared

to helper NLRs, supporting the notion that sensor NLRs require diversification to recognise
various pathogens, while helper NLRs are involved in mediating a conserved cell death pathway.

The NRC network in Ericales is simple

To characterise the NRC superclade in Ericales, we performed phylogenetic analysis on NLRs
identified from *A. chinensis*, *C. sinensis*, and *S. dulcificum* using tomato NRC superclade members
as a reference (Fig. 2, A and B). The Ericales NRC superclade can be divided into one NRC-H
clade and one NRC-S clade. To understand the phylogenetic relationship of the Ericales NRCs
with the NRC0 subclade described by Sakai et al. (2023), we performed phylogenetic analyses
encompassing all the predicted NRC-H from asterids. We found the Ericales NRC-H clustered
together with the NRC0 subclade, but formed its lineage-specific group (Supplemental Fig S2A).
Therefore, to differentiate this group of NRCs from the NRC0 subclade, we named the members
of this subclade NRC0-Ericales-specific (NRC0-Eri). In several plant species, NRC0 orthologs are
physically clustered with the NRC0-dependent sensor NLRs on the chromosome (Sakai et al.
2023). We found a putative NRC-S linked to one of the NRC0-Eri in *C. sinensis* (CsNRC0b-Eri,
CSS0008112.1) (Supplemental Fig. S2B and C), but no sensor-helper linkage was found in *A.
chinensis* or *S. dulcificum*.

To validate that the members of NRC0-Eri function as helper NLRs for the putative sensor NLRs,
we cloned NRC0 from *A. chinensis* (AcNRC0-Eri, Achn230571) and *C. sinensis* (CsNRC0-Eri)
and performed transient expression assays in *N. benthamiana* together with putative sensor NLRs
from the same species. As our cloned CsNRC0b-Eri (CSS0008112.1) contains a premature stop
codon in the NB-ARC domain and expression of AcNRC0-Eri alone induced cell death in wild
type and *nrc2/3/4*_KO *N. benthamiana,* we focused on CsNRC0a-Eri (CSS0010223.1) for
validation of the Ericales NRC network (Supplemental Fig. S2D). Out of the 7 putative NRC-
dependent sensor NLRs annotated in the genome of *C. sinensis*, two contained full-length NLR
sequence signatures, and the remaining five were truncated, either missing the coiled-coil domains
or part of the LRR domains. We cloned the two full-length *C. sinensis* NLRs (Cs0021741 and
Cs0024074) and introduced a D to V mutation into their MHD motifs to generate constitutively
active variants (Cs0021741^{DV} and Cs0024074^{DV}). Co-expression of Cs0024074^{DV}, but not
Cs0021741^{DV}, with CsNRC0a-Eri induced cell death in *N. benthamiana*. Cell death was not
observed when either Cs0024074^{DV} or CsNRC0a-Eri were expressed alone, indicating that
CsNRC0a-Eri is required for Cs0024074 cell death activity (Fig. 2C). Overall, our data suggests
that CsNRC0a-Eri functions as a helper NLR. Furthermore, the NRC network of Ericales is simple,
with few sensor NLRs signalling through one or two NRC0 homologs (Fig. 2D).

NRCs duplicated in some campanulids, forming an NLR network with partially redundant nodes

To characterise the NRC superclade in campanulids, we performed phylogenetic analyses of NLRs identified from seven species, including *Chrysanthemum nankingense*, lettuce (*Lactuca sativa*), sunflower (*Helianthus annuus*), *S. rebaudiana,* carrot (*Daucus carota*), celery (*Apium graveolens*), and *P. ginseng*, across two major orders (Asterales and Apiales) (Supplemental Fig. S3A). In line with our previous findings, the campanulids NLRs were classified into several highly supported clades consisting of TNLs, RNLs, CC_{G10}-NLRs and CC_{Rx}-NLRs, which include the NRC superclade (Fig. 3A and Supplemental Fig. S3A). Phylogenetic analysis revealed that the campanulids NRC superclade is divided into multiple clades, including one NRC-H clade and several putative NRC-S clades (Fig. 3B). The NRC-H clade can be further divided into the NRC0 subclade, which contains sequences from both Asterales, Apiales and the reference tomato NRC0, as well as another subclade which contains sequences only from the Asterales (Fig. 3B). Based on the phylogenetic tree of NRC-H from asterids, this subclade clusters together with NRC0-Eri and the NRC0 subclade defined by Sakai et al. (2023) (Supplemental Fig. S2A). Therefore, we named this group the NRC0-Asterales-specific (NRC0-Ast) subclade.

To determine the genetic structure of the campanulids NRC network, we selected *D. carota* from Apiales and *L. sativa* from Asterales as representative species. The carrot NRC superclade is comprised of DcNRC0 (DCAR_023561) and eight sensor NLRs falling into two subgroups (Fig. 3B and Supplemental Fig. S3B). Out of the eight sensor NLRs, two of them (Dc23557 and Dc23650) are located in a gene cluster together with DcNRC0 (Supplemental Fig. S3, C and D). We cloned four putative sensor NLRs, including Dc23557 and Dc23650, and introduced a D to V mutation into their MHD motifs and then co-infiltrated them independently with DcNRC0. While two of the sensor NLRs (Dc11400^{DV} and Dc12621^{DV}) did not induce cell death with DcNRC0, Dc23650^{DV} and Dc23557^{DV} induced different intensities of cell death in the presence of DcNRC0 (Supplemental Fig. S3E). The lettuce NRC superclade is composed of LsNRC0 (Lsat_1_v5_gn_3_132141.1), LsNRC0-Ast (Lsat_1_v5_gn_8_124621), and 15 sensor NLRs. Most of these sensor NLRs are clustered into two sub-groups (Supplemental Fig. S4A). Interestingly, LsNRC0-Ast is located in proximity to some of the sensor NLRs whereas LsNRC0 is located on a different chromosome (Supplemental Fig. S4, B and C). Similar to the previous experiment, we introduced autoactive mutations (D to V) into the MHD motif of the sensor NLRs, and then performed cell death assays by co-agroinfiltration with either LsNRC0 or LsNRC0-Ast on *N. benthamiana* leaves. Three members of the sensor NLRs within the same sub-group (Ls35940^{DV}, Ls28800^{DV}, and Ls36021^{DV}) induced cell death when co-expressed with LsNRC0 but not LsNRC0-Ast. Two putative sensor NLRs within another sub-group (Ls123301^{DV} and Ls124100^{DV}) induced cell death when co-expressed with LsNRC0-Ast but not LsNRC0. (Fig. 3C). Interestingly, one of these putative sensor NLRs (Ls124601^{DV}) induced cell death when coexpressed with either LsNRC0 or LsNRC0-Ast (Fig. 3C). These results indicate that the sensor NLRs of lettuce are divided into the NRC0-dependent and NRC0-Ast-dependent groups, with

some sensor NLRs capable of signalling through both NRCs (Fig. 3, C and D). Altogether, these
results suggest that, in campanulids, some orders such as Apiales (carrot) harbour a simple NRC
network composed of NRC0 and matching sensor NLRs, whereas the NRC network in other orders
such as Asterales (lettuce) contain at least two partially redundant NRCs, namely NRC0 and
NRC0-Ast, and the matching sensor NLRs (Fig. 3D).

To gain insight into the interchangeability of NRC networks across different species of
campanulids, we conducted cross-species comparisons using NRC helper and sensor NLR from
both carrot and lettuce. We found that two LsNRC0-dependent sensor NLRs (Ls36021[DV] and
Ls124601[DV]) trigger cell death when co-expressed with DcNRC0 (Supplemental Fig. S5, A and
B). The sensor NLR Ls28800[DV] triggered weak but statistically significant cell death when co-
expressed with DcNRC0 (Supplemental Fig. S5, A and B). The lettuce sensor NLRs Ls124041[DV]
and Ls123301[DV], which triggered cell death when co-expressed with LsNRC0-Ast, did not trigger
cell death when co-expressed with DcNRC0 (Supplemental Fig. S5, A and B). Interestingly, carrot
sensor NLR Dc23560[DV], which induces cell death together with DcNRC0, triggers clear cell death
with LsNRC0-Ast and weak but statistically significant cell death with LsNRC0 (Supplemental
Fig. S5, C and D). These results revealed that DcNRC0, LsNRC0 and LsNRC0-Ast are partially
interchangeable, displaying some compatibility with sensor NLRs across the two species.

The NRC network is highly expanded in most lamiids

To compare the NRC superclade of different lamiids, we selected 5 lamiids species, including
monkeyflower (*Erythranthe guttata*), European ash trees (*Fraxinus excelsior*), coffee (*Coffea
canephora*), *S. lycopersicum*, and water spinach (*Ipomoea aquatica*), and generated a phylogenetic
tree encompassing their NRC superclade members. This tree showed that, within the NRC
superclade, the NRC-H clade was the only well-supported clade that contained NLR sequences
from all five species (Fig. 4A). Most of the well-supported NRC-S clades only contained
sequences from single species. This is consistent with the view that sensor NLRs are more
diversified than helper NLRs in different lamiids lineages. Further phylogenetic analyses revealed
that the NRC family is also highly diverse, with only NRC0 from the five selected species forming
a well-supported clade (Fig. 4B). The other NRCs across the five species were rarely found to
cluster together, indicating that the NRC family has extensively diversified across different lamiids
species.

We then sought to reconstitute the NRC network using sequences identified from *Ipomoea* species.
The *I. aquatica* NRC superclade consists of 18 NRC-H and 31 NRC-S, which is less expanded
than that of *I. triloba*, *I. trifida* and *I. batatas* (Fig. 4C, and Supplemental Data Set 2). Given that
many of the orthologs of *I. triloba* NRC-H and NRC-S can be identified from *I. aquatica* in the
phylogenetic analysis, we reasoned that *I. aquatica* harbours a simplified NRC network compared
to that of *I. triloba* (Supplemental Fig. S6, A and B). Therefore, we used *I. aquatica* (Ia) as our
primary species for cloning *Ipomoea* NRC-H/S, and *I. triloba* (Itb) as an alternative species when

our attempts at cloning failed. Since most of these NRCs are not orthologous to the previously
described solanaceous NRCs, we name these 18 NRCs Convolvulaceae(Con)-Ia/ItbNRCs to
differentiate them from NRCs identified from the Solanaceae (Sol). In addition to the IaNRC0
(GWHPABKX015475), some of the Con-IaNRCs are located in gene clusters together with
putative NRC-S (Supplemental Fig. S6, C and D). We successfully cloned 10 NRCs and 20
putative NRC-S from *I. aquatica* or *I. triloba* and performed cell death assays as described
previously (Supplemental Data Set 3). As expected, the sensor NLR Ia15471^{DV} induced cell death
when co-expressed with its physically linked helper NLR, IaNRC0, but not with other Con-
IaNRC/ItbNRCs (Fig. 4C and Supplemental Fig. S7). An additional 13 sensor NLR and Con-
IaNRC/ItbNRCs pairings were shown to function together, displaying strong or weak cell death
(see method for definition) when co-expressed in *N. benthamiana* (Fig. 4C and Supplemental Fig.
S7). These results indicate that *I. aquatica* contains two independent NRC networks: one simple
network with NRC0, similar to what was observed in Ericales and campanulids; and one complex
network with multiple NRCs showing varying degrees of genetic redundancy, similar to the
phenomenon observed in Solanaceae (Supplemental Fig. S8).

Family-specific NRC subclades of lamiids show features of diversifying selection

To gain further insights into the evolution of NRC helper NLRs across asterids, we performed
phylogenetic analyses using full-length NRC sequences, with the tomato NRC0-dependent sensor
NLR (Sl08230) as an outgroup. This phylogenetic tree classified the NRCs into the NRC0
subclades (NRC0, NRC0-Eri, NRC0-Ast) and several lineage-specific NRC subclades of lamiids,
consistent with the previous analysis using only the NB-ARC domain (Fig. 5A, Supplemental Fig.
S9, and Supplemental Fig. S2). In lamiids, the NRC subclades grouped together in a manner
largely consistent with their respective family taxonomy (Fig. 5A, Supplemental Fig. S9, and
Supplemental Fig. S2). Consequently, we referred to these clusters as family-specific NRC
subclades. Next, we performed an adaptive branch-site REL test for episodic diversification
(aBSREL), and found signatures of diversifying selection in several of the lamiids family-specific
NRC subclades, but not in the NRC0 subclades (Fig. 5A, Supplemental Fig. S9, and Supplemental
Data Set 4). Furthermore, NRC0 subclades showed shorter branch lengths in general, whereas the
family-specific NRC subclades showed more variable and longer branch lengths (Fig. 5, B and C,
and Supplemental Fig. S9). These results suggest that the three NRC0 subclades are less diversified
than the family-specific NRC subclades of lamiids.

NRC0 subclades across asterids are partially interchangeable

To test the degree to which members of the NRC0 subclades (NRC0, NRC0-Eri, and NRC0-Ast)
are interchangeable, we conducted cell death experiments using NRC0 homologs cloned from *C.*
sinensis, D. carota, L. sativa, S. lycopersicum, I. aquatica. We found that tea NRC0-dependent
sensor NLR Cs24074^{DV} triggered clear cell death when co-expressed with DcNRC0, IaNRC0 and
SlNRC0, but no cell death with LsNRC0 or LsNRC0-Ast (Fig. 6, A and E). Out of the two carrot

NRC0-dependent sensor NLRs, only one (Dc23650^{DV}) induced cell death when co-expressed with
LsNRC0-Ast, LsNRC0, IaNRC0, and SlNRC0 (Fig. 6, B and E). The lettuce NRC0-dependent
sensor NLRs Ls28800^{DV}, Ls35940^{DV} and Ls36021^{DV}, which did not induce cell death when
expressed with LsNRC0-Ast, induced cell death when co-expressed with most of the NRC0 from
the other species, although the cell death with IaNRC0 tends to be weaker for Ls35940^{DV} and
Ls36021^{DV} (Fig. 6, C and E). The lettuce NRC0-Ast-dependent sensor NLRs Ls123301^{DV} and
Ls124100^{DV} failed to induce cell death with any of the other tested NRC0 variants. Interestingly,
the lettuce NRC-S Ls124601^{DV}, which can signal through both LsNRC0 and LsNRC0-Ast, is the
only NRC-S that activates cell death with all NRC0 variants tested (Fig. 6, C and E). The water
spinach sensor NLR Ia15471^{DV} only induced cell death with SlNRC0 or IaNRC0 but not any of
the other tested NRC0 variants. Similarly, the tomato NRC0-dependent sensor NLR Sl08230^{DV}
only induced cell death with IaNRC0 or SlNRC0 (Fig. 6, D and E). These results indicate that
NRC0 homologs exhibit partial interchangeability across different lineages of asterids. Pair-wise
sequence comparison of different NRC0 subclades members indicated that LsNRC0-Ast shows
lower similarity to other NRC0 subclades members, largely consistent with the results that
LsNRC0-Ast displays overall more different sensor compatibility compared to other NRC0 tested
(Supplemental Fig. S10A, and Fig. 6E). In the pairwise sensor NLR comparisons, apart from clear
homology among some *D. carota* and *L. sativa* sensor NLRs, Ls124601 displayed slightly higher
overall similarity to other sensor NLRs included in the analysis (Supplemental Fig. S10B).
Intriguingly, Ls124601 was the only sensor NLR that triggered cell death with all the NRC0
homologs tested (Fig. 6C). Together these results suggest that the NRC0 network, across different
asterids lineages, is broadly conserved, however, some of the sensor or helper NLRs have
diversified and lost their compatibility with each other (Fig. 6E).

Sensor and helper NLRs from Solanaceae and Convolvulaceae display some inter-family
cross-compatibility

We previously characterised the NRC network of Solanaceae, which consists of conserved NRC
homologs across most solanaceous plants (Wu et al. 2017). However, phylogenetic analyses of the
NRC superclade here revealed that NRC network members have diversified, and expanded
extensively in different families of lamiids (Fig. 4, A and B, and Fig. 5A). This has led us to
speculate that sensor and helper NLRs of NRC networks from distinct families in lamiids may
share very little or no compatibility. To test this, we co-expressed sensor NLRs from solanaceous
plants with NRCs from *Ipomoea* (representing the Convolvulaceae family), and sensor NLRs from
Ipomoea with NRCs from tomato (representing the Solanaceae family). As expected, most of the
solanaceous NLRs, including Bs2, R1, Prf (Pto/AvrPto), and Rpi-blb2, did not trigger cell death
when co-expressed with any *Ipomoea* NRCs tested (Supplemental Fig. S11). To our surprise, Rx
triggered cell death when co-expressed with Con-IaNRC3, 5, 9, and Con-ItbNRC8 with varying
degrees of intensity (Fig. 7, A and B, Supplemental Fig. S12), and Rpi-amr1 triggered cell death
when co-expressed with Con-IaNRC9 (Fig. 7, A and B, Supplemental Fig. S12). When we tested
the reverse combinations, we found that three of the tested water spinach sensor NLRs failed to

induce cell death with the Sol-SlNRCs (Supplemental Fig. S13). In addition to Ia15471[DV], which
we previously showed to signal through Sol-SlNRC0, two *Ipomoea* sensor NLRs (Itb05590[DV], and
Ia17067[DV]) induced cell death with solanaceous Sol-SlNRC1 but not the other Sol-SlNRCs tested
(Fig. 7, C and D, Supplemental Fig. S14). However, phylogenetic analysis reveals no clear
orthology between Sol-SlNRC1 and any Con-NRCs (Supplemental Fig. S6B), and Itb05590[DV] and
Ia17067[DV] signalled through different sets of Con-NRCs in the earlier analyses (Fig. 4C). The only
group of NRC-S that shows clear homology across these two species are the NRC0-dependent
sensor NLRs (Supplemental Fig. S15). This suggests that, despite having family-specific NRC
networks, some cross-compatibility between sensor and helper NLRs exists between Solanaceae
and Convolvulaceae. Further studies are required to provide mechanistic explanations of this NRC-
S and NRC-H inter-family cross-compatibility.

Discussion

The NRC network plays an essential role in disease resistance to multiple pathogens of
solanaceous plants (Wu et al. 2017). Nevertheless, functional studies of the NRC network beyond
solanaceous plants remained limited. We explored the evolutionary diversity of NRC networks
across lineages of angiosperms, with a particular focus on the three major lineages of asterids
(Ericales, campanulids, and lamiids). These lineages displayed distinct hierarchical structures
within their respective NRC networks (Fig. 8). Ericales represents one of the early branches of
asterids, predating the divergence of campanulids and lamiids. In our study, tea (*C. sinensis*) served
as a representative of Ericales and possessed a relatively simple NRC network. This network
involves the transmission of signals from NRC0-Eri-dependent sensor NLRs through NRC0-Eri
to induce cell death. Since the three examined Ericales species contained few NRC superclade
members, the NRC network in Ericales has likely undergone limited changes over millions of
years of evolution (Fig. 8). In campanulids, we identified two partially redundant NRC nodes:
NRC0 which is present in both Apiales and Asterales, and NRC0-Ast which exists exclusively in
Asterales (Fig. 8). We speculate that NRC0-Ast may have either emerged early in campanulids
and subsequently been lost in Apiales or arose via duplication of the ancestral NRC0 after the
divergence of Asterales and Apiales. The NRC superclade expanded and diversified significantly
in most lamiids (Fig. 1, Fig. 5, and Fig. 8). Apart from NRC0 which is very conserved, the
phylogenetic tree shows distinct lineage-specific clustering patterns of the NRC family members.
This specific diversification pattern appears to be unique for each plant family, implying that the
lineage-specific NRC network is likely largely conserved at the family level (Fig. 5, Fig. 8, and
Supplemental Fig S9). Our results are consistent with the view that the NRC superclade originated
from an ancestral NRC sensor-helper pair, and then differentially expanded across plant lineages
into complicated immune networks (Wu et al. 2017; Sakai et al. 2023).

We proposed that the ancestral state of the NRC network of superasterids (asterids together with
Caryophyllales) is likely a sensor-helper cluster resembling the NRC0-H and NRC0-S described
here and in Sakai et al., (2023) (Wu et al., 2017). Interestingly, the NRC-H identified in sugar beet

(Caryophyllales) does not cluster with the NRC0 clades, which consists of NRC0/NRC0-
Eri/NRC0-Ast, based on phylogenetic analyses (Supplemental Fig. S2A). Therefore, the ancestral
NRC0 may have emerged in asterids from an NRC network shared with Caryophyllales, but this
ancestral NRC network was subsequently lost in asterids. Alternatively, the ancestral NRC0 may
have appeared early in the ancestral superasterids and was later lost in Caryophyllales. More
phylogenetic and functional analyses using the NRC0 sequences identified from basal lineages of
superasterids/asterids should help to clarify the early evolutionary history of the NRC network.
Furthermore, the degree to which the phylogenetically defined Caryophyllales NRC-Hs function
as helper NLRs and their cross-compatibility with sensor NLRs from asterids are still yet to be
determined.

Tandem gene and whole-genome duplication (WGD), genetic drift, relaxed selection during
domestication, and adaptation to ecological niches have all been reported to influence the
expansion and contraction of *NLR* genes (Han and Tsuda 2022). While the total number of protein-
coding genes shows a weak correlation with the expansion of NLRs, it is even more weakly
correlated with the expansion of the NRC superclade (Fig. 1B and Supplemental Fig. S1).
Capsicum annuum, for example, has a large number of NRC superclade members despite having
relatively few total protein-coding genes (Fig. 1A and Supplemental Data Set 2) (Seo et al. 2016).
Conversely, ginseng has high total protein-coding gene numbers but few NRC superclade
members (Fig. 1A and Supplemental Data Set 2). Selection pressures placed on each plant lineage
in asterids may, therefore, have led to distinct expansion or contraction patterns of the NRC
superclade. Furthermore, the evolutionary events leading to the expansion of the NRC superclade
show no correlation to the overall genome size (Supplemental Fig. S1). Since the massive
expansion of the NRC superclade is a unique feature observed in lamiids, we speculate that
selection pressure in the ancestral lamiids species may have led to the initial expansion of the NRC
network. This ancient expansion provided advantages for the species to survive and was further
selected independently in the subsequent lamiids progenies. While most lamiids species encode
large numbers of NRC superclade members, *E. ulmoides* stands out as the only lamiids species
analysed that does not have NLRs belonging to the NRC superclade (Fig. 1A). Although we have
not yet been able to exclude the possibility that this is due to the insufficient quality of the
assembled genome, additional genome information from related species may help address whether
early gene loss events led to the absence of the NRC superclade in this species. Moreover, the
identification of NLRs and overall protein content is influenced by the quality of both assembly
and annotation. Future improvements in assembly and annotation should help to further verify the
expansion and contraction of the NRC superclade in the species analysed here. Notably, 89% of
NLRs predicted from the parasitic plant *S. asiatica* were classified within the NRC superclade,
whereas *Cuscuta* spp., and many other parasitic plants, have undergone substantial loss of their
NLRs (Fig. 1A) (Baggs et al. 2020; Liu et al. 2021). It would be interesting to understand whether
and how the unique lifestyle of these and other parasitic plants may have differentially shaped the
overall evolution of plant immunity, including the NRC superclade.

NLR genes are frequently located within gene clusters which can serve as a repository of genetic variation. Most of these clusters are results of tandem duplication leading to homogeneous NLR clusters, while heterogenous clusters are less prominent (Zhou et al. 2004; Meyers et al. 2005; Jupe et al. 2012). NRC0 stands out as the most conserved NRC homolog among asterids (Sakai et al., 2023). Furthermore, the NRC0 sensor-helper NLR gene cluster was identified in several species, likely representing the ancestral state of the asterids NRC network (Sakai et al., 2023). Based on the observation that most of the members in the NRC network do not remain as sensor-helper NLR gene clusters, we proposed that breaking the physical linkage was critical to enable the independent evolution of both sensor and helper NLRs. In Ericales, most of the sensor NLRs of the NRC superclade appeared to be truncated, and only those linked to NRC0-Eri retained full-length NLR sequence signatures (Fig. 2 and Supplemental Fig. S2). A similar phenomenon was observed in carrots (campanulids), where only the sensor NLRs that were closely linked to NRC0 remained functional (Supplemental Fig. S3). In lettuce, the three sensor NLRs that signal through LsNRC0-Ast are located on the same chromosome as LsNRC0-Ast, whereas LsNRC0 is physically disconnected from the sensor NLRs (Supplemental Fig. S4). The group of sensor NLRs that signal through LsNRC0 show longer branch lengths in the phylogenetic tree compared to the NRC0-Ast-dependent sensor NLRs group (Supplemental Fig. S4A). This supports the hypothesis that, after becoming physically unlinked from their helper NLR partners, the sensor NLRs are more prone to accumulating mutations, leading to diversification or non-functionalisation (Baumgarten et al. 2003; Leister 2004). In water spinach, we identified only a few sensor-helper NLR clusters in addition to the NRC0 cluster (Supplemental Fig. S6). Similar to what was observed in solanaceous crops, most of the water spinach NRC superclade members are dispersed on different chromosomes (Supplemental Fig. S6) (Seo et al. 2016; Wu et al. 2017). These results suggest that the expansion of the NRC superclade coincides with the transposition and tandem duplication of both sensor and helper NLR genes independently. Transposable elements, particularly the long terminal repeat retrotransposons, were shown to contribute to the tandem duplication and transposition of NLR genes in plant genomes (Wei et al. 2002; Kim et al. 2017; Seidl and Thomma 2017; Hao et al. 2023). Whether the associations with transposable elements correlate with the expansion of the NRC superclade in different plant lineages requires further investigation.

The asterids NRC superclade likely originated from an ancient sensor-helper NLR cluster that contained a one-to-one or two-to-one ratio of sensor and helper NLRs (Wu et al. 2017; Sakai et al. 2023). Our results indicate that NRC-S duplicated more times than NRC-Hs in most species analysed and generally showed longer branch lengths than that of helper NLRs in the phylogenetic trees (Fig. 1C, Fig. 2A, Fig. 3A, and Fig. 4A). This is consistent with the view that sensor NLRs are often under high dynamic and balancing selection, whereas helper NLRs show slower evolution rates and remain functionally conserved (Seo et al. 2016; Stam et al. 2019; Shimizu et al. 2022). Furthermore, an expanded and diversified sensor NLR repertoire offers a higher potential for conferring resistance to pathogens (Barragan and Weigel 2021). In campanulids, where the NRC superclade is not as extensively expanded, certain species such as *S. rebaudiana* encode a high sensor-to-helper ratio (Fig. 1C). This is perhaps due to several tandem duplication

and transposition events that occurred to NRC-S but not to NRC-H in these species. Although
sensor NLRs are generally highly expanded in lamiids, leading to a high sensor-to-helper ratio, *I.*
cairica, *I. aquatica*, and *N. benthamiana* are among the few species in which the sensor-to-helper
ratio remains close to the ancestral ratio (Fig. 1C). As NLR contraction events have been reported
in lineages of Brassicaceae and Apiaceae species (Shao et al. 2016; Zhang et al. 2016; Liang and
Dong 2023), one possible explanation is that NLR contraction contributes to the reduction of the
sensor-to-helper ratio, with many of sensor NLRs being lost during evolution. Additional
macroevolutionary analyses with diverse plant genomes may provide insights into how NLR
expansion and contraction events influence sensor-helper NLR ratios and provide information on
the factors that drive NLR contraction.

NRC0 stands out as the most conserved NRC across different plant lineages (Sakai et al. 2023).
We found two other NRC0 subclades, namely NRC0-Eri and NRC0-Ast, which show partial
interchangeability with the NRC0 clade defined by Sakai et al. (2023) (Fig. 6). The three NRC0
subclades show relatively short branch lengths, indicating that NRC0 homologs have not
undergone significant diversification since its split from the ancestral species (Fig. 5). Recent
macroevolutionary studies of ZAR1 homologs revealed that ZAR1 is the most conserved NLR
across angiosperms and evolved to function with partnered RLCKs early in its evolutionary history
(Adachi et al. 2023b). While RLCKs rapidly diversified to keep pace with fast-evolving effectors,
ZAR1 experienced relatively limited expansion and duplication (Adachi et al. 2023b). Whether
the conservation of NRC0 subclades is due to its critical role in immunity or simply a lack of
pathogen pressure remains to be investigated. The phylogenetic tree of the NRC family in lamiids
indicated that, apart from the highly conserved NRC0s, there are no clear orthologous NRCs across
different plant families (Fig. 4B, and Fig. 5A). Nevertheless, some NRC-S from solanaceous plants
can activate some *Ipomoea* NRCs to induce cell death, and some NRC-S from *Ipomoea* are capable
of inducing cell death through SlNRC1 (Fig. 7). Therefore, while NRCs function as helper NLRs
that mediate cell death for NRC-S, this group of genes has diversified to an extent that functional
orthologs (such as NRC0) are rarely found in plants from different families. Despite this,
compatibility between sensor and helper NLR across different plant families has been observed
(e.g. between Solanaceae and Convolvulaceae). Further studies focusing on understanding the
molecular mechanisms underpinning these interactions may help elucidate the critical changes that
determine compatibility between NRCs and their matching sensor NLRs.

We tested many NRC-S and NRC-H combinations using the heterologous expression system in *N.*
benthamiana. Although many of the NRC-S/H combinations induced cell death, some
combinations did not trigger cell death in *N. benthamiana*. In particular, NRC0-Eri was the only
NRC-H group identified in tea and carrot, but not all the NRC-S of tea and carrot can signal
through the tested NRC0-Eri. Similar phenomena were observed with other NRC-S cloned from
L. sativa and *I. aquatica* that did not trigger cell death with any NRC-H tested. While some of
these NRC-S may have accumulated loss-of-function mutations, we could not rule out that
additional putative NRC-H members from these species were missing in the annotated genome, or

an alternative mutation (such as H to R rather than D to V mutation in the MHD motif) in the NLR
is required to trigger cell death, such as the case for NbNRC2 (Derevnina et al., 2021). It is also
possible that an essential component for these NRC-S/H combinations to function is lacking in *N.*
benthamiana. Further studies with improved genome information, alternative mutations, or
expression in the original plant species may help to address this issue.

Our findings reveal the overall diversity and hierarchical structure of NRC networks in plants
belonging to different lineages of asterids. Except for a small number of NRC-H that can cooperate
with NRC-S from different plant lineages, the majority of NRC-H do not function with NRC-S
from divergent plant species. Knowledge of sensor-helper compatibility may be particularly useful
for overcoming restricted taxonomic functionality, a challenge when transferring resistance genes
across distantly related plants (Tai et al. 1999; Narusaka et al. 2013) In the cases of NLRs
belonging to the NRC network, it may be necessary to transfer both the matching sensor and helper
NLRs together. The recent discovery that CaRpi-blb2 specifically signals through the pepper
helper NLRs CaNRC8 and CaNRC9 further emphasizes the importance of understanding the
commonalities and differences in the immunity networks of various related species (Oh et al. 2023).
Among the *Ipomoea* species, *I. trifida, I. tiloba,* and *I. batatas* encode high numbers of overall
NLRs in their genomes. Given that the majority of NLRs in these three species belong to the NRC
superclade, delving into the NRC network of *Ipomoea* species might offer opportunities to utilize
their NLRome as a resource for conferring disease resistance against a variety of pathogens.
Further investigations into the evolution of the NRC network, along with validation using NRC-S
that confer resistance in combination with their corresponding NRC-H, could increase the
likelihood of successfully transferring disease resistance across distantly related crops.

Materials and methods

Prediction and phylogenetic analyses of NLRs

The annotated protein sequences of selected plant species used in this study were downloaded from
public databases (Supplemental Data Set 1). NLR prediction and phylogenetic classification were
performed according to previously established pipelines (Jupe et al. 2012; Andolfo et al. 2014).
To identify the proteins containing NLR features, sequences were scanned with MAST in MEME
v 5.4.0 using 20 previously defined motifs with default parameters (Bailey et al. 2009; Jupe et al.
2012). The extracted sequences were then aligned using MAFFT version 7 with the G-INS-1
progressive method and default settings (Katoh et al. 2019). The aligned amino acid sequences
were imported into MEGA 7 for manual trimming leaving only the NB-ARC domain. The NB-
ARC domains with at least three of the four major motifs (P-loop, GLPL, Kinase2 and MHD) were
considered intact. Sequences with truncated NB-ARC domains were excluded from further
analyses. The remaining alignments are usually around 250 amino acids in length depending on
the sequences included. Phylogenetic trees were constructed in MEGA 7 using Maximum-

likelihood phylogenetic analyses with the evolutionary model JTT+G+I and 500 bootstrap tests
(Kumar et al. 2016). The number of bootstrap tests required were conducted using a convergence
test in RAxML (Stamatakis 2014). Phylogenetic trees were further processed and visualized using
FigTree and iTOL (Rambaut 2018; Letunic and Bork 2021). NLR sequences from tomato were
included as references for the identification of the NRC superclade. The species tree of
angiosperms in Fig. 1 was modified from the published phylogenetic tree (The Angiosperm
Phylogeny Group 2016). All datasets used for phylogenetic analyses are in Supplemental Files 1.

Detection of positive selection

Full-length nucleotide sequences of the NRC family were analysed using MEGA 7. The nucleotide
sequences were translated into protein sequences and subsequently aligned using ClustalW. Gaps
in the alignment were manually removed. The resulting trimmed amino acid sequences of the NRC
family were subjected to phylogenetic analyses. Phylogenetic trees were constructed using
Maximum-likelihood phylogenetic analyses with the evolutionary model JTT+G+I and 200
bootstrap tests. We then generated codon-based alignment of the NRC family using the amino acid
sequences alignment mentioned above. Along with the phylogenetic tree, codon-based alignment
of the NRC family was subjected to aBSREL (An adaptive branch-site REL test for episodic
diversification) using the HyPhy (Hypothesis testing using Phylogenies) package on the
Datamonkey Adaptive Evolution Server (https://www.datamonkey.org/) (Smith et al. 2015;
Weaver et al. 2018; Kosakovsky Pond et al. 2020) SlNRC0-dependent sensor NLR Sl8230
(Solyc10g008230) was included as an outgroup.

RNA and DNA isolation

Materials of carrot (*Daucus carota*), kiwifruits (*Actinidia chinensis*), tea (*Camelia sinensis*),
lettuce (*Lactuca sativa*), and water spinach (*Ipomoea aquatica*) were obtained from local nurseries
or grocery stores. Littlebell (*Ipomoea triloba*) was collected from the campus of Academia Sinica
(Nankang, Taipei, Taiwan) and confirmed by Sanger sequencing using ITK and MatK primers.
DNA was extracted from young seedlings or mature leaves using DNeasy Plant Mini Kit (Qiagen).
RNA was extracted using Plant Total RNA Mini Kits (VIOGENE). Synthesis of cDNA was
performed using SuperScript™ III Reverse Transcriptase (Invitrogen) following the
manufacturer's instructions.

Cloning of NLR genes

Full-length NLRs were amplified from genomic DNA or cDNA with primers designed based on
the available genome sequences. The PCR amplicons were cloned into the pAGM9121 (Addgene
plasmid #51833), pICH41308 (Addgene plasmid #47998) using Golden Gate Cloning (Weber et
al. 2011) or pTA in T&A™ Cloning Kit from Yeastern Biotech (Taiwan). The putative sensor
NLRs of tea (Cs0021741 and Cs0024074), and *I. aquatica* NRC6 were synthesized in pUC57-Kan
as MoClo level 0 modules using the service provided by SynBio Technologies (New Jersey, USA).

Mutations (D to V) of the MHD motif were carried out using inverse PCR with primers containing
AarI (NEB) or BsmBI (NEB) enzyme sites for digestion followed by ligation using T4 DNA ligase
(Invitrogen). The constructs were then subcloned into binary vector pICSL86922OD using BsaI
(NEB) and T4 DNA ligase (Invitrogen). pICSL86922OD was kindly provided by Mark Youles
(The Sainsbury Laboratory, UK) (Addgene plasmid # 86181). The NLR gene expression
constructs were then transformed into *Agrobacterium tumefaciens* (GV3101) using electroporation.
The NLR gene sequences and cloning primers used in this research are listed in Supplemental Data
Set 3 and Supplemental Data Set 5. Several previously described NRC-dependent solanaceous
NLRs, including R1, Rx, Rpi-amr1, Rpi-blb2, Bs2, and Prf (Pto), and the corresponding effectors,
including AVR1, CP, AVRamr1, AVRblb2, AvrBs2, and AvrPto were used in the cell death assays
together with Con-IaNRCs (Wu et al. 2017; Witek et al. 2021).

Agroinfiltration and quantification of cell death in *N. benthamiana* leaves

Transient expression of NLRs were performed on 4-week-old *N. benthamiana* leaves (WT or
*nrc2/3/4*_KO) with *A. tumefaciens* (GV3101) carrying the indicated expression constructs (Wu et
al. 2017; Witek et al. 2021). *A. tumefaciens* suspensions in infiltration buffer (10 mM MES, 10
mM MgCl2, and 150 µM acetosyringone, pH 5.6) were adjusted to suitable OD_{600} and then
infiltrated into *N. benthamiana* leaves using needleless syringes. The agro-infiltrated plants were
kept in a walk-in growth chamber (temperature 24-26°C, humidity 45–65% and 16/8 hr light/dark
cycle) for 5 days before imaging and autofluorescence-based cell death quantification using UVP
ChemStudio PLUS (Analytik Jena). Raw cell death autofluorescence images were acquired using
blue LED light for excitation and the FITC filter (513 - 557 nm) as the emission filter. Areas
showing stronger visual cell death generally produce stronger autofluorescence at this wavelength
range. The exposure time was adjusted to 10 seconds to avoid saturation of the autofluorescence
signal. The mean signal intensity was calculated using VisionWorks v.11.2 software by manually
selecting the infiltrated areas and subtracting the background signal intensity. The mean signal
intensity value was further normalized with the maximum intensity (65535) to obtain the relative
intensity of cell death. Results of cell death assays were presented as relative cell death intensity
from at least 6 technical replicates. In some cases, co-expression of NRC-H and NRC-S induces
weak and variable cell death. Quantification using autofluorescence-based analyses may not yield
statistically significant results due to the variation observed for the weak phenotypes. We classified
these as weak cell death to distinguish them from those that completely lacked cell death (e.g.
where no visible cell death was observed).

Statistical analysis of cell death assays

706   Statistical analysis was performed using R (v4.3.3). Nonparametric Wilcoxon signed rank tests
707   were conducted to compare the negative controls (EV) and the tested groups using the rstatix
708   package in RStudio (Kassambara 2023). Statistical significance was defined as $p < 0.05$ and

709 represented in figures as follows: *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$. Detailed statistical data
710 are provided in Supplementary Data Set 6.

Acknowledgements

We thank Dr. Sophien Kamoun (The Sainsbury Laboratory, UK), Dr. Hiroaki Adachi (Laboratory
of Crop Evolution, Graduate School of Agriculture, Kyoto University) and Toshiyuki Sakai
(Laboratory of Crop Evolution, Graduate School of Agriculture, Kyoto University) for valuable
suggestions on the research and comments on the manuscript. We thank Dr. Tsai-Ming Lu
(Institute of Cellular and Organismic Biology, Academia Sinica) for helping with phylogenetic
analyses. We thank Mark Youles (SynBio, The Sainsbury Laboratory, UK) for sharing plasmids
for molecular cloning. We also thank Dr. Chih-Horng Kuo, Dr. Lay-Sun Ma, and Dr. Ting-Ying
Wu (Institute of Plant and Microbial Biology, Academia Sinica) for their suggestions on the
experimental design and data presentation. This research is funded by the Institute of Plant and
Microbial Biology, Academia Sinica. C.H.W. is funded by the 2030 Cross-Generation Young
Scholars Program of the National Science and Technology Council, Taiwan (NSTC 112-2628-B-
001-007). L.D. is funded by a National Institute of Agricultural Botany (NIAB) Fellowship.
Research in the LD lab is supported by the British Society of Plant Pathology, the Gatsby
Charitable Foundation and the Royal Society.

Author Contributions

727 F.J.G. and C.H.W. designed the research. F.J.G. and C.Y.H. conducted the experiments. F.J.G.
728 analysed the data. F.J.G., L.D., and C.H.W. wrote the manuscript.
729

References

Adachi H, Contreras MP, Harant A, Wu C-H, Derevnina L, Sakai T, Duggan C, Moratto E, Bozkurt TO, Maqbool A, et al. An N-terminal motif in NLR immune receptors is functionally conserved across distantly related plant species. eLife. 2019a:**8**:e49956. https://doi.org/10.7554/eLife.49956

Adachi H, Derevnina L, and Kamoun S. NLR singletons, pairs, and networks: evolution, assembly, and regulation of the intracellular immunoreceptor circuitry of plants. Curr Opin Plant Biol. 2019b:**50**:121–131. https://doi.org/10.1016/j.pbi.2019.04.007

Adachi H, Sakai T, Harant A, Pai H, Honda K, Toghani A, Claeys J, Duggan C, Bozkurt TO, Wu C-H, et al. An atypical NLR protein modulates the NRC immune receptor network in Nicotiana benthamiana. PLoS Genet. 2023a:**19**(1):e1010500. https://doi.org/10.1371/journal.pgen.1010500

Adachi H, Sakai T, Kourelis J, Pai H, Gonzalez Hernandez JL, Utsumi Y, Seki M, Maqbool A, and Kamoun S. Jurassic NLR: conserved and dynamic evolutionary features of the atypically ancient immune receptor ZAR1. Plant Cell. 2023b:koad175. https://doi.org/10.1093/plcell/koad175

Ahn H, Lin X, Olave-Achury AC, Derevnina L, Contreras MP, Kourelis J, Wu C, Kamoun S, and Jones JDG. Effector-dependent activation and oligomerization of plant NRC class helper NLRs by sensor NLR immune receptors Rpi-amr3 and Rpi-amr1. EMBO J. 2023:**42**(5):e111484. https://doi.org/10.15252/embj.2022111484

Andolfo G, Jupe F, Witek K, Etherington GJ, Ercolano MR, Jones JDG. Defining the full tomato NB-LRR resistance gene repertoire using genomic and cDNA RenSeq. BMC Plant Biol. 2014:14:120. doi: 10.1186/1471-2229-14-120.

Ashikawa I, Hayashi N, Yamane H, Kanamori H, Wu J, Matsumoto T, Ono K, and Yano M. Two adjacent nucleotide-binding site-leucine-rich repeat class genes are required to confer Pikm-specific rice blast resistance. Genetics. 2008:**180**(4):2267–2276. https://doi.org/10.1534/genetics.108.095034

Baggs EL, Monroe JG, Thanki AS, O'Grady R, Schudoma C, Haerty W, and Krasileva KV. Convergent loss of an EDS1/PAD4 signaling pathway in several plant lineages reveals coevolved components of plant immunity and drought response. Plant Cell. 2020:**32**(7):2158–2177. https://doi.org/10.1105/tpc.19.00903

Bailey TL, Boden M, Buske FA, Frith M, Grant CE, Clementi L, Ren J, Li WW, and Noble WS. MEME SUITE: tools for motif discovery and searching. Nucleic Acids Res. 2009:**37**:W202-208. https://doi.org/10.1093/nar/gkp335

Barragan AC and Weigel D. Plant NLR diversity: the known unknowns of pan-NLRomes. Plant Cell. 2021:**33**(4):814–831. https://doi.org/10.1093/plcell/koaa002

Baumgarten A, Cannon S, Spangler R, and May G. Genome-level evolution of resistance genes in *Arabidopsis thaliana*. Genetics. 2003:**165**(1):309–319. https://doi.org/10.1093/genetics/165.1.309

Bi G, Su M, Li N, Liang Y, Dang S, Xu J, Hu M, Wang J, Zou M, Deng Y, et al. The ZAR1 resistosome is a calcium-permeable channel triggering plant immune signaling. Cell. 2021:**184**(13):3528-3541 e12. https://doi.org/10.1016/j.cell.2021.05.003

Castel B, Ngou P-M, Cevik V, Redkar A, Kim D-S, Yang Y, Ding P, and Jones JDG. Diverse NLR immune receptors activate defence via the RPW8-NLR NRG1. New Phytol. 2019:**222**(2):966–980. https://doi.org/10.1111/nph.15659

Césari S, Kanzaki H, Fujiwara T, Bernoux M, Chalvon V, Kawano Y, Shimamoto K, Dodds P, Terauchi R, and Kroj T. The NB-LRR proteins RGA4 and RGA5 interact functionally and physically to confer disease resistance. EMBO J. 2014:**33**(17):1941–1959. https://doi.org/10.15252/embj.201487923

Cesari S, Thilliez G, Ribot C, Chalvon V, Michel C, Jauneau A, Rivas S, Alaux L, Kanzaki H, Okuyama Y, et al. The rice resistance protein pair RGA4/RGA5 recognizes the Magnaporthe oryzae effectors AVR-Pia and AVR1-CO39 by direct binding. Plant Cell. 2013:**25**(4):1463–1481. https://doi.org/10.1105/tpc.112.107201

Chen Z, Wu Q, Tong C, Chen H, Miao D, Qian X, Zhao X, Jiang L, and Tao X. Characterization of the roles of SGT1/RAR1, EDS1/NDR1, NPR1, and NRC/ADR1/NRG1 in Sw-5b-mediated resistance to *Tomato Spotted Wilt Virus*. Viruses. 2021:**13**(8):1447. https://doi.org/10.3390/v13081447

Chia K-S, Kourelis J, Vickers M, Sakai T, Kamoun S, and Carella P. The N-terminal executioner domains of NLR immune receptors are functionally conserved across major plant lineages. bioRxiv 2022: 2022.10.19.512840. https://doi.org/10.1101/2022.10.19.512840

Collier SM, Hamel L-P, and Moffett P. Cell death mediated by the N-terminal domains of a unique and highly conserved class of NB-LRR protein. Mol Plant-Microbe Interact. 2011:**24**(8):918–931. https://doi.org/10.1094/MPMI-03-11-0050

Contreras MP, Lüdke D, Pai H, Toghani A, and Kamoun S. NLR receptors in plant immunity: making sense of the alphabet soup. EMBO Rep. 2023a:**24**(10):e57495. https://doi.org/10.15252/embr.202357495

Contreras MP, Pai H, Tumtas Y, Duggan C, Yuen ELH, Cruces AV, Kourelis J, Ahn H-K, Lee K-T, Wu C-H, et al. Sensor NLR immune proteins activate oligomerization of their NRC helpers in response to plant pathogens. EMBO J. 2023b:**42**(5):e111519. https://doi.org/10.15252/embj.2022111519

Derevnina L, Contreras MP, Adachi H, Upson JL, Cruces AV, Xie R, Sklenar J, Menke FLH, Mugford ST, MacLean D, Ma W, Hogenhout S, Goverse A, Maqbool A, Wu CH, and Kamoun S. Plant pathogens convergently evolved to counteract redundant nodes of an

NLR immune receptor network. PLOS Biology. 2021:19(8): e3001136.
https://doi.org/10.1371/journal.pbio.3001136

Dodds PN and Rathjen JP. Plant immunity: towards an integrated view of plant-pathogen
interactions. Nat Rev Genet. 2010:**11**(8):539–48. https://doi.org/10.1038/nrg2812

Duggan C, Moratto E, Savage Z, Hamilton E, Adachi H, Wu CH, Leary AY, Tumtas Y,
Rothery SM, Maqbool A, et al. Dynamic localization of a helper NLR at the plant-
pathogen interface underpins pathogen recognition. Proc Natl Acad Sci U A. 2021:**118**(34).
https://doi.org/10.1073/pnas.2104997118

Duxbury Z, Wu CH, and Ding P. A Comparative Overview of the Intracellular Guardians of
Plants and Animals: NLRs in innate immunity and beyond. Annu Rev Plant Biol.
2021:**72**:155–184. https://doi.org/10.1146/annurev-arplant-080620-104948

Ence D, Smith KE, Fan S, Gomide Neves L, Paul R, Wegrzyn J, Peter GF, Kirst M, Brawner
J, Nelson CD, et al. NLR diversity and candidate fusiform rust resistance genes in loblolly
pine. G3. 2022:**12**(2):jkab421. https://doi.org/10.1093/g3journal/jkab421

Förderer A, Li E, Lawson AW, Deng Y-N, Sun Y, Logemann E, Zhang X, Wen J, Han Z,
Chang J, et al. A wheat resistosome defines common principles of immune receptor
channels. Nature. 2022:**610**(7932):532–539. https://doi.org/10.1038/s41586-022-05231-w

Gao Y, Wang W, Zhang T, Gong Z, Zhao H, and Han G-Z. Out of Water: the origin and early
diversification of plant *R* -genes. Plant Physiol. 2018:**177**(1):82–89.
https://doi.org/10.1104/pp.18.00185

Han X and Tsuda K. Evolutionary footprint of plant immunity. Curr Opin Plant Biol.
2022:**67**:102209. https://doi.org/10.1016/j.pbi.2022.102209

Hao Y, Pan Y, Chen W, Rashid MAR, Li M, Che N, Duan X, and Zhao Y. Contribution of
duplicated nucleotide-binding leucine-rich repeat (NLR) genes to wheat disease resistance.
Plants. 2023:**12**(15):2794. https://doi.org/10.3390/plants12152794

Jacob F, Vernaldi S, and Maekawa T. Evolution and conservation of plant NLR functions. Front
Immunol. 2013:**4**. https://doi.org/10.3389/fimmu.2013.00297

Jones JD, Vance RE, and Dangl JL. Intracellular innate immune surveillance devices in plants
and animals. Science. 2016:**354**(6316). https://doi.org/10.1126/science.aaf6395

Jupe F, Pritchard L, Etherington GJ, Mackenzie K, Cock PJA, Wright F, Sharma SK, Bolser
D, Bryan GJ, Jones JDG, et al. Identification and localisation of the NB-LRR gene family
within the potato genome. BMC Genomics. 2012:**13**:75. https://doi.org/10.1186/1471-
2164-13-75

Kassambara A. rstatix: pipe-friendly framework for basic statistical tests. 2023.

Katoh K, Rozewicki J, and Yamada KD. MAFFT online service: multiple sequence alignment,
interactive sequence choice and visualization. Brief Bioinform. 2019:**20**(4):1160–1166.
https://doi.org/10.1093/bib/bbx108

Kim S, Park J, Yeom S-I, Kim Y-M, Seo E, Kim K-T, Kim M-S, Lee JM, Cheong K, Shin H-
S, et al. New reference genome sequences of hot pepper reveal the massive evolution of
plant disease-resistance genes by retroduplication. Genome Biol. 2017:**18**(1):210.
https://doi.org/10.1186/s13059-017-1341-9

Kosakovsky Pond SL, Poon AFY, Velazquez R, Weaver S, Hepler NL, Murrell B, Shank SD,
Magalis BR, Bouvier D, Nekrutenko A, et al. HyPhy 2.5—A customizable platform for
evolutionary hypothesis testing using phylogenies. Mol Biol Evol. 2020:**37**(1):295–299.
https://doi.org/10.1093/molbev/msz197

Kourelis J, Contreras MP, Harant A, Pai H, Lüdke D, Adachi H, Derevnina L, Wu C-H, and
Kamoun S. The helper NLR immune protein NRC3 mediates the hypersensitive cell death
caused by the cell-surface receptor Cf-4. PLOS Genet. 2022:**18**(9):e1010414.
https://doi.org/10.1371/journal.pgen.1010414

Kourelis J, Sakai T, Adachi H, and Kamoun S. RefPlantNLR is a comprehensive collection of
experimentally validated plant disease resistance proteins from the NLR family. PLOS Biol.
2021:**19**(10):e3001124. https://doi.org/10.1371/journal.pbio.3001124

Kuang H, Woo S-S, Meyers BC, Nevo E, and Michelmore RW. Multiple genetic processes
result in heterogeneous rates of evolution within the major cluster disease resistance genes
in lettuce. Plant Cell. 2004:**16**(11):2870–2894. https://doi.org/10.1105/tpc.104.025502

Kumar S, Stecher G, and Tamura K. MEGA7: Molecular Evolutionary Genetics Analysis
Version 7.0 for bigger datasets. Mol Biol Evol. 2016:**33**(7):1870–1874.
https://doi.org/10.1093/molbev/msw054

Lee H, Mang H, Choi E, Seo Y, Kim M, Oh S, Kim S, and Choi D. Genome-wide functional
analysis of hot pepper immune receptors reveals an autonomous NLR clade in seed plants.
New Phytol. 2021:**229**(1):532–547. https://doi.org/10.1111/nph.16878

Leister D. Tandem and segmental gene duplication and recombination in the evolution of plant
disease resistance gene. Trends Genet. 2004:**20**(3):116–122.
https://doi.org/10.1016/j.tig.2004.01.007

Letunic I and Bork P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree
display and annotation. Nucleic Acids Res. 2021:**49**(W1):W293–W296.
https://doi.org/10.1093/nar/gkab301

Liang X and Dong J. Comparative-genomic analysis reveals dynamic NLR gene loss and gain
across Apiaceae species. Front Genet. 2023:**14**:1141194.
https://doi.org/10.3389/fgene.2023.1141194

Lin X, Olave-Achury A, Heal R, Pais M, Witek K, Ahn H-K, Zhao H, Bhanvadia S, Karki HS, Song T, et al. A potato late blight resistance gene protects against multiple Phytophthora species by recognizing a broadly conserved RXLR-WY effector. Mol Plant. 2022:**15**(9):1457–1469. https://doi.org/10.1016/j.molp.2022.07.012

Liu Y, Zeng Z, Zhang Y-M, Li Q, Jiang X-M, Jiang Z, Tang J-H, Chen D, Wang Q, Chen J-Q, et al. An angiosperm NLR Atlas reveals that NLR gene reduction is associated with ecological specialization and signal transduction component deletion. Mol Plant. 2021:**14**(12):2015–2031. https://doi.org/10.1016/j.molp.2021.08.001

Maqbool A, Saitoh H, Franceschetti M, Stevenson C, Uemura A, Kanzaki H, Kamoun S, Terauchi R, and Banfield M. Structural basis of pathogen recognition by an integrated HMA domain in a plant NLR immune receptor. eLife. 2015:**4**:e08709. https://doi.org/10.7554/eLife.08709

Marchal C, Michalopoulou VA, Zou Z, Cevik V, and Sarris PF. Show me your ID: NLR immune receptors with integrated domains in plants. Essays Biochem. 2022:**66**(5):527–539. https://doi.org/10.1042/EBC20210084

Martin EC, Ion CF, Ifrimescu F, Spiridon L, Bakker J, Goverse A, and Petrescu A-J. NLRscape: an atlas of plant NLR proteins. Nucleic Acids Res. 2023:**51**(D1):D1470–D1482. https://doi.org/10.1093/nar/gkac1014

Meyers BC, Kaushik S, and Nandety RS. Evolving disease resistance genes. Curr Opin Plant Biol. 2005:**8**(2):129–134. https://doi.org/10.1016/j.pbi.2005.01.002

Michelmore RW and Meyers BC. Clusters of resistance genes in plants evolve by divergent selection and a birth-and-death process. Genome Res. 1998:**8**(11):1113–1130. https://doi.org/10.1101/gr.8.11.1113

Narusaka M, Kubo Y, Hatakeyama K, Imamura J, Ezura H, Nanasato Y, Tabei Y, Takano Y, Shirasu K, and Narusaka Y. Breaking restricted taxonomic functionality by dual resistance genes. Plant Signal Behav. 2013:**8**(6):e24244. https://doi.org/10.4161/psb.24244

Narusaka M, Shirasu K, Noutoshi Y, Kubo Y, Shiraishi T, Iwabuchi M, and Narusaka Y. RRS1 and RPS4 provide a dual Resistance-gene system against fungal and bacterial pathogens. Plant J. 2009:**60**(2):218–226. https://doi.org/10.1111/j.1365-313X.2009.03949.x

Ngou BPM, Ding P, and Jones JDG. Thirty years of resistance: Zig-zag through the plant immune system. Plant Cell. 2022:**34**(5):1447–1478. https://doi.org/10.1093/plcell/koac041

Oh S, Kim S, Park H, Kim M, Seo M, Wu C, Lee H, Kim H, Kamoun S, and Choi D. Nucleotide-binding leucine-rich repeat network underlies nonhost resistance of pepper against the Irish potato famine pathogen *Phytophthora infestans*. Plant Biotechnol J. 2023:**21**(7):1361–1372. https://doi.org/10.1111/pbi.14039

Ortiz D and Dodds PN. Plant NLR Origins Traced Back to Green Algae. Trends Plant Sci.
2018:**23**(8):651–654. https://doi.org/10.1016/j.tplants.2018.05.009

Rambaut A. Figtree v1.4.4. 2018. http://tree.bio.ed.ac.uk/software/figtree/

Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
phylogenies. Bioinformatics. 2014:30(9):1312-1213.

Saile SC, Jacob P, Castel B, Jubic LM, Salas-Gonzáles I, Bäcker M, Jones JDG, Dangl JL,
and El Kasmi F. Two unequally redundant "helper" immune receptor families mediate
Arabidopsis thaliana intracellular "sensor" immune receptor functions. PLOS Biol.
2020:**18**(9):e3000783. https://doi.org/10.1371/journal.pbio.3000783

Sakai T, Martinez-Anaya C, Contreras MP, Kamoun S, Wu C-H, and Adachi H. The NRC0
gene cluster of sensor and helper NLR immune receptors is functionally conserved across
asterid   plants.   bioRxiv.   2023:2023.10.23.563533.
https://doi.org/10.1101/2023.10.23.563533

Sarris PF, Duxbury Z, Huh SU, Ma Y, Segonzac C, Sklenar J, Derbyshire P, Cevik V,
Rallapalli G, Saucet SB, et al. A plant immune receptor detects pathogen effectors that
target   WRKY   transcription   factors.   Cell.   2015:**161**(5):1089–1100.
https://doi.org/10.1016/j.cell.2015.04.024

Seidl MF and Thomma BPHJ. Transposable elements direct the coevolution between plants and
microbes. Trends Genet. 2017:**33**(11):842–851. https://doi.org/10.1016/j.tig.2017.07.003

Seo E, Kim S, Yeom S-I, and Choi D. Genome-wide comparative analyses reveal the dynamic
evolution of nucleotide-binding leucine-rich repeat gene family among Solanaceae plants.
Front Plant Sci. 2016:**7**:1205. https://doi.org/10.3389/fpls.2016.01205

Seong K, Seo E, Witek K, Li M, and Staskawicz B. Evolution of NLR resistance genes with
noncanonical N-terminal domains in wild tomato species. New Phytol. 2020:**227**(5):1530–
1543. https://doi.org/10.1111/nph.16628

Shao Z-Q, Xue J-Y, Wang Q, Wang B, and Chen J-Q. Revisiting the origin of plant NBS-LRR
genes. Trends Plant Sci. 2019:**24**(1):9–12. https://doi.org/10.1016/j.tplants.2018.10.015

Shao Z-Q, Xue J-Y, Wu P, Zhang Y-M, Wu Y, Hang Y-Y, Wang B, and Chen J-Q. Large-
scale analyses of angiosperm nucleotide-binding site-leucine-rich repeat genes reveal three
anciently diverged classes with distinct evolutionary patterns. Plant Physiol.
2016:**170**(4):2095–2109. https://doi.org/10.1104/pp.15.01487

Shimizu M, Hirabuchi A, Sugihara Y, Abe A, Takeda T, Kobayashi M, Hiraka Y, Kanzaki
E, Oikawa K, Saitoh H, et al. A genetically linked pair of NLR immune receptors shows
contrasting patterns of evolution. Proc Natl Acad Sci U S A. 2022:**119**(27):e2116896119.
https://doi.org/10.1073/pnas.2116896119

Smith MD, Wertheim JO, Weaver S, Murrell B, Scheffler K, and Kosakovsky Pond SL. Less
Is More: An adaptive branch-site random effects model for efficient detection of episodic
diversifying   selection.   Mol   Biol   Evol.   2015:**32**(5):1342–1353.
https://doi.org/10.1093/molbev/msv022

Sohn KH, Segonzac C, Rallapalli G, Sarris PF, Woo JY, Williams SJ, Newman TE, Paek KH,
Kobe B, and Jones JDG. The nuclear immune receptor RPS4 is required for RRS1SLH1-
dependent constitutive defense activation in *Arabidopsis thaliana*. PLoS Genet.
2014:**10**(10):e1004655. https://doi.org/10.1371/journal.pgen.1004655

Stam R, Silva-Arias GA, and Tellier A. Subsets of NLR genes show differential signatures of
adaptation during colonization of new habitats. New Phytol. 2019:**224**(1):367–379.
https://doi.org/10.1111/nph.16017

Tai TH, Dahlbeck D, Clark ET, Gajiwala P, Pasion R, Whalen MC, Stall RE, and Staskawicz
BJ. Expression of the Bs2 pepper gene confers resistance to bacterial spot disease in tomato.
Proc   Natl   Acad   Sci   U   S   A.   1999:**96**(24):14153–14158.
https://doi.org/10.1073/pnas.96.24.14153

Tarr DEK and Alexander HM. TIR-NBS-LRR genes are rare in monocots: evidence from
diverse monocot orders. BMC Res Notes. 2009:**2**(1):197. https://doi.org/10.1186/1756-
0500-2-197

Terefe-Ayana D, Kaufmann H, Linde M, and Debener T. Evolution of the Rdr1 TNL-cluster
in   roses   and   other   Rosaceous   species.   BMC   Genomics.   2012:**13**(1):409.
https://doi.org/10.1186/1471-2164-13-409

The Angiosperm Phylogeny Group. An update of the Angiosperm Phylogeny Group
classification for the orders and families of flowering plants: APG IV. Bot J Linn Soc.
2016:**181**(1):1–20. https://doi.org/10.1111/boj.12385

Thind AK, Wicker T, Müller T, Ackermann PM, Steuernagel B, Wulff BBH, Spannagl M,
Twardziok SO, Felder M, Lux T, et al. Chromosome-scale comparative sequence
analysis unravels molecular mechanisms of genome dynamics between two wheat cultivars.
Genome Biol. 2018:**19**(1):104. https://doi.org/10.1186/s13059-018-1477-2

Upson JL, Zess EK, Białas A, Wu C, and Kamoun S. The coming of age of EvoMPMI:
evolutionary molecular plant–microbe interactions across multiple timescales. Curr Opin
Plant Biol. 2018:**44**:108–116. https://doi.org/10.1016/j.pbi.2018.03.003

Van De Weyer A-L, Monteiro F, Furzer OJ, Nishimura MT, Cevik V, Witek K, Jones JDG,
Dangl JL, Weigel D, and Bemm F. A species-wide inventory of NLR genes and alleles
in   *Arabidopsis   thaliana*.   Cell.   2019:**178**(5):1260-1272.e14.
https://doi.org/10.1016/j.cell.2019.07.038

Van Ghelder C, Parent GJ, Rigault P, Prunier J, Giguère I, Caron S, Stival Sena J,
Deslauriers A, Bousquet J, Esmenjaud D, et al. The large repertoire of conifer NLR

resistance genes includes drought responsive and highly diversified RNLs. Sci Rep. 2019:**9**(1):11614. https://doi.org/10.1038/s41598-019-47950-7

Wang G, Roux B, Feng F, Guy E, Li L, Li N, Zhang X, Lautier M, Jardinaud M-F, Chabannes M, et al. The decoy substrate of a pathogen effector and a pseudokinase specify pathogen-induced modified-self recognition and immunity in plants. Cell Host Microbe. 2015:**18**(3):285–295. https://doi.org/10.1016/j.chom.2015.08.004

Wang J, Hu M, Wang J, Qi J, Han Z, Wang G, Qi Y, Wang H-W, Zhou J-M, and Chai J. Reconstitution and structure of a plant NLR resistosome conferring immunity. Science. 2019:**364**(6435):eaav5870. https://doi.org/10.1126/science.aav5870

Weaver S, Shank SD, Spielman SJ, Li M, Muse SV, and Kosakovsky Pond SL. Datamonkey 2.0: A modern web application for characterizing selective and other evolutionary processes. Mol Biol Evol. 2018:**35**(3):773–777. https://doi.org/10.1093/molbev/msx335

Weber E, Engler C, Gruetzner R, Werner S, and Marillonnet S. A modular cloning system for standardized assembly of multigene constructs. PloS One. 2011:**6**(2):e16765. https://doi.org/10.1371/journal.pone.0016765

Wei F, Wing RA, and Wise RP. Genome dynamics and evolution of the Mla (Powdery Mildew) resistance locus in barley. Plant Cell. 2002:**14**(8):1903–1917. https://doi.org/10.1105/tpc.002238

Witek K, Lin X, Karki HS, Jupe F, Witek AI, Steuernagel B, Stam R, van Oosterhout C, Fairhead S, Heal R, et al. A complex resistance locus in *Solanum americanum* recognizes a conserved *Phytophthora* effector. Nat Plants. 2021:**7**(2):198–208. https://doi.org/10.1038/s41477-021-00854-9

Woudstra Y, Tumas H, Ghelder C van, Hung TH, Ilska JJ, Girardi S, A'Hara S, McLean P, Cottrell J, Bohlmann J, et al. Conifers concentrate large numbers of NLR immune receptor genes on one chromosome. bioRxiv. 2023:2023.10.20.563305. https://doi.org/10.1101/2023.10.20.563305

Wu C-H, Abd-El-Haliem A, Bozkurt TO, Belhaj K, Terauchi R, Vossen JH, and Kamoun S. NLR network mediates immunity to diverse plant pathogens. Proc Natl Acad Sci U S A. 2017:**114**(30):8113–8118. https://doi.org/10.1073/pnas.1702041114

Wu CH, Belhaj K, Bozkurt TO, Birk MS, and Kamoun S. Helper NLR proteins NRC2a/b and NRC3 but not NRC1 are required for Pto-mediated cell death and resistance in *Nicotiana benthamiana*. New Phytol. 2016:**209**(4):1344–52. https://doi.org/10.1111/nph.13764

Wu C-H, Derevnina L, and Kamoun S. Receptor networks underpin plant immunity. Science. 2018:**360**(6395):1300–1301. https://doi.org/10.1126/science.aat2623

Wu CH and Kamoun S. Tomato *Prf* requires NLR helpers *NRC2* and *NRC3* to confer resistance against the bacterial speck pathogen *Pseudomonas syringae* pv. *tomato*. Acta Hortic. 2021:(1316):61–66. https://doi.org/10.17660/ActaHortic.2021.1316.9

Wu J-Y, Xue J-Y, and Van de Peer Y. Evolution of NLR resistance genes in Magnoliids:
Dramatic expansions of CNLs and multiple losses of TNLs. Front Plant Sci.
2021:**12**:777157. https://doi.org/10.3389/fpls.2021.777157

Xi Y, Cesari S, and Kroj T. Insight into the structure and molecular mode of action of plant
paired NLR immune receptors. Essays Biochem. 2022:**66**(5):513–526.
https://doi.org/10.1042/EBC20210079

Zhang N, Gan J, Carneal L, González-Tobón J, Filiatrault M, and Martin GB. Helper NLRs
Nrc2 and Nrc3 act codependently with Prf/Pto and activate MAPK signaling to induce
immunity in tomato. Plant J. 2024:**117**(1):7–22. https://doi.org/10.1111/tpj.16502

Zhang Y-M, Shao Z-Q, Wang Q, Hang Y-Y, Xue J-Y, Wang B, and Chen J-Q. Uncovering
the dynamic evolution of nucleotide-binding site-leucine-rich repeat (NBS-LRR) genes in
Brassicaceae: Evolution of NBS-LRR genes in Brassicaceae. J Integr Plant Biol.
2016:**58**(2):165–177. https://doi.org/10.1111/jipb.12365

Zhou T, Wang Y, Chen J-Q, Araki H, Jing Z, Jiang K, Shen J, and Tian D. Genome-wide
identification of NBS genes in japonica rice reveals significant expansion of divergent non-
TIR NBS-LRR genes. Mol Genet Genomics. 2004:**271**(4):402–415.
https://doi.org/10.1007/s00438-004-0990-z

1038

Figure 1. The expansion of the NRC superclade is a unique feature of lamiids. **A)** (Left) Phylogenetic tree of angiosperm species, modified from The Angiosperm Phylogeny Group (2016). (Right) Percentage and number of different types of NLRs identified from each species. Numbers in the stacked bar chart indicate the sizes of NRC superclade members in the plant species. **B)** Correlation between the size of the NRC superclade and the number of total protein-coding genes. **C)** Comparisons of the NRC sensor-helper ratio and the size of the NRC superclade. The highlighted area indicates the ancestral ratio of the NRC sensor and helper, which is close to 1:1 to 2:1. Plant species from different lineages are labelled with different colours.

Figure 2. Ericales encode a simple NRC network, with sensor NLRs signalling through NRC0-Eri to initiate cell death. **A)** Phylogenetic analysis of total NLRs from 3 Ericales species, including *A. chinensis*, *S. dulcificum*, and *C. sinensis*. NLRs from different species are indicated with lines of different colours. NLR sequences from the tomato NRC superclade were used as references. **B)** Phylogenetic analysis of the NRC superclade from the 3 Ericales species with tomato NRC0 as reference sequences. For both phylogenetic trees, major branches with bootstrap values higher than 70 are indicated with black dots. The scale bar indicates the evolutionary distance in amino acid substitution per site. **C)** Cell death assay was performed by co-expression of the putative NRC0-Eri-dependent sensor NLR from tea (*C. sinensis*) and tea CsNRC0a-Eri. All sensor NLRs carried the MHD motif (D to V) mutation. The intensity of cell death was analysed at 5 dpi. The dot plot represents the relative cell death intensity based on autofluorescence imaging using UVP ChemStudio PLUS. The bold line in the boxplots represents the medium, the box edges represent the 25th and 75th percentiles, and the whiskers extend to the most extreme data points no more than 1.5x of the interquartile range. Statistical differences were examined by paired Wilcoxon signed rank test (**** = $p < 0.0001$; * = $p < 0.05$). **D)** Putative NRC network structure in Ericales where few sensor NLRs signal through Ericales NRC0 (NRC0-Eri) to induce cell death.

Figure 3. Campanulids show an NRC network with two partially redundant NRC nodes. **A)** Phylogenetic analysis of CC$_{Rx}$-CNLs from 7 species of campanulids, including *C. nankingense*, *H. annuus*, *L. sativa*, *S. rebaudiana*, *A. graveolens*, *D. carota*, and *P. ginseng*. **B)** Phylogenetic analysis of the NRC superclade from 7 species of campanulids using tomato NRC0 as references. Major branches with bootstrap values > 70 are indicated with black dots in both (A) and (B). The scale bar indicates the evolutionary distance in amino acid substitution per site. **C)** Cell death assay results of NRC-dependent sensor NLRs co-expressed with the lettuce NRCs in *N. benthamiana* at 5 dpi. All sensor NLRs carried the MHD motif (D to V) mutation. As controls, helper NLR LsNRCs (LsNRC0 or LsNRC0-Ast) were infiltrated without sensor NLRs. The dot plot represents the cell death quantification analysed by UVP ChemStudio PLUS. The bold line in the boxplots represents the medium, the box edges represent the 25th and 75th percentiles, and the whiskers extend to the most extreme data points no more than 1.5x of the interquartile range. Statistical differences were examined by paired Wilcoxon signed rank test (* = p < 0.05, ** = p < 0.01, *** = p < 0.001, **** = p < 0.0001). **D)** Putative NRC network structure in campanulids where some NRC-S signal through NRC0, some NRC-S signal through NRC0-Ast, and some signal through both NRCs to induce HR.

Figure 4. The *Ipomoea* genus possesses a complex NRC network. **A)** Phylogenetic analysis of the NRC superclade of five plant species of lamiids. **B)** Phylogenetic analysis of the NRC family members across five plant species of lamiids. Major branches with bootstrap values > 70 are indicated with black dots in both (A) and (B). The scale bar indicates the evolutionary distance in amino acid substitution per site. **C)** Cell death matrix of *I. aquatica* putative NRC sensor NLRs co-expressed with NRC helper NLRs. All sensor NLRs carried the MHD motif (D to V) mutation. *Sequences cloned form *I. triloba* instead of *I. aquatica* were used in the analysis. Detailed cell death assay results are provided in Supplemental Fig. S7.

Figure 5. Lamiids family-specific NRC subclades, but not the three NRC0 subclades, show diversifying selection. **A)** Phylogenetic tree based on the full-length sequences of the NRC family from 15 selected asterids species. The aBSREL analysis indicated that 15 out of 81 selected internal branches show episodic diversifying selection. Diversifying selection on branch nodes was assessed using the Likelihood Ratio Test with a significance threshold set at $p \leq 0.05$. Detailed phylogeny and results of aBSREL analysis are provided in Supplemental Fig. S9 and Supplemental Data Set 3. The scale bar indicates the evolutionary distance in amino acid substitution per site. **B)** and **C)** Branch length distribution of the three NRC0 subclades and family-specific NRC subclades of lamiids based on the phylogenetic tree in (A). The red lines in both histograms indicate the predicted normal distribution.

Figure 6. NRC0 subclades from different plant lineages showed varying degrees of interchangeability. **A)** Cell death assay results of tea Cs24074 co-expressed with DcNRC0, LsNRC0, LsNRC0-Ast, IaNRC0, and SlNRC0 in *N. benthamiana* observed at 5 dpi. **B)** Cell death assay results of carrot sensor NLRs (Dc23557 and Dc23650) or lettuce sensor NLRs (Ls28800, Ls35940, Ls36021, Ls123301, Ls124100, and Ls124601) co-expressed with CsNRC0-Eri, IaNRC0, and SlNRC0 in *N. benthamiana* observed at 5 dpi. **C)** Cell death assay results of water spinach sensor NLR (Ia15471) or tomato sensor NLR (Sl08230) co-expressed with CsNRC0-Eri, DcNRC0, LsNRC0, LsNRC0-Ast, IaNRC0, and SlNRC0 in *N. benthamiana* observed at 5 dpi. All sensor NLRs carried the MHD motif (D to V) mutation. The dot plot represents cell death quantification analysed by UVP ChemStudio PLUS. The bold line in the boxplots represents the medium, the box edges represent the 25th and 75th percentiles, and the whiskers extend to the most extreme data points no more than 1.5x of the interquartile range. Statistical differences were examined by paired Wilcoxon signed rank test. **D)** Matrix of cell death assays of NRC0-dependent sensor NLRs co-expressed with NRC0 homologs from 5 different species of various lineages.

Figure 7. Sensor and helper NLRs in Solanaceae and Convolvulaceae exhibit a degree of cross-compatibility between these two plant families. **A)** Cell death assay results of solanaceous sensor NLRs (Sl8230, Rx, and Rpi-amr1) co-expressed with water spinach NRC helpers (IaNRC0, Con-IaNRC3, Con-IaNRC5, Con-IaNRC8, Con-IaNRC9) in *N. benthamiana* at 5 dpi. All sensor NLRs carried the MHD motif (D to V) mutation. The dot plot represents cell death quantification analysed by UVP ChemStudio PLUS. The bold line in the boxplots represents the medium, the box edges represent the 25th and 75th percentiles, and the whiskers extend to the most extreme data points no more than 1.5x of the interquartile range. Statistical differences were examined by paired Wilcoxon signed rank test. **B)** Matrix of cell death assays for solanaceous NRC-dependent sensor NLRs co-expressed with water spinach NRC helper NLRs, including information obtained from Supplemental Fig. S11 and S12. **C)** Cell death assay results of water spinach sensor NLRs (Ia15471, Ia04375, Ia14342, and Ia17067) co-expressed with solanaceous NRC helpers in *N. benthamiana* at 5 dpi. All sensor NLRs carried the MHD motif (D to V) mutation. The dot plot represents cell death quantification analysed by UVP ChemStudio PLUS. Statistical differences among the samples were analysed with Tukey's HSD test (p < 0.05) for each sensor NLR independently. **D)** Matrix of cell death assays for solanaceous NRC-dependent sensor NLRs co-expressed with water spinach NRC helper NLRs, including information obtained from Supplemental Fig. S13 and S14.

Figure 8. Evolution of the hierarchical structure of the NRC network in asterids. Ericales encode a simple NRC network where few sensor NLRs signal through NRC0-Eri to induce cell death. Campanulids show duplicated NRC nodes where some NRC-S signal through NRC0, some NRC-S signal through NRC0-Ast, and some signal through both NRCs to induce cell death. In addition to NRC0, lamiids display massively expanded and complex NRC networks that are likely plant family-specific.