Nuclear Pyruvate Dehydrogenase Complex Regulates Histone Acetylation and

Transcriptional Regulation in the Ethylene Response

Zhengyao Shao[1,2], Liangqiao Bian[3], Shyon K. Ahmadi[1], Tyler J. Daniel[1], Miguel A. Belmonte[1],

Jackson G. Burns[1], Prashanth Kotla[1], Yang Bi[4], Zhouxin Shen[5], Shou-Ling Xu[4], Zhi-Yong

Wang[4], Steven P. Briggs[5], and Hong Qiao[1,2*]

[1]Institute for Cellular and Molecular Biology and [2]Department of Molecular Biosciences, The

University of Texas at Austin, Austin, TX, 78712, USA; [3]Shimadzu Center for Advanced Analytical

Chemistry, University of Texas at Arlington, Arlington, TX, 76019, USA; [4]Department of Plant

Biology, Carnegie Institution for Science, Stanford, CA, 94305, USA; [5]Division of Biological

Sciences, University of California San Diego, La Jolla, CA ,92093, USA.

[*]Correspondence: hqiao@austin.utexas.edu

[*] To whom correspondence should be addressed:

Hong Qiao

Department of Molecular Biosciences, University of Texas at Austin

2506 Speedway, NMS 5.324, Austin, TX, 78712

Phone: (512) 471-0286; FAX: (512) 471-1218

Email: hqiao@austin.utexas.edu

Abstract

Ethylene plays its essential roles in plant development, growth, and defense responses by controlling the transcriptional reprograming, in which EIN2-C-directed regulation of histone acetylation is the first key-step for chromatin to perceive ethylene signaling. But how the nuclear acetyl coenzyme A (acetyl CoA) is produced to ensure the ethylene-mediated histone acetylation is unknown. Here we report that ethylene triggers the accumulation of the pyruvate dehydrogenase complex (PDC) in the nucleus to synthesize nuclear acetyl CoA to regulate ethylene response. PDC is identified as an EIN2-C nuclear partner, and ethylene triggers its nuclear accumulation. Mutations in PDC lead to an ethylene-hyposensitivity that results from the reduction of histone acetylation and transcription activation. Enzymatically active nuclear PDC synthesize nuclear acetyl CoA for EIN2-C-directed histone acetylation and transcription regulation. These findings uncover a mechanism by which PDC-EIN2 converges the mitochondrial enzyme mediated nuclear acetyl CoA synthesis with epigenetic and transcriptional regulation for plant hormone response.

Introduction

Ethylene is a pivotal plant hormone that plays pleiotropic roles in plant growth, development, stress response and defense response to pathogens by controlling downstream gene transcription, protein translation and posttranslational modifications, mitochondrial retrograde signaling, chromatin remodeling, and epigenetic regulation of gene expression[1-14]. Ethylene signal is perceived by ethylene receptors embedded on the endoplasmic reticulum (ER) membrane[15,16]. In the absence of ethylene, both the ethylene receptors and CONSTITUTIVE TRIPLE RESPONSE 1 (CTR1), an ER membrane associated kinase that directly interacts with one of ethylene receptors ETHYLENE RESPONSE1 (ETR1), are activated[17,18]. CTR1 phosphorylates ETHYLENE INSENSTIVE 2 (EIN2), the essential positive regulator of ethylene signaling, at its C-terminal end (EIN2-C), leading to the repression of the EIN2 activity[19,20]. Without ethylene

signaling activation, EIN2 is localized to the ER membrane at which it interacts with two F-box

proteins EIN2 TARGETING PROTEIN 1/2 (ETP1/2) that mediate its protein degradation via the

ubiquitin-proteasome pathway[21]. Upon the perception of ethylene, both ETR1 and CTR1 are

inactivated; the EIN2-C is dephosphorylated through an unknown mechanism[20]. The

dephosphorylated EIN2-C is cleaved and translocated into both the nucleus and the P-body[20,22,23].

In the P-body, EIN2-C mediates the translational repression of two F-box proteins EIN3-BINDING

F BOX PROTEIN 1/2 (EBF1/2) to promote the protein accumulation of ETHYLENE INSENSITIVE

3 (EIN3), the key transcription activator that is sufficient and necessary for activation of all

ethylene-response genes[6,23,24]. In the nucleus, EIN2-C mediates the direct regulation of histone

acetylation of H3K14 and H3K23 via histone binding protein EIN2 NUCLEAR ASSOCIATED

PROTEIN 1/2 (ENAP1/2), leading to an EIN3 dependent transcriptional regulation for ethylene

response[25-27].

Epigenetic regulation of gene expression plays critical roles in various plant developmental

processes and stress response[28-32]. Histone acetylation promotes the relaxation of nucleosome

between wrapped DNA and histone octamer by neutralizing the positive charges of lysine

residues in histones, which is crucial for all DNA-based processes, including DNA replication and

gene transcription[33,34]. Histone acetyltransferases (HATs) transfer acetyl group from acetyl

coenzyme A (acetyl CoA) to conserved lysine residues on histone N-terminal tails to acetylate

histones. In eukaryotes, the biosynthesis of acetyl CoA is thought to occur in the subcellular

compartment where it is required because of its membrane impermeability and high instability

due to the high-energy thioester bond that joins the acetyl and CoA groups[35]. The availability and

abundance of acetyl CoA are therefore crucial for HAT enzymatic activity and histone acetylation.

For most HATs, their Michaelis constants (Kms), the concentration of substrate which permits

the enzyme to achieve half Vmax, lie in the range within or greater than the cellular acetyl CoA

concentrations in order to sense and respond to the fluctuation of acetyl CoA production[36].

Therefore, the levels of histone acetylation are often linked with the availability of acetyl CoA.

Such metabolic regulation of histone acetylation by nuclear synthesis of acetyl CoA has been

reported to govern many major biological events including cell proliferation, cell fate determination,

and environmental response[37-43]. However, the relationship between the nuclear acetyl CoA and

histone acetylation in plant hormone response is completely unknown.

In this study, we report that functional pyruvate dehydrogenase complex (PDC) can translocate

from the mitochondria to the nucleus in the presence of ethylene, generating a nuclear pool of

acetyl CoA from pyruvate for the EIN2-directed acetylation of core histones for transcriptional

regulation in the ethylene response. In the presence of ethylene, the PDC complex translocates

to the nucleus where it interacts with EIN2-C. Mutations in PDC lead to a decreased ethylene

response at both at genetic level and transcriptional level. ChIP-seq assay reveals that ethylene-

induced elevation of histone acetylation is reduced in the PDC mutants. Biochemical assay

combined with Liquid Chromatography with mass spectrometry (LC-MS) demonstrates that PDC

regulates the nuclear acetyl CoA concentration and ethylene induced nuclear translocated PDC

is enzymatically active to generate acetyl CoA in the nucleus for histone acetylation, which is

necessary for EIN2-C's function in the histone acetylation regulation. Both genetics and molecular

evidence show that EIN2-C is required for the nuclear translocation of PDC in the ethylene

response. Altogether, this study reveals a previously unidentified mechanism by which the PDC

complex is translocated to the nucleus where it interacts with EIN2-C to provide acetyl CoA for

the elevation of histone acetylation at H3K14 and H3K23 to modulate the transcriptional regulation

in the ethylene response.

Results

Nuclear localized pyruvate dehydrogenase complex interacts with EIN2-C

By tandem co-immunoprecipitation coupled with mass spectrometry (Co-IP/MS) using anti-EIN2-

C native antibody and anti-HA antibody sequentially in the nuclear extracts from *EIN2-YFP-HA*

plants treated with 4 hours of ethylene gas, three subunits from pyruvate dehydrogenase complex

(PDC), E1-2, E2-2, and E3-2 (henceforth referred as E1, E2, and E3), were pulled down among

the top hits (Fig. S1A-1C). The direct interactions between EIN2-C and these three PDC subunits

identified in Co-IP/MS were confirmed by *in vitro* pull-down assays (Fig. 1A and Fig. S1D-F). In

prokaryotes and eukaryotes, PDC consists of three catalytic subunits: a pyruvate dehydrogenase

(E1), a dihydrolipoamide transacetylase (E2), and a dihydrolipoamide dehydrogenase (E3). This

complex is canonically localized in the mitochondrial matrix where it catalyzes the conversion

from pyruvate to acetyl CoA [44]. The interaction between EIN2-C and PDC complex led us to

examine whether it is possible that EIN2 is localized to the mitochondria. Co-localization assay of

EIN2 with MitoTracker Red, a cell-permeant mitochondria-selective dye, showed that EIN2 and

mitochondria were not co-localized either with or without the presence of ethylene (Fig. 1B). Given

that PDC complex was pulled down from the nuclear fraction by EIN2-C Co-IP/MS, we then

decided to confirm whether the interactions between EIN2-C and E1, E2, and E3 occur in the

nucleus *in vivo*. We generated *EIN2-YFP-HA/E1-FLAG-BFP*, *EIN2-YFP-HA/E2-FLAG-BFP,* and

EIN2-YFP-HA/E3-FLAG-BFP transgenic plants. By using EIN2-YFP-HA or PDC-FLAG-BFP as

bait, we performed reciprocal *in vivo* co-immunoprecipitation (Co-IP) assays both in the nuclear

fractions and in the cytosolic fractions from those 3-day-old etiolated transgenic seedlings treated

with or without 4 hours of ethylene gas. The efficiency of nuclear-cytoplasmic fractionation

procedure as well as the purity of cytosolic and nuclear fractions were assessed by multiple

marker proteins (Fig. S1G). The interactions between EIN2-C and all three PDC subunits were

detected only in the nuclear fractions with ethylene treatment, and no interactions were detected

from cytosolic fractions regardless of the ethylene treatment (Fig. 1C-1E). Similar results were

obtained by using the native EIN2-C antibody for the in vivo Co-IP assay in the *E1-YFP-HA*, *E2-*

FLAG-GFP, and *E3-FLAG-GFP* transgenic plants (Fig. S1H-1J). To further evaluate the

interaction between EIN2 and PDC in response to ethylene in the subcellular context, we

conducted bimolecular fluorescence complementation (BiFC) assays by transient co-expression

of EIN2-nYFP and E1-cYFP, E2-cYFP, and E3-cYFP, respectively, in tobacco epidermal cells.

The result showed that EIN2-C physically interacts with PDC subunits only in the nucleus after

ethylene treatment (Fig. 1F-1H). Furthermore, we examined the subcellular localization of EIN2-

C and PDC subunits in response to ethylene in 3-day-old etiolated seedlings of *EIN2-YFP-HA/E1-*

FLAG-BFP, *EIN2-YFP-HA/E2-FLAG-BFP,* and *EIN2-YFP-HA/E3-FLAG-BFP* plants. Clearly,

EIN2-C was co-localized with E1, E2, and E3 in the nucleus respectively with the ethylene

treatment (Fig. S1K-1P). Altogether, these results provide compelling evidence that PDC can

accumulate in the nucleus to interact with EIN2-C in response to ethylene.

Ethylene treatment induces the nuclear accumulation of PDC

We have observed the nuclear localization of PDC complex with the presence of ethylene (Fig.

1). To further confirm that the PDC nuclear translocation is induced by ethylene, we examined

the nuclear appearance of E1, E2, and E3 proteins over a time series of ethylene gas treatments.

E1 had a basal nuclear distribution in the absence of ethylene, but its nuclear levels were

significantly elevated by ethylene treatment and were positively correlated with the duration of

ethylene treatment while E1 cytoplasmic levels showed gradual decrease as the ethylene

treatment prolongs (Fig. 2A, Fig. S2A, and Fig. S2B). Neither E2 nor E3 proteins were detected

in the nucleus in the absence of ethylene, but these two proteins were accumulated in the nuclear

fraction after 4 hours of ethylene treatment and their levels were also positively correlated with

the duration of ethylene treatments (Fig. 2B, Fig. 2C, and Fig. S2C-2F). Similar to cytoplasmic E1,

we also observed that cytoplasmic E2 and E3 protein levels decrease in response to ethylene

(Fig. 2B, Fig. 2C, and Fig. S2C-2F). Notably, the total E1 and E2 subunit protein levels were not

altered by the ethylene treatments but E3 total protein level showed slight elevation after 12h

ethylene treatment. We then monitored the nuclear localization of PDC over a time series of

ethylene treatments in living cells. In the absence of ethylene, a basal level of E1 was observed

in the nucleus, but no nuclear E2 and E3 were observed (Fig. 2D-2F and Fig. S2G-2L). Upon

ethylene treatment, E1, E2, and E3 accumulated in the nucleus, and their accumulation was

positively associated with the duration of ethylene treatment (Fig. 2D-2F and Fig. S2G-2L). This

provides an additional piece of cellular evidence that ethylene induces PDC nuclear accumulation.

PDC subunits are required for ethylene response

Given the interaction between PDC and EIN2-C occurs in the nucleus in response to ethylene,

we decided to investigate the functions of PDC in the ethylene response. We first obtained two T-

DNA insertion lines for *E1* (*e1-2-2* and *e1-2-4)* and two for *E3* (*e3-2-1* and *e3-2-2)*. RT-qPCR

assays showed that the expression levels of these two genes were drastically reduced in their T-

DNA insertion mutants (Fig. S3A-3D). No *E2* T-DNA homozygous plants were obtained due to

homozygous lethality [45]. Using CRISPR-Cas9 mutagenesis, we generated two weak alleles of *e2*

mutants (*e2-2 #17* and *e2-2 #215*) that were variable (Fig. S3E-3G). Phenotypical assay showed

that the *e2* single mutants, but not *e1* nor *e3* single mutant, displayed a mild ethylene insensitivity

both in roots and in hypocotyls (Fig. 3A-3C). We then generated a variety of their higher order

mutants and examined their ethylene responses (Fig. S3E-3G). The mild ethylene insensitive

phenotype of the *e2* single mutant was significantly enhanced in *e1e2* (*e1-2-2 e2-2 #17* and *e1-*

2-4 e2-2 #215) and *e2e3* (*e2-2 e3-2-1 #99* and *e2-2 e3-2-2 #67*) double mutants and in *e1e2e3*

triple mutants (*e1-2-2 e2-2 e3-2-1 #167* and *e1-2-2 e2-2 e3-2-2 #67*) (Fig. 3D-3F). No significant

difference was observed between the *e1e3* (*e1-2-2 e3-2-1* and *e1-2-2 e3-2-2*) double mutants

and *e1* or *e3* single mutants (Fig. 3A-3F). The *e1e2*, *e2e3*, and *e1e2e3* ethylene insensitive

phenotype were complemented by introducing *proE2:gE2-FLAG-GFP* into each mutant

background, respectively (Fig. S3H and S3I). These genetic results demonstrated that PDC is

involved in the ethylene response, E2 is the most important subunit genetically, and E1 and E3

enhance the function of E2 in the ethylene response.

To further understand the functions of E1, E2, and E3 in the ethylene response, we generated

their gain-of-function mutants and selected two individual transgenic lines of each with similar

protein expression levels for further ethylene response analysis (Fig. S4A-4C). Compared to Col-

0, all the plants that overexpressed *PDC E1, E2,* and *E3* subunits had enhanced ethylene

responsive phenotype in the presence of ACC, but the phenotypes of *E2ox* and *E3ox* did not

differ from Col-0 in the absence of ACC (Fig. S4D). This showed that the overexpression of *PDC*

E2 and *E3* alone was not sufficient to trigger ethylene response in the absence of the hormone

when they were not accumulated in the nucleus. Interestingly, *PDC E1*ox displayed a mild

ethylene hypersensitivity in the absence of ACC compared to Col-0, which is in line with the

observation that there was a basal E1 in the nucleus without ethylene treatment (Fig. S4D). Next,

we expressed *E1* and *E2* fused with nuclear localization signal peptide (NLS) derived from EIN2-

C (*E1-NLS-GFP* and *E2-NLS-GFP*) in Col-0 (Fig. S4E). E1-NLS-GFP and E2-NLS-GFP fusion

proteins could be localized to the nucleus in the absence of ethylene; more importantly, the *E1-*

NLS-GFP and *E2-NLS-GFP* plants displayed a clear ethylene hypersensitivity phenotype even in

the absence of ethylene (Fig. S4F and S4G). Together, these data demonstrate that the nuclear

localization of PDC can trigger the ethylene response.

In order to evaluate the function of PDC in ethylene response at a molecular level, we performed

RNA sequencing (mRNA-seq) using *e1-2-2 e2-2 #17* and *e2-2 e3-2-2 #67* double mutants treated

with or without 4 hours of ethylene gas (Fig. S5). We found that about 50% genes up-regulated

by ethylene in Col-0 were not differentially expressed in either of the two mutants in response to

ethylene (Fig. S6A and S6B). For further statistical quantification, we plotted the $\log_2$ fold change

($\log_2$FC) of each gene up-regulated by ethylene in Col-0 against that in each double mutant.

These genes were then divided into three groups: elevated group ($\log_2$FC in mutants is 30%

greater than in Col-0), unchanged group ($\log_2$FC in mutants is within 30% of that in Col-0), and

compromised group (log$_2$FC in mutants is 30% less than that in Col-0). We found that the elevation

of more than half of ethylene-up regulated genes in Col-0 was significantly compromised in both

double mutants (Fig. 3G-3I). Importantly, most of the compromised genes identified from *e1-2-2*

e2-2 #17 seedlings were also compromised in the *e2-2 e3-2-2 #67* seedlings (Fig. 3J), and the

log$_2$FC profile showed a high similarity (Fig. 3I and Fig. S6C). Further gene ontology analysis

using those upregulation compromised genes showed that the ethylene signaling genes were

overrepresented (Fig. 3K and Fig. S6D). Together, these data provide genetic and molecular

evidence that E1, E2, and E3 function coordinately to regulate ethylene response.

PDC regulates ethylene-dependent elevation of histone acetylation at H3K14 and H3K23

PDC catalyzes the synthesis of acetyl CoA, the substrate of histone acetylation. Given that the

acetylation of histone at H3K14 and H3K23 is induced by ethylene, we compared their global

levels in Col-0 and in different *pdc* mutants. The ethylene-induced global elevation of H3K14ac

and H3K23ac in Col-0 was still detected in *e1-2-2 e2-2 #17* but with lower levels. Whereas,

ethylene-induced global elevation was not detected in *e2-2 e3-2-2 #67*, nor in *e1-2-2 e2-2 e3-2-*

2 #67 (Fig. 4A). In contrast, the H3K14ac and H3K23ac levels were elevated in the plants that

overexpress *E2* with ethylene treatment, and in the *E2-NLS-GFP* transgenic plants even without

ethylene treatment (Fig. 4B). We then conducted H3K14ac and H3K23ac chromatin

immunoprecipitation sequencing (ChIP-seq) in *e1-2-2 e2-2 #17* (Fig. S7A-7D). H3K14ac and

H3K23ac levels were reduced in *e1-2-2 e2-2 #17* compared to that in Col-0 over genes that were

transcriptionally compromised in both *e1-2-2 e2-2 #17* and *e2-2 e3-2-2 #67* plants, and the

ethylene-induced elevation was significantly impaired (Fig. 4C, 4D, and Fig. S7E-7H). When the

ChIP signals within the first 500bp downstream of transcription starts sites was evaluated, an

even more significant reduction was detected in *e1-2-2 e2-2 #17* both with and without ethylene

treatments (Fig. 4E and 4F). We then conducted H3K14ac and H3K23ac ChIP-qPCR assays in

the selected targets, and the results further validated the ChIP-seq data (Fig. S7I and S7J). Given

that *e2-2 e3-2-2 #67* and *e1-2-2 e2-2 e3-2-2 #67* has a similar ethylene responsive phenotype as

e1-2-2 e2-2 #17, we also conducted ChIP-qPCR assays in those two mutants, and a similar result

was obtained (Fig. S7I and S7J). Altogether, these results suggest that PDC regulates the histone

acetylation H3K14ac and H3K23ac in response to ethylene.

Enzymatically active nuclear PDC regulates acetyl CoA production in the nucleus

Because the biological function of PDC is to generate acetyl CoA, we hypothesized that the

reduced histone acetylation level in the plants that lack PDC subunits results from the decreased

acetyl CoA in the nucleus. We first measured the nuclear acetyl CoA concentrations in Col-0 and

in the *e1-2-2 e2-2 e3-2-2 #67* triple mutant treated with 4 hours of ethylene gas by liquid

chromatography–(LC) coupled to MS (LC–MS), and the levels of nuclear acetyl CoA were

significantly decreased in the *pdc* triple mutant compared to levels in Col-0 (Fig. 5A and 5B). We

then decided to examine whether the nuclear PDC was functional to synthesize acetyl CoA. It has

been shown that dephosphorylated E1 is required for the PDC activity [44,46], therefore, we first

examined the phosphorylation status of E1 in the nucleus in response to ethylene. Western blot

assay by anti-pSer antibody showed that most E1 was in a dephosphorylated state in the nucleus

after ethylene treatment (Fig. 5C). Phos-tag electrophoresis assay confirmed this result (Fig. 5D).

Next, we conducted LC-MS/MS analysis of E1 phosphorylation. We detected phosphorylation on

S292 (Fig. S8A), a residue that is evolutionary conserved with the reported inhibitory

phosphorylation site in human E1 (Fig. S8B)[46]. Further quantification of E1 phosphopeptides by

MS showed that the ratio of phosphorylated E1 to non-phosphorylated E1 in the nucleus was

significantly lower than that in the cytosol after 4 hours of ethylene treatment (Fig. 5E), suggesting

that the dephosphorylated E1 is the main species in the nucleus to function in the presence of

ethylene. Next, we assessed E1 activities in the nucleus using a pyruvate dehydrogenase (PDH)

activity assay. E1 activity was detected in the nucleus in Col-0 and its activity was elevated by the

ethylene treatment (Fig. 5F). But the ethylene-induced E1 activity was not detected in *e1-2-2 e2-*

2#17 or in *e1-2-2 e2-2 e3-2-2#67* (Fig. 5F and Fig. S8C). To further directly monitor PDC acetyl

CoA biosynthesis enzyme activity in the nucleus in response to ethylene, we employed a ^{13}C

isotopic tracing experiment. By feeding 2,3-^{13}C$_2$ pyruvate to the nuclear extracts and followed by

the LC-MS/MS, we are able to measure 1,2-^{13}C$_2$ labeled-acetyl CoA (1,2-^{13}C$_2$ acetyl CoA) that

will be converted from 2,3-^{13}C$_2$ pyruvate by PDC complex (Fig. 5G and Fig. S8D). This experiment

assesses the nuclear PDC activity since PDC is the only enzyme that synthesizes acetyl CoA

from pyruvate. As shown in Fig. 5H, we detected 1,2-^{13}C$_2$ acetyl CoA in the nuclear extracts of

Col-0 treated with 4 hours of ethylene gas, and the 1,2-^{13}C$_2$ acetyl CoA levels were significantly

increased in the presence of ethylene (Fig. 5H and Fig. S8E). This suggests that functional PDC

complex is accumulated in the nucleus in response to ethylene in Col-0. Further comparison

showed no significant differences in the levels of 1,2-^{13}C$_2$ acetyl CoA in the nuclei of Col-0

compared to that of *e1-2-2 e2-2 #17* or of *e1-2-2 e2-2 e3-2-2 #67* in the absence of ethylene (Fig.

5H and Fig. S8E). However, the ethylene-induced elevation of 1,2-^{13}C$_2$ acetyl CoA detected from

Col-0 nucleus was significantly reduced from the nucleus of *e1-2-2 e2-2 #17* or *e1-2-2 e2-2 e3-*

2-2#67 (Fig. 5H), providing further biochemical evidence that functional PDC is translocated into

the nucleus in response to ethylene to synthesize acetyl CoA.

EIN2 is required for PDC nuclear translocation and function in response to ethylene

Given the fact that EIN2-C interacts with PDC in the nucleus, we next explored the genetic

relationship between EIN2-C and PDC subunits. By introducing *E1ox, E2ox*, and *E3ox* into the

ein2-5 null mutant, we obtained *E1ox/ein2-5, E2ox/ein2-5*, and *E3ox/ein2-5* plants with

comparable protein expression levels in *E1ox/*Col-0*, E2ox/*Col-0, and *E3ox/*Col-0 separately (Fig.

S9A-9C). We then compared their ethylene response with their parental plants and Col-0. The

ethylene hypersensitivity induced by PDC overexpression was entirely abolished in all *E1ox/ein2-*

5, E2ox/ein2-5, and *E3ox/ein2-5* plants; these plants displayed the same complete ethylene

insensitivity as *ein2-5* mutant (Fig. 6A). Similarly, the ethylene hypersensitivity induced by PDC

overexpression was entirely abolished in all *E1ox/ein3-1eil1-1, E2ox/ein3-1eil1-1*, and *E3ox/ein3-*

1eil1-1 plants (Fig. S9A-9D). These two genetic results suggest that both EIN2 and EIN3/EIL1

are required for the function of PDC in the ethylene response.

To further confirm the EIN2 dependency of PDC in ethylene response, we examined the cellular

localization of PDC in response to ethylene in the *ein2-5* mutant. We found that the ethylene-

induced nuclear accumulation of E1, E2, and E3 was abolished in *E1ox/ein2-5, E2ox/ein2-5*, and

E3ox/ein2-5 (Fig. 6B-6J). Similarly, no obvious ethylene-induced nuclear accumulation of PDC

proteins was observed in *E1ox/ein3-1 eil1-1, E2ox/ein3-1 eil1-1*, and *E3ox/ein3-1 eil1-1* (Fig. S9E-

9J). ChIP-qPCR analyses in selected ethylene-responsive target genes (Fig. S7E-7H) showed

that the elevation of H3K14ac and H3K23ac levels detected in *E2ox* were not detected in the

E2ox/ein2-5 plants, and the levels were similar to those in the *ein2-5* mutant both with and without

ethylene treatments (Fig. S9K). In addition, the enhanced transcriptional activation in response to

ethylene in *E2ox* was not detected in *E2ox/ein2-5* plants (Fig. S9L). Thus, our genetic and

molecular evidence supports the conclusion that PDC functions in the ethylene response in an

EIN2 dependent manner.

Acetyl CoA is important for histone acetylation and EIN2-C is essential for the ethylene-induced

histone acetylation elevation; therefore, we further investigated whether PDC regulates the

function of EIN2-C. We crossed $EIN2^{S645A}$, in which a point mutation at Ser645 mimics EIN2

constitutive dephosphorylation resulting in EIN2-C cleavage and nuclear accumulation, into *e1-*

2-2 e2-2 e3-2-2 #67 (*e1e2e3*) background to generate $EIN2^{S645A}$/*e1e2e3* plants (Fig. S10A).

Phenotypic analysis of $EIN2^{S645A}$/Col-0 and $EIN2^{S645A}$/*e1e2e3* showed that the ethylene

hypersensitivity caused by $EIN2^{S645A}$ was partially rescued in the *e1e2e3* mutant (Fig. 6H). But

the nuclear accumulation of EIN2-C was comparable in the $EIN2^{S645A}$/Col-0 and in

$EIN2^{S645A}$/*e1e2e3* (Fig. S10B and S10C), showing that PDC participates in the downstream

ethylene signaling and cellular response mediated by EIN2-C rather than the initial steps of EIN2-

C cleavage and translocation into the nucleus in ethylene signaling pathway. We then compared

the H3K14ac and H3K23ac levels at selected ethylene-responsive genes in $EIN2^{S645A}$/Col-0 and

$EIN2^{S645A}$/e1e2e3 plants by ChIP-qPCR. We found that the enhanced H3K14ac and H3K23ac in

$EIN2^{S645A}$/Col-0 was reduced in $EIN2^{S645A}$/e1e2e3, although levels were still higher than that in

Col-0 (Fig. 6I). Consistently, the enhancement of target gene expression in $EIN2^{S645A}$/Col-0 was

also partially rescued in $EIN2^{S645A}$/e1e2e3 (Fig. S10D). These findings suggest that PDC in the

nucleus produces acetyl CoA that is necessary for EIN2-C-regulated histone acetylation to

mediate the ethylene response.

Discussion

This study reveals a novel molecular mechanism by which mitochondria resided PDC complex

translocates into the nucleus to provide acetyl CoA necessary for EIN2-C mediated histone

acetylation and subsequent transcription activation in response to ethylene (Fig. S11). In the

presence of ethylene, PDC subunits E1, E2, and E3 accumulate in the nucleus to provide acetyl

CoA for EIN2 mediated histone acetylation elevation at H3K14 and H3K23 through their

interaction with EIN2-C, initiating the downstream ethylene-induced transcriptional cascade (Fig.

S11). Our research establishes a direct link between cell metabolism and histone modification in

the ethylene response that is mediated by the key factor EIN2, opening a new avenue for the

study of how plant hormone and metabolisms function to ensure the chromatin and transcription

regulation.

Cellular metabolism is a series of important biochemical reactions fueling development with

energy and biomass, and chromatin is a mighty consumer of cellular energy generated by

metabolism. The balance between metabolism and chromatin activities ensures cell homeostasis

and normal cell growth. PDC functioning as a partner of EIN2-C to regulate histone acetylation

levels at H3K14 and H3K23 marks in response to ethylene treatment suggests that EIN2-C

recruits translocated PDC to the ethylene-responsive gene loci to provide a sufficient level of local

acetyl CoA to achieve histone acetylation at H3K14 and H3K23 by HATs. Production of acetyl CoA

within the nucleus may promote its availability to histone acetyltransferase to facilitate a rapid ethylene

response. Thus, identification of the histone acetyl transferases that function together with EIN2-

C and PDC complex will be an immediate goal.

In the nucleus, acetyl CoA production machineries are finely tuned to control the local metabolite

levels at certain gene loci to regulate chromatin modification by interacting with different partners.

For instance, PDC E2 subunit has been reported to function in the same complex with pyruvate

kinase M2 (PKM2), histone acetyltransferase p300, and the transcription factor AhR to regulate

histone acetylation for transcription regulation to facilitate cell proliferation[47]. Nuclear metabolic

enzyme Acetyl CoA Synthetase 2 (ACSS2) associates with the histone acetyltransferase CREP

binding protein (CBP) to regulate the expression of key neuronal genes through histone

acetylation in the mouse hippocampus[48]. In response to glucose deprivation, ACSS2 can also

constitute a complex with the transcription factor EB to activate lysosomal and autophagosomal

genes after its nuclear translocation[42]. In ethylene response, EIN3 is the key transcription factor

that determines the target gene for transcription regulation[25,27]; thus, it will be interesting to explore

the connection between EIN3 and PDC complex.

In mammal, three main cytosolic enzymes that synthesize acetyl CoA, including PDC, Acyl-CoA

Synthetase Short Chain Family Member (ACSS2) and ATP Citrate Lyase (ACLY or ACL), were

shown to localize to the nucleus for different biological functions[38,40,42,43,47-49]. PDC complex was

first reported to localize to the nucleus in cell division and synthesize acetyl CoA to regulate cell

proliferation through histone acetylation[40]. ACSS2 catalyzes the conversion of acetyl CoA from

acetate in the nucleus for the regulation of long-term spatial memory and glucose starvation

response[42,48]; the nuclear ACLY utilizes citrate to produce acetyl CoA to promote histone

acetylation at double strand break sites for DNA repair[38]. Recently, ACL subunit A2 (ACLA2) has

been reported to function with HAT1 in rice to regulate cell division in developing endosperm,

suggesting the evolutionarily conserved mechanism of metabolic regulation on epigenetic

modification in both plant and mammal species[50]. We have noticed that although the ethylene

induced elevation of H3K14ac and H3K23ac is compromised in the *pdc* mutant, some extent of

ethylene response still remains. It is possible that since we could not obtain the *e2* null mutant

because of its lethality, the weak alleles we obtained still have partially functional E2 and therefore

manifest a rather weakened ethylene response. However, we could not exclude the possibility

that the other acetyl CoA producing enzymes will be functioning in the nucleus to modulate histone

acetylation in response to ethylene in the absence of the nuclear PDC in *pdc* mutants.

Investigating the involvement of other acetyl CoA synthesis pathways, such as ACSS[42,48] and

ACLY[38], in the ethylene response will potentially address the question.

In this research, we found that ethylene treatment triggers the translocation of PDC from the

cytosol to the nucleus to facilitate H3K14ac and H3K23ac elevation for ethylene responsive

transcriptional regulation. However, the question remains how the translocation of PDC complex

occurs at molecular level because PDC is a large protein complex without known nuclear

localization signal (NLS). Recent studies have shown that specialized contact points between

mitochondria and nucleus aid in the exchanges of metabolites and proteins between these two

organelles, and more importantly, mammalian PDC directly enters the nucleus at those contact

points across the nuclear envelop through mitofusin 2 (MFN2)-mediated mitochondria-nucleus

tethering, which is independent from the nuclear pore complex (NPC)[51,52]. However, whether it is

possible that nuclear PDC proteins are transported into the nucleus from the cytosol following

their protein synthesis is unknown. Investigating how *Arabidopsis* PDC enters the nucleus and

whether its nuclear entry follows the similar NPC-independent mechanism to regulate histone

acetylation in response to ethylene will provide more insights into the mitochondria and nucleus

retro-communication in the establishment of the proper ethylene response, providing a new

perspective of metabolic regulation in plant hormone response research.

Acknowledgements

We thank D. Hernandez for lab maintenance. We thank Dr. C. Petucci and Metabolomics Core of

Cardiovascular Institute at University of Pennsylvania for nuclear acetyl CoA LC/MS quantification

and Dr. K.S. Browning for nuclear-cytoplasmic fractionation setup. We thank P. Oliphint, A. Webb,

and Microscopy and Imaging Facility at The University of Texas at Austin for the support of

confocal microscope imaging. E1 phosphorylation MS analysis was supported by the Carnegie

endowment fund to the Carnegie mass spectrometry facility, and we thank Andres V. Reyes for

his assistance in MS analysis. We also thank *Arabidopsis* Biological Resource Center for the

Arabidopsis T-DNA insertion mutants. Z.S. was supported by the Research Awards from

Department of Integrative Biology and the Graduate Continuing Fellowship from The University

of Texas at Austin. This work was supported by grants from the National Institute of Health to H.Q.

(NIH-2R01 GM115879).

Author contributions

Z.S. and H.Q. designed the study. Z.S. performed most of the experiments and analysis. L.B.

performed LC-MS/MS analysis of 1,2-$^{13}C_2$ acetyl CoA and Y.B. performed MS analysis of E1

phosphorylation. Z. Shen performed Co-IP/MS. J.G.B. and P.K. helped genetic material

generation and imaging. S.K.A., T.J.D., and M.A.B. helped biochemical assays and high-

throughput sequencing and contributed to manuscript editing. S.-L.X., Z.-Y.W., S.P.B. and H.Q.

supervised the study. Z.S. and H.Q. wrote the paper. **Competing interests**: The authors declare

no competing interests. **Data and materials availability**: Further information and requests for all

unique materials generated in this study may be directed to and will be fulfilled by the

corresponding author Hong Qiao (hqiao@austin.utexas.edu). The high-throughput sequencing

data generated in this study have been deposited in the Gene Expression Omnibus (GEO)

database (accession no. GSE212540).

421
422

References

Achard, P. *et al.* The plant stress hormone ethylene controls floral transition via DELLA-dependent regulation of floral meristem-identity genes. *Proceedings of the National Academy of Sciences* **104**, 6484, doi:10.1073/pnas.0610717104 (2007).

Boutrot, F. *et al.* Direct transcriptional control of the Arabidopsis immune receptor FLS2 by the ethylene-dependent transcription factors EIN3 and EIL1. *Proceedings of the National Academy of Sciences* **107**, 14502, doi:10.1073/pnas.1003347107 (2010).

Chang, C. & Bleecker, A. B. Ethylene Biology. More Than a Gas. *Plant Physiology* **136**, 2895, doi:10.1104/pp.104.900122 (2004).

Dubois, M. *et al.* The ETHYLENE RESPONSE FACTORs ERF6 and ERF11 Antagonistically Regulate Mannitol-Induced Growth Inhibition in Arabidopsis. *Plant Physiol* **169**, 166-179, doi:10.1104/pp.15.00335 (2015).

Ecker, J. R. The ethylene signal transduction pathway in plants. *Science* **268**, 667-675, doi:10.1126/science.7732375 (1995).

Guo, H. & Ecker, J. R. Plant Responses to Ethylene Gas Are Mediated by SCF[EBF1/EBF2]-Dependent Proteolysis of EIN3 Transcription Factor. *Cell* **115**, 667-677, doi:10.1016/S0092-8674(03)00969-3 (2003).

Hartman, S. *et al.* Ethylene-mediated nitric oxide depletion pre-adapts plants to hypoxia stress. *Nature Communications* **10**, 4020, doi:10.1038/s41467-019-12045-4 (2019).

Haydon, M. J., Mielczarek, O., Frank, A., Román, Á. & Webb, A. A. R. Sucrose and Ethylene Signaling Interact to Modulate the Circadian Clock. *Plant Physiology* **175**, 947-958, doi:10.1104/pp.17.00592 (2017).

Mersmann, S., Bourdais, G., Rietz, S. & Robatzek, S. Ethylene signaling regulates accumulation of the FLS2 receptor and is required for the oxidative burst contributing to plant immunity. *Plant Physiol* **154**, 391-400, doi:10.1104/pp.110.154567 (2010).

Wang, L., Ko, E. E., Tran, J. & Qiao, H. TREE1-EIN3–mediated transcriptional repression inhibits shoot growth in response to ethylene. *Proceedings of the National Academy of Sciences* **117**, 29178-29189, doi:10.1073/pnas.2018735117 (2020).

Vaseva, I. I. *et al.* The plant hormone ethylene restricts Arabidopsis growth via the epidermis. *Proceedings of the National Academy of Sciences* **115**, E4130-E4139, doi:10.1073/pnas.1717649115 (2018).

Shao, Z. *et al.* Phosphorylation status of Bβ subunit acts as a switch to regulate the function of phosphatase PP2A in ethylene-mediated root growth inhibition. *New Phytologist* **236**, 1762-1778, doi:https://doi.org/10.1111/nph.18467 (2022).

He, C. *et al.* The retrograde signaling regulator ANAC017 recruits the MKK9–MPK3/6, ethylene, and auxin signaling pathways to balance mitochondrial dysfunction with growth. *The Plant Cell* **34**, 3460-3481, doi:10.1093/plcell/koac177 (2022).

Jurdak, R., Launay-Avon, A., Paysant-Le Roux, C. & Bailly, C. Retrograde signalling from the mitochondria to the nucleus translates the positive effect of ethylene on dormancy breaking of Arabidopsis thaliana seeds. *New Phytologist* **229**, 2192-2205, doi:https://doi.org/10.1111/nph.16985 (2021).

Alonso, J. M. *et al.* Five components of the ethylene-response pathway identified in a screen for weak ethylene-insensitive mutants in Arabidopsis. *Proceedings of the National Academy of Sciences* **100**, 2992, doi:10.1073/pnas.0438070100 (2003).

Chang, C., Kwok, S. F., Bleecker, A. B. & Meyerowitz, E. M. Arabidopsis ethylene-response gene ETR1: similarity of product to two-component regulators. *Science* **262**, 539-544, doi:10.1126/science.8211181 (1993).

472	17	Clark, K. L., Larsen, P. B., Wang, X. & Chang, C. Association of the Arabidopsis CTR1 Raf-like kinase with the ETR1 and ERS ethylene receptors. *Proceedings of the National Academy of Sciences* **95**, 5401, doi:10.1073/pnas.95.9.5401 (1998).

17   Clark, K. L., Larsen, P. B., Wang, X. & Chang, C. Association of the
Arabidopsis CTR1 Raf-like kinase with the ETR1 and ERS ethylene
receptors. *Proceedings of the National Academy of Sciences* **95**, 5401,
doi:10.1073/pnas.95.9.5401 (1998).

18   Ju, C. *et al.* CTR1 phosphorylates the central regulator EIN2 to control ethylene hormone
signaling from the ER membrane to the nucleus in Arabidopsis. *Proc Natl Acad Sci U S A*
109, 19486-19491, doi:10.1073/pnas.1214848109 (2012).

19   Alonso, J. M., Hirayama, T., Roman, G., Nourizadeh, S. & Ecker, J. R. EIN2, a bifunctional
transducer of ethylene and stress responses in Arabidopsis. *Science* **284**, 2148-2152,
doi:10.1126/science.284.5423.2148 (1999).

20   Qiao, H. *et al.* Processing and Subcellular Trafficking of ER-Tethered EIN2 Control
Response to Ethylene Gas. *Science* **338**, 390, doi:10.1126/science.1225974 (2012).

21   Qiao, H., Chang, K. N., Yazaki, J. & Ecker, J. R. Interplay between ethylene, ETP1/ETP2
F-box proteins, and degradation of EIN2 triggers ethylene responses in Arabidopsis.
Genes Dev **23**, 512-521, doi:10.1101/gad.1765709 (2009).

22   Merchante, C. *et al.* Gene-Specific Translation Regulation Mediated by the Hormone-
Signaling Molecule EIN2. *Cell* **163**, 684-697, doi:https://doi.org/10.1016/j.cell.2015.09.036
(2015).

23   Li, W. *et al.* EIN2-directed translational regulation of ethylene signaling in Arabidopsis. *Cell*
163, 670-683, doi:10.1016/j.cell.2015.09.037 (2015).

24   Chang, K. N. *et al.* Temporal transcriptional response to ethylene gas drives growth
hormone cross-regulation in Arabidopsis. *eLife* **2**, e00675, doi:10.7554/eLife.00675 (2013).

25   Wang, L. *et al.* EIN2-directed histone acetylation requires EIN3-mediated positive
feedback regulation in response to ethylene. *The Plant Cell* **33**, 322-337,
doi:10.1093/plcell/koaa029 (2020).

26   Zhang, F. *et al.* EIN2-dependent regulation of acetylation of histone H3K14 and non-
canonical histone H3K23 in ethylene signalling. *Nature Communications* **7**, 13018,
doi:10.1038/ncomms13018 (2016).

27   Zhang, F. *et al.* EIN2 mediates direct regulation of histone acetylation in the ethylene
response. *Proceedings of the National Academy of Sciences* **114**, 10274,
doi:10.1073/pnas.1707937114 (2017).

28   Bieluszewski, T. *et al.* NuA4 and H2A.Z control environmental responses and autotrophic
growth in Arabidopsis. *Nature Communications* **13**, 277, doi:10.1038/s41467-021-27882-
5 (2022).

29   He, X. J. *et al.* An effector of RNA-directed DNA methylation in arabidopsis is an
ARGONAUTE 4- and RNA-binding protein. *Cell* **137**, 498-508,
doi:10.1016/j.cell.2009.04.028 (2009).

30   Crevillén, P. *et al.* Epigenetic reprograming that prevents transgenerational inheritance
of the vernalized state. *Nature* **515**, 587-590, doi:10.1038/nature13722 (2014).

31   Feng, S., Jacobsen, S. E. & Reik, W. Epigenetic reprogramming in plant and animal
development. *Science* **330**, 622-627, doi:10.1126/science.1190614 (2010).

32   Cortijo, S. *et al.* Transcriptional Regulation of the Ambient Temperature Response by
H2A.Z Nucleosomes and HSF1 Transcription Factors in Arabidopsis. *Mol Plant* **10**, 1258-
1273, doi:10.1016/j.molp.2017.08.014 (2017).

33   Tessarz, P. & Kouzarides, T. Histone core modifications regulating nucleosome structure
and dynamics. *Nat Rev Mol Cell Biol* **15**, 703-708, doi:10.1038/nrm3890 (2014).

34   Zentner, G. E. & Henikoff, S. Regulation of nucleosome dynamics by histone modifications.
Nat Struct Mol Biol **20**, 259-266, doi:10.1038/nsmb.2470 (2013).

35   Pietrocola, F., Galluzzi, L., Bravo-San Pedro, José M., Madeo, F. & Kroemer, G. Acetyl
Coenzyme A: A Central Metabolite and Second Messenger. *Cell Metabolism* **21**, 805-821,
doi:https://doi.org/10.1016/j.cmet.2015.05.014 (2015).

36 Fan, J., Krautkramer, K. A., Feldman, J. L. & Denu, J. M. Metabolic regulation of histone
post-translational modifications. *ACS Chem Biol* **10**, 95-108, doi:10.1021/cb500846u
(2015).
37 Peng, M. *et al.* Aerobic glycolysis promotes T helper 1 cell differentiation through an
epigenetic mechanism. *Science* **354**, 481-484, doi:10.1126/science.aaf6284 (2016).
38 Sivanand, S. *et al.* Nuclear Acetyl-CoA Production by ACLY Promotes Homologous
Recombination. *Mol Cell* **67**, 252-265.e256, doi:10.1016/j.molcel.2017.06.008 (2017).
39 Hsieh, W. C. *et al.* Glucose starvation induces a switch in the histone acetylome for
activation of gluconeogenic and fat metabolism genes. *Mol Cell* **82**, 60-74.e65,
doi:10.1016/j.molcel.2021.12.015 (2022).
40 Sutendra, G. *et al.* A nuclear pyruvate dehydrogenase complex is important for the
generation of acetyl-CoA and histone acetylation. *Cell* **158**, 84-97,
doi:10.1016/j.cell.2014.04.046 (2014).
41 Nagaraj, R. *et al.* Nuclear Localization of Mitochondrial TCA Cycle Enzymes as a Critical
Step in Mammalian Zygotic Genome Activation. *Cell* **168**, 210-223.e211,
doi:10.1016/j.cell.2016.12.026 (2017).
42 Li, X. *et al.* Nucleus-Translocated ACSS2 Promotes Gene Transcription for Lysosomal
Biogenesis and Autophagy. *Mol Cell* **66**, 684-697.e689, doi:10.1016/j.molcel.2017.04.026
(2017).
43 Li, W. *et al.* Nuclear localization of mitochondrial TCA cycle enzymes modulates
pluripotency via histone acetylation. *Nature Communications* **13**, 7414,
doi:10.1038/s41467-022-35199-0 (2022).
44 Patel, M. S., Nemeria, N. S., Furey, W. & Jordan, F. The pyruvate dehydrogenase
complexes: structure-based function and regulation. *J Biol Chem* **289**, 16615-16623,
doi:10.1074/jbc.R114.563148 (2014).
45 Song, L. & Liu, D. Mutations in the three Arabidopsis genes that encode the E2 subunit of
the mitochondrial pyruvate dehydrogenase complex differentially affect enzymatic activity
and plant growth. *Plant Cell Rep* **34**, 1919-1926, doi:10.1007/s00299-015-1838-1 (2015).
46 Rardin, M. J., Wiley, S. E., Naviaux, R. K., Murphy, A. N. & Dixon, J. E. Monitoring
phosphorylation of the pyruvate dehydrogenase complex. *Anal Biochem* **389**, 157-164,
doi:10.1016/j.ab.2009.03.040 (2009).
47 Matsuda, S. *et al.* Nuclear pyruvate kinase M2 complex serves as a transcriptional
coactivator of arylhydrocarbon receptor. *Nucleic Acids Res* **44**, 636-647,
doi:10.1093/nar/gkv967 (2016).
48 Mews, P. *et al.* Acetyl-CoA synthetase regulates histone acetylation and hippocampal
memory. *Nature* **546**, 381-386, doi:10.1038/nature22405 (2017).
49 Chueh, F. Y. *et al.* Nuclear localization of pyruvate dehydrogenase complex-E2 (PDC-E2),
a mitochondrial enzyme, and its role in signal transducer and activator of transcription 5
(STAT5)-dependent gene transcription. *Cell Signal* **23**, 1170-1178,
doi:10.1016/j.cellsig.2011.03.004 (2011).
50 Xu, Q. *et al.* ACL and HAT1 form a nuclear module to acetylate histone H4K5 and promote
cell proliferation. *Nature Communications* **14**, 3265, doi:10.1038/s41467-023-39101-4
(2023).
51 Zervopoulos, S. D. *et al.* MFN2-driven mitochondria-to-nucleus tethering allows a non-
canonical nuclear entry pathway of the mitochondrial pyruvate dehydrogenase complex.
Molecular Cell **82**, 1066-1077.e1067, doi:https://doi.org/10.1016/j.molcel.2022.02.003
(2022).
52 Desai, R. *et al.* Mitochondria form contact sites with the nucleus to couple prosurvival
retrograde response. *Science Advances* **6**, eabc9955, doi:doi:10.1126/sciadv.abc9955
(2020).

53 Zhao, B. *et al.* MYB44-ENAP1/2 restricts HDT4 to regulate drought tolerance in
Arabidopsis. *PLOS Genetics* **18**, e1010473, doi:10.1371/journal.pgen.1010473 (2022).
54 Wang, Z. P. *et al.* Egg cell-specific promoter-controlled CRISPR/Cas9 efficiently
generates homozygous mutants for multiple target genes in Arabidopsis in a single
generation. *Genome Biol* **16**, 144, doi:10.1186/s13059-015-0715-0 (2015).
55 Minkenberg, B., Zhang, J., Xie, K. & Yang, Y. CRISPR-PLANT v2: an online resource for
highly specific guide RNA spacers based on improved off-target analysis. *Plant Biotechnol
J* **17**, 5-8, doi:10.1111/pbi.13025 (2019).
56 Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21,
doi:10.1093/bioinformatics/bts635 (2013).
57 Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biology* **15**, 550, doi:10.1186/s13059-014-0550-
8 (2014).
58 Tian, T. *et al.* agriGO v2.0: a GO analysis toolkit for the agricultural community, 2017
update. *Nucleic Acids Res* **45**, W122-w129, doi:10.1093/nar/gkx382 (2017).
59 Ramírez, F., Dündar, F., Diehl, S., Grüning, B. A. & Manke, T. deepTools: a flexible
platform for exploring deep-sequencing data. *Nucleic Acids Research* **42**, W187-W191,
doi:10.1093/nar/gku365 (2014).
60 Guan, S., Price, J. C., Prusiner, S. B., Ghaemmaghami, S. & Burlingame, A. L. A data
processing pipeline for mammalian proteome dynamics studies using stable isotope
metabolic labeling. *Mol Cell Proteomics* **10**, M111.010728,
doi:10.1074/mcp.M111.010728 (2011).
61 MacLean, B. *et al.* Skyline: an open source document editor for creating and analyzing
targeted proteomics experiments. *Bioinformatics* **26**, 966-968,
doi:10.1093/bioinformatics/btq054 (2010).
62 Basu, S. S. & Blair, I. A. SILEC: a protocol for generating and using isotopically labeled
coenzyme A mass spectrometry standards. *Nature Protocols* **7**, 1-11,
doi:10.1038/nprot.2011.421 (2012).

604
605
606

Figures and Figure Legends

Figure 1. PDC interacts with EIN2-C in the nucleus in response to ethylene.

(**A**) *In vitro* pull-down experiment to validate the interaction between EIN2-C and PDC E1, E2,

and E3 subunits. (**B**) Subcellular localization of EIN2-YFP fusion protein in *Arabidopsis* leaf

epidermal cells with MitoTracker Red staining without (upper panel) and with (lower panel) the

presence of 10μM ACC. Red arrowhead indicates mitochondria signal by MitoTracker staining;

yellow arrowhead indicates EIN2-YFP fluorescence signal. Scale bar is 20 μm. (**C**-**E**) *In vivo* co-

immunoprecipitation assay to examine the interaction between EIN2-C and E1(**C**), E2 (**D**), and

E3 (**E**) in the indicated transgenic plants. 3-day-old etiolated seedlings carrying both PDC and

EIN2 fusion proteins treated with air or 4 hours of ethylene gas were fractionated to isolate cytosol

and nuclei for the immunoprecipitation with either Flag-Trap magnetic agarose (DYKDDDDK Fab-

Trap) or anti-HA magnetic beads, respectively. The immunoprecipitation with IgG beads serves

as a negative control. Phospho Enol Pyruvate Carboxylase (PEPC) was used as cytosolic marker

protein; Cytochrome C (CytC) is a mitochondrial marker protein, BiP is an ER marker to assess

nuclear extraction purity. Histone H3 is a loading control for nuclear fractions. (**F**-**H**) Confocal

microscopy images of BiFC assay showing the interaction between EIN2 and PDC E1, E2, and

E3 subunits in the nucleus after ethylene treatment. Agrobacteria containing indicated paired

constructs was co-infiltrated into tobacco leaves and the YFP fluorescence was observed two

days after infiltration with or without 4 hours of ACC treatment. NLS-BFP was used as nuclear

marker. Scale bars is 50μm.

629
630

Figure 2. PDC accumulates in the nucleus in response to ethylene.

(**A**-**C**) Fractionation western blot to examine the subcellular localization of PDC in of *E1-YFP-HA* (**A**), *E2-FLAG-GFP* (**B**), and *E3-FLAG-GFP* (**C**) transgenic plants with time series of ethylene gas treatments. PDC E1 was probed with anti-HA, and E2 and E3 were probed with anti-FLAG in total protein extracts, cytoplasmic fractions, and nuclear fractions. PEPC, ATPB, FUM1/2, Cyt C, and histone H3 were used to assess purities of nuclear and cytosolic fractionations and loading controls. Blue number indicates PDC band intensity that normalized to cytoplasmic PEPC signal. Red number indicates PDC band intensity that normalized to histone H3 signal. (**D**-**F**) Confocal microscopy images showing the subcellular localization of E1-YFP-HA (**D**), E2-FLAG-GFP (**E**), and E3-FLAG-GFP (**F**) with time series of ethylene gas treatments. DAPI staining labels nuclei. Scale bars is 20μm.

Figure 3. PDC is involved in the ethylene response.

(**A**) Photographs of seedlings with mutations in individual PDC subunits and Col-0 grown on MS

medium containing 1 μM, 2 μM, and 5 μM ACC or without ACC. (**B** and **C**) Measurements of

hypocotyl lengths (**B**) and root lengths (**C**) of indicated mutants. Values are means $\pm$ SD of at

least 30 seedlings. Different letters represent significant differences between each genotype

calculated by a One-way ANOVA test followed by Tukey's HSD test with $P \leq 0.05$. (**D**) Photographs

of representative double and triple PDC subunit mutants and Col-0 grown on MS medium

containing 1 μM, 2 μM, and 5 μM ACC or without ACC were selected for the photograph. (**E** and

F) Measurements of hypocotyl lengths and root lengths from indicated plants. Each value is

means $\pm$ SD of at least 30 seedlings. Different letters indicate significant differences between

different genotypes with $P \leq 0.05$ that calculated by a One-way ANOVA test and followed by

Tukey's HSD test for multiple comparisons. (**G** and **H**) Scatter plots to compare the $\log_2$(Fold

Change) of ethylene up-regulated genes in Col-0 with that in *e1-2-2 e2-2 #17* (**G**), or in *e2-2 e3-*

2-2 # 67 (**H**) in mRNA-seq. Each dot represents a gene that its transcription level is significantly

elevated by ethylene gas treatment in Col-0 ($\log_2$(Fold Change) > 1, *p*-adjust < 0.05). (**I**) Violin

plot of the distributions of $\log_2$(Fold Change) of ethylene up-regulated genes in Col-0 and in the

indicated plants. *P* values were determined by a two-tailed *t* test. (**J**) Venn diagram to show the

ethylene up-regulated genes that were compromised in *e1-2-2 e2-2 #17* and in *e2-2 e3-2-2 # 67*.

(**K**) Gene ontology analysis of the genes that are transcriptionally co-compromised in both *e1-2-*

2 e2-2 #17 and in *e2-2 e3-2-2 #67*. Top 15 GO categories ranked by *P* values were plotted. Dot

size represents the percentage of genes from each GO category of all co-compromised genes

(Gene ratio %) and dot colors indicate the *P* value of the GO categories.

Figure 4. PDC is involved in the ethylene-induced histone acetylation.

(**A**) Western blot analysis of total histone acetylation of H3K14 and H3K23 in response to ethylene

in different genetic backgrounds as indicated in the figure. Anti-H3 western blot served as a

loading control. (**B**) H3K14ac and H3K23ac levels in Col-0, *E2ox,* and *E2-NLS-GFP* treated with

air or 4 hours of ethylene gas. Histone H3 served as a loading control. Red number indicates the

quantification of H3K14ac and H3K23ac western blot signal intensity normalized with that of

histone H3. (**C** and **D**) Heatmaps of H3K14ac (**C**) and H3K23ac (**D**) ChIP-seq signal ($\log_2$ ChIP

signal) from the genes that their ethylene-induced expressions are co-compromised in *e1-2-2 e2-*

2 #17 and *e2-2 e3-2-2 #67*. TSS, transcription start site; TTS, transcription termination site. (**E**

and **F**) Violin plots illustrating $\log_2$ normalized H3K14ac ChIP signal (**E**) and H3K23ac ChIP signal

(**F**) from 500bp downstream of TSS that in the genes that were co-compromised in the *pdc* double

mutants in the indicated genotypes and conditions. *P* values were calculated by a two-tailed *t* test.

686
687

Figure 5. Nuclear PDC is enzymatically active to produce acetyl CoA.

(**A** and **B**) LC-MS detection of acetyl CoA (**A**) and malonyl CoA (**B**) in purified nuclei from Col-0

and *e1-2-2 e2-2 e3-2-2 #67* etiolated seedlings treated with 4 hours of ethylene gas. Total protein

mass was used to normalize metabolite concentration. *P* values were calculated by a two-tailed *t*

test and each data point was plotted as a dot in the bar graph. (**C**) Western blot analysis of the

phosphorylation status of E1 using anti-pSer antibody in the cytosolic fraction and nuclear

fractions from *E1ox* seedlings treated with 4 hours of ethylene gas. (**D**) Phos-tag phosphoprotein

mobility shift gel of E1 phosphorylation status in the cytosolic fraction and nuclear fractions of

E1ox etiolated seedlings treated with 4 hours of ethylene gas. Black arrow shows non-

phosphorylated E1; red arrow indicates phosphorylated E1 protein. PEPC, CytC, and H3 were

used to assess purities of nuclear and cytosolic fractionations. (**E**) The ratios of phosphorylated

to non-phosphorylated YHGHpSMSDPGSTYR E1 peptides in the cytosolic versus in the nuclear

fractions from *E1ox* with 4 hours of ethylene gas treatment. Peak area values were collected from

three replicates. *P* value was calculated by *t*-test. (**F**) Pyruvate dehydrogenase activity assays in

the nuclear fraction from Col-0 and indicated *pdc* mutants treated with 4 hours of ethylene gas.

(**G**) Schematic diagrams to show the conversion of 2,3-^{13}C$_2$ pyruvate to 1,2-^{13}C$_2$ acetyl CoA. (**H**)

Normalized LC MS/MS measured the concentration of 1,2-^{13}C$_2$ acetyl CoA that converted from

2,3-^{13}C$_2$ pyruvate by using the nuclear extracts from indicated plants with or without 4 hours of

ethylene gas treatment. Quantification from four replicates were normalized by total protein input.

Different letters represent significant differences between each group calculated by a one-way

ANOVA test followed by Tukey's HSD test.

Figure 6. PDC functions in the ethylene response in an EIN2-C-dependent manner.

(**A**) Epistasis analysis of PDC *E1ox*, *E2ox*, and *E3ox* in the *ein2-5* mutant. The seedlings were grown on MS medium containing with or without 10μM ACC in the dark for 3 days before being photographed. (**B-D**) Examination of the protein levels of E1 in *E1ox/*Col-0 and *E1ox/ein2-5* (**B**), E2 in *E2ox/*Col-0 and *E2ox/ein2-5* (**C**), and E3 in *E3ox/*Col-0 and *E3ox/ein2-5* (**D**) with 0 or 12-hour ethylene gas treatment. PEPC is cytoplasmic marker; CytC is a mitochondrial marker;

histone H3 is a nuclear marker for a loading control. (**E-J**) Subcellular localization of E1-YFP-HA

(**E** and **F**), E2-FLAG-GFP (**G** and **H**), and E3-FLAG-GFP (**I** and **J**) in Col-0 or *ein2-5* mutant

without or with 12 hours of ethylene gas treatment, respectively. Scale bars is 20µm. (**K**)

Photographs of 3-day old etiolated *EIN2^{S645A}/*Col-0 and *EIN2^{S645A}/e1e2e3* seedlings grown on MS

medium containing indicated ACC in the dark for 3 days before being photographed. (**L**) ChIP-

qPCR to evaluate H3K14ac (upper panel) and H3K23ac (lower panel) enrichment over selected

genes in the indicated etiolated seedlings treated with air or 4 hours of ethylene gas. Different

letters represent significant differences between each genotype and treatment condition that are

calculated by a one-way ANOVA test followed by Tukey's HSD test with $P \leq 0.05$.

731