Early antiviral CD4 and CD8 T cell responses and antibodies are associated with upper respiratory tract clearance of SARS-CoV-2

Sydney I. Ramirez[1, 2], Paul G. Lopez[1], Farhoud Faraji[3], Urvi M. Parikh[4], Amy Heaps[4], Justin Ritz[5], Carlee Moser[5], Joseph J. Eron[6], David A. Wohl[7], Judith S. Currier[7], Eric S. Daar[8], Alex L. Greninger[9], Paul Klekotka[10], Alba Grifoni[1], Daniela Weiskopf[1,2], Alessandro Sette[1,2], Bjoern Peters[1,2], Michael D. Hughes[5], Kara W. Chew[7], Davey M. Smith[2†], and Shane Crotty[1,2†], for the ACTIV-2/A5401 Study Team

[1]Center for Vaccine Innovation, La Jolla Institute for Immunology, La Jolla, CA, USA
[2]University of California San Diego, Department of Medicine, Division of Infectious Diseases and Global Public Health, La Jolla, CA, USA
[3]University of California San Diego, Department of Otolaryngology-Head and Neck Surgery, La Jolla, CA, USA
[4]Division of Infectious Diseases, Department of Medicine, University of Pittsburgh School of Medicine, Pittsburgh, PA, USA
[5]Center for Biostatistics in AIDS Research, Harvard T.H. Chan School of Public Health, Boston, MA, USA
[6]Department of Medicine, University of North Carolina at Chapel Hill School of Medicine, Chapel Hill, NC, USA
[7]Division of Infectious Diseases, Department of Medicine, David Geffen School of Medicine at University of California, Los Angeles, Los Angeles, CA, USA
[8]Lundquist Institute at Harbor-UCLA Medical Center, Torrance, CA, USA
[9]Department of Laboratory Medicine and Pathology, University of Washington Medical Center, Seattle, WA, USA
[10]Eli Lilly and Company, San Diego, CA, USA

[†]Correspondence:
Shane Crotty, 9420 Athena Circle, La Jolla, CA, 92037, +1 858-752-6500, shane@lji.org;
Davey M. Smith, 9500 Gilman Drive #0679, La Jolla, CA, 92093, +1 858-642-1620, d13smith@health.ucsd.edu.

Running title: Early T cell responses to SARS-CoV-2 associated with upper airway viral clearance

T cells are involved in protective immunity against numerous viral infections[1]. Data regarding functional roles of human T cells in SARS-CoV-2 (SARS2) viral clearance in primary COVID-19 are limited. To address this knowledge gap, samples were assessed for associations between SARS2 upper respiratory tract viral RNA levels and early virus-specific adaptive immune responses, for 95 unvaccinated clinical trial participants with acute primary COVID-19. Functionality and quantity of acute SARS2-specific CD4 and CD8 T cell responses were evaluated, in addition to antibody responses. Most individuals with acute COVID-19 developed SARS2-specific T cell responses within 6 days of symptom onset. Early CD4 T cell and CD8 T cell responses were strongly associated with reduced upper respiratory tract SARS2 viral RNA, independent of neutralizing antibody titers. Overall, these findings indicate a protective role for SARS2–specific T cells during acute COVID-19.

MAIN

COVID-19 still killed 73,513 Americans in 2023[2]. Worldwide death tolls in 2023 were substantial, with over 1,000,000 COVID-19 deaths estimated for mainland China alone[3,4]. Since the emergence of severe acute respiratory syndrome coronavirus 2 (SARS2) as a novel human pathogen, much has been learned about protective immunity to SARS2, both in the context of infection and COVID-19 vaccines. Serologic correlates of protection have been established for vaccines[5–12]. In the context of prophylaxis, virus neutralization by monoclonal antibodies (mAb) has been demonstrated as one mechanism of protective immunity[13–16]. In contrast, in the context of infection the relative importance of immune system compartments may differ, due to the substantial differences in kinetics of primary versus memory immune responses and lack of pre-existing antibodies. Studies of primary adaptive immunity to acute SARS2 infection offer a key opportunity to evaluate early humoral and cellular immune responses and their individual contributions to protection. Cellular immunity may be required for viral control and clearance during acute infection[1,17–20]. While acute and memory T cell responses clearly occur to SARS2 infection[21–27] and COVID-19 vaccines[28–35], evidence of functional protective roles of T cells have been limited in humans, particularly by assessing non-hospitalized COVID-19 cases[1,20]. Due to the combined difficulties of identifying early acute COVID-19 cases, recruiting those individuals, and the technical challenges of measuring virus-specific T cell responses and viral loads concomitantly, most acute T cell studies have been limited to small cohorts, including hospitalized cases, sometimes relatively late in disease, and often without viral load measurements[25,36–39]. There is evidence for cellular immunity contributing to SARS2 protection in non-human primate models[1,20]. Breakthrough infections represent a different context for studying protective immunity in humans, in which there is evidence for contributions of memory T cells[40,41]. Given expanded interest in next-generation COVID-19 and pan-sarbecovirus vaccines with T cell-specific components[42,43], or vaccines that are entirely T cell-based[33,44,45], future vaccine designs and clinical trial designs would benefit from better fundamental understanding of T cell protective immunity to COVID-19 in humans6/10/24 10:31:00 AM.

Longitudinal data were collected for 95 individuals with primary SARS2 infection in 2020 prior to the availability of COVID-19 vaccines, with sampling including nasopharyngeal (NP) swabs for SARS2 RNA levels, serum, and peripheral blood mononuclear cells (PBMCs), all collected in the context of a randomized, controlled clinical trial (ACTIV-2/A5401, NCT04518410). Herein, serum antibodies, SARS2-specific CD4 T cell, and SARS2-specific CD8 T cell response measurements are reported, which is the

largest and most comprehensive acute viral and antigen-specific immune response data set of its kind.

All participants were enrolled within 7 days from positive SARS2 testing and 10 days from COVID-19 symptom onset[24,47]. Individuals studied herein were among those randomized to receive either 700 mg of the mAb bamlanivimab or placebo (saline) intravenously (**Fig. 1A**) on study day 0 (median of 6 days post-symptom onset; **Fig. 1B**). The 46 bamlanivimab treatment and 49 placebo subjects were similar with respect to baseline characteristics such as age, sex at birth, risk for progression to severe COVID-19, and time post-symptom onset at randomization (**Fig. 1A and S1A**, and ref[24,47]). Peripheral blood was collected prior to mAb or placebo administration on study day 0. NP swabs were collected from all 95 participants by trained study staff prior to treatment on study day 0. SARS2 viral RNA was detectable in the NP swabs of nearly all individuals (89%) at study day 0 (**Fig. 1C**), and did not differ between the bamlanivimab (treatment) and placebo groups (**Fig. S1B**). SARS2 NP RNA at day 0 was not associated with participant age ($r = 0.18$, $p = 9.2 \times 10^{-2}$, **Fig. S1C**). SARS2 NP RNA levels declined over time (**Fig. 1D**), as expected[48–50], and as previously reported for the full ACTIV-2/A5401 cohort[47].

Acute adaptive immune responses were measured for all participants on study day 0 because both NP swab and peripheral blood samples were collected prior to any treatment. Additionally, longitudinal responses were measured in the 49 placebo group participants on study days 7 and 28. We previously reported day 28 antibody, memory CD4 T cell, and memory CD8 T cell outcomes[24]. Early (study day 0) SARS2-specific CD4 T cell response magnitude and functionality were evaluated by multiple techniques and using multiple SARS2 antigens. Activation induced marker (AIM) and hybrid AIM plus intracellular cytokine staining (AIM+ICS) T cell assays were employed (**Fig. 2A-B**, **Fig. S2A**). SARS2 Spike (S)- and non-Spike-specific CD4 T cells were measured by AIM assays following stimulation with S and non-S (CD4-RE[31]) peptide megapools (MP). Two sets of AIM phenotyping pairs were utilized (surface OX40+ and 41BB+, **Fig. 2A**; surface OX40+ and CD40L+, **Fig. S2B**). SARS2-specific CD4 T cell frequencies were comparable by AIM using OX40+41BB+ or OX40+CD40L+ surface marker expression (**Fig. 2A and S2B**). 77% of individuals were positive for S-specific CD4 T cells, and 79% of individuals were positive for non-S CD4 T cells (**Fig. 2A**). Combined responses to SARS2 S- and non-S epitopes were also quantified (**Fig. 2A and S2B. Fig. S2C-G**). Overall, substantial CD4 T cell responses were observed.

SARS2-specific CD4 T cells were also phenotyped for cytokine production by ICS following stimulation with the S and non-S MPs, including interferon gamma (IFNγ), granzyme B (GzmB), tumor necrosis factor (TNF), and interleukin-2 (IL-2) among CD40L+ cells (**Fig. 2B-F and S2H-J**). IFNγ was the most expressed cytokine at day 0 (**Fig. 2B**) and is known to be important in control of multiple viral infections in mouse models. 85-87% of individuals had combined (S plus non-S) SARS2–specific CD4 T cell responses by AIM or IFNγ ICS (**Fig. 2A-B**). SARS2-specific CD4 T cells were polyfunctional, with 69% producing at least 2 cytokines, and 53% producing 3 or more cytokines (**Fig. 2B-F**). SARS2-specific circulating T follicular helper cells (cT_FH, CXCR5+) were measured. Similar response rates were seen for SARS2-specific cT_FH as for total SARS2-specific CD4 T cells, with 79-84% of individuals had SARS2-specific T_FH cells at day 0 (**Fig. 2G-H**). Overall, SARS2-specific CD4 T cells with different functionalities and differentiation were present.

In parallel, study day 0 SARS2 S- and non-S-specific CD8 T cell responses were evaluated by AIM (surface CD69+41BB+) and ICS following stimulation with SARS2 S and non-S (CD8-RE[31]) MPs (**Fig. S3A, D-F**). 63% of individuals were positive for an acute SARS2–specific CD8 T cell response at study day 0 by

AIM (**Fig. S3B**). 62% of individuals were positive for an acute SARS2–specific CD8 T cell response by ICS (surface CD69$^+$ and intracellular IFNγ$^+$, **Fig. 3A**). As was observed for SARS2-specific CD4 T cells, IFNγ was the most commonly produced cytokine by SARS2-specific CD8 T cells (**Fig. 3A**). Background signals were lower for the CD69$^+$IFNγ$^+$ CD8 T cell assay (**Fig. 3A** LOQ) compared to CD8 AIM (**Fig. S3B** LOQ) and thus CD8 ICS was used as the primary quantification of SARS2-specific CD8 T cells in downstream analyses (see Methods for additional details). As expected, CD8 T cell responses did not vary on study day 0 when assessed by clinical trial group assignment (**Fig. S3C**). Most CD69$^+$ IFNγ-producing SARS2-specific CD8 T cells also produced GzmB (73%, **Fig. 3B and S3D**). Similar to SARS2-specific CD4 T cells, SARS2-specific CD8 T cells were predominantly polyfunctional, with 69% producing at least 2 cytokines and 58% producing 3 or more cytokines (**Fig. 3A-E**). Overall, polyfunctional virus-specific CD4 and CD8 T cell responses were detected in a substantial proportion of individuals during an acute primary SARS2 infection with COVID-19.

We examined the kinetics of T cell and antibody responses in the placebo group in more detail. Longitudinal assessment of SARS2-specific CD4 and CD8 T cell frequencies on study days 0 (**Fig. 2-3**), 7 (**Fig. 4A-C and S4A**), and 28 (**Fig. 4A-C, S4A**, and ref.[24]) showed that AIM$^+$ CD4 T cell frequencies and the fraction of positive responders increased over time (**Fig. 4A and S4A**). SARS2-specific IFNγ$^+$ CD4 and CD8 T cell frequencies remained stable (**Fig. 4B-C**). SARS2 serologic assessments, included nAb titers to ancestral SARS2 by lentiviral pseudovirus neutralization assays (PSV NT50) and receptor binding domain (RBD) binding IgG titers, were measured on study day 0 (**Fig. 4D-E** and ref.[24] for RBD IgG). 62% of participants had positive nAb titers (**Fig. 4D, Fig. 4SB**). As expected, nAb and RBD IgG titers increased between study days 0 and 7 (**Fig. S4D-E**).

To evaluate potential protective activities of early T cell responses, associations were examined between day 0 NP viral RNA levels and SARS2–specific CD4 and CD8 T cell responses. Early CD4 T cell responses were associated with lower levels of SARS2 NP RNA based on all three main CD4 T cell measurements (OX40$^+$41BB$^+$ AIM Spearman r = -0.51, p = 2.6x10^{-7}, **Fig. 5A**; OX40$^+$CD40L$^+$ AIM r = -0.45, p = 9.3X10^{-6}, **Fig. S5A**; IFNγ$^+$, r = -0.36, p = 5x10^{-4}, **Fig. 5B**). Early CD8 T cell responses were also associated with lower levels of SARS2 NP RNA (IFNγ$^+$ r = -0.44, p = 1.1x10^{-5}, **Fig. 5C**; CD69$^+$41BB$^+$ AIM r = -0.39, p = 1.4x10^{-4}, **Fig. S5C**). T cell and SARS2 NP RNA associations were strongest when using combined S plus non-S T cell response measurements, but the statistical associations were similar when S and non-S antigen-specific responses were evaluated separately (**Fig. S5D-J**). A positive correlation was observed between SARS2-specific CD4 and CD8 T cell responses (**Fig. S5K**). A positive correlation was observed between day 0 nAb and RBD IgG titers (**Fig. S5L**), as expected[48]. Notably, nAb titers measured at study day 0 were also associated with lower levels of SARS2 NP RNA, but the association with nAb was weaker than the association with either CD4 or CD8 T cells (r = -0.33, p = 1.2x10^{-3}, **Fig. 5D**).

Given the biological dependence of the antibody response on CD4 T cell help mediated by T$_{FH}$ cells[51,52], some statistical association between antibodies and viral load would always be expected, even if, hypothetically, viral control was exclusively mediated by T cells. S-specific cT$_{FH}$ cells had relatively modest associations with SARS2 NP RNA that were weaker than other AIM CD4 T cell response associations (Spearman r = -0.37 to 0.39, p = 1.6 to 2.8 x10^{-4}, **Fig. S5I-J**). Analysis of covariance (ANCOVA) testing was performed to better evaluate if the putative protective relationships observed between early SARS2-specific CD4 T cell responses and SARS2 NP RNA remained after accounting for antibodies, based

on day 0 nAb serostatus. The associations between SARS2-specific CD4 T cells and lower NP viral RNA did not show a dependence on nAb seropositivity (**Fig. S5M-O**). SARS2-specific CD8 T cell associations with lower SARS2 NP RNA also did not show a dependence on nAb seropositivity (**Fig. S5M**). Thus, early SARS2–specific CD4 and CD8 T cell responses were specifically associated with reduced upper airway SARS2 viral loads in unvaccinated, non-hospitalized cases of acute COVID-19.

Participants in the study enrolled at a range of days post-symptom onset (PSO), which provides richness to the dataset, but raised the possibility that the immune response associations with reduced NP viral RNA levels might only be surrogates for time from infection. To test this possibility, an analysis was performed modeling T cell responses and NP viral RNA levels adjusted to account for time PSO to study entry (day 0), using a log-linear regression model of the raw data (**Fig. S6A-E**, see Methods). If the original T cell associations with reduction of SARS2 RNA levels were only surrogates of time from infection (measured as days PSO), T cell associations should be lost upon adjustment for days PSO. If instead the T cell associations reflected a functional relationship, statistical associations between T cell responses and lower viral RNA would be expected to remain. After accounting for days PSO by extrapolating viral NP RNA values and adaptive immune responses to day 6 PSO to account for sample collections on different days PSO using a simple log-linear regression model (**Fig. S5A-E**), the relationships between CD4 and CD8 T cell responses and reduced viral NP RNA levels remained statistically significant (CD4 AIM r = -0.39, p = 1.9×10^{-4} **Fig. S6F**; CD4 IFNγ$^+$ r = -0.35, p = 1.0×10^{-3} **Fig. S6G**; CD8 IFNγ$^+$ r = -0.31, p = 4.4×10^{-3} **Fig. S6H**), consistent with a functional role for T cells in control of NP viral RNA levels. Associations between nAb titers and viral NP RNA levels also remained after accounting for days PSO (r = -0.32, p = 3.0×10^{-3} **Fig. S6I**). Altogether, these data are evidence of functional relationships between the speed of acute CD4 and CD8 T cell responses, and antibodies, in humans and control of SARS2 NP viral loads.

In summary, this study provides new insights into the relationship between acute adaptive immune responses to SARS2 and viral control. It has been hypothesized that SARS2–specific antiviral T cell responses may contribute to protection from severe disease by reducing viral burden, enhancing speed of viral clearance; however, few human studies have directly measured the role of cellular immunity on viral control of primary infection[39,53]. Here, using samples from 95 individuals from a randomized clinical trial, the largest and most comprehensive virus-specific acute immune response data set of its kind was generated. The data demonstrate that (i) SARS2-specific CD4 and CD8 T cells could be detected in the majority of adults with outpatient COVID-19 within days from symptom onset, (ii) the T cells were polyfunctional, (iii) stronger early SARS2-specific CD4 T cell responses were associated with lower viral RNA levels in the upper airway, (iv) stronger SARS2-specific CD8 T cell responses were associated with lower viral RNA levels in the upper airway, (v) the associations of T cell responses with lower viral RNA were independent of neutralizing antibody responses and (vi) remained significant after accounting for time from COVID-19 symptom onset. The conclusions from these analyses of T cell associations with viral control were reproducible by multiple different measurements of CD4 T cell and CD8 T cell functionalities and SARS2 antigen specificities. Overall, these findings suggest that SARS2–specific T cells play a role in protective immunity to SARS2 during acute infection.

In a smaller recent study (n=25 acute infections sampled), Eser et al. found a strong association between nucleocapsid-specific IFNγ$^+$ CD4 T cells and lower viral loads[39]. Those results are consistent with the CD4 T cell findings here, except the findings here were that T cell responses to S or non-S antigens

were both associated with reduced viral loads, and were approximately equally associated with reduced viral loads. Eser et al. observed less of an association for CD8 T cell responses and lower viral loads, perhaps because the study sample size was limited[39].

In a human controlled SARS2 infection study with a limited infectious dose (53% of individuals were infected), total CD38[+]Ki67[+] CD8 T cells in peripheral blood were the strongest association with increased viral clearance[53]. Those data are consistent with the CD8 T cell findings here. A strength of that study is the precisely defined time of infection. Several factors distinguish this study. The cohort size here is substantially larger (95 compared to 18) and presumably includes subjects infected by a range of viral exposure doses. The virus-specific T cell assays herein measured CD8 and CD4 T cell epitopes spanning S and non-S antigens. Lastly, in the controlled infection study[53] all subjects were young healthy adults, while the subjects herein were from a broad age range (18-86 years old) and preferentially enrolled from high-risk groups. Thus, the data herein represent a broad age spectrum and are also particularly relevant for those age groups and demographics most at risk for hospitalization or death due to COVID-19.

This study in no way excludes a complementary role for antibodies in control of SARS2 in unvaccinated individuals. Our study has limitations. It is intrinsically challenging to disentangle roles of T cells from roles of antibodies in combatting SARS2 in humans[19–21]. This study provides strong evidence that SARS2-specific T cell responses exhibit selective correlates with viral clearance, but it is entirely plausible that coordinated adaptive immunity of T cells and antibodies are important in protection[1,18,38]. Additionally, adaptive immune responses at the site of initial infection were not directly assessed in this study and may not be adequately captured by evaluation of circulating immune cells and antibodies from peripheral blood samples. Thus, a role for local, upper airway cellular and humoral immunity in protection from SARS2 cannot be excluded. The study was also underpowered to distinguish whether circulating CD4 and CD8 T cell responses were each independently associated with lower SARS2 NP viral loads. Lastly, our findings focused on primary infection with ancestral SARS2 and may not be reflective of all individuals with acute COVID-19, including SARS2 breakthrough infections.

Separately it has been observed that in COVID-19 vaccinated individuals who experience breakthrough infections, recall responses include rapid activation of memory S-specific CD4 and CD8 T cells within the first week, which precede humoral responses, and that S-specific CD8 T cells are associated accelerated viral clearance[40,41]. While T cell kinetics and phenotypes in response to SARS2 infection and COVID-19 vaccination may not be identical and are shaped by exposure history, the cumulative data now indicate that SARS2-specific T cells appear to play a protective role in both primary SARS2 infection and breakthrough infections. Given these findings, next generation SARS2 or pan-sarbecovirus vaccines may benefit from eliciting enhanced T cell responses, and from clinical trials that assess both baseline and vaccination-induced SARS2-specific CD4 and CD8 T cell responses.

METHODS

Study Population and Trial

ACTIV-2/A5401 is an ongoing, multicenter phase 2/3 randomized controlled trial designed to evaluate the safety and antiviral and clinical efficacy of therapeutics for acute COVID-19 in non-hospitalized adults. Individuals in this study represent a subset of participants from ACTIV-2/A5401 who received a single intravenous dose of 700 mg of bamlanivimab or normal saline placebo (comparator control group) and had available clinical data (e.g., demographics, days post-symptom onset, risk for severe COVID-19; **Fig. 1A** and ref[24,47]) and blood samples for immunologic testing. Samples from nearly half of the overall bamlanivimab 700 mg and comparator placebo groups were available for this study. Participants were from the United States and enrolled in ACTIV-2/A5401 between October-November 2020. At study entry, COVID-19 vaccines were not available to the general population in the United States and SARS2 infections could be attributed to ancestral SARS2 virus. The subjects in this study were comparable at study entry across treatment groups (**Fig. 1A** and ref[24,47]). Inclusion criteria for ACTIV-2/A5401 included adults 18 years or older with no more than 10 days of COVID-19 symptoms and documented SARS2 infection by FDA-authorized antigen or molecular testing within seven days prior to study entry. Subjects were assigned to the bamlanivimab or placebo groups at a 1:1 ratio, and randomization was stratified by time from symptom onset (less than or ≥5 days post-symptom onset) and risk of progression to severe COVID-19 based on age (less than or ≥ 55 years old) and the presence or absence of pre-defined comorbid medical conditions (e.g., body mass index > 35kg/m^2, hypertension, diabetes)[47]. Additional information for ACTIV-2/A5401 is available at ClinicalTrials.gov (Identifier: NCT04518410) and in the primary outcomes manuscript[47]. Three individuals were noted to have high NP viral loads despite high nAb titers at study day 0 (**Fig. 5D**). Notably, one of these three individuals was in the placebo group. It was subsequently re-confirmed that blood was drawn for all participants prior to bamlanivimab or placebo infusion, and that none of the placebo group participants received bamlanivimab or any other active drug treatment for COVID-19.

Partial data for day 0 RBD IgG titers by group, similar to the data shown in **Fig. S4C** but with smaller sample sizes and without using a strict mean fluorescence intensity (MFI)-based cutoff, was previously published[24]. Although combined day 28 AIM and AIM+ICS data was published to show T cell response rates between the treatment and placebo groups[24], it was not presented as longitudinal data (as shown in Fig. **4A-C**, **S4A**). Day 0 and 7 T cell data have not been previously published.

SARS2 Viral RNA

Samples were collected, processed, and analyzed as previously reported[47]. In brief, nasopharyngeal swab samples were collected by ACTIV-2/A5401 trial staff at designated study sites on assigned study days (days 0, 7, 14, and 28) using standardized swabs and validated collection and storage procedures. nasopharyngeal swabs were frozen upon collection and stored at −80 °C (−65 °C to −95 °C) until shipped on dry ice to a central laboratory (University of Washington) for extraction, amplification and quantitative detection testing using validated, previously published methods using the Abbott m2000sp/rt system with a validated controls and standards for correlation with cycle threshold (Ct) and viral load. The limit of detection (LOD) was 1.4 log10 copies/mL, lower limit of quantification (LLOQ) was 2 log10 copies/mL, and upper limit of quantification (ULOQ) was 7 log10 copies/mL for this assay. For samples with viral RNA levels above the ULOQ, samples were diluted, and the assay repeated to obtain a quantitative value.

PBMC and viability-based quality control

Peripheral blood was collected by ACTIV-2/A5401 trial staff at designated study sites on assigned study days. Serum and peripheral blood mononuclear cells (PBMC) for immunologic testing were isolated from whole blood using standard operating procedures. PBMC were cryopreserved and stored in liquid nitrogen until ready for use, then thawed at 37°C, resuspended in warm complete RPMI medium with 5% human AB serum (Gemini Bioproducts) and benzonase and centrifuged to remove cryopreservation medium. Following washing, cell counts were performed, and viability assessed using the Muse Count & Viability Kit (Muse Cell Analyzer; Luminex). A PBMC viability threshold was set at ≥ 75% for all samples for inclusion in data analysis. PBMC with viability below this threshold failed to appropriately respond to control stimuli. Four of 95 PBMC samples failed to meet the viability cutoff and were excluded from analyses. PBMC were resuspended to achieve a final concentration of 100,000 PBMC/100 µL for plating in 96-well format for T cell assays. Cryopreserved convalescent COVID-19 donor PBMC from known positive responders obtained from the Sette lab under an LJI IRB-approved protocol (VD-214) served as batch controls across T cell assays. These PBMC were handled in the same fashion as ACTIV-2/A5401 PBMC samples. Additional quality control metrics were also applied, as described below for the individual T cell assays.

Activation induced marker assay (AIM)

As previously described[24], PBMC were cultured in 96-well U-bottom plates for 24 hours at 37°C in an incubator with 5% CO_2 in the presence of a single stimulus per well: a negative control (equimolar amount of DMSO vehicle), a positive control (Staphylococcal enterotoxin B at 1 µg/mL), or a single SARS2 MP (1 µg/mL per MP) [23,31,54] containing Spike (S)[23,54] or paired SARS2 non-Spike epitope MPs (CD4-RE or CD8-RE dominant and subdominant MPs[31]). The Spike MP was comprised of overlapping peptides spanning the full-length ancestral Spike protein. The CD4-RE pools ("non-Spike") consisted of experimentally validated and optimized class II epitopes from the remainder of the SARS2 proteome (outside of the Spike open reading frame (ORF)). CD8 AIM responses were calculated using the AIM+ICS assay as PBMC counts were limited and did not allow for CD8-RE MP wells to be plated for both AIM and AIM+ICS assays (see AIM+ICS assay section below for additional experimental details).

PBMC were plated at 1 x 10^6 PBMC per MP stimulation well and between 0.5-1 x 10^6 PBMC per control well; negative controls were plated in duplicate. Prior to stimulation, PBMC were incubated at 37°C for 15 minutes with 0.5 µg/mL anti-human CD40 blocking antibody (Miltenyi Biotec) per well. Chemokine receptor antibodies were also added to each well on day 1 (see ref[24] for antibodies used in the AIM assay). After a 24-hour incubation, the plates were centrifuged, cells washed with PBS, then stained with LIVE/DEAD Fixable Blue (Invitrogen) 1:1000 in PBS with Fc block (5 µL/sample; BD Biosciences (BD)) for 15 minutes at room temperature, washed with FACS buffer (3% FBS in DPBS without calcium or magnesium), surface stained (see ref[24] for surface staining panel; 30 minutes at 4°C), fixed with BD Cytofix Fixation Buffer (4°C for 20 minutes), and stored at 4°C (up to overnight) until flow cytometric analysis was performed using a 5-laser Cytek Aurora (Cytek Biosciences). Flow cytometric data was acquired separately for each sample and stimulation condition; no samples were pooled. Analysis was performed using FlowJo (BD), and AIM+ gates were drawn based on MP-stimulated responses relative to negative (DMSO) and

positive (SEB) controls. T cell assays on two SARS2 convalescent control PBMC samples were run with each set of experimental PBMC samples to account for batch effects; aliquots from the same PBMC controls were used across batches. A single gating strategy was applied to all samples across all batches unless there was a clear batch-specific effect to justify modifying the gates for a specific batch of samples. Our standard acceptable background (% AIM$^+$ T cells) for DMSO gates is 0.1. DMSO gates were set to target average background values ≤ 0.1 using the same flow cytometry gating strategy for all samples across all batches. CD4 and CD8 T cell responses to MP versus DMSO stimulation are shown in **Fig. S2K-M** and **S3G-H**. Higher background was allowed for CD8 AIM (CD69$^+$41BB$^+$) assays (**Fig. S3G**). ICS (CD69$^+$IFNγ$^+$) was preferentially used as the main readout for SARS2-specific CD8 T cell responses given lower background (**Fig. S3H**) with comparable response rates to CD8 AIM.

PBMC quality was evaluated by measuring the median response to SEB for all samples, and samples with responses <50% of the overall median SEB response were excluded from downstream analyses. Two DMSO negative control well replicates were run for each sample and the average of the individual DMSO wells for each sample was calculated. T cell responses were verified by stimulation index (SI) as a quality control. SI was calculated for each sample as the fold change in responses to MP-stimulation versus the average response to DMSO for the same subject (as shown in **Fig. S2N-P** and **Fig. S3I-J**). An SI cutoff of 2 was used for determining AIM$^+$ CD4 T cell responses and 3 for AIM$^+$ CD8 T cell responses. The higher SI cutoff for CD8 T cell responses was applied based on higher background for CD8 AIM assays. When a non-zero DMSO value was needed for calculating the stimulation index, the minimum DMSO signal was set to 0.005%. For each MP-stimulated sample, the average DMSO value was subtracted from the MP response to calculate the background subtracted MP response. Background subtracted S and non-S values are shown for all figures. Combined AIM responses were calculated as the mathematical sum of the background subtracted responses to individual (S plus non-S) MPs. In the case of negative AIM responses to both S and non-S megapools, or summative responses that were less than the lower limit of quantitation (LOQ) for the assay, as described below, the combined AIM response was also considered negative. The geometric mean of all DMSO control wells for all samples for each assay was calculated to determine the LOQ for the assay. Positive responders were defined as those with background subtracted responses greater than the LOQ. The baseline for the y-axis was set to 0.5 x LOQ. All non-responders were set to this baseline.

Hybrid activation induced marker plus intracellular cytokine staining assay (AIM+ICS)

AIM+ICS assays were also performed as previously described[24]. For CD4 T cells, intracellular cytokine expression was measured in conjunction with surface CD40L. For CD8 T cells, intracellular cytokine staining was measured in conjunction with surface CD69. PBMC were thawed and plated in parallel and as AIM assays but with the addition of wells for CD8-RE MP (paired dominant and non-dominant epitopes[31]) stimulation. The CD8-RE pools ("non-Spike") consisted of optimized, experimentally validated class I epitopes from the remainder of the SARS2 proteome (outside of the Spike ORF)[31]. Cells were incubated with anti-CD40 blocking antibody as for AIM assays, but no chemokine receptor antibodies were added on day 1 except for CXCR5 for identification of circulating T$_{FH}$ cells (see ref[24] for antibodies used). After 20-22 hours, PMA (0.05 μg/mL) and ionomycin (0.25 μg/mL) were added to the positive (ICS) control wells. Two hours later, 0.25 μL/well of GolgiStop (BD) and GolgiPlug (BD) and the AIM marker antibodies (see

ref[24]) were added to all samples, and the plates were incubated for another 4 hours at 37°C (in a 5% CO_2 incubator). Cells were then washed, surface stained for 30 minutes at 4°C, fixed and washed using Cytofix/Cytoperm (BD) per the manufacturers' protocol. Intracellular cytokine staining was then performed using antibodies diluted in Perm/Wash Buffer (BD) for 30 minutes at 4°C. Cells were washed with Stain Buffer with FCS (BD) and stored in this buffer at 4°C until flow cytometric analysis was performed using a Cytek Aurora. Flow cytometric data was acquired separately for each sample and stimulation condition.

Gating and quality control evaluations with SEB responses and SI calculations were performed as described above for AIM assays. PMA + ionomycin served as an additional control for cytokine production (ICS). For each MP-stimulated sample, the average DMSO value was subtracted from the MP response to calculate the background subtracted MP response. Background subtracted values are shown for all figures unless indicated. Combined responses were calculated as the sum of the background subtracted responses to individual MPs (S and CD4-RE for CD4 T cells, S and CD8-RE for CD8 T cells). In the case of negative ICS responses to both S and non-S MPs, the combined response was also considered negative. The LOQ and baseline was calculated and plotted as described above for AIM assays.

Binding antibody titers

Serum SARS2-specific binding antibody assays were performed using the Bio-Plex Pro Human SARS2 IgG 4-Plex Panel serology assay (Bio-Rad 12014634) according to the manufacturer's protocol, as previously described[24]. MFI-based cutoffs for seropositivity are indicated in the figures.

Pseudovirus neutralizing antibody titers

Half-maximal neutralizing antibody titers (NT50) were generated using a lentiviral pseudovirus assay, as previously described[55]. Briefly, PSV was generated in HEK293T cells co-transfected with SARS2 Spike expression and pNL4–3.luc.R-E-mCherry-luciferase reporter packaging plasmids. Serum was isolated from the peripheral blood of study subjects using standard protocols, frozen and shipped to the testing site without thawing until ready for use. The serum was then serially diluted and incubated with SARS2 PSV, then HEK293T cells stably expressing human angiotensin converting enzyme 2. Serum dilutions were run in triplicate. The percent inhibition was calculated using relative light units emitted by the reporter at each dilution and plotted versus the log serum dilution. The NT50 was then derived by non-linear regression. The limit of detection for the assay was defined by the lowest dilution tested (1:3) as indicated in figures.

Statistics

Statistical analyses were performed in GraphPad Prism 10 (GraphPad Software) and Microsoft Excel. Comparisons of SARS2-specific T cell response magnitudes and antibody titers between treatment and placebo groups for equivalent conditions were determined by Mann-Whitney tests. Fisher's exact tests were used to compare participant characteristics (**Fig. 1A**) and to compare SARS2-specific T cell positive and negative response rates between treatment and control groups. Kruskal-Wallis nonparametric tests with post-hoc Dunn's multiple comparison tests were used for assessing T cell responses across more than one stimulation condition or cell type. Relationships between SARS2 nasopharyngeal RNA levels and immune responses were assessed using nonparametric Spearman correlations.

Individual variation in days from symptom onset to study entry (day 0) was accounted for using a log-linear regression model (**Fig. S6A-E**). Adjusted SARS2 NP RNA levels (**Fig. S6A**) and adaptive immune responses (CD4 and CD8 T cell responses, nAb titers; **Fig. S6B-E**) at 6 days post-symptom onset (the median time from symptom onset at study entry) were calculated for all participants. For the linear regression model shown in **Fig. S6A**, study day 0 NP RNA values from five individuals were excluded as outliers as these individuals had undetectable viral RNA but were 4-5 days PSO on study day 0. This included four values from individuals who consistently tested negative for viral RNA (from both nasal and NP swabs) on all study days, and a value for one individual who tested negative for viral NP RNA on study day 0 but who subsequently tested positive for viral NP RNA on study days 3, 7, and 14. The adjusted NP viral RNA and adaptive immunity data, calculated for day 6 post-symptom onset, was then analyzed using nonparametric Spearman correlations, including immune response data from all 95 subjects (**Fig. S6F-I**). One-way ANCOVA were performed to control for the presence/absence of nAb on the relationships between SARS2 NP RNA levels and SARS2-specific T cell responses at day 0 (**Fig. S5M-O**).

For all analyses, a two-sided 5% type I error rate was used, without adjustment for multiple comparisons, except as noted above. Additional details can be found in the results, figures and corresponding legends.

Correction in previous post-symptom onset analysis

A previous version of this manuscript (https://doi.org/10.1101/2023.10.25.564014) contained an error in the post-symptom onset analysis. An incorrect sign in the formula to adjust immunologic measurement values led to an increased rather than diminished correlation between immune responses and viral NP RNA levels after adjustment for sample collection times other than 6 days post-COVID-19 symptom onset. The analysis was conducted by B. Peters, who apologizes for identifying this error after the bioRxiv submission. This note is included for readers who accessed the previous version of the manuscript to explain differences in the figures and correlation coefficients since the prior bioRxiv submission.

Study approval

The ACTIV-2/A5401 clinical trial protocol (ClinicalTrials.gov Identifier: NCT04518410) was approved by a central institutional review board (IRB); Advarra (Pro00045266). The La Jolla Institute for Immunology (LJI) IRB provided additional oversight for this study. All individuals enrolled in ACTIV-2/A5401 provided written informed consent for participation.

AUTHOR CONTRIBUTIONS

Conceptualization, S.I.R., D.M.S., and S.C.; Investigation, S.I.R., P.G.L., F.F., A.H.; Formal Analysis, S.I.R., B.P, S.C.; Patient Recruitment and Samples D.M.S.; Material Resources, A.G., D.W., P.K., A.S.; Writing, S.I.R., F.F., D.M.S. and S.C.; Supervision D.M.S. and S.C.

FUNDING

This work was supported in part by the National Institute of Allergy and Infectious diseases (NIAID) of the National Institutes of Health (NIH), Department of Health and Human Services, under contract no.

75N93019C00065 (A.S, D.W.), under awards AI036214, AI131385, UM1AI068634, UM1AI068636, and UM1AI106701. Additional support was provided in part by the John and Mary Tu Foundation, NIH T32 AI007036 (S.I.R.) and an A.P. Giannini Foundation fellowship award (S.I.R.). Bamlanivimab was donated by Eli Lilly and Company. The content presented is the sole responsibility of the authors and may not represent the opinions of the NIH.

ACKNOWLEDGMENTS

We would like to thank the study participants, site staff and investigators, members of the study team and community advisory board, the AIDS Clinical Trials Group (ACTG), the Harvard Center for Biostatistics in AIDS Research (CBAR) and ACTG Statistical and Data Analysis Center (SDAC), the National Institute of Allergy and Infectious Diseases (NIAID) / Division of AIDS (DAIDS), Bill Erhardt, the Accelerating COVID-19 Therapeutic Interventions and Vaccines (ACTIV) partnership, and PPD for making this study possible. See Supplemental Acknowledgments for additional details.

DATA AVAILABILITY

The authors confirm that the source data underlying the findings are fully available. Due to ethical restrictions, additional ACTIV-2/A5401 clinical trial study data beyond what is presented in this manuscript and supplement are available upon request from the AIDS Clinical Trials Group (ACTG) and with the written agreement of ACTG and the manufacturer of the investigational product. Completion of a data use agreement may be required.

COMPETING INTERESTS

S.I.R. has no competing interests to declare. J.S.C. has consulted for Merck and Company. A.L.G. reports contract testing from Abbott, Cepheid, Novavax, Pfizer, Janssen and Hologic and research support from Gilead, outside of the described work. P.K. is an employee and shareholder of Eli Lilly. D.W. is a consultant for Moderna. A.S. is a consultant for AstraZeneca Pharmaceuticals, Calyptus Pharmaceuticals, Inc, Darwin Health, EmerVax, EUROIMMUN, F. Hoffman-La Roche Ltd, Fortress Biotech, Gilead Sciences, Gritstone Oncology, Guggenheim Securities, Moderna, Pfizer, RiverVest Venture Partners, and Turnstone Biologics. K.W.C. has received research funding to the institution from Merck Sharp & Dohme. D.M.S. has consulted for and has equity stake in Linear Therapies, Model Medicines and Vx Biosciences and consulted for Bayer, Kiadis, Signant Health and Brio Clinical. S.C. has consulted for GSK, JP Morgan, Citi, Morgan Stanley, Avalia NZ, Nutcracker Therapeutics, University of California, California State Universities, United Airlines, Adagio, and Roche. LJI has filed for patent protection for various aspects of T cell epitope and vaccine design work. The other authors have no competing interests to declare.

REFERENCES

1. Goldblatt, D., Alter, G., Crotty, S. & Plotkin, S. A. Correlates of protection against SARS-CoV-2 infection and COVID-19 disease. *Immunol Rev* **310**, 6–26 (2022).

2. Centers for Disease Control and Prevention. COVID Data Tracker. Atlanta, GA: U.S. Department of Health and Human Services, CDC; 2024, January 17. https://covid.cdc.gov/covid-data-tracker.

3. Du, Z. *et al.* Estimate of COVID-19 Deaths, China, December 2022–February 2023. *Emerg. Infect. Dis.* **29**, (2023).

4. Xiao, H., Wang, Z., Liu, F. & Unger, J. M. Excess All-Cause Mortality in China After Ending the Zero COVID Policy. *JAMA Netw Open* **6**, e2330877 (2023).

5. Gilbert, P. B. *et al.* Immune correlates analysis of the mRNA-1273 COVID-19 vaccine efficacy clinical trial. *Science* **375**, 43–50 (2022).

6. Corbett, K. S. *et al.* Immune correlates of protection by mRNA-1273 vaccine against SARS-CoV-2 in nonhuman primates. *Science* **373**, eabj0299 (2021).

7. Feng, S. *et al.* Correlates of protection against symptomatic and asymptomatic SARS-CoV-2 infection. *Nat Med* **27**, 2032–2040 (2021).

8. Gilbert, P. B. *et al.* A Covid-19 Milestone Attained — A Correlate of Protection for Vaccines. *N Engl J Med* **387**, 2203–2206 (2022).

9. Fong, Y. *et al.* Immune correlates analysis of the PREVENT-19 COVID-19 vaccine efficacy clinical trial. *Nat Commun* **14**, 331 (2023).

10. Fong, Y. *et al.* Immune correlates analysis of the ENSEMBLE single Ad26.COV2.S dose vaccine efficacy clinical trial. *Nat Microbiol* **7**, 1996–2010 (2022).

11. Regev-Yochay, G. *et al.* Correlates of protection against COVID-19 infection and intensity of symptomatic disease in vaccinated individuals exposed to SARS-CoV-2 in households in Israel (ICoFS): a prospective cohort study. *The Lancet Microbe* **4**, e309–e318 (2023).

12. Hertz, T. *et al.* Correlates of protection for booster doses of the SARS-CoV-2 vaccine BNT162b2. *Nat Commun* **14**, 4575 (2023).

13. Stadler, E. *et al.* Monoclonal antibody levels and protection from COVID-19. *Nat Commun* **14**, 4545 (2023).

14. Levin, M. J. *et al.* Intramuscular AZD7442 (Tixagevimab–Cilgavimab) for Prevention of Covid-19. *N Engl J Med* **386**, 2188–2200 (2022).

15. Isa, F. *et al.* Repeat subcutaneous administration of casirivimab and imdevimab in adults is well-tolerated and prevents the occurrence of COVID-19. *International Journal of Infectious Diseases* **122**, 585–592 (2022).

16. O'Brien, M. P. *et al.* Subcutaneous REGEN-COV Antibody Combination to Prevent Covid-19. *N Engl J Med* **385**, 1184–1195 (2021).

17. Wilkinson, T. M. *et al.* Preexisting influenza-specific CD4+ T cells correlate with disease protection against influenza challenge in humans. *Nat Med* **18**, 274–280 (2012).

18. Sette, A. & Crotty, S. Adaptive immunity to SARS-CoV-2 and COVID-19. *Cell* **184**, 861–880 (2021).

19. Follmann, D. *et al.* Examining protective effects of SARS-CoV-2 neutralizing antibodies after vaccination or monoclonal antibody administration. *Nat Commun* **14**, 3605 (2023).

20. Sette, A., Sidney, J. & Crotty, S. T Cell Responses to SARS-CoV-2. *Annu. Rev. Immunol.* **41**, 343–373 (2023).

21. Rydyznski Moderbacher, C. *et al.* Antigen-Specific Adaptive Immunity to SARS-CoV-2 in Acute COVID-19 and Associations with Age and Disease Severity. *Cell* (2020) doi:10.1016/j.cell.2020.09.038.

22. Dan, J. M. *et al.* Immunological memory to SARS-CoV-2 assessed for up to 8 months after infection. *Science* **371**, (2021).

23. Grifoni, A. *et al.* Targets of T Cell Responses to SARS-CoV-2 Coronavirus in Humans with COVID-19 Disease and Unexposed Individuals. *Cell* **181**, 1489-1501.e15 (2020).

24. Ramirez, S. I. *et al.* Bamlanivimab therapy for acute COVID-19 does not blunt SARS-CoV-2–specific memory T cell responses. *JCI Insight* **7**, e163471 (2022).

25. Weiskopf, D. *et al.* Phenotype and kinetics of SARS-CoV-2–specific T cells in COVID-19 patients with acute respiratory distress syndrome. *Sci. Immunol.* **5**, eabd2071 (2020).

26. Notarbartolo, S. *et al.* Integrated longitudinal immunophenotypic, transcriptional and repertoire analyses delineate immune responses in COVID-19 patients. *Sci Immunol* **6**, eabg5021 (2021).

27. Gao, Y. *et al.* Ancestral SARS-CoV-2-specific T cells cross-recognize the Omicron variant. *Nat Med* **28**, 472–476 (2022).

28. Zhang, Z. *et al.* Humoral and cellular immune memory to four COVID-19 vaccines. *Cell* **185**, 2434-2451.e17 (2022).

29. Mateus, J. *et al.* Low-dose mRNA-1273 COVID-19 vaccine generates durable memory enhanced by cross-reactive T cells. *Science* **374**, eabj9853 (2021).

30. Rydyznski Moderbacher, C. *et al.* NVX-CoV2373 vaccination induces functional SARS-CoV-2-specific CD4+ and CD8+ T cell responses. *J Clin Invest* **132**, e160898 (2022).

31. Yu, E. D. *et al.* Development of a T cell-based immunodiagnostic system to effectively distinguish SARS-CoV-2 infection and COVID-19 vaccination status. *Cell Host & Microbe* **30**, 388-399.e3 (2022).

32. Tarke, A. *et al.* SARS-CoV-2 vaccination induces immunological T cell memory able to cross-recognize variants from Alpha to Omicron. *Cell* **185**, 847-859.e11 (2022).

33. Wherry, E. J. & Barouch, D. H. T cell immunity to COVID-19 vaccines. *Science* **377**, 821–822 (2022).

34. Sahin, U. *et al.* BNT162b2 vaccine induces neutralizing antibodies and poly-specific T cells in humans. *Nature* **595**, 572–577 (2021).

35. Müller, T. R. *et al.* Additive effects of booster mRNA vaccination and SARS-CoV-2 Omicron infection on T cell immunity across immunocompromised states. *Sci Transl Med* **15**, eadg9452 (2023).

36. Tan, A. T. *et al.* Early induction of functional SARS-CoV-2-specific T cells associates with rapid viral clearance and mild disease in COVID-19 patients. *Cell Reports* **34**, 108728 (2021).

37. Swadling, L. *et al.* Pre-existing polymerase-specific T cells expand in abortive seronegative SARS-CoV-2. *Nature* **601**, 110–117 (2022).

38. Rydyznski Moderbacher, C. *et al.* Antigen-Specific Adaptive Immunity to SARS-CoV-2 in Acute COVID-19 and Associations with Age and Disease Severity. *Cell* **183**, 996-1012.e19 (2020).

39. Eser, T. M. *et al.* Nucleocapsid-specific T cell responses associate with control of SARS-CoV-2 in the upper airways before seroconversion. *Nat Commun* **14**, 2952 (2023).

40. Koutsakos, M. *et al.* SARS-CoV-2 breakthrough infection induces rapid memory and de novo T cell responses. *Immunity* **56**, 879-892.e4 (2023).

41. Painter, M. M. *et al.* Prior vaccination promotes early activation of memory T cells and enhances immune responses during SARS-CoV-2 breakthrough infection. *Nat Immunol* **24**, 1711–1724 (2023).

42. Arieta, C. M. *et al.* The T-cell-directed vaccine BNT162b4 encoding conserved non-spike antigens protects animals from severe SARS-CoV-2 infection. *Cell* **186**, 2392-2409.e21 (2023).

43. Tai, W. *et al.* An mRNA-based T-cell-inducing antigen strengthens COVID-19 vaccine against SARS-CoV-2 variants. *Nat Commun* **14**, 2962 (2023).

44. Sette, A. & Saphire, E. O. Inducing broad-based immunity against viruses with pandemic potential. *Immunity* **55**, 738–748 (2022).

45. Kingstad-Bakke, B. *et al.* Vaccine-induced systemic and mucosal T cell immunity to SARS-CoV-2 viral variants. *Proc. Natl. Acad. Sci. U.S.A.* **119**, e2118312119 (2022).

46. Moss, P. The T cell immune response against SARS-CoV-2. *Nat Immunol* **23**, 186–193 (2022).

47. Chew, K. W. *et al.* Antiviral and clinical activity of bamlanivimab in a randomized trial of non-hospitalized adults with COVID-19. *Nat Commun* **13**, 4931 (2022).

48. Puhach, O., Meyer, B. & Eckerle, I. SARS-CoV-2 viral load and shedding kinetics. *Nat Rev Microbiol* (2022) doi:10.1038/s41579-022-00822-w.

49. Killingley, B. *et al.* Safety, tolerability and viral kinetics during SARS-CoV-2 human challenge in young adults. *Nat Med* **28**, 1031–1041 (2022).

50. Chen, P. *et al.* SARS-CoV-2 Neutralizing Antibody LY-CoV555 in Outpatients with Covid-19. *N Engl J Med* **384**, 229–237 (2021).

51. Crotty, S. T Follicular Helper Cell Biology: A Decade of Discovery and Diseases. *Immunity* **50**, 1132–1148 (2019).

52. Lederer, K. *et al.* Germinal center responses to SARS-CoV-2 mRNA vaccines in healthy and immunocompromised individuals. *Cell* **185**, 1008-1024.e15 (2022).

53. Wagstaffe, H. R. *et al.* Mucosal and Systemic Immune Correlates of Viral Control following SARS-CoV-2 Infection Challenge in Seronegative Adults. (2023) doi:10.1101/2023.07.21.23292994.

54. Grifoni, A. *et al.* SARS-CoV-2 human T cell epitopes: Adaptive immune response against COVID-19. *Cell Host and Microbe* **29**, 1076–1092 (2021).

55. Heaps, A. L. *et al.* Rapid determination of SARS-CoV-2 antibody neutralization titer using Bio-Rad Bio-Plex correlates strongly with pseudovirus-determined neutralization titer. *Journal of Virological Methods* **316**, 114726 (2023).

Figure 1. **Study cohort characteristics and SARS2 NP RNA levels during acute COVID-19 and longitudinally.**
A. Demographics and other pertinent characteristics of the matched bamlanivimab (Treatment) and placebo group participants included in this study. Total n = 95 subjects. See methods for additional details. **B.** Median days post-symptom onset (PSO) from start of COVID-19 symptoms to study entry (study day 0) for all participants. **C.** SARS-CoV-2 NP RNA by quantitative reverse transcription polymerase chain reaction (RT-qPCR) for all participants prior to treatment on study day 0. Dotted black line limit of detection (LOD). Dotted red line limit of quantification (LOQ). Values >LOQ were considered positive. Bar = median. **D.** Longitudinal SARS-CoV-2 NP RNA data for the placebo group (n = 49) for days 0, 7, 14, and 28. Median for each time point in red.

Figure 2. Antigen-specific CD4 T cell responses to primary SARS2 infection and acute COVID-19.
A-B. Representative flow cytometry plots and frequency of SARS2-specific CD4 T cells to DMSO (negative control), Spike and CD4-RE MP stimulation conditions (Combined = sum of Spike + CD4-RE responses; see Methods for additional details) by (**A**) AIM using surface OX40 and 41BB co-expression, and (**B**) IFNγ ICS among surface CD40L⁺ cells. **C-E.** Study day 0 SARS2-specific CD4 T cell intracellular cytokine production (**C**) GzmB, (**D**) TNF, or (**E**) IL-2 among surface CD40L⁺ cells. **F.** Parts of a whole donut plot summary of intracellular cytokine (IFNγ, GzmB, TNF, IL-2) production by SARS2-specific CD4 T cells expressing 0 to 4 cytokines. **G-H.** As in **A** and **Fig. S2B** but for SARS2-specific circulating T_FH (cT_FH) cells. Bars = geometric mean. Dotted lines = LOQ. Flow cytometry gates display frequency (%).

Figure 3. Antigen-specific CD8 T cell responses to primary SARS2 infection and acute COVID-19.
A. Representative flow cytometry plots and frequency of SARS2-specific CD8 T cells to DMSO (negative control), Spike and CD8-RE MP stimulation conditions (Combined = sum of Spike + CD8-RE responses; see Methods for additional details) by IFNγ+ ICS among CD69+ cells. **B-D.** (**B**) IFNγ and GzmB, (**C**) TNF, (**D**) IL-2 production. **E.** Parts of a whole donut plot summary of intracellular cytokine (IFNγ, GzmB, TNF, IL-2) production by SARS2-specific CD8 T cells expressing 0 to 4 cytokines. Bars = geometric mean. Dotted lines = LOQ. Flow cytometry gates display frequency (%).

Figure 4. Longitudinal SARS2-specific T cell responses and Ab responses to primary SARS2 infection and acute COVID-19.
A-C. Longitudinal combined (Spike plus non-Spike) (**A**) CD4 AIM (**B**) CD4 AIM+ICS, (**C**) CD8 IFNγ+ T cell responses in placebo group (n = 49) participants at study days 0, 7, and 28. Red dots and lines represent median. Dotted line = LOQ. **D.** Study day 0 pre-treatment nAb titers for all participants. Dotted black line indicates LOD; seropositivity defined by values >LOD. Bar is median. **E.** Study day 0 pre-treatment RBD IgG binding titers for all participants. Dotted black line indicates cut off for seropositivity. Bar is geometric mean titer (GMT).

Figure 5. Correlative relationships between SARS2-specific adaptive immunity and upper airway viral RNA in acute COVID-19.
A-D. Relationships between study day 0 SARS2 NP RNA and combined day 0 SARS2-specific T cell responses by (**A**) CD4 AIM+, (**B**) CD4 IFNγ, (**C**) CD8 IFNγ+ and (**D**) nAb titers.