Title: Recombinant chimeric Horsepox Virus (TNX-801) is attenuated relative to Vaccinia Virus Strains in Human Primary Cell Lines and in Immunocompromised Mice

Stephanie V Trefry[1], Christy N Raney[1], Amy L Cregger[1], Chase A Gonzales[1], Brittney L Layton[1], Robert N Enamorado[1], Nelson A Martinez[1], Deborah S Gohegan[1], Tinoush Moulaei[1], Natasza E Ziółkowska[1], Scott J Goebel[1], Seth Lederman[1], Sina Bavari[1], Farooq Nasar[1*]

[1]*Tonix Pharmaceuticals, Frederick, MD 21701, USA*

*Corresponding author:

Farooq Nasar

Mailing address: Tonix Pharmaceuticals, Frederick, MD 21701

Phone: (240)-741-2974.

Email: farooq.nasar@tonixpharma.com, fanasar@icloud.com

Abstract: Recombinant chimeric Horsepox virus (TNX-801) is a preclinical vaccine in development against Monkeypox and smallpox. In this brief report, we investigated the potential phenotypic differences in *in vitro* and *in vivo* models between TNX-801 and older VACV-based vaccine strains (VACV-IHD, VACV-Lis, VACV-NYC) used in the eradication of smallpox virus. TNX-801 displayed a small plaque phenotype (~1-2 mm) in both BSC-40 and Vero-E6 cells and yielded >10- to 100-fold lower infectious titers than the VACV strains in multiple-step replication kinetics. Growth kinetics in primary human cell lines from two main routes of poxvirus transmission, respiratory and dermal tracts, yielded ~10- to 119-fold lower infectious titers of TNX-801. Intranasal infection of immunocompromised mice (C56BL/6 $Ifnar^{-/-}/Ifngr^{-/-}$) with VACV strains at 6.0 and 5.0 $\log_{10}$ PFU produced uniform lethal disease. In contrast, TNX-801 at 8.0 $\log_{10}$ PFU was unable to produce any clinical disease in mice. These data demonstrate that TNX-801 is >10- to 1,000-fold more attenuated than older VACV-based smallpox vaccines in human primary cell lines and immunocompromised mice.

Introduction.

Horsepox virus (HPXV) is a member of the genus *Orthopoxvirus* in the family

Poxviridae. The genome is comprised of double-stranded DNA ~212 kb in length [1].

Little is known about the biology of HPXV; however, the genus is also home to four

prominent members, Variola, monkeypox, cowpox, and vaccinia viruses, which are

important in human health. Variola, the causative agent of smallpox disease, and

monkeypox viruses are important human pathogens that can cause fatal infection. Case

fatality rates with smallpox and monkeypox infection range from ~30% to ~10%,

respectively [1]. Following a world-wide vaccination campaign, smallpox was eradicated

in 1980. The smallpox vaccine was the product of Edward Jenner's seminal observation

in milkmaids who contracted a mild disease from the cows and in turn were protected

from lethal smallpox. For 130 years following his observation, the virus in the

vaccination efforts was thought to be of Cowpox origin [1-3]. However, in 1939, it was

shown that the virus in the vaccine was comprised of Vaccinia Virus (VACV), a virus

serological related but distinct from Cowpox. The origin of VACV, its natural cycle, and

how it was introduced remain unknown. Recent genomic studies of American Civil War

Era smallpox vaccine scab kits showed the presence of both VACV and HPXV [4]. One

of the isolates displayed a 99.7% nucleotide identity to HPXV [4]. These data strongly

support Jenner's hypothesis and suggest that HPXV was used as a human vaccine.

Utilizing modern genetic techniques, we generated an infectious clone of Horsepox

virus (TNX-801) and are investigating it as a next generation vaccine against smallpox

and monkeypox disease [5]. One critical hurdle in the development of any live-

attenuated vaccine platform is ensuring safety in relevant model systems. Preliminary

studies demonstrated that TNX-801 readily grew in mammalian cells, however, it

displayed an attenuated phenotype both *in vitro* and *in vivo* models [5]. In this study, we

aimed to extend the previous findings by investigating the attenuation of TNX-801 in

immortalized non-human primate (NHP) cell lines, primary human cells, and in an

immunocompromised mouse model relative to the older Vaccinia virus-based smallpox

vaccines (VACV-IHD, VACV-Lister, and VACV-NYC) utilized to eradicate smallpox.

Results.

TNX-801 plaque phenotype. The plaque phenotype of TNX-801 was compared

relative to VACV-WR, VACV-IHD, VACV-Lis, VACV-NYC, and MVA on immortalized

non-human primate cell lines, BSC-40 and Vero-E6 cells (Figs. 1 and 2). In both cell

lines, VACV strains displayed large plaque phenotype plaques ~3-4 mm in diameter by

4 (BSC-40) and 6 dpi (Vero-E6). In contrast, TNX-801 and MVA displayed small plaque

phenotype with ~1-2 mm and ~0.5 mm in diameter on both cell lines, respectively.

Multiple-step replication kinetics. The replication kinetics of TNX-801, VACV-WR,

VACV-IHD, VACV-Lis, VACV-NYC, and MVA was investigated in BSC-40 and Vero-E6

cells at a multiplicity of infection (MOI) of 0.01 (Figure 3A and B). All VACV strains

reached peak titers of ~8.0 $\log_{10}$ PFU/mL between 48 and 72 hpi. In contrast, TNX-801

and MVA displayed delayed replication kinetics reaching peak titers between 72 and 96

hpi. The peak titer of TNX-801 was ~7.7 and ~6.6 $\log_{10}$ PFU/mL in BSC-40 and Vero-E6

cells, respectively. The TNX-801 titers were up to ~10- to 119-fold lower than VACV

strains timepoints. MVA infection yielded peak titers of ~5.6 and ~4.8 $\log_{10}$ PFU/mL in

BSC-40 and Vero-E6 cells, respectively.

Multiple-step replication studies were also performed in human primary cell lines

representing two main routes for poxvirus transmission from the dermal (melanocytes,

keratinocytes, dermal fibroblasts, and skeletal muscle) and respiratory

(bronchial/tracheal epithelial, small airway epithelial, lung fibroblasts, and

bronchial/tracheal smooth muscle) tracts. Cells were infected at an MOI of 0.01 and

samples were collected at 24-hour (hrs) intervals (Fig. 4A). The TNX-801 infection of

primary melanocytes, keratinocytes, and fibroblasts displayed delayed replication

kinetics. The peak infectious titer of TNX-801 and VACV strains ranged from ~3.8 to 5.9

in $\log_{10}$ PFU/mL and ~4.5 and 7.5 $\log_{10}$ PFU/mL, respectively. The infectious titers of

TNX-801 were ~5- to 112-fold lower in all three primary cell lines. In contrast, TNX-801

and VACV strains yielded similar infection kinetics and infectious titers in skeletal

muscle cells with peak titers of ~7.1 and ~7.7 $\log_{10}$ PFU/mL. The MVA infection did not

produce any infectious virus in any dermal tract primary cells throughout the 96-hr

sampling period.

Infection of primary cells from the respiratory tract yielded similar results. The

TNX-801 infection of fibroblasts, bronchial tracheal epithelial and small airway epithelial

cells produced peak infectious titers ranging from ~5.6 to 6.0 $\log_{10}$ PFU/mL (Fig 4B).

VACV strains yielded peak infectious titers of ~6.0 to 7.2 $\log_{10}$ PFU/mL. TNX-801 titers

were ~5- to 28-fold lower than VACV strains. In contrast, TNX-801 and VACV strains

displayed similar infection kinetics and infectious titers in bronchial tracheal smooth

muscle cells with peak titers of ~7.0 to 7.5 $\log_{10}$ PFU/mL. Lastly, infection with MVA did

not produce any infectious virus in any respiratory tract primary cells throughout the 96

hr sampling period.

Intranasal infection studies in C56BL/6 *Ifnar$^{-/-}$/Ifngr$^{-/-}$* mice. To further investigate phenotypic differences between TNX-801 and VACV strains, susceptibility to virus infection was determined in C56BL/6 *Ifnar$^{-/-}$/Ifngr$^{-/-}$* mice via the intranasal route (Fig 5A). Cohorts of 10 mice/group were infected at 6.0 and 5.0 log$_{10}$ PFU/mouse with VACV-WR, VACV-IHD, VACV-Lis, and VACV-NYC. Mice were also infected with TNX-801 at 8.0 log$_{10}$ PFU/mouse or mock infected with phosphate buffered saline (PBS) (Fig 5A).

Following infection with VACV-WR, VACV-IHD, VACV-Lis, and VACV-NYC, VACV-WR and VACV-IHD infected mice exhibited accelerated clinical disease. All mice exhibited clinical disease by 3 days post-infection (dpi) with peak scores of 2.3 and 2.0 for VACV-WR and VACV-IHD, respectively (Fig 5B). Body temperature began to decline between 1 to 2 dpi then accelerated with average peak decline ranging from -11°F to -14°F at 3 to 4 dpi (Fig 5C). Similar kinetics were observed for weight loss with peak average weight loss of ~17% (Fig 5D). All mice met the euthanasia criteria between 4 and 5 dpi regardless of dose (Fig 5E). Mice infected with VACV-Lis and VACV-NYC exhibited delayed disease kinetics. The onset of clinical disease was between 5 and 6 dpi, and the peak average scores ranged from 2.2 to 3 (Fig 5B). Body temperature declined between 5 and 6 dpi with peak average decline ranging from -2°F to -12°F (Fig 5C). Similarly, weight loss began to decline at 5 dpi and the peak average weight loss ranged from ~22% to 32% (Fig 5D). All mice met the euthanasia criteria between 5 and 10 dpi (Fig 5E). In contrast to all VACV infected mice, TNX-801 and mock infected mice did not exhibit any clinical disease or change in body temperature and weight loss, and all mice survived the study (*p<0.0001*).

Discussion. Recombinant chimeric HPXV (TNX-801) vaccine is a preclinical stage

vaccine against monkeypox and smallpox infection. One critical hurdle in the

development of replication competent vaccines is demonstration of vaccine safety in

relevant *in vitro* and *in vivo* models. In this study, we investigated the potential

attenuation of TNX-801 relative to VACV-based smallpox vaccines used to eradicate

smallpox. In immortalized non-human primate, TNX-801 displayed a small plaque

phenotype, delayed replication kinetics, and >10- to 100-fold lower infectious titers.

Human primary cells from dermal and respiratory tracts also showed similar levels of

attenuation. Lastly, in the infection of immunocompromised mice, TNX-801 was unable

to cause any clinical disease, alteration in body temperature and weight, and all mice

survived the study period. Taken together, these data demonstrate that TNX-801 is 10-

to 1,000-fold more attenuated than older Vaccinia virus-based vaccines used in

smallpox eradication.

TNX-801 displayed an attenuated phenotype in majority of the human primary

cells from two main poxvirus transmission routes, dermal and respiratory tracts.

However, in skeletal muscle and bronchial tracheal smooth muscle cells TNX-801

displayed comparable fitness to VACV. These data suggest that the TNX-801

attenuation is more prominent in cells that would serve as initial site of replication during

virus infection and will limit the spread of TNX-801 by either dermal or respiratory route.

The lack of clinical disease in mice supports this hypothesis.

Surprisingly, TNX-801 displayed comparable replication kinetic and infectious

titers in primary human skeletal muscle cells. This suggest that the intramuscular route

may provide superior immunogenicity following vaccination with TNX-801. Vaccination

studies in rabbits and NHPs with TNX-801 has focused on scarification route and have

yielded robust humoral responses [6]. Future studies will investigate multiple

vaccination routes and determine the optimal route of vaccination in human clinical

trials.

Interferon alpha and gamma play a critical role in host response and defense

against viral pathogens. The disruption of either or both pathways enable susceptibility

in mice to virus infection. We investigated an immunocompromised mouse model

comprising of interferon alpha and gamma receptor knockout to enable increased

susceptibility to poxviruses. The mouse model was extremely susceptible to infection

with VACV strains with all mice rapidly succumbing to virus infection at both 6.0 and 5.0

log_{10} PFU doses. In contrast, mice infected with TNX-801 at a 1,000-fold higher dose

were unable to exhibit any clinical disease. These data strongly suggests that the

vaccine platform is significantly more attenuated than VACV strains. Studies are

underway to determine the lethal dose (LD_{50}) of all VACV strains. In addition, future

studies will explore virus dissemination and virus induced pathology.

The central core of poxvirus genome ~90 kb in length encodes genes

responsible for virus replication and consequently is highly conserved [1, 7]. However,

the left and right arms of the genome are more flexible and encode for host range

restriction and/or host immune antagonism genes. TNX-801 genome is ~20 kb larger

than VACV genome, and yet it displays an attenuated phenotype than VACV [1, 7]. The

gene/s that underlie this phenotype are under investigation and future studies will aim to

determine their effect on various stages of poxvirus replication cycle.

In conclusion, the potential phenotypic difference of TNX-801 and VACV strains

was investigated in *in vitro* and *in vivo* models. TNX-801 displayed an attenuated

phenotype relative to VACV with a small plaque morphology and >~10- to 100-fold

lower infectious titers in immortalized and primary cell models. In immunocompromised

mice, TNX-801 was >1,000-fold more attenuated than VACV-based vaccines utilized for

the eradication of smallpox.

Materials and Methods.

Cells. Vero (CRL-1586) and BSC-40 (CRL-2761) cells were obtained from American

Tissue Culture Collection (ATCC, Manassas, VA). Cells were cultured in Dulbecco's

MEM (DMEM) supplemented with 10% fetal bovine serum (FBS) and gentamicin (50

µg/mL) at 37°C and 5% CO_2.

Normal adult human primary cells were obtained from American Tissue Culture

Collection (ATCC, Manassas, VA). Cells selected were representative of two

transmission routes, respiratory or dermal tracts. Primary dermal tracts cells comprised

of Epidermal Melanocytes (HEMa, PCS-200-013), Epidermal Keratinocytes (HEKa,

PCS-200-011), Dermal Fibroblasts (HDFa, PCS-201-012) and Skeletal Muscle

(HSkMC, PCS-950-010). Primary respiratory tract cells were Bronchial/Tracheal

Epithelial (PCS-300-010), Small Airway Epithelial (HSAEC, PCS-301-010), Lung

Fibroblasts (HLF, PCS-201-013), and Bronchial/Tracheal Smooth Muscle (PCS-130-

011). Cells were cultured in their respective basal mediums and growth kit supplements

per manufacturer recommendations. All primary cells were cultured at 37°C and 5%

CO_2. Cells were seeded overnight to achieve appropriate confluence for the multi-step

growth kinetics.

Viruses and Virus Amplification. Vaccina virus strains [Lister (Lis), International

Health Department (IHD), Western Reserve (WR), New York City (NYC), and Modified

Vaccinia Ankara (MVA) were obtained from BEI Resources (Manassas, VA). HPXV

(TNX-801) stock rescued as described previously. Virus stocks of each strain were

generated on BSC-40 cells. Briefly, cells were seeded overnight in flasks to achieve

70% confluence. Growth media was removed, and monolayers were infected at an MOI

of 0.01. Following an hour incubation at 37°C and 5% CO_2, 20 mL growth media was

added. Monolayers were harvested via cell scraping 72 hpi. Samples were subjected to

three cycles of freeze/thaw and sonication and were stored at -80°C. All virus stocks

were titrated by plaque assay on BSC-40 cells.

For mouse studies, all virus stocks were sucrose purified. Briefly, SW-28 tubes

were filled with 16 mL of 36% sucrose (w/v) in 10mM Tris pH 7.5 followed by overlay of

16 mL of virus isolates. Tubes were spun at 4°C at 33,000 x g for 80 mins. Following

centrifugation, all supernatants were discarded, and the virus pellet was resuspended in

5 mL of 1X PBS. Sucrose purified virus was titrated on BSC-40 cells.

Virus Titration. BSC-40 cells were seeded overnight in 6-well plates to achieve 95-

100% confluence. Each virus stock was serially diluted 10-fold in 1X DMEM.

Monolayers were infected with 100 µL of each dilution and incubated at 37°C and 5%

CO_2 for one hour. Following incubation, monolayers were overlaid with 2 mL of 0.5%

methylcellulose in MEM, GlutaMAX™ (Gibco) supplemented with 5% FBS and

gentamicin (50 µg/mL). Following 72 hours at 37°C and 5% CO_2, plates were fixed in

10% formalin overnight. Plates were stained with 2% crystal violet in 70% ethanol for 5-

10 minutes. Stain was removed with water, plates were dried, and plaques were

counted. For plaque phenotype comparison, the identical protocol was performed on

BSC-40 and Vero-E6 cells and representative 6-wells are shown in figures.

Multi-Step Growth Kinetics. BSC-40, Vero-E6, and human primary cells were seeded

overnight in 6-well plates to achieve 50% confluence. Growth media was removed, and

monolayers were infected at an MOI of 0.01 with TNX-801, VACV-WR, VACV-IHD,

VACV-Lis, VACV-NYC, and MVA. Following an hour incubation at 37°C and 5% CO_2,

monolayers were washed with 10 mL PBS, and 2 mL of growth media was added.

Monolayers were harvested via cell scraping at 0, 12- (BSC-40 and Vero-E6 cells only),

24-, 48-, 72-, and 96-hpi in triplicate. Samples were subjected to three cycles of

freeze/thaw and sonication, and were stored at -80°C. All samples were titrated on

BSC-40 cells.

Ethics statement. This work was supported by an approved Tonix Pharma Animal

Care and Use Committee (IACUC) animal research protocol. Research conducted

under an IACUC approved protocol and was in compliance with the Animal Welfare Act,

PHS Policy, and other Federal statutes and regulations relating to animals and

experiments involving animals. The facility where this research was conducted is

accredited by the Association for Assessment and Accreditation of Laboratory Animal

Care (AAALAC International) and adheres to principles stated in the Guide for the Care

and Use of Laboratory Animals, National Research Council, 2011 [8].

Mouse Studies. Breeding pairs of C57BL/6 *ifnar* [-/-]/*ifngr* [-/-] were purchased from

Jackson Laboratories to generate mouse colonies at Tonix Pharma. A ten-day study

was performed to investigate virulence of TNX-801, VACV-WR, VACV-IHD, VACV-Lis,

and VACV-NYC. Cohorts of ten (5 male and 5 female) four-week-old mice were infected

via intranasal route. Mice were infected with 50 µL (25 µL/nare) of virus. Mock infected

mice received 50 µL (25 µL/nare) of 1X PBS. C57BL/6 *ifnar* $^{-/-}$/*ifngr* $^{-/-}$ mice were

infected with VACV-WR, VACV-IHD, VACV-Lis, and VACV-NYC at 6.0, and 5.0 $\log_{10}$

PFU/mouse. Mice were infected with TNX-801 at 8.0 $\log_{10}$ PFU/mouse. Following

infection, mice were monitored daily for clinical disease, body temperature, weight loss,

and survival. The clinical disease score criteria: 0 = normal, 1 = rough coat and/or

hunched posture, 20% body weight loss from baseline, 2 = mild lethargy and/or mild

dyspnea, 25% body weight loss from baseline, and 3 = moderate lethargy and/or

moderate dyspnea and/or 30% body weight loss from baseline. All target doses were

verified via plaque assay and actual dose is shown in respective figures.

Statistical Analysis. GraphPad Prism version 9 for Windows (GraphPad Software, La

Jolla, California, USA) software was utilized for statistical analysis. Significant

differences in each parameter were determined using one-way ANOVA followed by

multiple comparisons between groups. Significance between TNX-801 and VACV

strains were reported in the graphs.

Disclaimer: TNX-801 vaccine is not approved for any indication.

Conflicts of Interest: S.L. is a co-inventor of the TNX-801 vaccine described in this
study.

References:

1. Moss B and Smith GL. 2020. The Viruses and Their Replication. Howley PM, Knipe DM, and Whelan SPJ. Fields virology, 7th ed. Lippincott Williams & Wilkins, Philadelphia, PA.

2. Fenner F, Henderson A, Arita I, Jezek Z, Ladnyi ID. Smallpox and its eradication. Geneva: World Health Organization; 1988. 1460 p.

3. Baxby D. Jenner's smallpox vaccine: the riddle of vaccinia virus and its origin. London: Heinemann Educational; 1981.

4. Brinkmann A, Souza ARV, Esparza J, Nitsche A, Damaso CR. Re-assembly of nineteenth-century smallpox vaccine genomes reveals the contemporaneous use of horsepox and horsepox-related viruses in the USA. Genome Biol. 2020 Dec 4;21(1):286.

5. Noyce RS, Lederman S, Evans DH (2018) Construction of an infectious horsepox virus vaccine from chemically synthesized DNA fragments. PLoS ONE 13(1): e0188453.

6. Noyce RS, Westfall LW, Fogarty S, Gilbert K, Mpanju O, Stillwell H, Esparza J, Daugherty B, Koide F, Evans DH, Lederman S. Single Dose of Recombinant Chimeric Horsepox Virus (TNX-801) Vaccination Protects Macaques from Lethal Monkeypox Challenge. Viruses. 2023 Jan 26;15(2):356.

7. Tulman ER, Delhon G, Afonso CL, Lu Z, Zsak L, Sandybaev NT, Kerembekova UZ, Zaitsev VL, Kutish GF, Rock DL. Genome of horsepox virus. J Virol. 2006 Sep;80(18):9244-58.

292   8. National Research Council (U.S.). Committee for the Update of the Guide for the

293   Care and Use of Laboratory Animals., Institute for Laboratory Animal Research

294   (U.S.), National Academies Press (U.S.). Guide for the care and use of

295   laboratory animals. 8th ed. Washington, D.C.: National Academies Press; 2011.

Figure 1. Plaque phenotype of VACV-WR, VACV-IHD, VACV-Lis, VACV-NYC, TNX-801, and MVA was investigated in BSC-40 cells. Representative wells from 6-well plates are shown in each photograph.

Figure 2. Plaque phenotype of VACV-WR, VACV-IHD, VACV-Lis, VACV-NYC, TNX-801, and MVA was investigated in Vero-E6 cells. Representative wells from 6-well plates are shown in each photograph.

316 **Figure 3.** Multi-step growth kinetics of VACV-WR, VACV-IHD, VACV-Lis, VACV-NYC,

317 TNX-801, and MVA in BSC-40 (A) and Vero-E6 (B) cells. The limit of detection is shown

318 with black dashed line.

Figure 4. Multi-step growth kinetics of VACV-WR, VACV-IHD, VACV-Lis, VACV-NYC,

TNX-801, and MVA in primary human cell lines from dermal (A) and respiratory (B)

tracts. The limit of detection is shown with black dashed line.

Figure 5. Infection of C57BL/6 *ifnar $^{-/-}$/ifngr $^{-/-}$* mice with VACV-WR, VACV-IHD, VACV-

Lis, VACV-NYC, TNX-801, and mock treated via intranasal route (A). Mice were

infected with VACV-WR, VACV-IHD, VACV-Lis, and VACV-NYC at 6.0 and 5.0 $\log_{10}$

PFU/mouse. Mice were infected with TNX-801 at 8.0 $\log_{10}$ PFU/mouse. Post-infection,

338   mice were monitored for clinic disease (B), body temperature (C), weight loss (D), and

339   survival (E). *p*-values comparing TNX-801 to VACV strains are provided in each graph.