Human ADA2 deficiency is characterized by the absence of an intracellular hypoglycosylated form of adenosine deaminase 2

Short title: DADA2 cells lack lysosomal LMW-ADA2

Authors:

Lisa Ehlers[1,2,3,4,5] (ORCID: 0000-0001-8737-001X), Marjon Wouters[1] (ORCID: 0000-0002-4675-1665), Anneleen Hombrouck[1] (ORCID: 0009-0003-5236-6063), Bethany Pillay[1] (ORCID: 0000-0001-7964-544X), Selket Delafontaine[1,6] (ORCID: 0000-0002-8985-8155), Giorgia Bucciol[1,6] (ORCID: 0000-0001-5004-0738), Marco Baggio[1,7] (ORCID: 0000-0001-6708-9925), Mariia Dzhus[1] (ORCID: 0000-0002-2077-4295), Emma Francés Rabanal[1], Alexandra Damerau[5,8] (0000-0001-6651-2963), Julia Neugebauer[9] (ORCID: 0000-0002-2576-1012), Frédéric Ebstein[10] (ORCID: 0000-0002-3729-7878), Maarten Jacquemyn[11] (ORCID: 0000-0003-2626-4189), Lien De Somer[12,13] (ORCID: 0000-0002-8488-5090), Rik Schrijvers[14] (ORCID: 0000-0002-4261-6220), Steven Vanderschueren[15] (ORCID: 0000-0002-4003-5465), David Cassiman[16] (ORCID: 0000-0002-6154-0970), Marieluise Kirchner[17] (ORCID: 0000-0002-7049-534X), Philipp Mertins[17] (ORCID: 0000-0002-2245-528X), Mir-Farzin Mashreghi[4,5], Tilmann Kallinich[2,3,4,5] (ORCID: 0000-0003-2404-9397), Dirk Daelemans[7] (ORCID: 0000-0001-7092-1153), Patrizia Agostinis[18] (ORCID: 0000-0003-1314-2115), Leen Moens[1] (ORCID: 0000-0002-5347-6526), Isabelle Meyts[1] (ORCID: 0000-0003-1214-0302)

Affiliations:

[1] Department of Microbiology, Immunology and Transplantation, Laboratory for Inborn Errors of Immunity, KU Leuven, Leuven, Belgium

[2] Department of Pediatric Respiratory Medicine, Immunology and Critical Care Medicine, Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Berlin, Germany

[3] Berlin Institute of Health at Charité – Universitätsmedizin Berlin, Berlin, Germany

[4] German Center for Child and Adolescent Health (DZKJ), partner site Berlin, Berlin, Germany

[5] Deutsches Rheuma-Forschungszentrum, an Institute of the Leibniz Association, Berlin, Germany

[6] Department of Pediatrics, University Hospitals Leuven, KU Leuven, Leuven, Belgium

[7] Laboratory of Computational and Developmental Biology, Berlin Institute for Medical Systems Biology (BIMSB), Max-Delbrück-Centrum for Molecular Medicine in the Helmholtz Association (MDC), Berlin, Germany

[8] Department of Rheumatology and Clinical Immunology, Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Berlin, Germany

[9] Department of Pediatric Gastroenterology, Nephrology and Metabolic Diseases, Center of Chronically Sick Children, Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Berlin, Germany

[10] Nantes University, CHU Nantes, CNRS, INSERM, Nantes, France

[11] KU Leuven, Department of Microbiology, Immunology and Transplantation, Rega Institute for Medical Research, Molecular Genetics and Therapeutics in Virology and Oncology Research Group, Leuven, Belgium

[12] Department of Pediatric Rheumatology, University Hospitals Leuven, Leuven, Belgium

[13] Laboratory of Immunobiology, Department of Microbiology and Immunology, KU Leuven,

Leuven, Belgium

[14] Department of Microbiology, Immunology and Transplantation, Allergy and Clinical immunology

Research group, KU Leuven, Leuven, Belgium

[15] Department of General Internal Medicine, Research Department Microbiology, Immunology, and

Transplantation, Laboratory of Clinical Infectious and Inflammatory Disorders, University Hospitals

Leuven, Leuven, Belgium

[16] Department of Gastroenterology-Hepatology and Metabolic Center, University Hospital Leuven,

Leuven, Belgium

[17] Core Unit Proteomics, Berlin Institute of Health at Charité - Universitätsmedizin Berlin and Max-

Delbrück-Center for Molecular Medicine, Berlin, Germany

[18] Cell Death Research & Therapy (CDRT) Lab, Department of Cellular & Molecular Medicine, Center

for Cancer Biology, VIB-KU Leuven, Leuven, Belgium

Corresponding author:

Isabelle Meyts, Laboratory for Inborn Errors of Immunity and Department of Pediatrics UZ Leuven,

Herestraat 49, 3000 Leuven, Belgium; Tel +32 16 343841; Fax +32 16 343842;

isabelle.meyts@uzleuven.be

Abstract

Human deficiency of adenosine deaminase 2 (DADA2) is an autoinflammatory disease caused by pathogenic variants in *ADA2* that lead to impaired deaminase activity. Recently, a lysosomal function of ADA2 has been proposed but an intracellular form of the protein has not yet been characterized. Here, we analyze protein expression of mutant ADA2 in human monocyte-derived macrophages from 10 DADA2 patients. We identify an intracellular low-molecular-weight (LMW) form of ADA2 that undergoes glycan trimming by α-mannosidases and is absent in DADA2 macrophages. Subcellular fractionation and immunofluorescence microscopy demonstrate that LMW-ADA2 is localized in the lysosomes. By overexpression of 34 *ADA2* variants in HEK293T and U-937 cells, we show that absence of LMW-ADA2 strongly correlates with reduced deaminase activity and predicts variant pathogenicity. In conclusion, we describe a previously unreported intracellular hypoglycosylated form of ADA2 and establish the absence of this LMW-ADA2 as a cellular characteristic of DADA2. Thereby, we introduce a protein correlate of the recently described lysosomal form of ADA2.

Summary

Ehlers et al. demonstrate that mutant ADA2 fails to undergo glycan processing beyond the ER leading to the absence of intracellular low-molecular-weight (LMW) ADA2 in DADA2 patients' macrophages. LMW-ADA2 localizes to the lysosomes and can be generated from extracellular wild-type ADA2.

Introduction

Human deficiency of adenosine deaminase 2 (DADA2) is an inborn error of immunity caused by biallelic deleterious mutations in the *ADA2* gene *(1, 2)*. The disease manifests with a diverse phenotype displaying features of both autoinflammation and immunodeficiency *(3)*. More than 600 missense or predicted loss-of-function variants in *ADA2* are listed in the Genome Aggregation Database (gnomAD v4.1.0) *(4)*. In the Infevers registry, 116 *ADA2* variants are classified as (likely) pathogenic *(5)*. Complete loss-of-function variants are more likely to cause bone marrow failure. Missense variants which account for 80% of pathogenic variants are found in patients with a primarily vasculitis phenotype as well as in those presenting with cytopenias and hypogammaglobulinemia *(6, 7)*. Our group recently demonstrated that specific monoallelic *ADA2* variants can cause ADA2 deficiency through a dominant negative mechanism *(8)*.

How mutations in *ADA2* give rise to this diverse phenotype, even in patients with identical genotypes, is not understood. Inflammatory symptoms of the disease typically respond well to tumor necrosis factor-alpha inhibitors (TNFi) *(9)*. Insufficient resolution of bone marrow failure in response to this treatment however often necessitates hematopoietic stem cell transplantation and disease lethality still is 8% *(10)*. To develop improved treatment options, we require a better understanding of the pathomechanisms underlying the immunological and clinical phenotype of DADA2.

ADA2 is a 59-kD glycoprotein with a signal peptide that mediates secretion of the protein via the endoplasmic reticulum (ER), and exhibits extracellular adenosine deaminase activity as a homodimer *(11)*. The protein has four N-glycosylation sites and insufficient glycosylation leads to ER retention and intracellular aggregate formation of ADA2 transfected into HEK293T cells *(12, 13)*. Impaired protein secretion and absent serum ADA2 enzyme activity have been established as characteristics shared by the majority of pathogenic *ADA2* variants *(14)*. Mechanistic approaches to the pathophysiology of ADA2-deficient cells have therefore been based on impaired extracellular

adenosine deamination driving inflammation *(15, 16)*. This assumption fails to acknowledge the weak

affinity of ADA2 for adenosine and the presence of functional ADA1 in DADA2 patients *(17)*.

Research into alternative functions of the ADA2 protein and DADA2-causing variants is therefore

warranted. Biochemical characterization of pathogenic *ADA2* variants has until now mostly been

performed in a HEK293T overexpression system with supraphysiological intracellular levels of ADA2

while expression of the mutant protein has been shown to be minimal in DADA2 CD14$^+$ monocytes

(14, 18, 19). Disease-associated *ADA2* variants have to date hardly been studied in their physiological

milieu.

In this study, we analyze trafficking and glycosylation of endogenous ADA2 expressed in primary

human macrophages from healthy donors and DADA2 patients. We provide evidence of an

intracellular glycoform of ADA2 that results from glycan trimming by Golgi α-mannosidases and show

that DADA2 macrophages do not express this low-molecular-weight-form of ADA2. Absence of

LMW-ADA2 correlates with reduced ADA2 enzyme activity and serves as a biomarker of variant

pathogenicity.

Results

***In vitro* macrophage differentiation partially restores ADA2 protein expression in DADA2**

monocytes

Of the circulating leukocytes, monocytes express ADA2 most abundantly. We analyzed ADA2 protein

expression and *ADA2* mRNA levels in peripheral blood monocytes of 10 DADA2 patients and three

heterozygous carriers (**Table 1** and **Fig. S1A+B**). CD14$^+$ monocytes from DADA2 patients showed

very low or absent ADA2 protein expression despite normal *ADA2* mRNA levels in patients with

missense mutations which account for 80% of pathogenic variants found in DADA2 patients (**Fig. 1A**

and **Fig. S2A**). Upon macrophage differentiation, we observed an increase in mutant ADA2 protein

expression in the patient cells while ADA2 protein expression was reduced in healthy control (HC)

human monocyte-derived macrophages (HMDM) (**Fig. 1B**). Successful differentiation of DADA2

HMDM was achieved both with granulocyte–macrophage colony-stimulating factor (GM-CSF) and

macrophage colony-stimulating factor (M-CSF) (**Fig. S2B**), which confirms the reported restoration

of macrophage differentiation of monocytes from DADA2 patients under treatment with TNFi *(9)*.

Interestingly, western blotting of pathogenic ADA2 protein variants expressed in DADA2 HMDM

revealed the presence of only a high-molecular-weight (HMW) band of around 60 kD compared to

HC HMDM displaying two bands at 60 and 57 kD, with the low-molecular-weight (LMW) band being

more pronounced (**Fig. 1B** and **Fig. S2C**). This protein expression pattern of pathogenic *ADA2*

variants was confirmed in ADA2$^{-/-}$ U-937 cells transduced with the variants p.G47R, p.G47V and

p.R169Q (**Fig. 1C** and **Fig. S2D**). In order to verify our finding in a larger number of *ADA2* variants,

we analyzed the protein expression of mutant ADA2 by transfection of HEK293T cells.

Overexpression of tagged WT ADA2 and several confirmed pathogenic variants uniformly led to

predominant intracellular expression of HMW-ADA2 (63 kD) (**Fig. 1D**). In line with our findings in

DADA2 HMDM, intracellular LMW-ADA2 was absent after overexpression of mutant ADA2 in

HEK293T cells (**Fig. 1D**). Interestingly, this band was still present – albeit at lower intensity – in the

variants p.N127Q and p.W326R, two variants not reported in DADA2 patients but created for the

study of ADA2 glycosylation and dimerization, respectively *(11, 12)*. The molecular weight of secreted

wild-type (WT) ADA2 in the supernatant of HC HMDM or transfected HEK293T cells corresponded

to the intracellular HMW form – the only form expressed in DADA2 HMDM (**Fig. 1B+D**). We

confirmed the secretion of HMW-ADA2 in the monocytic cell lines U-937 and THP-1 (**Fig. S2E**).

HMDM from DADA2 patient P8 showed residual secretion of HMW-ADA2 with a molecular weight

equivalent to secreted WT ADA2 (**Fig. S2F**). This HMW-form of ADA2 also corresponded to the

ADA2 form detected in human serum (**Fig. 1E**). Collectively, we show that HC HMDM intracellularly

express a form of ADA2 that is lower in molecular weight than the secreted protein. This LMW-form

is absent in DADA2 HMDM which only express the HMW-form intracellularly.

Human monocyte-derived macrophages intracellularly express a hypoglycosylated low-

molecular-weight form of ADA2 that is absent in DADA2 macrophages

Differences in the molecular weight of a protein can be caused by posttranslational protein

modifications and N-glycosylation of ADA2 has previously been confirmed *(11, 13, 18)*. Therefore,

we initially hypothesized that the LMW-form of ADA2 unique to cells expressing WT ADA2

corresponded to glycan-free ADA2. To test this, we performed glycan removal on lysates and

supernatants containing ADA2 protein (**Fig. 2A**). Complete removal of N-glycans by Peptide-N-

Glycosidase F (PNGase F) yielded a difference in molecular weight of 7 kD between HMW-ADA2

and glycan-free ADA2 both overexpressed in HEK293T cells and endogenously expressed in the

monocytic cell lines U-937 and THP-1 as well as HMDM (**Fig. 2B+C**), revealing that LMW-ADA2

was in fact slightly larger than glycan-free ADA2. Importantly, the anti-ADA2 antibody clone

EPR25430-131 (#ab288296; abcam) was superior in detecting LMW-ADA2 but poorly detected

glycan-free ADA2 (**Fig. S3A**). Glycan removal by PNGase F eliminated the differences in molecular

weight between the LMW-form and the HMW-form in HMDM, yielding a single band of glycan-free

ADA2 at 53 kD (**Fig. 2C-E**). Hence, the difference in molecular weight between intracellular WT

ADA2 (LMW) and secreted WT ADA2 or intracellular pathogenic ADA2 (both HMW) was due to

differential protein glycosylation. Of note, glycan removal yielded a smaller band of 53 kD in addition

to the glycan-removed 56 kD protein for tagged WT ADA2 overexpressed in HEK293T cells that was

absent for all tested pathogenic variants and did not appear after inhibition of N-glycosylation by

tunicamycin (**Fig. 2B**). As this band was absent after glycan removal of endogenous WT ADA2, its

physiological relevance is unclear.

Overall, we showed that the difference in size between HMW- and LMW-ADA2 is due to differential

protein glycosylation.

N-glycosylated ADA2 undergoes glycan trimming in the ER as part of protein quality control

In order to better characterize the glycosylation of endogenous ADA2, we inhibited N-glycosylation

in HMDM and the monocytic cell lines U-937 and THP-1 by 24-hour treatment with tunicamycin. We

found that this treatment only acted on the fainter HMW-band while LMW-ADA2 remained

unchanged in size (**Fig. 2D+E**). In accordance with previous findings *(13)*, we showed that adenosine

deaminase activity of ADA2 was preserved after glycan removal while treatment with tunicamycin

reduced enzymatic activity. This was true for ADA2 overexpressed in HEK293T cells as well as

endogenous ADA2 from HC HMDM (**Fig. S3B+C**). These findings indicated that N-glycosylation

was required for adequate folding of the ADA2 protein. The ER glucosidases I and II form part of

the ER quality control by removing glycosyl residues from accurately folded proteins prior to transfer

to the Golgi apparatus *(20)*. Treatment with castanospermine, an inhibitor of the ER glucosidases I

and II, caused an additional increase in the size of HMW-ADA2 in both HC and DADA2 HMDM

(**Fig. 3A**), confirming trimming of N-glycosylated ADA2 in the ER. Since the same shift in molecular

weight was observed in DADA2 HMDM, we concluded that processing of WT and mutant does not

yet diverge at this early stage of the secretory pathway. As seen previously upon incubation with

tunicamycin, the LMW-form of ADA2, which is absent in DADA2 HMDM, was unaffected by this

treatment (**Fig. 3A**). We confirmed this finding in the three ADA2-expressing cell lines U-937, THP-

1 and Jurkat (**Fig. 3B**). Since hypoglycosylated LMW-ADA2 was still detected after 24-hour treatment

with tunicamycin – an inhibitor of the initial step of N-glycosylation – we concluded that glycosylation

and trimming of the residual LMW-form must have occurred prior to the initiation of the incubation

period, suggesting high stability of LMW-ADA2. Accordingly, LMW-ADA2 was responsive to

prolonged incubation with tunicamycin at a lower concentration (**Fig. 3B**). The same was true for

inhibition of ER glucosidases: the more stable LMW-ADA2 was unaffected by 24-hour incubation

with castanospermine and the detected LMW-form had likely already matured before the beginning of

the incubation period. We confirmed the increased stability of LMW-ADA2 compared to HMW-

ADA2 by cycloheximide chase assay (**Fig. 3C** and **Fig. S3D**). Besides, this experiment revealed that

HMW-ADA2 protein expressed by HEK293T cells transfected with pathogenic *ADA2* variants

showed increased intracellular stability compared with the WT protein. This was also true after

inhibition of N-glycosylation and after an increased incubation period of 48 hours (**Fig. 3C** and

Fig. S3E). Since ADA2 is a secreted protein, protein secretion must be taken into account when

assessing the intracellular signal reduction in this assay over time (**Fig. S3F**). Impaired secretion has

previously been established as a characteristic of pathogenic ADA2 variants *(2, 14)*. We confirmed the

secretory defect in HMDM and serum from our DADA2 cohort (**Fig. 3D**). Overexpression in

HEK293T cells recapitulated the endogenous behavior of the respective variants well (**Fig. 3D**).

Comparing the levels of increased ADA2 stability observed by cycloheximide chase assay with residual

protein secretion, the variants p.T129P and p.R169Q that exhibited complete absence of secretion did

indeed show the highest levels of intracellular ADA2 24 hours after inhibition of protein synthesis

(**Fig. 3C** and **Fig. S3D**). However, intracellular levels of the secreted mutant p.P344L were still higher

compared with WT.

We next blocked the secretory pathway to impair secretion of WT ADA2 and thereby confirmed that

the secretory defect strongly contributed to the increased intracellular protein levels of mutant ADA2

(**Fig. 3E** and **Fig. S4**). While the secretory defect of mutant ADA2 was phenocopied by WT ADA2

upon incubation with brefeldin A (BFA), we also found that this treatment caused increased

degradation of the mutants p.G47V, p.T129P and p.R169Q (**Fig. 3E** and **Fig. S4**). BFA inhibits ER

to Golgi trafficking. We observed that treatment of WT ADA2-expressing cells with BFA yielded a

single ADA2 band with a molecular weight between that of HMW- and LMW-ADA2 (**Fig. 3B+E**),

suggesting that both HMW- and LMW-ADA2 are the result of further glycan processing in the Golgi.

Indeed, inhibition of trans Golgi transport by monensin *(21, 22)* did not impair the generation of

LMW-ADA2 in cells overexpressing WT ADA2 (**Fig. 3E**). The differential behavior in response to

the secretory pathway inhibitors BFA and monensin was confirmed in DADA2 HMDM (**Fig. 3F**).

The absence of LMW-ADA2 upon inhibition of ER to Golgi trafficking and the increase in LMW-

ADA2 upon inhibition of trans Golgi transport supported our hypothesis that LMW- and HMW-

ADA2 are generated from a common precursor glycoform after transport to the Golgi apparatus

(**Fig. 3G**).

In summary, we showed that N-glycosylation is required for folding and secretion of ADA2 and that

both WT and mutant ADA2 are subject to processing by the ER glucosidases I and II while only WT

ADA2 undergoes further glycan trimming in the Golgi apparatus.

The pathogenic ADA2 variant p.R169Q shows increased binding to proteins of the protein

folding machinery

After verifying the expression of LMW- and HMW-ADA2 in cells expressing WT ADA2 and an

impairment of glycan-processing in DADA2 cells, our aim was to further characterize the different

glycoforms of ADA2 in the second part of this study: (i) retained mutant HMW-ADA2, (ii) secreted

WT HMW-ADA2 and (iii) intracellular WT LMW-ADA2.

First, we performed immunoprecipitation followed by mass spectrometry (IP-MS) to analyze the

interactome of WT ADA2 and the pathogenic ADA2 variant p.R169Q transfected into HEK293T

cells. p.R169Q displayed increased interaction with proteins involved in the ER quality control

machinery for glycoproteins (**Fig. 4A** and **Table S1**). Upon inhibition of N-glycosylation by

tunicamycin, we observed a similar increase also for the WT protein (**Fig. 4A** and **Table S1**),

highlighting the importance of adequate N-glycosylation for successful folding and secretion of ADA2.

Additionally, we observed an increase in lysosomal proteins in the ADA2 interactome after treatment

with tunicamycin, suggesting autophagic degradation of non-glycosylated ADA2. Among the induced

interactors were several proteins of the chaperone-mediated autophagy pathway, e.g. LAMP2 and

HSPA13, and we verified the presence of a KFERQ-related motif (QKFVE, amino acids 274-278) in

the sequence of ADA2 *(23)*. IP-MS of HC and DADA2 HMDM did not yield any specific binders,

which was possibly due to low protein abundance in the pull down of endogenous ADA2 (**Fig. S5A**).

Notably, IP preferentially pulled down the HMW-form of ADA2 (**Fig. S5B+C**). The experiment

consequently did not allow us to draw conclusions about the interactome and potential distinctive

function of LMW-ADA2.

Given the increased binding of mutant ADA2 to the ER folding machinery, we verified whether

pathogenic *ADA2* variants induced an increased ER stress response. In line with previous reports *(14)*,

we observed that some pathogenic *ADA2* variants caused signs of activation of an ER stress response,

measured by increased levels of CHOP (*DDIT3*) and BiP (*HSPA5*) (**Fig. S5D**). These differences

were however no longer present after down-titrating the amount of transfected DNA to achieve near-

physiological protein levels (**Fig. 4B** and **Fig. S5E+F**) and neither was there a difference in BiP

expression in CD14[+] monocytes from DADA2 patients compared to healthy controls (**Fig. 4C** and

Fig. S6). In line with other studies *(8, 24)*, we showed that protein misfolding led to intracellular

aggregate formation of mutant ADA2 (**Figure 4D+E**). In summary, our experiments revealed the

interaction of the ADA2 variant p.R169Q overexpressed in HEK293T cells with mediators of the

folding machinery and ER quality control despite lack of increased ER stress in the presence of

pathogenic ADA2 variants. The interactome analyses were inconclusive regarding the characterization

of LMW-ADA2.

Wild-type HMW-ADA2 is subject to additional glycan maturation prior to secretion

Next, we aimed to further explore the glycan structure of HMW-ADA2. We hypothesized that the ER retention of most pathogenic protein variants of ADA2 leads to incomplete glycan processing. To examine the glycosylation pattern of WT and mutant ADA2, we performed additional glycan removal experiments in ADA2$^{-/-}$ U-937 cells transduced pathogenic *ADA2* variants. We chose a monocytic cell line for these experiments to minimize errors due to differential expression of glycosylation enzymes in different cell types.

First, we confirmed that complete N-glycan removal by PNGase F cancelled out the differences in molecular weight between HMW- and LMW-ADA2, confirming that they are due to N-glycosylation (**Fig. 5A**). Endo H is another endoglycosidase that cleaves high mannose glycans and – unlike PNGase F – does not remove complex glycans. We found that WT and mutant ADA2 were both sensitive to glycan removal by Endo H (**Fig. 5A+B**), hinting at the presence of high mannose oligosaccharides in both LMW- and HMW-ADA2 glycans. In contrast with PNGase F treatment, glycan removal by Endo H revealed differences between WT and mutant ADA2: Compared with mutant ADA2, Endo H-treated WT ADA2 visualized as a smear on western blot as opposed to the clean single band observed for mutant ADA2 in U-937 cells and HMDM (**Fig. 5A+B**). This suggests that WT ADA2 undergoes additional glycan processing as complex glycans are less sensitive to Endo H. Since the size of LMW-ADA2 makes the presence of additional carbohydrate residues unlikely, we concluded that in WT ADA2-expressing cells, the glycan trees of HMW-ADA2 mature into complex glycans, generating HMW(c)-ADA2, the HMW-form observed in HC HMDM (**Fig. 5C**). For all tested pathogenic variants, glycan removal by Endo H yielded a clean band very close in molecular weight to complete removal of N-glycans by PNGase F (**Fig. 5A**). Consequently, mutant HMW-ADA2 is likely not subject to any glycan maturation beyond the ER, which is in agreement with the postulated ER retention of pathogenic ADA2 protein variants. This finding also implies that WT HMW(c)-ADA2 and mutant

HMW-ADA2 – while very similar in molecular weight –differ in their glycan structures and do not

represent the same ADA2 glycoform.

Collectively, our findings suggest that the N-glycan structures of ADA2 contain high-mannose

oligosaccharides and that WT HMW(c)-ADA2 contains complex glycan branches while mutant

HMW-ADA2 does not undergo advanced glycan maturation.

LMW-ADA2 is generated by glycan trimming through alpha-mannosidases

The addition of complex glycan structures is usually preceded by the removal of α-linked mannose

residues in the late ER and Golgi by a set of α-mannosidases *(25)*. We therefore hypothesized that

mutant ADA2 is inaccessible to glycan-trimming by endogenous α-mannosidases, accounting for the

absence of complex glycan structures as well as the absence of LMW-ADA2. To test this hypothesis,

we incubated WT and mutant ADA2-expressing U-937 cells with different inhibitors of glycan-

processing enzymes. The corresponding western blots were performed in a capillary-based separation

and detection system to allow for a more precise distinction of the different molecular weight peaks.

After synthesis and N-glycan transfer in the ER, proteins are initially trimmed by the ER α-glucosidases

I and II that are inhibited by castanospermine *(20)*. As shown in HMDM, castanospermine caused an

increase in the molecular weight of HMW-ADA2 (**Fig. S7**). By increasing the incubation time

≥ 48 hours we observed a shift also for LMW-ADA2, confirming the superior stability of LMW-

ADA2. The change in molecular weight of both HMW- and LMW-ADA2 upon inhibition of the ER

α-glucosidases I and II supported our hypothesis that the processing of the two glycoforms only

diverges after exiting the ER. After passing the quality control mechanisms in the ER, proteins can be

trimmed further by different α-mannosidases in the ER and Golgi. Kifunensine inhibits class I α-

mannosidases that are typically found in the late ER and cis-Golgi and cleave α-1,2-linked mannose

residues *(20)*. Incubation of U-937 cells with kifunensine led to an increase in the molecular weight of

LMW-ADA2 (**Fig. 6A** and **Fig. S7**). After 96 hours, LMW-ADA2 almost overlay HMW-ADA2

(**Fig. 6A**). Concomitant incubation with swainsonine led to similar results. However, incubation with

swainsonine alone also increased the molecular weight of LMW-ADA2, suggesting that multiple

mannosidases are involved in the trimming of LMW-ADA2. Swainsonine inhibits both α-mannosidase

II in the medial Golgi and the lysosomal α-mannosidase *(26)*. To better understand whether also the

lysosomal enzyme is involved in the trimming of LMW-ADA2, we analyzed HMDM from a patient

with α-mannosidosis, harbouring biallelic mutations in *MAN2B1* encoding the lysosomal α-

mannosidase *(27)*. We found that the patient's cells phenocopied HC cells treated with the class II α-

mannosidase inhibitor swainsonine and that swainsonine treatment did not have an additional effect

on the molecular weight of LMW-ADA2 expressed in the lysosomal α-mannosidase-deficient cells

(**Fig. 6B**). This finding suggests that LMW-ADA2 undergoes processing not only in the ER and Golgi

but also in the lysosomes. Finally, as a proof of principle, we performed glycan removal by α1-2,3,6

mannosidase on WT ADA2. This experiment yielded a single band slightly lower but very close in

molecular weight to LMW-ADA2 (**Fig. 6C**). This is in line with the findings of the previous

experiments since the inhibited glycan-processing enzymes typically do not cleave α-1,6-linked

mannose residues, which accounts for the lower molecular weight of α1-2,3,6 mannosidase-treated

ADA2 compared with LMW-ADA2 (**Fig. 6D+E**). Importantly, the glycan structures of LMW-ADA2

predicted by this set of experiments corresponded with the glycan profile determined in another study

(**Fig. S8A**) *(28)*.

In summary, we demonstrated that LMW-ADA2 is generated from HMW-ADA2 by glycan-

processing through multiple α-mannosidases in the ER, Golgi and lysosomal compartments.

LMW-ADA2 localizes to the lysosomes and can be generated from WT HMW(c)-ADA2 in

ADA2-deficient cells

To further support the processing of LMW-ADA2 by the lysosomal α-mannosidase, we next aimed to

verify the lysosomal localization of the LMW-glycoform in monocytic cells. To this end, we performed

immunofluorescence microscopy of U-937 cells expressing WT ADA2 or the pathogenic ADA2

variant p.R169Q. In cells expressing the WT protein, ADA2 colocalized with the lysosomal marker

LAMP1 (**Fig. 7A** and **Fig. S8B**). Moreover, the experiment confirmed ER retention of the mutant

protein while WT ADA2 was found in the ER, Golgi and lysosomal compartments (**Fig. 7A**). To

verify that the intracellular glycoform of ADA2 localizing to the lysosomes was indeed LMW-ADA2,

we then performed subcellular fractionation experiments. We showed that the lysosomal fraction

contained only LMW-ADA2 (**Fig. 7B**). LMW-ADA2 was also present in the cytosolic and membrane

fractions. HMW-ADA2 localized exclusively to the membranous fraction that also contains ER and

Golgi membrane proteins (**Fig. 7B**).

Besides intracellular transport, proteins can also enter the lysosomes via the endocytotic pathway. To

evaluate whether this route allowed the uptake and processing of ADA2 from the extracellular space,

we incubated ADA2$^{-/-}$ U-937 cells with the supernatant of transfected HEK293T cells containing WT

HMW(c)-ADA2. We first confirmed that exogenous HMW(c)-ADA2 protein was taken up by the

ADA2-deficient cells (**Fig. 7C**). Next, we analyzed whether HMW(c)-ADA2 underwent intracellular

glycan processing after uptake. Within the first 6 hours of incubation, HMW-ADA2 was the only

glycoform found intracellularly (**Fig. 7D**). After 24 hours, we observed a strong signal for LMW-

ADA2 in the whole cell lysate of the ADA2-deficient cells suggesting that exogenous HMW(c)-ADA2

was subject to intracellular glycan trimming after uptake (**Fig. 7D**). In accordance with our previous

findings, LMW-ADA2 was not found in the supernatant of the incubated cells. This finding also

mechanistically adds to our previous observation that LMW-ADA2 exhibits increased stability:

Generation of LMW-ADA2 from internalized previously secreted extracellular HMW(c)-ADA2

represents an additional mechanism accounting for the prolonged presence of intracellular LMW-

ADA2 after inhibition of protein synthesis or N-glycosylation.

Finally, we confirmed that the secreted pathogenic *ADA2* variant p.G47A was also taken up into the

monocytic cells. Unlike the WT protein, the mutant protein was however not processed into LMW-

ADA2 (**Fig. 7D**). These experiments confirm that LMW-ADA2 can be generated from exogenous

extracellular ADA2 and that ADA2-deficient cells possess the glycan-processing machinery to generate

LMW-ADA2 from the WT protein (**Fig. 7E**).

Absence of LMW-ADA2 correlates with impaired ADA2 enzyme activity and predicts the

pathogenicity of *ADA2* variants

After characterizing LMW-ADA2 as the intracellular glycoform of ADA2 and establishing its absence

as a feature of DADA2 cells, we validated our findings on a larger set of *ADA2* variants. For this, we

analyzed a total of 34 *ADA2* variants, including 18 variants found in homozygous state in DADA2

patients and two variants predicted to be benign with homozygous individuals documented in the

gnomAD database (p.H335R and p. R230Q) (gnomAD v4.1.0, https://gnomad.broadinstitute.org).

Our data set also included all the *ADA2* variants previously described as DADA2-associated despite

an ADA2 enzyme activity above 25% of WT ADA2 *(29)*. The included variants cover all the domains

of the ADA2 protein (**Fig. 8A**). The variants were overexpressed in HEK293T cells and studied for

LMW-ADA2 expression, ADA2 protein secretion and both extracellular and intracellular ADA2

enzyme activity (**Fig. 8B**). In line with our studies on the trafficking of LMW-ADA2, this glycoform

was exclusively processed from variants that also showed residual ADA2 secretion (**Fig. 8B**). The

greatest part of the pathogenic variants with residual secretion lacking enzymatic activity was also

characterized by the absence of LMW-ADA2 (**Fig. 8B**). Overall, we found an excellent correlation

between the expression of LMW-ADA2 and extracellular ($r = 0.91$, $p < 0.0001$) or intracellular ADA2

enzyme activity (r = 0.89, p < 0.0001) (**Fig. 8C**). Two pathogenic variants lacking ADA2 enzyme

activity – p.H112Q and p.T360A – showed moderate expression of LMW-ADA2. By contrast,

p.P334L was characterized by absent LMW-ADA2 and a residual ADA2 enzyme activity > 25%

(**Fig. 8C**). All the DADA2-associated variants previously reported to show comparatively high

enzymatic activity also showed intermediate levels of LMW-ADA2. Four of those variants were

reported in homozygous state in DADA2 patients (**Table 2**). Several of the variants were classified as

benign by different pathogenicity scores despite having been reported in DADA2 patients (**Table 2**).

Overall, the REVEL, CADD and PolyPhen scores correlated best with pathogenicity by lack of LMW-

ADA2 (**Fig. 8D** and **Table S2+3**). SIFT classified the majority of variants as pathogenic despite

preserved LMW-ADA2 expression and enzymatic activity while with alpha missense even many clearly

pathogenic variants were categorized as benign.

In summary, we confirmed that LMW-ADA2 expression of ADA2 protein variants strongly correlates

with their adenosine deaminase activity and verified absence of LMW-ADA2 as a robust indicator of

pathogenicity.

Discussion

In this study, we showed that pathogenic protein variants of ADA2 expressed in HMDM from

DADA2 patients do not undergo glycan processing beyond the ER and that DADA2 HMDM

therefore lack lysosomal LMW-ADA2 and secreted HMW(c)-ADA2. We reported that the LMW-

form represents an intracellular glycoform of ADA2 that is generated by α-mannosidase-mediated

glycan cleavage after transport to the Golgi apparatus.

The importance of N-glycosylation in the trafficking and function of ADA2 has previously been

established *(12, 13)*. Yet, the behavior of mutant ADA2 at the cellular level has to date only been

examined in overexpression systems of cells that do not express the protein endogenously *(13, 14, 24)*.

Importantly, expression of glycosyltransferases and glycosidases and therefore glycosylation patterns

of individual proteins show striking variation even between different immune cell populations *(30)*. It

is therefore essential to understand the behavior of ADA2 in the patients' primary cells. In our cohort

of 10 DADA2 patients, we identified the intracellular predominance of unprocessed HMW-ADA2 in

the absence of LMW-ADA2 as a feature that was shared by all missense variants, independently of the

patients' clinical phenotype. Differential patterns of glycosylation of the same protein being associated

with distinctive protein functions have previously been described: IL-24 is a signal peptide-containing

glycoprotein that is secreted via the ER. A partially glycosylated cytosolic form of IL-24 has been

shown to mediate protein kinase R activation independently of its extracellular function *(31)*. The

predominance of the partially glycan-trimmed LMW-form of ADA2 in HC HMDM raises the question

of whether, in a similar way, this form differs in function from secreted HMW(c)-ADA2. In fact, the

function of ADA2 has repeatedly been subject to discussion for several reasons: (i.) Pathogenic ADA2

variants display a wide spectrum of residual adenosine deaminase function which does not clearly

correlate with disease severity *(29)*. (ii.) ADA2 has evolved towards lower affinity to adenosine in higher

species *(11)*. (iii.) ADA1 exhibits superior adenosine deaminase activity (Km of 20-50 μM vs. 2-3 mM

for ADA2) and extracellular ADA1 bound to CD26 theoretically compensates for the absence of

ADA2 at physiological levels of adenosine *(17)*. Yet, the majority of studies proposing possible

pathomechanisms underlying the immunological phenotype of DADA2 rely on the role of ADA2 as

an extracellular adenosine deaminase *(15, 16)*. In contrast, two groups introduced the possibility of an

intracellular function of ADA2 as a lysosomal protein *(28, 32)*. Greiner-Tollersrud and colleagues

characterized the glycan structures of ADA2 isolated from porcine brain and identified a high

mannose-6-phosphate content, indicating targeting of ADA2 to the lysosome. Indeed, they showed

an overlap between the glycosylation pattern of ADA2 with several known lysosomal proteins *(28)*.

We showed that the glycan structures of LMW- and HMW-ADA2 expressed in HMDM were sensitive

to glycan removal by Endo H, suggesting the presence of high-mannose oligosaccharides *(33)*. This is

in line with a potential lysosomal localization of ADA2 that we also confirmed by imaging. Moreover,

we showed that multiple – also lysosomal – α-mannosidases are involved in the processing of LMW-

ADA2. Combined inhibition of class I and II α-mannosidases caused an increase in the molecular

weight of LMW-ADA2 yielding a glycoform almost identical to HMW-ADA2. The current lack of an

antibody that sufficiently pulls down the LMW-glycoform by immunoprecipitation prevents a

thorough glycan and interactome analysis of LMW-ADA2 at this point in time. We were however able

to show that the predicted glycan structures derived from our inhibition experiments aligned perfectly

with glycan analyses performed on porcine ADA2 (**Fig. S8A**) *(28)*. Combined with the previous studies

(28, 32), our data provide a strong hint that LMW-ADA2 represents the lysosomal glycoform of ADA2

and might be functionally distinct from secreted HMW(c)-ADA2. In addition, we show for the first

time that exogenous WT ADA2 can be taken up by ADA2-deficient cells and – even more importantly

– processed into the intracellular LMW-glycoform (**Fig. 7C+D**). This finding is crucial from a clinical

perspective given current endeavors to develop new therapeutic approaches including the use of

PEGylated ADA2. Our study provides the first hint at the capacity of externally applied ADA2 to

successfully carry out also intracellular functions of ADA2.

In the HEK293T model system, transfected ADA2 was predominantly expressed as the HMW-form

corresponding to secreted ADA2 in these cells, highlighting the limitations of this system to

understand the behavior of endogenous ADA2. Discrepancies between the two systems are also

evident from the fact that ADA2 in whole cell lysates from transfected HEK293T cells was equally

sensitive to glycan removal by PNGase F or Endo H *(13)*, while we observed some residual glycan

structures upon Endo H treatment of ADA2 in lysates from U-937 cells and HMDM (**Fig. 5A+B**).

A caveat regarding the relevance of macrophage LMW-ADA2 in the pathogenesis of DADA2 is that

in primary CD14[+] monocytes, the cells with the highest endogenous expression of ADA2, differentially

glycosylated forms of ADA2 cannot be clearly distinguished (**Fig. 1A+B**). Moreover, patient

monocytes – both from patients with missense variants with normal *ADA2* mRNA expression and

from patients with variants that exhibit nonsense-mediated mRNA decay – show complete absence of

mutant ADA2 protein expression. This implies that mutations in the *ADA2* gene might manifest

differently on a cellular level depending on the expressing cell.

This notion needs to be taken into account when examining the potential induction of ER stress by

pathogenic variants of ADA2. Intracellular retention of ADA2 has only been reported after

overexpression in HEK293T cells *(2, 14)*. While primary monocytic cells also show impaired secretion

of mutant ADA2, intracellular protein levels are very low in HMDM and close to absent in CD14[+]

monocytes. In our cohort of DADA2 patients, we did not observe an increased ER stress response in

CD14[+] monocytes. This was in line with single-cell RNA-sequencing data reported by Watanabe et al.

that showed downregulation of eIF2 signaling – a main pathway of the integrated stress response *(34,*

35).

Although we could not confirm that the presence of misfolded mutant ADA2 contributes to the

pathophysiology of the disease by induction of an unfolded protein response, our data highlight the

need for a better understanding of the folding and trafficking mechanisms of WT and mutant ADA2.

We verified that the presence of pathogenic ADA2 protein variants as well as inhibition of N-

glycosylation provoked the formation of intracellular protein aggregates, thereby extending previous

reports *(13, 24)*.

Next to Sanger sequencing of the *ADA2* gene, the measurement of serum ADA2 enzyme activity

currently represents the diagnostic gold standard for DADA2. Assuming an intracellular function of

the protein, it is possible that this strategy may overlook a proportion of patients. In this article, we

validated the absence of LMW-ADA2 as a new characteristic of pathogenic *ADA2* variants. We

showed that the pathogenic variant p.P344L exhibited complete absence of LMW-ADA2 despite a

residual enzyme activity above 25% of WT ADA2. This highlights the potential benefit of additionally

checking new variants for LMW-ADA2 expression in the validation process, especially when enzyme

activity is inconclusive. Overall, we did however find an excellent correlation between LMW-ADA2

expression and enzyme activity, emphasizing that in most cases pathogenic variants affect both glycan

processing and catalytic activity. This is in line with a similar correlation shown by Greiner-Tollersrud

and colleagues in the study of the lysosomal function of ADA2 *(28)*. It is important to note that both

ADA2 enzyme activity and LMW-ADA2 expression are imperfect biomarkers of pathogenicity. We

identified 11 DADA2-associated variants with comparatively high LMW-ADA2 expression and

enzyme activity (**Table 2**). Described in patients with a DADA2-phenotype, they have all been

classified as likely disease-causing. However, given our incomplete understanding of the

pathophysiology of DADA2, the pathogenicity of these variants has to be considered uncertain. For

p.F355L, Nihira et al. concluded from their analyses that this variant may not be pathogenic, also

considering its high allele frequency. Moreover, for ambiguous variants only found in compound

heterozygous state in DADA2 patients, dominant-negative effects of the other pathogenic variant also

need to be taken into account *(8)*. Lastly, it has to be pointed out that for the variant p.H91R that

exhibits close-to-normal enzymatic activity and LMW-ADA2 expression when expressed in HEK293T

cells, a homozygous patient has been described that showed strongly reduced serum ADA2 enzyme

activity *(36)*. In our study, we showed that – unlike HEK293T cells – CD14$^+$ monocytes expressing

pathogenic *ADA2* variants show absent ADA2 expression. Therefore, it is conceivable that the

phenotype of certain *ADA2* variants only becomes apparent in the patients' primary cells.

Until now, monocytes as the main producers of endogenous ADA2 have received the most attention

in deciphering the pathophysiology of the disease. Considering the role of glycosylation in protein

folding and the potential impact of this modification on protein function, we cannot exclude that the

main ADA2 function might vary even between different immune cells. This might also account for

the diverse clinical phenotype of the disease and the varying involvement of different immune cell

populations. The large number of pathogenic variants and their diversity with respect to biochemical

characteristics are additional factors contributing to the difficulties in delineating a single all-

encompassing pathomechanism that explains the complexity of the disease.

In conclusion, we demonstrate for the first time the differential N-glycosylation of pathogenic ADA2

in patient-derived macrophages. We establish LMW-ADA2 as a hypoglycosylated non-secreted form

of ADA2 in healthy controls. This LMW-ADA2 is generated by α-mannosidases after transfer to the

Golgi and is absent in DADA2. Further studies will be needed to confirm the functional relevance of

these results in the pathogenesis of the disease.

Materials and Methods

Experimental design

This exploratory study included patients treated at University Hospitals Leuven and their parents were

included as heterozygous carriers. The diagnosis of DADA2 was made based on the clinical phenotype

followed by a combination of serum ADA2 enzyme activity and sequencing of the *ADA2* gene

according to the DADA2 management guidelines *(37)*. Healthy donors were recruited at University

Hospitals Leuven and KU Leuven. This study was approved by the Ethics Committee for Research of

Leuven University Hospitals (project numbers S63077, S63807). The patient with α-mannosidosis was

recruited at Charité University Hospital in Berlin. The inclusion of the patient from Berlin was

approved by the Ethics Committee of Charité University Medicine Berlin (project number

EA2/1178/22). The patients provided written informed consent prior to participation. The study was

performed in accordance with the ethical standards as laid down in the 1964 Declaration of Helsinki

and its later amendments. Our study examined male and female individuals, and similar findings are

reported for both sexes.

The aim of this study was to determine the protein characteristics of ADA2 endogenously expressed in human immune cells carrying pathogenic *ADA2* variants. To compensate for the limited sample size in the study of patients with rare diseases, our results were validated in two separate *in vitro* models of ADA2 deficiency examining a total of 34 pathogenic *ADA2* variants. The results were confirmed using two different anti-ADA2 antibodies both validated with the help of ADA2 knockout samples and reproduced in two technically distinct blotting systems to exclude technical artefacts.

Sanger sequencing

Genomic DNA samples were prepared from heparinized peripheral blood following the instructions of the QIAamp DNA Blood Mini kit (#51104; QIAGEN). Primers were designed with the help of Oligo Primer Analysis Software version 7 (Molecular Biology Insights). ADA2-specific gDNA amplification was performed using Platinum™ SuperFi™ PCR Master Mix (#12358010; Thermo Fisher Scientific). PCR products were purified using the QIAquick PCR purification kit (#28106; QIAGEN). Sanger sequencing was performed on an ABI 3730 XL Genetic Analyzer (Applied Biosystems) at LGC Genomics (Berlin, Germany). Sequencing data were analyzed using Chromas 2.6.5 (http://www.technelysium.com.au).

Cell culture

Peripheral blood mononuclear cells (PBMCs) were isolated by density gradient centrifugation using Lymphoprep™ (#1114546; PROGEN) in SepMate™ isolation tubes (#85450; STEMCELL Technologies) according to the manufacturer's instructions. Prior to monocyte isolation, the final centrifugation step was performed at 300xg for 10 minutes at 4 °C. CD14$^+$ monocytes were isolated magnetically by positive selection using human CD14 MicroBeads (#130-050-201; Miltenyi Biotec) on LS columns (#130-042-401; Miltenyi Biotec) according to the manufacturer's protocol. The cells were

eluted into complete medium (RPMI 1640 medium (#61870044; Gibco) supplemented with 10% fetal

calf serum (FCS) (#S181BH-500, Biowest) and 1% penicillin-streptomycin (#15140122; Gibco).

Purity > 97% of CD14$^+$ monocytes after magnetic sorting was verified by flow cytometry (anti-CD14-

FITC clone MφP9, 1:40, #345784; BD Biosciences). For macrophage differentiation, 3x10^5 CD14$^+$

monocytes were seeded in a 12-well plate in 1 mL complete medium containing 20 ng/mL GM-CSF

(#300-03; Peprotech) or 50 ng/mL M-CSF (#300-25; Peprotech). The cells were differentiated for ten

days, with medium changes every three days. U-937, THP-1 and Jurkat cells (all purchased from

ATCC) were cultured in complete RPMI medium. HEK293T cells (ATCC) were cultured in DMEM

supplemented with 10% FCS and 1% penicillin-streptomycin. For transfection, the cells were seeded

in medium containing no antibiotics.

Inhibition of the secretory pathway was achieved by 24-hour incubation with 1 μg/mL brefeldin A

(#00-4506-51; Thermo Fisher Scientific) or 2 μg/mL monensin (GolgiStop™, #554724, BD

Biosciences). For analysis of N-glycosylation and glycan trimming, the cells were incubated with

tunicamycin (#T7765; Sigma-Aldrich) at 2.5 μg/mL (HEK293T cells) or 5 μg/mL (HMDM) and

castanospermine (#BML-S107-0100; Enzo Life Sciences) at 100 μg/mL for 24h or 48h. α-

mannosidases were inhibited by 72-96h incubation with 375 nM kifunensine (#K1140; Sigma-Aldrich)

and/or 10 μM swainsonine (#S8195; Sigma-Aldrich). Monocytes were treated with 1 μg/mL

tunicamycin for 24h for ER stress induction.

Transfection

The plasmid expressing myc-DDK-tagged wild-type ADA2 (transcript variant 3, NM_001282225) was

purchased from OriGene Technologies (#RC238645). Pathogenic *ADA2* variants were created by

site-directed mutagenesis using the Q5® Site-Directed Mutagenesis Kit (#E0554; New England

Biolabs) according to the manufacturer's instructions. Stable competent E. coli (#C3040H, New

England Biolabs) were transformed with the generated constructs. Plasmid DNA was purified with

the help of the QIAprep Spin Miniprep Kit (#27104, QIAGEN). Successful mutagenesis was verified

by Sanger sequencing (LGC Genomics). HEK293T cells were seeded at $5x10^4$ cells/well in 1 mL in a

24-well plate or $2.5x10^5$ cells/well in 2 mL in a 6-well plate 24h prior to transfection. The cells were

transfected with 5 ng and 25 ng plasmid DNA, respectively, using Lipofectamine™ 2000 Transfection

Reagent (#11668019; Invitrogen) according to the manufacturer's instructions. The medium was

changed after 24h. Cells and supernatants were collected 48h after transfection. For the reuptake

experiments, HEK293T cells were transfected with 1 μg plasmid expressing untagged wild-type ADA2

or tagged wild-type and mutant as indicated. Supernatants were collected after 48 hours and transferred

to U-937 cells for a 6-24-hour incubation period.

595   Generation of ADA2$^{-/-}$ cell lines by CRISPR/Cas9

Single-guide RNAs targeting ADA2 from the human CRISPR Brunello library (#73179; addgene) *(38)*

were cloned into the lentiCRISPRv2 puro plasmid. lentiCRISPRv2 puro was a gift from Brett Stringer

(plasmid #98290; addgene; http://n2t.net/addgene:98290; RRID: Addgene_98290). U-937 cells were

transfected by electroporation using the Neon™ Transfection System (#MPK5000; Thermo Fisher

Scientific) according to the manufacturer's instructions: $2x10^5$ cells were resuspended in Resuspension

Buffer R and mixed with 1 μg plasmid DNA. Electroporation was achieved with the following

conditions: 1325V, 10ms, 3 pulses. After electroporation, the cells were directly transferred into 500 μL

prewarmed complete RPMI 1640 medium. On day 1 after transfection, puromycin (#ant-pr-1;

InvivoGen) was added at a final concentration of 1 μg/mL. After 36h, cells were seeded at 1 cell/well

in 100 μL RPMI 1640 medium supplemented with 20% FCS in a round bottom 96-well plate. After

14 days, the clones were screened for knock out of ADA2 by western blot.

608 **Transduction**

609 Pathogenic *ADA2* variants were created by site-directed mutagenesis as described above. Fragments

610 containing the sequence coding for the untagged ADA2 protein were then generated using the

611 following primers and CloneAmp™ HiFi PCR Premix (#639298; Takara Bio):

612 ORF_ADA2_PSPXI_F_GGTGGTACTCGAGTATGTTGGTGGATGGCCCATCTGA;

613 ORF_ADA2_PmeI_R_GGTGGTGTTTAAACCTTTGTAGCCACATCTGCTATGAACTT.

614 The fragments were cleaned up using the QIAquick PCR Purification Kit (#28104; QIAGEN) and

615 cloned into the p3D lentiviral expression vector as follows: Restriction of the expression vector was

616 performed in CutSmart buffer (#B7204; New England Biolabs) with PspX1 (#R0656; New England

617 Biolabs) and PmeI (#R0560; New England Biolabs) at 37 °C overnight followed by 20-minute heat

618 inactivation at 65 °C. Dephosphorylation of 5´-ends was performed using rSAP (#M0371; New

619 England Biolabs) at 37 °C overnight followed by 20-minute heat inactivation at 65 °C. The PCR

620 fragments were subsequently inserted into the expression vector using T4 ligase (#M0202; New

621 England Biolabs) for 10 minutes at room temperature followed by 10-minute heat inactivation at

622 65 °C. After DNA cleanup (#28104; QIAGEN), stable competent E. coli (#C3040H, New England

623 Biolabs) were transformed with the generated plasmids. Plasmid DNA was purified with the help of

624 the QIAprep Spin Miniprep Kit (#27104, QIAGEN). Sanger sequencing (LGC Genomics) was

625 performed to confirm that the genetic information for ADA2 was intact and that the respective

626 mutations were present. Lentiviral particles were generated by transfecting HEK293T cells with the

627 plasmids p3A (Gag-Pol), p3B (Rev), p3C (VSV-G) and p3D (expression vector) at a ratio of 1:1:1:3

628 using Lipofectamine™ 2000 Transfection Reagent (#11668019; Invitrogen). Supernatants containing

629 lentiviral particles were harvested 48h and 72h after transfection. The viral titer was determined using

630 the qPCR Lentivirus Titer Kit (#LV900; Applied Biological Materials). Viral supernatants were

631 prepared at MOI=40 with 8 μg/mL polybrene and 10^6 U-937 cells were transduced by spinoculation

at 700*xg* for 2 hours at 32 °C. After 4 hours at 37 °C, the supernatant was replaced by fresh complete

RPMI medium. Selection with 1 μg/mL puromycin (#ant-pr-1; InvivoGen) was started 48 hours after

the transduction.

Subcellular fractionation

Membrane and cytosolic fractions were generated from $10x10^6$ U-937 cells using the Mem-PER™

Plus Membrane Protein Extraction Kit (#89842; Thermo Fisher Scientific). Incubation times were

optimized to increase protein yield to 45 minutes for the permeabilization buffer and 90 minutes for

the solubilization buffer. The lysosomal fraction was produced with the help of the Lysosome

Enrichment Kit for Tissues and Cultured Cells (#89839; Thermo Fisher Scientific) according to the

manufacturer's protocol.

Glycan removal

Glycan removal was performed on whole cell lysates generated with NP-40 lysis buffer or precipitated

supernatant protein reconstituted in NP-40 lysis buffer. Glycan removal was performed on 20 μg,

10 μg or 25 μg total protein using PNGase F (#P0704; New England Biolabs), Endo H (#P0702; New

England Biolabs), α1-2,3,6 Mannosidase (#P0768; New England Biolabs) or PNGase F (#A39245;

Gibco) according to the manufacturer's instructions. Glycan removal with PNGase F (#P0704; New

England Biolabs) was completed under denaturing conditions for 1 h at 37 °C or under non-

denaturing conditions for 4 h at 37 °C as indicated.

Immunoblotting

Adherent cells were detached with trypsin-EDTA (0.05%) (#25300062; Gibco) for 5 minutes at 37 °C

and the reaction was blocked by addition of complete medium. Cells were pelleted by centrifugation

at 400xg for 5 minutes at 4 °C. Whole cell lysates were obtained by lysing 1x10^6 cells in 25 μL RIPA

buffer (150 mM NaCl, 1% Triton X-100, 0.5% sodium deoxycholate, 0.1% SDS, pH 8.0) or NP-40

lysis buffer (150 mM NaCl, 50 mM Tris-HCl, 1% NP-40, pH 7.4) containing protease inhibitor

(#78429; Thermo Fisher Scientific) for 30 minutes on ice, followed by centrifugation at 13,500xg for

20 minutes at 4 °C.

To collect supernatants, cells were cultured in serum-free medium for 24 h. For immunoblotting,

supernatants were concentrated by adding 1200 μL ice-cold acetone to 300 μL supernatant and

overnight storage at -20°C. Supernatant proteins were then precipitated by centrifugation at 15,000xg

for 15 minutes at 4°C. The protein pellet was resuspended in 25 μL NP-40 buffer containing protease

inhibitor. Bolt™ LDS sample buffer (#B0007; Thermo Fisher Scientific) mixed with Bolt™ Sample

Reducing Agent (#B0009; Thermo Fisher Scientific) was added to the samples prior to gel

electrophoresis. For analysis of intracellular dimer formation, the reducing agent was omitted. Samples

were heated for 5 minutes at 70 °C. Supernatants from HEK293T cells transfected with ADA2 were

diluted 1:10 in 2X Bolt™ LDS sample buffer (#B0007; Thermo Fisher Scientific), serum was diluted

1:200. SeeBlue™ Plus2 Pre-stained Protein Standard (#LC5925; Thermo Fisher Scientific) was used

as protein molecular weight marker.

Proteins were transferred onto PVDF transfer membranes. The membranes were probed with the

following primary antibodies: anti-ADA2 (clone EPR25430-131, #ab288296, 1:1000; abcam), anti-

ADA2 (#HPA007888, 1:500; Sigma-Aldrich; RRID: AB_1078495), Clone OTI4C5, anti-DDK

(FLAG) (clone OTI4C5, #TA50011, 1:1000; OriGene Technologies; RRID: AB_2622345), anti-BiP

(#3183, 1:1000; Cell Signaling; RRID: AB_10695864), anti-LAMP-1 (clone H4A3, #sc-20011; 1:500;

Santa Cruz Biotechnology; RRID: AB_626853), anti-CD98 (clone E-5, #sc-376815, 1:500; Santa Cruz

Biotechnology; RRID: AB_2938854), anti-GAPDH (clone 7B, #sc-69778, 1:500; Santa Cruz

Biotechnology; RRID: AB_1124759), anti-β-actin (clone: AC-15, #A5441, 1:9,000; Sigma-Aldrich;

RRID: AB_476744) at 4°C overnight or at room temperature for two hours. The membranes were

washed and incubated with the respective HRP-coupled secondary antibodies for one hour at room

temperature: goat anti-rabbit IgG H&L (#ab205718, 1:5000; abcam; RRID: AB_2819160), goat anti-

mouse IgG (H + L) (#71045, 1:5000; Sigma-Aldrich; RRID: AB_11211441) or VeriBlot Detection

Reagent (#ab131366, 1:1000 in tris saline/5% skim milk; abcam).

The anti-ADA2 antibody (clone EPR25430-131, #ab288296, 1:1000; abcam) was used in the following

figures of the article: 1A, 1B, 1C, 1D, 1E, 3A, 3B, 3C (left), 3D, 3E, 3F, 4D, 4E, 5B (right), 6A, 6B, 6C

(top), 7C, 7D, S2C, S2D, S2E, S2F, S3A, S3D, S3F, S4, S5C, S5E, S5F, S6 and S7. The anti-ADA2

antibody (#HPA007888, 1:500; Sigma-Aldrich; RRID: AB_1078495) was applied in the remaining

blots: 2B, 2C, 2D, 2E, 3C (right), 5A, 5B (left), 6C (bottom), 7B, S3B, S3C, S3E, S5B.

Protein expression was visualized by enzymatic chemiluminescence using Pierce™ ECL western

blotting substrate (#32106; Thermo Fisher Scientific) or SuperSignal™ West Pico PLUS

Chemiluminescent Substrate (#34580; Thermo Fisher Scientific) in a ChemiDoc XRS+ Imaging

System (Bio-Rad).

The experiments examining the glycan-processing of LMW-ADA2 were performed in a Wes

Automated Western Blot System (ProteinSimple / bio-techne) according to the manufacturer's

instructions. Samples were loaded at a final protein concentration of 0.4 mg/mL.

Immunoprecipitation

Whole cell lysates were prepared from HEK293T cells transfected with wild-type *ADA2* or the variant

p.R169Q or HMDM from HC or DADA2 patients in NP-40 lysis buffer at a protein concentration of

1 mg/mL. Primary antibodies were added to 1000 μg lysate or 500 μg lysate, respectively, and

incubated rotating at 4 °C overnight. Alternatively, 1 mL supernatant was directly incubated with the

primary antibodies. After overexpression, tagged ADA2 was pulled down using 1 μg anti-DDK

(FLAG) clone OTI4C5 (#TA50011, OriGene Technologies). An isotype control sample was

incubated with 1 μg mouse IgG1 (#02-6100; Invitrogen) in parallel. Endogenous ADA2 was pulled

down using 1 μg anti-ADA2 (clone EPR25430-131, #ab288296, abcam) with rabbit monoclonal IgG

(clone EPR25A, #ab172730; abcam) used as isotype control. For immunoprecipitation, the samples

were rotated with 100 μL SureBeads™ Protein G Magnetic Beads (#1614023, Bio-Rad) for 1h at room

temperature. The beads were magnetized and washed four times with PBS/0.1% Tween 20. Samples

for mass spectrometric analysis were washed twice with NP-40 lysis buffer and twice with PBS. For

western blot, the samples were eluted from the beads in Bolt™ LDS sample buffer (#B0007; Thermo

Fisher Scientific).

Protein digest

Beads from immunoprecipitation experiments were resuspended in 20 μL denaturation buffer (6 M

Urea, 2 M Thiourea, 10 mM HEPES, pH 8.0), reduced for 30 minutes at 25 °C in 12 mM

dithiothreitol, followed by alkylation with 40 mM chloroacetamide for 20 minutes at 25 °C. Samples

were first digested with 0.5 μg endopeptidase LysC (Wako, Osaka, Japan) for 4 h. After diluting the

samples with 80 μL 50 mM ammonium bicarbonate (pH 8.5), 1 μg sequence grade trypsin (Promega)

was added overnight at 25 °C. The peptide-containing supernatant was collected and acidified with

formic acid (1% final concentration) to stop the digestion. Peptides were desalted and cleaned up using

Stage Tip protocol *(39)*. After elution with 80% acetonitrile/0.1% formic acid, samples were dried

using speedvac, resolved in 3% acetonitrile/0.1% formic acid and analyzed by LC-MS/MS.

LC-MS/MS analyses

Peptides were separated on a reversed-phase column (20 cm fritless silica microcolumns with an inner

diameter of 75 μm, packed with ReproSil-Pur C18-AQ 3 μm resin (Dr. Maisch GmbH) using a 90-

minute gradient with a 250 nL/min flow rate of increasing Buffer B concentration (from 2% to 60%)

on a High-Performance Liquid Chromatography (HPLC) system (Thermo Fisher Scientific) and

ionized using an electrospray ionization (ESI) source (Thermo Fisher Scientific) and analyzed on a

Thermo Q Exactive HF-X instrument (HEK293T) or on an Q Exactive Plus instrument (HMDM).

The Q Exactive HF-X instrument was run in data dependent mode selecting the top 20 most intense

ions in the MS full scans, selecting ions from 350 to 2000 m/z, using 60 K resolution with a $3x10^6$ ion

count target and 10 ms injection time. Tandem MS was performed at a resolution of 15 K. The MS2

ion count target was set to $1x10^5$ with a maximum injection time of 100 ms. Only precursors with

charge state 2–6 were selected for MS2. The dynamic exclusion duration was set to 30 s with a 10-ppm

tolerance around the selected precursor and its isotopes. The Q Exactive Plus instrument was run in

data dependent mode selecting the top 10 most intense ions in the MS full scans, selecting ions from

350 to 2000 m/z, using 70 K resolution with a $3x10^6$ ion count target and 50 ms injection time. Tandem

MS was performed at a resolution of 17.5 K. The MS2 ion count target was set to $5x10^4$ with a

maximum injection time of 250 ms. Only precursors with charge state 2–6 were selected for MS2. The

dynamic exclusion duration was set to 30 s with a 10-ppm tolerance around the selected precursor and

its isotopes.

Raw data were analyzed using MaxQuant software package (v1.6.3.4) *(40)*. The internal Andromeda

search engine was used to search MS2 spectra against a human UniProt database (HUMAN.2020-06),

including ADA2 mutant sequence, containing forward and reverse sequences. The search included

variable modifications of methionine oxidation, N-terminal acetylation and fixed modification of

carbamidomethyl cysteine. The FDR was set to 1% for peptide and protein identifications. Unique

and razor peptides were considered for quantification. The LFQ (label-free quantitation) algorithm

was activated. The resulting text file was filtered to exclude reverse database hits, potential

contaminants, and proteins only identified by site.

Microscopy

HMDM were differentiated with GM-CSF or M-CSF for 10 days in a 12-well plate as described above.

Cell morphology was evaluated in the culture plate by brightfield microscopy on an EVOS M7000

Microscope Imaging System (#AMF7000; Invitrogen) using the 20X objective.

The immunofluorescence staining was performed in V-shaped 96-well plates in the dark at room

temperature (RT). Cells were harvested, fixed in 4% (v/v) paraformaldehyde (Electron Microscopy

Sciences) for 10 minutes, and permeabilized with PBS containing 0.1% Tween®20 (Qbiogene Inc.)

for 10 minutes. To prevent nonspecific antibody binding, cells were incubated in blocking buffer (1x

PBS supplemented with 5% FCS and 5 mg/mL human IgG; IgG1 66.6%, IgG2 28.5%, IgG3 2.7%,

IgG4 2.2%; Grifols) for 30 minutes.

For the primary staining, an unconjugated rabbit anti-ADA2 antibody (clone EPR25430-131,

#ab288296, 1:100; abcam) was applied, followed by incubation with a secondary antibody (donkey

anti-rabbit AF488, 1:500, #A-21206, Invitrogen or goat anti-rabbit AF546, 1:500, #A-11035,

Invitrogen). In addition, co-staining was performed using rabbit anti-GM130 AF647 (clone D6B1,

1:100, #59890, Cell Signaling Technology) in combination with either rabbit anti-LAMP1 AF488

(clone D2D11, 1:100, #58996, Cell Signaling Technology) or rabbit anti-calnexin AF555 (clone C5C9,

1:100, #23198, Cell Signaling Technology). All antibodies were diluted in 1X PBS supplemented with

5% FCS, 5 mg/mL human IgG, and 0.1% Tween®20, and incubated for 2 hours at RT. After each

incubation step, cells were washed three times with PBS. Nuclear staining was performed as the final

step using DAPI (1 μg/mL in PBS / 0.1% Tween®20) for 15 minutes. For imaging, stained cells were

transferred onto glass slides using a Cytospin 4 centrifuge (Thermo Fisher Scientific) at 800 rpm for 3

minutes with medium acceleration. Images were acquired with a laser scanning confocal fluorescence

microscope (LSM 710, Carl Zeiss).

ADA2 enzyme activity

Adenosine deaminase 2 enzyme activity was determined in whole cell lysates from HEK293T cells overexpressing ADA2 or HMDM with and without glycan removal as well as in supernatants from transfected HEK293T cells and human serum. Deaminase activity was measured in a colorimetric assay adapted from Giusti and Galanti *(41)*. Erythro-9-(2-hydroxy-3-nonyl) adenine (EHNA) (#E114; Sigma-Aldrich) was used to inhibit ADA1 activity. Triplicate measurements were performed for all samples. Enzymatic activity of pathogenic ADA2 variants overexpressed HEK293T cells was normalized to the activity of wild-type ADA2.

qPCR

5-10x10^5 cells were lysed in TRIzol™ Reagent (#15596018; Thermo Fisher Scientific) for 3 minutes at room temperature and homogenized before storage at -80 °C. To measure the ER stress response, HEK293T cells were seeded at 5x10^4 cells/ml in a 24-well plate 24 h prior to transfection. TRIzol™ samples were prepared 48h after transfection. RNA was extracted using the PureLink™ RNA Mini Kit (#12183018A; Thermo Fisher Scientific) according to the manufacturer's instruction. cDNA was generated from 20 ng RNA using the SuperScript™ VILO™ cDNA Synthesis Kit (#11754050; Thermo Fisher Scientific). Quantitative polymerase chain reaction (qPCR) analysis was performed with SsoAdvanced™ Universal SYBR® Green Supermix (#1725271; Bio-Rad Laboratories) and the following primers:

ADA2(929-1038)_F_GGCTCCGAATCAAGTTCCCC;

ADA2(929-1038)_R_TAAGGCAGCTTAACGCCATCC;

ADA2(1138-1257)_F_ACCAGAATCGGCCATGGA;

ADA2(1138-1257)_R_CTACAGGGTGGTTCCTCAAGTCA;

HSPA5_F_CAAGCAACCAAAGACGCTGGA;   HSPA5_R_GCCACCCAGGTCAAACACCA;

DDIT3_F_AGAGGAAGAATCAAAAATCT;   DDIT3_R_AGCTCTGACTGGAATCTGGA;

GAPDH_F_GTCTCCTCTGACTTCAACAGCG;

GAPDH_R_ACCACCCTGTTGCTGTAGCCAA.

For all conditions, three technical replicates were measured. The experiment was run on a

QuantStudio™ 3 Real-Time PCR System (Thermo Fisher Scientific) and analyzed using the

QuantStudio™ Design & Analysis Software v1.5.2. The relative abundance of *ADA2* was normalized

to the expression level of *GAPDH*. and different conditions were compared using the $2^{-\Delta\Delta Ct}$ method

(42).

Statistical analysis

Statistical analysis was performed in R using the packages stats and Hmisc. Correlation was measured

using Pearson's correlation coefficient.

Statistical analysis of the mass spectrometry data was performed using Perseus software (v1.6.2.1) *(43)*.

Log2 transformed LFQ intensity values were filtered for minimum of 3 valid values in at least one

experimental group and missing values were imputed with random low intensity values taken from a

normal distribution. Differences in protein abundance between groups were calculated using two-

sample Student's t-test using an FDR-based significance cut-off of 5% (*HEK293T*) or 10% (*HMDM*)

in order to define enriched proteins.

References

1. Q. Zhou, D. Yang, A. K. Ombrello, A. V. Zavialov, C. Toro, A. V. Zavialov, D. L. Stone, J. J.

Chae, S. D. Rosenzweig, K. Bishop, K. S. Barron, H. S. Kuehn, P. Hoffmann, A. Negro, W. L.

Tsai, E. W. Cowen, W. Pei, J. D. Milner, C. Silvin, T. Heller, D. T. Chin, N. J. Patronas, J. S. Barber,

C.-C. R. Lee, G. M. Wood, A. Ling, S. J. Kelly, D. E. Kleiner, J. C. Mullikin, N. J. Ganson, H. H.

Kong, S. Hambleton, F. Candotti, M. M. Quezado, K. R. Calvo, H. Alao, B. K. Barham, A. Jones,

J. F. Meschia, B. B. Worrall, S. E. Kasner, S. S. Rich, R. Goldbach-Mansky, M. Abinun, E. Chalom,

A. C. Gotte, M. Punaro, V. Pascual, J. W. Verbsky, T. R. Torgerson, N. G. Singer, T. R. Gershon,

S. Ozen, O. Karadag, T. A. Fleisher, E. F. Remmers, S. M. Burgess, S. L. Moir, M. Gadina, R.

Sood, M. S. Hershfield, M. Boehm, D. L. Kastner, I. Aksentijevich, Early-onset stroke and

vasculopathy associated with mutations in ADA2. *N. Engl. J. Med.* **370**, 911–920 (2014).

2.   P. Navon Elkan, S. B. Pierce, R. Segel, T. Walsh, J. Barash, S. Padeh, A. Zlotogorski, Y. Berkun, J.

J. Press, M. Mukamel, I. Voth, P. J. Hashkes, L. Harel, V. Hoffer, E. Ling, F. Yalcinkaya, O.

Kasapcopur, M. K. Lee, R. E. Klevit, P. Renbaum, A. Weinberg-Shukron, E. F. Sener, B.

Schormair, S. Zeligson, D. Marek-Yagel, T. M. Strom, M. Shohat, A. Singer, A. Rubinow, E. Pras,

J. Winkelmann, M. Tekin, Y. Anikster, M.-C. King, E. Levy-Lahad, Mutant adenosine deaminase

2 in a polyarteritis nodosa vasculopathy. *N. Engl. J. Med.* **370**, 921–931 (2014).

3.   I. Meyts, I. Aksentijevich, Deficiency of Adenosine Deaminase 2 (DADA2): Updates on the

Phenotype, Genetics, Pathogenesis, and Treatment. *J. Clin. Immunol.* **38**, 569–578 (2018).

4.   K. J. Karczewski, L. C. Francioli, G. Tiao, B. B. Cummings, J. Alföldi, Q. Wang, R. L. Collins, K.

M. Laricchia, A. Ganna, D. P. Birnbaum, L. D. Gauthier, H. Brand, M. Solomonson, N. A. Watts,

D. Rhodes, M. Singer-Berk, E. M. England, E. G. Seaby, J. A. Kosmicki, R. K. Walters, K.

Tashman, Y. Farjoun, E. Banks, T. Poterba, A. Wang, C. Seed, N. Whiffin, J. X. Chong, K. E.

Samocha, E. Pierce-Hoffman, Z. Zappala, A. H. O'Donnell-Luria, E. V. Minikel, B. Weisburd, M.

Lek, J. S. Ware, C. Vittal, I. M. Armean, L. Bergelson, K. Cibulskis, K. M. Connolly, M.

Covarrubias, S. Donnelly, S. Ferriera, S. Gabriel, J. Gentry, N. Gupta, T. Jeandet, D. Kaplan, C.

Llanwarne, R. Munshi, S. Novod, N. Petrillo, D. Roazen, V. Ruano-Rubio, A. Saltzman, M.

Schleicher, J. Soto, K. Tibbetts, C. Tolonen, G. Wade, M. E. Talkowski, B. M. Neale, M. J. Daly,

D. G. MacArthur, The mutational constraint spectrum quantified from variation in 141,456

humans. *Nature* **581**, 434–443 (2020).

5. Infevers: an online database for autoinflammatory mutations. Copyright. Available at

https://infevers.umai-montpellier.fr/ Accessed 14/08/2023. (available at https://infevers.umai-

montpellier.fr/web/instructions_for_use.php).

6. P. Y. Lee, E. S. Kellner, Y. Huang, E. Furutani, Z. Huang, W. Bainter, M. F. Alosaimi, K. Stafstrom,

C. D. Platt, T. Stauber, S. Raz, I. Tirosh, A. Weiss, M. B. Jordan, C. Krupski, D. Eleftheriou, P.

Brogan, A. Sobh, Z. Baz, G. Lefranc, C. Irani, S. S. Kilic, R. El-Owaidy, M. R. Lokeshwar, P.

Pimpale, R. Khubchandani, E. P. Chambers, J. Chou, R. S. Geha, P. A. Nigrovic, Q. Zhou,

Genotype and functional correlates of disease phenotype in deficiency of adenosine deaminase 2

(DADA2). *J. Allergy Clin. Immunol.* **145**, 1664-1672.e10 (2020).

7. M. Dzhus, L. Ehlers, M. Wouters, K. Jansen, R. Schrijvers, L. De Somer, S. Vanderschueren, M.

Baggio, L. Moens, B. Verhaaren, R. Lories, G. Bucciol, I. Meyts, A Narrative Review of the

Neurological Manifestations of Human Adenosine Deaminase 2 Deficiency. *J. Clin. Immunol.*

(2023), doi:10.1007/s10875-023-01555-y.

8. M. Wouters, L. Ehlers, W. Van Eynde, M. E. Kars, S. Delafontaine, V. Kienapfel, M. Dzhus, R.

Schrijvers, P. De Haes, S. Struyf, G. Bucciol, Y. Itan, A. Bolze, A. Voet, A. Hombrouck, L. Moens,

B. Ogunjimi, I. Meyts, Dominant negative ADA2 mutations cause ADA2 deficiency in

heterozygous carriers. *J. Exp. Med.* **222**, e20250499 (2025).

9. N. T. Deuitch, D. Yang, P. Y. Lee, X. Yu, N. S. Moura, O. Schnappauf, A. K. Ombrello, D. Stone,

H. S. Kuehn, S. D. Rosenzweig, P. Hoffmann, C. Cudrici, D. M. Levy, E. Kessler, J. B. Soep, A.

D. Hay, A. Dalrymple, Y. Zhang, L. Sun, Q. Zhang, X. Tang, Y. Wu, K. Rao, H. Li, H. Luo, Y.

Zhang, J. M. Burnham, M. Boehm, K. Barron, D. L. Kastner, I. Aksentijevich, Q. Zhou, TNF

inhibition in vasculitis management in adenosine deaminase 2 deficiency (DADA2). *J. Allergy Clin.*

Immunol. **149**, 1812-1816.e6 (2022).

10. H. Hashem, A. R. Kumar, I. Müller, F. Babor, R. Bredius, J. Dalal, A. P. Hsu, S. M. Holland, D.

D. Hickstein, S. Jolles, R. Krance, G. Sasa, M. Taskinen, M. Koskenvuo, J. Saarela, J. van

Montfrans, K. Wilson, B. Bosch, L. Moens, M. Hershfield, I. Meyts, Hematopoietic stem cell

transplantation rescues the hematological, immunological, and vascular phenotype in DADA2.

Blood **130**, 2682–2688 (2017).

11. A. V. Zavialov, X. Yu, D. Spillmann, G. Lauvau, A. V. Zavialov, Structural basis for the growth

factor activity of human adenosine deaminase ADA2. *J. Biol. Chem.* **285**, 12367–12377 (2010).

12. P. Y. Lee, Y. Huang, Q. Zhou, O. Schnappauf, M. S. Hershfield, Y. Li, N. J. Ganson, N. Sampaio

Moura, O. M. Delmonte, S. S. Stone, M. J. Rivkin, S.-Y. Pai, T. Lyons, R. P. Sundel, V. W. Hsu, L.

D. Notarangelo, I. Aksentijevich, P. A. Nigrovic, Disrupted N-linked glycosylation as a disease

mechanism in deficiency of ADA2. *J. Allergy Clin. Immunol.* **142**, 1363-1365.e8 (2018).

13. M. Ito, Y. Maejima, K. Nishimura, Y. Nakae, A. Ono, S. Iwaki-Egawa, A role for N-glycosylation

in active adenosine deaminase 2 production. *Biochim. Biophys. Acta BBA - Gen. Subj.* **1866**, 130237

(2022).

14. L. Chen, A. Mamutova, A. Kozlova, E. Latysheva, F. Evgeny, T. Latysheva, K. Savostyanov, A.

Pushkov, I. Zhanin, E. Raykina, M. Kurnikova, I. Mersiyanova, C. D. Platt, H. Jee, K. Brodeur, Y.

Du, M. Liu, A. Weiss, G. S. Schulert, J. Rodriguez-Smith, M. S. Hershfield, I. Aksentijevich, Q.

Zhou, P. A. Nigrovic, A. Shcherbina, E. Alexeeva, P. Y. Lee, Comparison of disease phenotypes

and mechanistic insight on causal variants in patients with DADA2. *J. Allergy Clin. Immunol.* (2023),

doi:10.1016/j.jaci.2023.04.014.

15. C. Carmona-Rivera, S. S. Khaznadar, K. W. Shwin, J. A. Irizarry-Caro, L. J. O'Neil, Y. Liu, K. A.

Jacobson, A. K. Ombrello, D. L. Stone, W. L. Tsai, D. L. Kastner, I. Aksentijevich, M. J. Kaplan,

P. C. Grayson, Deficiency of adenosine deaminase 2 triggers adenosine-mediated NETosis and

TNF production in patients with DADA2. *Blood* **134**, 395–406 (2019).

16. R. Dhanwani, M. Takahashi, I. T. Mathews, C. Lenzi, A. Romanov, J. D. Watrous, B. Pieters, C.

C. Hedrick, C. A. Benedict, J. Linden, R. Nilsson, M. Jain, S. Sharma, Cellular sensing of

extracellular purine nucleosides triggers an innate IFN-β response. *Sci. Adv.* **6**, eaba3688 (2020).

17. T. K. Tarrant, S. J. Kelly, M. S. Hershfield, Elucidating the pathogenesis of adenosine deaminase

2 deficiency: current status and unmet needs. *Expert Opin. Orphan Drugs* **9**, 257–264 (2021).

18. O. K. Greiner-Tollersrud, V. Boehler, E. Bartok, M. Krausz, A. Polyzou, J. Schepp, M. Seidl, J. O.

Olsen, C. R. Smulski, S. Raieli, K. Hübscher, E. Trompouki, R. Link, H. Ebersbach, H. Srinivas,

M. Marchant, D. Staab, D. Guerini, S. Baasch, P. Henneke, G. Kochs, G. Hartmann, R. Geiger, B.

Grimbacher, M. Warncke, M. Proietti, *ADA2 is a lysosomal DNase regulating the type-I interferon response*

(2020; https://www.biorxiv.org/content/10.1101/2020.06.21.162990v2), p. 2020.06.21.162990.

19. L. Ehlers, M. Wouters, B. Pillay, S. Delafontaine, M. Dzhus, M. Baggio, T. Niehues, G. Dückers,

L. Sevenants, K. Casteels, L. D. Somer, R. Schrijvers, S. Vanderschueren, M. Jacquemyn, D.

Daelemans, A. Hombrouck, E. P. Chambers, T. Tousseyn, G. Bucciol, P. Agostinis, L. Moens, I.

Meyts, Inhibition of lysosomal degradation increases expression of mutant ADA2 in DADA2

monocytes. *J. Allergy Clin. Immunol.* **0** (2025), doi:10.1016/j.jaci.2025.06.009.

20. A. Varki, R. D. Cummings, J. D. Esko, P. Stanley, G. W. Hart, M. Aebi, D. Mohnen, T. Kinoshita,

N. H. Packer, J. H. Prestegard, R. L. Schnaar, P. H. Seeberger, Eds., *Essentials of Glycobiology* (Cold

Spring Harbor Laboratory Press, Cold Spring Harbor (NY), ed. 4th, 2022;

http://www.ncbi.nlm.nih.gov/books/NBK579918/).

21. T. Fujiwara, K. Oda, S. Yokota, A. Takatsuki, Y. Ikehara, Brefeldin A causes disassembly of the

Golgi complex and accumulation of secretory proteins in the endoplasmic reticulum. *J. Biol. Chem.*

263, 18545–18552 (1988).

22. H. H. Mollenhauer, D. James Morré, L. D. Rowe, Alteration of intracellular traffic by monensin;

mechanism, specificity and relationship to toxicity. *Biochim. Biophys. Acta BBA - Rev. Biomembr.* **1031**,

225–246 (1990).

23. P. Kirchner, M. Bourdenx, J. Madrigal-Matute, S. Tiano, A. Diaz, B. A. Bartholdy, B. Will, A. M.

Cuervo, Proteome-wide analysis of chaperone-mediated autophagy targeting motifs. *PLOS Biol.*

17, e3000301 (2019).

24. S. M. Bowers, M. Sundqvist, P. Dancey, D. A. Cabral, K. L. Brown, Pathogenic variant c.1052T>A

(p.Leu351Gln) in adenosine deaminase 2 impairs secretion and elevates type I IFN responsive gene

expression.   *Front.   Immunol.   **13**   (2022)   (available   at

https://www.frontiersin.org/articles/10.3389/fimmu.2022.995191).

25. K. Legler, R. Rosprim, T. Karius, K. Eylmann, M. Rossberg, R. M. Wirtz, V. Müller, I. Witzel, B.

Schmalfeldt, K. Milde-Langosch, L. Oliveira-Ferrer, Reduced mannosidase MAN1A1 expression

leads to aberrant N-glycosylation and impaired survival in breast cancer. *Br. J. Cancer* **118**, 847–856

(2018).

26. D. R. Rose, Structure, mechanism and inhibition of Golgi α-mannosidase II. *Curr. Opin. Struct. Biol.*

22, 558–562 (2012).

27. D. Malm, Ø. Nilssen, Alpha-mannosidosis. *Orphanet J. Rare Dis.* **3**, 21 (2008).

28. O. K. Greiner-Tollersrud, M. Krausz, V. Boehler, A. Polyzou, M. Seidl, A. Spahiu, Z. Abdullah,

K. Andryka-Cegielski, F. I. Dominick, K. Huebscher, A. Goschin, C. R. Smulski, E. Trompouki,

R. Link, H. Ebersbach, H. Srinivas, M. Marchant, G. Sogkas, D. Staab, C. Vågbø, D. Guerini, S.

Baasch, E. Latz, G. Hartmann, P. Henneke, R. Geiger, X. P. Peng, B. Grimbacher, E. Bartok, I.

Alseth, M. Warncke, M. Proietti, ADA2 is a lysosomal deoxyadenosine deaminase acting on DNA

involved in regulating TLR9-mediated immune sensing of DNA. *Cell Rep.* **43** (2024),

doi:10.1016/j.celrep.2024.114899.

29. H. Jee, Z. Huang, S. Baxter, Y. Huang, M. L. Taylor, L. A. Henderson, S. Rosenzweig, A. Sharma,

E. P. Chambers, M. S. Hershfield, Q. Zhou, F. Dedeoglu, I. Aksentijevich, P. A. Nigrovic, A.

O'Donnell-Luria, P. Y. Lee, Comprehensive analysis of ADA2 genetic variants and estimation of

carrier frequency driven by a function-based approach. *J. Allergy Clin. Immunol.* (2021),

doi:10.1016/j.jaci.2021.04.034.

30. M. A. Daniels, K. A. Hogquist, S. C. Jameson, Sweet "n" sour: the impact of differential

glycosylation on T cell responses. *Nat. Immunol.* **3**, 903–910 (2002).

31. S. Davidson, C.-H. Yu, A. Steiner, F. Ebstein, P. J. Baker, V. Jarur-Chamy, K. Hrovat Schaale, P.

Laohamonthonkul, K. Kong, D. J. Calleja, C. R. Harapas, K. R. Balka, J. Mitchell, J. T. Jackson,

N. D. Geoghegan, F. Moghaddas, K. L. Rogers, K. D. Mayer-Barber, A. A. De Jesus, D. De Nardo,

B. T. Kile, A. J. Sadler, M. C. Poli, E. Krüger, R. Goldbach Mansky, S. L. Masters, Protein kinase

R is an innate immune sensor of proteotoxic stress via accumulation of cytoplasmic IL-24. *Sci.*

Immunol. **7**, eabi6763 (2022).

32. L. Dong, W. Luo, S. Maksym, S. C. Robson, A. V. Zavialov, Adenosine deaminase 2 regulates the

activation of the toll-like receptor 9 in response to nucleic acids. *Front. Med.* (2024),

doi:10.1007/s11684-024-1067-5.

33. M.-S. Kim, D. Leahy, in *Methods in Enzymology*, Laboratory Methods in Enzymology: Cell, Lipid

and Carbohydrate. J. Lorsch, Ed. (Academic Press, 2013), vol. 533, pp. 259–263.

34. N. Watanabe, S. Gao, Z. Wu, S. Batchu, S. Kajigaya, C. Diamond, L. Alemu, D. Q. Raffo, P.

Hoffmann, D. Stone, A. K. Ombrello, N. S. Young, Analysis of deficiency of adenosine deaminase

2 pathogenesis based on single-cell RNA sequencing of monocytes. *J. Leukoc. Biol.* **110**, 409–424

(2021).

35. K. Pakos-Zebrucka, I. Koryga, K. Mnich, M. Ljujic, A. Samali, A. M. Gorman, The integrated

stress response. *EMBO Rep.* **17**, 1374–1395 (2016).

36. S. Cooray, E. Omyinmi, Y. Hong, C. Papadopoulou, L. Harper, E. Al-Abadi, R. Goel, S. Dubey,
M. Wood, S. Jolles, S. Berg, M. Ekelund, K. Armon, D. Eleftheriou, P. A. Brogan, Anti-tumour
necrosis factor treatment for the prevention of ischaemic events in patients with deficiency of
adenosine deaminase 2 (DADA2). *Rheumatol. Oxf. Engl.* **60**, 4373–4378 (2021).

37. P. Y. Lee, B. A. Davidson, R. S. Abraham, B. Alter, J. I. Arostegui, K. Bell, A. Belot, J. R. E.
Bergerson, T. J. Bernard, P. A. Brogan, Y. Berkun, N. T. Deuitch, D. Dimitrova, S. A. Georgin-
Lavialle, M. Gattorno, B. Grimbacher, H. Hashem, M. S. Hershfield, R. N. Ichord, K. Izawa, J. A.
Kanakry, R. P. Khubchandani, F. C. C. Klouwer, E. A. Luton, A. W. Man, I. Meyts, J. M. Van
Montfrans, S. Ozen, J. Saarela, G. C. Santo, A. Sharma, A. Soldatos, R. Sparks, T. R. Torgerson, I.
L. Uriarte, T. A. B. Youngstein, Q. Zhou, I. Aksentijevich, D. L. Kastner, E. P. Chambers, A. K.
Ombrello, DADA2 Foundation, Evaluation and Management of Deficiency of Adenosine
Deaminase 2: An International Consensus Statement. *JAMA Netw. Open* **6**, e2315894 (2023).

38. J. G. Doench, N. Fusi, M. Sullender, M. Hegde, E. W. Vaimberg, K. F. Donovan, I. Smith, Z.
Tothova, C. Wilen, R. Orchard, H. W. Virgin, J. Listgarten, D. E. Root, Optimized sgRNA design
to maximize activity and minimize off-target effects of CRISPR-Cas9. *Nat. Biotechnol.* **34**, 184–191
(2016).

39. J. Rappsilber, Y. Ishihama, M. Mann, Stop and go extraction tips for matrix-assisted laser
desorption/ionization, nanoelectrospray, and LC/MS sample pretreatment in proteomics. *Anal.*
Chem. **75**, 663–670 (2003).

40. S. Tyanova, T. Temu, J. Cox, The MaxQuant computational platform for mass spectrometry-based
shotgun proteomics. *Nat Protoc* **11**, 2301–2319 (2016).

41. J. Illingworth, Methods of enzymatic analysis: Third edition: Editor-in-Chief: Hans Ulrich
Bergmeyer. Verlag Chemie, 1983 (vols I–III), 1984 (vols IV & V) DM258 each volume or DM2240
vols I–X inclusive. *Biochem. Educ.* **13**, 38–38 (1985).

42. K. J. Livak, T. D. Schmittgen, Analysis of relative gene expression data using real-time quantitative

PCR and the 2(-Delta Delta C(T)) Method. *Methods San Diego Calif* **25**, 402–408 (2001).

43. S. Tyanova, T. Temu, P. Sinitcyn, A. Carlson, M. Y. Hein, T. Geiger, M. Mann, J. Cox, The Perseus

computational platform for comprehensive analysis of (prote)omics data. *Nat. Methods* **13**, 731–740

(2016).

44. A. Sharma, G. Naidu, V. Sharma, S. Jha, A. Dhooria, V. Dhir, P. Bhatia, V. Sharma, S. Bhattad, K.

G. Chengappa, V. Gupta, D. P. Misra, P. P. Chavan, S. Malaviya, R. Dudam, B. Sharma, S. Kumar,

R. Bhojwani, P. Gupta, V. Agarwal, K. Sharma, M. Singhal, M. Rathi, R. Nada, R. W. Minz, V.

Chaturvedi, A. Aggarwal, R. Handa, A. Grossi, M. Gattorno, Z. Huang, J. Wang, R. Jois, V. S.

Negi, R. Khubchandani, S. Jain, J. I. Arostegui, E. P. Chambers, M. S. Hershfield, I. Aksentijevich,

Q. Zhou, P. Y. Lee, Deficiency of Adenosine Deaminase 2 in Adults and Children: Experience

From India. *Arthritis Rheumatol. Hoboken NJ* **73**, 276–285 (2021).

45. R. Caorsi, F. Penco, A. Grossi, A. Insalaco, A. Omenetti, M. Alessio, G. Conti, F. Marchetti, P.

Picco, A. Tommasini, S. Martino, C. Malattia, R. Gallizzi, R. A. Podda, A. Salis, F. Falcini, F.

Schena, F. Garbarino, A. Morreale, M. Pardeo, C. Ventrici, C. Passarelli, Q. Zhou, M. Severino, C.

Gandolfo, G. Damonte, A. Martini, A. Ravelli, I. Aksentijevich, I. Ceccherini, M. Gattorno, ADA2

deficiency (DADA2) as an unrecognised cause of early onset polyarteritis nodosa and stroke: a

multicentre national study. *Ann. Rheum. Dis.* **76**, 1648–1656 (2017).

46. J. Schepp, M. Proietti, N. Frede, M. Buchta, K. Hübscher, J. Rojas Restrepo, S. Goldacker, K.

Warnatz, J. Pachlopnik Schmid, A. Duppenthaler, V. Lougaris, I. Uriarte, S. Kelly, M. Hershfield,

B. Grimbacher, Screening of 181 Patients With Antibody Deficiency for Deficiency of Adenosine

Deaminase 2 Sheds New Light on the Disease in Adulthood. *Arthritis Rheumatol. Hoboken NJ* **69**,

1689–1700 (2017).

47. S. Özen, E. D. Batu, E. Z. Taşkıran, H. A. Özkara, Ş. Ünal, N. Güleray, A. Erden, Ö. Karadağ, F.

Gümrük, M. Çetin, H. E. Sönmez, Y. Bilginer, D. Ç. Ayvaz, I. Tezcan, A Monogenic Disease with

a Variety of Phenotypes: Deficiency of Adenosine Deaminase 2. *J. Rheumatol.* **47**, 117–125 (2020).

48. A. Betrains, F. Staels, L. Moens, S. Delafontaine, M. S. Hershfield, D. Blockmans, A. Liston, S.

Humblet-Baron, I. Meyts, R. Schrijvers, S. Vanderschueren, Diagnosis of deficiency of adenosine

deaminase type 2 in adulthood. *Scand. J. Rheumatol.* , 1–4 (2021).

49. O. Akgun-Dogan, P. O. Simsek-Kiper, E. Taskiran, C. Lissewski, J. Brinkmann, D. Schanze, R.

Göçmen, D. Cagdas, Y. Bilginer, G. E. Utine, M. Zenker, S. Ozen, İ. Tezcan, M. Alikasifoglu, K.

Boduroğlu, ADA2 deficiency in a patient with Noonan syndrome-like disorder with loose anagen

hair: The co-occurrence of two rare syndromes. *Am. J. Med. Genet. A.* **179**, 2474–2480 (2019).

50. M. Rama, C. Duflos, I. Melki, D. Bessis, A. Bonhomme, H. Martin, D. Doummar, S. Valence, D.

Rodriguez, E. Carme, D. Genevieve, K. Heimdal, A. Insalaco, N. Franck, V. Queyrel-Moranne,

N. Tieulie, J. London, F. Uettwiller, S. Georgin-Lavialle, A. Belot, I. Koné-Paut, V. Hentgen, G.

Boursier, I. Touitou, G. Sarrabay, A decision tree for the genetic diagnosis of deficiency of

adenosine deaminase 2 (DADA2): a French reference centres experience. *Eur. J. Hum. Genet. EJHG*

26, 960–971 (2018).

51. A. K. Ombrello, J. Qin, P. M. Hoffmann, P. Kumar, D. Stone, A. Jones, T. Romeo, B. Barham,

G. Pinto-Patarroyo, C. Toro, A. Soldatos, Q. Zhou, N. Deuitch, I. Aksentijevich, S. L. Sheldon, S.

Kelly, A. Man, K. Barron, M. Hershfield, W. A. Flegel, D. L. Kastner, Treatment Strategies for

Deficiency of Adenosine Deaminase 2. *N. Engl. J. Med.* **380**, 1582–1584 (2019).

52. K. M. Gibson, K. A. Morishita, P. Dancey, P. Moorehead, B. Drögemöller, X. Han, J. Graham, R.

E. W. Hancock, D. Foell, S. Benseler, R. Luqmani, R. S. M. Yeung, S. Shenoi, M. Bohm, A. M.

Rosenberg, C. J. Ross, D. A. Cabral, K. L. Brown, Identification of Novel Adenosine Deaminase

2 Gene Variants and Varied Clinical Phenotype in Pediatric Vasculitis. *Arthritis Rheumatol. Hoboken*

NJ **71**, 1747–1755 (2019).

1039

Abbreviations

ADA2   Adenosine deaminase 2

BFA   Brefeldin A

DADA2   Deficiency of adenosine deaminase 2

Endo H   Endoglycosidase H

ER   Endoplasmic reticulum

GM-CSF   Granulocyte–macrophage colony-stimulating factor

HC   Healthy control

HMDM   Human monocyte-derived macrophages

HMW   High-molecular-weight

LMW   Low-molecular-weight

M-CSF   Macrophage colony-stimulating factor

PNGase F   Peptide-N-Glycosidase F

TNFi   TNF inhibitors

WT   Wild type

1055

Acknowledgments

We thank Prof. Jaak Jaeken, Prof. Daisy Rymen and Dr. Eva Morava-Kozicz for kindly sharing their

expertise and insight on protein glycosylation. We thank Lotte Bral for her assistance in the production

of the knock-out cell lines. Above all, we would like to express our gratitude to our patients and their

parents for their willingness to participate in our study and to their treating physicians and nurses for

their efforts in the collection of research samples. The figures Fig. 2A, Fig. 3G, Fig. 5C, Fig. 6E, Fig.

7E, Fig. 8A and Fig. S8A were created with BioRender.com.

1063

Funding

This project has received funding from the following sources:

European Research Council (ERC) under the European Union's Horizon 2020 research and

innovation program GA No. 948959 (MORE2ADA2) (LE, MW, BP, MD, IM)

KU Leuven C1 Grant C16/18/007 (IM)

Research Foundation-Flanders (FWO) grant G0B5120N (IM)

Jeffrey Modell Foundation (IM)

ERN-RITA (IM)

Research Foundation – Flanders (FWO) grant 11E0123N (LE)

Charité – Universitätsmedizin Berlin and the Berlin Institute of Health BIH Charité Junior Clinician

Scientist Fellowship (LE)

Research Foundation – Flanders (FWO) grant 11F4421N (SD)

Research Foundation – Flanders (FWO) grant 1805523N (RS)

Research Foundation – Flanders (FWO) grant G0A3320N (PA)

KU Leuven C14/21/095 InterAction consortium (PA)

EOS MetaNiche consortium N° 40007532 (PA)

iBOF/21/053 ATLANTIS network (PA)

EOS DECODE consortium N° 30837538 (PA)

1082

Author contributions

Conceptualization and design: LE, LM, IM

Material preparation and data collection: LE, AH, MW, BP, SD, MD, EFR, AD, JN, MK, LM

Data analysis: LE, AH, MB, MK, LM

Generation of ADA2 knock-out cell lines: LE, MJ, DD

Proteomics analyses: MK, PM, MFM, TM

Design of trafficking experiments: LE, LM, IM, FE

Design of ER stress experiments: LE; LM, IM, PA

Collection of clinical data and patient materials: LE, SD, GB, LDS, RS, SV, DC, IM

Supervision: IM

Writing – original draft: LE

Writing – review & editing: all authors

Competing interests

IM receives research funding from CSL-Behring outside this project. Other grant support was received from Octapharma, as well as Advisory Board honoraries from Takeda and Boehringer-Ingelheim - paid to the institution and outside the submitted work.

Data and materials availability

All data needed to evaluate the conclusions of our study are provided in the manuscript or the Supplementary Materials. The mass spectrometry data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository with the dataset identifier PXD046128. The authors provide a supplementary file containing the original blots of all the presented data.

Figures

Figure 1

 Fig. 1: Expression of pathogenic ADA2 protein variants. (A) mRNA and intracellular protein

 expression of *ADA2* in CD14⁺ monocytes from healthy controls (HC) and DADA2 patients (P) by

 qPCR or bulk RNA-sequencing and western blot of whole cell lysates, respectively. The left plot shows

mRNA expression determined by qPCR. *ADA2* mRNA expression was normalized to *GAPDH* and

is depicted relative to the mean of HC samples. The right plot shows counts determined by RNA-

sequencing. Dot plots show median and interquartile range. **(B)** ADA2 protein expression in whole

cell lysates and supernatant (SN) of CD14$^+$ monocytes and GM-CSF differentiated human monocyte-

derived macrophages (HMDM) from HC, carriers (CR) and DADA2 patients (P) was determined by

western blot. **(C)** ADA2 protein expression in whole cell lysates of transduced U-937 cells and HMDM

from HC and DADA2 patients (P) was determined by western blot. **(D)** ADA2 protein expression in

whole cell lysates of HEK293T cells after transfection with wild-type ADA2 and different pathogenic

variants (*left*) and comparing whole cell lysates (WCE) and supernatant (SN) by western blot (*right*). **(E)**

ADA2 protein expression in supernatants (SN) from healthy control human monocyte-derived

macrophages (HMDM) and serum samples from healthy controls (HC) and DADA2 patients (P) by

western blot using VeriBlot for detection. ADA2 detection using anti-ADA2 antibody clone

EPR25430-131 (#ab288296; abcam) in all depicted western blots. Triangles indicate HMW-ADA2

(dark blue) and LMW-ADA2 (light blue). Legend: Ig, immunoglobulin; WT, wild type.

1126

Figure 2

Fig. 2: N-glycosylation of ADA2. (A) Schematic illustration of inhibition of glycosylation by

tunicamycin and glycan removal by PNGase F. **(B)** Western blot of wild-type (WT) ADA2 and the

pathogenic variants p.R169Q and p.T129P transfected into HEK293T cells after incubation with or

without 2.5 µg/mL tunicamycin for 24 h and/or glycan removal by PNGase F. **(C)** ADA2 protein

expression by western blot in whole cell lysates (WCE) and supernatants (SN) of human monocyte-

derived macrophages (HMDM), U-937 cells and THP-1 cells. **(D)** ADA2 protein expression in whole

cell lysates and supernatants (SN) from GM-CSF-differentiated HMDM of healthy control HC8 and

DADA2 patient P8 left untreated or treated with 5 μg/mL tunicamycin or 1 μg/mL brefeldin A for

24 h. **(E)** ADA2 protein expression in whole cell lysates and supernatants (SN) from U-937 and THP-

1 cells left untreated or treated with 2.5 μg/mL tunicamycin or 1 μg/mL brefeldin A for 24 h. Glycan

removal was performed by incubation with PNGase F for 1 hour at 37°C under denaturing conditions

in all experiments. ADA2 detection using anti-ADA2 (#HPA007888, Sigma-Aldrich) in all depicted

western blots. Triangles indicate HMW-ADA2 (dark blue), LMW-ADA2 (light blue) and glycan-free

ADA2 (black).

1143

Figure 3

Fig. 3: Glycan processing of ADA2 in the ER. (A) ADA2 protein expression by western blot in

GM-CSF-differentiated HMDM from healthy control HC4 and DADA2 patient P4 after incubation

with 100 μg/mL castanospermine for 24 h. **(B)** ADA2 protein expression upon inhibition of N-

glycosylation, ER glucosidases I and II, or ER to Golgi trafficking. U-937, THP-1 and Jurkat cells were

incubated with 2.5 μg/mL tunicamycin (TM), 100 μg/mL castanospermine (CAS) or 1 μg/mL

brefeldin A (BFA) for 24 h or 48 h. **(C)** HEK293T cells were transfected with wild-type (WT) ADA2

or the pathogenic variant p.R169Q and treated with 500 μg/mL cycloheximide (CHX) or 2.5 μg/mL

tunicamycin (TM) over 24-48 h. Whole cell lysates were generated at the indicated time points and

immunoblotted for ADA2 expression. **(D)** ADA2 protein levels were determined in supernatants of

M-CSF-differentiated human monocyte-derived macrophages (HMDM) from healthy controls (HC)

or DADA2 patients (P) or HEK293T cells transfected with wild-type (WT) *ADA2* or different

pathogenic *ADA2* variants. The patients' genotypes are indicated above. Western blot of serum

samples from healthy controls (HC), heterozygous DADA2 carriers (CR) and DADA2 patients (P)

detected by anti-ADA2 antibody clone EPR25430-131 (#ab288296; abcam) and VeriBlot.

(E) HEK293T cells were transfected with wild-type (WT) ADA2 or different pathogenic variants and

treated with 500 μg/mL cycloheximide (CHX) ± 1 μg/mL brefeldin A or 2 μg/mL monensin over

24 h. Whole cell lysates were generated at the indicated time points and immunoblotted for ADA2

expression. **(F)** ADA2 protein expression in GM-CSF-differentiated human monocyte-derived

macrophages from healthy controls (HC) or DADA2 patient P4 left untreated or treated with 1 μg/mL

brefeldin A (BFA) or 2 μg/mL monensin (MN). **(G)** Schematic overview of trafficking and processing

of ADA2 in healthy control and DADA2 cells. ADA2 detection using anti-ADA2 antibody clone

EPR25430-131 (#ab288296; abcam) in panels A, B, C (*left*), D, E and F and using anti-ADA2

(#HPA007888, Sigma-Aldrich) in panel C (*right*). Triangles indicate HMW-ADA2 (dark blue), LMW-

ADA2 (light blue) and glycan-free ADA2 (black). Legend: BFA, brefeldin A, CAS, castanospermine,

EV, empty vector; MN, monensin, MWM, molecular weight marker; TM, tunicamycin; WT, wild type.

1171

Figure 4

Fig. 4: ER stress in DADA2. (A) Interactome analysis of wild-type (WT) ADA2 and the pathogenic

variant p.R169Q overexpressed in HEK293T cells treated with or without 2.5 μg/mL tunicamycin

(TM) for 24 h was performed by immunoprecipitation followed by mass spectrometry. Heatmap

shows z-score of protein abundance. Binders significantly pulled down compared to isotype control

with a false discovery rate of 5% are highlighted by colored circles. Localization of the interacting

proteins to the ER/Golgi (blue) or lysosome (green) is indicated by color. **(B)** ER stress response in

HEK293T cells transfected with wild-type (WT) *ADA2* and different pathogenic variants. Protein

expression of BiP (*top*) and mRNA expression of *HSPA5* and *DDIT3* (*bottom*) were determined 48h

after transfection. A representative western blot is shown in **Fig. S5F**. mRNA expression was

normalized to *GAPDH* and is depicted as $2^{-\Delta\Delta Ct}$ relative to non-transfected control samples. Results

from 6 independent experiments are shown. Dot plots show median. **(C)** ER stress response was

evaluated by protein expression of BiP in whole cell lysates of CD14$^+$ monocytes at baseline or after

24 h incubation with or without 1 μg/mL tunicamycin (TM). BiP expression was normalized to β-

actin. Dot plots show median and interquartile range. The corresponding western blots are shown in

Fig. S6. **(D)** Formation of intracellular ADA2 dimers in HEK293T cells transfected with wild-type

(WT) ADA2 and different pathogenic *ADA2* variants. The blot is equivalent to the one shown in

Fig. 1D. **(E)** ADA2 protein expression and intracellular dimer formation in GM-CSF differentiated

human monocyte-derived macrophages from healthy controls (HC) and DADA2 patients (P). ADA2

detection using anti-ADA2 antibody clone EPR25430-131 (#ab288296; abcam) in all depicted western

blots. Legend: EV, empty vector; FC, fold change; NT, non-transfected; WT, wild type.

1194

1195 **Figure 5**

1196

Fig. 5: Glycan processing of HMW(c)-ADA2. ADA2 protein expression by western blot after

glycan removal in ADA2$^{WT/WT}$ and ADA2$^{-/-}$ U-937 cells transduced with pathogenic *ADA2* variants

(A) and CD14$^+$ monocytes or GM-CSF-differentiated HMDM from healthy controls (HC) and

DADA2 patient P4 **(B)** by incubation with PNGase F or Endo H for 1 hour at 37 °C under denaturing

conditions. **(C)** Cutting sites of PNGase F and Endo H. Endo H does not cleave complex glycans

(top). Carbohydrate residues are depicted according to the Symbol Nomenclature for Glycans (SNFG)

(20). ADA2 detection using anti-ADA2 antibody clone EPR25430-131 (#ab288296; abcam) in panel

B (*right*) and using anti-ADA2 (#HPA007888, Sigma-Aldrich) in panels A and B (*left*). Triangles indicate

HMW-ADA2 (dark blue), LMW-ADA2 (light blue) and glycan-free ADA2 (black).

1206

Figure 6

Fig. 6: Glycan processing of LMW-ADA2. (A) α-mannosidases were inhibited by 72-96 h

incubation of ADA2$^{WT/WT}$ and ADA2$^{-/-}$ U-937 cells transduced with R169Q with 375 nM kifunensine

and/or 10 μM swainsonine. ADA2 protein expression was determined by western blot. The

histograms indicate molecular weight peaks as determined by Simple Western. The original graphs are

provided in **Fig. S7**. **(B)** ADA2 protein expression by western blot GM-CSF- and M-CSF-

differentiated HMDM from a healthy control (HC) and a patient with α-mannosidosis (α-man) after

incubation with 10 μM swainsonine (SW) for 96 h. **(C)** ADA2 protein expression by western blot after

glycan removal in ADA2$^{WT/WT}$ U-937 cells by incubation with PNGase F, Endo H or α1-2,3,6

Mannosidase for 1 hour at 37 °C. **(D)** Cleavage sites of PNGase F, Endo H and α-Mannosidase. **(E)**

Schematic overview of the trafficking and glycan modifications of ADA2 expressed in healthy control

(*left*) and DADA2 (*right*) human monocyte-derived macrophages based on the findings presented in

this article. *MOGS* encodes ER α-glucosidase I, *GANAB* encodes ER α-glucosidase II, *MAN1B1*

encodes ER α-1,2-mannosidase, *MAN1A1*, *MAN1A2* and *MAN1C1* encode Golgi α-1,2-

mannosidases, *MAN2A1* encodes Golgi α-mannosidase II, and *MAN2B1* encodes lysosomal α-

mannosidase. Mechanisms of action of the inhibitors used for the *in vitro* study of glycan processing

are depicted in orange. Carbohydrate residues are depicted according to the Symbol Nomenclature for

Glycans (SNFG) *(20)*. ADA2 detection using anti-ADA2 antibody clone EPR25430-131 (#ab288296;

abcam) in panels A, B and C (*top*) and using anti-ADA2 (#HPA007888, Sigma-Aldrich) in panel C

(*bottom*). Triangles indicate HMW-ADA2 (dark blue), LMW-ADA2 (light blue) and glycan-free ADA2

(black). Legend: ECS, extracellular space; ER, endoplasmic reticulum.

1229

Figure 7

Fig. 7: Localization and generation of LMW-ADA2. (A) Immunofluorescence microscopy of

ADA2$^{WT/WT}$ U-937 cells and ADA2$^{-/-}$ U-937 transduced with the pathogenic variant p.R169Q stained

for ADA2 and markers of lysosomal (LAMP1) and Golgi (GM130) compartments. **(B)** ADA2 protein

expression in cytosolic, membranous and lysosomal fractions of U-937 cells by western blot. **(C)**

Intracellular ADA2 uptake into ADA2$^{-/-}$ U-937 cells after incubation with supernatants (SN) from

HEK293T cells transfected with empty vector (EV) or wild-type (WT) *ADA2* for the indicated time.

(D) Generation of intracellular LMW-ADA2 in ADA2$^{WT/WT}$ or ADA2$^{-/-}$ U-937 cells after incubation

with supernatants (SN) from HEK293T cells transfected with empty vector (EV), wild-type (WT)

ADA2 or the pathogenic variant p.G47A for the indicated time. **(E)** Schematic illustration of uptake

and processing of exogenous ADA2. ADA2 detection using anti-ADA2 antibody clone EPR25430-

131 (#ab288296; abcam) in panels C and D and using anti-ADA2 (#HPA007888, Sigma-Aldrich) in

panel B. Triangles indicate HMW-ADA2 (dark blue), LMW-ADA2 (light blue) and glycan-free ADA2

(black). Legend: cl, clone.

1245

Figure 8

Fig. 8: Validation of absent LMW-ADA2 as a pathogenicity marker of *ADA2* variants. (A) Distribution of all evaluated ADA2 variants (red lines) across the ADA2 protein domains. The N-glycosylation sites are indicated in blue. **(B)** ADA2 secretion and LMW-ADA2 expression in HEK293T cells transfected with *ADA2* variants were determined by western blot using anti-ADA2 antibody clone EPR25430-131 (#ab288296; abcam). Enzyme activity was measured in whole cell lysates and supernatants of the transfected cells. Dot plots show median. Each variant was examined in n=3-4 independent experiments. **(C)** Correlation of LMW-ADA2 and intracellular or extracellular ADA2 enzyme activity of *ADA2* variants expressed in HEK293T cells. The dots show median of n=3-4 independent experiments. The dotted lines indicate 25% expression/activity of WT ADA2. The blue line represents the fitted linear regression line with its 95% confidence interval shaded in light blue. **(D)** Correlation of LMW-ADA2 expression and ADA2 enzyme activity with different pathogenicity scores. Red shading indicates (likely) pathogenic scores while grey shading represents ambiguous scores. The blue line represents the fitted linear regression line with its 95% confidence interval shaded in light blue. The corresponding Pearson correlation coefficients are shown in Table S3. Legend: CD, catalytic domain; DD, dimerization domain; EV, empty vector; NT, non-transfected; PRB D, putative receptor binding domain; SP, signal peptide; WT, wild type.

 Tables

ID	Age [years]	Sex	Genotype	Predominant phenotype
P1	12	M	c.973-2A>G/c.del1240-1442; splice site (intron 6)/del exon 9	Vasculitis, hepatosplenomegaly, hypogammaglobulinemia
P2	14	F	c.973-2A>G/c.del1240-1442; splice site (intron 6)/del exon 9	Vasculitis, warts, arthralgia, hypo-gammaglobulinemia,
P3	26	M	c.140G>T/c.del1240-1442; p.G47V/del exon 9	Bone marrow failure, hypogam-maglobulinemia, hepatosplenomegaly
P4	26	M	c.140G>T/c.140G>T; p.G47V/p.G47V	Hypogammaglobulinemia, hepatosplenomegaly
P5	6	F	c.140G>T/c.506G>A; p.G47V/p.R169Q	Hypogammaglobulinemia, hepatosplenomegaly, stroke
P6	9	F	c.140G>T/c.506G>A; p.G47V/p.R169Q	Bone marrow failure, hypogam-maglobulinemia, hepatosplenomegaly
P7	32	F	c.973-2A>G/c.506G>A; splice site (intron 6)/p.R169Q	Stroke, hypertension
P8	16	M	c.139G>A/c.139G>A; p.G47R/p.G47R	Stroke, vasculitis

P9	10	F	c.973-2A>G/c.973-2A>G; splice site (intron 6)/splice site (intron 6)	Vasculitis, hypogammaglobulinemia
P10	50	F	c.973-2A>G/c.506G>A; splice site (intron 6)/p.R169Q	Vasculitis, leukopenia
CR1	39	F	WT/c. del1240-1442; WT/del exon 9	
CR2	38	F	WT/c.140G>T; WT/p.G47V	
CR3	42	M	WT/c.506G>A; WT/p.R169Q	

1266 **Table 1: Cohort characteristics.** Genotype and clinical phenotype of the included DADA2 patients

1267 (P) and heterozygous carriers (CR).

Protein	LMW-ADA2	ADA2 enzyme activity	REVEL	CADD	PolyPhen	SIFT	Alpha missense	Infevers	ClinVar	Homo-zygous patients	Hetero-zygous patients	Homozygous individuals in gnomAD
p.G25C	81.8	64.0	0.076	0.498	benign (0)	tolerated (0.61)	likely benign (0.0748)	likely pathogenic	uncertain significance	2	0	-
p.H91R	91.6	90.5	0.248	0.086	benign (0.001)	tolerated (0.38)	likely benign (0.0686)	not classified	uncertain significance	1	0	2
p.H112Q	63.9	**1.1**	**0.956**	**19.6**	**probably damaging (0.987)**	**deleterious (0)**	likely pathogenic (0.9015)	pathogenic	pathogenic/likely pathogenic	1	11	-
p.L188V	61.8	45.7	**0.657**	**21.8**	**possibly damaging (0.841)**	**deleterious (0)**	likely benign (0.1571)	likely pathogenic	conflicting	0	2	2
p.R230Q	129.2	89.3	0.232	0.437	benign (0.007)	tolerated (1)	likely benign (0.0611)	not classified	likely benign	0	0	2
p.D329N	98.7	67.3	**0.694**	**25.7**	**probably damaging (1)**	**deleterious (0.02)**	likely benign (0.1525)	likely pathogenic	uncertain significance	1	0	-

p.H335P	82.4	86.4	0.175	**11.27**	benign (0.012)	**deleterious (0)**	likely benign (0.1858)	not classified	uncertain significance	0	1	-
p.H335R	138.8	99.6	0.176	8.752	benign (0.015)	**deleterious (0.04)**	likely benign (0.0727)	benign	benign	0	0	94070
p.P344L	**0**	36.3	0.52	**22.3**	**probably damaging (0.979)**	**deleterious (0)**	ambiguous (0.4429)	likely pathogenic	NA	0	3	-
p.F355L	108.1	90.5	0.502	**17.34**	benign (0.25)	**deleterious (0.04)**	likely pathogenic (0.6625)	likely pathogenic	conflicting	0	2	-
p.A357T	44.5	35.5	**0.919**	**23.8**	**probably damaging (0.993)**	**deleterious (0)**	likely benign (0.2518)	likely pathogenic	conflicting	1	3	-
p.T360A	33.8	**10.0**	**0.774**	**22.8**	**probably damaging (0.994)**	**deleterious (0)**	ambiguous (0.465)	likely pathogenic	pathogenic	10	15	-
p.P435A	66.6	32.0	**0.706**	**22.8**	**probably damaging (0.997)**	**deleterious (0)**	likely benign (0.1426)	likely pathogenic	likely benign	0	2	-

Table 2: Variant characteristics. Classification of *ADA2* variants exhibiting inconclusive (> 25% of WT) levels of LMW-ADA2 expression or ADA2 enzyme activity. LMW-ADA2 expression and extracellular ADA2 enzyme activity are displayed as % of WT ADA2.