A Reinterpretation of the Relationship Between Persistent and Resurgent Sodium Currents

Condensed Title: Na$^+$ currents role in repetitive neuronal firing

Samuel P. Brown[1], B.S.; Ryan J. Lawson[1], B.S.; Jonathan D. Moreno, M.D., Ph.D.[2]; Joseph L.
Ransdell[1,4], Ph.D.

[1]Department of Biology
Miami University, Oxford, OH 45056

[2]Division of Cardiology, Department of Medicine
Washington University in St. Louis, St. Louis, MO 63130

[4]Correspondence: ransdejl@miamioh.edu

Number of Figures: 8

16   Tables: 2

Supp. Figures: 4

Abstract: 151 words

Introduction: 587 words

Discussion: 1694 words

Declaration of interests

The authors declare no competing interests.

Acknowledgments

This work was supported by startup funds from the Miami University College of Arts and Sciences
and by the NINDS at the National Institutes of Health: Award 1R15NS125560

Abstract

The resurgent sodium current (I_{NaR}) activates on membrane repolarization, such as during the downstroke of neuronal action potentials. Due to its unique activation properties, I_{NaR} is thought to drive high rates of repetitive neuronal firing. However, I_{NaR} is often studied in combination with the persistent or non-inactivating portion of sodium currents (I_{NaP}). We used dynamic clamp to test how I_{NaR} and I_{NaP} individually affect repetitive firing in adult cerebellar Purkinje neurons. We learned I_{NaR} does not scale repetitive firing rates due to its rapid decay at subthreshold voltages, and that subthreshold I_{NaP} is critical in regulating neuronal firing rate. Adjustments to the Nav conductance model used in these studies revealed I_{NaP} and I_{NaR} can be inversely scaled by adjusting occupancy in the slow inactivated kinetic state. Together with additional dynamic clamp experiments, these data suggest the regulation of sodium channel slow inactivation can fine-tune I_{NaP} and Purkinje neuron repetitive firing rates.

Significance Statement

Across neuronal cell types, the resurgent sodium current (I_{NaR}) is often implicated in driving high rates of repetitive firing. Using dynamic clamp, we determined I_{NaR} is ineffective at driving subsequent action potentials, and that the subthreshold persistent sodium current is the critical parameter for scaling repetitive firing rates. We propose I_{NaR} measured in native neurons may reflect a mechanism by which the magnitude of I_{NaP} is fine-tuned. We propose I_{NaR} measured in native neurons may reflect a mechanism by which the magnitude of I_{NaP} is fine-tuned.

Introduction

Purkinje neurons are the exclusive output cells of the cerebellar cortex, delivering GABA-mediated inhibitory signals to neurons in the deep cerebellar nuclei (Ito, 1984; Ito et al., 1964; Obata et al., 1967, 1970). In the absence of synaptic drive, Purkinje neurons have a unique capacity to fire action potentials repetitively at high frequencies (Häusser & Clark, 1997; Raman & Bean, 1997; Thach, 1968), properties which rely on voltage-gated sodium (Nav) channels and the depolarizing currents these channels mediate (Llinás & Sugimori, 1980; Raman & Bean, 1999; Ransdell & Nerbonne, 2018). Beyond the fast-inactivating transient sodium current (I_{NaT}), Nav channels can generate a persistent sodium current (I_{NaP}) component, which was first measured and described in cerebellar Purkinje neurons (Llinás & Sugimori, 1980). Nav channels may also generate a resurgent Nav current (I_{NaR}) component, also first characterized in cerebellar Purkinje neurons, which has now been measured in over twenty neuronal cell types throughout the central and peripheral nervous systems (Lewis & Raman, 2014; Raman & Bean, 1997). I_{NaR} is distinct from the transient and persistent Nav currents in that it activates on membrane repolarization and involves the recovery of fast-inactivated Nav channels into an open/conducting state (see Figure 1A). In the context of neuronal firing, this type of repolarization would occur during the downstroke of the action potential.

On its discovery, I_{NaR} was proposed to support high rates of neuronal firing by reducing the Nav channel refractory period (Khaliq et al., 2003; Raman & Bean, 1997, 1999), and this assumption continues to be widely accepted (Quattrocolo et al., 2021; Raman, 2023). During an action potential repolarization, if a subset of Nav channels recover into the open state allowing resurgent Na$^+$ influx, the depolarizing current should drive subsequent action potentials. After its discovery, perturbations to I_{NaR} were also associated with changes in repetitive firing (Khaliq et al., 2003; Lewis & Raman, 2014; Raman et al., 1997; Ransdell et al., 2017). For instance, targeted knockout of Navβ4, a sodium channel accessory subunit, was found to result in significantly reduced I_{NaR}

in mouse Purkinje neurons (Ransdell et al., 2017) as well as striatal medium spiny neurons

(Miyazaki et al., 2014). The loss of Navβ4 was additionally associated with reduced repetitive

firing rates compared to wild type controls in these cell types (Miyazaki et al., 2014; Ransdell et

al., 2017, 2022). The isolated contribution of I_{NaR} to neuronal repetitive firing, however, has not

been directly tested in native neurons.

We use dynamic clamp and multiple (10) Nav conductance models to systematically assess how

adjusting properties of I_{NaR} and I_{NaP} affect the firing properties of mouse cerebellar Purkinje

neurons. We learned that scaling the proportion and/or kinetic properties of I_{NaR} does not

significantly affect Purkinje neuron firing frequency, nor does the addition of I_{NaR} meaningfully

impact the action potential waveform in Purkinje neurons. Using voltage-clamp simulation studies,

we determined I_{NaR} does not influence repetitive firing rates due to its rapid decay at subthreshold

voltages. Subthreshold I_{NaP}, however, was found to be a critical parameter in tuning Purkinje

neuron repetitive firing rates. In examining the Nav Markov models, we determined peak I_{NaR} and

I_{NaP} can be inversely scaled by adjusting occupancy in the slow-inactivated kinetic state, an

inactivation pathway which is distinct from the conventional fast-inactivation pathway. Combined,

these data suggest I_{NaR} may reflect a population of Nav channels which would, in the absence of

slow inactivation, contribute to I_{NaP}; suggesting Nav channel slow-inactivation may function to fine

tune the amount of Nav channels that contribute to I_{NaP} in order to regulate repetitive firing rates.

Methods

Animals

All animal experiments were performed in accordance with protocols approved by Miami

University Institutional Animal Care and Use Committee (IACUC) guidelines. All experiments

utilized C57BL/6J wild type mice 4-8 weeks old.

Preparation of acute cerebellar slices

Mice were anaesthetized using an intraperitoneal injection of 1mL/kg ketamine (10

mg/ml)/xylazine (0.25 mg/ml) cocktail and perfused transcardially with 25 ml cutting solution

containing (in mM) 240 sucrose, 2.5 KCl, 1.25 NaH_2PO_4, 25 $NaHCO_3$, 0.5 $CaCl_2$, and 7 $MgCl_2$.

The brain was rapidly dissected, superglued to a specimen tube, and submerged in warmed

agarose dissolved in cutting solution. The cerebellum was sliced sagittally in ice-cold cutting

solution saturated with 95% O_2/5% CO_2 using a Compresstome VF-300 (Precisionary

Instruments) at 350 µM thickness. Slices were placed in artificial cerebrospinal fluid (ACSF)

containing (in mM) 125 NaCl, 2.5 KCl, 1.25 NaH_2PO_4, 25 $NaHCO_3$, 2 $CaCl_2$, 1 $MgCl_2$, and 25

dextrose (pH 7.4, ~300 mOsm) on stretched nylon for 25 minutes at 33°C and then at room

temperature for at least 35 minutes before electrophysiological recordings. Purkinje neurons were

identified in the Purkinje neuron layer of sagittal cerebellar sections using a SliceScope Pro 3000

(Scientifica).

Electrophysiological recordings

Whole-cell current-clamp and dynamic clamp recordings were obtained from adult mouse

Purkinje neurons (4-8 weeks old) in acutely isolated sagittal cerebellar slices that were constantly

perfused with ACSF warmed to ~34°C and continuously bubbled with 95% O_2/5% CO_2. Recording

pipettes (borosilicate standard wall, 1.5 mm outer diameter, 0.86 mm inner diameter) were made

using a P-1000 Flaming/Brown Micropipette Puller (Sutter Instruments) and filled with internal

solution containing (in mM) 0.2 EGTA, 3 $MgCl_2$, 10 HEPES, 8 NaCl, 4 Mg-ATP, and 0.5 Na-GTP.

Patch pipette resistances were 2-4 MΩ. Prior to patching, a junction potential correction of 17.5

mV was applied and pipette tip potentials were zeroed. After a gigaseal was obtained,

spontaneous action potentials were recorded using a dPatch Amplifier and SutterPatch software

(Sutter Instruments). Input resistance and capacitance calculations were obtained through a

whole-cell voltage clamp protocol in which the membrane voltage was stepped from a -80 mV

holding potential to -90 mV (for 100 ms) and to -70 mV (for 100 ms) in a second sweep. This

voltage-clamp protocol was applied in each cell prior to current-clamp or dynamic clamp

measurements. For membrane capacitance, the integrated area of the capacitive transient current

was divided by the change in membrane voltage (10 mV) during each sweep. Action potential

properties were analyzed using SutterPatch software (Sutter Instruments). Action potential events

were detected if a transient change in membrane potential exceeded 20 mV. Threshold potential

was calculated in SutterPatch as the membrane potential prior to an action potential event when

the dV/dt is >1 mV/100 μs, or when 25% of the maximum action potential dV/dt is reached,

whichever is smaller. Action potential duration was calculated as the time interval from the pass

of threshold voltage during action potential upstroke until the pass of threshold voltage during

action potential downstroke. Each action potential measurement from a given cell was recorded

as the average of the measurement across 20 spontaneous action potentials. Dynamic clamp

current injection values, measured during the interspike interval (presented in Figures 5D, 6B),

were obtained by measuring the dynamic clamp current injection value 3 ms before the action

potential threshold was reached for the subsequent action potential. These measurements were

made 1 second into each dynamic clamp recording. Recordings were acquired and analyzed

using SutterPatch (Sutter Instruments), Microsoft Excel (Microsoft), and Prism (GraphPad)

software.

Markov modeling

A previously developed Markov kinetic state model of Nav channel gating (Ransdell et al., 2022)

was used as the base/control model, and is referred to as the 'nominal model'. To develop the

'I_{NaR} increased', 'I_{NaR} decreased', 'τdecay increased', 'τdecay decreased', 'I_{NaR} reduced', and 'I_{NaP}

reduced' models (Figures 2-4), rate constants in the nominal model were altered to cause

selective changes in Nav current properties. Each of these models consist of a nine state Markov

matrix that includes three closed states (C1, C2, C3), two inactivated-closed states (IC1, IC2),

two fast inactivated states (IF1, IF2), a slow inactivated state (IS), and an open state (O). The

hand-tuned 'I_{NaR} increased' and 'I_{NaR} decreased' models, presented in Figure 2, were generated

by altering rate constant equations (see Table 1) which resulted in voltage dependent rate

constant alterations. For instance, the b3s rate constant changed from 0.94767 to 3.76696 at -45

mV in the 'I_{NaR} increased' model and in the 'I_{NaR} decreased' model, the b2 rate constant changed

from 0.1081 to 3.884 at -45 mV. In order to generate additional models, properties of the originally

developed (nominal) model were modified by adjusting values in a parameter optimization

algorithm scripted in MATLAB (MathWorks) and previously described in (Moreno et al., 2016;

Ransdell et al., 2022). The rate constant values for the nominal and each manipulated model are

described in Table 1. To generate the 'I_{NaR} reduced' model (Figure 4), state transition rate

calculations a3s, b3s, a2s, and b2s were removed from the Q_Matrix.m script prior to parameter

optimization. To generate the 'I_{NaP} reduced' model (Figure 4), the variable (ResT_line) within the

Na_Matrix_DrugFree.m script was changed from 0.18678*V + 22.92 to 0.18678*V + 38.00.

Additionally, in line 113 the Err_Late minimization variable was multiplied by 200. To generate the

'τ_{decay} decreased' model, the ResT_line variable was changed to 0.18678*V + 38.00. Both

Err_ACT and Err_SSA variables in line 113 of the error minimization equation (part of the

parameter optimization script) were also multiplied by a value of 10. To generate the 'τ_{decay}

increased' model, the ResT_line variable was changed to 0.18678*V + 15.00 and the error

minimization variables Err_ACT and Err_SSA were multiplied by 90.

With the above alterations, each model was generated by running a minimization routine using a

Nelder-Mead "amoeba" algorithm-based optimization procedure (Lagarias et al., 1998; Moreno et

al., 2016; Nelder & Mead, 1965) to determine the final transition rate constants associated with

each manipulated model (presented in Table 1). All models were simulated using a sodium

reversal potential of 71.5 mV. All kinetic state occupancy simulations and optimizations were

completed in MATLAB 2022a. Simulations and current measurements from each model were

performed in SutterPatch using a model cell (Sutter Instruments) to verify the Nav currents

generated by each model (I_{NaT}, I_{NaR}, and I_{NaP}) had appropriate voltage-dependent and kinetic

properties. Currents were evoked in each model using simulated voltage clamp protocols applied

to the model cell while the simulated conductance was applied via dynamic clamp software

(Sutterpatch) using a Sutter Instruments dPatch amplifier. The voltage dependance of activation

of I_{NaT} and I_{NaP} were obtained by stepping the model cell (from a -90 mV holding potential) to -70

mV – 0 mV in 5 mV increments for 80 ms. I_{NaR} voltage-dependence of activation was obtained by

stepping the model cell from a holding potential of -90 mV to 0 mV for 5 ms, and subsequently

applying a repolarizing step to -70 mV - 0mV (in 5 mV increments) for 80 ms. Prior to analysis of

I_{NaT} or I_{NaR}, I_{NaP} was digitally subtracted. MATLAB scripts used for the optimization procedure and

the   nominal   Markov   state   matrix   are   available   on   GitHub   at

https://github.com/morenomdphd/Resurgent_INa.

Action potential clamp gating simulations

To visualize gating state probability over time with an action potential as the voltage command,

the MATLAB script Fig_4H_4I.m was adjusted (Ransdell et al., 2022). Prior to plotting, the

parameter optimization procedure (see previous section) was run to populate the MATLAB

workspace with the required variables. An action potential recording, captured from an adult

mouse Purkinje neuron firing spontaneous action potentials (at 60.8 Hz) was exported to a

Microsoft Excel spreadsheet that contained the time and voltage values of the action potential

record. The AP-clamp script was utilized as follows: The script loaded the excel file, which was

iterated through the Q_Matrix.m script using the voltage values as the voltage command and the

time values as the length of the protocol. Through this script, the values for each gating state as

well as I_{Na} and the voltage command were written to the NASIM matrix and then were plotted over

time.

Hodgkin-Huxley modeling

A previously developed Hodgkin-Huxley (HH) model of I_{NaP} in Golgi cells was adopted for dynamic

clamp experiments (Solinas et al., 2007) and is available on SenseLab ModelDB (ID: 112685).

The conductance value of 3.4 nS was selected for this model because peak HH-generated I_{NaP}

values were similar to the peak I_{NaP} from nominal model-generated values at the 400 nS

conductance used in other dynamic clamp experiments. The alpha and beta equations for the HH

I_{NaP} model are presented below.

α	β
5/((0.0421*(V+50)/(1-exp(-1*(V+50)/5)))+(-0.287*(V+50)/(1-exp((V+50)/5))))	1/(1+exp(-1*(V+53)/5))

Dynamic clamp

Model conductances were applied to whole-cell-patched adult mouse Purkinje neurons through

SutterPatch software and a dPatch amplifier. Markov model conductances were applied in

dynamic clamp experiments using a sodium reversal potential of 71.5 mV and a conductance

value of 400 nS. The Hodgkin Huxley model, described above, was applied using the same

reversal potential and a 3.4 nS conductance. In all dynamic clamp experiments, a model

conductance was only applied after a gap-free current clamp recording (10 second duration) of

action potential firing (with dynamic clamp turned off). These intermittent recordings (between

applications of model Nav conductances) served as control measurements of Purkinje neuron

firing rate without the addition of a model Nav conductance. In all cells included in these analyses,

baseline repetitive firing frequencies, with no dynamic clamp current injection, did not vary by

more than 10% throughout the duration of the experiment.

Results

Dynamic clamp addition of a modeled Nav conductance increases the spontaneous firing

frequency of adult cerebellar Purkinje neurons

Purkinje neuron voltage-gated sodium (Nav) currents include fast-transient (I_{NaT}), persistent (I_{NaP}),

and resurgent (I_{NaR}) components, which can be isolated and measured using steady state voltage-

clamp steps. Voltage-clamp records presented in Figure 1A (*middle traces*) were acquired from

an acutely isolated Purkinje neuron (P16) and reflect the I_{NaT} and I_{NaR} components. A brief

depolarizing voltage step to 0 mV (from a -90 mV holding potential) activate Nav channels, the

majority of which quickly inactivate resulting in the transient Nav current (I_{NaT}), and a repolarization

voltage step to more negative potentials enable a portion of the fast-inactivated Nav channels to

recovery into an open/conducting state, which results in 'resurgent' sodium influx (I_{NaR}). I_{NaR} (at

fixed voltages) decays much slower than I_{NaT}. At voltages that enable peak I_{NaR} activation (typically

around -45 mV), the time constant of I_{NaR} decay is ~12 ms, and at more hyperpolarized voltages,

the time constant of I_{NaR} decay falls to 2-3 ms (Raman & Bean, 1997; Zemel et al., 2021). I_{NaR}

decay eventually results in a steady-state depolarizing Nav current in which a population of Nav

channels remain in the open/conducting state. This remaining persistent current component

reflects I_{NaP}. I_{NaR} was first measured in cerebellar Purkinje neurons (Raman & Bean, 1997) and

because I_{NaR} is a depolarizing current activated by membrane repolarization, such as the

downstroke of an action potential, this depolarizing Nav current is hypothesized to be associated

with supporting high rates of repetitive firing. To test this role directly, we used a Markov kinetic

state model (depicted in Figure 1B), which accurately reproduces the voltage and time dependent

properties of Purkinje neuron Nav currents (Ransdell et al., 2022), to test directly how I_{NaR}

properties affect Purkinje neuron repetitive firing. Development of the Nav conductance model is

published in Ransdell et al. (2022) and is based on Nav current properties recorded from C57bl/6

mouse Purkinje neurons. The kinetic state Markov model has a single open/conducting state

along with two parallel inactivation pathways which are separated as fast (IF1+IF2) and slow (IS)

inactivation kinetic states.

To test and verify the dynamic clamp configuration (depicted in Figure 1C), we used SutterPatch

dynamic clamp software to apply the model Nav conductance to a passive model cell (Sutter

Instruments). Voltage-clamp commands were applied to the model cell with the Markov model

conductance added using dynamic clamp software, and the model-generated currents were

recorded. Nav currents generated by this conductance model have similar voltage-dependent and

kinetic properties as those measured in mouse Purkinje neurons (Ransdell et al., 2022). Figure

1A (bottom) current traces show the performance of the Nav conductance model under an

identical voltage command. The nominal conductance model produces I_{NaT}, I_{NaR}, and I_{NaP} current

components. We utilized this model throughout our dynamic clamp experiments as a baseline

addition of Nav conductance and refer to it as the 'nominal' model throughout the rest of this

manuscript. The nominal model conductance was applied to adult (4-7-week-old) Purkinje

neurons in acute sagittal brain slices at physiological temperatures. The nominal model applied

at 400 nS (see Methods) significantly (P < 0.0001, paired Student's t-test, n = 30) increased the

spontaneous firing frequency of Purkinje neurons, compared to spontaneous firing without

dynamic clamp (Figure 1E). Addition of the nominal model conductance also resulted in significant

(P < 0.001, RM one-way ANOVA, n = 24) increases in evoked firing frequency during depolarizing

current injections (see Supplementary Figure 1).

Adjusting peak I_{NaR} does not scale the repetitive firing frequency of Purkinje neurons

We developed and tested Nav conductance models that have varying levels of peak I_{NaR}. To alter

peak I_{NaR}, individual rate constants in the nominal Markov model (presented in Figure 1) were

adjusted. By increasing the rate constant b2, which is responsible for transiting channel

occupancy from the fast-inactivated state into the open state, peak I_{NaR} was increased to a peak

value of 1.38 nA (measured at -45 mV), compared to 680 pA in the nominal model (see Figure

2A, B, *green*). Alternatively, by increasing the b3s rate constant, which is responsible for transiting

channel occupancy from the closed state into the slow inactivated state, peak I_{NaR} was reduced

to 350 pA (measured at -45 mV) (Figure 2A, B, *red*). Voltage-clamp records of Nav currents

produced by these models, along with peak I_{NaR} I-V plots for these models are shown in Figures

2B, and C. To test how scaling I_{NaR}, using these new models, affects the repetitive firing properties

of Purkinje neurons, we added the model variants to Purkinje neurons using dynamic clamp. The

nominal model, along with the 'I_{NaR} decreased' (*red*) and 'I_{NaR} increased' (*green*) models were

applied to the same Purkinje neurons using 400 nS conductance for each model. Consistent with

the hypothesis that I_{NaR} supports high rates of repetitive firing, we found adding the 'I_{NaR}

decreased' model conductance resulted in significantly reduced repetitive firing frequencies

compared to addition of the nominal model conductance (Figure 2D, *left*). However, when the 'I_{NaR}

increased' conductance model was added, there was no significant change in the repetitive firing

rates compared to addition of the nominal model conductance (Figure 2D, *right*). The 'I_{NaR}

increased' conductance model has a peak I_{NaR} value that is 2.3 times larger than the nominal

model (see Figure 2C). We reasoned that additional Nav current properties, other than peak I_{NaR},

were likely altered in creating the 'I_{NaR} increased' and 'I_{NaR} decreased' models, which may have

influenced the effects of each model conductance on repetitive firing frequencies. Analysis of the

voltage-clamp simulation traces revealed the time constant of I_{NaR} decay was reduced in both the

'I_{NaR} decreased' and 'I_{NaR} increased' models, compared to the nominal model. Additionally, we

noted I_{NaP} was much smaller in the 'I_{NaR} reduced' model compared to the nominal model.

Measures of these Nav current components are presented under the I-V plot in Figure 2C.

Purkinje neuron repetitive firing rates are not affected by adjusting the time constant of

I_{NaR} decay

Because altering a single rate constant was found to alter multiple Nav conductance parameters,

notably the tau of I_{NaR} decay and the proportion of I_{NaP}, we utilized a numerical optimization

procedure (see schematic in Fig. 3A and description in Methods) to develop additional Nav

conductance Markov models that exhibit more targeted changes to the nominal model properties,

while working to conserve other Nav conductance parameters in the nominal model. Our first goal

was to test if the rate of I_{NaR} decay influences how a Nav conductance affects repetitive firing. To

develop models with either a longer or shorter I_{NaR} decay, compared to the nominal model, we

adjusted the constraint for the decay of I_{NaR} before applying the rate constant optimization

(amoeba method) procedure (Figure 3A). This strategy allowed us to develop two new

conductance models that, compared to the nominal model in Figure 3B-C, have a tau of I_{NaR} decay

that is either longer ('τdecay increased', *blue*) or shorter ('τdecay decreased', *green*). The time

constant of I_{NaR} decay is plotted against voltage for each model in Figure 3B. While these models

have different peak I_{NaT}, I_{NaP}, and I_{NaR} values (using a 400 nS conductance) than the nominal

model (see Figures 3C1-C3), the values of these current components were similar to one another.

This enabled us to test the effect of altered I_{NaR} decay rates directly using dynamic clamp

experiments.

When the two newly developed models, 'τdecay increased' and 'τdecay decreased' were added

at 400 nS to adult Purkinje neurons, both models were found to cause significant increases in

repetitive firing, compared to no model (see Supplemental Figure 2). However, the increased

repetitive firing rates (under both models) were not significantly different from one another (Figure

3D, E), indicating that while adding the modeled Nav conductance increases repetitive firing rates,

applying substantial changes to the tau of I_{NaR} decay does not affect these firing rates (see Figure

3D). Both of the new models resulted in significantly lower firing rates than the nominal model

(Figure 3E), however, we also noted that both the 'τdecay increased' and 'τdecay decreased'

models generate lower peak I_{NaP} values than the nominal model (Figure 3C2).

Peak I_{NaP}, not peak I_{NaR}, is critical in scaling Purkinje neuron repetitive firing rates.

We hypothesized that I_{NaP} may be the critical parameter for scaling Purkinje neuron repetitive

firing rates. To test this hypothesis, we again used the constrained numerical optimization method,

(presented in Figure 3A) to develop a Nav conductance with reduced I_{NaP} This model is referred

to as the 'I_{NaP} reduced' model. Even utilizing numerical optimization methods, it was difficult to

develop the 'I_{NaP} reduced' model without also significantly reducing peak I_{NaR} values (found in the

nominal model, see Figure 4A1, *red* and *black*). For this reason, we developed an additional Nav

model which has similar levels of I_{NaP} and I_{NaT} as the nominal model (Figure 4A2, 4A3, *orange*),

but that also has a reduced peak I_{NaR} compared to the nominal model (Figure 4A1, *orange*). We

refer to this model as 'I_{NaR} reduced'. Dynamic clamp-mediated addition of these models could

then be used to compare the effects of reduced I_{NaP} or reduced I_{NaR} on Purkinje neuron firing.

Results from these dynamic clamp studies, presented in Figure 4B, C, reveal addition of the 'I_{NaR}

reduced' model has no effect on repetitive firing rate compared to the nominal model (Figure 4C,

left) and addition of the 'I_{NaP} reduced' model results in significantly ($P < 0.0001$) lower repetitive

firing rates than addition of the nominal model (Figure 4C, *right*). Similar to previous experiments,

all three models caused a significant increase in repetitive firing frequency when compared to no

model/zero nS conductance (see Supplementary Figure 2).

Applying or subtracting I_{NaP} conductance alone scales repetitive firing

The nominal model, and subsequent iterations of the Nav conductance models developed in

Figures 3 and 4, have a persistent component that activates maximally at around -60 mV (see

Figures 3C2, 4A2). However, I_{NaP} measured in Purkinje neurons typically activates at more

depolarized voltages, with peak persistent currents evoked at around -40 mV (Ransdell et al.,

2017). To test how this shift in the voltage-dependence of I_{NaP} activation affects the contribution

of I_{NaP} to repetitive firing in Purkinje neurons, and to also examine how sole application of I_{NaP}

affects repetitive firing in Purkinje neurons, we utilized an I_{NaP} Hodgkin-Huxley (HH) model

(previously published in Solinas et al., 2007) for dynamic clamp experiments. We tested the I_{NaP}

HH model using voltage-clamp simulations and adjusted the conductance of the model to match

the peak I_{NaP} values measured from the nominal model (see Figure 5A, B). The HH I_{NaP}

conductance model applied in dynamic clamp using 3.4 nS conductance was found to be

sufficient to increase Purkinje neuron repetitive firing frequency (Figure 5C, *left*). Additionally, we

tested if subtracting this conductance via dynamic clamp reduces repetitive firing. We found that

dynamic clamp-mediated subtraction of I_{NaP} results in a significant reduction in repetitive firing

frequency (Figure 5C, *right*). As is evident in Figure 5C, the HH I_{NaP} conductance, applied across

cells at either 3.4 nS or -3.4 nS, has variable effects on firing frequency. In some cells, adding 3.4

nS I_{NaP} caused only slight increases in firing frequency, while in other cells, adding this

conductance drove a nearly 3-fold increase in firing frequency.

We investigated if passive membrane properties might influence the contribution of I_{NaP} to firing

frequency. No correlation was found between the input resistance of the cell or capacitance with

the increase or decrease in firing frequency after applying 3.4 nS or -3.4 nS, respectively, of the

I_{NaP} conductance model (data not shown). These passive membrane properties also did not

correlate with the change in cell firing after applying the nominal model conductance (see

Supplemental Figure 3). However, dynamic clamp current injection values during the interspike

interval, measured 3 ms prior to action potential threshold (see Methods), were found to correlate

with the percent increase or decrease in firing frequency after applying the 3.4 nS I_{NaP} model (r^2

= 0.68) or -3.4 nS I_{NaP} model (r^2 = 0.36), respectively. These data suggest the steady-state

depolarizing current provided by I_{NaP} during interspike intervals is a critical determinant in scaling

the repetitive firing rates of Purkinje neurons.

When applied using dynamic clamp, peak Nav current components (I_{NaT}, I_{NaR}, and I_{NaP}) do

not correlate with changes in the action potential waveform.

Application of the nominal model to spontaneously firing Purkinje neurons resulted in significant

alterations to the action potential duration (increased), peak (increased), amplitude (increased),

and threshold voltage (decreased) (see Table 2). The amplitude of the action potential after-

hyperpolarization and the maximum dV/dt, taken from a first derivative phase plot of the action

potential waveform, were not significantly affected by applying the nominal model conductance

(Table 2). Other Markov conductance models had varying effects on these action potential

properties. To test if the magnitude of the Nav current components (I_{NaT}, I_{NaR}, and I_{NaP}), measured

across models, scales or consistently alters a given action potential property, we plotted the

current value of I_{NaT}, I_{NaR}, and I_{NaP} from each Markov model against the mean change in the action

potential parameter after addition of the respective model. These plots (shown in Supplemental

Figure 4) indicate the size of these Nav current components across models do not consistently

affect or scale any of the measured parameters of the action potential waveform.

Across dynamic clamp Nav conductance models, I_{NaP} is positively correlated with changes

in repetitive firing frequency.

Using a similar approach as Figure 5D, we examined how dynamic clamp current injection values

relate to changes in repetitive firing frequency across Markov Nav conductance models. We

reasoned that during application of a model conductance, the contribution of I_{NaP} is most localized

during the interspike interval immediately preceding the subsequent action potential. For each of

the seven Markov conductance models, I_{DynC} was measured during the interspike interval (3 ms

prior to the threshold potential of the subsequent action potential), and the mean I_{DynC} was

calculated for each model. Across the seven Markov models, these mean I_{DynC} values were plotted

against the mean percent increase in repetitive firing (Figure 6B) for each model. This plot

revealed a strong positive correlation ($r^2 = 0.93$), suggesting the steady-state Nav current, which

occurs at subthreshold voltages, is a critical parameter for Nav conductance mediated increases

in repetitive firing frequency. In contrast, if the I_{DynC} value was measured at the peak of the action

potential waveform (see Figure 6A schematic), there was no correlation (r^2 = 0.004) with each

model's effect on repetitive firing frequency (Figure 6C).

To examine more directly how each current component affected firing, we also plotted the peak

I_{NaT}, I_{NaR}, and I_{NaP} values from each model (as it was measured at a single voltage-step), against

the mean increase in repetitive firing frequency after the model was applied in dynamic clamp.

Again, these plots reveal a strong correlation between I_{NaP} measured at -65 mV and the change

in repetitive firing rate (Figure 6D), while I_{NaR} (Figure 6E), and I_{NaT} (Figure 6F) values were not

significantly correlated to the mean increases in repetitive firing rate.

Markov model open state occupancy reveals steady-state I_{NaP} is the primary depolarizing

drive of subsequent action potentials

Across models, we hoped to determine why scaling peak I_{NaR} does not affect the repetitive firing

properties of Purkinje neurons. To analyze this more directly, we used voltage-clamp simulation

experiments in which the voltage command consisted of two consecutive action potential

waveforms previously recorded from a Purkinje neuron firing at ~60 Hz. Under this voltage-clamp

paradigm, we analyzed the performance of the nominal model in addition to the 'I_{NaR} reduced'

model (orange), and the 'I_{NaP} reduced model' (red) presented in Figure 4. In addition to examining

the current evoked by this voltage command for each model, we measured and plotted occupancy

in the open/conducting kinetic state. Time-locked traces from this examination are presented in

Figure 7. In comparing the 'I_{NaR} reduced' model (*orange*) traces with the nominal (*black*) traces,

there was only a slight difference in the open-state occupancy (middle traces) and evoked steady-

state I_{Na} traces during the interspike interval. This is consistent with the 'I_{NaR} reduced' model

having a slightly smaller I_{NaP} component than the nominal model (see Figure 4A2). In the 'I_{NaR}

reduced' model's current trace and open-state occupancy traces (*orange traces*), we found no

evidence of the reduced I_{NaR} component. This is likely because I_{NaR} is very small at subthreshold

voltages and because the decay of I_{NaR} is relatively rapid at hyperpolarized membrane potentials.

However, the 'I_{NaP} reduced' model (*red traces*) has significantly less open-state-occupancy

corresponding to less steady-state evoked I_{Na} during the interspike interval, revealing why the 'I_{NaP}

reduced' model, when applied using dynamic clamp, results in significantly lower firing

frequencies than the nominal model and the 'I_{NaR} reduced' model (see Figure 4C).

Peak I_{NaP} and I_{NaR} can be inversely adjusted by adjusting occupancy in the slow inactivated

kinetic state

In developing the various models with alterations in I_{NaP} and I_{NaR}, it became evident that increased

occupancy in the slow inactivated kinetic state, shown as IS in the Markov kinetic state models

(Figure 8A), was necessary to increase peak I_{NaR}. On reflection, this is unsurprising because the

decay of the I_{NaR} current is simulated in these Markov models by transiting open channels into the

IS state. Any simulated channels that fail to transit from the open state into IS, which is absorbing

(Ransdell et al., 2022), are available to contribute to I_{NaP}. According to these model properties,

we reasoned that we should be able to inversely scale peak I_{NaP} and I_{NaR} by adjusting the rate

constants that affect Open and IS state occupancy. These rate constants are labeled b2s and a2s

and are highlighted in panel 8A. We conducted voltage-clamp simulation studies to test this

hypothesis and indeed found that reducing the b2s rate constant results in an increase in the

proportion of I_{NaR} and reduces I_{NaP} (Figure 8B). Alternatively, if the a2s rate constant is reduced,

peak I_{NaP} is increased and I_{NaR} is diminished. As current traces in Figure 8B traces reveal,

inversely adjusting I_{NaR} and I_{NaP} in this way does not affect recovery of fast-inactivated channels

into the open state, i.e., channels are still able to recover from fast-inactivation into an

open/conducting state on repolarization of the membrane.

Previous dynamic clamp experiments suggest peak I_{NaR} has no significant impact on the firing

rate or action potential waveform properties of cerebellar Purkinje neurons. However, in adjusting

rate constants that affect occupancy in the slow inactivated state, we inversely scale I_{NaR} and I_{NaP}.

These data indicate the presence of I_{NaR} may reflect the degree of Nav channel slow inactivation,

which could regulate or fine-tune the population of Nav channels available to participate in I_{NaP}.

We hypothesized that adjusting slow inactivation occupancy, and increasing or decreasing I_{NaP},

as we have done if Figure 8B, will result in increased or decreased repetitive firing rates. To test

this hypothesis, we applied the models developed in Figure 8B in dynamic clamp experiments,

using application of the nominal model in dynamic clamp as a baseline. Indeed, as panels 8C and

8D show, adjusting the transiting of open channels into the IS state can be used to inversely scale

I_{NaR} and I_{NaP}, which results in significant (compared to the nominal model) increases or decreases

in both spontaneous (Fig. 8C) and evoked (Fig. 8D) repetitive firing frequencies. Plotting a2s (Fig.

8E) or b2s (Fig 8F) rate constants against the resulting Nav conductance models' peak I_{NaP} and

I_{NaR} further reveal this relationship; with rate constant changes which reduce occupancy in the

slow-inactivated kinetic state (IS) causing an increase in I_{NaP} and reduction in I_{NaR}.

Discussion

We used dynamic clamp to apply several Nav conductance models (10), which have varying

resurgent (I_{NaR}) and persistent (I_{NaP}) Nav current properties, to test how these current components

contribute to the firing of adult cerebellar Purkinje neurons. Results from these experiments

indicate peak I_{NaR} does not directly contribute to the repetitive firing rates of Purkinje neurons.

Instead, the amplitude of I_{NaP}, particularly at subthreshold voltages, reliably scales firing frequency

regardless of I_{NaR} amplitude. Modeling work revealed I_{NaR} and I_{NaP} can be inversely scaled by

adjusting occupancy in the slow inactivated kinetic state, whereby slow inactivation removes Nav

channels from contributing to I_{NaP}.

I_{NaR} and excitability

In our initial experiments, we constructed two Markov kinetic state models with decreased or

increased I_{NaR} by adjusting the properties of two pairs of state transition rate constants of a

previously developed Nav conductance model (Ransdell et al., 2022). These hand-tuned Nav

models were individually added to Purkinje neurons via dynamic clamp. Surprisingly, after adding

a model Nav conductance with an increased I_{NaR} component, there were no significant changes

in firing frequency compared to the base (nominal) model (Figure 2). Additional dynamic clamp

studies utilized Nav models which differ in the peak amplitude of I_{NaR} and I_{NaP}, or that vary in the

time constant of I_{NaR} decay. These studies revealed I_{NaP} is the critical Nav conductance parameter

for tuning Purkinje neuron repetitive firing frequency, while peak I_{NaR} has very little effect on firing

frequency or the action potential waveform. Of course, this provokes an important question- what

role does I_{NaR} serve in regulating Purkinje neuron function if it has no effect on action potential

firing? We propose I_{NaR}, or more specifically the decay of I_{NaR} (which is rapid at subthreshold

voltages), may reflect the removal of non-inactivating Nav channels into a slow inactivated kinetic

state, and thus, I_{NaR} reflects a mechanism by which Purkinje neuron I_{NaP} is regulated.

There is a long-standing hypothesis that I_{NaR} increases membrane excitability and scales neuronal

repetitive firing rates since its discovery by Raman and Bean in 1997. *Scn8a* mutant mice (*Scn8a*

null) have reduced I_{NaR} (by 90%) and reduced Purkinje neuron firing frequencies, which suggests

I_{NaR} contributes to repetitive action potential generation (Raman et al., 1997). However, it was also

noted that the targeted deletion of *Scn8a* results in reduced (by ~70%) I_{NaP}, which may have

caused the attenuated repetitive firing rates (see Figures 4, 8). *Scn8a* null mice were also found

to have hyperpolarized interspike voltages, which also might reflect attenuated subthreshold I_{NaP}

(Raman et al., 1997).

The decay of I_{NaR} at subthreshold voltages is rapid (~3 ms) (Raman and Bean, 1997), suggesting

the majority of depolarizing Nav current during interspike intervals, and the Nav current

immediately preceding action potentials during repetitive firing, is a result of subthreshold I_{NaP}.

Subthreshold Nav current has previously been shown to play a critical role in setting the rate of

interspike interval depolarization as well as the action potential threshold voltage in mouse

Purkinje neurons as well as CA1 hippocampal neurons (Carter et al., 2012). The Markov

conductance models developed here and dynamic clamp studies presented in Figure 3 reveal

that even if the tau of I_{NaR} decay is increased (nearly doubled at -65 mV) or decreased, compared

to the nominal model (Figure 3B), the effects of I_{NaR} on repetitive firing are similar (Figure 3E),

revealing I_{NaR} does not directly influence repetitive firing frequency even with much slower decay

kinetics.

Targeted deletion of *Scn4b*, which encodes the Nav channel β4 (Navβ4) subunit, results in

reduced firing rates, as well as reduced (by ~50%) peak I_{NaR} in Purkinje neurons (Ransdell et al.,

2017). Importantly, this result was consistent with an earlier study which found the deletion of

Scn4b causes reduced I_{NaR} and evoked firing rates in rat striatal medium spiny neurons (Miyazaki

et al., 2014). To test if the deficits in *Scn4b*[-/-] Purkinje neuron firing could be rescued, Ransdell et

al. (2017) applied a modeled I_{NaR} conductance via dynamic clamp to these cells and found that

firing rates of *Scn4b*[-/-] Purkinje neurons could be rescued to wild type levels (Ransdell et al., 2017).

Importantly, however, the model Nav conductance utilized in Ransdell et al., (2017) included a

persistent Nav component and the effect of targeted *Scn4b* deletion on I_{NaP} was not investigated.

In RA projection neurons (RAPNs) of the adult zebra finch, Zemel and colleagues present a

collection of data showing that RAPN maturation includes paired increases in firing frequency and

peak I_{NaR}, which corresponded with increased levels of Navβ4 expression (Zemel et al., 2021). In

these studies, dynamic clamp was again used to test the role of I_{NaR} on RAPN firing frequencies,

however, similar to Ransdell et al. (2017), the I_{NaR} model used included a persistent Nav current

component.

Scaling I_{NaR} across conductance models did not drive consistent changes in action potential

duration (Supplemental Figure 4). Previous work has shown Purkinje neurons express large and

fast-activating voltage-gated potassium currents which quickly repolarize the membrane during

the action potential downstroke (Raman & Bean, 1999). These large hyperpolarizing

conductances likely shunt I_{NaR} depolarization and prevent any consistent effect of I_{NaR} on action

potential duration. This point underscores the reasonable assumption that I_{NaR} contribution to

membrane excitability will vary across neuronal cell types with unique firing and ion channel

expression properties (Lewis & Raman, 2014).

As with any dynamic clamp investigation which focuses on intact neurons, complex architecture

and distinct neuronal compartments result in testing the effects of a conductance on a non-

isopotential membrane. This may cause erroneous voltage measures, and as a result, erroneous

dynamic clamp current injections. In our studies, we patched the broad end of the somatic Purkinje

neuron membrane toward the axonal projection. Previously, it was shown through focal TTX

application to either the peri-somatic region of the cell or at the first node of Ranvier along the

axon, that action potential initiation and spontaneous firing in Purkinje neurons relies on somatic

and peri-somatic Nav channels (Khaliq & Raman, 2006), suggesting the voltage measurements

and corresponding dynamic-clamp current injections in our studies were accurate affecting the

action potential initiation zone of Purkinje neurons. Across cells included in the study, passive

membrane properties did not correlate with the effect of the nominal Nav conductance on

repetitive firing (Supplementary Figure 3), suggesting firing was not impacted by variations in

Purkinje neuron morphology.

Persistent current and its relationship with resurgent current

Since the discovery of the persistent sodium current in Purkinje neurons (Llinás & Sugimori,

1980), I_{NaP} has been implicated in promoting repetitive and complex firing in several neuronal cell

types (Carter et al., 2012; Han et al., 2015; Lee & Heckman, 2001; Taddese & Bean, 2002;

Yamanishi et al., 2018). Experiments here demonstrate subthreshold I_{NaP} is critical in scaling

Purkinje neuron firing rate (Figure 6D), but also that I_{NaR} may reflect an important mechanism by

which Nav channels are removed from contributing to I_{NaP}. Increased levels of I_{NaP} and I_{NaR} have

often been identified as a result of gain-of-function mutations in Nav channel α subunits which are

associated with human diseases such as paroxysmal extreme pain disorder (PEPD), long QT

syndrome, and epilepsies (Eberhardt et al., 2014; Hargus et al., 2013; Jarecki et al., 2010; Pan &

Cummins, 2020). In addition to the increased levels of I_{NaR} and I_{NaP}, several of the cells in these

studies were found to have increased action potential firing (Estacion et al., 2008; Hargus et al.,

2013; Pan & Cummins, 2020). Mutations in Nav channel pore-forming α subunits, which drive

singular or dual changes in I_{NaR} and/or I_{NaP}, provide important clues regarding how changes in

Nav channel activation, fast-inactivation, and slow-inactivation processes affect these current

components, and additionally, how alterations in I_{NaR} and/or I_{NaP} affect membrane excitability. For

instance, PEPD-associated mutations in Nav1.7 typically impact the voltage-dependence of fast-

inactivation resulting in enhanced I_{NaP} (Dib-Hajj et al., 2013; Eberhardt et al., 2014). It will be

interesting to examine how mutations in Nav channel pore-forming or accessory subunits affect

I_{NaR} and I_{NaP} along with Nav channel slow inactivation processes in native neuronal cell types.

Slow inactivation and adjusting the ratio of I_{NaR} and I_{NaP}

Nav channel slow inactivation is distinct from conventional fast-inactivation and occurs during

prolonged periods of Nav channel activation, generally tens to hundreds of milliseconds (Aman &

Raman, 2007; Silva, 2014). The Markov models (Ransdell et al., 2022) used here have parallel

inactivation pathways, a fast-inactivation pathway and slow inactivation pathway, used to

accurately reproduce the fast-inactivation kinetics of I_{NaT} and slower decay of I_{NaR} at steady-state

voltages (see Figures 1A, C). Adjusting rate constants that determine entry into and out of the

slow inactivated kinetic state readily tunes the portion of I_{NaR} and I_{NaP}, which determines the

model's effect on Purkinje neuron firing (Figure 8). From these results, it seems Nav channels

can be regulated in at least two domains in order to regulate repetitive firing frequency: 1. The

proportion of channels that are activated and fail to undergo fast-inactivation (I_{NaR} + I_{NaP}) and 2.

The proportion of these non-inactivating channels which undergo slow inactivation (I_{NaR}). These

domains of Nav channel regulation would regulate the level of subthreshold I_{NaP} during interspike

intervals, which, as our dynamic clamp studies show, scales Purkinje neuron firing rates. Slow

inactivation of Nav currents has previously been shown to be essential in the regulation of

pacemaking activity in STN neurons (Do & Bean, 2003) and interspike voltage changes (Carter

et al., 2012). Our results further suggest the magnitude of I_{NaP}, may be regulated by Nav channel

slow inactivation in Purkinje neurons. Enhancing cAMP-dependent protein kinase and protein

kinase C activity has been shown to rapidly enhance Nav channel slow inactivation in

heterologously expressed Nav1.2 channels (Chen et al., 2006), suggesting post-translational

regulation of Nav channels can quickly modify this inactivation pathway. Reduced extracellular

Na^+ concentrations have also been found to increase entry into the slow inactivated kinetic state

(Afshari et al., 2004; Aman & Raman, 2007; Townsend & Horn, 1997). Moving forward, it will be

interesting to test how the regulation of slow inactivation processes in native neuronal cell types

affect Nav current components, particularly I_{NaR} and I_{NaP}, and to determine if post-translational

regulation of Nav channel slow inactivation is capable/necessary for the fine-tuning of I_{NaP} as well

as firing rates in spontaneously active cells.

Author Contributions

SPB and JLR conceived and planned the studies; SPJ and RJL performed dynamic clamp

experiments; SPJ and JDM developed Nav conductance models for dynamic clamp; SPJ and

JLR wrote the manuscript; SPJ, JLR and JDM revised the manuscript.

References

Afshari, F. S., Ptak, K., Khaliq, Z. M., Grieco, T. M., Slater, N. T., McCrimmon, D. R., & Raman, I. M. (2004). Resurgent Na Currents in Four Classes of Neurons of the Cerebellum. *Journal of Neurophysiology*, *92*(5), 2831–2843. https://doi.org/10.1152/jn.00261.2004

Aman, T. K., & Raman, I. M. (2007). Subunit dependence of Na channel slow inactivation and open channel block in cerebellar neurons. *Biophysical Journal*, *92*(6), 1938–1951. https://doi.org/10.1529/biophysj.106.093500

Carter, B. C., Giessel, A. J., Sabatini, B. L., & Bean, B. P. (2012). Transient sodium current at subthreshold voltages: Activation by EPSP waveforms. *Neuron*, *75*(6), 1081–1093. https://doi.org/10.1016/j.neuron.2012.08.033

Chen, Y., Yu, F. H., Surmeier, D. J., Scheuer, T., & Catterall, W. A. (2006). Neuromodulation of Na+ channel slow inactivation via cAMP-dependent protein kinase and protein kinase C. *Neuron*, *49*(3), 409–420. https://doi.org/10.1016/j.neuron.2006.01.009

Dib-Hajj, S. D., Yang, Y., Black, J. A., & Waxman, S. G. (2013). The Na(V)1.7 sodium channel: From molecule to man. *Nature Reviews. Neuroscience*, *14*(1), 49–62. https://doi.org/10.1038/nrn3404

Do, M. T. H., & Bean, B. P. (2003). Subthreshold sodium currents and pacemaking of subthalamic neurons: Modulation by slow inactivation. *Neuron*, *39*(1), 109–120. https://doi.org/10.1016/s0896-6273(03)00360-x

Eberhardt, M., Nakajima, J., Klinger, A. B., Neacsu, C., Hühne, K., O'Reilly, A. O., Kist, A. M., Lampe, A. K., Fischer, K., Gibson, J., Nau, C., Winterpacht, A., & Lampert, A. (2014). Inherited pain: Sodium channel Nav1.7 A1632T mutation causes erythromelalgia due to a shift of fast inactivation. *The Journal of Biological Chemistry*, *289*(4), 1971–1980. https://doi.org/10.1074/jbc.M113.502211

Estacion, M., Dib-Hajj, S. D., Benke, P. J., Te Morsche, R. H. M., Eastman, E. M., Macala, L. J., Drenth, J. P. H., & Waxman, S. G. (2008). NaV1.7 gain-of-function mutations as a

continuum: A1632E displays physiological changes associated with erythromelalgia and paroxysmal extreme pain disorder mutations and produces symptoms of both disorders. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *28*(43), 11079–11088. https://doi.org/10.1523/JNEUROSCI.3443-08.2008

Han, C., Estacion, M., Huang, J., Vasylyev, D., Zhao, P., Dib-Hajj, S. D., & Waxman, S. G. (2015). Human Nav1.8: Enhanced persistent and ramp currents contribute to distinct firing properties of human DRG neurons. *Journal of Neurophysiology*, *113*(9), 3172–3185. https://doi.org/10.1152/jn.00113.2015

Hargus, N. J., Nigam, A., Bertram, E. H., & Patel, M. K. (2013). Evidence for a role of Nav1.6 in facilitating increases in neuronal hyperexcitability during epileptogenesis. *Journal of Neurophysiology*, *110*(5), 1144–1157. https://doi.org/10.1152/jn.00383.2013

Häusser, M., & Clark, B. A. (1997). Tonic synaptic inhibition modulates neuronal output pattern and spatiotemporal synaptic integration. *Neuron*, *19*(3), 665–678. https://doi.org/10.1016/s0896-6273(00)80379-7

Ito, M. (1984). *The Cerebellum and Neural Control* (1st ed.). Raven Press.

Ito, M., Yoshida, M., & Obata, K. (1964). Monosynaptic inhibition of the intracerebellar nuclei induced from the cerebellar cortex. *Experientia*, *20*(10), 575–576. https://doi.org/10.1007/BF02150304

Jarecki, B. W., Piekarz, A. D., Jackson, J. O., & Cummins, T. R. (2010). Human voltage-gated sodium channel mutations that cause inherited neuronal and muscle channelopathies increase resurgent sodium currents. *The Journal of Clinical Investigation*, *120*(1), 369–378. https://doi.org/10.1172/JCI40801

Khaliq, Z. M., Gouwens, N. W., & Raman, I. M. (2003). The contribution of resurgent sodium current to high-frequency firing in Purkinje neurons: An experimental and modeling study. *The Journal of Neuroscience: The Official Journal of the Society for*

Neuroscience, *23*(12), 4899–4912. https://doi.org/10.1523/JNEUROSCI.23-12-04899.2003

Khaliq, Z. M., & Raman, I. M. (2006). Relative Contributions of Axonal and Somatic Na Channels to Action Potential Initiation in Cerebellar Purkinje Neurons. *The Journal of Neuroscience*, *26*(7), 1935–1944. https://doi.org/10.1523/JNEUROSCI.4664-05.2006

Lagarias, J. C., Reeds, J. A., Wright, M. H., & Wright, P. E. (1998). Convergence Properties of the Nelder—Mead Simplex Method in Low Dimensions. *SIAM Journal on Optimization*, *9*(1), 112–147. https://doi.org/10.1137/S1052623496303470

Lee, R. H., & Heckman, C. J. (2001). Essential role of a fast persistent inward current in action potential initiation and control of rhythmic firing. *Journal of Neurophysiology*, *85*(1), 472–475. https://doi.org/10.1152/jn.2001.85.1.472

Lewis, A. H., & Raman, I. M. (2014). Resurgent current of voltage-gated Na+ channels. *The Journal of Physiology*, *592*(Pt 22), 4825–4838. https://doi.org/10.1113/jphysiol.2014.277582

Llinás, R., & Sugimori, M. (1980). Electrophysiological properties of in vitro Purkinje cell somata in mammalian cerebellar slices. *The Journal of Physiology*, *305*, 171–195. https://doi.org/10.1113/jphysiol.1980.sp013357

Miyazaki, H., Oyama, F., Inoue, R., Aosaki, T., Abe, T., Kiyonari, H., Kino, Y., Kurosawa, M., Shimizu, J., Ogiwara, I., Yamakawa, K., Koshimizu, Y., Fujiyama, F., Kaneko, T., Shimizu, H., Nagatomo, K., Yamada, K., Shimogori, T., Hattori, N., … Nukina, N. (2014). Singular localization of sodium channel β4 subunit in unmyelinated fibres and its role in the striatum. *Nature Communications*, *5*, 5525. https://doi.org/10.1038/ncomms6525

Moreno, J. D., Lewis, T. J., & Clancy, C. E. (2016). Parameterization for In-Silico Modeling of Ion Channel Interactions with Drugs. *PLOS ONE*, *11*(3), e0150761. https://doi.org/10.1371/journal.pone.0150761

Nelder, J. A., & Mead, R. (1965). A Simplex Method for Function Minimization. *The Computer*

Journal, *7*(4), 308–313. https://doi.org/10.1093/comjnl/7.4.308

Obata, K., Ito, M., Ochi, R., & Sato, N. (1967). Pharmacological properties of the postsynaptic

inhibition by Purkinje cell axons and the action of γ-aminobutyric acid on Deiters

neurones. *Experimental Brain Research*, *4*(1), 43–57.

https://doi.org/10.1007/BF00235216

Obata, K., Takeda, K., & Shtnozaki, H. (1970). Further study on pharmacological properties of

the cerebellar-induced inhibition of Deiters neurones. *Experimental Brain Research*,

11(4), 327–342. https://doi.org/10.1007/BF00237907

Pan, Y., & Cummins, T. R. (2020). Distinct functional alterations in SCN8A epilepsy mutant

channels. *The Journal of Physiology*, *598*(2), 381–401. https://doi.org/10.1113/JP278952

Quattrocolo, G., Dunville, K., & Nigro, M. J. (2021). Resurgent Sodium Current in Neurons of the

Cerebral Cortex. *Frontiers in Cellular Neuroscience*, *15*, 760610.

https://doi.org/10.3389/fncel.2021.760610

Raman, I. M. (2023). The Hodgkin-Huxley-Katz Prize Lecture: A Markov model with permeation-

dependent gating that accounts for resurgent current of voltage-gated Na channels. *The*

Journal of Physiology. https://doi.org/10.1113/JP285166

Raman, I. M., & Bean, B. P. (1997). Resurgent sodium current and action potential formation in

dissociated cerebellar Purkinje neurons. *The Journal of Neuroscience: The Official*

Journal of the Society for Neuroscience, *17*(12), 4517–4526.

https://doi.org/10.1523/JNEUROSCI.17-12-04517.1997

Raman, I. M., & Bean, B. P. (1999). Ionic currents underlying spontaneous action potentials in

isolated cerebellar Purkinje neurons. *The Journal of Neuroscience: The Official Journal*

of the Society for Neuroscience, *19*(5), 1663–1674.

https://doi.org/10.1523/JNEUROSCI.19-05-01663.1999

Raman, I. M., Sprunger, L. K., Meisler, M. H., & Bean, B. P. (1997). Altered subthreshold

sodium currents and disrupted firing patterns in Purkinje neurons of Scn8a mutant mice.

Neuron, *19*(4), 881–891. https://doi.org/10.1016/s0896-6273(00)80969-1

Ransdell, J. L., Dranoff, E., Lau, B., Lo, W.-L., Donermeyer, D. L., Allen, P. M., & Nerbonne, J.

M. (2017). Loss of Navβ4-Mediated Regulation of Sodium Currents in Adult Purkinje

Neurons Disrupts Firing and Impairs Motor Coordination and Balance. *Cell Reports*,

19(3), 532–544. https://doi.org/10.1016/j.celrep.2017.03.068

Ransdell, J. L., Moreno, J. D., Bhagavan, D., Silva, J. R., & Nerbonne, J. M. (2022). Intrinsic

mechanisms in the gating of resurgent Na+ currents. *eLife*, *11*, e70173.

https://doi.org/10.7554/eLife.70173

Ransdell, J. L., & Nerbonne, J. M. (2018). Voltage-gated sodium currents in cerebellar Purkinje

neurons: Functional and molecular diversity. *Cellular and Molecular Life Sciences:*

CMLS, *75*(19), 3495–3505. https://doi.org/10.1007/s00018-018-2868-y

Silva, J. (2014). Slow inactivation of Na(+) channels. *Handbook of Experimental Pharmacology*,

221, 33–49. https://doi.org/10.1007/978-3-642-41588-3_3

Solinas, S., Forti, L., Cesana, E., Mapelli, J., De Schutter, E., & D'Angelo, E. (2007).

Computational reconstruction of pacemaking and intrinsic electroresponsiveness in

cerebellar Golgi cells. *Frontiers in Cellular Neuroscience*, *1*, 2.

https://doi.org/10.3389/neuro.03.002.2007

Taddese, A., & Bean, B. P. (2002). Subthreshold Sodium Current from Rapidly Inactivating

Sodium Channels Drives Spontaneous Firing of Tuberomammillary Neurons. *Neuron*,

33(4), 587–600. https://doi.org/10.1016/S0896-6273(02)00574-3

Thach, W. T. (1968). Discharge of Purkinje and cerebellar nuclear neurons during rapidly

alternating arm movements in the monkey. *Journal of Neurophysiology*, *31*(5), 785–797.

https://doi.org/10.1152/jn.1968.31.5.785

Townsend, C., & Horn, R. (1997). Effect of alkali metal cations on slow inactivation of cardiac

Na+ channels. *The Journal of General Physiology*, *110*(1), 23–33.

https://doi.org/10.1085/jgp.110.1.23

Yamanishi, T., Koizumi, H., Navarro, M. A., Milescu, L. S., & Smith, J. C. (2018). Kinetic

properties of persistent Na+ current orchestrate oscillatory bursting in respiratory

neurons. *The Journal of General Physiology*, *150*(11), 1523–1540.

https://doi.org/10.1085/jgp.201812100

Zemel, B. M., Nevue, A. A., Dagostin, A., Lovell, P. V., Mello, C. V., & von Gersdorff, H. (2021).

Resurgent Na+ currents promote ultrafast spiking in projection neurons that drive fine

motor control. *Nature Communications*, *12*(1), 6762. https://doi.org/10.1038/s41467-021-

26521-3

Figure Legends

Figure 1. Dynamic clamp addition of a modeled Nav conductance increases the spontaneous

firing frequency of adult cerebellar Purkinje neurons **A.** Voltage clamp step protocol (top) and

current traces of the evoked fast-transient (I_{NaT}) and resurgent (I_{NaR}) Nav currents in an acutely

dissociated Purkinje neuron (*middle*). The same voltage protocol was used to evoke and measure

Nav currents generated by a Markov Nav conductance model. **B.** This simulated Nav current was

generated by a nine-state gating Markov matrix that includes three closed states (C1, C2, C3),

two inactivated-closed states (IC1, IC2), two fast inactivated states (IF1, IF2), a slow inactivated

state (IS), and an open state (O). Transition rate constant values are denoted by variables

between states (see Methods). Panel **C.** shows a schematic of the dynamic clamp setup used to

interface Nav conductance models with a current injection electrode that is patched on an adult

Purkinje neuron in a sagittal brain section. A computer running SutterPatch acquisition and

dynamic clamp software is interfaced with a Sutter digidata/amplifier, which is connected to the

patch electrode. The computer running SutterPatch software uses voltage signals measured in

the Purkinje neuron to calculate, in real-time (via the Nav conductance model) current injection

values. Current injections representing the model Nav conductance are then applied to the

patched cell. **D.** Spontaneous action potentials recorded from an adult Purkinje neuron before

and after (**D1**) the dynamic clamp-mediated addition of the Nav conductance represented in panel

B. Dynamic clamp-mediated current injections are shown in red. **D2.** Shows an expanded time

trace of a single action potential (*black*) and its corresponding dynamic clamp current injection

(*red*). **E.** Firing frequency of adult Purkinje neurons (*open circles*) was significantly (P < 0.0001,

paired Student's t-test, n = 30) increased after dynamic clamp-mediated addition of the nominal

Nav conductance (*closed circles*).

Figure 2. Adjusting peak I_{NaR} does not scale the repetitive firing frequency of Purkinje neurons **A.**

To generate the 'I_{NaR} decreased' and 'I_{NaR} increased' models, transition rate constant values b2

(*green box*) and b3s (*red box*) were selectively adjusted. Values of the nominal and manipulated

state transition variables at -45 mV are listed below the matrix. **B.** A simulated voltage clamp

recording of evoked I_{NaR} (at -45 mV) is shown for the nominal model (black), the 'I_{NaR} increased'

model (*green*), and the 'I_{NaR} decreased' model (*red*). **C.** Peak I_{NaR} values were measured at various

repolarizing voltage steps after a 5 ms depolarization step to 0 mV. These peak current values

are plotted against voltage for the nominal model (black), 'I_{NaR} increased' model (*green*), and the

'I_{NaR} decreased' model (*red*). I_{NaP} is digitally subtracted prior to all peak I_{NaR} measurements. All

models assessed using a 400 nS conductance (see Methods). **D.** Spontaneous firing frequency

of adult Purkinje neurons after dynamic clamp-mediated addition of the nominal and 'I_{NaR}

decreased' model (*left, red squares*), and the nominal and 'I_{NaR} increased' model (*right, green*

triangles). Spontaneous firing frequency of the 'I_{NaR} decreased' model is significantly reduced

compared to spontaneous firing frequencies measured during application of the nominal model.

Applying the 'I_{NaR} increased' model resulted in no significant change in firing frequency (paired

Student's t-test, 'I_{NaR} decreased' vs nominal: P = 0.0042, n = 13; 'I_{NaR} increased' vs nominal: P =

0.21, n = 13).

Figure 3. Purkinje neuron repetitive firing rates are not affected by adjusting the time constant of

I_{NaR} decay. An automated optimization (script) procedure utilizing the Nelder-Mead "amoeba"

algorithm (Nelder and Mead, 1965; Lagarias et al., 1998) was used to apply targeted changes to

Nav conductance models while preserving other aspects of the nominal Nav conductance model.

A schematic of this process is shown in panel **A.** A MATLAB script was used to apply the amoeba

algorithm, checking the difference between the simulation prediction and the target experimental

values with each iteration, and 'crawling' through potential rate constant values in order to

minimize total error. **B.** This amoeba method was used to generate two new Markov conductance

models with targeted changes in the time constant of I_{NaR} decay. The time constant of I_{NaR} decay

(τdecay) is plotted against voltage for the original nominal model and the two newly generated

models, 'τdecay decreased' (*green squares*) and 'τdecay increased' (*blue triangles*). **C.** A

simulated voltage clamp recording of model generating I_{NaR} is shown for the nominal (*black*),

'τdecay decreased' (*green*), and 'τdecay increased' (*blue*) models. **C1.** For each model, peak I_{NaR}

values are normalized to the max I_{NaR} generated by the nominal model and are plotted against

voltage. Similar plots of normalized I_{NaP} and I_{NaT} are presented in panels **C2.** and **C3.**, respectively.

D. Representative traces showing dynamic clamp-mediated addition of the 'τdecay reduced'

(*green*) and 'τdecay increased' (*blue*) conductances in the same cell. Voltage records are shown

in black with the corresponding dynamic clamp current injections shown below. **E.** Spontaneous

firing frequencies measured in Purkinje neurons after the addition of the nominal (*black circles*),

'τdecay decreased' (*green squares*), and 'τdecay increased' (*blue triangles*) model conductances

(400 nS) are plotted with lines connecting measurements taken from the same cell. Adding the

'τdecay decreased' and the 'τdecay increased' model conductances resulted in spontaneous firing

frequencies that were significantly reduced compared to firing frequencies measured after adding

the nominal model conductance, however, the addition of 'τdecay decreased' and the 'τdecay

increased' model conductances resulted in similar repetitive firing frequencies (paired Student's

t-test- τdecay decreased vs nominal, $P < 0.0001$, n = 29; τdecay increased vs nominal, $P < 0.0001$,

n = 29; τdecay increased vs τdecay decreased, $P = 0.94$, n = 29).

Figure 4. Reducing the amount of I_{NaP} modeled conductance significantly decreases spontaneous

firing frequency compared to the nominal model, while reducing I_{NaR} does not. **A.** Voltage clamp

simulation traces showing Nav currents generated by the nominal (*black*), 'I_{NaP} reduced' (*red*), and

'I_{NaR} reduced' (*orange*) models are shown with the common voltage command shown above. **A1-**

A3. Peak I_{NaR} (A1), I_{NaP} (A2), and I_{NaT} (A3) were normalized to the peak Nav currents generated

by the nominal model and are plotted against voltage. **B.** Representative action potential firing

during dynamic clamp-mediated addition of the 'I_{NaR} reduced' (*upper*) or 'I_{NaP} reduced' (*lower*)

conductance models is shown from the same Purkinje neuron. Dynamic clamp current injections

are presented as orange or red traces. **C.** The spontaneous firing frequency of Purkinje neurons

with addition of the nominal (*black circles*) conductance model was not significantly changed after

addition of the 'I_{NaR} reduced' conductance model (*left, orange triangles*). However, this firing

frequency was significantly decreased after the addition of the 'I_{NaP} reduced' conductance model

(*right, red diamonds*). (Paired Student's t-test. nominal vs 'I_{NaR} reduced', P = 0.21, n = 29; nominal

vs 'I_{NaP} reduced', P < 0.0001, n = 29).

Figure 5. Applying or subtracting an isolated I_{NaP} conductance scales repetitive firing in Purkinje

neurons **A.** A Hodgkin-Huxley (HH) modeled conductance (using 3.4 nS conductance) was used

to simulate isolated I_{NaP}. The current traces presented in panel A. show simulated voltage clamp

of this modeled conductance. **B.** Simulated peak I_{NaP} values versus voltage steps generated by

the nominal conductance model (*black*, 400 nS) are compared directly to the HH model (*blue*, 3.4

nS). The HH model's conductance values were adjusted so that peak I_{NaP} measurements would

match the peak I_{NaP} generated by 400 nS of the nominal model. An example of applying the HH

model in dynamic clamp experiments is shown to the right; the dynamic clamp current injection is

presented in blue. **C.** Spontaneous firing frequency of Purkinje neurons with no dynamic clamp

(*open circles*) was significantly increased after dynamic clamp addition of 3.4 nS of the HH

conductance model (*left, blue squares*) and significantly decreased after addition of -3.4 nS of the

HH conductance model (*right, red triangles*) (Paired Student's t-test. no model vs 3.4 nS, P =

0.0009, n = 13; no model vs -3.4 nS, P = 0.047, n = 14). **D.** The percent increase in repetitive

firing frequency after addition (*left*, 3.4 nS) or subtraction (*right*, -3.4 nS) of the HH conductance

model is plotted against the amplitude of dynamic clamp current injected during the interspike

interval (see Methods) (3.4 nS, r^2 = 0.68, n = 16; -3.4 nS, r^2 = 0.36, n = 15).

Figure 6. Across models used for dynamic clamp, I_{NaP} is positively correlated with the changes in

repetitive firing frequency. The action potential waveform shown in panel **A.** depicts how dynamic

clamp current measurements were obtained. For interspike values, dynamic clamp current was

recorded 3 ms before action potential threshold was reached. Note that the arrow that points to

steady-state voltage is not to scale, and the measurement was not taken at this point. For peak

values, dynamic clamp current was recorded during the peak (max voltage) of the action potential.

B. A strong correlation exists between the percent increase in spontaneous firing frequency,

driven by the addition of each Markov conductance model, and the dynamic clamp current

injection during the interspike interval of an action potential (r^2 = 0.93). If the dynamic clamp

current injection is measured during the peak of the action potential, the correlation coefficient (r^2)

falls to 0.0035 (**C.**). **D.** The correlation between the percent increase in spontaneous firing

frequency after addition of each Markov conductance model and the peak I_{NaP} (measured using

voltage clamp at a -65 mV voltage step, is also strong (r^2 = .96; P = 0.0012,), while this positive

correlation is absent if measurements of I_{NaR} (at -45 mV) (**E.**) or I_{NaT} (at 0 mV) (**F.**) are used (r^2 =

0.41, P = 0.12; r^2 = 0.38, P = 0.77, respectively). Pearson's correlation test was used to determine

significance.

Figure 7. Action potential clamp experiments were conducted using a voltage command (dashed

trace, *upper*) that includes two action potentials that were originally recorded from a Purkinje

neuron firing at ~60 Hz. The nominal (*shown in black*), 'I_{NaR} reduced' (*shown in orange*), and 'I_{NaP}

reduced' (*shown in red*) Markov conductance models (see Figure 4) were applied to a Sutter

Model cell during these simulated action potential clamp experiments. Kinetic state occupancy

plots, showing occupancy in the open state, are shown below (*middle traces*) and are time-locked

with the voltage command. The bottom traces show the time-locked evoked currents produced

by the voltage command for each of these models.

Figure 8. Peak I_{NaP} and I_{NaR} can be inversely adjusted by changing rate constants responsible for

slow inactivation occupancy **A.** To develop models with inversely scaled I_{NaP} and I_{NaR}, the state

transition rate constants between the open (O) and slow inactivated (IS) states were individually

altered. Decreasing the b2s (*blue box*) transition rate constant (in the nominal model) resulted in

a conductance model with increased I_{NaR} and decreased I_{NaP}. Decreasing the a2s (*red box*)

transition rate constant resulted in a conductance model with decreased I_{NaR} and increased I_{NaP}.

These changes resulted in the simulated voltage clamp traces that elicit the b2s reduced (*blue*)

and a2s reduced (red) models, which are compared to the nominal model (*black*) in panel **B. C.**

When compared to the nominal conductance model (black circles), spontaneous firing frequency

of adult Purkinje neurons was significantly *increased* after addition of the a2s reduced model (*red*

squares, left), and significantly *decreased* after addition of the b2s reduced model (*blue triangles,*

right) (Paired Student's t-test. 400 nS a2s reduced, P = 0.0004, n = 8; b2s reduced, P = 0.0029,

n = 9;). **D.** Representative recordings of action potentials during addition of the nominal model

(*black, top*), and the a2s decreased (*red, middle*), and b2s decreased (*blue, bottom*) models (*left*)

reveal scaling in spontaneous and evoked (see F-I plots, *right*) firing rates in a single Purkinje

neuron. **E. F.** 3D plots were created in which the a2s (E.) or b2s (F.) state transition rate constants

are decreased across 14 iterations and plotted as log10 against the resulting peak I_{NaP} and I_{NaR}

values (measured at -45 mV). By decreasing the a2s state transition rate, and thereby decreasing

accumulation into the slow inactivated state, peak I_{NaP} is increased while peak I_{NaR} is decreased.

Alternatively, if the b2s state transition rate is decreased, accumulation into the slow inactivated

state is increased, resulting in increased peak I_{NaR} and reduced I_{NaP}.

Figure 1. Brown et al.

A.

	Nominal	I_{NaR} decreased
b3s	0.948	3.884

	Nominal	I_{NaR} increased
b2	0.108	3.787

C.

τdecay of I_{NaR}	I_{NaT} at -45mV	I_{NaP} at -45mV
11.9ms	-3531.1pA	-119.7pA
4.3ms	-2496.6pA	-27.2pA
4.5ms	-3840.5pA	-127.8pA

Figure 2. Brown et al.

Figure 3. Brown et al.

Figure 4. Brown et al.

Figure 5. Brown et al.

Figure 6. Brown et al.

Figure 7. Brown et al.

Figure 8. Brown et al.

Table 1. Table of rate constants for Markov Nav conductance models

Variable	Nominal	τdecay Reduced	τdecay Increased	I_{NaR} Reduced	I_{NaP} Reduced
a11 var1	0.023989	0.02288	0.0234922	0.024273	0.017821
a11 var2	961.08	962.9323	955.288069	961.1444	955.74551
a12	856.13	861.1349	855.83844	855.6402	943.691619
a13	72.682	70.27949	75.02708	72.05751	45.1006
b11 var1	0.17233	0.13391	0.1957689	0.145184	0.162969
b11 var2	19.691	18.23159	18.08988	20.02451	17.5219955
b12	88.549	87.66816	90.404	88.7198	77.35325
b13	148.41	162.7141	230.9623	174.5183	160.84845
a3 var1	0.36734	0.514279	0.391971	0.308533	1.006506
a3 var2	980.34	964.714	979.85264	980.6689	985.713
b3 var1	53.241	53.5899	53.2644	53.28371	74.161099
b3 var2	14.204	13.08476	13.65378	14.39238	14.43037
a2 var1	87.852	84.88214	99.3117	87.57206	83.96448
a2 var2	999.72	998.5897	998.65724	999.7462	999.9939
a6 var1	609.21	589.9252	621.0904	610.1094	619.312417
a6 var2	86.49	84.00415	88.18935	101.533	96.412457
b6 var1	156.45	224.2441	150.19414	156.3997	147.127477
b6 var2	30.317	28.8311	39.98108	58.02996	56.26988
a2s var 1	15.817	37.25738	12.52364	15.71892	22.593552
a2s var 2	999.82	999.9025	999.95304	999.7914	987.9528
b2s var1	0.001001	0.001207	0.001026	0.001003	0.001002
b2s va2	11.963	10.81732	13.212289	11.96024	15.16399
a3s var1	0.00447733	0.001042	0.003335	0.005083	0.00100109
a3s var2	999.93	999.2872	999.97387	999.9429	999.97908

Transition rates are of the form: a11 = Tfactor*1/(Input(1)*exp(-V/Input(2)));
a12=Input(3)*a11; a13= Input(4)*a11; b11=Tfactor*1/(Input(5)*exp(V/Input(6)));
b12=Input(7)*b11; b13=Input(8)*b11; a3 = Tfactor*Input(9)*exp(-V/ Input(10));
b3=Tfactor*Input(11)*exp((V)/Input(12)); a2=Tfactor*(Input(13)*exp(V/Input(14)));
b2=((a13*a2*a3)/(b13*b3)); a6=Tfactor*(Input(15)*exp(V/ Input(16)));
b6=Tfactor*Input(17)*exp(-V/Input(18)); a2s = Tfactor*(Input(19)*exp(V/Input(20)));
b2s=Tfactor*(Input(21)*exp(-V/Input(22))); a3s=Tfactor*Input(23)*exp(-V/Input(24));
b3s = (a2s*a3s*a13)/ (b2s*b13);
Q10 = 3; T = 295;
Tfactor = 1.0/(Q10^((37.0-(T-273))/10.0))

Table 1. Brown et al.

Table 2. Action potential waveform properties after adding each Markov Nav conductance

Model	Mean AP duration (100%) (µs)	Mean AHP (mV)	Mean AP Peak (mV)	Mean AP Threshold (mV)	Mean AP Amplitude (mV)	Max dV/dt (V/µs)
No model (n = 29)	411.8 ± 11.95	-73.10 ± 0.561	1.024 ± 1.089	-59.11 ± 0.546	60.14 ± 1.242	587.7 ± 20.35
Nominal (n = 29)	475.3 ± 13.20 (p < 0.0001)	-72.70 ± 0.615 (p = 0.4499)	5.662 ± 1.044 (p < 0.0001)	-56.86 ± 0.578 (p = 0.0026)	62.52 ± 1.329 (p = 0.0011)	582.2 ± 25.62 (p = 0.7291)
I_{NaP} Reduced (n = 37)	488.6 ± 11.82 (p = 0.0015)	-73.41 ± 0.588 (p = 0.9242)	5.544 ± 1.342 (p < 0.0001)	-57.08 ± 0.628 (p < 0.0001)	62.62 ± 1.720 (p < 0.0001)	573.7 ± 26.18 (p = 0.0001)
I_{NaR} Reduced (n = 35)	523.9 ± 33.22 (p = 0.0153)	-72.93 ± 0.755 (p = 0.6177)	5.769 ± 1.213 (p = 0.8196)	-56.60 ± 0.807 (p = 0.0022)	62.37 ± 1.513 (p = 0.6351)	555.3 ± 24.62 (p = 0.0282)
τdecay decreased (n = 34) *Figure 3*	513.4 ± 17.43 (p = 0.0009)	-74.34 ± 0.501 (p < 0.0001)	10.93 ± 1.245 (p < 0.0001)	-58.84 ± 0.463 (p = 0.1665)	69.78 ± 1.410 (p = 0.0003)	670.6 ± 24.89 (p < 0.0001)
τdecay increased (n = 36) *Figure 3*	497.9 ± 13.32 (p < 0.0001)	-73.48 ± 0.700 (p = 0.0601)	7.036 ± 1.320 (p < 0.0001)	-56.98 ± 0.717 (p = 0.0318)	64.02 ± 1.484 (p < 0.0001)	589.5 ± 24.63 (p < 0.0001)
I_{NaR} decreased (n = 17) *Figure 2*	488.3 ± 28.81 (p = 0.0183)	-72.50 ± 1.321 (p = 0.6783)	1.364 ± 1.046 (p = 0.9883)	-56.29 ± 1.812 (p = 0.2675)	57.65 ± 2.402 (p = 0.5275)	502.7 ± 33.32 (p = 0.0968)
I_{NaR} increased (n = 15) *Figure 2*	508.9 ± 50.16 (p = 0.2171)	-72.14 ± 1.498 (p = 0.3844)	3.745 ± 1.112 (p = 0.0281)	-52.79 ± 1.872 (p = 0.0088)	56.53 ± 1.817 (p = 0.2323)	478.4 ± 20.75 (p = 0.0005)

Mean ± standard deviation is presented across models for each action potential (AP) waveform parameter. P values are the result of paired Student's t-tests which compare action potential data with no dynamic clamp and action potential data after adding the Markov conductance model (see Methods). AHP refers to action potential after hyperpolarization (see Methods).

Table 2. Brown et al.