Quantifying the performance benefits of middle distance running super

spikes

Montgomery Bertschy[1*], Victor Rodrigo-Carranza[1,2*], Ethan W.C. Wilkie[3*], Laura A.

Healey[4], Jeremy Noble[3], Wayne J. Albert[3], Wouter Hoogkamer[1]

[1] Department of Kinesiology, University of Massachusetts, Amherst, MA, USA

[2] Sports Performance Research Group (GIRD) University of Castilla-La Mancha, Toledo,

Spain

[3] University of New Brunswick, Fredericton, NB, Canada

[4] PUMA SE, Innovation, Somerville, MA, USA

* These authors contributed equally to this work

Corresponding Author:

Wouter Hoogkamer

Integrative Locomotion Laboratory

Department of Kinesiology

University of Massachusetts, Amherst

Amherst, MA, 01003-9258 USA

Email: whoogkamer@umass.edu

ORCID:

Wouter Hoogkamer 0000-0003-0524-0196

Victor Rodrigo-Carranza 0000-0003-1637-7550

Laura Healey 0000-0003-0613-1629

Abstract

Background Quantifying the potential benefits of advanced footwear technology (AFT)

track spikes in middle-distance events is challenging, because repeated maximal effort

trials (as in sprinting) or aerobic running economy trials (as in long-distance running) are

not feasible.

Objective We propose and evaluate a novel approach to assess the benefits of AFT spikes

that consists of a series of 200m intervals at self-perceived middle-distance race pace.

Methods We conducted four experiments to evaluate the validity, sensitivity,

reproducibility, and utility of this novel method.

Results (1) Participants ran 1.26% slower (p=0.0188) in spikes with 200g added mass

than in control spikes, exactly equal to the known effects of shoe mass on running

performance. (2) Participants ran significantly faster in AFT prototype spikes than in

traditional spikes (p=0.0296). (3) Participants performed our interval-based method to

compare two other AFT prototype spikes against traditional spikes, on three separate

days. Group-level results were consistent across days, but to evaluate individual

responses at least two separate sessions are needed. (4) Participants ran significantly

faster in two AFT spike models than in traditional spikes (2.22% and 1.74%). Speed was

similar between a third AFT spike model and the traditional spikes. These speed results

were mirrored by changes in step length, as participants took significantly longer steps in

the two faster AFT spike models (2.31% and 1.92%), while step length was similar

between the other spikes.

Conclusion Our novel interval-based approach is a valid and reliable method to quantify

differences between spikes at middle-distance running intensity.

Key points

● A novel, effort-based methodology allows for valid and reliable comparison of

spikes for middle-distance running events.

● AFT track spikes are found to be significantly faster than traditional track spikes,

primarily due to increases in step length.

● Group-level results were consistent across days, but at least two separate sessions

are needed to evaluate individual responses.

Declarations

Compliance with Ethical Standards

Ethical approval The study was performed in accordance with the ethical standards of

the Declaration of Helsinki. Ethics approval was obtained from the University of

Massachusetts Institutional Review Board (Protocol# 2647), from the University of New

Brunswick Research Ethics Board (REB #2021-079) and the University of Castilla-La

Mancha University Ethics Committee (Protocol CEIC924)

Informed consent Written informed consent was obtained from all individual

participants included in the study.

Funding

This research was partly supported by a research contract from PUMA SE with the

University of Massachusetts, Amherst. All spikes were provided by PUMA SE.

Conflicts of interest / Competing interests

Montgomery Bertschy, Victor Rodrigo-Carranza, Ethan W.C. Wilkie, Jeremy Noble and

Wayne J. Albert have no conflicts of interest relevant to the content of this article.

Wouter Hoogkamer is a paid consultant to PUMA and has received research grants from

PUMA and Saucony. Laura A. Healey is an employee of PUMA. No footwear company

had any influence on the conceptualization of this study or results presented in this

publication.

Data availability

All data discussed in this manuscript are provided in the tables.

Authors' contributions

Conceptualization: Wouter Hoogkamer; Experimental design: Montgomery Bertschy,

Laura A. Healey, Wouter Hoogkamer; Data collection: Montgomery Bertschy, Victor

Rodrigo-Carranza, Ethan W.C. Wilkie, Jeremy Noble; Data analysis: Montgomery

Bertschy, Victor Rodrigo-Carranza, Ethan W.C. Wilkie, Jeremy Noble, Wayne J. Albert;

Writing - original draft preparation: Montgomery Bertschy, Victor Rodrigo-Carranza,

Ethan W.C. Wilkie, Laura A. Healey, Wouter Hoogkamer; Writing - review and editing:

Montgomery Bertschy, Victor Rodrigo-Carranza, Ethan W.C. Wilkie, Laura A. Healey,

Jeremy Noble, Wayne J. Albert, Wouter Hoogkamer

1   Introduction

Once thought to be the last unbreakable barrier on the track [1], the magical sub-4-

minute mile mark stood strong for decades, until Sir Roger Bannister in 1954 completed

the unthinkable and ran the 1609 meters in 3:59.4 seconds. In February 2023, 52 male

professional, club, and collegiate level athletes broke the 4-minute mile barrier all within

the span of an hour at Boston University's indoor track facility. A similar phenomenon

was observed in the women's section, as three athletes broke the 4:30 mile barrier, and 17

ran sub-4:36, a mark more equivalent to the men's barrier.[2] Over the past decades, the

number of athletes breaking these barriers had been increasing gradually but there has

been a substantial increase since 2021.While the Boston University track has been known

to be fast, another important common feature among these performances was the use of

advanced footwear technology (AFT) track spikes, colloquially known as "super spikes".

Similar to the now common AFT marathon racing shoes (e.g., Nike Vaporfly), AFT track

spikes feature a lightweight, compliant, and resilient midsole foam, often combined with

a stiff midsole plate. While we have argued before that anecdotal performances should

not be used as evidence on the efficacy of specific shoes [3], with the widespread

improvement across middle distance running performances, this warrants a closer look.

The recent improvements on the track are similar to improvements in road

running performances since 2017 which have been attributed to advances in road running

shoes.[4–6] While the adoption of AFT road shoes happened in parallel with the publication

of a randomized crossover study showing 4% improvements in running economy,

scientific evidence on potential benefits of AFT track spikes is currently limited to race

results analyses (preprint: [7]) and preliminary laboratory data.[8] Willwacher et al., [7] found

in an observational study that AFT track spikes had a greater impact on performance in

middle-distance events (0.31 - 1.33%) than in sprint events (0.22 - 0.66%) for both men

and women.

Traditionally, the effects of footwear on middle distance running performance

have been studied in the lab by measuring running economy, or on the track with time

trials. Barnes and Kilding [9] observed that running economy in traditional lightweight

track spikes was ~3% worse than in AFT road shoes during treadmill running at speeds

between 3.9 and 5.0 m/s. Oehlert et al.,[8] showed that different AFT track spikes improved

running economy by ~2% compared to a traditional spike model during treadmill running

at 14 km/h in women and at 16 km/h in men. Lab-measured changes in running economy

have been shown to translate to changes in running performance [10], but for speeds faster

than 3 m/s, performance improvements can be expected to be smaller than observed

improvements in running economy.[11] Furthermore, running economy measures are only

valid when quantified during running at a sustainable, aerobic, steady-state intensity. For

very fit, elite participants, this could involve running as fast as 5.8 m/s.[12,13] However,

world record pace for middle distance events ranges from 5.9 m/s (women's 5000m) to

7.9 m/s (men's 800m). How footwear affects running economy in high-caliber runners at

5.0 m/s [14] might not be indicative of how this footwear affects running performance at a

speed of 7.9 m/s, almost 60% faster. At these high middle-distance running speeds, at

intensities above aerobic steady-state capacity, quantifying physiological responses to

different spikes is not feasible or valid (see [3]).

An alternative approach to quantify the benefits of spikes could be to have

athletes perform a series of time trials in different spikes. Indeed, we showed earlier that

3,000m time trial performances are slower in shoes with added mass.[10] Participants ran

three time trials on an indoor track, each at the same time of day at the same day of the

week, a week apart. Performing race-distance time trials at maximal effort across

multiple days can seriously interfere with training programming, while doing multiple

high effort time trials on a single day, on the other hand, will affect the validity of the

results through fatigue building up.

In this paper we propose, evaluate, and use a novel approach to quantify potential

benefits of AFT middle-distance spikes. Rather than comparing time trial performance

using race distance time trials at maximal race effort over multiple days, we asked

athletes to run a series of 200m intervals at self-perceived race pace (800m or 1500m,

based on their specialty), within a single session. Series of 200m intervals at race pace are

a common component of middle-distance training, [15,16] and all participants were familiar

with this type of workout. This is important as the premise of the proposed protocol is to

measure differences in performance, while controlling effort. In a series of experiments,

we compared up to four different spike models, while participants ran at least two

intervals in each model. First, we evaluated the validity of our novel approach by

comparing traditional spike models that were nearly identical and only differed in mass

(control vs. control + 200g) by verifying the interval time differences against the

theoretically expected differences. Second, we assessed the sensitivity of this novel

protocol in comparing new prototype AFT spikes against traditional spikes. Third, we

evaluated the reliability between three different days for two AFT spikes against

traditional spikes, and finally we used this protocol to compare several commercially

available AFT spikes against traditional spikes. We used an inertial measurement unit

(IMU) placed at the pelvis to assess if any differences in running speed were

predominantly driven by changes in step frequency or by changes in step length.

2. Materials and Methods

2.1 A Series of Experiments

This study involved four separate experiments. The inclusion criteria were

generally similar for all four experiments: fitting US women's size 7, 8.5 or 9, or US

men's size 9 or 11 track spikes (Experiment 1) or fitting US men's size 9 track spikes

(Experiments 2, 3 and 4); being 16 years of age or older (Experiment 1 and 2) or 19 years

of age or older (Experiment 3 and 4); currently participating in track and field training;

having at minimum two years of experience training in track spikes; being free of any

orthopedic, cardiovascular, or neuromuscular conditions; and having not sustained injury

nor undergone any surgery within the 3 months prior to participation. Recruitment of

participants was done through means of personal contacts, local track clubs, and local

university track teams. Participants were Tier 3: highly trained, national level, but not

elite,[17] with average personal bests of 2:02min in 800 to 8:55min in 3000m across the

different experiments (average IAAF scores for each experiment: 815, 762, 817, 761,

respectively).[18] Because of the different timelines of the experiments and the increasing

availability of AFT spikes with time, participants experience running with AFT varied

between experiments: all participant had experience running in spikes, for Experiment 2

only two participants had run in AFT road shoes, none in AFT spikes, for Experiments 3

and 4 all participants had experience running with AFT shoes and/or spikes (Experiment

1 only used traditional spikes). The characteristics of the participants in each experiment

are shown in Table 1. The experiments were performed in accordance with the ethical

standards of the Declaration of Helsinki. Ethics approval was obtained from the

University of Massachusetts, Amherst Institutional Review Board (Protocol # 2647), the

University of New Brunswick, Fredericton Research Ethics Board (REB # 2021-079),

and the University of Castilla-La Mancha University Ethics Committee (Protocol

CEIC924). Before taking part in the study, participants provided informed written

consent.

2.1.1 Spike Conditions

For all experiments we used the PUMA EvoSpeed Distance 9 (TRAD) as the

control condition; a typical traditional track spike, lightweight in construction, while

providing only minimal cushioning and no added bending stiffness elements. It has a

low-profile EVA based foam with a flat geometry and a stack height of 17mm (Fig 1). In

Experiment 1, we compared the standard version of this spike (TRAD, 137-169g) against

the same spikes with 200g of added mass (TRAD+, 331-368g) (Table 2).

In Experiment 2 we compared two track spike conditions: TRAD and a PUMA

AFT prototype middle-distance spike (AFTP1, 179g). The AFTP1 spike combines a

compliant, highly resilient midsole foam with an embedded carbon fiber plate. It features

PUMA's Nitro foam, a lightweight PEBA-based foam compound, forming a thicker,

rockered midsole with midsole thickness of 19mm (Fig 1; Table 2).

In Experiment 3 we compared three track spike conditions: TRAD, a PUMA AFT

prototype long-distance spike (AFTP2, 151g) and a PUMA AFT prototype middle-

distance spike (AFTP3, 149g, different from AFTP1). The AFTP2 spike features

PUMA's Nitro foam, a lightweight PEBA-based foam compound, forming a rockered

midsole with midsole thickness of 18mm, but no carbon fiber plate. The AFTP3 spike

combines a compliant, highly resilient midsole foam with an embedded carbon fiber

plate. It features PUMA's Nitro foam, a lightweight PEBA-based foam compound,

forming a thicker, rockered midsole with midsole thickness of 19mm (Fig 1; Table 2).

In Experiment 4 we compared four commercially available track spike conditions:

TRAD, a commercially available PUMA AFT spike (AFTC1) and two competitor brand

AFT spikes (AFTC2 and AFTC3). The AFTC1 was the PUMA evoSPEED Long

Distance Nitro Elite+ (154g). It features PUMA's Nitro foam, a lightweight PEBA-based

foam compound, forming a rockered midsole with midsole thickness of 18mm, but no

carbon fiber plate. The AFTC2 (140g) and AFTC3 (163g) were commercially available

competitor brand AFT spike models, featuring similar, but different, AFT properties such

as increased compliance, resilience, bending stiffness, stack height and rocker geometry

(Fig 1; Table 2).

2.1.3 Experiment 1 - Validation
To evaluate the external validity of our novel approach, we compared 200m

interval time in control spikes versus spikes that were nearly identical and only differed

in mass (control vs. control + 200g). Similar to Hoogkamer et al., [10], we used a mass

intervention to predictably induce a change in running energetics.[19,20] Frederick et al., [19]

showed that, across a range of speed, adding 100g mass to each shoe increases the energy

cost of running by 1%. By how much a 1% increase in energy cost will slow down a

runner depends on how fast they are running. Kipp et al., [11] provided a theoretical

framework supported by observations in Hoogkamer et al., [10] that predicts that when

running at 6.67m/s (200m in 30s), a 1% increase in energy cost will lead to a 0.6% slower

interval. Therefore, with 200g added to each spike, we hypothesized that participants

would run 1.2% slower.

2.1.4 Experiment 2 - Sensitivity

Next, we evaluated if this approach had the sensitivity to find differences in what

may be seen as "anecdotal" benefits of AFT spikes over traditional spikes. For this newly

developed prototype middle distance spike, we anticipated that the metabolic savings

from the modern midsole foam and the embedded plate would add up to about 2%, based

on preliminary running economy results from Oehlert et al., [8]. Therefore, we

hypothesized that participants would run 1.2% faster in the prototype spikes.

2.1.5 Experiment 3 – Reliability

We then evaluated the reliability of our novel approach to quantify the potential

benefits of different AFT spikes technologies. The objective was to assess the

consistency of the performance benefits from AFT spikes across different days. We

hypothesized that different AFT spikes would result in faster times than TRAD, similarly

and consistently across the 3 measurement days.

2.1.6 Experiment 4 - Commercially Available Spikes

Finally, we used our novel approach to quantify potential benefits of a set of

different commercially available AFT spikes over traditional spikes. The aim was to

evaluate if potential performance gains from AFT spikes differed between models. We

hypothesized that the different AFT spikes all would result in 1.2% faster times than

TRAD, with no detectable differences between AFT models. Further, we hypothesized

that step frequency would be similar between conditions, while step length would be

about 1% longer in the AFT spikes [14,21].

2.2 Generic Experimental Protocol

2.2.1 General protocol

A single-blinded crossover methodology was implemented as part of the protocol

for Experiments 1, 2 and 4. During Experiment 3 the participants were able to see the

spikes but were not aware of the spike properties. All experiments took place at an

outdoor 400-meter track facility on days with little to no wind. The middle-distance track

runners enrolled in the study performed a series of 200-meter intervals at self-perceived

mile or 800m race pace. For middle-distance running we could not use maximal effort

trials and the effectiveness of the protocol depended on the participants' ability to run

every trial at a similar sub-maximal effort. To minimize bias, it is important that the

participants are unaware of the spike intervention and that they are able to control their

effort at a consistent submaximal level. Aiming to achieve the latter, we opted for 200m

intervals at self-perceived race pace, since this is a common component of middle-

distance training,[15,16] and all participants, like most middle-distance runners, were

familiar with this type of workout. To minimize any confounding effects of participants

running the first (excitement) or last ("emptying the tank") intervals at a higher effort, the

number of intervals that the researchers told the participants to run was higher than the

number of intervals they would actually run. Their first interval was considered a

habituation trial and not included in the evaluation, and then after running all the needed

experimental intervals, the researchers informed the participants that the data collection

was complete. Experiment 1 consisted of 6 experimental trials (3 intervals in both

conditions), and we asked participants to run 8 intervals, but then informed them that the

experiment was complete after 7 intervals. Similarly, for Experiments 2 and 4 we asked

participants to run 10 intervals, but we only included 4 and 8 experimental trials,

respectively (two intervals in each spike condition). For Experiment 3 the participants

performed the same protocol on three separate visits, so they ran all 8 intervals, but we

only analyzed the 6 experimental intervals.

The order of spike conditions was randomized for each participant prior to each

session. Participants performed each 200m interval from a standing start (similar to a

middle-distance race start). Participants were unaware of the times they were running for

each interval as they were not using a watch and researchers did not disclose the times

they recorded until after the data collection was complete. Rest between intervals was ten

minutes, to allow for changing spikes, filling out questionnaires (Experiments 2 and 4),

and walking to the start of the next interval.

2.2.2 Experimental Setup

When the participant arrived at the track facility, we explained the study protocol

to them, and they signed all the necessary forms [i.e., the Get Active Questionnaire

(CSEP, 2017) or the Physical Activity Readiness Questionnaire (ePARmed-X+) and

informed consent document]. We asked participants to run a warmup similar to as they

would for a 200m interval session, followed by drills and stretches as they saw fit. For

Experiment 4, after their warmup, we placed an inertial measurement unit (IMU; Vicon

Blue Trident sensor, Vicon, Oxford, UK) at the participant's sacrum with a clip attached

to the back of the waistband on their shorts.[22] Next, the participant removed their shoes

and researchers put fabric spike blinders on the participant's legs (Experiments 1, 2 and

4). Then the researchers put the spikes on the participant's feet and tightened the laces.

Following this, spike blinders were pulled down over the spikes. Spike blinders fit around

the participants spikes and lower legs in a non-restrictive fashion, providing a negligible

addition to spike mass.

At the 200m line the researcher started the IMU recording (Experiment 4) and

then moved towards the finish line. For the timing of each interval, researchers started

their timers when the participant started running, indicated by them quickly lowering one

hand from a raised position, and stopped their stopwatches when the participant crossed

the finish line. Once a trial was complete, the 200m time was recorded and the IMU

capture was stopped (Experiment 4). This was repeated for all intervals.

2.3 Data Analyses and Statistics

We calculated running speed in m/s by dividing the 200m distance by the interval

time. To calculate step frequency and step length in Experiment 4, we exported the raw

IMU data from the Capture.U Desktop. Orientation of the IMU was verified from

acceleration signal by finding the gravitational component ($-9.81 m/s^2$) during quiet

standing before the running trial began. The vertical component was filtered using a 5Hz

cut-off frequency and multiplied by subject mass to create a vertical ground reaction

force estimation, as demonstrated by Day et al., [22]. Peaks in the vertical ground reaction

force estimate were used to calculate step time and its inverse, step frequency. Average

speed for each interval was calculated as interval distance (i.e., 200m) divided by interval

time. Average step length was calculated as the product of average speed and average

step time. Next, we averaged the values across the two intervals that were performed in

the same spike condition.

We used linear-mixed effect models for each experiment [23]. For Experiments 1

and 2, we created models with spike as a fixed effect, subject as a random intercept, trial

as a random effect, and speed as the output variable. For Experiment 3, the model had

spike and day as fixed effects, subject as a random intercept, trial as a random effect, and

speed as the output variable. Additionally, for each day of this experiment, we created a

model with spike as a fixed effect, subject as a random intercept, trial as a random effect,

and speed was the output variable to verify that the results did not change per day. For

Experiment 4, we created three models with spike as a fixed effect, subject as a random

intercept, and speed, step length, and step frequency as the output variables, respectively.

A traditional level of significance ($\alpha = 0.05$) was employed. Tukey HSD tests were

conducted for each pairwise comparison of spike conditions. Additionally, we calculated

effect size (ES) as Cohen's d and reported percent differences for each measure with the

intention of providing relevant context for our determined p values. ES with values of

0.2, 0.5, and above 0.8 were considered as small, medium, and large, respectively [24]. To

evaluate the reliability, in Experiment 3 we calculated the coefficient of variation (CV =

SD / mean $\times$ 100%) within each day and for each spike, across days. The ranges of values

considered for categorizing CV% were as follows: CV < 10 as very good, 10–20 as good,

20–30 as acceptable, and CV > 30 as unacceptable [24]. Further, for each participant we

calculated the average speeds in each spike across days 1 and 2, and across days 1, 2 and

3. Then, we calculated the speed difference between the TRAD and the AFTP3 spikes.

Next, we calculated Pearson's correlation coefficients, for these differences, between

each individual day and the 3-day average, and between the 2-day average and the 3-day

average. Correlation coefficients with values of 0.1, 0.3, and 0.5 were considered as

small, medium, and large, respectively, with 1.0 being a perfect correlation [24]. Statistical

analyses were performed in R using lme4 package[25] for the linear mixed effect models

and emmeans package[26] for post hoc Tukey HSD tests. Descriptive statistics are reported

as mean ± standard deviation (SD). For visual display in the figures, we averaged the

200m interval times that were performed in the same spike condition for each

experimental session (three trials for Experiment 1; two trials for Experiments 2, 3 and

4).

3 Results

In Experiment 1, participants ran significantly slower in the TRAD+ spikes (5.96

± 0.53 m/s) than in the TRAD spikes (6.04 ± 0.50 m/s) (1.26%, ES = 0.16, $p = 0.019$;

Fig. 2). In Experiment 2, participants ran significantly faster in the AFTP1 spikes (6.42 ±

0.41 m/s) than in the TRAD spikes (6.28 ± 0.49 m/s) (2.45%, ES = 0.31, $p = 0.030$; Fig.

2).

In Experiment 3 participants ran significantly faster in the AFTP2 spikes (6.52 ±

0.36 m/s) and the AFTP3 spikes (6.60 ± 0.36 m/s) than in the TRAD spikes (6.40 ± 0.34

m/s) (1.87%, ES=0.34, $p < 0.0001$ and 3.12%, ES= 0.57, $p < 0.0001$, respectively).

Running speeds in AFTP3 spikes were significantly faster than in AFTP2 spikes (1.22%,

ES=0.19, p = 0.0004). The between-spike results were consistent between days. The

CV% values were classified as very good for inter-session (5.53%, 5.44% and 5.69% for

day 1, 2 and 3 respectively) and intra-session (5.29%, 5.53% and 5.46% for TRAD,

AFTP2 and AFTP3 respectively). The linear mixed effects model analyses for each day

separately all showed similar differences between spikes, both in magnitude (% change

and effect size) and in significance (Table 3).

The speed difference between TRAD and AFTP3 for each individual participant

varied across days. Pearson's correlation coefficient between day 1, day 2 and day 3, and

the 3-day average was large: 0.85, 0.77, and 0.88, respectively. However, the Pearson's

correlation coefficient between the 2-day average and the 3-day average was approaching

a perfect correlation: 0.97.

In Experiment 4 participants ran significantly faster in the AFTC1 spikes (6.44 ±

0.46 m/s) and the AFTC2 spikes (6.41 ± 0.51 m/s) than in the TRAD spikes (6.30 ± 0.47

m/s) (2.22%, ES= 0.31, p = 0.002 and 1.74%, ES= 0.22, p = 0.035, respectively; Fig. 2).

Running speeds in the AFTC3 spikes (6.36 ± 0.48 m/s) were not significantly different

from the TRAD spikes (0.95%, ES=0.13, p = 0.493). Running speeds were not

significantly different between the AFTC spikes (AFTC1 vs. AFTC2 p = 0.800; AFTC1

vs. AFTC3 p = 0.116; AFTC2 vs. AFTC3 p = 0.534), but the numerical difference

between AFTC1 and AFTC3 was more than 1% (1.24%, ES = 0.17).

Step frequency was not different between spikes (all p > 0.52; Fig. 2), but

participants ran with significantly longer steps in AFTC1 spikes (1.93 ± 0.15 m) and

AFTC2 spikes (1.92 ± 0.16 m) than in the TRAD spikes (1.88 ± 0.15 m) (2.31%, ES =

0.33, p = 0.001; 1.92%, ES = 0.26, p = 0.008; Fig. 2). Step lengths in the AFTC3 spikes

(1.91 ± 0.15 m) were not significantly different from the TRAD spikes (1.50%, ES =

0.20, p = 0.0512). Step lengths were not significantly different between the AFTC spikes

(AFTC1 vs. AFTC2 p = 0.893; AFTC1 vs. AFTC3 p = 0.472; AFTC2 vs. AFTC3 p =

0.875).

395 **4 Discussion**

In this paper we proposed, evaluated and used a novel approach to quantify

potential benefits of middle-distance AFT spikes. The combined results indicate that our

novel interval-based approach has the validity, sensitivity and reliability to quantify the

benefits of middle-distance AFT spikes. The results of Experiment 1 provide insights into

the validity of our interval-based approach. On average, participants ran 1.26% slower in

the spikes with 200g added mass, similar to the 1.2% difference we hypothesized. The

results of Experiment 2 indicate that our approach has the sensitivity to confirm what so

far has been considered "anecdotal" benefits of AFT spikes over traditional spikes.

Participants ran significantly (2.45%) faster in AFT prototype spikes than in traditional

spikes. The results of Experiment 3 indicate that our approach is reliable. On three

separate days between spike differences in running speed were similar, both in magnitude

(% change and effect size) and in significance. However, at the level of individual

runners, the responses to AFT spikes varied between days. Our data suggests that two

separate sessions are needed to obtain reliable individual data. Finally, we used our

interval-based approach to compare several commercially available AFT spikes against

traditional spikes. In two AFT spike models participants ran significantly (2.22% and

1.74%) faster than in the traditional spikes. Speed was similar between a third AFT spike

model and the traditional spikes. These speed results were mirrored by changes in step

length, as participants took significantly (2.31% and 1.92%) longer steps in the two faster

AFT spike models, while step length was similar between the third AFT spike model and

the traditional spikes.

4.1 Construct Validity

419   Shoe mass is one of the most studied features of running shoes [10,19,20,27]. There is

currently a strong consensus that there is a linear relationship between the increase in

shoe mass and the increase in running energy cost, with an increment in the

oxygen/energy cost of 1% per added 100g of shoe mass. Kipp et al., [11] provided a

theoretical framework to translate changes in energy cost into changes in running

performance, based the known curvilinear relationship between running speed and energy

cost and the cost of overcoming air resistance, and supported by shoe-mass time trial

results from Hoogkamer et al., [10]. Using this framework, we anticipated that at 6.67m/s

participants would run 1.2% slower in the TRAD+ spikes with 200g added mass. Our

participants ran their intervals at 6.0m/s; at this speed the framework predicts a 1.26%

slower speed in the TRAD+ spikes. We found that on average our participants ran 1.26%

slower, confirming the construct validity of our interval-based approach.

4.2 Sensitivity

433   Next, we evaluated if our interval-based approach is sensitive enough to find

differences between AFT spikes and traditional spikes. We anticipated that speed

differences between AFT spikes and traditional spikes would be similar to the effect of

200g added mass and we hypothesized that participants would run 1.2% faster in the AFT

prototype spikes. Intriguingly, participants ran 2.45% faster in the AFT prototype spikes

than in the traditional spikes, which is equivalent to the effect of 400g added mass.

4.3 Reliability

We evaluated the reliability of this protocol by asking 12 participants to run in the

same 3 spike conditions with the same protocol on 3 separate days. We found that, at the

group level, results were similar between all three days (Table 2) and similar to the 3-day

average values (significant speed improvements of 1.87 and 3.12% of the AFTP2 and

AFTP3, compared TRAD, respectively). The Pearson correlation coefficient for Day 1, 2,

and 3 vs. the 3-day average was large: 0.85, 0.77, and 0.88, respectively. However, the

correlation coefficient for 2-day vs. 3-day average results was substantially higher (0.97),

suggesting that a minimum of 2 days is needed to quantify dependable individual

responses to AFT spikes. Quantifying individual responses to AFT shoes is common

when aiming to identify biomechanical mechanisms underlying improvements from AFT

[28], or characteristics of runners who experience more or less benefits[29–31]. However, as

discussed by Barrons et al., [32], such analyses critically depend on the accuracy of the

response measurement. Note that both for running economy measures [32] and for speed

outcomes from our interval-based approach, multiple sessions with repeated conditions

are recommended, when attempting to quantify individual responses to AFT.

Interestingly, even though the differences between spikes varied between days for some

participants, the participants who ran faster in the AFTP3 spikes (with a curved carbon

fiber plate) than the AFTP2 spikes (no plate) did consistently (on each day) run faster in

the AFTP3 spikes than in the TRAD spikes, while the participants who ran faster in the

AFTP2 spikes than the AFTP3 spikes did consistently (on each day) run faster in the

AFTP2 spikes than in the TRAD spikes (Fig 3). This suggests that when only interested

in determining which spike is best suited for an individual, a single session in one day

(with repeated trials) may be sufficient.

4.4 Quantifying the Benefits of Super Spikes

Finally, we used our novel approach to quantify potential benefits of a set of

different commercially available AFT spikes over traditional spikes. Participants ran

significantly faster in the AFTC1 (2.22%) and AFTC2 (1.74%) than in the TRAD spikes.

However, running speeds in the AFTC3 spikes were not significantly different from the

TRAD spikes (0.95%), nor between the AFTC spikes. The improvements from the

commercially available AFT spikes compared to the traditional spikes were similar, but

slightly less than those of the prototype AFT spikes evaluated in Experiments 2 and 3

(2.22 - 3.12%) of this study. Over 800 or 1500m at these speeds athletes would run 2.5 or

5 seconds faster, respectively, in these AFT spikes than in traditional spikes.

While the effects of spikes on sprinting performance have been studied for at least

50 years, with an inquiry into the potential benefits of the banned Puma brush spikes in

the early seventies,[33] here we specifically focused on middle-distance running spikes.

Given the novelty of AFT spikes, and the challenges of quantifying their potential

benefits on middle-distance running performance,[3] we are not able to directly compare

our findings to the literature.

Willwacher et al., [7] performed in a longitudinal, observational and retrospective

study design evaluating the 100 best performances per year for men and women in

outdoor track events from 2010 to 2022. They observed more pronounced improvements

for the 1500m and longer distances longer (0.66 - 1.84%) than shorter distances (100m to

800m: 0.31-0.63%) for women, and less of an effect of AFT spikes in men. When

comparing running economy results for AFT spikes at 4.4 and 5.0 m/s, Oehlert et al., [8]

found 1.8 and 2.1% improvements for two AFT spike models as compared to traditional

spikes. Such running economy improvements can be expected to improve middle-

distance running performance by ~1.2% [11], which is less than the interval speed

improvements we observed in the current study at 6.4 m/s. Although, not specific to AFT

spikes, some earlier studies have evaluated 3,000m time trial performance on the track in

AFT road shoes and reported improvements ranging from 1 to 5%, similar to our spike

results [30,34].

Across Experiment 2, 3 and 4, participants consistently ran faster in the AFT

spikes than in the traditional spikes (with the exception of AFTC3). The three

experiments each used slightly different models of AFT spikes and the group-level

benefits varied from 2.45% in Experiment 2, 1.87 and 3.12% in Experiment 3 to 2.22 and

1.74% in Experiment 4. The consistency of these findings, while these experiments were

performed by different research teams in three different countries (US, Canada and

Spain) is a testament of the robustness of our interval-based approach and further

supports the validity of the observed benefits of AFT spikes.

Intriguingly, the faster speeds in the AFTC1 (2.22%) and AFTC2 (1.74%) spikes

as compared to the TRAD spikes in Experiment 4, were accompanied by 2.31% (i.e.,

4.3cm) and 1.92% (i.e., 3.6cm) longer steps in the two faster AFTC1 and AFTC2 spikes,

respectively. Step lengths were similar between the other spike models. Furthermore, step

frequency was similar across all spikes (~3.35 Hz, or 201 steps per minute). Combined,

this suggests that the performance benefits from AFT spikes in middle-distance running

originate from the spikes allowing runners to take longer step lengths without requiring

any compensation in step frequency. With average step lengths of 1.93m and 1.92m in

the AFTC1 and AFTC2 spikes, the participants took only 103.6 steps and 104.1 steps

during their 200m, respectively, about 2.5 steps less than in the traditional spikes. Over a

1500m race this adds up to 21 or 17 less steps in the AFTC1 and AFTC2 spikes than in

traditional spikes, respectively.

514   These biomechanical changes are similar to those reported in treadmill studies on

AFT road shoes but different from preliminary findings for AFT spikes. Treadmill

studies on road shoes consistently report that runners take ~1% longer steps in AFT shoes

than in traditional shoes [9,14,21,35]. Since running speed is fixed between conditions in such

studies, these longer steps are accompanied by reductions in step frequency. In a case

study on overground, track running, Russo et al., [36] evaluated biomechanics during 80m

trials at 95% of maximal speed comparing AFT spikes with a carbon-fiber plate and a

modern midsole foam against traditional spikes. The athlete ran at the same speed and did

not substantially change step length or step frequency, even though contact time was

longer in the AFT spikes. The fixed speed and single subject nature of Russo et al., [36]

might explain the differences with our study.

4.5 Limitations and Future Directions

While our study shows that our novel interval-based approach to quantify

potential benefits of middle-distance AFT spikes is valid and reliable at the group level,

we would caution against using the results of a single session to label an individual as a

responder/non-responder to AFT spikes, or to compare differences in responses between

individuals. Specific to that purpose, we found that at least two days are necessary to be

able to evaluate individual responses to AFT spikes.

Additionally, there are limitations with the implementation of this protocol. In this

current study, interval times were recorded by hand, which is subject to human error.

Step lengths were calculated as the average of the 200m run, based on the average step

frequency and the interval time. Future studies could place IMUs at the feet, which

allows to accurately calculate stride length at a stride-by-stride level [37]. Evaluating

between-spike differences in stride length at a stride-by-stride level could provide

additional information about how benefits of AFT spikes potentially vary from the

acceleration phase to the top speed phase.

For researchers looking to implement this protocol in future studies, great care

must be taken to limit the bias of the researchers and participants. Participants should be

blinded to their shoe condition. During collections in this present study, participants were

typically unaware of how many different spike conditions they were running in, what

spikes they were currently wearing, or that they were wearing other spikes than they

thought. However, with AFT spikes becoming more common, especially when

comparing commercially available spikes, some participants might be able to feel /

recognize differences in midsole cushioning or longitudinal bending stiffness between

spikes, which could bias their performances in our effort-based protocol.

Moreover, caution should be taken when interpreting the results on the benefits of

AFT spikes in Experiment 4. On average the male participants in this experiment ran

their intervals at ~6.4m/s, which is equivalent to a 2:05min 800m, or 3:54min 1500m. To

evaluate the benefits at elite or even world record level, this experiment should be

repeated with elite runners. Still, our findings are more informative than running

economy measurements for running at 16km/h (4.44m/s; equivalent to a 3:00min 800m).

Elite and/or female runners might have different plantar flexor strength or body mass

than our male participants, factors are likely to affect responses to AFT [38,39]. At the start

of our study, we only had spikes available in the men's sample size US9 (except for

Experiment 1), but follow-up studies should include more female participants.

5 Conclusions

To overcome the challenges in quantifying the potential benefits of advanced

footwear technology spikes related to the physiology of middle-distance running

intensity, we introduced a novel interval-based approach. In a series of experiments, we

showed the validity and reliability of our method and then used it to compare

commercially available spikes. Our reliability experiment showed that group-level results

were consistent across days, but that at least two separate sessions are needed to evaluate

individual responses. Participants ran ~2% faster and with ~2% longer steps in two

commercially available AFT spike models than in traditional spikes, while speed and step

length was similar between a third AFT spike model and the traditional spikes.

Acknowledgements

Running Title: Quantifying the performance benefits of super spikes

We would like to thank the athletes of Athletics Canada and the UNB Reds Varsity Track

and Cross Country team, and of the running communities in Amherst, Boston, San

Antonio, Castilla La-Mancha and Madrid for their participation in our study. We would

like to thank Rodger Kram and Jose van der Veen for energetic(s) discussions to spark

the ideas presented in this paper. We would also like to thank Dan Feeney for guidance

regarding our statistical analysis. Lastly, we would like to thank our undergraduate

researchers, Elizabeth Hoenes, Caleb Zylka, Violeta Muñoz de la Cruz and Robert Carew

for assisting with data collections and recruitment. All spikes were provided by PUMA

SE.

References

1.   Taylor B. What breaking the 4-minute mile taught us about the limits of
conventional thinking. *Harv Bus Rev* 2018.
2.   Martin Fritz Huber. The four-minute mile is still worth celebrating. *Outside*
2023;
3.   Healey L, Bertschy M, Kipp S, Hoogkamer W. Can we quantify the benefits of
"super spikes" in track running? *Sports Med* 2022;**52**:1211–8.
https://doi.org/10.1007/s40279-022-01657-4.
4.   Senefeld JW, Haischer MH, Jones AM, Wiggins CC, Beilfuss R, Joyner MJ, et al.
Technological advances in elite marathon performance. *J Appl Physiol*
2021;**130**:2002–8. https://doi.org/10.1152/japplphysiol.00002.2021.
5.   Rodrigo-Carranza V, González-Mohíno F, Santos-Concejero J, González-Ravé
JM. Impact of advanced footwear technology on elite men's in the evolution of
road race performance. *J Sports Sci* 2023:1–8.
https://doi.org/10.1080/02640414.2023.2183103.
6.   Rodrigo-Carranza V, González-Mohíno F, Santos del Cerro J, Santos-Concejero
J, González-Ravé JM. Influence of advanced shoe technology on the top 100
annual performances in men's marathon from 2015 to 2019. *Sci Rep*
2021;**11**:1–9. https://doi.org/https://doi.org/10.1038/s41598-021-01807-0.
7.   Willwacher S, Mai P, Helwig J, Hipper M, Utku B, Robbin J. Does advanced
footwear technology improve track and road racing performance? An
explorative analysis based on the 100 best yearly performances in the world
between 2010 and 2022: The effects of super-footwear on world-class track
and road racing performances. *sportrxiv* 2024;**Preprint**.
https://doi.org/https://doi.org/10.51224/SRXIV.297.
8.   Oehlert GM, Jones EJ, Burns GT, Joubert DP. Comparative effects of advanced
footwear technology on running economy in track spikes and racing shoes:
161. *Med Sci Sports Exerc* 2023;**55**:41–2.
9.   Barnes KR, Kilding AE. A randomized crossover study investigating the
running economy of highly-trained male and female distance runners in
marathon racing shoes versus track spikes. *Sports Med* 2019;**49**:331–42.
https://doi.org/10.1007/s40279-018-1012-3.
10.   Hoogkamer W, Kipp S, Spiering BA, Kram R. Altered running economy directly
translates to altered distance-running performance. *Med Sci Sports Exerc*
2016;**48**:2175–80. https://doi.org/10.1249/MSS.0000000000001012.
11.   Kipp S, Kram R, Hoogkamer W. Extrapolating metabolic savings in running:
Implications for performance predictions. *Front Physiol* 2019;**10**:1–8.
https://doi.org/10.3389/fphys.2019.00079.
12.   Lucia A, Esteve-Lanao J, Olivan J, Gomez-Gallego F, San Juan AF, Santiago C, et
al. Physiological characteristics of the best Eritrean runners—exceptional
running economy. *Appl phy, nutri, and metab* 2006;**31**:530–40.
13.   Joyner, Hunter SK, Lucia A, Jones AM. Physiology and fast marathons. *J Appl*
Physiol 2020;**128**:1065–8. https://doi.org/10.1152/japplphysiol.00793.2019.

14. Hoogkamer W, Kipp S, Frank JH, Farina EM, Luo G, Kram R. A comparison of
the energetic cost of running in marathon racing shoes. *Sports Med*
2018;**48**:1009–19. https://doi.org/10.1007/s40279-017-0811-2.
15. Daniels J. Daniels' running formula. Human Kinetics, 2013.
16. Haugen T, Sandbakk Ø, Seiler S, Tønnessen E. The training characteristics of
world-class distance runners: an integration of scientific literature and
results-proven practice. *Sports Med-Open* 2022;**8**:1–18.
17. McKay AKA, Stellingwerff T, Smith ES, Martin DT, Mujika I, Goosey-Tolfrey VL,
et al. Defining training and performance caliber: a participant classification
framework. *Int J Sports Physiol Perform* 2022;**17**:317–31.
https://doi.org/10.1123/IJSPP.2021-0451.
18. Spiriev B, Spiriev A. IAAF scoring tables of athletics. International Association
of Athletics Federations Monaco, 2011.
19. Frederick EC. Physiological and ergonomics factors in running shoe design.
Appl Ergon 1984;**15**:281–7.
20. Franz JR, Wierzbinski CM, Kram R. Metabolic cost of running barefoot versus
shod: Is lighter better? *Med Sci Sports Exerc* 2012;**44**:1519–25.
https://doi.org/10.1249/MSS.0b013e3182514a88.
21. Hoogkamer W, Kipp S, Kram R. The biomechanics of competitive male
runners in three marathon racing shoes: a randomized crossover study.
Sports Med 2019;**49**:133–43. https://doi.org/10.1007/s40279-018-1024-z.
22. Day EM, Alcantara RS, McGeehan MA, Grabowski AM, Hahn ME. Low-pass
filter cutoff frequency affects sacral-mounted inertial measurement unit
estimations of peak vertical ground reaction force and contact time during
treadmill running. *J Biomech* 2021;**119**:110323.
23. Wilkinson RD, Mazzo MR, Feeney DF. Rethinking the statistical analysis of
neuromechanical data. *Exerc Sport Sci Rev* 2023;**51**:43–50.
24. Cohen J. Statistical Power Analysis. *Curr Dir Psychol Sci* 1992;**1**:98–101.
https://doi.org/10.1111/1467-8721.ep10768783.
25. Bates D, Mächler M, Bolker B, Walker S. Fitting linear mixed-effects models
using lme4. *arXiv preprint arXiv:14065823* 2014.
26. Lenth R, Singmann H, Love J, Buerkner P, Herve ME. Estimated marginal
means, aka least-squares means R package version 1.8. 4-1. 2019.
27. Rodrigo-Carranza V, González-Mohíno F, Santos-Concejero J, González-Ravé
JM. Influence of shoe mass on performance and running economy in trained
runners. *Front Physiol* 2020;**11**:1–8.
https://doi.org/10.3389/fphys.2020.573660.
28. Chollet M, Michelet S, Horvais N, Pavailler S, Giandolini M. Individual
physiological responses to changes in shoe bending stiffness : a cluster
analysis study on 96 runners. *Eur J Appl Physiol* 2022;**123**:189–177.
https://doi.org/10.1007/s00421-022-05060-9.
29. Knopp M, Muñiz-Pardos B, Wackerhage H, Schönfelder M, Guppy F, Pitsiladis
Y, et al. Variability in running economy of kenyan world-class and european
amateur male runners with advanced footwear running technology:

experimental and meta-analysis results. *Sports Med* 2023.
https://doi.org/10.1007/s40279-023-01816-1.

30. Rodrigo-Carranza V, Hoogkamer W, Salinero JJ, Rodríguez-Barbero S,
González-Ravé JM, González-Mohíno F. Influence of running shoe longitudinal
bending stiffness on running economy and performance in trained and
national level runners. *Med Sci Sports Exerc* 2023.

31. Rodrigo-Carranza V, Hoogkamer W, González-Ravé JM, González-Mohíno F.
Relationship between advanced footwear technology longitudinal bending
stiffness and energy cost of running . *Med Sci Sports Exerc* 2024;**In press**.

32. Barrons Z, Rodrigo-Carranza V, Bertschy M, Hoogkamer W. The Fallacy of
Single Trials: The Need for Multiple Trials in Assessing Running Economy
Responses in Advanced Footwear Technology. *Sport Med* 2024;**Under
review**.

33. Wood DW. An experimental test of the Puma model number 296 brush spike
shoe and the Mexico Puma model number 295 standard four spike shoe as to
their effect on sprint running. [Ohio]: Ohio University, 1972.

34. Hébert-Losier K, Finlayson SJ, Driller MW, Dubois B, Esculier JF, Beaven CM.
Metabolic and performance responses of male runners wearing 3 types of
footwear: Nike Vaporfly 4%, Saucony Endorphin racing flats, and their own
shoes. *J Sport Health Sci* 2021;**00**:1–10.
https://doi.org/10.1016/j.jshs.2020.11.012.

35. Hunter I, McLeod A, Valentine D, Low T, Ward J, Hager R. Running economy,
mechanics, and marathon racing shoes. *J Sports Sci* 2019;**37**:2367–73.
https://doi.org/10.1080/02640414.2019.1633837.

36. Russo L, Montagnani E, Buttari D, Ardigò LP, Melenco I, Larion A, et al. Track
running shoes: a case report of the transition from classical spikes to "super
spikes" in track running. *Appl Sci* 2022;**12**:10195.

37. Aristizabal-Pla G. Assessing the validity of the zero-velocity update method
for sprinting speeds. [Amherst]: University of Massachusetts Amherst, 2022.

38. Willwacher S, König M, Braunstein B, Goldmann JP, Brüggemann GP. The
gearing function of running shoe longitudinal bending stiffness. *Gait Post*
2014;**40**:386–90. https://doi.org/10.1016/j.gaitpost.2014.05.005.

39. Ortega JA, Healey LA, Swinnen W, Hoogkamer W. Energetics and
biomechanics of running footwear with increased longitudinal bending
stiffness: a narrative review. *Sports Med* 2021;**51**:873–94.
https://doi.org/10.1007/s40279-020-01406-5.

Figures

Fig. 1 Spike conditions for each experiment. From top to bottom, Experiment 1: TRAD

(PUMA EvoSpeed Distance 9), TRAD+ (PUMA EvoSpeed Distance 9 + 200g);

Experiment 2: TRAD, AFTP1 (PUMA AFT middle-distance prototype spike);

Experiment 3: TRAD, AFTP2 (PUMA AFT long distance prototype spike), AFTP3

(PUMA AFT middle distance prototype spike); Experiment 4: TRAD, AFTC1 (PUMA

evoSPEED Long Distance Nitro Elite+), AFTC2 (Competitor brand AFT spike), AFTC3

(Competitor brand AFT spike).

Fig. 2 Spike conditions significantly affect running speed and step length, but not step

frequency. Panel **A** shows average running speeds in different spike conditions, for

Experiment 1, 2, 3 and 4, from left to right. Panel **B** shows significantly longer step

lengths for AFTC1 and AFTC2 than TRAD. Panel **C** shows step frequency was similar

across spike conditions. Grey lines represent individual data, black lines represent group

mean values, error bars represent standard error of the mean. * indicates significant

difference between spike conditions.

Running Title: Quantifying the performance benefits of super spikes

729   **Fig. 3** Individual response of each participant in prototype AFT spikes vs. traditional

730   spikes (TRAD) in Experiment 3 for each of the three days and the 3-day average.

732 **Tables**

733   Table 1. Participant characteristics for each experiment (mean ± standard deviation).

Experiment	Participants (men, women)	Age (years)	Mass (kg)	Height (m)	Location
1	8, 4	20.3 ± 2.3	66.2 ± 10.3	1.76 ± 0.09	US
2	12, 0	21.3 ± 1.8	62.8 ± 2.8	1.75 ± 0.04	US, Canada
3	12, 0	27.0 ± 5.7	64.8 ± 3.3	1.76 ± 0.06	Spain
4	12, 0	22.0 ± 2.7	69.1 ± 8.3	1.81 ± 0.07	Canada

737   Table 2: Characteristics of experimental conditions used in each Experiment.

Spike	Sizes	Mass (g)	Midsole thickness (mm)	Experiments used
TRAD	5.5, 7, 9, 11	131, 149, 158, 170	17	1,2,3,4
TRAD+	5.5, 7, 9, 11	331, 341, 355, 369	17	1
AFTP1	9	179	19	2
AFTP2	9	151	18	3
AFTP3	9	149	19	3
AFTC1	9	154	18	4
AFTC2	9	140	12	4
AFTC3	9	163	17	4

741 Table 3: Magnitude and significance of between spike differences were consistent across
742 days.

Comparison		Day 1	Day 2	Day 3	3-Day average
TRAD vs. AFTP2	% change	2.02	1.72	1.74	1.87
	effect size	0.36	0.32	0.31	0.34
	p value	0.003	0.004	0.0003	<0.0001
TRAD vs. AFTP3	% change	3.11	3.13	2.93	3.12
	effect size	0.57	0.56	0.51	0.57
	p value	<0.0001	<0.0001	<0.0001	<0.0001
AFTP2 vs. AFTP3	% change	1.06	1.38	1.23	1.22
	effect size	0.19	0.25	0.21	0.19
	p value	0.124	0.015	0.035	0.0004

743