Self-perceived middle-distance race pace is faster

in Advanced Footwear Technology spikes

Montgomery Bertschy[1]*, Victor Rodrigo-Carranza[1,2]*, Ethan W.C. Wilkie[3]*, Laura A.

Healey[4], Jeremy Noble[3], Wayne J. Albert[3], Wouter Hoogkamer[1]

[1] Department of Kinesiology, University of Massachusetts, Amherst, MA, USA

[2] Sports Performance Research Group (GIRD) University of Castilla-La Mancha, Toledo,

Spain

[3] University of New Brunswick, Fredericton, NB, Canada

[4] PUMA SE, Innovation, Somerville, MA, USA

* These authors contributed equally to this work

<u>Corresponding Author:</u>

Wouter Hoogkamer

Integrative Locomotion Laboratory

Department of Kinesiology

University of Massachusetts, Amherst

Amherst, MA, 01003-9258 USA

Email: whoogkamer@umass.edu

ORCID:

Wouter Hoogkamer 0000-0003-0524-0196

Victor Rodrigo-Carranza 0000-0003-1637-7550

Laura Healey 0000-0003-0613-1629

Abstract

Background: Quantifying the potential benefits of advanced footwear technology (AFT)

track shoes (i.e., "spikes") in middle-distance events is challenging, because repeated

maximal effort trials (as in sprinting) or aerobic running economy trials (as in long-

distance running) are not feasible.

Methods: We introduce a novel approach to assess the benefits of AFT spikes, consisting

of a series of 200 m runs at self-perceived middle-distance race pace with 10 min

recovery and conducted four experiments to evaluate its validity, sensitivity,

reproducibility, and utility.

Results: In experiment 1, participants ran 1.2% slower in spikes with 200 g added mass

vs. control spikes, exactly equal to the known effects of shoe mass on running

performance. In experiment 2, participants ran significantly faster in AFT prototype

spikes vs. traditional spikes. In experiment 3, we compared two other AFT prototype

spikes against traditional spikes, on three separate days. Group-level results were

consistent across days, but our data indicates that at least two separate sessions are

needed to evaluate individual responses. In experiment 4, participants ran significantly

faster in two AFT spike models vs. traditional spikes (2.1% and 1.6%). Speed was similar

between a third AFT spike model and the traditional spikes. These speed results were

mirrored by changes in step length, as participants took significantly longer steps in the

two faster AFT spike models (2.3% and 1.9%), while step length was similar between the

other spikes.

Conclusion: Our novel, interval-based approach is a valid and reliable method to quantify

differences between spikes at middle-distance running intensity.

Key words: super spikes, track and field, running performance, carbon fiber plate,

athletics.

Declarations

Compliance with Ethical Standards

Ethical approval The study was performed in accordance with the ethical standards of

the Declaration of Helsinki. Ethics approval was obtained from the University of

Massachusetts Institutional Review Board (Protocol# 2647), from the University of New

Brunswick Research Ethics Board (REB #2021-079) and the University of Castilla-La

Mancha University Ethics Committee (Protocol CEIC924)

Informed consent Written informed consent was obtained from all individual

participants included in the study.

Funding

This research was partly supported by a research contract from PUMA SE with the

University of Massachusetts, Amherst. All spikes were provided by PUMA SE.

Conflicts of interest / Competing interests

Montgomery Bertschy, Victor Rodrigo-Carranza, Ethan W.C. Wilkie, Jeremy Noble and

Wayne J. Albert have no conflicts of interest relevant to the content of this article.

Wouter Hoogkamer is a paid consultant to PUMA and has received research grants from

PUMA and Saucony. Laura A. Healey is an employee of PUMA. No footwear company

had any influence on the conceptualization of this study or results presented in this

publication.

Data availability

All data discussed in this manuscript are provided in the tables.

Authors' contributions

Conceptualization: Wouter Hoogkamer; Experimental design: Montgomery Bertschy,

Laura A. Healey, Wouter Hoogkamer; Data collection: Montgomery Bertschy, Victor

Rodrigo-Carranza, Ethan W.C. Wilkie, Jeremy Noble; Data analysis: Montgomery

Bertschy, Victor Rodrigo-Carranza, Ethan W.C. Wilkie, Jeremy Noble, Wayne J. Albert;

Writing - original draft preparation: Montgomery Bertschy, Victor Rodrigo-Carranza,

Ethan W.C. Wilkie, Laura A. Healey, Wouter Hoogkamer; Writing - review and editing:

Montgomery Bertschy, Victor Rodrigo-Carranza, Ethan W.C. Wilkie, Laura A. Healey,

Jeremy Noble, Wayne J. Albert, Wouter Hoogkamer

1   Introduction

Once thought to be an unbreakable barrier on the track [1], the magical sub-4-

minute mile mark stood strong for decades, until Sir Roger Bannister in 1954 completed

the unthinkable and ran the 1609.344 meters in 3:59.4 seconds. In February 2023, 52

male professional, club, and collegiate level athletes broke the 4-minute mile barrier all

within the span of an hour at Boston University's indoor track facility. A similar

phenomenon was observed in the women's section, as three athletes broke the 4:30 mile

barrier, and 17 ran sub-4:36, a mark some consider to be more equivalent to the men's

barrier.[2] Over the past decades, the number of athletes breaking these barriers had been

increasing gradually, but there has been a substantial increase since 2021. While the

Boston University track has been known to be fast, another important common feature

among these performances was the use of advanced footwear technology (AFT) track

shoes (i.e., "spikes"), colloquially known as "super spikes". Similar to the now common

AFT marathon racing shoes (e.g., Nike Vaporfly), AFT track spikes feature a lightweight,

compliant, and resilient midsole foam, often combined with a stiff midsole plate. While

we have argued before that anecdotal performances should not be used as evidence on the

efficacy of specific shoes [3], the widespread improvement across middle-distance running

performances warrants a systematic, scientific controlled investigation.

The recent improvements on the track are similar to improvements in road

running performances since 2017 which have been partly attributed to advances in road

running shoes.[4–6] While the adoption of AFT road shoes happened in parallel with the

publication of a randomized crossover study showing 4% improvements in running

economy,[7] scientific evidence on potential benefits of AFT track spikes is currently

limited to race results analyses (preprint: [8]) and preliminary laboratory data.[9] In an

observational study, Willwacher et al.[8] found that AFT track spikes had a greater effect

on performance in middle-distance events (0.31 - 1.33%) than in sprint events (0.22 -

0.66%) for both men and women.

Traditionally, the effects of footwear on middle-distance running performance

have been studied in the lab by measuring running economy, or on the track with time

trials. Barnes and Kilding[10] observed that running economy in traditional lightweight

track spikes was ~3% worse than in AFT road shoes during treadmill running at speeds

between 3.9 and 5.0 m/s. Oehlert et al.[9] showed that different AFT track spikes improved

running economy by ~2% compared to a traditional spike model during treadmill running

at 14 km/h in women and at 16 km/h in men. Lab-measured changes in running economy

have been shown to translate to changes in running performance,[11] but for speeds faster

than 3 m/s, percentage improvements in running speed can be expected to be smaller than

observed percentage improvements in running economy.[12] Furthermore, running

economy measures are only valid when quantified during running at a sustainable,

aerobic, steady-state intensity. For very fit, elite participants, this could involve running

as fast as 5.8 m/s.[13,14] However, world record pace for middle-distance events ranges

from 6.2 m/s (women's 3000 m) to 7.9 m/s (men's 800 m). How footwear affects running

economy in high-caliber runners at 5.0 m/s (e.g., [7]) might not be indicative of how this

footwear affects running performance at a speed of 7.9 m/s, almost 60% faster. Such fast

middle-distance speeds require metabolic energy rates above aerobic steady-state

capacity and thus quantifying physiological responses to different spikes is not feasible or

valid (see [3]).

An alternative approach to quantifying the benefits of spikes could be to have

athletes perform a series of time trials in different spikes. Indeed, it has been shown

previously that 3,000 m time trial performances are slower in shoes with added mass.[11] In

that study, participants ran three-time trials on an indoor track, each at the same time of

day at the same day of the week, a week apart. Performing race-distance time trials at

maximal effort across multiple days can profoundly interfere with athletes' training

programming and alternatively doing multiple high-effort time trials on a single day,

would affect the validity of the protocol due to fatigue.

In this paper, we propose, evaluate, and use a novel approach to quantify the

potential benefits of middle-distance AFT spikes. We asked athletes to perform a series

of 200 m runs at self-perceived race pace (800 m or 1500 m, based on their specialty),

with 10 min recovery in between, within a single session. Such 200 m race pace efforts

are a common component of middle-distance training,[15,16] and all participants were

familiar with this type of workout. This is important because the premise of the proposed

protocol is to measure differences in performance, while controlling effort. In a series of

experiments, we compared up to four different spike models, while participants ran at

least two trials in each model. First, we evaluated the validity of our novel protocol by

comparing traditional spike models that were nearly identical and only differed in mass

(control vs. control + 200 g) by verifying the 200 m run time differences against the

theoretically expected differences. Second, we assessed the sensitivity of this novel

protocol by comparing new prototype AFT spikes against traditional spikes. Third, we

evaluated the reliability between three different days for two AFT spikes against

traditional spikes, and finally we used this protocol to compare several commercially

available AFT spikes against traditional spikes. We used an inertial measurement unit

(IMU) placed at the pelvis to assess if any differences in running speed were

predominantly driven by changes in step frequency or by changes in step length.

158 2. Materials and Methods

159 2.1 A Series of Experiments

This study involved four separate experiments across 3 different countries and

research groups, each with 12 participants. The inclusion criteria were generally similar

for all four experiments: fitting US women's size 7, 8.5 or 9, or US men's size 9 or 11

track spikes (Experiment 1) or fitting US men's size 9 track spikes (Experiments 2, 3 and

4); being 16 years of age or older (Experiment 1 and 2) or 19 years of age or older

(Experiment 3 and 4); currently participating in track and field training; having at

minimum two years of experience training in track spikes; being free of any orthopedic,

cardiovascular, or neuromuscular conditions; and having not sustained injury nor

undergone any surgery within the 3 months prior to participation. Recruitment of

participants was done through means of personal contacts, local track clubs, and local

university track teams. Participants were Tier 3: highly-trained, national level, but not

elite,[17] with average personal bests of 2:02 in 800 to 8:55 in 3000 m across the different

experiments (average World Athletics scoring tables points for each experiment: 815,

762, 817, 761, respectively (Table 1)).[18] Because of the different timelines of the

experiments and the increasing availability of AFT spikes with time, participants'

experience running with AFT varied between experiments: all participants had

experience running in spikes, for Experiment 1 we only used traditional spikes, for

Experiment 2 only two participants had run in AFT road shoes, none in AFT spikes, for

Experiments 3 and 4 all participants had experience running with AFT road shoes and/or

spikes. The characteristics of the participants in each experiment are shown in Table 1.

The experiments were performed in accordance with the ethical standards of the

Declaration of Helsinki. Ethics approval was obtained from the University of

Massachusetts, Amherst Institutional Review Board (Protocol # 2647), the University of

New Brunswick, Fredericton Research Ethics Board (REB # 2021-079), and the

University of Castilla-La Mancha University Ethics Committee (Protocol CEIC924).

Before taking part in the study, participants provided informed written consent.

2.1.1 Spike Conditions

For all experiments, we used the PUMA EvoSpeed Distance 9 (TRAD) as the

control condition; a typical traditional track spike, lightweight in construction, while

providing only minimal cushioning and no added bending stiffness elements. It has a

low-profile ethylene vinyl acetate foam with a flat geometry, a midsole thickness of

17mm and 5 spike pins per shoe (Fig 1). For all spike conditions, midsole thickness was

measured at the heel and the forefoot, and reported as the largest number among those

locations. In Experiment 1, we compared the standard version of this spike (TRAD, 137-

169 g) against the same spikes with 200 g of added mass (TRAD+, 331-368 g) (Table 2).

Mass was added to the spikes by means of seven 28.3 g (1 oz) weights glued to the sides

of the shoes (Fig. 1).

In Experiment 2, we compared two track spike conditions: TRAD and a PUMA

AFT prototype middle-distance spike (AFTP1, 179 g). The AFTP1 spike combines a

compliant, highly resilient midsole foam with an embedded carbon fiber plate. It features

PUMA's Nitro foam, a lightweight polyether block amide (PEBA) foam compound,

forming a rockered midsole with a thickness of 19 mm, and 6 spike pins per shoe (Fig 1;

Table 2).

In Experiment 3, we compared three track spike conditions: TRAD, a PUMA

AFT prototype long-distance spike (AFTP2, 151 g) and a PUMA AFT prototype middle-

distance spike (AFTP3, 149 g, different from AFTP1). The AFTP2 spike features Nitro

foam, forming a rockered midsole with midsole thickness of 18 mm, but no carbon fiber

plate. The AFTP3 spike combines a compliant, highly resilient midsole foam with an

embedded carbon fiber plate. It features Nitro foam, forming a rockered midsole with a

thickness of 19 mm, and 6 spike pins per shoe (Fig 1; Table 2).

In Experiment 4, we compared four commercially available track spike

conditions: TRAD, a commercially available PUMA AFT spike (AFTC1) and two

competitor brand AFT spikes (AFTC2 and AFTC3). The AFTC1 was the PUMA

evoSPEED Long Distance Nitro Elite+ (154 g). It features Nitro foam, forming a

rockered midsole with a thickness of 18 mm, but no carbon fiber plate, and 6 spike pins

per shoe. The AFTC2 was the Nike ZoomX Dragonfly (140 g), featuring ZoomX foam, a

lightweight PEBA foam compound, forming a rockered midsole with a thickness of 12

mm, but no carbon fiber plate, and 6 spike pins per shoe. The AFTC3 was the Adidas

adizero Avanti TYO (163 g), featuring Lightstrike Pro foam, a lightweight thermoplastic

polyether elastomer foam, forming a rockered midsole with a thickness of 17 mm, with

embedded fiber glass rods for increased bending stiffness, and 6 spike pins per shoe (Fig

1; Table 2).

2.1.2 Blinding Procedures

A single-blinded crossover methodology was implemented as part of the protocol

for Experiments 1, 2 and 4 (i.e., participants were blinded, researchers were not). For all

experiments the participants were unaware of any differences between spike conditions

and which properties were modified, since the informed consent documents and

researchers did not mention spike mass, foam or plate properties, only that there were

several different conditions. Participants were also kept unaware of the number of

different conditions or that they ran multiple times in the same condition. In none of the

experiments the participants were allowed to manipulate the spikes.[11,19] Further, for

Experiments 1, 2, and 4, we used fabric boot covers (Fig. 1) such that participants did not

see the spikes. These spike blinders fit around the participants spikes and lower legs in a

non-restrictive fashion, providing a negligible addition to spike mass. Through an

oversight, we did not use these spike blinders in Experiment 3. For this experiment, spike

conditions were of a similar color/visual design, and similar to the other experiments,

participants were unaware of potential differences between the spikes, and were not

allowed to manipulate the spikes.

2.1.3 Experiment 1 - Validation
To evaluate the external validity of our novel approach, we compared 200 m time

in control spikes versus spikes that were nearly identical and only differed in mass

(control vs. control + 200 g). Similar to Hoogkamer et al.,[11] we used a mass intervention

to predictably induce a change in running energetics.[20,21] Frederick et al.[20] showed that,

across a range of speed, adding 100 g mass to each shoe increases the energy cost of

running by ~1%. The amount that a 1% increase in energy cost will slow down a runner

depends on how fast they are running. Kipp et al.[12] provided a theoretical framework

supported by observations in Hoogkamer et al.[11] that predicts that when running at

6.67m/s (200 m in 30 s), a 1% increase in energy cost will lead to a 0.6% slower run.

Therefore, with 200 g added to each spike, we hypothesized that participants would run

1.2% slower.

2.1.4 Experiment 2 - Sensitivity

Next, we evaluated if our protocol had the sensitivity to detect differences in what

may be seen as "anecdotal" benefits of AFT spikes over traditional spikes. For this newly

developed prototype middle-distance spike, we anticipated that the metabolic savings

from the modern midsole foam and the embedded plate would add up to about 2%, based

on preliminary running economy results from Oehlert et al.[9] Therefore, we hypothesized

that participants would run 1.2% faster in the prototype spikes.

2.1.5 Experiment 3 – Reliability

We then evaluated the reliability of our novel approach to quantify the potential

benefits of different AFT spikes technologies. The objective was to assess the

consistency of the benefits from two AFT spikes over TRAD across three separate days.

We hypothesized that the benefits of both AFT spikes would be consistent across the

three measurement days.

2.1.6 Experiment 4 - Commercially Available Spikes

Finally, we used our novel approach to quantify potential benefits of a set of

different commercially available AFT spikes vs. traditional spikes. The aim was to

evaluate if potential performance gains from AFT spikes differed between models. We

hypothesized that the different AFT spikes all would result in 1.2% faster times than

TRAD, with no detectable differences between AFT models. Further, we hypothesized

that step frequency would be similar between conditions, while step length would be

about 1% longer in the AFT spikes.[7,22]

2.2 Generic Experimental Protocol

2.2.1 General protocol

All experiments took place at an outdoor 400 m track facility on days with little to

no wind. The middle-distance track runners enrolled in the study performed a series of

200-meter runs at self-perceived 800 m or 1500 m race pace with 10 min recovery (to

allow for changing spikes (seated) and walking to the start of the next trial). The

effectiveness of the protocol depended on the participants' ability to run every trial at a

similar sub-maximal effort. To minimize bias, it is important that the participants are

unaware of the spike intervention and that they are able to control their effort at a

consistent submaximal level. Aiming to achieve the latter, we opted for a series of 200 m

runs at self-perceived race pace, since this is a common component of middle-distance

training,[15,16] and all participants, like most middle-distance runners, were familiar with

this type of workout, although recovery times in this experiment (10 min) were likely

longer than during a real workout.

To minimize any confounding effects of participants running the first

(excitement) or last ("emptying the tank") trials at a higher effort, the number of trials

that the researchers told the participants to run was greater than the number of trials they

would actually run. Their first trial was considered a habituation trial and not included in

the evaluation, and then after running all the needed experimental trials, the researchers

revealed to the participants that the data collection was complete. Experiment 1 consisted

of six experimental trials (three trials in both conditions), and we initially told

participants that they would be expected to run eight, 200 m trials with 10 min recovery,

but then we revealed to them that the experiment was complete after seven trials.

Similarly, for Experiments 2 and 4 we asked participants to run ten trials, but we only

included four and eight experimental trials, respectively (two trials in each spike

condition). For Experiment 3, the participants performed the same protocol on three

separate visits, so they ran all eight trials, but we only analyzed the last six experimental

trials.

The order of spike conditions was randomized for each participant prior to each

session. Participants performed each 200 m trial from a standing start (similar to a

middle-distance race start). Participants were unaware of the times they were running for

each trial as they were not using a watch and researchers did not disclose the times they

recorded until after the data collection was complete. Rest between trials was 10 minutes,

to allow for changing spikes, filling out questionnaires (Experiments 2 and 4), and

walking to the start of the next trial. Limited pilot data (n = 2) showed that venous blood

lactate concentration did not increase over the course of an experimental session, with

consistent values of ~1.5 mmol/l before the start of the next trial.

2.2.2 Experimental Setup

When the participant arrived at the track facility, we explained the study protocol

to them, and they signed all the necessary forms [i.e., the Get Active Questionnaire[23] or

the Physical Activity Readiness Questionnaire (ePARmed-X+) and the informed consent

document]. We asked participants to run a warmup similar to how they would for a 200

m interval workout, followed by drills and stretches as they saw fit. For Experiment 4,

after their warmup, we placed an inertial measurement unit (IMU; Vicon Blue Trident

sensor, Vicon, Oxford, UK) at the participant's sacrum with a clip attached to the back of

the waistband on their shorts.[24] Next, the participant removed their shoes and researchers

put fabric spike blinders on the participant's legs (Experiments 1, 2 and 4). Then the

researchers put the spikes on the participant's feet and tightened the laces. Following this,

spike blinders were pulled down over the spikes.

At the 200 m line, the researcher started the IMU recording (Experiment 4) and

then moved towards the finish line. For the timing of each trial, researchers started their

timers when the participant started running, indicated by them quickly lowering one hand

from a raised position, and stopped their stopwatches when the participant crossed the

finish line. After a trial was complete, the 200 m time was recorded and the IMU

recording was stopped (Experiment 4). This was repeated for all trials.

2.3 Data Analyses and Statistics

We calculated average running speed in m/s by dividing the 200 m distance by the

200 m time. Raw IMU data was used to calculate spatiotemporal variables in Experiment

4. Orientation of the IMU was verified from acceleration signal by finding the

gravitational component (-9.81 m/s^2) during quiet standing before the running trial began.

The vertical component was filtered using a 5 Hz cut-off frequency and multiplied by

participant mass to create a vertical ground reaction force estimation, as demonstrated by

Day et al.[24] The time between peaks in the vertical ground reaction force estimate were

used to calculate step time and its inverse, step frequency over the course of the full

running trial. Here, step refers to the period from one foot strike to the contralateral foot

strike. Average step length was calculated as the product of average speed and average

step time.

We used linear-mixed effect models (LMEM) for each experiment.[25] For all

experiments, we created a LMEM with trial number as a fixed effect, participant as a

random intercept, and speed as an output variable to determine if there was a significant

effect of trial number on speed (i.e., slowing down or speeding up during the

experiment). This was only significant for Experiment 1 (p = 0.006), so for this

experiment we included trial number as a fixed effect: we created a LMEM with spike

condition and trial number as fixed effects, participant as a random intercept, and speed

as the output variable. For Experiment 2, we created a LMEM with spike condition as a

fixed effect, participant as a random intercept, trial as a random effect, and speed as the

output variable. For Experiment 3, we created a LMEM with day as a fixed effect,

participant as a random intercept, and speed as an output variable to determine if there

was a significant effect of day on speed. Since this was significant (p = 0.032), day was

included as a fixed effect: we created a LMEM with spike condition and day as fixed

effects, participant as a random intercept, trial as a random effect, and speed as the output

variable. Additionally, for each day of this experiment, we created a LMEM with spike

conditions as a fixed effect, participant as a random intercept, trial as a random effect,

and speed was the output variable to verify that the results did not change per day. For

Experiment 4, we created three LMEM with spike condition as a fixed effect, participant

as a random intercept, and speed, step length, and step frequency as the output variables,

respectively. A traditional level of significance ($\alpha = 0.05$) was employed. Tukey HSD

tests were conducted for each pairwise comparison of spike conditions. Additionally, we

calculated effect size (ES) as Cohen's d and reported percent differences for each

measure with the intention of providing relevant context for our determined p values. ES

with values of 0.2, 0.5, and above 0.8 were considered as small, medium, and large,

respectively.[26] To evaluate the reliability, in Experiment 3 we calculated the coefficient

of variation (CV = SD / mean × 100%) within each day and for each spike, across days.

The ranges of values considered for categorizing CV% were as follows: CV < 10 as very

good, 10–20 as good, 20–30 as acceptable, and CV > 30 as unacceptable.[26] Further, for

each participant we calculated the average speeds in each spike across days 1 and 2, days

1 and 3, days 2 and 3, and across days 1, 2 and 3. Then, we calculated the speed

difference between the TRAD and the AFTP3 spikes. Next, we calculated Pearson's

correlation coefficients, for these differences, between each individual day and the 3-day

average, and between each 2-day average and the 3-day average. Correlation coefficients

with values of 0.1, 0.3, and 0.5 were considered as small, medium, and large,

respectively, with 1.0 being a perfect correlation[26]. Statistical analyses were performed in

R using lme4 package[27] for the linear mixed effect models and emmeans package[28] for

post hoc Tukey HSD tests. Descriptive statistics are reported as mean ± standard

deviation (SD). For visual display in the figures, we averaged the 200 m trial times that

were performed in the same spike condition for each experimental session (three trials for

Experiment 1; two trials for Experiments 2, 3 and 4).

3 Results

In Experiment 1, participants ran significantly slower in the TRAD+ spikes (5.97

± 0.52 m/s) than in the TRAD spikes (6.04 ± 0.50 m/s) (1.2%, ES = 0.14, p = 0.016, n =

12; Fig. 2). Participants significantly slowed down during the experimental session (0.05

m/s per set of two trials, p = 0.004). In Experiment 2, participants ran significantly faster

in the AFTP1 spikes (6.35 ± 0.42 m/s) than in the TRAD spikes (6.28 ± 0.50 m/s) (1.1%,

ES = 0.15, p = 0.030, n = 12; Fig. 2).

In Experiment 3, participants ran significantly faster in the AFTP2 spikes (6.52 ±

0.37 m/s) and the AFTP3 spikes (6.60 ± 0.36 m/s) than in the TRAD spikes (6.40 ± 0.34

m/s) (1.8%, ES=0.34, p < 0.0001 and 3.1%, ES= 0.57, p < 0.0001, respectively, n = 12;

Fig. 2). Running speeds in AFTP3 spikes were significantly faster than in AFTP2 spikes

(1.2%, ES=0.19, p = 0.0003). The between-spike results were consistent between days.

The CV% values were classified as very good for inter-session (5.5%, 5.4% and 5.7% for

day 1, 2 and 3 respectively) and intra-session (5.3%, 5.5% and 5.5% for TRAD, AFTP2

and AFTP3 respectively). The linear mixed effects model analyses for each day

separately all showed similar differences between spikes, both in magnitude (% change

and effect size) and in significance (Table 3).

The speed difference between TRAD and AFTP3 for each individual participant

varied across days. Pearson's correlation coefficients between day 1, day 2 and day 3,

and the 3-day average were large: 0.80, 0.77, and 0.89, respectively. However, the

Pearson's correlation coefficients between each 2-day average and the 3-day average

were substantially higher: 0.97, 0.90 and 0.96 for day 1 and 2, day 1 and 3, and day 2 and

3, respectively.

In Experiment 4, participants ran significantly faster in the AFTC1 spikes (6.44 ±

0.45 m/s) and the AFTC2 spikes (6.41 ± 0.51 m/s) than in the TRAD spikes (6.31 ± 0.47

m/s) (2.1%, ES= 0.31, $p = 0.002$ and 1.6%, ES= 0.22, $p = 0.035$, respectively, n = 12;

Fig. 2). Running speeds in the AFTC3 spikes (6.36 ± 0.48 m/s) were not significantly

different from the TRAD spikes (0.8%, ES=0.13, $p = 0.493$). Running speeds were not

significantly different between the AFTC spikes (AFTC1 vs. AFTC2 $p = 0.800$; AFTC1

vs. AFTC3 $p = 0.116$; AFTC2 vs. AFTC3 $p = 0.534$), but the numerical difference

between AFTC1 and AFTC3 was more than 1% (1.3%, ES = 0.17).

Step frequency was not different between spikes (all $p > 0.52$; Fig. 2), but

participants ran with significantly longer steps in AFTC1 spikes (1.93 ± 0.15 m) and

AFTC2 spikes (1.92 ± 0.16 m) than in the TRAD spikes (1.88 ± 0.15 m) (2.3%, ES =

0.33, $p = 0.001$; 1.9%, ES = 0.26, $p = 0.008$; Fig. 2). Step lengths in the AFTC3 spikes

(1.91 ± 0.15 m) were not significantly different from the TRAD spikes (1.5%, ES = 0.20,

$p = 0.0512$). Step lengths were not significantly different between the AFTC spikes

(AFTC1 vs. AFTC2 $p = 0.893$; AFTC1 vs. AFTC3 $p = 0.472$; AFTC2 vs. AFTC3 $p =$

0.875).

4 Discussion

In this paper, we proposed, evaluated and used a novel approach to quantify the

potential benefits of middle-distance AFT spikes. The combined results indicate that our

novel approach has the validity, sensitivity and reliability to quantify the benefits of

middle-distance AFT spikes. The results of Experiment 1 provide insights into the

validity of our interval-based approach. On average, participants ran 1.2% slower in the

spikes with 200 g added mass, similar to the 1.2% difference we hypothesized. The

results of Experiment 2 indicate that our approach has the sensitivity to confirm what so

far has been considered "anecdotal" benefits of AFT spikes over traditional spikes.

Participants ran significantly (1.1%) faster in AFT prototype spikes than in traditional

spikes. The results of Experiment 3 indicate that our approach is reliable. Across three

separate days, the between-spike differences in running speed were similar, both in

magnitude (% change and effect size) and in significance. However, at the level of

individual runners, the responses to AFT spikes varied between days. Our data suggests

that two separate sessions are needed to obtain reliable individual data. Finally, we used

our interval-based approach to compare several commercially available AFT spikes

against traditional spikes. In two AFT spike models participants ran significantly (2.1%

and 1.6%) faster than in the traditional spikes. Speed was similar between a third AFT

spike model and the traditional spikes. These speed results were mirrored by changes in

step length, as participants took significantly (2.3% and 1.9%) longer steps in the two

faster AFT spike models, while step length was similar between the third AFT spike

model and the traditional spikes.

4.1 Construct Validity

Shoe mass is one of the most studied features of running shoes.[11,20,21,29] There is currently a strong consensus that there is a linear relationship between the increase in shoe mass and the increase in running energy cost, with an increment in the oxygen/energy cost of ~1% per added 100 g of shoe mass.[20] Kipp et al.[12] provided a theoretical framework to translate changes in energy cost into changes in running performance, based on the known curvilinear relationship between running speed and energy cost and the cost of overcoming air resistance, and supported by shoe-mass time trial results from Hoogkamer et al.[11] Using this framework, we anticipated that participants would run 1.2% slower in the TRAD+ spikes with 200 g added mass. We found that on average our participants ran 1.2% slower, with 75% of the participants (9 of 12) running slower in the TRAD+ spikes. Our participants ran slightly slower than anticipated (~6.0 m/s vs. 6.67 m/s) and at that speed the framework predicts a 1.3% slower speed in the TRAD+ spikes (not 1.2% slower). While our results are close to our prediction, it should be acknowledged that our predictions are based on the 1% for 100 g shoe mass rule of thumb and that this 1% number slightly varies between studies,[11,20,21,29] and that it might be less than 1% at faster speeds.[20]

4.2 Sensitivity

Next, we evaluated if our interval-based approach is sensitive enough to detect differences between AFT spikes and traditional spikes. We anticipated that speed differences between AFT spikes and traditional spikes would be similar to the effect of 200 g added mass and we hypothesized that participants would run 1.2% faster in the

AFT prototype spikes. Participants ran 1.1% faster in the AFT prototype spikes than in

the traditional spikes, which is equivalent to the effect of reducing the mass of the

traditional spikes by ~180 g per spike (which is not possible, considering that the

traditional spikes weigh ~150 g each).

4.3 Reliability

We evaluated the reliability of this protocol by asking 12 participants to run in the

same three spike conditions with the same protocol on three separate days. We found

that, at the group level, results were similar between all three days (Table 3) and similar

to the 3-day average values (significant speed improvements of 1.8 and 3.1% of the

AFTP2 and AFTP3, compared to TRAD, respectively). The Pearson correlation

coefficients for Day 1, 2, and 3 vs. the 3-day average were large (0.77-0.89), but the

correlation coefficients for 2-day average vs. 3-day average results were substantially

higher (0.90-0.97), suggesting that a minimum of two days is needed to quantify

dependable individual responses to AFT spikes. Quantifying individual responses to AFT

shoes is common when aiming to identify biomechanical mechanisms underlying

improvements from AFT,[30,31] or characteristics of runners who experience more or less

benefits.[32–34] However, as discussed by Barrons et al.,[35] such analyses critically depend

on the accuracy of the response measurement. Note that both for running economy

measures[35] and for speed outcomes from our interval-based approach, multiple sessions

with repeated conditions are recommended, when attempting to quantify individual

responses to AFT. In our study, 75% of the participants ran consistently faster in AFTP2

vs. TRAD across the three days, and similarly 75% of the participants ran consistently

faster in AFTP3 vs. TRAD across the three days (Fig. 3). For AFTP2 vs. TRAD, two

participants (17%) ran faster in two of the three days, and one participant (8%) only ran

faster on one day. For AFTP3 vs. TRAD, three participants (25%) ran faster on two of

the three days. Nine participants (75%) ran faster in AFTP3 vs. AFTP2, and eight of them

did this consistently across the three days (day 1 for participant #6 being the exception).

For the three participants who ran faster in AFTP2 vs. AFTP3 (#8, 9 and 10), two of them

did this consistently across the three days (day 1 for participant #8 being the exception).

Noteworthy, the footwear condition with plate (AFTP3) lead to significant improvements

over the condition with similar construction without plate (AFTP2). This is in contrast to

Healey and Hoogkamer[36] who observed that making medio-lateral cuts in the carbon-

fiber plates of AFT shoes (i.e., effectively reducing the shoes' longitudinal bending

stiffness) did not significantly worsen running economy. This discrepancy might be

related to the different speeds run in each experiment (~6.50 m/s for Experiment 3 vs.

3.89 m/s in Healey and Hoogkamer [36]), suggesting that increased longitudinal bending

stiffness is more beneficial at faster running speeds.

4.4 Quantifying the Benefits of AFT Spikes

Finally, we used our novel approach to quantify potential benefits of several

different commercially available AFT spikes over traditional spikes. Participants ran

significantly faster in the AFTC1 (2.1%) and AFTC2 (1.6%) than in the TRAD spikes.

However, running speeds were not significantly different in the AFTC3 spikes versus the

TRAD spikes (0.8%), nor between the AFTC spikes. The improvements from the

commercially available AFT spikes compared to the traditional spikes were similar to,

but slightly less than, those of the prototype AFT spikes evaluated in Experiment 3 of this

study (1.8 - 3.1%). Over 800 or 1500 m at these speeds, athletes would run 2.5 or 5 sec

faster, respectively, in these commercially available AFT spikes than in traditional spikes.

While the effects of spikes on sprinting performance have been studied for at least

50 years, with an inquiry into the potential benefits of the banned Puma brush spikes in

the early seventies[37], here we specifically focused on middle-distance running spikes.

Given the novelty of AFT spikes, and the challenges of quantifying their potential

benefits on middle-distance running performance,[3] we are not able to directly compare

our findings to any previous studies in the literature.

Willwacher et al.[8] performed a longitudinal, observational, and retrospective

study design evaluating the 100 best performances per year for men and women in

outdoor track events from 2010 to 2022. They observed more pronounced improvements

for the 1500 m and longer distances (0.7 - 1.8%) than shorter distances (100 m to 800 m:

0.3-0.6%) for women, and less of an effect of AFT spikes in men. The differences at the

group level in the 800 m were ~0.7 sec for both men and women, and ~1.3 and ~2.2 sec

in the 1500 for men and women, respectively. Substantially smaller than the 2.5 and 5 sec

time savings that our data indicate. When comparing running economy results for AFT

spikes at 4.4 and 5.0 m/s, Oehlert et al.[9] found 1.8 and 2.1% improvements for two AFT

spike models as compared to traditional spikes. Such running economy improvements

can be expected to improve middle-distance running performance by ~1.2%,[12] which is

less than the speed improvements we observed in the current study at ~6.4 m/s. Although,

not specific to AFT spikes, some earlier studies have evaluated 3,000 m time trial

performance on the track in AFT road shoes and reported improvements ranging from 1

to 5%, similar to our spike results.[33,38] Without mechanical characterization of midsole

compliance and resilience, and longitudinal bending stiffness of the different commercial

AFT spike models (beyond the scope of this study), we can only speculate why

participants only ran significantly faster in AFTC1 and AFTC2 vs. TRAD, and not in

AFTC3. It might be that the compliance and resilience properties of the thermoplastic

polyether elastomer foam in the AFTC3 are less than that of the PEBA-based foams in

AFTC1 and AFTC2, or that the bending stiffness of AFTC3 is suboptimal. Importantly,

the exact benefits of increased longitudinal bending stiffness within AFT shoes are still

under debate,[39,40] with some evidence that the improved foam properties are more

important.[22,36,41]

Across Experiment 2, 3 and 4, participants consistently ran faster in the AFT

spikes than in the traditional spikes (with the exception of AFTC3). The three

experiments each used slightly different models of AFT spikes and the group-level

benefits varied from 1.1% in Experiment 2, 1.8 and 3.1% in Experiment 3 to 2.1 and

1.6% in Experiment 4. The consistency of these findings, while these experiments were

performed by different research teams in three different countries (US, Canada and

Spain) is a testament of the robustness of our interval-based approach and further

supports the validity of the observed benefits of AFT spikes.

Intriguingly, the faster speeds in the AFTC1 (2.1%) and AFTC2 (1.6%) spikes as

compared to the TRAD spikes in Experiment 4, were accompanied by 2.3% (i.e., 4.3 cm)

and 1.9% (i.e., 3.6 cm) longer steps in the two faster AFTC1 and AFTC2 spikes,

respectively. Step lengths were similar between the other spike models. Furthermore, step

frequency was similar across all spikes (~3.35 Hz, or 201 steps per minute). Combined,

this suggests that the performance benefits from AFT spikes in middle-distance running

originate from the spikes allowing runners to take longer step lengths without requiring

any compensation in step frequency. This is supported by observations of runners taking

longer steps at fixed speeds in AFT road shoes vs. traditional shoes.[7,22,42] To confirm this,

future studies could compare step lengths in AFT spikes vs. traditional spikes for running

at a fixed, middle-distance running speed. With average step lengths of 1.93 m and 1.92

m in the AFTC1 and AFTC2 spikes, the participants took only 103.6 steps and 104.1

steps during their 200 m, respectively, about 2.5 steps less than in the traditional spikes.

Over a 1500 m race this would add up to 21 or 17 less steps in the AFTC1 and AFTC2

spikes than in traditional spikes, respectively.

These biomechanical changes are similar to those reported in treadmill studies on

AFT road shoes but different from preliminary findings for AFT spikes. Treadmill

studies of road shoes consistently report that runners take ~1% longer steps in AFT shoes

than in traditional shoes.[7,10,22,42] Since running speed is fixed between conditions in such

studies, these longer steps are accompanied by reductions in step frequency. In a case

study on overground, track running, Russo et al.[43] evaluated biomechanics during 80 m

trials at 95% of maximal speed comparing AFT spikes with a carbon-fiber plate and a

modern midsole foam against traditional spikes. The athlete ran at the same speed (~8.0

m/s) and did not substantially change step length or step frequency, even though contact

time was longer in the AFT spikes. The fixed speed and single subject nature of Russo et

al.[43] might explain the differences with our study.

4.5 Limitations and Future Directions

While our approach of evaluating a series of 200 m runs at a self-perceived race

pace, interferes less with athletes' training programming and is less sensitive to day-to-

day variability than performing race-distance time trials at maximal effort across multiple

days, self-perceived sub-maximal efforts are not a direct measure of running performance

and sensitive to bias. For Experiments 2-4, it may be possible that the participants ran

faster in the AFT spikes for 200 m simply because the AFT spikes felt bouncier or more

comfortable, or because of their belief in performance enhancement from AFT spikes.

These faster sub-maximal 200 m times might not translate into any true performance

gains over a full 800 m or 1500 m race at maximal effort. However, results for

Experiment 1 show that participants quite precisely adjust their speed according to the

expected metabolic energy increase from the added mass, despite the identical underfoot

construction of the two spike conditions (TRAD and TRAD+). Furthermore, our

participant blinding procedures were effective in that participants were typically unaware

of how many different spike conditions they were running in, and what spikes they were

currently wearing. In general, any speculation by the participants about their shoe

conditions were inaccurate, and researchers neither confirmed nor denied their comments.

Future further proof of validity could come from experiments that directly compare

between-spike differences observed with our interval-based approach to between-spike

differences in race distance time trials.

For researchers looking to implement our interval-based protocol in future

studies, great care must be taken to limit the bias of the researchers and participants.

Participants should be blinded to the shoe conditions, not just to what is currently on their

feet. If, for example, the participant knows that they will be wearing one traditional spike

and one AFT spike, the participant might be able to use this information to correctly

deduce the spike condition they are currently wearing. With AFT spikes becoming more

common, especially when comparing commercially available spikes, some participants

might be able to feel or recognize differences in midsole cushioning or longitudinal

bending stiffness between spikes, which could bias their performances in our effort-based

protocol. We recommend that investigators mirror the order of spike conditions (e.g.,

ABCCBA) and counterbalance condition orders across participants (i.e., each possible

order is done, and all orders are done by a same number of participants), to minimize any

order effects. Participants ran their 200 m trials at consistent speeds throughout the test

session for Experiments 2-4, but in Experiment 1 they slowed down significantly from

the first set of trials in each condition to the third set of trials in each condition.

While our study shows that our novel interval-based approach to quantify

potential benefits of middle-distance AFT spikes is valid and reliable at the group level,

we caution against using the results of a single session to label an individual as a

responder/non-responder to AFT spikes, or to compare differences in responses between

individuals. Specific to that purpose, we found that at least two days are necessary to be

able to evaluate individual responses to AFT spikes.

Additionally, there are limitations with the implementation of this protocol. In this

current study, running times were recorded by hand, which is subject to human error.

Step lengths were calculated as the average of the 200 m run, based on the average step

frequency and the running time. Future studies could place IMUs at the feet, which would

allow accurate calculation of stride length at a stride-by-stride resolution.[44] Evaluating

between-spike differences in stride length at a stride-by-stride resolution could provide

additional information about how benefits of AFT spikes potentially vary from the

acceleration phase to the top speed phase.

Finally, caution should be taken when interpreting the results on the benefits of

AFT spikes in Experiment 4. On average, the male participants in this experiment ran

their trials at ~6.4 m/s, which is equivalent to a 2:05 min 800 m, or 3:54 min 1500 m. To

evaluate the benefits at elite or even world record level, this experiment should be

repeated with elite runners. Still, our findings at ~6.4 m/s are likely more informative

than running economy measurements for running at 16 km/h (4.44 m/s; equivalent to a

3:00 min 800 m). Elite and/or female runners might have different plantar flexor strength

or body mass than our male participants, factors that are likely to affect responses to

AFT.[45,46] At the start of our study, we only had spikes available in the men's sample size

US9 (except for Experiment 1), but follow-up studies should include more female

participants.

5 Conclusions

To overcome the challenges in quantifying the potential benefits of advanced

footwear technology spikes related to the physiology of middle-distance running

intensity, we introduced a novel interval-based approach consisting of a series of 200 m

runs at self-perceived middle-distance race pace. In a series of experiments, we showed

the validity and reliability of our method and then used it to compare commercially

available spikes. Our reliability experiment showed that group-level results were

consistent across days, but that at least two separate sessions are needed to evaluate

individual responses. Participants ran ~2% faster and with ~2% longer steps in two

commercially available AFT spike models than in traditional spikes, while speed and step

length were similar between a third AFT spike model and the traditional spikes.

Acknowledgements

We thank the athletes of Athletics Canada and the UNB Reds Varsity Track and Cross

Country team, and of the running communities in Amherst, Boston, San Antonio, Castilla

La-Mancha and Madrid for their participation in our study. We thank Rodger Kram and

Jose van der Veen for energetic(s) discussions to spark the ideas presented in this paper.

We thank Dan Feeney for guidance regarding our statistical analysis. Lastly, we thank

our undergraduate researchers, Elizabeth Hoenes, Caleb Zylka, Violeta Muñoz de la Cruz

and Robert Carew for assisting with data collections and recruitment. All spikes were

provided by PUMA SE.

References

1.   Taylor B. What breaking the 4-minute mile taught us about the limits of
674   conventional thinking. *Harv Bus Rev* 2018.
675   2.   Martin Fritz Huber. The four-minute mile is still worth celebrating. *Outside*
676   2023;
677   3.   Healey L, Bertschy M, Kipp S, Hoogkamer W. Can we quantify the benefits of
678   "super spikes" in track running? *Sports Medicine* 2022;**52**:1211–8.
679   https://doi.org/10.1007/s40279-022-01657-4.
680   4.   Senefeld JW, Haischer MH, Jones AM, Wiggins CC, Beilfuss R, Joyner MJ, et al.
681   Technological advances in elite marathon performance. *J Appl Physiol*
682   2021;**130**:2002–8. https://doi.org/10.1152/japplphysiol.00002.2021.
683   5.   Rodrigo-Carranza V, González-Mohíno F, Santos-Concejero J, González-Ravé
684   JM. Impact of advanced footwear technology on elite men's in the evolution of
685   road race performance. *J Sports Sci* 2023;**40**:2661–8.
686   https://doi.org/10.1080/02640414.2023.2183103.
687   6.   Rodrigo-Carranza V, González-Mohíno F, Santos del Cerro J, Santos-Concejero
688   J, González-Ravé JM. Influence of advanced shoe technology on the top 100
689   annual performances in men's marathon from 2015 to 2019. *Sci Rep*
690   2021;**11**:1–9. https://doi.org/10.1038/s41598-021-01807-0.
691   7.   Hoogkamer W, Kipp S, Frank JH, Farina EM, Luo G, Kram R. A comparison of
692   the energetic cost of running in marathon racing shoes. *Sports Medicine*
693   2018;**48**:1009–19. https://doi.org/10.1007/s40279-017-0811-2.
694   8.   Willwacher S, Mai P, Helwig J, Hipper M, Utku B, Robbin J. Does advanced
695   footwear technology improve track and road racing performance? An
696   explorative analysis based on the 100 best yearly performances in the world
697   between 2010 and 2022: The effects of super-footwear on world-class track
698   and road racing performances. *Sports Medicine-Open* 2024;**10**.
699   https://doi.org/10.1186/s40798-024-00683-y.
700   9.   Oehlert GM, Jones EJ, Burns GT, Joubert DP. Comparative effects of Advanced
701   Footwear Technology on running economy in track spikes and racing shoes.
702   *Med Sci Sports Exerc* 2023;**55**:41–2.
703   10.   Barnes KR, Kilding AE. A randomized crossover study investigating the
704   running economy of highly-trained male and female distance runners in
705   marathon racing shoes versus track spikes. *Sports Med* 2019;**49**:331–42.
706   https://doi.org/10.1007/s40279-018-1012-3.
707   11.   Hoogkamer W, Kipp S, Spiering BA, Kram R. Altered running economy directly
708   translates to altered distance-running performance. *Med Sci Sports Exerc*
709   2016;**48**:2175–80. https://doi.org/10.1249/MSS.0000000000001012.
710   12.   Kipp S, Kram R, Hoogkamer W. Extrapolating metabolic savings in running:
711   Implications for performance predictions. *Front Physiol* 2019;**10**:1–8.
712   https://doi.org/10.3389/fphys.2019.00079.
713   13.   Lucia A, Esteve-Lanao J, Olivan J, Gomez-Gallego F, San Juan AF, Santiago C, et
714   al. Physiological characteristics of the best Eritrean runners—exceptional
715   running economy. *Appl Physio Nutr Metab* 2006;**31**:530–40.

716   14.   Joyner, Hunter SK, Lucia A, Jones AM. Physiology and fast marathons. *J Appl*
717   *Physiol* 2020;**128**:1065–8. https://doi.org/10.1152/japplphysiol.00793.2019.
718   15.   Daniels J. Daniels' running formula. Human Kinetics, 2013.
719   16.   Haugen T, Sandbakk Ø, Seiler S, Tønnessen E. The training characteristics of
720   world-class distance runners: an integration of scientific literature and
721   results-proven practice. *Sports Med Open* 2022;**8**:1–18.
722   17.   McKay AKA, Stellingwerff T, Smith ES, Martin DT, Mujika I, Goosey-Tolfrey VL,
723   et al. Defining training and performance caliber: a participant classification
724   framework. *Int J Sports Physiol Perform* 2022;**17**:317–31.
725   https://doi.org/10.1123/IJSPP.2021-0451.
726   18.   Spiriev B. Scoring tables of athletics. World Athletics 2022 revised edition.
727   2022.
728   19.   Slade SJ, Greenya JG, Kliethermes CL, Senchina DS. Somatosensory perception
729   of running shoe mass. *Ergonomics* 2014;**57**:912–20.
730   20.   Frederick EC. Physiological and ergonomics factors in running shoe design.
731   *Appl Ergon* 1984;**15**:281–7.
732   21.   Franz JR, Wierzbinski CM, Kram R. Metabolic cost of running barefoot versus
733   shod: Is lighter better? *Med Sci Sports Exerc* 2012;**44**:1519–25.
734   https://doi.org/10.1249/MSS.0b013e3182514a88.
735   22.   Hoogkamer W, Kipp S, Kram R. The biomechanics of competitive male
736   runners in three marathon racing shoes: a randomized crossover study.
737   *Sports Med* 2019;**49**:133–43. https://doi.org/10.1007/s40279-018-1024-z.
738   23.   CSEP. 2017. Get Active Questionnaire.Canadian Society for Exercise
739   Physiology, Ottawa, Ont., Canada. Available from http://www.csep.ca/home.
740   24.   Day EM, Alcantara RS, McGeehan MA, Grabowski AM, Hahn ME. Low-pass
741   filter cutoff frequency affects sacral-mounted inertial measurement unit
742   estimations of peak vertical ground reaction force and contact time during
743   treadmill running. *J Biomech* 2021;**119**:110323.
744   25.   Wilkinson RD, Mazzo MR, Feeney DF. Rethinking the statistical analysis of
745   neuromechanical data. *Exerc Sport Sci Rev* 2023;**51**:43–50.
746   26.   Cohen J. Statistical power analysis. *Curr Dir Psychol Sci* 1992;**1**:98–101.
747   https://doi.org/10.1111/1467-8721.ep10768783.
748   27.   Bates D, Mächler M, Bolker B, Walker S. Fitting linear mixed-effects models
749   using lme4. *arXiv preprint arXiv:14065823* 2014.
750   28.   Lenth R, Singmann H, Love J, Buerkner P, Herve ME. Estimated Marginal
751   Means, aka Least-Squares Means R package version 1.8. 4-1. 2019.
752   29.   Rodrigo-Carranza V, González-Mohíno F, Santos-Concejero J, González-Ravé
753   JM. Influence of shoe mass on performance and running economy in trained
754   runners. *Front Physiol* 2020;**11**:1–8.
755   https://doi.org/10.3389/fphys.2020.573660.
756   30.   Chollet M, Michelet S, Horvais N, Pavailler S, Giandolini M. Individual
757   physiological responses to changes in shoe bending stiffness: a cluster
758   analysis study on 96 runners. *Eur J Appl Physiol* 2022;**123**:189–177.
759   https://doi.org/10.1007/s00421-022-05060-9.

31. Rodrigo Carranza V. Running footwear matters: decoding the influence of
running shoe characteristics on physiology, biomechanics and running
performance (PhD Academy Award). *Br J Sports Med* 2023;**57**:1581–2.
https://doi.org/10.1136/bjsports-2023-107452.
32. Knopp M, Muñiz-Pardos B, Wackerhage H, Schönfelder M, Guppy F, Pitsiladis
Y, et al. Variability in running economy of kenyan world-class and European
amateur male runners with advanced footwear running technology:
experimental and meta-analysis results. *Sports Med* 2023;**53**:1255–71.
https://doi.org/10.1007/s40279-023-01816-1.
33. Rodrigo-Carranza V, Hoogkamer W, Salinero JJ, Rodríguez-Barbero S,
González-Ravé JM, González-Mohíno F. Influence of running shoe longitudinal
bending stiffness on running economy and performance in trained and
national level runners. *Med Sci Sports Exerc* 2023;**55**:2290–8.
34. Rodrigo-Carranza V, Hoogkamer W, González-Ravé JM, González-Mohíno F.
Relationship between advanced footwear technology longitudinal bending
stiffness and energy cost of running. *Biorxiv* 2024.
https://doi.org/10.1101/2023.12.27.573423
35. Barrons Z, Rodrigo-Carranza V, Bertschy M, Hoogkamer W. The fallacy of
single trials: the need for multiple trials in assessing running economy
responses in Advanced Footwear Technology. *Sport Med* 2024.
https://doi.org/10.1007/s40279-023-01991-1.
36. Healey LA, Hoogkamer W. Longitudinal bending stiffness does not affect
running economy in Nike Vaporfly shoes. *J Sport Health Sci* 2022;**11**:285–92.
37. Wood DW. An experimental test of the Puma Model Number 296 Brush Spike
Shoe and the Mexico Puma Model Number 295 Standard Four Spike shoe as to
their effect on sprint running. [Ohio]: Ohio University, 1972.
38. Hébert-Losier K, Finlayson SJ, Driller MW, Dubois B, Esculier JF, Beaven CM.
Metabolic and performance responses of male runners wearing 3 types of
footwear: Nike Vaporfly 4%, Saucony Endorphin racing flats, and their own
shoes. *J Sport Health Sci* 2022;**11**:275-284.
https://doi.org/10.1016/j.jshs.2020.11.012.
39. Kram R. Ergogenic distance running shoes: how do we think they work and
how can we understand them better? *Footwear Sci* 2022;**14**:139–46.
40. Nigg BM, Cigoja S, Nigg SR. Teeter-totter effect: A new mechanism to
understand shoe-related improvements in long-distance running. *Br J Sports
Med* 2021;**55**:462-463
41. Rodrigo-Carranza V, Hoogkamer W, González-Ravé JM, Horta-Muñoz S, Serna-
Moreno M del C, Romero-Gutierrez A, et al. Influence of different midsole
foam in advanced footwear technology use on running economy and
biomechanics in trained runners. *Scand J Med Sci Sports* 2023;**34**:e14526.
https://doi.org/10.1111/sms.14526.
42. Hunter I, McLeod A, Valentine D, Low T, Ward J, Hager R. Running economy,
mechanics, and marathon racing shoes. *J Sports Sci* 2019;**37**:2367–73.
https://doi.org/10.1080/02640414.2019.1633837.

43.   Russo L, Montagnani E, Buttari D, Ardigò LP, Melenco I, Larion A, et al. Track
running shoes: A case report of the transition from classical spikes to "super
spikes" in track running. *App Sci* 2022;**12**:10195.
44.   Aristizabal Pla G, Martini DN, Potter M, Hoogkamer W. Assessing the validity
of the zero-velocity update method for sprinting speeds. *PLoS One*
2024;**19**:e0288896.
https://doi.org/https://doi.org/10.1371/journal.pone.0288896.
45.   Willwacher S, König M, Braunstein B, Goldmann JP, Brüggemann GP. The
gearing function of running shoe longitudinal bending stiffness. *Gait Posture*
2014;**40**:386–90. https://doi.org/10.1016/j.gaitpost.2014.05.005.
46.   Ortega JA, Healey LA, Swinnen W, Hoogkamer W. Energetics and
biomechanics of running footwear with increased longitudinal bending
stiffness: A narrative review. *Sports Med* 2021;**51**:873–94.
https://doi.org/10.1007/s40279-020-01406-5.
818
819

Figures

Fig. 1 Spike shoe models used in the experiments. From top to bottom, Experiment 1:

TRAD (PUMA EvoSpeed Distance 9), TRAD+ (PUMA EvoSpeed Distance 9 + 200g);

Experiment 2: TRAD, AFTP1 (PUMA AFT middle-distance prototype spike);

Experiment 3: TRAD, AFTP2 (PUMA AFT long-distance prototype spike), AFTP3

(PUMA AFT middle-distance prototype spike); Experiment 4: TRAD, AFTC1 (PUMA

evoSPEED Long Distance Nitro Elite+), AFTC2 (Nike ZoomX Dragonfly), AFTC3

(Adidas adizero Avanti TYO).

Fig. 2 Spike conditions significantly affect running speed and step length, but not step

frequency. Panel **A** shows average running speeds in different spike conditions, for

Experiment 1, 2, 3 and 4, from left to right. Panel **B** shows step frequency was similar

across spike conditions. Panel **C** shows significantly longer step lengths for AFTC1 and

AFTC2 than for TRAD. Grey lines represent individual data, black lines represent group

mean values, error bars represent standard error of the mean. * indicates significant

difference between spike conditions.

Fig. 3 Individual response of each participant in prototype AFT spikes vs. traditional

spikes (TRAD) in Experiment 3 for each of the three days and the 3-day average.

Tables

Table 1. Participant characteristics for each experiment (mean ± standard deviation).

Experiment	Participants	Age (years)	Mass (kg)	Height (m)	World Athletics Score[18]	Location
1	8 men	20.3 ± 2.7	70.9 ± 9.4	1.80 ± 0.06	787 ± 125	US
	4 women	20.0 ± 1.4	56.8 ± 2.4	1.67 ± 0.02	872 ± 110	
2	12 men	21.3 ± 1.8	62.8 ± 2.8	1.75 ± 0.04	772 ± 118	US, Canada
3	12 men	27.0 ± 5.7	64.8 ± 3.3	1.76 ± 0.06	817 ±126	Spain
4	12 men	22.0 ± 2.7	69.1 ± 8.3	1.81 ± 0.07	761 ± 120	Canada

850   Table 2: Characteristics of experimental conditions used in each Experiment.

Spike	Sizes	Mass (g)	Midsole thickness (mm)	Experiments used
TRAD	5.5, 7, 9, 11	131, 149, 158, 170	17	1,2,3,4
TRAD+	5.5, 7, 9, 11	331, 341, 355, 369	17	1
AFTP1	9	179	19	2
AFTP2	9	151	18	3
AFTP3	9	149	19	3
AFTC1	9	154	18	4
AFTC2	9	140	12	4
AFTC3	9	163	17	4

854 Table 3: Magnitude and significance of between spike differences were consistent across
855 days.

Comparison		Day 1	Day 2	Day 3	3-Day average
TRAD vs. AFTP2	% change	1.9	1.7	1.9	1.8
	effect size	0.36	0.32	0.31	0.34
	p value	0.003	0.004	0.0003	<0.0001
TRAD vs. AFTP3	% change	3.0	3.1	3.0	3.1
	effect size	0.57	0.56	0.51	0.57
	p value	<0.0001	<0.0001	<0.0001	<0.0001
AFTP2 vs. AFTP3	% change	1.1	1.4	1.1	1.2
	effect size	0.19	0.25	0.21	0.19
	p value	0.124	0.015	0.035	0.0003