Noise-induced changes in rat head stability and orientation correlate with dysfunction of otolith irregular afferents

Mamiko Niwa, David Bauer, Marie Anderson, Ariane Kanicki, Katharine Keller, Richard A. Altschuler, Courtney E. Stewart, and W. Michael King

Abstract

Exposure to intense noise damages both the cochlea and vestibular end organs. Our group previously reported attenuated vestibular short-latency evoked potentials (VsEP) and reduced numbers of calretinin-positive (CR+) calyces in the saccule following noise exposure. Here, we examined rats' resting head orientation with respect to gravity as well as head stability following a 4-hour exposure to 120 dB SPL noise. We also assessed how behavioral changes are related to changes in VsEP waveforms and calretinin expression in the otoliths to elucidate potential underlying mechanisms. We found significant delays in the VsEP N1 and P2 peak latencies and significant reductions in the N2P3 amplitudes following noise exposure. The number of CR+ calyces in both saccule and utricle were also significantly reduced. The size of the reduction in N2P3 amplitude was significantly correlated to the number of CR+ calyces in the utricle, but not to the saccule. Animals with larger delays in N1 peak latency and/or smaller numbers of CR+ calyces in the utricle following noise showed significant decreases in the average speed of y-axis rotational head motion, while those with smaller changes showed significant increases. In addition, animals with larger noise-induced changes held their heads motionless longer following noise exposure. We hypothesize that noise exposure is inherently destructive to an animal's head stability and thereby manifests as an increase in average head speed in mildly to moderately affected animals. But when the damage was large enough, animals exhibited reduced duration and head motion speed as a behavioral adaptation. The noise exposure also significantly altered the pitch angle of head orientation in animals who had the smallest number of CR+ calyces in the saccule, suggesting that saccular irregular afferents, including those that are CR+, are critical in control of head and body posture.

Introduction

Exposure to intense noise has been widely accepted as a hazard to our hearing, but less accepted is the effect of noise exposure on the well-being of vestibular function. Impairment of the vestibular system, whose canonical functions include stabilization of gaze, head motion, and posture, can lead to debilitating symptoms such as vertigo, blurred vision during head movement and postural instability. Furthermore, inputs originating from the vestibular periphery diffusely innervate various cortical areas and the limbic system; thus, vestibular dysfunction can have far-reaching impacts on various cognitive functions such as spatial navigation, memory, and attention (reviews in Shinder and Taube, 2010; Smith and Zheng, 2013). Because vestibular impairments are especially prevalent among the elderly, it is important to identify external factors and underlying mechanisms that lead to dysfunction of the vestibular periphery.

Clinical studies of industry and military workers with noise-induced hearing loss (NIHL) has long noted concomitant abnormalities in the vestibular periphery, implicating noise exposure as

damaging of both the auditory and vestibular organs in the inner ear(Collins, 1948). Abnormal nystagmus as well as reduced gain in the vestibulo-ocular reflex in the NIHL patients indicate dysfunction of the semi-circular canal organs, which encode angular acceleration of the head (Man et al., 1980; Oosterveld et al., 1982; Shupak et al., 1994). Hypofunction of the otolith organs, which encode linear acceleration of the head, were also indicated in the NIHL patients, as evidenced by reduced vestibular-evoked myogenic potentials of cervical and extraocular muscles (cVEMP and oVEMP; Akin et al., 2012; Tseng and Young, 2013; Viola et al., 2020; Wang and Young, 2007).

In animal models, noise exposure has been shown to induce molecular and morphological changes in the vestibular periphery including elevated oxidative stress markers (Tamura et al., 2012), reduced calcium-binding protein in afferent terminals (Stewart et al., 2018, 2020), hair bundle disruption(Stewart et al., 2016), vacuole formation and loss of nucleus in type-I hair cells (Hsu et al., 2008), degeneration of epithelial cells (Akdogan et al., 2009), and structural collapse and/or membrane rupture (Mangabeira-Albernaz et al., 1959; Mccabe and Lawrence, 1958) depending on the intensity, duration, and frequency profiles of the noise exposures (review by Stewart et al., 2020). Electrophysiological changes were also noted as reduced amplitude of the cVEMP (Hsu et al., 2008; Kumar et al., 2010) and reduced amplitude and prolonged latency in the vestibular short-latency evoked potential (VsEP; Perez et al., 2002; Stewart et al., 2018, 2020).

Behavioral or clinical consequences of noise-induced damage to the vestibular periphery are less well understood. Notably, among the NIHL patients with abnormal findings in the vestibular periphery, only a subset complained of symptoms such as vertigo and dizziness. However, the absence of clinical symptoms does not mean there is no behavioral consequence; studies have shown that the NIHL patients without clinically relevant vestibular symptoms show larger body sways compared to control subjects (Juntunen et al., 1987; Ylikoski et al., 1988). Animal studies also show that noise exposure reduced rodents' proficiency in motor tasks that require balance (rota-rod: Tamura et al., 2012; balance beam crossing: Bartikofsky et al., 2023). Potential mechanisms underlying these behavioral outcomes include reduced gain and/or increased variability in encoding the gravitational vector, reduced accuracy in an internal model of active vs passive head movements, and reduced speed/gain in the vestibulo-collic and -spinal reflexes, degrading stability of head and body posture.

A goal of our study was to provide a quantitative analysis of behavioral changes that follow noise exposure. Here, we examined animals' head orientation and stability during self-generated head movements before and after exposure to 120 dB noise. Our group recently reported that rats exposed to a similar noise profile exhibited significantly decreased proficiency in balance-beam crossing (Bartikofsky et al., 2023). We also showed that this noise exposure attenuated VsEP waveforms and reduced the number of countable calretinin-positive calyces in the saccule (Stewart et al., 2018, 2020). The present study identifies a relationship between electrophysiological and morphological changes in the periphery to changes in head stability and orientation and suggests potential underlying mechanisms of noise-induced dysfunction.

Results

The vestibular short-latency evoked potential (VsEP) was measured before and after exposing rats to 3-octave band (0.53 to 4.2 kHz) noise at 120 dB SPL for 4 hours. The VsEP is

thought to be a far field potential, generated by synchronous discharges of the action potentials of irregular vestibular afferents and vestibular neurons in the brain stem, and evoked by a linear acceleration jerk applied to the animal's head (Nazareth and Jones, 1998). The rate of change in linear acceleration, jerk, has been shown as a parameter on which the VsEP primarily depends (Jones et al., 2011).

A representative animal's VsEP responses to 5 jerk levels are shown in Figure 1A prior to noise exposure (black traces), 1 week (magenta traces) and 4 weeks after noise exposure (green traces). The VsEP has been shown to typically start with a positive peak, followed by multiple sets of negative and positive peaks (Jones and Jones, 1999; Plotnik et al., 1997). In the present study, the first positive peak, P1 (denoted with a grey arrow in Figure 1A) was not as prominent as the later peaks and in some animals undetectable. A less prominent P1 peak has also been seen in other studies that measured the VsEPs in rodents (Jones and Jones, 1999). The N1 (denoted with red star), P2 (red arrow), N2 (blue star), and P3 (blue arrow) peaks were prominent in all animals tested. In this animal, N1P2 amplitude, defined as the potential difference from N1 to P2 peak, was reduced at 1 and 4 weeks after noise compared to the baseline at all stimulus levels (Figure 1B). A similar reduction was seen for the N2P3 amplitude (Figure 1C). The N1 and P2 peak latencies were delayed both at 1 and 4 weeks after noise for most stimulus levels (Figure 1D and 1E).

Across the population, the N2P3 amplitude was significantly reduced at both 1 and 4 weeks after noise exposure compared to the baseline at all but the smallest stimulus level (p = 0.0040 or lower with t-test, Figure 2B). The population mean reduction across stimulus levels was 27.7% (SEM = 4.1) for week 1 and 27.8 % (SEM = 3.7) for week 4, showing no sign of recovery within this period. The N1P2 amplitude at week 4 was significantly different from the baseline when the change was collapsed across all stimulus levels (p = 0.019, Figure 2A). At week 1, there was no significant reduction at any individual stimulus level or at all-levels-collapsed (p = 0.080 or higher, Figure 2A) because a small subset of animals showed a large enhancement in the N1P2 amplitude following noise, although their N2P3 amplitudes were greatly reduced (Figure 2D). There was a significant correlation between % change in the N2P3 amplitude and the absolute value of the % change in the N1P2 amplitude at 1 week after exposure at all but the smallest stimulus level (stim level 5: r = -0.68, p = 0.0038; level 4: r = -0.56, p = 0.024; level 3: r = -0.72, p = 0.0016; level 2: r = -0.60, p = 0.014 with Spearman's correlation test, Figure 2E).

The N1 peak latency was significantly delayed at an intermediate stim level and at all-levels-collapsed (week 1: p = 0.007 at stim level 3, p = 0.0035 at all-levels-collapsed; week 4: p = 0.0047 at stim level 4, p = 0.0001 at all-levels-collapsed; Figure 2C). Percent change in the N1 peak latency was significantly correlated with % change in the N2P3 amplitude at 1 week after the exposure (stim level 5: r = -0.67, p = 0.0045; stim level 2: r = -0.582, p = 0.018, Figure 2F). We also saw a significant noise-induced delay in the P2 peak latency and a similar degree of correlation to the N2P3 amplitude (data not shown).

The reduced number of the calretinin-positive calyces in the utricle correlates with reductions in VsEP amplitude

Next, we examined the expression of a calcium-binding protein, calretinin, in the calyces of both the saccule and utricle at 5 weeks after noise exposure. Calretinin is expressed in the calyceal terminations of afferents contacting only type-1 hair cells (calyx-only afferents), but not

in the terminations of afferents contacting only on type-2 hair cells (bouton afferents) or those contacting both type-1 and type-2 hair cells (dimorphic afferents; Desai et al., 2005; Desmadryl and Dechesne, 1992; Leonard and Kevetter, 2002). Afferents with calretinin-positive (CR+) calyces in the otoliths are known to have irregular temporal firing patterns and large gain and phase leads to head motion (Goldberg et al., 1990) and to be sensitive to high-frequency changes in linear acceleration of the head, such as the jerk stimuli that cause the VsEP.

We found that the number of CR+ calyces in the saccule and utricle of noise-exposed animals was significantly reduced compared to control animals (Figure 3A and 3B). The population-average reduction was similar in both organs (~20%). While some animals showed similar numbers of CR+ calyces in the utricle and saccule, other animals showed a larger number in one organ and fewer in the other. Thus, we found no significant correlation between the number of CR+ calyces in the saccule and utricle in individual animals (r = 0.26, p = 0.30 with Pearson's correlation test, Figure 3C).

The reduction by ~20 % in the number of CR+ calyces in the utricle and saccule matched reasonably well with the population-average 28% reduction in the VsEP's N2P3 amplitude across stimulus levels (Figure 2B). Furthermore, at the level of individual animals, the degree of the N2P3 amplitude reduction was tightly linked to the number of CR+ calyces in the utricle (stim level 5: r = 0.56, p = 0.025 at week 1 and r = 0.78, p = 0.0027 at week 4; Figure 3D), but not to the saccule. The correlation was robust and consistent across all but the smallest stimulus levels. Thus, the reduction in N2P3 amplitude following noise exposure appears to reflect changes in the utricle more strongly than the saccule. The stronger dependence on the utricle may be because the VsEP stimulation in the animal's nasal-occipital axis aligns better with the activation direction of hair cells' stereocilia in the utricle than the saccule.

We also found that the change in the N1 peak latency is correlated with the number of CR+ calyces averaged between the saccule and utricle (Figure 3E). However, the correlation is statistically significant only when the number of CR+ calyces are averaged between the saccule and utricles, and only at week 4 (r = -0.62, p = 0.032 with Spearman's correlation test, bottom right panel in Figure 3E). Thus, the change in the number of CR+ calyces appears to be more tightly linked to the amplitude aspect of the VsEP, and less so to its temporal aspect.

Animal's head stability and orientation following noise exposure.

Next, we present how these electrophysiological and morphological changes affected head stability and orientation following noise exposure. Using a miniature motion sensor attached to an animal's head, we examined their head orientation and stability during inter-trial-intervals (ITIs) between abrupt, horizontal rotations given in our behavioral experiment (see Methods for behavioral test). During the ITIs, no external motion was applied to the rat, and it could freely move its head. The present study includes behavioral changes following exposure to sodium arsanilate, a vestibulo-toxic compound, as a positive control of bilateral vestibular loss (see Methods for trans-tympanic injection of sodium arsanilate).

Animals with a greater noise-induced damage keep their head motionless longer following noise exposure

To assess head stability, we first quantified how long an animal's head was stationary during ITIs. Figure 4A shows 'stable period duration', measuring how long each stationary interval lasted, over the course of the behavioral experiment in a representative noise-exposed

animal. The daily median of stable period duration significantly increased after noise exposure (p = 3.4 x 10^4 with rank-sum test, Figure 4B). Figure 4D and 4E show that within each ITI, the total amount of time an animal's head remained stationary, 'the proportion of stable time', also increased after noise exposure in this animal (p = 2.4 * 10^4, Figure 4E). We found that a subset of noise-exposed animals significantly increased stable period duration (5 out of 17 animals, Figure 4C) and proportion of stable time (2 out of 17 animals, Figure 4F), suggesting that noise exposure resulted in an apparent promotion of head stability in these animals.

The change in stable period duration was significantly correlated with % change in the VsEP N2P3 amplitude at week 4 (stim level 4: r = -0.72, p = 0.0080; stim level 3: r = -0.71, p = 0.010; stim level 2: r = -0.59, p = 0.045; Figure 4G). The change in stable period duration was also correlated with the number of CR+ calyces in the utricle with marginal significance (r = -0.45, p = 0.065; Figure 4G). Similarly, the change in the proportion of stable time after noise is correlated with % change in N2P3 amplitude with marginal significance (stim level 5: r = -0.54, p = 0.068; stim level 4: r = -0.57, p = 0.051; stim level 3: r = -0.51, p = 0.091; Figure 4H), and significantly correlated with the average number of CR+ calyces in the utricle and saccule (r = -0.51, p = 0.036; Figure 4H). Thus, the result indicates that animals that had larger noise-induced damage, as assessed by VsEP amplitude reduction and reduced CR+ calyces in the otoliths, also showed larger increases in head stability/motionlessness after noise exposure.

In contrast to noise exposure, all 3 animals exposed to arsanilate showed decreased head stability/motionlessness after lesion. There were significant decreases in the proportion of stable time for all 3 animals (Figure 4F). The stable period durations were also shorter, although the change was not statistically significant for any animal (Figure 4C). The decreased head stability in arsanilate-exposed animals was an expected result for animals with peripheral vestibular lesions.

The speed of the animal's rotational head motion increased or decreased depending on the degree of noise-induced dysfunction

Next, we examined the speed of the rats' voluntary head motion. We computed the root-mean-square (RMS) of the angular velocity measurements from the x- (roll), y- (pitch), and z- (yaw) axis gyros of the head sensor while the animal's head was in motion for any duration longer than 0.5 second. Because the average speed of motion tended to be higher for longer periods of motion, and because noise and arsanilate lesions affect active period duration (and stable period duration, see Figure 4), we transformed the RMS values into z-scores using the pre-exposure mean and standard deviation of RMS speeds calculated within short duration bins. In this way, we minimized the confounding effects of active period duration on RMS angular velocity.

Figure 5A shows the z-score-transformed RMS pitch-axis angular velocity over experimental days in a representative animal exposed to noise. Angular velocity of animals' head motion was inherently variable; we observed a large variability between segments of motion within a day (error bars) and large day-to-day variability in the daily median RMS (blue squares for pre- and red squares for post-exposure). Nonetheless, most of the daily median RMS values after exposure were below the pre-exposure average (black dashed line), so there was a significant reduction in the daily median RMS pitch-axis angular velocity following noise exposure (p = 0.0055 with Wilcoxon's rank-sum test, Figure 5B).

Out of 17 animals exposed to noise, we found 4 animals with a significant decrease in the daily median RMS pitch-axis angular velocity, and 5 animals with a significant increase (Figure 5D). For the roll- and yaw-axis motion, we found smaller numbers of animals with significant increases or decreases (Figure 5C and 5E). When the RMS values from all 3 axes were combined by vector averaging, 4 animals showed a significant increase in overall head speed after noise, while 2 animals showed a significant decrease (Figure 5F). Because of these opposed changes following noise exposure, the population-average was diminished to ~0 for all directions. This contrasts with the changes following arsanilate exposure, where all 3 animals showed a significant increase in head angular velocity in all 3 directions (Figure 5C~E).

We found that animals with significant decreases in the pitch-axis head velocity had larger delays in the N1 peak latency at 1 week after noise exposure (Figure 5H; stim level 5: r = -0.59, p = 0.026 with Pearson's correlation test: rho = -0.68, p =0.0073 with Spearman's correlation test; stim level 3: r = -0.68, p = 0.0074: rho = -0.58, p = 0.029). The relationship appears highly non-linear, as if there were a threshold of % change in the N1 peak latency (~5%) that divides individual animals into two distinct groups: increased vs decreased RMS velocity.

We also found a significant correlation between the change in the RMS pitch-axis angular velocity and the number of CR+ calyces in the utricle (r = 0.49, p = 0.048), but not in the saccule (r = 0.20, p = 0.45, Figure 5G). Although statistically significant, the number of CR+ calyces does not clearly separate individual animals into two distinct groups as seen with the change in N1 peak latency. Among the animals showing a small number of CR+ calyces in the utricle (< 36), about a half of them increased pitch-axis motion speed, while the other half showed a decrease. The number of CR+ calyces in the utricle is tightly linked to the VsEP amplitude change following noise exposure, and less so to the latency change (Figure 3D and 3E). Together with a weak correlation between the change in the pitch-axis angular velocity and % change in the VsEP N2P3 amplitude (data not shown), the result suggests that the change in the temporal aspect of the VsEPs is a stronger predictor of how the speed of the pitch-axis head motion changes following noise exposure in individual animals, compared to the amplitude aspect.

Examination of the frequency profile of the animal's head motion before vs after noise exposure hints at a possible stiffening of the animal's neck musculature as an adaptive strategy to decrease the speed of the y-axis rotational motion. The power spectrum of the pitch-axis angular velocity in Figure 5I shows a shallower slope in the low frequency region (< 5Hz) and a steeper slope in the mid frequency region (10~30Hz). The shape of the power spectrum generally agrees with a previous study, which has reported the power spectrum of rotational head motion deviates from a power law likely due to the biomechanical filtering (Carriot et al., 2014, 2017; Zobeiri et al., 2021). Indeed, when angular velocity is converted into angular acceleration by differentiation, the power spectrum for angular acceleration of the pitch-axis head motion shows a unimodal distribution with a peak around 8 Hz for all animals tested (see right panels in Figure 5I~K). In animals who showed head bobbing, the frequency of bobbing coincides at ~8 Hz (Figure 5J). These data suggest that ~8 Hz represents a resonance frequency for rotational motion of a rat's head.

In noise-exposed animals with a significant decrease in RMS pitch-axis angular velocity, the power at frequencies below 10 Hz decreased for all animals (Figure 5N). Intriguingly, the power at frequencies above 10 Hz increased in all but one animal. One interpretation of this

result is that these animals have stiffened neck musculature, effectively reducing the overall speed of head motion, yet slightly raising the power at higher frequencies (> 10 Hz) following noise exposure.

Note that animals with smaller noise effects in terms of the change in N1 peak latency and the number of CR+ calyces in the utricle still showed an increase in the speed of y-axis head motion. The arsanilate-exposed animals also showed an increase, although the effect size is much larger. Our interpretation of this result is that noise exposure degrades head stability. Together with the results shown in Figure 3, we hypothesize that animals with larger noise-induced damage employ adaptive strategies to reduce both the time duration and the speed of head motion to mitigate head instability brought on by noise exposure.

Noise-exposure alters animals' resting head orientation

Lastly, we present how the animal's head orientation, roll and pitch (Figure 6A and 6H, respectively), changed following noise or arsanilate exposure. Figure 6B shows a scatter plot of a noise-exposed animal's roll orientation against stable period duration (blue dots for pre- and red dots for post-exposure). Figure 5C shows the mean and standard deviation of roll orientation for pre- (blue line) and post-exposure (red line) as a function of stable period duration. The center of the distribution has shifted by ~5 degree (tilted toward the animal's left side) and the distribution broadened after noise, regardless of the stable period duration. In the rest of analyses, we only included head orientation from stable periods > 1 sec to examine the 'resting' head orientation.

The histograms of z-score-transformed roll orientation collapsed across durations >1 sec (Figure 6D) reiterate the shift and broadening in roll distribution after noise exposure. There was a marginally significant increase in the daily median roll orientation (p = 0.08, Figure 6F) and a significant increase in variability within a day (p = 0.0031 with rank-sum test, Figure 6G; see larger red error bars vs. blue in Figure 6E) after noise exposure.

The animal's pitch orientation was also altered after noise. The plots of pitch orientation as a function of duration on Figure 6I and 6J clearly show a shift and tightening of the distribution regardless of duration after noise exposure in a representative animal. The variability in pitch orientation within each day was significantly reduced after exposure (p = 0.028 with rank-sum test, Figure 6N). The day-to-day variability in median pitch orientation was also reduced, though not significantly (p = 0.105 with variance test, Figure 6M). Both these changes contributed to tightening the overall pitch distribution after noise-exposure in Figure 6K (red histogram). The daily median pitch orientation was also significantly decreased (head lowered toward chest; p = 0.0086 with rank-sum test, Figure 6M).

Of the 17 animals exposed to noise, 4 animals showed a significant change in median roll orientation, with 3 animals tilted to their right side and 1 to the left (Figure 7A). For pitch, 5 animals showed a significant change in median orientation, with 4 animals going downward and 1 upward, following noise exposure (Figure 7D). As a population, the shift in pitch orientation is biased downward, with a statistically marginal significance (p = 0.093 with signed-rank test), possibly due to gravity. There was a significant correlation between the sizes of the change in median roll and pitch (Figure 7G); an animal with a larger change in median roll orientation tends also to have a larger change in median pitch orientation.

Strikingly, the size of change in the median pitch orientation was significantly correlated to the number of CR+ calyces in the saccule (r = -0.67, p = 0.0033 with Pearson's correlation test; left panel in Figure 6H), but not to the utricle (r = -0.18, p = 0.49; right panel in Figure 6H). We did not find a significant correlation to VsEP changes, likely because the VsEP appears more strongly tied to utricular activity.

Discussion

Noise exposure and head stability

Animals with larger delays in N1 peak latency of the VsEP and/or smaller numbers of CR+ calyces in the utricle following noise exposure showed significant decreases in the speed of y-axis head motion, while those with smaller changes showed significant increases. In addition, animals with larger noise-induced changes held their heads motionless for longer time periods following noise exposure. Together, we interpret this result as (1) noise exposure was inherently challenging to an animal's head stability, and thereby manifests as an increase in average head speed in mildly to moderately affected animals, but (2) when the damage was large enough, animals exhibited reduced duration and speed of head motion as a behavioral adaptation to modified vestibular afference.

In patients with vestibular disturbances, a reduction in natural head movements is a common behavioral adaptation because the movement aggravates their vestibular symptoms (Colledge et al., 1994; Heidenreich et al., 2010; Paul et al., 2017). This supports our finding of prolonged head motionlessness in animals with a larger noise-induced damage. Also, an adaptive strategy that includes co-contractions of antagonistic neck muscles to stabilize head upon body perturbations has been reported in elderly subjects (Keshner, 2000). Our result is consistent with the human findings since (1) the animals in our study were also subjected to rapid, horizontal body perturbations that required them to stabilize their unsupported heads and (2) the power of rotational head motion over high frequencies (> 10 Hz) increased following noise exposure in animals which had an overall reduction in RMS speed of motion. Thus, we hypothesize that animals with larger noise-induced damage in the vestibular periphery likely stiffen their neck musculature as an adaptive strategy.

Alternatively, increased power in high-frequency head motion could be interpreted as the functional consequence of the dysfunction of irregular afferents, which are better at encoding natural head movement at higher frequencies than regular afferents (Jamali et al., 2019). In a study of patients with unilateral vestibular loss, significant increases in high-frequency head movements were seen when patients performed challenging tasks, such as walking on foam, that likely required online updating of an internal model of self-motion (Zobeiri et al., 2021). Because there was no change in the power of low-frequency head movements, for which vision and proprioception could aid stabilization, the increased power of high-frequency head motions in the patients was interpreted to be due to impairment of the vestibular system, which contributes to damping high-frequency head motion.

Noise exposure alters animal's resting head orientation

Noise exposure changed the animal's median head orientation in both roll and pitch. One interpretation is that the change is specific to the behavioral context of the present study,

because head orientation was measured during the inter-trial-intervals of imposed horizontal rotations. As mentioned above, an adaptive strategy of co-contraction of antagonistic neck muscles during body perturbations could contribute to a change in head orientation following noise exposure. Resting head orientation during natural movements will need to be examined to confirm this interpretation.

An alternative interpretation is that noise exposure degrades the directional estimate of gravity. Otolith regular afferents likely encode the best estimate of static head position (Jamali et al., 2019); however, afference from the semi-circular canals must also be combined to distinguish head tilt re gravity from linear translational motion (Angelaki et al., 1999). Therefore, the change in resting head orientation following noise exposure suggests that otolith regular afferents and/or canal afference is compromised. Indeed, results from previous studies are consistent with this conjecture (Fetoni et al., 2009); for example, exposure to a broadband white noise at 116 dB SPL for 3 hours significantly reduced the number of the hair bundles on the hair cells in the striolar and extra-striolar regions of the saccules and utricles as well as the anterior and horizontal semicircular canal cristae (Stewart et al., 2016).

We found a significant correlation between the size of the post-noise change in pitch and the number of CR+ calyces in the saccule but not the utricle. This points to a strong contribution of saccular irregular afferents to the animal's resting head orientation regardless of whether it is specific to the behavioral context of the present study. Because the saccules are anatomically more aligned with the earth vertical axis, a greater correlation between saccular damage and the change in an animals' pitch position is reasonable. In addition, vestibular nucleus neurons that project to the upper cervical cord predominantly receive irregular afferent inputs, whereas those neurons that project to the extraocular muscles (VO and VOC neurons; Cullen et al., 2001) receive higher proportions of regular afferent inputs (Boyle et al., 1992; Highstein et al., 1987). The difference in the ratio of regular vs irregular inputs to neurons involved in vestibuloocular vs vestibulospinal pathways may reflect the higher inertia of the head as compared to the eyeball, since the large phase-lead exhibited by irregular afferents' responses to head motion may be necessary to control head position. Thus, irregular afferents, including those that are CR+, are critical to control static and dynamic posture of the head and body, and a change in resting head position may reflect their loss.

Noise exposure delays the early components of the VsEP

We found that the latency of the VsEP N1 peak was significantly delayed following noise exposure. In the chicken, the P1 and N1 peaks were suggested to reflect compound action potentials of vestibular irregular afferents based on their resistance to brain stem lesions and a small dependence on body temperature (Nazareth and Jones, 1998). Thus, a delay in the N1 peak latency may represent either an increased delay in irregular afferents' spike discharges (e.g., first spike latency) or less of synchrony in irregular afferents' discharges.

A very recent study has shown strong evidence that vestibular irregular afferents' short-latency response to bone conducted vibration is generated through non-quantal transmission at the synapse between vestibular type-1 hair cells and calyx-bearing afferents (Pastras et al., 2023). The basolateral membrane of type-1 hair cells is enclosed by a calyceal membrane at close apposition, and there is a large resting membrane conductance due to low-voltage activated K+ channels and HNC channels on both sides of the synapse (Contini et al., 2020, 2017). The electrical coupling due to this arrangement enables hair-cell depolarization by

mechano-transduction currents to trigger action potentials in calyx-bearing afferent fibers with extremely short latencies (nearly instantaneously) and without activation of glutamate receptors (Contini et al., 2020, 2017; Govindaraju et al., 2023).

Assuming the earliest components of the VsEP represent irregular afferents' compound action potentials driven by non-quantal transmission, then delayed latency in the N1 peak in the present study may be interpreted as reduced speed (greater latency) or less synchrony in non-quantal spike activation following noise exposure. A computational model by Govindaraju et al., (2023) has shown that the height of the calyceal enclosure on the hair cell affects the speed of non-quantal transmission. Retraction of calretinin-positive calyces has been noted following exposure to vestibulo-toxic gentamicin (Sultemeier and Hoffman, 2017), which can lead to greater AP discharge latencies according to the model. Whether or not noise exposure also causes calyceal retraction is an interesting hypothesis that must be tested in future experiments.

Noise exposure affects the expression of calcium-binding protein in the calyx

In our study, the saccule and utricle appeared similarly susceptible to noise exposure, at least regarding the changes manifested in the number of CR+ calyces (~20 % reduction). Otolith irregular afferents, consisting of calyx-only and striolar dimorphic afferents, can respond to air conducted sound with intensity thresholds 80~100 dB SPL for the saccule and 100~120 dB SPL for the utricle over the frequency range of noise used in the present study (Curthoys et al., 2016). Their firing rate steeply increases with intensity above threshold, suggesting that a noise intensity of 120 dB SPL as used in the present study caused excessive firing in otolith irregular afferents, leading to noise-induced damage.

We found a striking correlation between the number of CR+ calyces in the utricle and the reduction in the VsEP N2P3 amplitude following noise exposure. This points to a dominant role for CR+ (calyx-only) afferents in generation of the VsEP, and the utricle as more likely the major contributor than the saccule when stimulus jerks in the naso-occipital axis are used. At the same time, it raises a question of the degree of contribution by irregular dimorphic afferents in generation of the VsEP as well as their susceptibility to noise exposure. Calyx-only afferents in the utricle, on average, have higher irregularity in their firing pattern and larger phase leads to head motion stimulation compared to irregular dimorphic afferents (Goldberg et al., 1990). This may make calyx-only afferents more sensitive to high-frequency changes in linear acceleration during jerks and be a dominant source of the VsEP response. A future examination of the expression of calbindin in calretinin-negative calyces would address the susceptibility of irregular dimorphic afferents to noise.

Our previous report showed that reduction in CR+ calyces occurred without a significant change in the total number of calyces in the saccule following noise exposure (C. E. Stewart et al., 2020). This suggests that the reduction in the number of CR+ calyces is due to down-regulation of calretinin in a subset of calyx-only afferents, and not due to the loss of these afferent terminations.

Noise-induced down-regulation of calretinin would alter the dynamics of the intracellular free Ca^{2+} in calyx-only afferents, which in turn would influence the activity of calcium-dependent K+ channels, HCN channel via cAMP, KCNQ K+ channels via calmodulin, and efferent modulation via G-protein coupled mechanism (Holt et al., 2017, 2015; Lee et al., 2017; Meredith et al., 2011; Seseña et al., 2014; Ventura and Kalluri, 2019). The calix-only afferents contain calbindin, calcium pumps, and many mitochondria in addition to calretinin (Lysakowski et al.,

2022; Lysakowski and Goldberg, 1997), indicating that their intracellular free Ca^{2+} must be intricately regulated through mechanisms having different spatiotemporal dynamics. We found a robust correlation between the number of CR+ calyces and a change in head stability and orientation. This points to the relevance of calcium regulation for calyx-only afferent function and ultimately to vestibular behavior. Future studies involving intracellular recording of calyx-bearing afferents would be instrumental in revealing the role of calcium regulation on the functional properties of these afferents in the context of noise-induced damage.

Methods

Animals: We used adult Long-Evans rats of both sexes (Charles River Laboratories, MA) with weight ranging between 600 ~ 900 g in males and between 250 to 380 g in females. They were caged in pairs or trios with free access to food and water. The housing facility had a 12/12-hour light/dark cycle, and all experiments were done between 8am-5pm during the light cycle. Twenty rats were the subjects in the behavioral tests, VsEP recordings, and the immunohistological study. Seventeen rats (8 males and 9 females) received noise exposure while three male rats received sodium arsanilate. Thirteen additional rats served as control subjects in the immunohistological study. Two other rats received noise exposure and served in VsEP recordings and immunohistological study. All procedures were carried out in accordance with National Institutes of Health guidelines and were approved by the Institutional Animal Care and Use Committee at the University of Michigan.

Noise exposure: Rats were anesthetized with ketamine (University of Michigan B2 Pharmacy, MI and Covetrus, OH) and xylazine (Covetrus, OH and MWI Animal Health, ID). They received 3-octave band noise (530 ~ 4240 Hz) bilaterally for 4 hours and were given Atipamezole (Covetrus, OH) as reversal after the completion of the exposure. The waveform of the noise stimulus was created using Adobe Audition version 1.5 software (Adobe, CA). The sound was amplified with an amplifier (DA134, Rolls, UT) and delivered through inserted earphones (ER 3C, Etymotic Research Inc., IL). The sound level was calibrated with a fast Fourier transform spectrum analyzer (SR760; Stanford Research Systems, CA) by setting the peak of the frequency spectrum to 120 dB SPL. The SPL of the spectrum analyzer was calibrated with the sound level calibrator (R8090, Reed Instruments, NC). Bruel & Kjaer type 4136 microphone (Bruel & Kjaer Sound and Vibration Measurement, Denmark) was used in calibration.

Arsanilate exposure: The stock solution was prepared by dissolving sodium arsanilate in 0.9% sterile saline at 150 mg/mL concentration. An animal was given oral administration of Meloxicam and then anesthetized with isoflurane (MWI Animal Health, ID). Under surgical loupe, 20 µl of the stock solution was injected through the tympanic membrane with a Hamilton syringe. The animal was held on its side for 5 min to allow absorption before repeating the injection in the other ear.   Because arsanilate is known as a vestibulo-toxic chemical that destroys the epithelium of the vestibular end organs and degrades vestibular behavior ((Hunt et al., 1987; Ossenkopp et al., 1990; Vignaux et al., 2012), we treated arsanilate-exposed animals as a positive control of bilateral vestibular loss.

Surgical implantation: The rat was anesthetized with isoflurane (MWI Animal Health, ID) and its head was held in a stereotaxic frame. A midline incision was made to expose the bregma

and lambda on the dorsal surface of the skull. The stereotaxic positioning was further adjusted so that the Z-axis coordinates of the bregma and lambda were equal. Three anchor screws were placed into the skull. A custom, small bolt was placed vertically against the bregma and cemented with C&B Metabond (Perkell, NY). The attachment of the head bolt to the skull was strengthened with dental acrylic fusing the bolt and the skull via the anchor screws. The incision was sutured around the implantation. The animal was placed on post-operative recovery for 10~14 days.

Vestibular short-latency Evoked Potential (VsEP): VsEP was recorded as described previously (Stewart et al., 2020). Briefly, a rat was anesthetized with ketamine and xylazine and its head was affixed to a shaker (ET-132, Labworks Inc., CA) via the surgically implanted bolt so that the direction of the stimulus aligned with the animal's naso-occipital axis. Jerk stimuli with 5 different amplitudes (~ 0.3 to 6.4 g/ms) were produced by applying linear voltage ramps of different slopes and durations to the shaker. Jerk is the rate of change in linear acceleration. The stimulus duration ranged between 0.65 ~1.35 ms. Electrodes (stainless needles) were placed subcutaneously at the vertex (noninverting), mastoid (reference), and hip (ground). Stimuli were delivered as successive positive and negative jerks (~200 each) and averaged together to minimize electromagnetic artifacts from the shaker.

Immunohistochemistry: The number of calretinin-positive calyces was determined as described previously (Stewart et al., 2020). Briefly, animals were given a lethal dose of sodium pentobarbital and perfused with 4% paraformaldehyde in PBS. Otic capsules were removed, immersed in the fixative for 2~4 hours, and decalcified in 5% EDTA in PBS for 3~4 days at room temperature. The utricle and saccule were dissected, then blocked and permeabilized in 0.3% Triton X-100, 5% normal donkey serum in PBS with Ca^{2+} and Mg^{2+} for 2 hours at room temperature. The primary antibodies were mouse anti-calretinin (CR; 1:1000, Millipore MAB1568), chicken anti-neurofilament H (NF; 1:1,000, Millipore AB5539) or rabbit anti-tubulin beta 3 (Tubb3, 1:1000, BioLegend Poly18020), and rabbit anti-myosin 7a (Myo7a; 1:100, Proteus Bioscience, PAB 25–6790) or Phalloidin Alexa Fluor 647(1:100, Millipore Sigma A22287). A 40X z-series with 1-μm steps were collected with a confocal microscope (Stellaris, Leica, Germany) at a region of interest (Stewart et al., 2018, 2020). Images were analyzed for the presence of CR+ calyces with MetaMorph software. CR+ calyces are defined as CR-positive, Myo7a- (or Phalloidin-) negative, and NF- (or Tubb3) positive circles of diameter greater than 5 μm that appeared below the cuticular plate of the sensory epithelium.

Motion sensor: A miniature motion sensor (3-Space™ LX Embedded, Yost Labs Inc., Ohio) was placed on the animal's head via the surgically implanted bolt. The motion sensor consisted of 3 accelerometers, 3 gyroscopes, and 3 magnetometers along 3 orthogonal axes. The sensor was oriented so that the x-axis is aligned with the midline of the animal, the y-axis with the interaural line, and the z-axis with the animal's head bolt. The motion sensor was connected to a personal computer through a USB connection. A custom C++ program along with the sensor manufacturer's API protocols was run on Visual Studio (Microsoft, WA) to record accelerometer and gyroscope measurements on the PC. The sampling rate was 500 or 1000 Hz. The dynamic range was at +/- 2 G for accelerometers and +/- 500 or 1000 ° /s for gyroscopes.

Behavioral test: The rat and motion sensor were placed in a holding device that restrained the body with harnesses. The rat supported its own weight on 4 limbs and was able to move its head freely. The rat and restraint device were then placed on a horizontal, servo-controlled table in a dark booth. The table rotated about an earth vertical axis. In each trial of the behavioral test,

the rotator generated an abrupt velocity step of 25, 50, 75, or 100 °/s amplitude in clockwise or counterclockwise direction. After the velocity-step rotation, the table paused for ~0.2 sec and rotated back to the starting position in a reverse direction at 10 °/sec. The table then paused for a random period (3 ~ 5.5 sec) before the next trial. The data in the present study was taken during these inter-trial-intervals (ITIs).

Determination of the period of head stability and head orientation: The animal's head was characterized as motionless when the following term, A(t), was less than an empirically determined threshold;

A(t) = $\frac{d}{dt}\left(g_x^2 + g_y^2 + g_z^2\right)$, where g_x, g_y, and g_z are angular velocity in x-, y-, and z-, axis, respectively.

Euler angles of the animal's head position were calculated for time periods where the animal's head was motionless, using the following;

$\text{roll} = \text{atan}\left(\frac{a_y}{a_z}\right)$, $\quad \text{pitch} = \text{atan}\left(\frac{-a_x}{(a_z*\cos(roll)+ a_y*\sin(roll))}\right)$, where a_x, a_y, and a_z are linear acceleration in x-, y-, and z-, axis, respectively.

This pair of Euler angles are based on intrinsic rotations in the 'yaw-pitch-roll' order (Kuipers, 2002), which is equivalent to extrinsic rotations in the 'roll-pitch-yaw' order.

References

Akdogan O, Selcuk A, Take G, Erdoğan D, Dere H. 2009. Continuous or intermittent noise exposure, does it cause vestibular damage?. An experimental study. *Auris Nasus Larynx* **36**:2–6. doi:10.1016/J.ANL.2008.03.003

Akin FW, Murnane OD, Tampas JW, Clinard C, Byrd S, Kelly JK. 2012. The effect of noise exposure on the cervical vestibular evoked myogenic potential. *Ear Hear* **33**:458–465. doi:10.1097/AUD.0B013E3182498C5F

Angelaki DE, McHenry MQ, Dickman JD, Newlands SD, Hess BJM. 1999. Computation of Inertial Motion: Neural Strategies to Resolve Ambiguous Otolith Information. *Journal of Neuroscience* **19**:316–327. doi:10.1523/JNEUROSCI.19-01-00316.1999

Bartikofsky D, Hertz MJ, Bauer DS, Altschuler R, King WM, Stewart CE. 2023. Balance beam crossing times are slower after noise exposure in rats. *Front Integr Neurosci* **17**:1196477. doi:10.3389/FNINT.2023.1196477

Boyle R, Goldberg JM, Highstein SM. 1992. Inputs from regularly and irregularly discharging vestibular nerve afferents to secondary neurons in squirrel monkey vestibular nuclei. III. Correlation with vestibulospinal and vestibuloocular output pathways. *J Neurophysiol* **68**:471–484. doi:10.1152/JN.1992.68.2.471

Carriot J, Jamali M, Chacron MJ, Cullen KE. 2017. The statistics of the vestibular input experienced during natural self-motion differ between rodents and primates. *Journal of Physiology* **595**:2751–2766. doi:10.1113/JP273734

Carriot J, Jamali M, Chacron MJ, Cullen KE. 2014. Statistics of the vestibular input experienced during natural self-motion: Implications for neural processing. *Journal of Neuroscience* **34**:8347–8357. doi:10.1523/JNEUROSCI.0692-14.2014

Colledge NR, Wilson JA, Macintyre CCA, Maclennan WJ. 1994. The Prevalence and Characteristics of Dizziness in an Elderly Community.

Collins EG. 1948. Aural trauma caused by gunfire:Report on a clinical investigation of 108 soldiers exposed to gunfire who, on some occasion, had complained of injury to their ears. *J Laryngol Otol* **62**:358–390. doi:10.1017/S0022215100009099

Contini D, Holstein GR, Art JJ. 2020. Synaptic cleft microenvironment influences potassium permeation and synaptic transmission in hair cells surrounded by calyx afferents in the turtle. *J Physiol* **598**:853–889. doi:10.1113/JP278680

Contini D, Price SD, Art JJ. 2017. Accumulation of K+ in the synaptic cleft modulates activity by influencing both vestibular hair cell and calyx afferent in the turtle. *J Physiol* **595**:777–803. doi:10.1113/JP273060

Cullen KE, Roy JE, Sylvestre PA. 2001. Signal processing by vestibular nuclei neurons is dependent on the current behavioral goal. *Ann N Y Acad Sci* **942**:345–363. doi:10.1111/J.1749-6632.2001.TB03759.X

Curthoys IS, Vulovic V, Burgess AM, Sokolic L, Goonetilleke SC. 2016. The response of guinea pig primary utricular and saccular irregular neurons to bone-conducted vibration (BCV) and air-conducted sound (ACS). *Hear Res* **331**:131–143. doi:10.1016/J.HEARES.2015.10.019

Desai SS, Zeh C, Lysakowski A. 2005. Comparative morphology of rodent vestibular periphery. I. Saccular and utricular maculae. *J Neurophysiol* **93**:251–266. doi:10.1152/JN.00746.2003/ASSET/IMAGES/LARGE/Z9K0010542921007.JPEG

Desmadryl G, Dechesne CJ. 1992. Calretinin immunoreactivity in chinchilla and guinea pig vestibular end organs characterizes the calyx unit subpopulation. *Exp Brain Res* **89**:105–108. doi:10.1007/BF00229006

Fetoni AR, Ferraresi A, Picciotti P, Gaetani E, Paludetti G, Troiani D. 2009. Noise induced hearing loss and vestibular dysfunction in the guinea pig. *Int J Audiol* **48**:804–810. doi:10.3109/14992020903023140

Goldberg JM, Desmadryl G, Baird RA, Fernandez C. 1990. The vestibular nerve of the chinchilla. V. Relation between afferent discharge properties and peripheral innervation patterns in the utricular macula. *https://doi.org/101152/jn1990634791* **63**:791–804. doi:10.1152/JN.1990.63.4.791

Govindaraju AC, Quraishi IH, Lysakowski A, Eatock RA, Raphael RM. 2023. Nonquantal transmission at the vestibular hair cell–calyx synapse: KLV currents modulate fast electrical and slow K+ potentials. *Proc Natl Acad Sci U S A* **120**. doi:10.1073/pnas.2207466120

Heidenreich KD, Carender WJ, Heidenreich MJ, Telian SA. 2010. Strategies to distinguish benign paroxysmal positional vertigo from rotational vertebrobasilar ischemia. *Ann Vasc Surg* **24**:553.e1-553.e5. doi:10.1016/J.AVSG.2009.09.018

Highstein SM, Goldberg JM, Moschovakis AK, Fernandez C. 1987. Inputs from regularly and irregularly discharging vestibular nerve afferents to secondary neurons in the vestibular nuclei of the squirrel monkey. II. Correlation with output pathways of secondary neurons. *J Neurophysiol* **58**:719–738. doi:10.1152/JN.1987.58.4.719

Holt JC, Jordan PM, Lysakowski A, Shah A, Barsz K, Contini D. 2017. Muscarinic Acetylcholine Receptors and M-Currents Underlie Efferent-Mediated Slow Excitation in Calyx-Bearing Vestibular Afferents. *Journal of Neuroscience* **37**:1873–1887. doi:10.1523/JNEUROSCI.2322-16.2017

Holt XJC, Kewin K, Paivi X, Jordan M, Cameron P, Klapczynski M, Mcintosh JM, Crooks PA, Dwoskin LP, Lysakowski A. 2015. Pharmacologically Distinct Nicotinic Acetylcholine Receptors Drive Efferent-Mediated Excitation in Calyx-Bearing Vestibular Afferents. doi:10.1523/JNEUROSCI.3388-14.2015

Hsu WC, Wang J Der, Lue JH, Day AS, Young YH. 2008. Physiological and morphological assessment of the saccule in guinea pigs after noise exposure. *Archives of Otolaryngology - Head and Neck Surgery* **134**:1099–1106. doi:10.1001/ARCHOTOL.134.10.1099

Hunt MA, Miller SW, Nielson HC, Horn KM. 1987. Intratympanic injection of sodium arsanilate (atoxyl) solution results in postural changes consistent with changes described for labyrinthectomized rats. *Behavioral neuroscience* **101**:427–428. doi:10.1037//0735-7044.101.3.427

Jamali M, Carriot J, Chacron MJ, Cullen KE. 2019. Coding strategies in the otolith system differ for translational head motion vs. static orientation relative to gravity. *Elife* **8**. doi:10.7554/eLife.45573

Jones TA, Jones SM. 1999. Short latency compound action potentials from mammalian gravity receptor organs. *Hear Res* **136**:75–85. doi:10.1016/S0378-5955(99)00110-0

Jones TA, Jones SM, Vijayakumar S, Brugeaud A, Bothwell M, Chabbert C. 2011. The adequate stimulus for mammalian linear vestibular evoked potentials (VsEPs). *Hear Res* **280**:133–140. doi:10.1016/j.heares.2011.05.005

Juntunen J, Ylikoski J, Ojala M, Matikainen E, Ylikoski M, Vaheri E. 1987. POSTURAL BODY SWAY AND EXPOSURE TO HIGH-ENERGY IMPULSE NOISE. *The Lancet* **330**:261–264. doi:10.1016/S0140-6736(87)90840-3

Keshner EA. 2000. Modulating active stiffness affects head stabilizing strategies in young and elderly adults during trunk rotations in the vertical plane, Gait and Posture.

Kumar K, Vivarthini C, Bhat J. 2010. Vestibular evoked myogenic potential in noise-induced hearing loss. *Noise Health* **12**:191–194. doi:10.4103/1463-1741.64973

Lee C, Holt JC, Jones TA. 2017. Effect of M-current modulation on mammalian vestibular responses to transient head motion. *J Neurophysiol* **118**:2991–3006. doi:10.1152/JN.00384.2017

Leonard RB, Kevetter GA. 2002. Molecular probes of the vestibular nerve I. Peripheral termination patterns of calretinin, calbindin and peripherin containing fibers. *Brain Res* **928**:8–17. doi:10.1016/S0006-8993(01)03268-1

Lysakowski A, Goldberg JM. 1997. A Regional Ultrastructural Analysis of the Cellular and Synaptic Architecture in the Chinchilla Cristae Ampullares. *J Comp Neurol* **389**:419–443. doi:10.1002/(SICI)1096-9861(19971222)389:3<419::AID-CNE5>3.0.CO;2-3

Lysakowski A, Govindaraju C, Raphael RM. 2022. Structural and functional diversity of mitochondria in vestibular/cochlear hair cells and vestibular calyx afferents. *Hear Res* **426**:108612. doi:10.1016/j.heares.2022.108612

Man A, Naggan L, Segal S. 1980. Vestibular involvement in acoustic trauma. (An electronystagmographic study). *J Laryngol Otol* **94**:1395–1400. doi:10.1017/S0022215100090228

Mangabeira-Albernaz PL, Covell WP, Eldredge DH. 1959. Changes in the vestibular labyrinth with intense sound. *Laryngoscope* **69**:1478–1493. doi:10.1288/00005537-195912000-00002

Mccabe BF, Lawrence M. 1958. The effects of intense sound on the non-auditory labyrinth. *Acta Otolaryngol* **49**:147–157. doi:10.3109/00016485809134738

Meredith FL, Li GQ, Rennie KJ. 2011. Postnatal expression of an apamin-sensitive K(Ca) current in vestibular calyx terminals. *Journal of Membrane Biology* **244**:81–91. doi:10.1007/S00232-011-9400-8/FIGURES/6

Nazareth AM, Jones TA. 1998. Original Contribution CENTRAL AND PERIPHERAL COMPONENTS OF SHORT LATENCY VESTIBULAR RESPONSES IN THE CHICKEN, Journal of Vestibular Research.

Oosterveld WJ, Polmant AR, Schoonheytt J. 1982. Vestibular Implications of Noise-Induced Hearing Loss. *Br J Audiol* **16**:227–232. doi:10.3109/03005368209081467

Ossenkopp KP, Prkacin A, Hargreaves EL. 1990. Sodium arsanilate-induced vestibular dysfunction in rats: Effects on open-field behavior and spontaneous activity in the automated digiscan monitoring system. *Pharmacol Biochem Behav* **36**:875–881. doi:10.1016/0091-3057(90)90093-W

Pastras CJ, Curthoys IS, Asadnia M, McAlpine D, Rabbitt RD, Brown DJ. 2023. Evidence that ultrafast non-quantal transmission underlies synchronized vestibular action potential generation. *J Neurosci* JN-RM-1417-23. doi:10.1523/JNEUROSCI.1417-23.2023

Paul SS, Walther RG, Beseris EA, Dibble LE, Lester ME. 2017. Feasibility and Validity of Discriminating Yaw Plane Head-on-Trunk Motion Using Inertial Wearable Sensors. *IEEE Trans Neural Syst Rehabil Eng* **25**:2347–2354. doi:10.1109/TNSRE.2017.2740945

Perez R, Freeman S, Cohen D, Sohmer H. 2002. Functional impairment of the vestibular end organ resulting from impulse noise exposure. *Laryngoscope* **112**:1110–1114. doi:10.1097/00005537-200206000-00032

Plotnik M, Elidan J, Mager M, Sohmer H. 1997. Short latency vestibular evoked potentials (VsEPs) to linear acceleration impulses in rats. *Electroencephalography and Clinical Neurophysiology - Evoked Potentials* **104**:522–530. doi:10.1016/S0168-5597(97)00062-2

Seseña E, Vega R, Soto E. 2014. Activation of μ-opioid receptors inhibits calcium-currents in the vestibular afferent neurons of the rat through a cAMP dependent mechanism. doi:10.3389/fncel.2014.00090

Shinder ME, Taube JS. 2010. Differentiating ascending vestibular pathways to the cortex involved in spatial cognition. *Journal of Vestibular Research* **20**:3–23. doi:10.3233/VES-2010-0344

Shupak A, Bar-el E, Podoshin L, Spitzer O, Gordon CR, Ben-david J. 1994. Vestibular findings associated with chronic noise induced hearing impairment. *Acta Otolaryngol* **114**:579–585. doi:10.3109/00016489409126109

Smith PF, Zheng Y. 2013. From ear to uncertainty: Vestibular contributions to cognitive function. *Front Integr Neurosci* **7**:64621. doi:10.3389/FNINT.2013.00084/BIBTEX

Stewart C, Yu Y, Huang J, Maklad A, Tang X, Allison J, Mustain W, Zhou W, Zhu H. 2016. Effects of high intensity noise on the vestibular system in rats. *Hear Res* **335**:118–127. doi:10.1016/j.heares.2016.03.002

Stewart C. E., Bauer DS, Kanicki AC, Altschuler RA, King WM. 2020. Intense noise exposure alters peripheral vestibular structures and physiology. *J Neurophysiol* **123**:658–669. doi:10.1152/JN.00642.2019

Stewart Courtney Elaine, Holt AG, Altschuler RA, Cacace AT, Hall CD, Murnane OD, King WM, Akin FW. 2020. Effects of Noise Exposure on the Vestibular System: A Systematic Review. *Front Neurol* **11**:1327. doi:10.3389/FNEUR.2020.593919/BIBTEX

Stewart CE, Kanicki AC, Altschuler RA, King WM. 2018. Vestibular short-latency evoked potential abolished by low-frequency noise exposure in rats. *J Neurophysiol* **119**:662–667. doi:10.1152/JN.00668.2017

Sultemeier DR, Hoffman LF. 2017. Partial Aminoglycoside Lesions in Vestibular Epithelia Reveal Broad Sensory Dysfunction Associated with Modest Hair Cell Loss and Afferent Calyx Retraction. *Front Cell Neurosci* **11**. doi:10.3389/FNCEL.2017.00331

Tamura H, Ohgami N, Yajima I, Iida M, Ohgami K, Fujii N, Itabe H, Kusudo T, Yamashita H, Kato M. 2012. Chronic exposure to low frequency noise at moderate levels causes impaired balance in mice. *PLoS One* **7**. doi:10.1371/JOURNAL.PONE.0039807

Tseng CC, Young YH. 2013. Sequence of vestibular deficits in patients with noise-induced hearing loss. *Eur Arch Otorhinolaryngol* **270**:2021–2026. doi:10.1007/S00405-012-2270-6

Ventura CM, Kalluri R. 2019. Enhanced activation of HCN channels reduces excitability and spike-timing regularity in maturing vestibular afferent neurons. *Journal of Neuroscience* **39**:2860–2876. doi:10.1523/JNEUROSCI.1811-18.2019

Vignaux G, Chabbert C, Gaboyard-Niay S, Travo C, Machado ML, Denise P, Comoz F, Hitier M, Landemore G, Philoxène B, Besnard S. 2012. Evaluation of the chemical model of vestibular lesions induced by arsanilate in rats. *Toxicol Appl Pharmacol* **258**:61–71. doi:10.1016/J.TAAP.2011.10.008

Viola P, Scarpa A, Pisani D, Petrolo C, Aragona T, Spadera L, De Luca P, Gioacchini FM, Ralli M, Cassandro E, Cassandro C, Chiarella G. 2020. Sub-Clinical Effects of Chronic Noise Exposure on Vestibular System. *Transl Med UniSa* **22**:19.

Wang YP, Young YH. 2007. Vestibular-evoked myogenic potentials in chronic noise-induced hearing loss. *Otolaryngology - Head and Neck Surgery* **137**:607–611. doi:10.1016/j.otohns.2007.05.005

Ylikoski J, Juntunen J, Matikainen E, Ylikoski M, Ojala M. 1988. Subclinical vestibular pathology in patients with noise-induced hearing loss from intense impulse noise. *Acta Otolaryngol* **105**:558–563. doi:10.3109/00016488809119520

Zobeiri OA, Ostrander B, Roat J, Agrawal Y, Cullen KE. 2021. Loss of peripheral vestibular input alters the statistics of head movement experienced during natural self-motion. *J Physiol* **599**:2239–2254. doi:10.1113/JP281183

Figure 1. Noise exposure attenuates VsEP responses. **A**: Representative VsEP recordings before noise exposure (Baseline, black traces), 1-week post-exposure (magenta traces), and 4-week post-exposure (green traces). Stimulus jerk levels were (1) 0.32, (2) 1.1, (3) 2.2, (4) 3.2, and (5) 5.5 G/ms. VsEP peaks P1, N1, P2, N2, and P3 peaks are denoted with a gray arrow, red star, red arrow, blue star, and blue arrow, respectively, on the left bottom VsEP trace. **B**: N1P2 amplitude; **C**: N2P3 amplitude; **D**; N1 peak latency; **E**; N2 peak latency as a function of stimulus level for baseline (black), 1 week (magenta), and 4 weeks (green) post noise exposure for the animal shown in **A**.

Figure 2. **A**, **B**, and **C**: The population-mean % change in the N1P2 amplitude (**A**), N2P3 amplitude (**B**), and N1 latency (**C**) at 1 week (magenta) and 4 weeks (green) after noise exposure compared to the baseline as a function of stimulus level. The % change values are collapsed across stimulus levels and shown at the column denoted as 'All' to show the overall effect. Each dot denotes an individual animal's data point, the cross symbols denote population means, and error bars denote standard error of the mean (SEM). Asterisks denote significance levels at $p < 0.01$ and $p < 0.05$ by t-test for individual stimulus levels and all-levels collapsed, respectively. **D**: % change in N2P3 amplitude as a function of % change in N1P2 amplitude at 1 week after noise exposure for stimulus levels 1 through 5. **E**: Same as **D**, except the absolute value of % change in N1P2 amplitude is plotted on x-axis. **F**: % change in N2P3 amplitude as a function of % change in N1 peak latency 1 week after noise exposure. In panels **D~F**, linear fits are shown as a red solid line for significant correlations ($p < 0.05$) with Pearson's correlation test. Red, dotted lines denote marginally significant correlations ($p < 0.10$). Each combination of symbol and color designates an individual animal's data point, and is consistently used for all other figures.

Figure 3: Noise exposure reduces the number of CR+ calyces in otolith striolar regions in relation to changes in VsEP waveforms **A**: Number of CR+ calyces in the saccule of control animals (black) and noise-treated animals (red). **B**: The same as **A**, but for the utricle. **C**: CR+ calyces in saccule against CR+ calyces in utricle in individual animals. **D**: % change in N2P3 amplitude against the number of CR+ calyces in the saccule (left panels) and the utricle (right panels). **E**: % change in N1 latency against the number of CR+ calyces in the saccule (left panels), the utricle (middle panels), and averaged between saccule and utricle (right panels). In **D** and **E**, the VsEP changes at 1 week and those at 4 weeks after noise exposure are shown in top rows and bottom rows, respectively.

Figure 4: Noise exposure causes reduced head motion durations in relation to VsEP changes and numbers of reduced CR+ calyces in the otolith organs. **A**: The stable period duration versus days before or after noise exposure for a representative animal. Blue squares, daily medians pre-noise; red squares, daily medians post-noise. Error bars show the standard deviation for each day. **B**: Circles: daily median stable period durations (squares) averaged across pre- and post-exposure days for pre- (blue) and post-noise exposure (red). Error bars, standard deviation. **C**: The change in the stable period duration, defined as average daily median duration of post-exposure days (red circles in **B**) minus average daily median duration of pre-exposure days (blue circles in **B**), is shown for the arsanilate- (n = 3) and noise-exposed animals (n = 17). Filled circles denote a significant change (p<0.05 by t-test) in individual animals. **D**, **E**, and **F**: The same as **A**, **B**, and **C**, but for the proportion of stable time. **G**: Left three panels: change in the stable period duration versus the number of CR+ calyces in saccule, in utricle, and their average. Right five panels: change in the stable period duration and change in VsEP N2P3 amplitude 4 weeks after noise exposure, for stimulus levels 1 through 5. **H**: The same as **G**, but for change in the proportion of stable time.

Figure 5: Noise exposure altered head rotational motion in pitch in some animals. **A**: Speed of pitch-axis rotational motion plotted against days before or after noise exposure. RMS speeds were calculated from pitch-axis angular velocity during any segment of motion longer than 0.5 sec, then transformed into z-scores based on mean and standard deviation of RMS speeds in narrow bins determined by the durations of active periods from pre-exposure dates. As in Fig. 3, Blue symbols denote pre-noise and & red post noise exposure medians. **B**: daily median z-score of RMS speeds (squares) averaged across pre- and post-exposure days. The asterisk denotes significance ($p < 0.05$, t-test). **C~F**: Changes in z-score transformed RMS angular velocity in roll-axis (**C**), pitch-axis (**D**), yaw-axis (**E**), and vector-average of all 3 axes (**F**) are shown for arsanilate- (n = 3) and noise-exposed animals (n = 17). Change is defined as average daily median z-score transformed RMS of post-exposure days (red circle in **B**) minus that of pre-exposure days (blue circle in **B**). Filled circles indicate statistical significance for individual animals. **G**: Correlation between change in RMS of the pitch-axis rotational motion and number of CR+ calyces in saccule and utricle. **H**: Correlation between change in RMS pitch-axis rotational motion and change in the VsEP N1 peak latency 1 week after noise exposure, at stimulus levels 1 through 5 (from right to left panel). I: Frequency profiles of y-axis rotational motion of before and after arsanilate or noise was obtained by Fourier transform of angular velocity (left panels) and angular acceleration (right panels) when the rat's head was in motion for durations > 0.5 sec. The power spectrum before (blue) and after (red) exposure is shown for a representative animal exposed to arsanilate. **J**: the same as I, for a rat whose pitch-axis RMS velocity increased following noise exposure. **K**: the same as I, for a rat whose pitch-axis RMS velocity decreased after noise exposure. **L**: The ratio of the post- to the pre-noise power spectrums for 3 animals exposed to arsanilate. **M** and **N**: The same as **L**, but for noise-exposed animals with increases (**M**) or decreases (**N**) in RMS y-axis velocity ($p < 0.10$) following noise.

Figure 6: Noise exposure caused changes in head orientation about the roll and pitch axes. **A**: Roll orientation with respect to gravity and the rat's head post. **B**: Distribution of roll angles as a function of motionless period duration for pre- (blue) and post-exposure (red) in a representative rat. **C**: The mean and standard deviation of roll angle as a function of the duration of motionless period. **D**: Probability histogram of Z-score transformed roll angles collapsed across durations > 1 sec. **E**: Z-score transformed roll orientation from motionless periods > 1 sec plotted against days before or after noise exposure. Squares: daily median z-scores of roll; error bars: daily standard deviation. **F**: Daily median z-score of roll (squares) averaged across pre- and post-exposure days: blue circle, pre-, red circle, post-exposure, error bars, standard deviation. Asterisk denotes statistical significance of the mean and variance (p < 0.05). **G**: The daily standard deviation of z-score transformed roll (squares) averaged across pre- and post-exposure days. Asterisk denotes significance of the mean (p < 0.05). **H~N**: Same **A~G** except these show pitch instead of roll.

Figure 7: Changes in the daily median z-score transformed roll (**A**) and pitch orientations (**D**) for arsanilate (n = 3) and noise-exposed rats (n = 17). Change is defined as the average daily median z-score transformed orientation of post-exposure days (e.g., the red circle in Figure 5F) minus that of pre-exposure days (e.g., the blue circle in Figure 5F). **B** and **E**: Changes in variance of daily median z-score transformed roll (**B**) and pitch (**E**). **C** and **F**: Changes in daily standard deviation of z-score transformed roll (**C**) and pitch (**F**). In **A ~ F** filled circles indicate statistical significance in individual animals. **G**: Relationship between absolute values of the change in median pitch and roll. **H**: Relationship between absolute value of the change in median pitch and number of CR+ calyces in saccule (left panel) and utricle (right panel).