Metabolic and behavioral alterations associated with viral vector-mediated toxicity in the paraventricular hypothalamic nucleus

Rohan Savani[1,2], Erin Park[1,2], Nidhi Busannagari[1,2], Yi Lu[1], Hyook Joon Kwon[3], Le Wang[1, *],

Zhiping P. Pang[1, *]

[1]The Child Health Institute of New Jersey, Robert Wood Johnson Medical School, Rutgers, The State University of New Jersey, New Brunswick, NJ, USA
[2]Department of Cell Biology and Neuroscience, Undergraduate Program, Rutgers, The State University of New Jersey, New Brunswick, NJ, USA
[3]Department of Medicine, Rutgers-Robert Wood Johnson Medical School, New Brunswick, NJ, 08901, USA

Main text Words: 3467

Figures: 3

*Corresponding authors:

Le Wang: lw611@rwjms.rutgers.edu

Zhiping P. Pang: pangzh@rwjms.rutgers.edu

ABSTRACT

Objective: Combining adeno-associated virus (AAV)-mediated expression of Cre recombinase with genetically modified floxed animals is a powerful approach for assaying the functional role of genes in regulating behavior and metabolism. Extensive research in diverse cell types and tissues using AAV-Cre exemplifies advantages over crossing with Cre driver lines in several aspects, including saving time and avoiding developmental/compensatory issues. We initially sought to study the impact of ablation of corticotropin-releasing hormone (CRH) in the paraventricular hypothalamic nucleus (PVN) using intracranial AAV8-hSyn-Cre injection in adult animals.

Methods: In this study, we stereotactically injected AAV8-hSyn-Cre or a control viral vector (1E13 and 1E12 vg/ml, respectively, both at 150 nL) in both Crh-floxed and wild-type mouse PVN to assess behavioral and metabolic impacts. We then used immunohistochemical markers to systematically evaluate the density of hypothalamic peptidergic neurons and glial cells.

Results: We found that delivery of one preparation of AAV8-hSyn-Cre in the PVN led to the development of obesity, hyperphagia, and anxiety-like behaviors. This effect occurred independent of sex and in both floxed and wild-type mice. We subsequently found that AAV8-hSyn-Cre led to neuronal cell death and gliosis at the site of viral vector injections. These behavioral and metabolic deficits were dependent on injection into the PVN specifically.

Conclusions: Our findings reveal that delivery of AAV8-hSyn-Cre could lead to cellular toxicity and lesions in the PVN that cause robust metabolic and behavioral impacts. These alterations can complicate the interpretation of AAV-Cre-mediated gene knockout and highlight the need for rigorous controls.

Keywords: AAV, toxicity, paraventricular hypothalamus, obesity

1. INTRODUCTION

Located in the ventral forebrain, the paraventricular nucleus of the hypothalamus (PVN) acts as a major integrative center for diverse regulatory functions, including neuroendocrine response, autonomic tone, and complex behaviors including food intake and social interaction (1-3). Dissecting the molecular, morphological, and functional heterogeneity of the PVN has thus far given rise to several distinct cell types involved in the control of these processes.

Refining our understanding of how the PVN integrates and outputs multimodal information across neuroendocrine and autonomic pathways may inform several disorders, including cardiovascular, metabolic, and neuropsychiatric disorders (2, 4, 5). However, the PVN contains spatially intermingled but functionally distinct populations that can only be selectively targeted through genetic access, such as conditional knockout (KO) by Cre-lox recombination (6, 7). While much work relies on promoter-driven Cre expression, another standard method relies on adeno-associated virus-mediated Cre expression (AAV-Cre) (8-11). Methods such as AAV-mediated Cre delivery in adult mice are valuable approaches to bypass transient developmental transcription and ensure high-fidelity expression in specific brain regions (8, 12). However, emerging evidence suggests that there may be deleterious effects associated with AAV-Cre, resulting in morphological and behavioral abnormalities even in the absence of loxP sequences (13, 14).

In this study, we assessed the striking phenotype that emerged from the delivery of an AAV-Cre-prepared virus in the PVN of adult *Crh-flox* ($Crh^{f/f}$) mice. We found that $Crh^{f/f}$ mice with putatively knocked out corticotropin-releasing hormone (CRH) via AAV-mediated expression of Cre recombinase in the PVN rapidly developed obesity and metabolic syndrome, which was an intriguing and unexpected finding. However, subsequent rigorous analysis revealed that both CRH- and oxytocin-expressing neuron densities significantly decreased, likely due to cellular toxicity at the injection sites. Our study confirms the importance of the PVN in the control of metabolism and highlights the potential for toxicity due to a commonly used AAV, which necessitates careful interpretation and proper controls for experiments utilizing viral vectors.

2. MATERIALS AND METHODS

2.1 Animals

All studies and procedures involving mice were approved by the Rutgers University Institutional Animal Care and Use Committee (IACUC) and by the National Institute of Health (NIH) guidelines. The animals used in this study were 5-7 weeks old at the time of stereotactic surgery. $Crh^{f/f}$ mice

(15), *Glp1r^{f/f}* mice (16), and wild-type (C57BL/6) mice were used in this study. For each

experimental paradigm, littermate mice were randomized to each experimental group on the basis

of body weight. Both males and females were used in this study. Mice of both groups were housed

together. Investigators were blinded to treatment. The sample size required (n=5-7) for each group

was estimated from previous studies (17).

2.2 AAV vectors and stereotactic surgery

The AAVs used in this study included pAAV-hSyn-EGFP (Addgene, #50465-AAV8, titer: titer

7×10^{12} vg/mL) and pENN.AAV.hSyn.HI. eGFP-Cre.WPRE.SV40 (Addgene, #105540-AAV8, titer:

1×10^{13} vg/mL). For surgical injections, mice were anesthetized with isoflurane and placed in a

stereotactic frame. Burr holes were drilled above the injection sites. 150 nL of the viruses were

bilaterally stereotactically injected in the PVN (AP: -0.7, ML: ±0.18, DV: -4.7) at a rate of 1 nL/s

(eGFP for control group, Cre for experimental group). For experiments in the dorsolateral septum

(dLS), 100 nL of viruses were bilaterally injected in the dLS (AP: +0.5, ML: ±0.45, DV: -2.7) at a

rate of 1 nL/second. Mice recovered for at least 3 weeks prior to conducting behavioral

experiments. Injection sites were confirmed by *post hoc* histological examinations.

2.3 Metabolic phenotyping

All experiments were conducted during the light cycle unless otherwise stated. Investigators were

blinded to conditions during experiments.

Plasma corticosterone measurements: Blood was collected either at the beginning of the ITT test

or during transcardiac perfusion. At least 10 µL was collected into EDTA-coated tubes and then

centrifuged at 8000rpm at 4°C for 5 minutes. Extracted plasma was stored at -80° C until further

processing. Corticosterone measurements were obtained with an ELISA as per manufacturer's

instructions (Enzo Life Sciences, ADI-900-097).

Food intake: Mice were singly housed before the experiment with ad libitum access to normal

chow (Purina Mouse Diet 5058) and water. Chow remaining in the cage after each day (12 hr.

light/dark cycle) was measured for 5 days and averaged together.

EchoMRI/fat mass measurements: Whole body composition and fat mass was measured by

EchoMRI (Echo Medical Systems, Houston, Texas).

Comprehensive Lab Animal Monitoring System (CLAMS) assays: *Crh^{f/f}* mice were placed into

metabolic phenotyping cages (Columbus Instruments, OH, USA). Data was recorded every 15

minutes. Data was collected for 3 days (both light and dark cycles) after 2 days of habituation to
single housing within the cages.

Glucose tolerance test (GTT) and insulin tolerance test (ITT): Mice were fasted overnight (GTT)
or for 6 hours (ITT). To minimize stress, 0.5 grams of chow was provided. For GTT, mice were
intraperitoneally (IP) administered 20% glucose. Blood glucose measurements were collected at
the 0, 15-, 30-, 60-, and 120-minute time points. For ITT, mice were IP-administered insulin.
Injection volumes were 0.1 mL/10 g body weight. Blood glucose measurements were collected at
the 0-, 15-, 30-, and 60-minute time points.

2.4 Behavioral assays
Open field test: Mice were placed in the middle of a custom-made 45 x 45 cm square open field
arena for 10 minutes. Analysis was performed using DeepLabCut v2.2 (18). We labeled 150-200
frames taken from 14 or 10 videos, as resolution changed between recording sessions (95% of
frames were used for training, 5% for testing). We used a ResNet50-based network with default
parameters for 225,000 or 450,000 training iterations (until loss plateaued). After analyzing open-
field videos, we then calculated the time spent in the inner zone (inner 3/5 of the arena, ~25x25cm)
using a region of interest tool. Distance traveled was calculated with custom codes. Investigators
were blinded to the condition during analysis.

Light-dark box test: Mice were placed in the SmartCage system (AfaSci) light side of a commercial
light-dark box apparatus consisting of one dark and one illuminated compartment connected by
a door. Time spent in each zone, latency to enter the dark zone, and number of entries were
measured for 10 minutes. Data were analyzed using Excel.

2.5 Histology and immunohistochemistry (IHC) analysis
Mice used for IHC analysis of CRH and oxytocin (OXT) were intracerebroventricularly
administered colchicine a day prior to perfusion. Mice were anesthetized with isoflurane and
perfused with PBS (pH 7.4) and then 4% paraformaldehyde (PFA). Coronal brain slices were
cryosectioned (50 μm). For IHC analyses, brain slices were first incubated in blocking buffer (4%
BSA, 1% goat serum in PBS) for 1 hour, then in primary antibody overnight at 4°C. Slices were
washed with blocking buffer 3x15 min prior to incubation with a secondary antibody at RT for 2
hours, followed by further washing with PBS. Sections were mounted with Fluoroshield with DAPI
(Sigma-Aldrich, # F6057). Images were collected with a Zeiss LSM700 confocal microscope. A Z-
stack captured the entire thickness of the section with fluorescence. NeuN-stained slices were

imaged at a 5 µm-thickness interval for a total range of 25 µm. Imaging analysis was performed
using NIH-ImageJ and with the Allen Brain Atlas as a reference. The number of OXT- and CRH-
cells in the PVN was counted with the Cell Counter plugin. To quantify NeuN density, images were
background subtracted (20-pixel radius) before converting to a binary image. Then, segmented
particles were filtered by despeckling twice and watershed transforming, followed by final counting
(minimum size of 7 µm^2). GFAP and Iba1 fluorescence area was calculated by dividing the area
of thresholded signal by total PVN area. The quantifications were performed by two independent
investigators who were blinded to the condition during analysis.

2.6: Antibodies
The primary antibodies used in this study were rabbit Anti-CRH (1:500, BMA Biomedicals, #T-
4037, RRID: AB_2314240), mouse Anti-OXT (1:500 Sigma-Aldrich, #MABN844), rabbit Anti-
NeuN (1:500, Millipore, #ABN78, RRID: AB_10807945), rabbit Anti-GFAP (1:1000, Dako, #Z0334,
RRID: AB_10013382), and chicken Anti-Iba1 (1:500, Aves Labs, #IBA1-0020).

2.7 Statistical analysis
Statistical analyses were performed using GraphPad Prism 9. Data were analyzed using
Student's two-tailed t-tests and two-way ANOVA followed by Bonferroni's multiple comparisons
tests. Data are presented as mean ± SEM. A p-value < 0.05 was considered statistically
significant.

3. RESULTS

3.1 Delivery of AAV-mediated Cre in the PVN rapidly induces behavioral changes and
metabolic syndrome in Crh$^{f/f}$ animals.
Our initial goal was to apprise the functional role of CRH by knocking it out in the PVN in adult
animals. To this end, we bilaterally stereotactically injected AAV8-hSyn-eGFP-Cre in the PVN of
5-7-week-old *Crh$^{f/f}$* mice, with AAV-hSyn-eGFP (Control) injected in littermate controls (**Fig. 1A**).
After four weeks of recovery, we found that plasma corticosterone levels in Cre-injected *Crh$^{f/f}$* mice
were markedly reduced, which was highly suggestive of possible loss of function of CRH (**Fig.
1B**). Even with normal chow feeding, the Cre-injected mice showed robust and rapid body weight
gain within days. Cre-injected mice reached an average of 50.5 ± 0.7 grams in 6 weeks, while the
average body weight of control animals was 28.2 ± 0.7 grams (**Fig. 1C**). The increased body
weight in the Cre-injected group might be, at least in part, due to elevated food intake (**Fig. 1D**).
We repeated these experiments in an independent cohort of *Crh$^{f/f}$* female mice (**Fig. S1A**). In this,

the Cre-injected group fed with normal chow weighed 45.2 ± 4.2 grams 6 weeks after viral injection

(**Fig. S1B**), indicating that the changes observed thus far occur in a sex-independent manner.

As PVN-derived CRH is known to be involved in stress-related behaviors, coping, and relief (19-

22), we hypothesized that reduced CRH release and corticosterone levels (**Fig. 1B**) in the Cre-

injected animals would be anxiolytic. To test this hypothesis, we examined anxiety-like behavior

in AAV-Cre injected $Crh^{f/f}$ male mice in the open-field and light-dark box assays (**Fig. S2**). Contrary

to what we hypothesized, we found that the Cre-injected mice spent less time in the center zone

in the open-field assay (**Fig. S2B**), and they spent more time in the dark zone during the light-

dark box test (**Fig. S2D**). These tests suggested that AAV-Cre in the adult PVN may lead to

heightened anxiety-like behaviors.

Nevertheless, the drastic increase in body weight of AAV-Cre-injected animals prompted us to

further assess their metabolic phenotype. We used CLAMS to measure energy intake and

expenditure, as well as to identify any changes in nutrient utilization. Compared to AAV-GFP-

injected animals, AAV-Cre-injected mice consumed less oxygen throughout the entire day (**Fig.**

1E-F). The respiratory exchange ratio was equal between both groups, indicating no shift in

nutrient utilization accompanying increases in weight (**Fig. 1G**). Energy expenditure was

significantly decreased in AAV-Cre-injected mice in both light and dark cycles (**Fig. 1H**). This may

be in part due to a decrease in locomotor activity (**Fig. 1I**). These data suggest that alterations in

energy expenditure may contribute to the development of obesity alongside hyperphagia in AAV-

Cre-injected mice. Consistent with these findings, Cre-injected mice display significantly

increased body fat composition (**Fig. 1J**) and significantly increased fasting plasma glucose levels

measured (**Fig. 1K**). To evaluate any changes in glucose and insulin homeostasis, we also

performed GTT and ITT. GTT revealed that Cre-injected mice had consistently elevated glucose

levels, indicative of higher glucose tolerance (**Fig. 1L**), while ITT revealed that they had insulin

resistance (**Fig. 1M**). These results indicate that Cre recombinase expression in the PVN region

of $Crh^{f/f}$ mice leads to obesity development and maintenance through changes in food intake

behavior and energy expenditure and causes metabolic syndrome resembling type 2 diabetes

mellitus.

Having observed the development of type 2 diabetes and obesity in the PVN Cre-injected $Crh^{f/f}$

mice, we sought to validate these findings as attributable to the depletion of hypothalamic CRH.

We examined CRH expression using immunohistochemistry and found decreased CRH-positive

cell density in the PVN of AAV-Cre-injected mice (**Fig. 2A & 2C**). This suggested that CRH

expression was suppressed, which supports the hypothesis that PVN CRH participates in regulating metabolism. However, this hypothesis is complicated by the observation of decreased OXT expression in brain sections of the same animals with Cre injections (**Fig. 2B & 2D**). Decreased expression of both CRH and OXT was accompanied with a trend toward reduced cell density as measured by DAPI staining (**Fig. 2E**). Based on these data, we caution that virally expressed Cre recombinase may be sufficient to deplete CRH from the PVN of *Crh^{f/f}* animals but may also have an off-target impact on PVN neuronal cells.

3.2 Delivery of AAV-mediated Cre in the PVN rapidly alters behavior and causes metabolic syndrome in wild-type animals.

While OXT expression may have decreased due to the development of obesity, the trend of decreased cell density in the AAV-Cre group led us to hypothesize that the AAV-Cre virus may impact PVN neuronal survival. The toxicity of Cre expression on cellular health has been reported previously (13, 14, 23, 24).

To this end, we examined the impacts of AAV-Cre in C57BL/6 wild-type (WT) mice (**Fig. 3A**). Similar to the AAV-Cre injected *Crh^{f/f}* animals, we found a marked reduction in plasma corticosterone levels in WT mice injected with AAV-Cre (**Fig. 3B**). They also gained significant body weight potentially due to hyperphagia (**Fig. 3C-D**). Given that our observations closely mirrored those found in *Crh^{f/f}* mice, we also assessed changes in anxiety-like behaviors (**Fig. S3A & S3C**). AAV-Cre-injected mice spent less time in the center zone during the open field test (**Fig. S3B**). Consistent with this, Cre mice spent more time in the dark zone of the light-dark box (**Fig. S3D**). These data suggest that the behavioral, metabolic, and cellular alterations observed with Cre injection in the PVN may be due to the expression of the Cre-encoding viral vector.

We next systematically evaluated the number of neurons in the PVN of WT mice following AAV-Cre or AAV-GFP injections using neuronal nuclei (NeuN) staining (**Fig. 3E**). Expression of NeuN was significantly decreased across most of the PVN with the administration of AAV-Cre, likely due to diffusion (**Fig. 3F**). Again, accompanied by the enlarged visual appearance of the third ventricle, the widespread decline in neuronal density in AAV-Cre-expressing mice lends support to the idea of toxic and nonspecific AAV-Cre action, resembling the effects of a chemical lesion for PVN neuronal cells (25, 26). This neuronal cell loss is accompanied by astrogliosis and microglial cell activation (**Fig. 3G-I**).

231 Together, these data suggest that AAV-Cre injection may cause non-specific cell death at the
232 injection site, with PVN neuronal ablation causing obesity and type 2 diabetes. This is consistent
233 with previous reports that PVN lesions result in severe obesity and dysregulated metabolism (25-
234 29).

3.3 Delivery of AAV-mediated Cre in the dorsolateral septum induces neuronal cell lesions
without causing metabolic syndrome in Glp1r $^{f/f}$ animals.

To further confirm the apparent toxicity of the AAV-Cre that we used, we tested it in another brain
region: the dorsal lateral septum (dLS). The glucagon-like peptide-1 receptor (GLP-1R) is highly
expressed in dLS and is implicated in feeding and metabolism (30-33). We assessed if a diluted
AAV-Cre could be used to determine endogenous GLP-1R function. To prevent the possibility of
cellular toxicity due to high AAV titers, we diluted our AAV to 5 x 10^{12} vg/mL and only injected 100
nL. We injected this AAV-Cre or a control vector in *Glp1r $^{f/f}$* mouse dLS (**Fig S4A**), but we did not
observe significant changes in body weight or metabolism within 4 weeks of injection (**Fig S4B**).
However, we found significantly decreased NeuN positive cell density (**Fig S4C-D**), suggesting
that Cre viral vector injection in the mouse brain causes neuronal cell loss.

4. DISCUSSION

In this study, we showed that delivery of one preparation of AAV-Cre recombinase in the PVN of
adult mice can lead to the development of metabolic syndrome and alterations in food-related and
anxiety-like behaviors, as well as metabolic phenotype. We demonstrated that *Crh$^{f/f}$* mice
expressing AAV-Cre in PVN develop obesity largely mediated by hyperphagia and changes in
energy expenditure. Moreover, this develops alongside glucose tolerance and insulin resistance,
as well as decreased locomotion. We investigated whether CRH was truly and specifically
depleted and found off-target effects on OXT expression (**Fig. 2**) and cell loss at sites of Cre viral
vector injections in either *Crh$^{f/f}$* or WT mice. These data suggested off-target effects of our viral
vector and indicate that caution should be used when interpreting the physiological consequences
of AAV injection in adult animals.

Specific depletion of PVN-derived CRH has thus far not been shown to induce massive metabolic
dysregulation when knocked out using Cre expression driven by either *Crh$^{f/f}$* crossed with *Sim1-*
Cre mice or CRISPR (17, 34). Our initial goal was to assess the impact of ablation of PVN CRH
in adult animals by using stereotactic injection of AAV-Cre. We observed massive obesity and
metabolic syndromes that starkly differ from other models of CRH knockout (17, 34-38). Despite

the observation of reduction of CRH expression and plasma corticosterone levels, which may be
an indication of impaired regulation of the hypothalamic-pituitary-adrenal axis (HPA), we surmise
that any metabolic or behavioral deficits reflected off-target effects of AAV-Cre. Indeed, we
confirmed that neuronal cell death and reactive gliosis occurred in the PVN with AAV-Cre
expression, resembling a response to central nervous system injury. Therefore, the study cannot
conclude whether adult ablation of CRH in the PVN leads to metabolic or behavioral changes in
adult animals.

The PVN is speculated to provide top-down control of feeding-related information processing via
intrahypothalamic connections or dense projections to midbrain and hindbrain regions (2). These
results could not provide insights on CRH impact on metabolism but highlight the necessity of the
PVN in regulating energy homeostasis and body weight. These findings are in accord with prior
ablation experiments and more recent work on genetic knockout models specific to the PVN (25,
26). It has been shown that unilateral PVH electrolytic lesions would increase lipid accumulation
in white adipose tissue ipsilateral to the side of the PVH (39). Chronically inhibiting PVN
MC4R/PDYN neuron synaptic release by expressing tetanus toxin or ablating those neurons by
expressing caspase-3 resulted in pronounced hyperphagia, obesity, and a significant elevation in
food consumption (26). Our metabolic phenotyping data revealed that cellular lesions of PVN
develop a strong obesity phenotype largely mediated by hyperphagia and changes in energy
expenditure. Moreover, dysfunction of the PVN induces hypersomnia and mediates the diurnal
rhythm of metabolism in mice (40, 41). Consistent with this, we also found that changes in energy
expenditure were more pronounced in AAV-Cre-injected mice during dark cycles (**Fig. 1H**).

Manipulating gene expression on and off using AAVs in genetically engineered animals is one of
the most powerful tools to bypass transient developmental transcription for studying gene function
in vivo (8, 12). For example, it has been shown that knockout of *Glp1r* via AAV-Cre in PVN
demonstrated a significant obesity phenotype, which was not observed in *Glp1r^{f/f}* mice cross with
SIM1- or NKX2.1-Cre mice (8, 42). Similarly, AAV-CRISPR–Cas9-mediated ablation of leptin
receptor in arcuate nucleus demonstrated that agouti-related peptide (AgRP) neurons but not
proopiomelanocortin (POMC) neurons are required for the primary action of leptin to regulate both
energy balance and glucose homeostasis (12). The present results add to a growing body of
literature suggesting that toxicity occurs irrespective of serotype and brain region, though a
systematic comparison is lacking. Moreover, this effect may occur independent of the specific
virus used; while we used one viral preparation in this study, previous studies have reported
varying levels of success with this AAV-Cre and others (9, 13, 14, 41, 43). In the mesolimbic

reward system, it has been reported that Cre expression in the ventral tegmental area and
substantia nigra can induce toxicity associated with perturbations of dopamine-related behaviors
(13, 14). Further work delineating different viral preparations and optimal titers for neuronal
manipulations will be needed. Inducible systems of Cre expression, such as those using
tamoxifen, offer an appealing alternative to constitutively expressed Cre systems; however,
administration of tamoxifen has also been reported to lead to cytotoxicity and off-target effects
(44).

Our data confirm that the PVN is an important brain region in the regulation of metabolism;
dysfunction of the PVN, such as via neuronal death, causes obesity and type 2 diabetes. We also
conclude that AAV-Cre vector toxicity ablates PVN neurons and results in a phenotype that is not
associated with the gene of interest. These results underscore the importance of rigorous controls
to ensure virus quality and fidelity of action. Caution must especially be exercised when the
observed phenotype closely matches that of lesions in the area of use. Ensuring that expression
in wild-type mice results in no observable changes or that neuronal cell densities are unaltered (if
they are not expected to be) are two such methods of validating that the AAV is not toxic on its
own. Together, viral vectors represent a class of worthwhile tools to manipulate the nervous
system; however, scientists must take care to ensure that the potential for toxicity in commonly
used AAVs is ruled out in experimental design and data interpretation.

AUTHOR CONTRIBUTIONS

R.S. conducted animal behavior assays, histology, and data analysis. E.P. helped with histology and imaging analysis. N.B. helped with histology and imaging analysis. Y.L. conducted stereotactic surgeries, animal behavior analysis, and histology. H.J.K. conducted CLAMS experiments and corresponding data analysis. L.W. conducted stereotactic surgeries. L.W. and Z.P.P. conceived the project, and designed the experiments and wrote the paper with R.S. with input from all authors.

ACKNOWLEDGMENTS

This study was supported by grants from the Robert Wood Johnson Foundation to the Child Health Institute of New Jersey (RWJF grant #74260), and the NIH NIDDK R01DK131452. L.W. was supported by the New Jersey Governor's Council for Medical Research and Treatment of Autism Postdoctoral Fellowship (CAUT24DFP) and the NExT-Metabolism Pilot Award (500301). R.S. was supported by Rutgers Aresty Research Center's independent research award. The authors would like to thank Dr. Mark Rossi (Rutgers University) for critical reading of the manuscript and insightful comments. The graphical abstract was created with BioRender.

DATA AVAILABILITY

Code used to analyze open field test data can be found at https://github.com/RohanSavani/OpenFieldAnalysis. The data from the current study are available from the corresponding authors upon reasonable request.

SUPPLEMENTARY DATA
Supplementary data to this article: Supplemental Figures S1-S4

REFERENCES

1. Ferguson AV, Latchford KJ, Samson WK. The paraventricular nucleus of the hypothalamus - a potential target for integrative treatment of autonomic dysfunction. Expert Opin Ther Targets. 2008;12(6):717-27.

2. Watts AG, Kanoski SE, Sanchez-Watts G, Langhans W. The physiological control of eating: signals, neurons, and networks. Physiological Reviews. 2021;102(2):689-813.

3. Qin C, Li J, Tang K. The Paraventricular Nucleus of the Hypothalamus: Development, Function, and Human Diseases. Endocrinology. 2018;159(9):3458-72.

4. Manaye KF, Lei DL, Tizabi Y, Dávila-García MI, Mouton PR, Kelly PH. Selective neuron loss in the paraventricular nucleus of hypothalamus in patients suffering from major depression and bipolar disorder. J Neuropathol Exp Neurol. 2005;64(3):224-9.

5. Benarroch EE. Paraventricular nucleus, stress response, and cardiovascular disease. Clinical Autonomic Research. 2005;15(4):254-63.

6. Sauer B, Henderson N. Site-specific DNA recombination in mammalian cells by the Cre recombinase of bacteriophage P1. Proceedings of the National Academy of Sciences. 1988;85(14):5166-70.

7. Tsien JZ, Chen DF, Gerber D, Tom C, Mercer EH, Anderson DJ, et al. Subregion- and Cell Type–Restricted Gene Knockout in Mouse Brain. Cell. 1996;87(7):1317-26.

8. Liu J, Conde K, Zhang P, Lilascharoen V, Xu Z, Lim BK, et al. Enhanced AMPA Receptor Trafficking Mediates the Anorexigenic Effect of Endogenous Glucagon-like Peptide-1 in the Paraventricular Hypothalamus. Neuron. 2017;96(4):897-909.e5.

9. Deng Y, Dickey JE, Saito K, Deng G, Singh U, Jiang J, et al. Elucidating the role of Rgs2 expression in the PVN for metabolic homeostasis in mice. Mol Metab. 2022;66:101622.

10. Souza LAC, Worker CJ, Li W, Trebak F, Watkins T, Gayban AJB, et al. (Pro)renin receptor knockdown in the paraventricular nucleus of the hypothalamus attenuates hypertension development and AT(1) receptor-mediated calcium events. Am J Physiol Heart Circ Physiol. 2019;316(6):H1389-h405.

11. Xia H, de Queiroz TM, Sriramula S, Feng Y, Johnson T, Mungrue IN, et al. Brain ACE2

overexpression reduces DOCA-salt hypertension independently of endoplasmic reticulum

stress. Am J Physiol Regul Integr Comp Physiol. 2015;308(5):R370-8.

12. Xu J, Bartolome CL, Low CS, Yi X, Chien C-H, Wang P, et al. Genetic identification of leptin

neural circuits in energy and glucose homeostases. Nature. 2018;556(7702):505-9.

13. Erben L, Welday JP, Murphy R, Buonanno A. Toxic and Phenotypic Effects of AAV_Cre

Used to Transduce Mesencephalic Dopaminergic Neurons. Int J Mol Sci. 2022;23(16).

14. Rezai Amin S, Gruszczynski C, Guiard BP, Callebert J, Launay J-M, Louis F, et al. Viral

vector-mediated Cre recombinase expression in substantia nigra induces lesions of the

nigrostriatal pathway associated with perturbations of dopamine-related behaviors and

hallmarks of programmed cell death. Journal of Neurochemistry. 2019;150(3):330-40.

15. Sanford CA, Soden ME, Baird MA, Miller SM, Schulkin J, Palmiter RD, et al. A Central

Amygdala CRF Circuit Facilitates Learning about Weak Threats. Neuron. 2017;93(1):164-78.

16. Sisley S, Gutierrez-Aguilar R, Scott M, D'Alessio DA, Sandoval DA, Seeley RJ. Neuronal

GLP1R mediates liraglutide's anorectic but not glucose-lowering effect. The Journal of Clinical

Investigation. 2014;124(6):2456-63.

17. Zhang R, Asai M, Mahoney CE, Joachim M, Shen Y, Gunner G, et al. Loss of hypothalamic

corticotropin-releasing hormone markedly reduces anxiety behaviors in mice. Molecular

Psychiatry. 2017;22(5):733-44.

18. Nath T, Mathis A, Chen AC, Patel A, Bethge M, Mathis MW. Using DeepLabCut for 3D

markerless pose estimation across species and behaviors. Nature Protocols. 2019;14(7):2152-

76.

19. Daviu N, Bruchas MR, Moghaddam B, Sandi C, Beyeler A. Neurobiological links between

stress and anxiety. Neurobiology of Stress. 2019;11:100191.

20. Sterley T-L, Baimoukhametova D, Füzesi T, Zurek AA, Daviu N, Rasiah NP, et al. Social

transmission and buffering of synaptic changes after stress. Nature Neuroscience.

2018;21(3):393-403.

21. Yuan Y, Wu W, Chen M, Cai F, Fan C, Shen W, et al. Reward Inhibits Paraventricular CRH
Neurons to Relieve Stress. Curr Biol. 2019;29(7):1243-51.e4.

22. Füzesi T, Daviu N, Wamsteeker Cusulin JI, Bonin RP, Bains JS. Hypothalamic CRH neurons
orchestrate complex behaviours after stress. Nat Commun. 2016;7:11937.

23. Semprini S, Troup TJ, Kotelevtseva N, King K, Davis JR, Mullins LJ, et al. Cryptic loxP sites
in mammalian genomes: genome-wide distribution and relevance for the efficiency of BAC/PAC
recombineering techniques. Nucleic Acids Res. 2007;35(5):1402-10.

24. Correa-da-Silva F, Kalsbeek MJ, Gadella FS, Oppersma J, Jiang W, Wolff SEC, et al.
Reduction of oxytocin-containing neurons and enhanced glymphatic activity in the hypothalamic
paraventricular nucleus of patients with type 2 diabetes mellitus. Acta Neuropathologica
Communications. 2023;11(1):107.

25. Sims JS, Lorden JF. Effect of paraventricular nucleus lesions on body weight, food intake
and insulin levels. Behavioural Brain Research. 1986;22(3):265-81.

26. Li MM, Madara JC, Steger JS, Krashes MJ, Balthasar N, Campbell JN, et al. The
Paraventricular Hypothalamus Regulates Satiety and Prevents Obesity via Two Genetically
Distinct Circuits. Neuron. 2019;102(3):653-67.e6.

27. Leibowitz SF, Hammer NJ, Chang K. Hypothalamic paraventricular nucleus lesions produce
overeating and obesity in the rat. Physiology & Behavior. 1981;27(6):1031-40.

28. Fukushima M, Tokunaga K, Lupien J, Kemnitz JW, Bray GA. Dynamic and static phases of
obesity following lesions in PVN and VMH. American Journal of Physiology-Regulatory,
Integrative and Comparative Physiology. 1987;253(3):R523-R9.

29. Gold RM, Jones AP, Sawchenko PE, Kapatos G. Paraventricular area: Critical focus of a
longitudinal neurocircuitry mediating food intake. Physiology & Behavior. 1977;18(6):1111-9.

30. Terrill SJ, Jackson CM, Greene HE, Lilly N, Maske CB, Vallejo S, et al. Role of lateral
septum glucagon-like peptide 1 receptors in food intake. Am J Physiol Regul Integr Comp
Physiol. 2016;311(1):R124-32.

31. Rizzi-Wise CA, Wang DV. Putting Together Pieces of the Lateral Septum: Multifaceted
Functions and Its Neural Pathways. eNeuro. 2021;8(6).

32. Bales MB, Centanni SW, Luchsinger JR, Fathi P, Biddinger JE, Le TDV, et al. High fat diet

blunts stress-induced hypophagia and activation of Glp1r dorsal lateral septum neurons in male

but not in female mice. Molecular Metabolism. 2022;64:101571.

33. Terrill SJ, Holt MK, Maske CB, Abrams N, Reimann F, Trapp S, et al. Endogenous GLP-1 in

lateral septum promotes satiety and suppresses motivation for food in mice. Physiology &

Behavior. 2019;206:191-9.

34. Li S-B, Borniger JC, Yamaguchi H, Hédou J, Gaudilliere B, de Lecea L. Hypothalamic

circuitry underlying stress-induced insomnia and peripheral immunosuppression. Science

Advances.6(37):eabc2590.

35. Weninger SC, Muglia LJ, Jacobson L, Majzoub JA. CRH-deficient mice have a normal

anorectic response to chronic stress. Regul Pept. 1999;84(1-3):69-74.

36. Weninger SC, Dunn AJ, Muglia LJ, Dikkes P, Miczek KA, Swiergiel AH, et al. Stress-induced

behaviors require the corticotropin-releasing hormone (CRH) receptor, but not CRH.

Proceedings of the National Academy of Sciences. 1999;96(14):8283-8.

37. Muglia L, Jacobson L, Dikkes P, Majzoub JA. Corticotropin-releasing hormone deficiency

reveals major fetal but not adult glucocorticoid need. Nature. 1995;373(6513):427-32.

38. Dunn AJ, Swiergiel AH. Behavioral responses to stress are intact in CRF-deficient mice.

Brain Research. 1999;845(1):14-20.

39. Foster MT, Song CK, Bartness TJ. Hypothalamic paraventricular nucleus lesion involvement

in the sympathetic control of lipid mobilization. Obesity (Silver Spring). 2010;18(4):682-9.

40. Chen CR, Zhong YH, Jiang S, Xu W, Xiao L, Wang Z, et al. Dysfunctions of the

paraventricular hypothalamic nucleus induce hypersomnia in mice. Elife. 2021;10.

41. Kim ER, Xu Y, Cassidy RM, Lu Y, Yang Y, Tian J, et al. Paraventricular hypothalamus

mediates diurnal rhythm of metabolism. Nat Commun. 2020;11(1):3794.

42. Burmeister MA, Ayala JE, Smouse H, Landivar-Rocha A, Brown JD, Drucker DJ, et al. The

Hypothalamic Glucagon-Like Peptide 1 Receptor Is Sufficient but Not Necessary for the

Regulation of Energy Balance and Glucose Homeostasis in Mice. Diabetes. 2017;66(2):372-84.

43. Bozadjieva-Kramer N, Ross RA, Johnson DQ, Fenselau H, Haggerty DL, Atwood B, et al.

The Role of Mediobasal Hypothalamic PACAP in the Control of Body Weight and Metabolism.

Endocrinology. 2021;162(4).

44. Lee CM, Zhou L, Liu J, Shi J, Geng Y, Liu M, et al. Single-cell RNA-seq analysis revealed

long-lasting adverse effects of tamoxifen on neurogenesis in prenatal and adult brains. Proc

Natl Acad Sci U S A. 2020;117(32):19578-89.

FIGURES AND FIGURE LEGENDS

Figure 1. AAV-Cre administration in PVN of *Crh^{f/f}* mice results in the development of

metabolic syndrome. (A) Experimental paradigm for virus delivery. AAV-Cre or AAV-eGFP was

stereotactically injected into the adult murine PVN. **(B)** Plasma corticosterone decreased following

AAV-Cre injection (t (12) = 4.011, p = 0.0017). **(C)** Changes in body weight following virus delivery

at week 0 (two-way ANOVA, main effect of Group: F (1, 12) = 294.1, p < 0.0001; main effect of

Time: F (1.378, 16.53) = 228.5, p < 0.0001; interaction between Group and Time: F (9, 108) =

138.4, p < 0.0001). **(D)** Daily chow intake following virus injection increased with AAV-Cre (t (12)

= 3.152, p = 0.0083). **(E)** Changes in oxygen consumption were observed over the course of 96

hours. **(F)** Quantification of the decrease in oxygen consumption in AAV-Cre-injected mice (two-

way ANOVA with Bonferroni's multiple comparisons test, main effect of Group: F (1, 12) = 25.11,

p = 0.0003; main effect of Time: F (1, 12) = 208.4, p < 0.0001; interaction between Group and

Time: F (1, 12) = 43.58, p < 0.001). **(G)** The respiratory exchange ratio (RER) was unaltered (two-

way ANOVA, no main effect of Group: F (1, 12) = 0.02021, p = 0.8893; no main effect of Time: F

(1, 12) = 2.970, p = 0.1105; no interaction between Group and Time: F (1, 12) = 3.566, p = 0.0834).

(H) Energy expenditure decreased following AAV-Cre delivery (two-way ANOVA with Bonferroni's

multiple comparisons test, main effect of Group: F (1, 12) = 26.31, p = 0.0002; main effect of Time:

F (1, 12) = 207.9, p < 0.0001; interaction between Group and Time: F (1, 12) = 45.10, p < 0.0001).

(I) Locomotion in the x-axis in the CLAMS cage decreased during the dark cycle (two-way ANOVA

with Bonferroni's multiple comparisons test, main effect of Group: F (1, 12) = 8.461, p = 0.0131;

main effect of Time: F (1, 12) = 41.86, p < 0.0001, interaction between Group and Time: F (1, 12)

= 10.17, p = 0.0078). **(J)** Proportion of fat mass as determined by EchoMRI increased in AAV-

Cre-injected mice (t (12) = 12.67, p < 0.0001). **(K)** Changes in average blood glucose after

overnight fasting (t (12) = 5.888, p < 0.0001). **(L)** The glucose tolerance test reveals increased

baseline blood glucose levels (two-way ANOVA with Bonferroni's multiple comparisons test, main

effect of Group: F (1, 12) = 22.30, p = 0.0005; main effect of Time: F (2.717, 32.60) = 24.72, p <

0.0001; interaction between Group and Time: F (4, 48) = 3.111, p = 0.0235). **(M)** The insulin

tolerance test reveals insulin tolerance (two-way ANOVA with Bonferroni's multiple comparisons

test, main effect of Group: F (1, 11) = 11.30, p = 0.0063; main effect of Time: F (1.712, 18.84) =

55.72, p < 0.0001; no interaction between Group and Time: F (2, 22) = 0.6622, p = 0.5257).

Sample sizes are as shown on graphs (n = 7 mice per group). Data are presented as individual

points and mean ± or + SEM.

Figure 2. AAV-Cre action in PVN of _Crh^{f/f}_ mice depletes CRH and OXT. (A) Example comparison of CRH expression in PVN between both groups. **(B)** Example of OXT expression in PVN. **(C)** CRH-positive cell density decreased following AAV-Cre expression in _Crh^{f/f}_ PVN ($t_{(4)}$ = 3.176, p = 0.0337). **(D)** OXT-positive cell density decreased as well ($t_{(4)}$ = 2.807, p = 0.0236). **(E)** Overall cell density trended toward a decline ($t_{(4)}$ = 2.511, p = 0.0660). Sample sizes are as shown on graphs (n = 3 mice per group). Data are presented as individual points and mean + SEM.

Figure 3: AAV-Cre action in wild-type PVN induces metabolic changes, neuronal cell death, and recruitment of glia. (A) Experimental paradigm for virus delivery in wild-type mice. **(B)** Plasma corticosterone levels decreased after AAV-Cre injection ($t_{(7)}$ = 3.260, p = 0.0139). **(C)** Body weight changes followed virus delivery (two-way ANOVA, main effect of Group: $F_{(1, 7)}$ = 22.37, p = 0.0021; main effect of Time: $F_{(1.134, 7.940)}$ = 191.5, p < 0.0001; interaction between Group and Time: $F_{(6, 42)}$ = 79.71, p < 0.0001). **(D)** Daily food intake increased in AAV-Cre-injected mice (Mann-Whitney test, U = 0, p = 0.0159). **(E)** Example images of neuronal nuclei (NeuN) across the PVN following injection in wild-type mice. **(F)** Quantification of neuron density at different stereotactic coordinates: -0.34 ($t_{(5)}$ = 18.72, p < 0.0001), -0.58 ($t_{(5)}$ = 5.343, p = 0.0031), -0.94 ($t_{(5)}$ = 3.053, p = 0.0283), and -1.22 ($t_{(5)}$ = 3.372, p = 0.0199). **(G)** Example comparison of glial fibrillary acidic protein (GFAP) and ionized calcium binding adaptor molecule 1 (Iba1) expression in the wild-type PVN between the two groups. **(H)** Expression of astrocytes increased as quantified by GFAP fluorescence area ($t_{(6)}$ = 3.017, p = 0.0235) **(I)** Expression of microglia increased as quantified by Iba1 fluorescence area ($t_{(6)}$ = 2.917, p = 0.0267). Sample sizes are as shown on graph: **(F)** n = 3 control, 4 AAV-Cre **(H, I)** n = 4 per group. Data are presented as individual points and mean ± or + SEM.

SUPPLEMENTARY INFORMATION

Figure S1: AAV-Cre expression in the PVN of female *Crh^{f/f}* mice recapitulates body weight increases. (A) Experimental paradigm for virus delivery. **(B)** Body weight increases following AAV-Cre injection (two-way ANOVA, main effect of Group: $F_{(1, 4)} = 59.08$, $p = 0.0015$; main effect of Time: $F_{(1.782, 7.129)} = 77.74$, $p < 0.0001$; interaction between Group and Time: $F_{(6, 24)} = 58.41$, $p < 0.0001$). Sample sizes are as shown on graphs (n = 3 mice per group). Data are presented as mean ± SEM.

Figure S2: *Crh^{f/f}* mice expressing AAV-Cre in PVN display elevated anxiety-like behaviors. (A) Representative trajectories of AAV-GFP- (top) and AAV-Cre-injected (bottom) mice during the open field test. **(B)** AAV-Cre-injected mice spent less time in the center zone of the open field test ($t_{(12)} = 2.181$, $p = 0.0498$), and changes in distance traveled were not significant ($t_{(12)} = 1.811$, $p = 0.0952$) during the open field test. **(C)** Representative trajectories of both groups during the light-dark box test. **(D)** Time spent in the dark zone increased ($t_{(12)} = 2.416$, $p = 0.0326$), and entries into the light zone decreased (Mann-Whitney test, $U = 0$, $p = 0.0006$) during the light-dark box test. Sample sizes are as shown on graphs (n = 7 mice per group). Data are presented as individual points and mean + SEM.

Figure S3. Wild-type mice injected with AAV-Cre in PVN exhibit increased anxiety-like behaviors. (A) Representative traces of wild-type mice during the open field test. **(B)** During the open field test, time spent in the center zone decreased ($t_{(7)} = 2.817$, $p = 0.0259$), and total distance traveled decreased ($t_{(7)} = 3.275$, $p = 0.0136$). **(C)** Representative movement during the light-dark box test for both groups. **(D)** During the light-dark box test, time spent in the dark zone increased ($t_{(7)} = 2.798$, $p = 0.0266$), and entries into the light zone decreased ($t_{(7)} = 3.142$, $p = 0.0163$). Sample sizes are as shown on graphs (n = 5 AAV-GFP-injected mice, n = 4 AAV-Cre-injected mice). Data are presented as individual points and mean + SEM.

Figure S4. AAV-Cre expression in the dLS of male *Glp1r^{f/f}* mice results in cell death without metabolic deficits. (A) Experimental paradigm for virus delivery. **(B)** Body weight was unchanged (two-way ANOVA, main effect of Group: $F_{(1, 16)} = 0.07282$, $p = 0.7907$; main effect of Time: $F_{(1.826, 29.22)} = 364.6$, $p < 0.0001$; interaction between Group and Time: $F_{(4, 64)} = 3.406$, $p = 0.0138$). **(C)** Representative images of neuronal nuclei in the dLS after AAV-GFP or AAV-Cre injection. **(D)** Neuron density decreased in the dLS after AAV-Cre expression ($t_{(4)} = 4.631$, $p = 0.0098$). Sample sizes are as shown on graphs: **(B)** n = 9 mice per group; **(D)** n = 3 mice per group. Data are presented as individual points and mean ± or + SEM.

Savani et al. Figure 1

Savani et al. Figure 2

Savani et al. Figure 3

Savani et al. Supplementary Figure 1

Savani et al. Supplementary Figure 3

A

AAV-Control/
AAV-Cre-eGFP

dLS

Glp1r flox/flox

B

Control (n=9)
AAV-Cre (n=9)

Body Weight (g)

0   1   2   3   4
Week

C

NeuN

Control

cc
dLS

100 μm

AAV-Cre

cc
dLS

D

NeuN+ cells/mm²

**

Control   AAV-Cre

Savani et al. Supplementary Figure 4