Title: **On the road to losing connectivity: Fecal samples provide genome-wide**

insights into anthropogenic impacts on two large herbivore species in

central India

Authors: Abhinav Tyagi*[1,2], Nidhi Yadav[1], Awadhesh Pandit[1] and Uma Ramakrishnan*[1]

Affiliation: 1. National Centre for Biological Sciences, Tata Institute of Fundamental

Research, Bangalore, India.

2. SASTRA Deemed to be University, Thanjavur, India.

* *Corresponding authors*: tyagiabhinav2008@gmail.com; uramakri@ncbs.res.in

National Centre for Biological Sciences, Tata Institute of Fundamental Research, Bellary

Road, Bangalore 560065. Phone number: +91 80 23666030/31

Short running title: Connectivity of large herbivores

Abstract:

Humans have impacted most of the planet, and the ensuing fragmentation results in small,

isolated habitat patches posing a risk of genetic diversity loss, inbreeding and genetic load.

Understanding how natural and anthropogenic landscape features affect gene flow among

habitat patches is critical for maintaining connectivity. Genome-wide data is required to

comprehend the impacts of recent fragmentation, which can be challenging when only non-

invasive samples are available. Here, we build upon advancements in conservation genomics

to address connectivity of two large herbivores, gaur (*Bos gaurus*) and sambar (*Rusa

unicolor*) in central India. Given their habitat associations, we expected these species to

respond similarly to habitat fragmentation. We used fecal-DNA and methylation-based host-

DNA enrichment with modified ddRAD protocol to generate genome-wide single nucleotide

polymorphism (SNP) data for 124 Gaur and 99 Sambar individuals. Our findings reveal that

gaur populations in central India are fragmented, displaying high genetic differentiation,

with drift significantly affecting small populations like Umred-Karhandala Wildlife Sanctuary.

Although Sambar shows low genetic structure, another small population, Bor Tiger Reserve

is genetically differentiated. Our results suggest that although landcover change and roads

restrict animal movement, the extent of this impact varies across the two species. We show

that different species, respond differently to landscape features, even with similar habitat

associations. We highlight small and isolated populations requiring urgent conservation

intervention. Such multi-species approaches enhance our understanding of cross-species

connectivity patterns. We suggest shifting from single-species to multi-species holistic

conservation approach in rapidly developing landscapes to better manage co-occurring

endangered species.

Keywords: ddRAD, Non-invasive samples, Gaur, Sambar, Wildlife genomics, Wildlife

connectivity

1. Introduction:

The ongoing wildlife crisis leading to local and global species extinctions is primarily driven by anthropogenic impacts such as habitat loss and fragmentation (Dirzo et al., 2014; Young et al., 2016). In particular, large mammals bear the brunt of vulnerability due to their life history traits, including large home ranges, migratory behaviour, slow reproductive rates, and sparse population densities (Cardillo et al., 2004). While large carnivores have received frequent spotlight as threatened keystone species, herbivores are often neglected. However, large herbivores play a critical role in shaping ecosystem structure and function by controlling wildfires, cycling nutrients, and maintaining vegetation dynamics (le Roux et al., 2020; Rouet-Leduc et al., 2021; Sandom et al., 2014). Large herbivores also make habitats suitable for carnivores and scavengers as they are the primary food source (Ripple et al., 2015).

Unfortunately, despite their ecological significance, most large herbivore species are experiencing declines in abundance and distribution due to anthropogenic pressures such as land-use change, hunting, competition, and disease transmission (Bluhm et al., 2023; Craigie et al., 2010; Ripple et al., 2015). Land-use change, such as habitat loss, fragmentation, and deforestation, restricts movement and disrupts gene flow (Crego et al., 2021; Ripple et al., 2015). The development and presence of other anthropogenic landscape features like road and railway networks can lead to increase in habitat fragmentation and reduces landscape permeability (Bennett, 2017; Hepenstrick et al., 2012). There have been several incidences of local extinction and extirpation of large herbivores from fragmented habitat patches (Sankar et al., 2013). Therefore,

maintaining gene flow and functional connectivity among these fragmented habitat

patches is critical for long-term species persistence (Thatte et al., 2018).

Most connectivity studies, especially in South Asia, have focused on a single species,

mostly charismatic carnivores (Thatte et al., 2021). Connectivity is often species-specific

and depends on species traits and ecology like body size, dispersal ability and habitat

requirement (McClure et al., 2016; Thatte et al., 2020). This is also relevant for large

herbivores as they have specific habitat associations and requirements, and can exhibit

high association with the migration routes or corridors that might not necessarily

support the movement of other species (Sawyer et al., 2019). Therefore, a multi-species

and comparative approach is essential to comprehensively understand how connectivity

varies across species. This comparative analysis, especially among species with similar

habitat associations, sheds light on the role of species ecology, beyond just habitat, in

shaping connectivity (Brodie et al., 2015; Thatte et al., 2020, 2021).

Landscape genetics integrates methods from population genetics and landscape

ecology to examine the impact of landscape features on gene flow and genetic diversity

(Storfer et al., 2007). These methods are instrumental for identifying barriers to animal

movement and have been applied to a diverse range of species (Dudaniec et al., 2016;

Fedorca et al., 2020; Schoen et al., 2022; Thatte et al., 2020; Tyagi et al., 2022). The use

of molecular markers and landscape genetic approaches aid in quantifying regional

patterns of genetic structure, providing insights into how different landscape features

influence connectivity (Epps & Keyghobadi, 2015; Holderegger & Wagner, 2008; Tyagi et

al., 2022). In many places including India, non-invasive samples, such as fecal samples,

are the only feasible means of studying large herbivores in their natural habitats.

However, the DNA quality of these samples is often compromised, as fecal DNA extracts

tend to contain exogenous DNA and PCR inhibitors (Khan & Tyagi, 2021; Kohn & Wayne,

1997; Putman, 1984; Schrader et al., 2012).

Recent developments in conservation genomics enable the use of DNA obtained

from poor-quality non-invasive samples to generate genome-wide data using next-

generation sequencing (NGS) (Chiou & Bergey, 2018; Hayward et al., 2022; Natesh et al.,

2019; Orkin et al., 2020; Tyagi et al., 2022). The use of single nucleotide polymorphism

(SNP) markers has led to better estimates of genetic diversity and population structure

(Bohling et al., 2019; Lemopoulos et al., 2019; Ling et al., 2020) and inference of

landscape features on animal movement (Tyagi et al., 2022). The use of a higher number

of SNP markers provides more power in comparison to increasing the number of

samples or individuals in quantifying fine-scale genetic variation and functional

connectivity, as evidenced from both simulations and empirical data (Landguth et al.,

2012; Tyagi et al., 2022).

Further, double-digest restriction-associated DNA (ddRAD) sequencing has emerged

as a promising approach for generating genome-wide data for non-model species.

Despite the promising potential, there is still a need to develop library preparation

methods that complement the use of poor-quality, low-concentration input DNA and

enable the generation of a comparable amounts of data per sample. Most available

protocols are not designed to deal with the specific issues related to analysing samples

with poor-quality and varied DNA concentrations (Peterson et al., 2012; Peterson et al.,

2014). To address this, we developed a modified ddRAD sequencing protocol that allows

us to generate genome-wide data, suitable for poor-quality DNA samples and minimizes

the impact of input host DNA concentration on the data generated.

In order to address gaps in our understanding of how fragmentation impacts large

herbivores, we utilized genome-wide SNPs obtained from fecal samples to investigate

connectivity patterns of two large vulnerable herbivore species, gaur (*Bos gaurus*) and

sambar (*Rusa unicolor*), in central India. Our main objectives were to (a) determine the

genetic diversity and population structure of these species, (b) assess the impact of

landscape heterogeneity on gene flow, and (c) compare the response of two species

with similar habitat dependency, but different body size and dispersal-related functional

traits, to habitat fragmentation and different landscape features. We predict that both

species may reveal similar patterns of connectivity as they share similar habitat

associations. Additionally, we aimed to identify priority areas for restoring connectivity

for these species. This study represents the first genome-wide investigation of

landscape-level connectivity for both species and underscores the importance of

studying genetic variation and connectivity among different species.

2. Materials and Methods

2.1. Study area

The central Indian landscape (CIL) is a well-studied landscape in terms of connectivity

studies in India, but the focus of these studies has largely been on carnivores,

specifically, tigers (Schoen et al., 2022; Thatte et al., 2021). It is a priority tiger

conservation landscape and is also inhabited by various other endangered wild species

(Jhala et al., 2008). Our study landscape was spread across two central Indian states—

Madhya Pradesh and Maharashtra. The forest is primarily tropical dry and moist

deciduous, harbouring rich biodiversity (Champion & Seth, 1968). Central India has

around 33% of the area under forest cover, which consists of multiple protected areas

(PAs), reserved and territorial forests (Dutta et al., 2018). A varied and heterogeneous

landscape matrix with multiple land-use types, such as agricultural fields and human

built-ups, surrounds these protected areas.

Central India, like other areas of conservation concern, faces threats including

habitat fragmentation, degradation, and loss of connectivity due to rapid

transformations driven by human activities such as transportation network expansion,

mining projects, and development (Crooks et al., 2011; Habib et al., 2016; Ibisch et al.,

2016; Schoen et al., 2022; Venter et al., 2016). Major state and national highways

bisecting the landscape further jeopardizes wildlife connectivity and integrity (Thatte et

al., 2018, 2020).

2.2. Study species and sample collection

Gaur (*Bos gaurus*) and sambar (*Rusa unicolor*) are endemic to South and Southeast

Asia and rank among the top prey species for large carnivores like tigers (Andheria et al.,

2007). Gaur is the largest wild cattle species and one of the few mega-herbivore species

that inhabit the Indian subcontinent. Although Gaur was once found throughout the

forested areas of India, its current distribution is restricted to fragmented habitat

pockets, primarily in the Western Ghats, the central Indian highlands, and Northeast

India (Musser & Koop-man, 1994).

On the other hand, Sambar is a meso-herbivore distributed across a wide range of

habitats, from tropical rainforests to subtropical mixed forests in South and Southeast

Asia. In India, it can be found from the foothills of the Himalayas in the north to the

Western Ghats in the south and the Peninsular and north-eastern regions (Timmins et

al., 2015). Although sambar exhibits a wide range, the population is very patchy,

especially in central India (Kushwaha et al., 2004; Timmins et al., 2015).

Both species have similar habitat association and prefer habitats with high tree and

shrub densities, and generally avoid anthropogenic disturbances (Sankar et al., 2013;

Singh et al., 2021). Both species face threats like habitat fragmentation, habitat loss,

illegal poaching and other anthropogenic impacts responsible for population decline and

local extirpation. Gaur has experienced several local extinctions in central India,

including from Bor Tiger Reserve, Bandhavgarh Tiger Reserve, and Kengar Valley

National Park (Sankar et al., 2013). While there are no published reports of local

extinctions for Sambar, the lack of monitoring and attention towards large herbivores

raises concerns about its status in many areas.

To assess the genetic connectivity of the two species, we collected fecal samples

during two field seasons (October 2017 to February 2018 and December 2020 to March

2021) from inside and outside six protected areas: 1) Kanha Tiger Reserve (KTR), 2)

Pench Tiger Reserve (PTR), 3) Nagzira-Nawagaon Tiger Reserve (NNTR), 4) Bor Tiger

Reserve (BOR), 5) Tadoba-Andhari Tiger Reserve (TATR), and 6) Umred-Karhandla

Wildlife Sanctuary (UMR). Outside protected area sampling was done in forested areas

like the Kanha-Pench corridor (KPC) (Figure 1).

We sampled in areas with frequent use by study species to maximize the collection

of fresh samples (less than 3 days old). Fresh dung (for gaur) and pellets (for sambar)

were sampled using a sterilized swab and stored in Longmire's buffer in 2 ml

microcentrifuge tubes (Longmire et al., 1997). Each sample was collected in duplicate,

and we targeted a large sample size to compensate for a potential loss of samples due to

the low quality and quantity of DNA. To maximize the area covered and avoid recaptures

of individuals, each forest beat (i.e., administrative divisions) was sampled only once, and

adjoining beats were sampled simultaneously. We recorded GPS location, species

identity based on morphology, and a visually estimated sample age for all samples.

2.3. DNA extraction, species identification and quantification

All samples were extracted using QIAamp DNA Blood Mini Kit (Qiagen) following the

manufacturer protocol. Species identification was done for all the samples using species-

specific primers designed in this study. The gaur (Bg_cytb1.1F *5'-*

CACAGCAATTGCCATAGTCC-3' & Bg_cytb1.1R *5'-AAGGGCTAGAATTAGTAACAAGGTT-3'*)

and sambar (CommF_CytB1 *5'-ATTGTAAACAACGCATTCAT-3'* & SambarR_CytB1 *5'-*

GGAATAGGCCTGTGATG-3') specific primers amplified 151 and 115 base pair sequences,

respectively, of mitochondrial cytochrome b region (Table S1 and Supplementary

material S1). No cross-amplification, with any of the other sympatric species with which

the fecal samples could be misidentified based on morphology, was observed.

Therefore, any positive amplification observed using fecal DNA corresponds to

conclusive species identity (Supplementary material S1).

Total DNA concentrations were measured post species identification, using Qubit

(Invitrogen). Host DNA (hDNA) was estimated using quantitative PCR (qPCR) (Morin et

al., 2001). All qPCR reactions were performed in triplicates along with positive and

negative controls and evaluated against the standard curve constructed using serial

dilutions of known tissue DNA.

2.4. DNA enrichment

Host DNA enrichments were performed using the NEBNext Microbiome DNA

enrichment kit (Feehery et al., 2013). The samples and MBD2-Fc magnetic beads were

prepared following the procedure described in Tyagi et al., (2022). To maximize the

hDNA capture, the DNA extracts were concentrated using Cenrivap DNA vacuum

concentrator (Labconco). Total and hDNA were also quantified post-enrichment.

2.5. Library preparation and ddRAD sequencing

We prepared ddRAD libraries using the enriched fecal DNA samples. Our protocol

was based on modifications of existing methods (Peterson et al., 2012; Peterson et al.,

2014; White et al., 2018) and tailored to our sample types. We digested all samples with

a combination of Sph1 and MluC1 restriction enzymes and performed adapter ligation,

purification using AMPure XP magnetic beads (Agencourt), and indexing. To reduce PCR

bias, we performed indexing PCR in four replicates and pooled all replicates before

conducting dual-size selection using AMPure XP magnetic beads. We quantified and

assessed the size distribution of individual sample libraries using Qubit and Tapestation

(Agilent Technologies), respectively, before pooling them in equimolar concentration.

The final pooled library was also quantified and checked for size distribution before

sequencing on the HiSeq2500 platform using PE 2X100 V2 kit. Please see supplementary

material S2 for detailed protocol, adapter, index and, modified read 1 and read 2

sequences (Table S3 and S4).

2.6. Quality control, SNP calling and filtering

Raw reads were first demultiplexed and then FastQC was used for quality checking

(http://www. bioinformatics.babraham.ac.uk/projects/fastqc/). The reads were trimmed

using TrimGalore (v.0.5.0 Babraham Bioinformatics) and mapped to respective reference

genomes using the BWA mem algorithm (Li, 2013). Cow reference genome (ARS-UCD1.2;

RefSeq accession: GCF_002263795.1) was used to align gaur data and red deer

reference assembly (mCerEla1.1; RefSeq accession: GCF_910594005.1) was used for

mapping sambar reads. We selected these reference genomes for mapping as they

represented the best available reference assemblies for the species most closely related

to our study species (Mackiewicz et al., 2022; Wu et al., 2018). Post mapping, read-

group information was added to each BAM file using Picard tools

(http://broadinstitute.github.io/ picard).

Variant calling was done using Freebayes (Garrison & Marth, 2012) and vcftools

(Danecek et al., 2011) was used to filter the sites based on quality (≥10) and minor allele

count (≥3). Sites supported by three or more reads were considered for identifying

genotypes (Arafa et al., 2017; Chang et al., 2023; Chen et al., 2021; Shirasawa et al.,

2016). Only biallelic variants were retained, and sites that were present in less than 60%

of the individuals were discarded. Samples with more than 40% missing data were

removed from further analysis.

2.7. Individual identification, isolation by distance and outlier loci detection

We generated relatedness estimates using PLINK (Purcell et al., 2007) to find

recaptures and related individuals. For testing Isolation by distance (IBD), we calculated

pairwise genetic distance based on the proportion of shared alleles (D_{PS}=1-proportion of

shared alleles) and Euclidian distance using *adegenet* package (Jombart, 2008) in R

version 4.1.2. We tested for correlations between genetic and geographic distances

using Mantel tests using package *adegenet* (Jombart, 2008) with randomization with 999

replicates. Genalex 6.503 (Peakall & Smouse, 2012) was used to calculate and plot a

multi-locus spatial correlogram using genetic and geographic distances using 1000

bootstraps, and 1000 permutations of the data were used to evaluate significance. We

also used Bayescan software (Foll & Gaggiotti, 2008) to detect and remove loci under

selection, using a false discovery rate of 5%.

2.8. Genetic differentiation and population structure

Expected and observed heterozygosity was computed using R packages *adegenet*

(Jombart, 2008) and *Hierfstat* (Goudet, 2005). We used multiple approaches to

understand genetic differentiation comprehensively. We assessed each species' genetic

structure using discriminant analysis of principal components (DAPC) and STRUCTURE

(Pritchard et al., 2000). DAPC (Jombart et al., 2010) was conducted using the *adegenet*

package in r version 4.1.2. The number of principal components (PC) axes and

discriminant functions (DFs) to be retained were determined using the optim.a.score

function. We also used the program STRUCTURE (Pritchard et al., 2000) as a Bayesian

approach to determine the most likely number of genetic clusters (K). STRUCTURE was

run for 200,000 Markov Chain Monte Carlo (MCMC) iterations after a burn-in of 50,000

iterations, with ten replicates for K ranging from one to eight, using the correlated allele

frequencies and admixture model. The best K was selected by likelihood and delta K

values and methods described by Evanno et al., (2005). The results were analysed and

plotted using a web-based server, CLUMPAK (Kopelman et al., 2015).

2.9. Landscape genetic analysis

2.9.1. Landscape variables

To build resistance models in this study, we used four landscape variables: land use

land cover (LULC), roads with traffic, human population density (Hpop), and density of

linear features (Linden). These variables have been previously used to understand the

connectivity of various species in the same landscape (Thatte et al., 2020; Tyagi et al.,

2022). Each layer was reclassified and ranked in order of the resistance offered.

Landcover data was reclassified into four categories: forest, scrub and degraded forest,

agriculture, and built-up areas. The road layer was reclassified into seven different

classes based on traffic intensity (measured in Passenger Car Units (PCU)) and road

width (e.g., 1,2,4, or 6 lanes). To calculate the density of linear features, we combined

roads (without traffic), railway lines, and irrigation canals into a single polylines layer.

Roads and LULC were discrete variables, whereas human population density and density

of linear features were continuous landscape variables used to generate the resistance

surfaces. More details can be found in Thatte et al., (2020) and Tyagi et al., (2022).

Previous studies have highlighted that the spatial scale of a landscape variable can affect

the species–landscape relationship (Jackson & Fahrig, 2015). To address this issue, we

analysed the landscape variables at seven different resolutions (0.25, 0.5, 1, 2.5, 5, 10,

25 km) for continuous variables (Hpop and Linden) and one categorical variable (LULC).

While optimising the resistance landscape, a buffer of 20 km around all sample points

was also incorporated.

2.9.2. Multi-model optimization

To evaluate how landscape variables affect animal movement, we utilized a multi-

model optimization approach to determine the maximum resistance offered (Rmax) and

the relationship (x) between each variable and resistance. Genetic distance based on the

proportion of shared alleles, D_{PS} (Bowcock et al., 1994), was used as the response

variable. We used different parameter values for Rmax and x to identify the best

combination that explained the genetic data. Seven values of maximum resistance

(Rmax= 2, 5, 10, 50, 100, 500, 1000) and shape parameter (x = 0.01, 0.1, 0.5, 1, 2, 5, 10)

were used. We calculated the cost distance for each pair of individuals for each

combination of Rmax and x and utilized the MLPE.lmm() function from the *ResistanceGA*

package (Peterman, 2018) to fit Linear mixed-effects (LME) models with maximum

likelihood population effects (MLPE) parameterization. We identified the best model for

each landscape variable based on the lowest AIC score from the MLPE mixed effects

model (Shirk et al., 2010). In univariate optimization, we determined the scale and

combination of parameters (Rmax and x) of the best-supported model for each

landscape variable.

Best-fitting univariate models were combined additively and further optimized in a

multivariate framework. We individually varied the model parameters (Rmax and x) for

each landscape variable while holding the other variables' parameter values constant

(Thatte et al., 2020; Tyagi et al., 2022). If the optimal parameters changed for a

landscape variable in the multivariate context, we held the new model parameters

constant and varied the parameters of the next variable. We repeated this process until

the parameterization of all the variables stabilized. This multivariate optimization

approach enabled us to estimate resistance offered by each variable in combination

with others for both species (Shirk et al., 2010).

2.10. Bootstrapping of landscape analysis

We used the resist.boot() function for bootstrapping within the *ResistanceGA*

package (Peterman, 2018), with 10,000 iterations. Bootstrapping was performed using

all four univariate optimization models, all the best multivariate combination models,

and a null model for both species (Tables 3 and 4).

3. Results

3.1. Data summary

A total of 1,144 gaur and 756 sambar fecal samples (based on morphology) were

collected from the field, and post-genetic species identity, 921 samples were identified

as gaur, while 504 samples were determined to be of sambar. Subsequently, all

identified samples underwent hDNA quantification before being selected for MBD

enrichment. We selected 356 gaur and 288 sambar samples for MBD enrichment. After

enrichment, the samples were quantified again, and those yielding a hDNA

concentration of more than 0.06 ng/μl were selected for the preparation of ddRAD

libraries (Tyagi et al., 2022). 171 gaur samples and 120 sambar samples were chosen for

library preparation and sequencing, along with positive controls, extraction controls,

PCR and library prep controls. The landscape and individual locations are shown in

Figure 1.

3.2. Sequencing and SNP calling

Post demultiplexing and quality filtering, the average number of reads obtained per

sample was 10.21 million for gaur (n=171) and 6.98 million for sambar (n=120). The

higher number of reads in gaur samples was due to the data being sequenced twice and

subsequently merged per sample (Fig S1). Mapping to the reference genome showed a

higher percentage in sambar (average of 44.56%) compared to gaur (average of 32.63%)

(Fig S1 and S2). Sample-wise detailed breakdown can be found in supplementary Tables

S5 and S6.

The raw VCF file for gaur contained 30,47,450 variant sites, while for sambar, the

number of variant sites in the raw VCF was 41,19,472. Post filtering, we excluded 47

gaur and 21 sambar samples, respectively due to high missing data. Individual

identification was performed using relatedness values, and no recaptures were found in

our dataset (Fig S3 and S4). The final putatively neutral dataset included 124 gaur

individuals genotyped at 1186 SNPs and 2909 SNPs for 99 sambar individuals.

3.3. Genetic diversity, Isolation by distance and spatial autocorrelation

The genetic diversity of gaur was found to be low, with observed heterozygosity (Ho)

of 0.129 and expected heterozygosity (He) of 0.149. In comparison, sambar also

exhibited an overall low genetic diversity (Ho = 0.191; He = 0.22), although higher than

that of gaur. Population-wise heterozygosity did not vary significantly for gaur, while

there was variation observed for sambar (Table 1). Notably, both species exhibited a

significant positive correlation between genetic and Euclidean distance, according to

Mantel's test (Gaur: Mantel's R = 0.112, p-value = 0.004; Sambar: Mantel's R = 0.117, p-

value = 0.001; see Fig 2c and 3c). Spatial autocorrelation analysis revealed a significant

positive spatial correlation up to 65 km and 30 km for gaur and sambar respectively (Fig

2d and 3d).

3.4. Population genetic structure

Discriminant Analysis of Principal Components (DAPC) was conducted using the optimal number of PCs. For gaur, 14 PCs were retained based on the optim.a.score, revealing differentiation between Umred Karhandala Wildlife Sanctuary (UMR) and all other populations. Tadoba-Andhari Tiger Reserve (TATR) and Nagzira Nawegaon Tiger Reserve (NNTR) also exhibited differentiation (Fig 2a). STRUCTURE analyses indicated that the optimal number of genetic clusters for gaur was K = 3 (Fig 2b and S7). At K = 2, NNTR and UMR were segregated from the remaining populations, while at K = 3, UMR and NNTR formed separate clusters. At K = 4, UMR, NNTR, and TATR formed distinct clusters, while KTR, KPC, and PTR clustered together, displaying admixture (Fig 2b). Pairwise Fst also supported the results obtained from both DAPC and STRUCTURE analyses (Fig 2e). The highest pairwise Fst value was observed between UMR and TATR (0.134), whereas the lowest was found between KTR and KPC (0.019). UMR exhibited the highest pairwise Fst with all other populations, suggesting significant genetic differentiation and limited gene flow (Fig 2e).

In the case of sambar, a low level of genetic differentiation was observed. DAPC analysis using 22 PCs revealed overlap among all populations (Fig 3a). The most supported K value in STRUCTURE analysis was K = 2 (Evanno's method, Fig 3b and S7). Pairwise Fst analysis also indicated low differentiation among sambar populations in central India (Fig 3e). The highest pairwise Fst value was observed between BOR and KTR (0.056), while the lowest was between NNTR and TATR (0.009).

3.5. Landscape connectivity and resistance analysis

Land use land cover (LULC) and roads with traffic were identified as the most influential factors impacting the genetic connectivity of both species. However, the resistance offered by different land use classes and traffic intensities were found to be different, as were the strengths of the correlation.

The impact of landscape modification on gaur movement was more pronounced than sambar, as indicated by the higher model fit (Table 2). In univariate analysis, all four landscape variables impact connectivity. Univariate analysis identified a spatial resolution of 25 km for Linden and Hpop, whereas LULC had the highest impact at a scale of 1 km (Table S2). LULC, roads with traffic, and the density of linear features exhibited the greatest resistance to gaur movement based on our multivariate analysis. Human population density did not contribute significantly to observed resistance, based on the best-fit model of multivariate analysis (Table 2; Fig 4). Moderate to high traffic roads offer high resistance to gaur movement. Areas with high linear feature density offered maximum resistance. Resistance increased as land cover changed to agriculture and human habitation (Fig S5). The selected model exhibited a marginal R^2 of 0.503, explaining around 50% of the variation in the data (Table 2).

Univariate analysis for sambar (when landscape variables were tested individually) revealed that roads, human population density, and linear infrastructure density impacted connectivity, followed by land cover (Table S2). Specifically, roads strongly influenced sambar movement (x = 0.5; Table S2). Resistance demonstrated a linear

increase with higher human population density (x = 1), and the shape parameter for

linear feature density indicated that areas with high density offered greater resistance

(Fig S6). Univariate analysis identified a spatial resolution of 25 km as optimal for all

three layers (LULC, Hpop, Linden), suggesting their significance at a broader scale for

sambar movement and connectivity. After combining the layers based on univariate

optimization, the maximum resistance offered by the variables changed. In the final

multivariate model, LULC emerged as the most influential variable for explaining

connectivity, followed by roads with traffic and human population density (Table 2; Fig 4

& S5). However, the model fit was relatively low (mR^2 = 0.173). Final resistance maps for

both species were created using QGIS (Version 3.22.3) by employing optimized

resistance values (Fig 5).

Bootstrapping analysis for gaur, with best models for all the possible combinations

(n=15) along with the null model (Euclidian distance), indicated that the model

consisting of Roads, LULC, and Linden achieved the highest mean log-likelihood score

and had a selection probability of approximately 65% (Table 3). For sambar, the

selection probability of the top model (including LULC, Roads, and Hpop) was 38%,

followed by models incorporating Hpop and LULC, and one including Roads and Hpop

(Table 4).

4. Discussion

Despite over two decades of research on connectivity, little is known about how

large herbivores respond to various landscape features (Thatte et al., 2021). Addressing

this gap, our study investigated the population structure and functional connectivity of

gaur and sambar in central India. Using a customized ddRAD sequencing protocol for

poor-quality DNA, we provide novel insights into genetic structure, impact of landscape

features, and comparative connectivity of the two species.

Through a comprehensive analysis of thousands of genome-wide SNPs, we reveal

distinct population structure for each species. Our landscape genetics analyses

demonstrated that land use and linear infrastructure like roads hinder animal

movement, highlighting differential species responses to landscape features. We identify

small, isolated populations of both species in the landscape, and recommend that these

need targeted conservation and management attention. Furthermore, our findings shed

light on species-specific interactions with landscape elements, potentially influenced by

life history traits and current population attributes, shaping spatial genetic structures.

Such comparative datasets are important for understanding differential species

response enabling holistic landscape-level conservation planning. Critically, these

insights required the implementation of novel protocol optimizations.

4.1. ddRAD using fecal samples

We collected a substantial number of fecal samples for both species to mitigate loss

due to poor DNA quality and quantity (Khan & Tyagi, 2021). Working with fecal DNA of

herbivores can be challenging due to plant metabolites acting as PCR inhibitors,

potentially limiting downstream processing for NGS data generation (Schrader et al.,

2012). While we also observed sample loss at the enrichment stage, our ddRAD library

preparation protocol minimized loss post-sequencing (due to insufficient data

generation or missing data), an advancement over already available methods from

previous studies (Tyagi et al., 2022).

465   Our protocol is robust and cost-effective. With the given protocol and designed

indexes, 468 samples can be pooled and sequenced on any Illumina platform (in our

case HiSeq2500, but can also be sequenced on NextSeq or NovaSeq). Most of the

available ddRAD protocols are designed to use around 200 ng of DNA as starting

material (Magbanua et al., 2023; Peterson et al., 2012; Peterson et al., 2014). We used

as little as 5 ng of enriched host DNA for data generation. Next-generation sequencing

has allowed creative approaches to access genome-wide or genotyping data from a wide

array of non-invasive samples (including fecal samples). Such approaches include SNP

panels, capture baits, and enrichment methods, among others (Chiou & Bergey, 2018;

Hayward et al., 2022; Khan et al., 2020; Natesh et al., 2019; Orkin et al., 2020; Snyder-

Mackler et al., 2016; Tyagi et al., 2022). Our study adds to the growing body of literature

enhancing conservation genomics application for wildlife studies and significantly

improving applicability, cost and DNA concentration requirements.

4.2. Population genetic structure

480   Our study spotlights low genetic diversity in gaur and sambar populations,

underscoring habitat degradation and fragmentation threats. We document low

diversity and high population differentiation among gaur populations in central India,

which aligns with previous findings from another protected area in the same region

using mitochondrial and microsatellite data (Atkulwar et al., 2020; Farah et al., 2021).

Our results highlight that the small population of Umred-Karhandala Wildlife Sanctuary

(UMR) is the most genetically differentiated, despite being geographically central to the

three large protected areas (Fig 1 and Fig 2a & b). This pattern could be driven by

isolation and genetic drift, as this is the smallest population among the sampled

populations. In STRUCTURE analysis, at higher K values of 3 and 4, populations like NNTR

and TATR also separate into distinct clusters (Fig 2b). This was unlike KTR, KPC and PTR,

where we always observed admixture. This could result from a well-maintained corridor

(KPC) between these two protected areas (KTR and PTR). Many studies have highlighted

the importance and functionality of this corridor using data from different species like

tiger (Schoen et al., 2022; Thatte et al., 2018), leopard (Dutta et al., 2013) and sloth bear

(Dutta et al., 2015). Our study highlights the significance and functionality of existing

corridors in maintaining connectivity for large herbivores like gaur.

However, sambar populations exhibit low differentiation and genetic diversity

(relatively higher as compared to gaur). This contrasts with our expectations based on

the species' similar habitat requirements. A previous study on sambar in the western

Himalayas, also reported similar genetic diversity using mitochondrial and microsatellite

datasets (Singh et al., 2021). The population in Bor Tiger Reserve, which is close to cities

and separated by national highways (Fig 1), displays the highest Fst value and the least

genetic diversity (Fig 3b & 3e; Table 1), highlighting limited animal movement. Overall,

low genetic differentiation in sambar might suggest that this species can utilize a variety

of land covers to disperse across the landscape.

Both species showed a weak but significant correlation between genetic distance

and geographical distance (Fig 2c and 3c). The spatial autocorrelation relationship broke

down at distances beyond ~65 km for gaur and ~30 km for sambar, which is in

agreement with dispersal abilities of these species. The weak correlation indicated that

isolation-by-distance has a weak effect, and other factors might be more important in

shaping the population structure. For example, the geographic distance between KTR

and TATR is around 300 km, while that between UMR and TATR is around 100 km, but

pairwise Fst is higher for the later pair of populations (0.065 versus 0.134 respectively).

This highlights dispersal limitation caused by landscape factors other than geographic

distance, also documented in various other species (Rengefors et al., 2021; Thatte et al.,

2020).

4.3. Impact of anthropogenic disturbance on functional connectivity

Land use land cover (LULC) and roads with traffic emerged as key factors influencing

genetic connectivity for both species. While agriculture and human habitation greatly

impacted gaur movement, agriculture had a variable impact on sambar gene flow. This

aligns with connectivity trends observed in other large mammals like tiger, leopard

(Thatte et al., 2020) and mule deer (Fraser et al., 2019). High density of linear features

(roads, railway tracks, canals) had high resistance for gaur movement, known to

influence gene flow of other species like roe deer (Coulon et al., 2006), and can

potentially lead to genetic differentiation (Thatte et al., 2020; Weckworth et al., 2013).

Numerous studies have shown that road networks affect the connectivity and hinder

gene flow of various wildlife populations (Epps et al., 2005; Ernest et al., 2014; Fraser et

al., 2019; Laurance & Balmford, 2013; Thatte et al., 2018, 2020; Tyagi et al., 2022). Our

findings reveal varying impacts of road networks and traffic intensities on our study

species. Medium to high traffic roads significantly affect gaur movement in the

landscape, with resistance increasing non-linearly (x=2), and was highest with high

traffic. In contrast, sambar experiences lower resistance on small and low-traffic roads

compared to gaur. Roads seem to impact large herbivores (gaur and sambar) more

severely than large carnivores like tigers (Thatte et al., 2020), which are primarily

affected by high-traffic roads rather than low to mid-traffic intensities. India has the

second largest road network globally and central India has a complex road network with

varying traffic intensities. Our results indicate that while these networks may not act as

absolute barriers to movement for species like tigers, they have a significant negative

impact on herbivores like gaur, among others (suggested in Thatte et al., 2020). This

emphasizes the need for wildlife connectivity mitigation measures on smaller roads in

addition to major state and national highways.

Human population density was found to have a low impact on gaur movement in our

analysis (maybe due to forest-restricted habitat association) but impacts sambar

movement significantly in the landscape. The magnitude of resistance offered was not as

high as LULC, but was similar to the road networks (Rmax = 100). Our findings using

genome-wide data resonate with a previous study in the same landscape (Jayadevan et

al., 2020), which suggested that human land use, population density, and linear

infrastructure could hinder both species' movement. Our study provides a more detailed

analysis and empirical evidence supporting these landscape responses.

The univariate analyses indicated differing optimal scales for the same landscape

variable between the two species. The optimal LULC scale for gaur was 1 km, while for

sambar, it was 25 km. In contrast, both species were impacted by human population

density and linear feature density at a 25 km scale. This distinct response highlights

gaur's susceptibility to minor habitat modifications and landscape changes, whereas

sambar appears more resilient to small-scale alterations in land use patterns.

Our multivariate optimization approach offered comprehensive insight into the

combined impacts of multiple landscape variables on genetic connectivity. Landscape

genetic analysis is particularly reliable for species or populations with significant genetic

differentiation and when landscape variables strongly affect gene flow (Shirk et al.,

2017, 2018). In central India, gaur populations exhibit high differentiation, with our

analysis demonstrating a significant role of landscape in shaping genetic structure. The

model fit was strong, with an mR^2 of 0.503. Bootstrapping analysis supports the

robustness of our findings, confirming the best-fit model's selection probability at

approximately 65%.

However, for sambar, the genetic differentiation is low, possibly due to a large

effective population size and wide distribution, mitigating detectable genetic structure

(Whitlock & McCauley, 1999). Potential past translocations may also contribute to

reduced genetic structure. Our landscape genetics results imply lower landscape impact

on sambar movement. The model fit for sambar was lower (mR^2 = 0.173), with

bootstrapping findings suggesting selection probabilities for multiple models instead of a

single best-fit model (Table 2 & 4; Fig 4). After 10,000 iterations, the top model had a

selection probability of around 38% (Table 4). Lower model fit may not always imply lack

of impact of landscape on connectivity, but also the inability of the methods to detect

those. More samples from the broader geographic range or an even higher number of

SNPs might improve landscape genetic inferences for sambar. This underscores the need

to develop novel methods for assessing landscape feature impact on species with low-

genetic-differentiation. There are a few other caveats worth mentioning here, including

the need for good quality reference genomes for effective conservation of endangered

species (Paez et al., 2022). In our study, we utilized the reference genomes of closely

related species due to the unavailability of high-quality reference assembly for both of

our study species. Another crucial point for future studies adopting our approach is the

quality and quantity of field samples. We recommend collecting fresh, and a higher

number of samples to mitigate potential losses during enrichment, library preparation,

and filtering.

4.4. Conservation implications

This study presents the first landscape-wide connectivity assessment for large

herbivores using genomic tools in India. Our methods and models can be applied to

other data-deficient endangered species in other landscapes. Urban areas and road

networks are expected to grow substantially by 2030 (Seto et al., 2012), and human

populations are also expected to increase, especially in biodiversity hotspots (Cincotta et

al., 2000). The combined effects of human demographic growth, land use change, and

climate change pose a serious threat to wildlife connectivity globally (Sanderson et al.,

2019). In India, infrastructure development like the expansion of cities, construction of

new roads and expansion of existing highways (http://morth.nic.in/BharatMala), railway

networks and interlinking river projects, and construction of canals and dams, are

prominent threats to wild species and their habitats. Despite the need for development

to meet India's economic goals, it is imperative to align development with conservation

objectives. While the Wildlife Protection Act of India restricts major developments in

and around Tiger Reserves (which are designated as eco-sensitive zones), the provisions

for construction and implementation of mitigation measures on roads and railway lines,

which bisect crucial wildlife movement areas, are limited, partly due to lack of data on

animal movement. Our results tried to bridge this gap by providing information on how

different species with similar habitat associations can respond to various landscape

alterations.

Central India is a priority conservation landscape and stronghold of multiple

endangered species (Jhala et al., 2011), but is exposed to anthropogenic pressures like

land use change, mining, and road network expansion. Along with this wildlife also faces

pressures like human-wildlife conflict, tourism pressure, risk of hybridization and disease

transfer from domestic species (Miller et al., 2015; Srivathsa et al., 2019; Tyagi et al.,

2019, 2023). While most connectivity focused on single charismatic species like tigers

(Schoen et al., 2022; Thatte et al., 2018; Yumnam et al., 2014), we highlight that it is

crucial to consider the connectivity requirements of different species in the region, as

they exhibit varied responses to landscape features based on their habitat associations

and requirements. This could eventually aid in the conservation and maintenance of

populations for top carnivores such as tigers. Carnivore populations might not solely rely

on their connectivity but may also depend on connected, viable populations of wild prey

species. In the future, detailed understanding of priority species coupled with broader

scale attributes (e.g., threats and habitats) will help with on-ground landscape

prioritization and species management, and aid in conservation planning.

In conclusion, our study contributes to the growing body of literature on landscape

genetics and the conservation of large herbivores in the face of anthropogenic pressures

using cutting-edge genomic and landscape genetic methods. We identified common and

different responses of two study species to landscape elements. Our work identified

priority areas, including smaller populations like UMR and BOR, vital for gene flow and

restoration, which could help secure the long-term survival of these species. Our work

lays the groundwork for holistic conservation and prioritization strategies by combining

genetic data with landscape resistance. Our results offer valuable insights for wildlife

managers and policymakers, to mitigate the impacts of habitat loss, fragmentation, and

impending development on large mammals, to enhance biodiversity management.

Author contributions

Conceptualization: A.T., and U.R., Data collection: A.T., Laboratory work: A.T., N.Y., and

A.P., Data analyses: A.T., Project administration: A.T., N.Y., and U.R., Data Curation: A.T.,

N.Y., and A.P., Supervision: A.T. and U.R., Funding acquisition: A.T. and U.R., Writing -

Original Draft: A.T., Writing - Review & Editing: A.T., N.Y., A.P., and U.R.

Acknowledgements

We thank the forest departments of Maharashtra (No: Desk-22(8)/CR- 12(17-

18)/928/2017-2018, dated 29/6/2017 and Desk-22(8)/WL/research/ CR-12(17 -18)/2660

/20-21/2nd February 2021) and Madhya Pradesh (S.No./Technical-I/5104, dated:

27/07/2017 and S.No./M.CH-II/Research/7130, dated: 26/11/2020) for their support in

granting the necessary permits and logistics during the sample collection. We would also

like to thank the dedicated forest department staff and our field assistants for their

steady assistance throughout the sample collection phase. We extend our thanks to all

the remarkable volunteers, namely Varun Taneja, Basawaraj Mulage, Tanvi Gurjar,

Himani Khangwal, Sourav Mehra, Hrishikesh Wagh, Varad Bansod, Shardul Joshi,

Purnima Mohite, Abhilasha Fulzele, Abhishek Shukla, Srishti Manna, Neha More, Ashish

Thoke, Mrunali Rout, Akshay Narnaware, and Saurav Jagtap for their immense efforts

and support during the fieldwork, which was instrumental for the sample collection. We

thank Ankit Kacchwaha from The Corbett Foundation for his invaluable help in

organizing logistics and providing support throughout the fieldwork. We also thank Tejali

Naik and Gokul Nair for their invaluable contributions during the data generation. We

thank Abishek Harihar and Divya Vasudev for providing crucial comments on the

manuscript. We would like to acknowledge the Wildlife Conservation Trust (WCT) for

generously providing a small grant to A.T., which not only supported the fieldwork but

also aided in preliminary laboratory analysis, including DNA extractions. Additionally, we

acknowledge The Defries Bajpai Foundation (DBF) for awarding a small grant to A.T.,

which covered the second field season. For institutional support, we thank National

Centre for Biological Sciences (NCBS). Furthermore, we would like to acknowledge the

Next Generation Genomics Facility (NGGF, Bangalore Life Science Cluster, BLiSC) for

assistance during data generation. NCBS data cluster used is supported under project

no. 12-R&D-TFR-5.04-0900, Department of Atomic Energy, Government of India. AT was

supported by NCBS and the Council for Industrial and Scientific Research-University

Grant Commission fellowship.

Conflict of interest statement

The authors declare no conflict of interest.

Data availability statement

Raw demultiplexed sequence data for both the species, generated in this study has

been made available on NCBI Sequence Read Archive (SRA) database and can be found

under the BioProject accession numbers PRJNA1121501 and PRJNA1121523 for gaur and

sambar respectively.

ORCHID

Abhinav Tyagi: orcid.org/0000-0002-5370-7056

Uma Ramakrishnan: orcid.org/0000-0002-5370-5966

References

Andheria, A. P., Karanth, K. U., & Kumar, N. S. (2007). Diet and prey profiles of three sympatric large

carnivores in Bandipur Tiger Reserve, India. *Journal of Zoology*, *273*(2), 169–175.

https://doi.org/10.1111/j.1469-7998.2007.00310.x

Arafa, R. A., Rakha, M. T., Soliman, N. E. K., Moussa, O. M., Kamel, S. M., & Shirasawa, K. (2017).

Rapid identification of candidate genes for resistance to tomato late blight disease using

next-generation sequencing technologies. *PLOS ONE*, *12*(12), e0189951.

https://doi.org/10.1371/journal.pone.0189951

Atkulwar, A., Farah, S., Gadhikar, Y., & Baig, M. (2020). Mitochondrial DNA diversity in wild gaur (Bos

gaurus gaurus): Evidence from extant and historical samples. *Mitochondrial DNA Part B*, *5*(2),

1556–1560. https://doi.org/10.1080/23802359.2020.1742589

Bennett, V. J. (2017). Effects of Road Density and Pattern on the Conservation of Species and

Biodiversity. *Current Landscape Ecology Reports*, *2*(1), 1–11.

https://doi.org/10.1007/s40823-017-0020-6

Bluhm, H., Diserens, T. A., Engleder, T., Heising, K., Heurich, M., Janík, T., Jirků, M., Klich, D., König, H.

J., Kowalczyk, R., Kuijper, D., Maślanko, W., Michler, F.-U., Neumann, W., Oeser, J., Olech,

W., Perzanowski, K., Ratkiewicz, M., Romportl, D., … Kuemmerle, T. (2023). Widespread

habitat for Europe's largest herbivores, but poor connectivity limits recolonization. *Diversity*

and Distributions, *29*(3), 423–437. https://doi.org/10.1111/ddi.13671

Bohling, J., Small, M., Von Bargen, J., Louden, A., & DeHaan, P. (2019). Comparing inferences derived

from microsatellite and RADseq datasets: A case study involving threatened bull trout.

Conservation Genetics, *20*(2), 329–342. https://doi.org/10.1007/s10592-018-1134-z

Bowcock, A. M., Ruiz-Linares, A., Tomfohrde, J., Minch, E., Kidd, J. R., & Cavalli-Sforza, L. L. (1994).

High resolution of human evolutionary trees with polymorphic microsatellites. *Nature*,

368(6470), 455–457. https://doi.org/10.1038/368455a0

Brodie, J. F., Giordano, A. J., Dickson, B., Hebblewhite, M., Bernard, H., Mohd-Azlan, J., Anderson, J.,

& Ambu, L. (2015). Evaluating multispecies landscape connectivity in a threatened tropical

mammal community. *Conservation Biology: The Journal of the Society for Conservation*

Biology, *29*(1), 122–132. https://doi.org/10.1111/cobi.12337

Cardillo, M., Purvis, A., Sechrest, W., Gittleman, J. L., Bielby, J., & Mace, G. M. (2004). Human

Population Density and Extinction Risk in the World's Carnivores. *PLOS Biology*, *2*(7), e197.

https://doi.org/10.1371/journal.pbio.0020197

Champion, S. H. G., & Seth, S. K. (1968). A revised survey of the forest types of India. *A Revised*

Survey of the Forest Types of India.

https://www.cabdirect.org/cabdirect/abstract/19720603193

Chang, J.-T., Nakamura, K., Chao, C.-T., Luo, M.-X., & Liao, P.-C. (2023). Ghost introgression facilitates

genomic divergence of a sympatric cryptic lineage in Cycas revoluta. *Ecology and Evolution*,

13(8), e10435. https://doi.org/10.1002/ece3.10435

Chen, M.-Z., Cao, L.-J., Li, B.-Y., Chen, J.-C., Gong, Y.-J., Yang, Q., Schmidt, T. L., Yue, L., Zhu, J.-Y., Li,

H., Chen, X.-X., Hoffmann, A. A., & Wei, S.-J. (2021). Migration trajectories of the

diamondback moth Plutella xylostella in China inferred from population genomic variation.

Pest Management Science, *77*(4), 1683–1693. https://doi.org/10.1002/ps.6188

Chiou, K. L., & Bergey, C. M. (2018). Methylation-based enrichment facilitates low-cost, noninvasive

genomic scale sequencing of populations from feces. *Scientific Reports*, *8*(1), 1–10.

https://doi.org/10.1038/s41598-018-20427-9

Cincotta, R. P., Wisnewski, J., & Engelman, R. (2000). Human population in the biodiversity hotspots.

Nature, *404*(6781), Article 6781. https://doi.org/10.1038/35010105

Coulon, A., Guillot, G., Cosson, J.-F., Angibault, J. M. A., Aulagnier, S., Cargnelutti, B., Galan, M., &

Hewison, A. J. M. (2006). Genetic structure is influenced by landscape features: Empirical

evidence from a roe deer population. *Molecular Ecology*, *15*(6), 1669–1679.

https://doi.org/10.1111/j.1365-294X.2006.02861.x

Craigie, I. D., Baillie, J. E. M., Balmford, A., Carbone, C., Collen, B., Green, R. E., & Hutton, J. M.

(2010). Large mammal population declines in Africa's protected areas. *Biological*

Conservation, *143*(9), 2221–2228. https://doi.org/10.1016/j.biocon.2010.06.007

Crego, R. D., Wells, H. B. M., Ndung'u, K. S., Evans, L., Njeri Nduguta, R., Chege, M. A., Brown, M. B.,

Ogutu, J. O., Ojwang, G. O., Fennessy, J., O'Connor, D., Stacy-Dawes, J., Rubenstein, D. I.,

Martins, D. J., Leimgruber, P., & Stabach, J. A. (2021). Moving through the mosaic:

Identifying critical linkage zones for large herbivores across a multiple-use African landscape.

Landscape Ecology, *36*(5), 1325–1340. https://doi.org/10.1007/s10980-021-01232-8

Crooks, K. R., Burdett, C. L., Theobald, D. M., Rondinini, C., & Boitani, L. (2011). Global patterns of

fragmentation and connectivity of mammalian carnivore habitat. *Philosophical Transactions*

of the Royal Society B: Biological Sciences, *366*(1578), 2642–2651.

https://doi.org/10.1098/rstb.2011.0120

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., Handsaker, R. E., Lunter,

G., Marth, G. T., Sherry, S. T., McVean, G., & Durbin, R. (2011). The variant call format and

VCFtools. *Bioinformatics*, *27*(15), 2156–2158. https://doi.org/10.1093/bioinformatics/btr330

Dirzo, R., Young, H. S., Galetti, M., Ceballos, G., Isaac, N. J. B., & Collen, B. (2014). Defaunation in the

Anthropocene. *Science*, *345*(6195), 401–406. https://doi.org/10.1126/science.1251817

Dudaniec, R. Y., Wilmer, J. W., Hanson, J. O., Warren, M., Bell, S., & Rhodes, J. R. (2016). Dealing with

uncertainty in landscape genetic resistance models: A case of three co-occurring marsupials.

Molecular Ecology, *25*(2), 470–486. https://doi.org/10.1111/mec.13482

Dutta, T., Sharma, S., & DeFries, R. (2018). Targeting restoration sites to improve connectivity in a

tiger conservation landscape in India. *PeerJ*, *6*, e5587. https://doi.org/10.7717/peerj.5587

Dutta, T., Sharma, S., Maldonado, J. E., Panwar, H. S., & Seidensticker, J. (2015). Genetic Variation,

Structure, and Gene Flow in a Sloth Bear (Melursus ursinus) Meta-Population in the Satpura-

Maikal Landscape of Central India. *PLOS ONE*, *10*(5), e0123384.

https://doi.org/10.1371/journal.pone.0123384

Dutta, T., Sharma, S., Maldonado, J. E., Wood, T. C., Panwar, H. S., & Seidensticker, J. (2013). Fine-scale population genetic structure in a wide-ranging carnivore, the leopard (Panthera pardus fusca) in central India. *Diversity and Distributions*, *19*(7), 760–771. https://doi.org/10.1111/ddi.12024

Epps, C. W., & Keyghobadi, N. (2015). Landscape genetics in a changing world: Disentangling historical and contemporary influences and inferring change. *Molecular Ecology*, *24*(24), 6021–6040. https://doi.org/10.1111/mec.13454

Epps, C. W., Palsbøll, P. J., Wehausen, J. D., Roderick, G. K., Ramey II, R. R., & McCullough, D. R. (2005). Highways block gene flow and cause a rapid decline in genetic diversity of desert bighorn sheep. *Ecology Letters*, *8*(10), 1029–1038. https://doi.org/10.1111/j.1461-0248.2005.00804.x

Ernest, H. B., Vickers, T. W., Morrison, S. A., Buchalski, M. R., & Boyce, W. M. (2014). Fractured Genetic Connectivity Threatens a Southern California Puma (Puma concolor) Population. *PLOS ONE*, *9*(10), e107985. https://doi.org/10.1371/journal.pone.0107985

Evanno, G., Regnaut, S., & Goudet, J. (2005). Detecting the number of clusters of individuals using the software structure: A simulation study. *Molecular Ecology*, *14*(8), 2611–2620. https://doi.org/10.1111/j.1365-294X.2005.02553.x

Farah, S., Atkulwar, A., Nahid, R., Gadhikar, Y., & Baig, M. (2021). Recent population expansion in wild gaur (Bos gaurus gaurus) as revealed by microsatellite markers. *Mammalian Biology*, *101*(5), 695–707. https://doi.org/10.1007/s42991-021-00145-y

Fedorca, A., Popa, M., Jurj, R., Ionescu, G., Ionescu, O., & Fedorca, M. (2020). Assessing the regional landscape connectivity for multispecies to coordinate on-the-ground needs for mitigating linear infrastructure impact in Brasov – Prahova region. *Journal for Nature Conservation*, *58*, 125903. https://doi.org/10.1016/j.jnc.2020.125903

Feehery, G. R., Yigit, E., Oyola, S. O., Langhorst, B. W., Schmidt, V. T., Stewart, F. J., Dimalanta, E. T., Amaral-Zettler, L. A., Davis, T., Quail, M. A., & Pradhan, S. (2013). A Method for Selectively

Enriching Microbial DNA from Contaminating Vertebrate Host DNA. *PLOS ONE*, *8*(10),

e76096. https://doi.org/10.1371/journal.pone.0076096

Foll, M., & Gaggiotti, O. (2008). A Genome-Scan Method to Identify Selected Loci Appropriate for

Both Dominant and Codominant Markers: A Bayesian Perspective. *Genetics*, *180*(2), 977–

993. https://doi.org/10.1534/genetics.108.092221

Fraser, D. L., Ironside, K., Wayne, R. K., & Boydston, E. E. (2019). Connectivity of mule deer

(Odocoileus hemionus) populations in a highly fragmented urban landscape. *Landscape*

Ecology, *34*(5), 1097–1115. https://doi.org/10.1007/s10980-019-00824-9

Garrison, E., & Marth, G. (2012). Haplotype-based variant detection from short-read sequencing.

arXiv:1207.3907 [q-Bio]. http://arxiv.org/abs/1207.3907

Goudet, J. (2005). Hierfstat, a package for r to compute and test hierarchical F-statistics. *Molecular*

Ecology Notes, *5*(1), 184–186. https://doi.org/10.1111/j.1471-8286.2004.00828.x

Habib, B., Rajvanshi, A., Mathur, V. B., & Saxena, A. (2016). Corridors at Crossroads: Linear

Development-Induced Ecological Triage As a Conservation Opportunity. *Frontiers in Ecology*

and Evolution, *4*, 132. https://doi.org/10.3389/fevo.2016.00132

Hayward, K. M., Clemente-Carvalho, R. B. G., Jensen, E. L., de Groot, P. V. C., Branigan, M., Dyck, M.,

Tschritter, C., Sun, Z., & Lougheed, S. C. (2022). Genotyping-in-thousands by sequencing (GT-

seq) of noninvasive faecal and degraded samples: A new panel to enable ongoing monitoring

of Canadian polar bear populations. Molecular Ecology Resources, 22(5), 1906–1918.

https://doi.org/10.1111/1755-0998.13583

Hepenstrick, D., Thiel, D., Holderegger, R., & Gugerli, F. (2012). Genetic discontinuities in roe deer

(Capreolus capreolus) coincide with fenced transportation infrastructure. *Basic and Applied*

Ecology, *13*(7), 631–638. https://doi.org/10.1016/j.baae.2012.08.009

Holderegger, R., & Wagner, H. H. (2008). Landscape Genetics. *BioScience*, *58*(3), 199–207.

https://doi.org/10.1641/B580306

Ibisch, P. L., Hoffmann, M. T., Kreft, S., Pe'er, G., Kati, V., Biber-Freudenberger, L., DellaSala, D. A.,

Vale, M. M., Hobson, P. R., & Selva, N. (2016). A global map of roadless areas and their

conservation status. *Science*, *354*(6318), 1423–1427.

https://doi.org/10.1126/science.aaf7166

Jackson, H. B., & Fahrig, L. (2015). Are ecologists conducting research at the optimal scale? *Global*

Ecology and Biogeography, *24*(1), 52–63. https://doi.org/10.1111/geb.12233

Jayadevan, A., Nayak, R., Karanth, K. K., Krishnaswamy, J., DeFries, R., Karanth, K. U., &

Vaidyanathan, S. (2020). Navigating paved paradise: Evaluating landscape permeability to

movement for large mammals in two conservation priority landscapes in India. *Biological*

Conservation, *247*, 108613. https://doi.org/10.1016/j.biocon.2020.108613

Jhala, Y. V., Gopal, R., & Qureshi, Q. (Qamar). (2008). *Status of tigers, co-predators, and prey in India*.

http://agris.fao.org/agris-search/search.do?recordID=US201300137392

Jombart, T. (2008). adegenet: A R package for the multivariate analysis of genetic markers.

Bioinformatics (Oxford, England), *24*(11), 1403–1405.

https://doi.org/10.1093/bioinformatics/btn129

Khan, A., Patel, K., Bhattacharjee, S., Sharma, S., Chugani, A. N., Sivaraman, K., Hosawad, V., Sahu, Y.

K., Reddy, G. V., & Ramakrishnan, U. (2020). Are shed hair genomes the most effective

noninvasive resource for estimating relationships in the wild? *Ecology and Evolution*, *10*(11),

4583–4594. https://doi.org/10.1002/ece3.6157

Khan, A., & Tyagi, A. (2021). Considerations for Initiating a Wildlife Genomics Research Project in

South and South-East Asia. *Journal of the Indian Institute of Science*, *101*(2), 243–256.

https://doi.org/10.1007/s41745-021-00243-3

Kohn, M. H., & Wayne, R. K. (1997). Facts from feces revisited. *Trends in Ecology & Evolution*, *12*(6),

223–227. https://doi.org/10.1016/S0169-5347(97)01050-1

Kopelman, N. M., Mayzel, J., Jakobsson, M., Rosenberg, N. A., & Mayrose, I. (2015). Clumpak: A

program for identifying clustering modes and packaging population structure inferences

837 across K. *Molecular Ecology Resources*, *15*(5), 1179–1191. https://doi.org/10.1111/1755-

838 0998.12387

839 Landguth, E. L., Fedy, B. C., OYLER-McCANCE, S. J., Garey, A. L., Emel, S. L., Mumma, M., Wagner, H.

840 H., Fortin, M.-J., & Cushman, S. A. (2012). Effects of sample size, number of markers, and

841 allelic richness on the detection of spatial genetic pattern. *Molecular Ecology Resources*,

842 *12*(2), 276–284. https://doi.org/10.1111/j.1755-0998.2011.03077.x

843 Laurance, W. F., & Balmford, A. (2013). Land use: A global map for road building. *Nature*, *495*, 308–

844 309. https://doi.org/10.1038/495308a

845 le Roux, E., van Veenhuisen, L. S., Kerley, G. I. H., & Cromsigt, J. P. G. M. (2020). Animal body size

846 distribution influences the ratios of nutrients supplied to plants. *Proceedings of the National*

847 *Academy of Sciences*, *117*(36), 22256–22263. https://doi.org/10.1073/pnas.2003269117

848 Lemopoulos, A., Prokkola, J. M., Uusi-Heikkilä, S., Vasemägi, A., Huusko, A., Hyvärinen, P., Koljonen,

849 M.-L., Koskiniemi, J., & Vainikka, A. (2019). Comparing RADseq and microsatellites for

850 estimating genetic diversity and relatedness—Implications for brown trout conservation.

851 *Ecology and Evolution*, *9*(4), 2106–2120. https://doi.org/10.1002/ece3.4905

852 Li, H. (2013). Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.

853 *arXiv:1303.3997 [q-Bio]*. http://arxiv.org/abs/1303.3997

854 Ling, C., Lixia, W., Rong, H., Fujun, S., Wenping, Z., Yao, T., Yaohua, Y., Bo, Z., & Liang, Z. (2020).

855 Comparative analysis of microsatellite and SNP markers for parentage testing in the golden

856 snub-nosed monkey (Rhinopithecus roxellana). *Conservation Genetics Resources*.

857 https://doi.org/10.1007/s12686-020-01147-7

858 Longmire, J. L., Maltbie, M., Baker, R. J., Texas Tech University, & Museum. (1997). *Use of "Lysis*

859 *Buffer" in DNA isolation and its implication for museum collections*. Museum of Texas Tech

860 University.

861 Mackiewicz, P., Matosiuk, M., Świsłocka, M., Zachos, F. E., Hajji, G. M., Saveljev, A. P., Seryodkin, I.

862 V., Farahvash, T., Rezaei, H. R., Torshizi, R. V., Mattioli, S., & Ratkiewicz, M. (2022). Phylogeny

and evolution of the genus Cervus (Cervidae, Mammalia) as revealed by complete

mitochondrial genomes. *Scientific Reports*, *12*(1), Article 1. https://doi.org/10.1038/s41598-

022-20763-x

Magbanua, Z. V., Hsu, C.-Y., Pechanova, O., Arick, M., Grover, C. E., & Peterson, D. G. (2023).

Innovations in double digest restriction-site associated DNA sequencing (ddRAD-Seq)

method for more efficient SNP identification. *Analytical Biochemistry*, *662*, 115001.

https://doi.org/10.1016/j.ab.2022.115001

McClure, M. L., Hansen, A. J., & Inman, R. M. (2016). Connecting models to movements: Testing

connectivity model predictions against empirical migration and dispersal data. *Landscape*

Ecology, *31*(7), 1419–1432. https://doi.org/10.1007/s10980-016-0347-0

Miller, J. R. B., Jhala, Y. V., Jena, J., & Schmitz, O. J. (2015). Landscape-scale accessibility of livestock

to tigers: Implications of spatial grain for modeling predation risk to mitigate human–

carnivore conflict. *Ecology and Evolution*, *5*(6), 1354–1367.

https://doi.org/10.1002/ece3.1440

Morin, P. A., Chambers, K. E., Boesch, C., & Vigilant, L. (2001). Quantitative polymerase chain

reaction analysis of DNA from noninvasive samples for accurate microsatellite genotyping of

wild chimpanzees (Pan troglodytes verus). *Molecular Ecology*, *10*(7), 1835–1844.

https://doi.org/10.1046/j.0962-1083.2001.01308.x

Musser, G. G., & Koop-man, K. F. (1994). Corbet, G. B., and J. E. Hill. 1992. The Mammals of the

Indomalayan Region: A Systematic Review. Oxford University Press, Oxford, United Kingdom

viii + 488 pp., 45 figs. ISBN 0-19-854693-9. Price (hardcover) $90.00. *Journal of Mammalogy*,

75(3), 799–803. https://doi.org/10.2307/1382538

Natesh, M., Taylor, R. W., Truelove, N. K., Hadly, E. A., Palumbi, S. R., Petrov, D. A., & Ramakrishnan,

U. (2019). Empowering conservation practice with efficient and economical genotyping from

poor quality samples. *Methods in Ecology and Evolution*, *10*(6), 853–859.

https://doi.org/10.1111/2041-210X.13173

Orkin, J. D., Montague, M. J., Tejada-Martinez, D., Manuel, M. de, Campo, J. del, Hernandez, S. C.,

Fiore, A. D., Fontsere, C., Hodgson, J. A., Janiak, M. C., Kuderna, L. F. K., Lizano, E., Martin, M.

P., Niimura, Y., Perry, G. H., Valverde, C. S., Tang, J., Warren, W. C., Magalhães, J. P. de, …

Melin, A. D. (2020). The evolution of ecological flexibility, large brains, and long lives:

Capuchin monkey genomics revealed with fecalFACS. *bioRxiv*, 366112.

https://doi.org/10.1101/366112

Paez, S., Kraus, R. H. S., Shapiro, B., Gilbert, M. T. P., Jarvis, E. D., & VERTEBRATE GENOMES PROJECT

CONSERVATION GROUP. (2022). Reference genomes for conservation. *Science*, *377*(6604),

364–366. https://doi.org/10.1126/science.abm8127

Peakall, R., & Smouse, P. E. (2012). GenAlEx 6.5: Genetic analysis in Excel. Population genetic

software for teaching and research—an update. *Bioinformatics*, *28*(19), 2537–2539.

https://doi.org/10.1093/bioinformatics/bts460

Peterman, W. E. (2018). ResistanceGA: An R package for the optimization of resistance surfaces

using genetic algorithms. *Methods in Ecology and Evolution*, *9*(6), 1638–1647.

https://doi.org/10.1111/2041-210X.12984

Peterson, B. K., Weber, J. N., Kay, E. H., Fisher, H. S., & Hoekstra, H. E. (2012). Double Digest RADseq:

An Inexpensive Method for De Novo SNP Discovery and Genotyping in Model and Non-

Model Species. *PLOS ONE*, *7*(5), e37135. https://doi.org/10.1371/journal.pone.0037135

Peterson, G. W., Dong, Y., Horbach, C., & Fu, Y.-B. (2014). Genotyping-By-Sequencing for Plant

Genetic Diversity Analysis: A Lab Guide for SNP Genotyping. *Diversity*, *6*(4), Article 4.

https://doi.org/10.3390/d6040665

Pritchard, J. K., Stephens, M., & Donnelly, P. (2000). Inference of population structure using

multilocus genotype data. *Genetics*, *155*(2), 945–959.

Putman, R. J. (1984). Facts from faeces. *Mammal Review*, *14*(2), 79–97.

https://doi.org/10.1111/j.1365-2907.1984.tb00341.x

Rengefors, K., Gollnisch, R., Sassenhagen, I., Aloisi, K. H., Svensson, M., Lebret, K., Čertnerová, D.,

Cresko, W. A., Bassham, S., & Ahrén, D. (2021). Genome-wide single nucleotide

polymorphism markers reveal population structure and dispersal direction of an expanding

nuisance algal bloom species. *Molecular Ecology*, *30*(4), 912–925.

https://doi.org/10.1111/mec.15787

Ripple, W. J., Newsome, T. M., Wolf, C., Dirzo, R., Everatt, K. T., Galetti, M., Hayward, M. W., Kerley,

G. I. H., Levi, T., Lindsey, P. A., Macdonald, D. W., Malhi, Y., Painter, L. E., Sandom, C. J.,

Terborgh, J., & Van Valkenburgh, B. (2015). Collapse of the world's largest herbivores.

Science Advances, *1*(4), e1400103. https://doi.org/10.1126/sciadv.1400103

Rouet-Leduc, J., Pe'er, G., Moreira, F., Bonn, A., Helmer, W., Shahsavan Zadeh, S. A. A., Zizka, A., &

van der Plas, F. (2021). Effects of large herbivores on fire regimes and wildfire mitigation.

Journal of Applied Ecology, *58*(12), 2690–2702. https://doi.org/10.1111/1365-2664.13972

Sanderson, E. W., Moy, J., Rose, C., Fisher, K., Jones, B., Balk, D., Clyne, P., Miquelle, D., & Walston, J.

(2019). Implications of the shared socioeconomic pathways for tiger (Panthera tigris)

conservation. *Biological Conservation*, *231*, 13–23.

https://doi.org/10.1016/j.biocon.2018.12.017

Sandom, C. J., Ejrnæs, R., Hansen, M. D. D., & Svenning, J.-C. (2014). High herbivore density

associated with vegetation diversity in interglacial ecosystems. *Proceedings of the National*

Academy of Sciences, *111*(11), 4162–4167. https://doi.org/10.1073/pnas.1311014111

Sankar, K., Pabla, H. S., Patil, C. K., Nigam, P., Qureshi, Q., Navaneethan, B., Manjreakar, M., Virkar, P.

S., & Mondal, K. (2013). Home Range, Habitat Use and Food Habits of Re-Introduced Gaur

(Bos Gaurus Gaurus) in Bandhavgarh Tiger Reserve, Central India. *Tropical Conservation*

Science, *6*(1), 50–69. https://doi.org/10.1177/194008291300600108

Sawyer, H., Merkle, J. A., Middleton, A. D., Dwinnell, S. P. H., & Monteith, K. L. (2019). Migratory

plasticity is not ubiquitous among large herbivores. *Journal of Animal Ecology*, *88*(3), 450–

460. https://doi.org/10.1111/1365-2656.12926

Schoen, J. M., Neelakantan, A., Cushman, S. A., Dutta, T., Habib, B., Jhala, Y. V., Mondal, I.,

Ramakrishnan, U., Reddy, P. A., Saini, S., Sharma, S., Thatte, P., Yumnam, B., & DeFries, R.

(2022). Synthesizing habitat connectivity analyses of a globally important human-dominated

tiger-conservation landscape. *Conservation Biology*, *36*(4), e13909.

https://doi.org/10.1111/cobi.13909

Schrader, C., Schielke, A., Ellerbroek, L., & Johne, R. (2012). PCR inhibitors – occurrence, properties

and removal. *Journal of Applied Microbiology*, *113*(5), 1014–1026.

https://doi.org/10.1111/j.1365-2672.2012.05384.x

Seto, K. C., Güneralp, B., & Hutyra, L. R. (2012). Global forecasts of urban expansion to 2030 and

direct impacts on biodiversity and carbon pools. *Proceedings of the National Academy of

Sciences*, *109*(40), 16083–16088. https://doi.org/10.1073/pnas.1211658109

Shirasawa, K., Hirakawa, H., & Isobe, S. (2016). Analytical workflow of double-digest restriction site-

associated DNA sequencing based on empirical and in silico optimization in tomato. *DNA

Research*, *23*(2), 145–153. https://doi.org/10.1093/dnares/dsw004

Shirk, A. J., Landguth, E. L., & Cushman, S. A. (2017). A comparison of individual-based genetic

distance metrics for landscape genetics. *Molecular Ecology Resources*, *17*(6), 1308–1317.

https://doi.org/10.1111/1755-0998.12684

Shirk, A. J., Landguth, E. L., & Cushman, S. A. (2018). A comparison of regression methods for model

selection in individual-based landscape genetic analysis. *Molecular Ecology Resources*, *18*(1),

55–67. https://doi.org/10.1111/1755-0998.12709

Shirk, A. J., Wallin, D. O., Cushman, S. A., Rice, C. G., & Warheit, K. I. (2010). Inferring landscape

effects on gene flow: A new model selection framework. *Molecular Ecology*, *19*(17), 3603–

3619. https://doi.org/10.1111/j.1365-294X.2010.04745.x

Singh, V. K., Joshi, B. D., Bhat, G. J., Singh, S. K., Chandra, K., Sharma, L. K., & Thakur, M. (2021).

Population genetics of Sambar (Rusa unicolor) from the Western Himalayas: Preliminary

findings. *Molecular Biology Reports*. https://doi.org/10.1007/s11033-021-06845-5

Snyder-Mackler, N., Majoros, W. H., Yuan, M. L., Shaver, A. O., Gordon, J. B., Kopp, G. H., Schlebusch,

S. A., Wall, J. D., Alberts, S. C., Mukherjee, S., Zhou, X., & Tung, J. (2016). Efficient Genome-

Wide Sequencing and Low-Coverage Pedigree Analysis from Noninvasively Collected

Samples. *Genetics*, *203*(2), 699–714. https://doi.org/10.1534/genetics.116.187492

Srivathsa, A., Puri, M., Karanth, K. K., Patel, I., & Kumar, N. S. (2019). Examining human–carnivore

interactions using a socio-ecological framework: Sympatric wild canids in India as a case

study. *Royal Society Open Science*, *6*(5), 182008. https://doi.org/10.1098/rsos.182008

Storfer, A., Murphy, M. A., Evans, J. S., Goldberg, C. S., Robinson, S., Spear, S. F., Dezzani, R.,

Delmelle, E., Vierling, L., & Waits, L. P. (2007). Putting the 'landscape' in landscape genetics.

Heredity, *98*(3), Article 3. https://doi.org/10.1038/sj.hdy.6800917

Thatte, P., Chandramouli, A., Tyagi, A., Patel, K., Baro, P., Chhattani, H., & Ramakrishnan, U. (2020).

Human footprint differentially impacts genetic connectivity of four wide-ranging mammals

in a fragmented landscape. *Diversity and Distributions*, *26*(3), 299–314.

https://doi.org/10.1111/ddi.13022

Thatte, P., Joshi, A., Vaidyanathan, S., Landguth, E., & Ramakrishnan, U. (2018). Maintaining tiger

connectivity and minimizing extinction into the next century: Insights from landscape

genetics and spatially-explicit simulations. *Biological Conservation*, *218*, 181–191.

https://doi.org/10.1016/j.biocon.2017.12.022

Thatte, P., Tyagi, A., Neelakantan, A., Natesh, M., Sen, M., & Thekaekara, T. (2021). Trends in Wildlife

Connectivity Science from the Biodiverse and Human-Dominated South Asia. *Journal of the*

Indian Institute of Science, *101*(2), 177–193. https://doi.org/10.1007/s41745-021-00240-6

Timmins, R., Group, W. D. (IUCN S. S. C. S., Group), I. S. D. S., Nepal), H. S. B. (Bird C., Mekong), T. G.

(WWF, History), S. M. (Salim A. C. for O. and N., & Sg), D. J. (Cat. (2014). IUCN Red List of

Threatened Species: Felis chaus. *IUCN Red List of Threatened Species*.

https://www.iucnredlist.org/en

Tyagi, A., Godbole, M., Vanak, A. T., & Ramakrishnan, U. (2023). Citizen science facilitates first ever

992   genetic detection of wolf-dog hybridization in Indian savannahs. *Ecology and Evolution*,

993   *13*(5), e10100. https://doi.org/10.1002/ece3.10100

994   Tyagi, A., Khan, A., Thatte, P., & Ramakrishnan, U. (2022). Genome-wide single nucleotide

995   polymorphism (SNP) markers from fecal samples reveal anthropogenic impacts on

996   connectivity: Case of a small carnivore in the central Indian landscape. *Animal Conservation*,

997   *25*(5), 648–659. https://doi.org/10.1111/acv.12770

998   Tyagi, A., Kumar, V., Kittur, S., Reddy, M., Naidenko, S., Ganswindt, A., & Umapathy, G. (2019).

999   Physiological stress responses of tigers due to anthropogenic disturbance especially tourism

in two central Indian tiger reserves. *Conservation Physiology*, *7*(1).

https://doi.org/10.1093/conphys/coz045

Venter, O., Sanderson, E. W., Magrach, A., Allan, J. R., Beher, J., Jones, K. R., Possingham, H. P.,

Laurance, W. F., Wood, P., Fekete, B. M., Levy, M. A., & Watson, J. E. M. (2016). Sixteen

years of change in the global terrestrial human footprint and implications for biodiversity

conservation. *Nature Communications*, *7*(1), Article 1.

https://doi.org/10.1038/ncomms12558

Weckworth, B. V., Musiani, M., DeCesare, N. J., McDevitt, A. D., Hebblewhite, M., & Mariani, S.

(2013). Preferred habitat and effective population size drive landscape genetic patterns in an

endangered species. *Proceedings of the Royal Society B: Biological Sciences*, *280*(1769),

20131756. https://doi.org/10.1098/rspb.2013.1756

White, L. C., Moseby, K. E., Thomson, V. A., Donnellan, S. C., & Austin, J. J. (2018). Long-term genetic

consequences of mammal reintroductions into an Australian conservation reserve. *Biological*

Conservation, *219*, 1–11. https://doi.org/10.1016/j.biocon.2017.12.038

Whitlock, M. C., & McCauley, D. E. (1999). Indirect measures of gene flow and migration: $F_{ST} \neq 1/(4$

$N_m +1)$. *Heredity*, *82*(2), Article 2. https://doi.org/10.1038/sj.hdy.6884960

Wu, D.-D., Ding, X.-D., Wang, S., Wójcik, J. M., Zhang, Y., Tokarska, M., Li, Y., Wang, M.-S., Faruque,

O., Nielsen, R., Zhang, Q., & Zhang, Y.-P. (2018). Pervasive introgression facilitated

domestication and adaptation in the Bos species complex. *Nature Ecology & Evolution*, *2*(7),

1139–1145. https://doi.org/10.1038/s41559-018-0562-y

Young, H. S., McCauley, D. J., Galetti, M., & Dirzo, R. (2016). Patterns, Causes, and Consequences of

Anthropocene Defaunation. *Annual Review of Ecology, Evolution, and Systematics*, *47*(1),

333–358. https://doi.org/10.1146/annurev-ecolsys-112414-054142

Yumnam, B., Jhala, Y. V., Qureshi, Q., Maldonado, J. E., Gopal, R., Saini, S., Srinivas, Y., & Fleischer, R.

C. (2014). Prioritizing Tiger Conservation through Landscape Genetics and Habitat Linkages.

PLOS ONE, *9*(11), e111207. https://doi.org/10.1371/journal.pone.0111207

1026

1027

1028

1029

1030

1031

1032

1033

1034

1035

1036

1037

1038

1039

1040

1041

1042

1043

Table 1: Population-wise summary of the number of individuals, observed and expected
heterozygosity for both species.

1046

Population	Gaur			Sambar		
	N	Ho	He	N	Ho	He
KTR	33	0.131	0.152	12	0.195	0.218
KPC	10	0.131	0.142	11	0.194	0.218
PTR	25	0.121	0.145	26	0.194	0.226
NNTR	31	0.124	0.152	21	0.187	0.226
TATR	14	0.129	0.147	15	0.209	0.224
BOR	–	–	–	14	0.165	0.209
UMR	11	0.122	0.148	–	–	–

1047

1048

1049

1050

1051

1052

1053

1054

1055

1056

1057

1060

1061 **Table 2:** Multivariate optimization results for both gaur and sambar. The optimum
1062 parameters, Rmax: maximum resistance; x: shape parameter, for the different landscape
1063 variables for the best model based on lowest AIC score. The AIC score of the model in
1064 comparison with AIC score of the null model (Euclidian distance) along with the marginal R^2
1065 values are presented.

1066

	Gaur					Sambar				
Layers	**Rmax**	**x**	**AIC**	**AIC null**	**mR²**	**Rmax**	**x**	**AIC**	**AIC null**	**mR²**
LULC	1000	2				1000	5			
Roads	1000	2	-48431	-47114.2	0.503	100	0.1	-24681	-24627.2	0.173
Linden	1000	5				50	5			
Hpop	50	10				100	0.5			

1067

1068

1069

1070

1071

1072

1073

1074

1075

1076

1077

1078

1079

1080

1081

1082

Table 3: Bootstrapping results for gaur with 15 models including all possible combinations of covariates along with the null (no resistance) model. The average AIC scores for the respective model is reported along with average weight, average rank and the average log-likelihood of the parameter combination. Percent top represents how many times the respective model was selected as the best model during 10,000 iterations.

1088

S.No.	Model with landscape variable combinations	Average AIC	Average weight	Average rank	Average LL	Percent top
1	Roads+Linden+LULC	-27095.87	0.38	1.84	13550.93	65.76
2	Hpop+Linden+LULC	-27095.76	0.36	2.39	13550.88	19.75
3	Roads+Linden+LULC+Hpop	-27093.85	0.13	3.67	13550.93	0
4	Roads+Linden	-27046.11	0.11	5.03	13525.06	11.86
5	Linden	-27065.30	0.01	5.65	13533.65	0.96
6	Hpop+LULC	-27029.47	0.01	7.17	13516.73	0.66
7	Roads+Hpop+Linden	-27029.97	0.00	7.68	13517.99	0.01
8	Roads+Linden	-27036.87	0.00	8.43	13520.44	0.06
9	Roads+Hpop+LULC	-27027.46	0.00	8.49	13516.73	0
10	Hpop+LULC	-27024.18	0.01	9.35	13514.09	0.9
11	LULC	-27024.24	0.00	9.63	13513.12	0
12	Roads+Hpop	-27021.94	0.00	10.22	13512.97	0.04
13	Linden+LULC	-27003.70	0.00	12.22	13503.85	0
14	Null	-26982.23	0.00	13.77	13491.12	0
15	Roads	-26965.79	0.00	15.17	13483.89	0
16	Hpop	-26965.22	0.00	15.27	13483.61	0

1089

1090

1091

1092

Table 4: Bootstrapping results for sambar with 15 models including all possible
combinations of covariates along with the null (no resistance) model. The average AIC
scores for the respective model is reported. The average weight, rank and the mean log-
likelihood of the parameter combination are also shown along with how many times the
respective model was selected as the best model during 10,000 iterations.

1098

S.No.	Model with landscape variable combinations	Average AIC	Average weight	Average rank	Average LL	Percent top
1	Roads+Hpop+LULC	-13723.30	0.34	2.36	6863.65	38.6
2	Hpop+LULC	-13723.10	0.30	2.51	6863.53	28
3	Roads+Hpop	-13719.30	0.15	3.90	6863.66	18.1
4	Roads+Hpop+LULC+Linden	-13718.80	0.05	4.79	6862.42	1.2
5	LULC+Linden	-13708.50	0.05	6.60	6856.27	11.2
6	Hpop+Linden	-13710.60	0.01	6.75	6857.29	0.8
7	Roads+LULC	-13708.50	0.05	7.46	6856.27	0.4
8	Hpop+Linden+LULC	-13711.10	0.01	7.68	6858.54	0
9	Roads+Hpop+Linden	-13708.60	0.00	8.69	6857.29	0
10	Roads+Linden+LULC	-13706.50	0.02	9.30	6856.27	0
11	Roads	-13705.10	0.01	9.44	6853.55	0.7
12	Roads+Hpop	-13705.10	0.00	9.70	6854.53	0.1
13	LULC+Linden	-13698.30	0.01	11.84	6849.15	0.9
14	Linden	-13670.00	0.00	14.50	6835.99	0
15	Hpop	-13668.80	0.00	15.13	6835.41	0
16	Null	-13668.70	0.00	15.36	6835.34	0

1099

1100

Figure 1: Study area with sampling sites. Red and yellow dots represent gaur (top right) and
sambar (bottom right) individuals respectively. Two major national highways (NH6 & 7),
bisecting the study landscape are also shown as bold black lines. Protected areas are
marked in dark green colour with black boundaries. KTR – Kanha Tiger Reserve; KPC – Kanha
Pench corridor; PTR – Pench Tiger Reserve; NNTR – Nagzira Nawegaon Tiger Reserve; BOR –
Bor Tiger Reserve; TATR – Tadoba Andhari Tiger Reserve and UMR – Umred Karhandala
Wildlife Sanctuary.

1108

1109

1110

1111

1112

1113

Figure 2: Population structure results of gaur described by (a) discriminant analysis of
principal components (DAPC) based on 1186 SNPs. Each point represents a unique individual
and each color corresponds to population, as assessed by STRUCTURE analysis. (b)
STRUCTURE analysis plot where each vertical line represents a unique individual. Samples
are colour-coded based on population. (c) Scatterplots for isolation by distance between
genetic and geographic distance, colours indicate the density of data points as high (red),
medium (yellow), and low (blue). (d) Spatial autocorrelation at distance classes of 20 km
each till 300 km with r being the autocorrelation coefficient. Red dotted lines represent a
95% confidence interval. (e) Pairwise Fst between all the populations sampled.

1124

1125

1126

1127

1128

Figure 3: Population structure results of sambar based on (a) DAPC and (b) STRUCTURE analyses. Samples are colour-coded based on population. (c) Scatterplot between genetic and geographic distance. The density of data points is depicted with colours: red (high) to blue (low). (d) Spatial autocorrelation at distance classes of 20 km each till 300 km with r being autocorrelation coefficient. The dotted red lines represent a 95% confidence interval. (e) Pairwise Fst between all the populations sampled.

1136

1137

1138

1139

1140

1141

1142

1143

1144

1145 **Figure 4:** Association between landscape variables and resistance offered along with the
1146 optimized model parameters, Rmax and x, for both species. A- depicts land use and land
1147 cover variable, whereas B- depicts roads with traffic. The density of linear features and
1148 human population is depicted as C and D respectively. The thickness of the arrows shows
1149 the magnitude of resistance offered by each landscape variable.

1150

1151

1152

1153

1154

Figure 5: Optimized resistance surfaces for (a) gaur and (b) sambar. Dark colour represents areas with high resistance and light colours depict areas with low resistance. Black dots are the sample locations of the respective species on each map with protected areas marked as green polygons.