Degradation of Fatty Acid Export Protein1 by Rhomboid-Like Protease11 Contributes to Cold Tolerance in Arabidopsis

Annalisa John[a], Moritz Krämer[b], Martin Lehmann[b], Hans-Henning Kunz[b], Fayezeh Aarabi[c], Saleh Alseekh[c], Alisdair Fernie[c], Frederik Sommer[d], Michael Schroda[d], David Zimmer[e], Timo Mühlhaus[e], Helga Peisker[f], Katharina Gutbrod[f], Peter Dörmann[f], Jens Neunzig[g], Katrin Philippar[g] and H. Ekkehard Neuhaus[a*]

[a] Plant Physiology, University of Kaiserslautern, Erwin-Schrödinger-Str., D-67653 Kaiserslautern, Germany

[b] Plant Biochemistry, Faculty of Biology, Ludwig-Maximilians-Universität Munich, 82152 Planegg-Martinsried, Germany

[c] Max Planck Institut for Molecular Plant Physiology, Wissenschaftspark Golm, D-14476 Potsdam, Germany

[d] Molecular Biotechnology and Systems Biology, University of Kaiserslautern, Erwin-Schrödinger-Str., D-67653 Kaiserslautern, Germany

[e] Computational Systems Biology, University of Kaiserslautern, Erwin-Schrödinger-Str., D-67653 Kaiserslautern, Germany

[f] Institute for Molecular Physiology and Biotechnology of Plants, IMBIO, University of Bonn, Karlrobert-Kreiten-Str. 13, D-53115 Bonn, Germany

[g] Plant Biology, Center for Human and Molecular Biology (ZHMB), Saarland University, D-66123 Saarbrücken, Germany

* Address correspondence to:

Ekkehard Neuhaus, University of Kaiserslautern, Plant Physiology, Gottlieb Daimler-Str. 47, D-67653 Kaiserslautern, Germany

Tel.: +49-631-2052372

Fax: +49-631-205-2600

E-mail: neuhaus@rhrk.uni-kl.de

Short title: RBL11 regulates FAX1 protein abundance in cold

One sentence summary: Degradation of the inner envelope protein Fatty Acid Export1 via Rhomboid Like Protease11 represents a critical process to achieve cold and frost tolerance in Arabidopsis

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors is: Ekkehard Neuhaus (neuhaus@rhrk.uni-kl.de).

Abstract

Plants need to adapt to different stresses to optimize growth under unfavorable conditions. The abundance of the chloroplast envelope located Fatty Acid Export Protein1 (FAX1) decreases after the onset of low temperatures. However, it was unclear how FAX1 degradation occurs and whether altered FAX1 abundance contributes to cold tolerance in plants. The rapid cold-induced increase in rhomboid-like protease11 (RBL11) transcript, the physical interaction of RBL11 with FAX1, the specific FAX1 degradation after RBL11 expression, and the absence of cold-induced FAX1 degradation in *rbl11* loss-of-function mutants suggest that this enzyme is responsible for FAX1 degradation. Proteomic analyses showed that *rbl11* mutants have higher levels of FAX1 and other proteins involved in membrane lipid homeostasis, suggesting that RBL11 is a key element in the remodeling of membrane properties during cold. Consequently, in the cold, *rbl11* mutants show a shift in lipid biosynthesis towards the eukaryotic pathway, which coincides with impaired cold tolerance. To demonstrate that cold sensitivity is due to increased FAX1 levels, FAX1 overexpressors were analyzed. *rbl11* and FAX1 overexpressor mutants show superimposable phenotypic defects upon exposure to cold temperatures. Our results show that the cold-induced degradation of FAX1 by RBL11 is critical for Arabidopsis to survive cold and freezing periods.

Introduction

The vast majority of vascular plants are sessile. One of their most remarkable characteristics is their ability to cope with a wide range of environmental conditions. When light intensity, temperature, or water and nutrient availability leave certain ranges, corresponding stress stimuli trigger systemic responses such as genetic, metabolic, and, to some extent, morphological changes. These processes lead to a new metabolic homeostasis that allows the plant to successfully cope with the environmental challenge (Obata and Fernie, 2012; Koevoets et al., 2016; Choudhury et al., 2017; Pommerrenig et al., 2018; Garcia-Molina et al., 2020; Wang et al., 2020).

Changes in growth temperature are usually more rapid than changes in water or nutrient availability. Accordingly, plants must rapidly initiate appropriate acclimation programs. These efficient molecular responses are essential because temperature affects the two main processes of photosynthesis, i.e., the light-driven electron transport across the thylakoid membrane and the subsequent enzyme-catalyzed Calvin-Benson cycle, in different ways. Thus, photosynthesis is markedly responsive to temperature changes, as demonstrated in several species representing a broad spectrum of CO_2-fixing organisms (Lin et al., 2012; Mackey et al., 2013; Walker et al., 2013; Song et al., 2014).

It is well-known that the composition of membrane lipids in plant cells exhibits a dynamic remodeling after onset of low temperatures (Moellering et al., 2010; Li et al., 2015; Barnes et al., 2016; Barrero-Sicilia et al., 2017). These structural changes comprise a higher degree of desaturation and altered abundancies of different phospho-, sulfo- or galacto-lipid species, which ensure sufficient extent of membrane fluidity under unfavorable environmental conditions (Smallwood and Bowles, 2002).

In plants, lipid biosynthesis represents a complex metabolic network in which initial metabolic steps in the plastids (e.g., chloroplasts) are connected to subsequent processes at the Endoplasmic Reticulum (ER) (Li-Beisson et al., 2010; Nakamura, 2017; Hölzl and Dörmann, 2019; Lavell and Benning, 2019). Generally, for the *de novo* synthesis of both classes of lipids, namely storage lipids (tri̲acyl-g̲lycerols, TAG) or membrane lipids (in chloroplasts mainly glyco-, phospho glycerolipids; and extraplastidic membranes, phosphoglycerolipids, sphingolipids, and sterol lipids) it is necessary that fatty acid synthe-

sis in plastids provides acyl-chains for subsequent steps located in both, the plastid and

the ER (Rawsthorne, 2002; Li-Beisson et al., 2010). Accordingly, the subsequent lipid

biosynthesis takes either place via the plastid located "prokaryotic pathway" or via the ER

located "eukaryotic pathway" (Roughan and Slack, 1982).

While TAG is synthesized in the ER, the biosynthesis of membrane lipid occurs in both

organelles the ER and in plastids (Li-Beisson et al., 2013). During membrane lipid

synthesis, the usage of fatty acids in chloroplasts or alternatively in the ER leads to

different types of structural lipids. Generally, plastids represent the major site for

phosphatidyl-glycerol (PG) synthesis and the exclusive site for mono- and digalactosyl-

diacylglycerol (MGDG and DGDG) biosynthesis, as well as for sulfoquinovosyl-

diacylglycerol (SQDG) assembly. The ER and plastids are responsible for the provision of

diacyl-glycerol (DAG) backbones which serve as a precursor for all ER-borne

phospholipids, including phosphatidylcholine (PC) and phosphatidylethanolamine (PE)

(Hagio et al., 2002; Andersson and Dörmann, 2009; Li-Beisson et al., 2010; Lavell and

Benning, 2019). (Hagio et al., 2002; Andersson and Dörmann, 2009; Li-Beisson et al.,

2010; Lavell and Benning, 2019). DAG synthesis in chloroplasts is particularly pronounced

under cold conditions because the Sensitive to Freezing2 (SFR2) enzyme transfers

galactosyl groups from MGDG to other galactolipids, resulting in the formation of

oligogalactolipids (Moellering et al., 2010; Barnes et al., 2016). Besides the plastid-

produced DAG, the ER-derived DAG moieties also act - after import into plastids - as

precursors for plastid lipid biosynthesis (Li-Beisson et al., 2010).

Each membrane type is defined by a characteristic lipid composition. This composition is

dynamic in response to changing environmental conditions with individual lipid mixtures

giving rise to specific membrane properties (van Meer et al., 2008; Li-Beisson et al., 2010;

Moellering et al., 2010). For example, at low temperatures, lipid remodeling maintains

membrane fluidity to prevent ion leakage, to keep carrier and receptor proteins functional

or to integrate novel protective proteins (Steponkus et al., 1977; Barrero-Sicilia et al.,

2017). Accordingly, plant mutants exhibiting altered activities of (i) selected fatty acid

biosynthesis enzymes, of (ii) fatty acid desaturases, of (iii) lipid transfer proteins or of (iv)

lipases might exhibit modified cold tolerance and photosynthesis properties (Miquel et al.,

1993; Welti et al., 2002; Khodakovskaya et al., 2006; Guo et al., 2013; Gao et al., 2020;

Schwenkert et al., 2023). The latter studies emphasize the impact of a proper membrane

lipid remodeling after the onset of low temperatures. In fact, it has been shown for various

species that low environmental temperatures lead to an upregulation of the chloroplast

located lipid biosynthesis pathway (Li et al., 2015). Although several proteins involved in

this process have been identified, the precise regulation responsible for this metabolic

shift is still unknown.

In contrast to chloroplasts, which are the site of fatty acid *de novo* synthesis in the cell (Li-

Beisson et al., 2010), the lipid biosynthesis pathway localized in the ER depends on import

of fatty acids from plastids. The molecular nature of transport proteins mediating export of

newly synthesized fatty acids from plastids had remained unknown for a long time (Wang

and Benning, 2012). However, with the identification of the protein Fatty Acid Export

Protein1 (FAX1) a first candidate was identified (Li et al., 2015) and the ability of FAX1 to

promote shuttling of fatty acids across membranes has been shown in recombinant

baker's yeast cells (Li et al., 2015). Furthermore, the absence of FAX1, which resides in

the plastid inner envelope membrane, leads apart from male sterility (Li et al., 2015; Zhu

et al., 2020) to decreased levels of ER-derived eukaryotic lipids, while the relative content

of PG, synthesized via the prokaryotic pathway, was increased. In contrast, FAX1

overexpressor lines exhibit an increased level of lipids assembled via the eukaryotic

pathway, e.g., more TAG (Li et al., 2015).

The chloroplast serves as a cellular hub coordinating genetic and molecular responses

required to acclimate to altered environmental conditions (Schwenkert et al., 2022). Thus,

all signals and each metabolite emitted from- or received by the chloroplast must pass the

inner-envelope membrane. Therefore, the inner-envelope proteome undergoes profound

changes in response to the onset of stress conditions (Nishimura et al., 2016; Pottosin

and Shabala, 2016; Wagner et al., 2016). Strikingly, FAX1 belongs to a group of inner

envelope proteins which exhibit a proteolytic degradation after exposure to low

temperatures, while other inner envelope proteins increase in their abundancy

(Trentmann et al., 2020). The mechanism responsible for the decrease of FAX1 in

response to cold temperatures is elusive. In addition, it is unknown whether this

phenomenon represents an important molecular process required for a maximal tolerance

against low temperatures and/or frost.

Chloroplasts contain more than 2000 different soluble- or membrane bound proteins
(Abdallah et al., 2000) and more than 20 different proteases ensure proteome homeo-
stasis within the organelle (Nishimura et al., 2016). So far, two major classes of proteases
have been identified to cleave intrinsic inner-envelope proteins, the metallo-dependent
AAA type FtsH proteases (with the isoforms FtsH 7, 9, 11 and 12) and the rhomboid-like
(RBL) proteases (with the isoforms RBL 10 and 11) (Knopf et al., 2012; Wagner et al.,
2016; Adam et al., 2019).

Thus, the function of FAX1 and the pronounced impact of lipid remodeling on plant
acclimation to cold raise two timely questions: First, what is the mechanism responsible
for the rapid decline of FAX1 upon onset of cold? Second, is the cold-induced
downregulation of FAX1 a pleotropic response or is it important for low temperature
tolerance? To answer both questions, we searched for an interaction of a selective
protease with FAX1. It turned out, that FAX1 and RBL11 can physically interact, and that
this protease is responsible for the cold-induced downregulation of FAX1. In addition, we
have shown that under cold conditions, *rbl11* mutants exhibit physiological and molecular
phenotypes similar to FAX1 overexpressors. Furthermore, we show that a lack of FAX1
degradation in response to cold impairs the ability of Arabidopsis plants to cope with low
temperature and freezing. These findings contribute to our understanding of how plants
tolerate adverse environmental conditions, one of the most impressive traits required to
ensure plant productivity.

Results

Transcript abundance of the plastid protease *RBL11* responds to cold treatments

It was previously shown that the protein Fatty Acid Export Protein1 belongs to a small
group of inner envelope proteins which are downregulated in their abundance upon onset
of cold temperatures (Trentmann et al., 2020). While FtsH12 is critical for chloroplast
development (Mielke et al., 2020), FtsH11 and RBL10 activity affect either, the
temperature response or the membrane lipid composition, respectively (Chen et al., 2006;
Wagner et al., 2011; Lavell et al., 2019). Thus, in addition to FtsH11 and RBL10, we
initially focused on RBL11 (the besides RBL10 only other rhomboid-like protease at the
inner envelope) which was shown to be active in fully developed mesophyll chloroplasts
(Knopf et al., 2012).

To gain first insight into possible molecular interactions between the proteases and FAX1
we analyzed the transcription level of corresponding genes after transfer to 4°C (Figure
1A), the standard temperature in our laboratory to study cold effects (Patzke et al., 2019;
Cvetkovic et al., 2021). Initially, transcript levels of all three envelope proteases
accumulated in response to the shift towards cold growth conditions. After one week at
4°C the most pronounced *RBL11* mRNA change was detected, which had about 14-fold
higher abundance when compared to the beginning of this treatment (Figure 1A). Both,
RBL10 and *FTSH11* mRNA amounts peaked at 4- or 7 times higher levels respectively,
after 7 days of cold treatment versus the start of the cold period (Figure 1A). The highest
levels of all three mRNAs were present after 14 days in cold conditions and from then on,
these three transcript levels declined in their abundance (Figure 1A).

***rbl11* mutants exhibit impaired growth at cold temperature since RBL11 controls**
FAX1 abundance in the cold

To search for specific responses in mutants lacking one of the three proteases, we grew
wild-type and respective loss-of-function lines under either, control conditions (21°C), or
for 7 or 56 days at 4°C.

When grown at 21°C, none of the mutants exhibited a phenotypic pattern distinct from

wild-type plants (Figure 1B). When grown at 4°C for 7 days all genotypes showed slightly

decreased chlorophyll levels, as indicated by brighter leaves (Figure 1B). However, after

56 days at 4°C all mutant plants exhibited impaired growth, i.e., smaller rosette sizes when

compared to the wild type (Figure 1B). We previously showed that plants with impaired

tolerance against cold temperatures exhibited increased numbers of wilted leaves when

exposed to 4°C (Trentmann et al., 2020). Interestingly, after 56 days at 4°C, *rbl11* mutants

exhibited wilted leaves (Figure 1B). This effect was further exacerbated in *rbl11* plants

after ten weeks at 4°C (Figure 1C, pictures from top and bottom of the rosette). A

quantification of this observation showed that after ten weeks at 4°C, wild types exhibited

1.25 wilted leaves/plant, while *rbl11* mutants displayed in five wilted leaves per plant on

average (Figure 1D).

The observation that from all three proteases tested, the *RBL11* mRNA is the fastest

responding transcript within 7 days after transfer to 4°C (Figure 1A), rendered the

corresponding enzyme a prime candidate for cold-induced FAX1 degradation. To test the

effect of RBL11 on FAX1 levels, we isolated chloroplast envelopes from wild types and

rbl11 loss-of-function mutant plants and conducted immunoblots using a previously

established FAX1 antibody (Li et al., 2015). The parallel immunoblotting of the inner

envelope protein TIC110 (Balsera et al., 2009) confirmed similar levels of total envelope

proteins in each lane (Figure 1E). Interestingly, already in *rbl11* mutants grown at control

temperature the FAX1 protein appeared increased by about 50% when compared to wild

type plants (Figure 1E). After 7 days in cold conditions, the FAX1 protein in wild type plants

decreased markedly (Figure 1E), which confirms our previous observations (Trentmann

et al., 2020). Such cold-induced FAX1 degradation was almost absent in *rbl11* mutants

compared to wild-type controls, while cold-induced FAX1 degradation occurred similarly

in *rbl10* and *ftsh11* mutants (Figure 1E).

Figure 1: Gene expression, phenotypic and immunoblot analysis of Arabidopsis protease loss-of-function mutants *rbl11*, *rbl10* and *ftsh11* grown under standard and cold (4°C) conditions. Plants were grown under standard conditions (21°C day and night temperature, 10h day length and 120µE light intensity) for 3 weeks and then treated with cold (4°C day and night temperature, 10h day length and 120µE light intensity). A) Gene expression levels of *RBL11*, *FTSH11* and *RBL10* in wild type by qRT-PCR under standard growth conditions (0 days at 4°C) and several days during cold treatment (7; 14; 28; 42, and 70 days at 4°C). Data represent relative mean expression levels of 3 biological replicates and are normalised to standard conditions (0 days at 4°C) using

UBQ as an internal control. B) Phenotypic analysis of *rbl11*, *rbl10* and *ftsh11* Arabidopsis plants. Images were taken from plants grown under standard conditions and after chilling for 7 and 56 days. C) Wt and *rbl11* rosettes were cut after 70 days of cold treatment and rotated to highlight the wilted leaves. D) Number of wilted leaves per plant in Wt and *rbl11* after 70 days of cold treatment. E) Immunoblot analysis via FAX1 antibody in isolated chloroplast envelopes of Wt, *rbl11*, *rbl10* and *ftsh11* from plants grown under standard conditions (21°C) and plants grown for 7 days at 4°C. Immunoblot detection via Tic110 antibody is used as a control and shows an equal loading of the protein samples at 3 µg per lane. Error bars in A) represent ± SEM. Error bars in D) are ± SD. Asterisks indicate significant differences between Wt and mutant using a t-test: p-value ≤0.001: *** (Supplemental file 1).

rbl11 mutants exhibit impaired frost tolerance

The observation that *rbl11* plants show an increased number of wilted leaves when grown at cold temperatures (Figure 1B-D) led us to test the freezing tolerance of this mutant. The ability of vascular plants to withstand freezing temperatures depends on a pretreatment of several days at temperatures above 0°C, called cold acclimation (Alberdi and Corcuera, 1991). To investigate whether the altered cold response in mutants affects freezing tolerance, we compared the ability of wild-types and *rbl11* plants to recover from exposure to freezing temperatures. To this end, plants were grown at 21°C and shifted to 4°C for four days to acclimate the cold. The temperature was then gradually reduced (2°C/h) until -10°C was reached. This temperature was maintained for 15h and then gradually increased (2°C/h) to 21°C before re-lighting (Trentmann et al., 2020). We assessed the phenotypic appearance of frost-treated plants after one and two additional weeks of growth at 21°C and quantified the number of wilted leaves as well as the survival rate after two weeks recovery phase.

As expected, unacclimated wild types and *rbl11* plants are unable to survive a freezing treatment (Figure 2A). As seen previously (Trentmann et al., 2020), wild-type plants are able to recover from a period of freezing with an efficiency of about 80% (Figure 2A,B) and exhibited 42% wilted leaves (Figure 2C). In contrast, *rbl11* mutants showed a significantly reduced ability to recover from freezing treatment, as only 28% of all plants survived the freezing period and more than 60% of the leaves were wilted (Figure 2A-C).

Figure 2: Recovery from freezing is impaired in *rbl11* loss-of-function mutants. Plants were grown under standard growth conditions for 3 weeks. Prior to freezing, the temperature was lowered to 4°C for 4 days (day and night temperature) for cold acclimation. The lowering of the temperature for freezing was done stepwise (2°C/h) and in complete darkness. The temperature for freezing was maintained at -10°C for 15 h before being gradually increased to 21°C (2°C/h). A) Representative Wt and *rbl11* plants recovered from -10°C freezing. Images were taken 7 and 14 days after freezing, from unacclimated and unfrozen (control) plants. B) Comparison of survival between Wt and *rbl11* mutants 7 days after -10°C treatment. Data represent the mean of four independent experiments with 10 to 20 plants per line and experiment. C) Quantification of wilted leaves from Wt and *rbl11* plants recovered from -10°C freezing for 7 days under standard growth conditions. Data are the mean of 4 independent experiments. Statistical differences between wild-type and overexpressor lines in B) and C) were analysed by Student's t-test followed by Welch's correction: p-value ≤0.05: *; p-value ≤0.01: ** (Supplemental file 1).

RBL11 and FAX1 interact physically at the inner envelope membrane of chloro-plasts

Having established that RBL11 activity is mandatory for cold-induced FAX1 degradation (Figure 1E), we were next interested to probe for a direct physical interaction between these two proteins. To this end, we exploited the Bimolecular Fluorescence Complementation (BiFC) analysis (Kerppola, 2008) based on the association of complementary yellow fluorescent protein (YFP) fragments fused to putative partner proteins. Only when both partners are in close proximity, a functional fluorescent protein can be formed, indicating the protein-protein interaction and its subcellular location within the living cell.

We infiltrated *Nicotiana benthamiana* leaves with the constructs FAX1::YFP^{C-n} and RBL11::YFP^{C-c} for synthesis of the respective YFP fragment proteins. Co-expression of both constructs resulted in yellow fluorescing spots in epidermis cells (Figure 3A), indicating that both proteins interact in corresponding plastids. To further visualize the complex formation of both proteins at the chloroplast envelope, we infiltrated *N. benthamiana* leaves with both constructs and subsequently isolated mesophyll protoplasts. The combined expression of FAX1::YFP^{C-n} and RBL11::YFP^{C-c} led to a yellow fluorescence of chloroplasts (Figure 3B). The ring-shaped fluorescence (Figure 3B) resembles the YFP fluorescence emitted by other inner envelope associated YFP fusion proteins (Witz et al., 2014; Patzke et al., 2019). To validate the BiFC data we performed further control experiments using inner envelope associated protease FtsH11. In contrast to the co-expression of FAX1::YFP^{C-n} and RBL11::YFP^{C-c}, the infiltration of the plasmids with FAX1::YFP^{C-n} and FTSH11::YFP^{C-c} did not yield fluorescence complementation (Figure 3C,D). In addition, the combined infiltration of the constructs FTSH11::YFP^{C-n} and RBL11::YFP^{C-c} also did not give rise to a fluorescence signal (Figure 3E,F). These two independent control experiments indicate that protein-protein interaction at the inner envelope does not occur by chance, highlighting the specificity of the RBL11 and FAX1 interaction (Figure 3A,B).

Figure 3: Probing of putative RBL11 targets. Constructs of full-length coding sequences were fused upstream to the C-terminus of yellow fluorescent protein (YFP) and transiently expressed in *Nicotiana benthamiana* leaves by agroinfiltration. YFP and chlorophyll fluorescence signals were recorded 4 days after infiltration in epidermis cells as well as in isolated protoplasts by confocal microscopy.

Induction of *RBL11* leads to a specific degradation of FAX1

In contrast to wild types, *rbl11* plants are almost unable to degrade FAX1 after the onset of cold (Figure 1E). To further support our hypothesis that RBL11 is responsible for cold-induced FAX1 degradation, we generated a stable RBL11-HA (HA = hemagglutinin) over-expressor line in which the recombinant *rbl11* gene was placed under the control of a dexamethasone (DEX)-inducible promoter (Aoyama and Chua, 1997) (Figure 4A). In the presence of DEX, the RBL11-HA protein is synthesized in mutant leaf discs at 21°C and at 4°C (Figure 4B,C), and neither DEX-induced overexpression nor the C-terminal HA tag alters the membrane localization of RBL11 (Figure 4B,C). Subsequent enrichment of total leaf membranes, followed by immunoblot analysis using the FAX1-specific antibody, showed that RBL11 overexpression does not lead to FAX1 degradation from leaf discs incubated with DEX at 22°C (Figure 4D), which might be due to some post-translational modification required for FAX1 activity. In contrast, DEX incubation at 4°C leads to FAX1 degradation (Figure 4D, note that extraction of total membrane proteins from whole membranes enriched from cold treated leaf discs generally resulted in a higher abundance of FAX1 (Figure 4D, left and right panels).

Figure 4: Cold induced degradation of FAX1 by RBL11 in Arabidopsis. A) Scheme of experimental setup. B) Immunoblot analysis via HA antibody of soluble and insoluble (membrane) protein extracts fom wildtype and transgenic plants expressing RBL11 tagged with a HA epitope (~ 32 kDa) under the control of an 35S promotor after induction with dexamethasone for 48h at 21°C. C) Immunoblot detection via HA antibody of soluble and insoluble (membrane) protein extracts fom wildtype and transgenic plants expressing RBL11 tagged with a HA epitope (~ 67 kDa) under the control of an 35S promotor after induction with dexamethasone for 48h at 4°C. Note that the size of the protein (~ 67 kDa) comes from the presence of an additional biotinylase (~35 kDa), which was not relevant in this experiment. D) Immunoblot of insoluble protein extracts from Wt and and transgenic plants overexpressing *RBL11* under the control of an 35S promotor after induction with DEX for 48h at 21°C or 4°C with a FAX1 antibody. Please note that the extraction of whole membranes enriched from cold treated leaves generally resulted in a higher abundance of FAX1 and is not compareable with results from leaves treated at 21°C. Ponceau staining in B), C) and D) is representing equal loading of protein samples with 18 µg per lane.

The membrane lipid composition of *rbl11* mutants indicates a shift to the eukaryotic biosynthesis pathway under low temperature

The abundance of FAX1 strongly decreases after transfer of wild type to cold conditions (Figure 1E and Trentmann et al. 2020). Because the absence of RBL11 prevents cold-induced FAX1 degradation (Figure 1E) we aimed to determine changes in lipid levels under these growth conditions. For this purpose, wild-type and *rbl11* mutant plants were first grown for 4 weeks under control conditions and then either, for further 14 days at 21°C or at 4°C, prior to extraction and quantification of leaf lipids.

Growth of plants at 21°C led to lower leaf levels of the major glycolipids monogalactosyl-diacylglycerol (MGDG) and digalactosyl-diacylglycerol (DGDG) in *rbl11* plants compared to wild types (Figure 5). In contrast, the levels of sulfoquinovosyl-diacylglycerol (SQDG) and the phospholipids phosphatidic acid (PA), phosphatidylserine (PS), phosphati-dylinositol (PI), phosphatidylglycerol (PG), phosphatidylethanolamine (PE) and phospha-tidylcholine (PC) were similar in leaves from the two plant lines (Figure 5). These absolute amounts summed up to 230 nmol mg^{-1} DW of phospho- and glycolipids in *rbl11* mutants and 274 nmol mg^{-1} DW of phospho- and glycolipids in wild-type plants (Figure 5A, inset). However, after recalculating the data from total levels into mol% no obvious changes in the relative individual lipid species between *rbl11* mutant and wild type plants were found (Figure 5B). Interestingly, growth for 14 days under cold temperature conditions led to higher total lipid levels in the *rbl11* plants (Figure 5C inset). This increase of total lipids in the *rbl11* mutants is due to higher levels of MGDG, DGDG and SQDG, but also to higher levels of all phospholipids (Figure 5C). Furthermore, after recalculating the data into mol% it appeared that growth at 4°C did not only lead to higher total lipid contents in *rbl11* plants, but that especially the relative proportions (in mol%) of the phospholipid species PA, PI, PE and PC to the total lipids was increased in the *rbl11* plants, while proportions of MGDG and DGDG are decreased (Figure 5D).

Figure 5. Analysis of galacto- and phospholipids in Wt and *rbl11* loss-of-function mutants. Changes in A) total contents and B) relative amounts of different galacto- and phospholipids in rosette leaves of 3-week-old Arabidopsis plants grown under standard conditions. The box plot in A) shows changes in the sum of the measured total lipid contents. Differences in C) total lipid contents and D) relative galacto- and phospholipid amounts in plants grown under standard conditions for 3 weeks before lowering the temperature to 4°C for 14 days. The box plot in C) shows the changes in the sum of the measured total lipid contents. Data represent the mean of 5 plants per row. Error bars indicate ± SD. Significance of differences between wild-type and mutant was analysed by Student's t-test: p-value ≤0.05:* (Supplemental file 1).

Identification of other putative RBL11 protease substrates

By analyzing the envelope proteome of *rbl10rbl11* double mutants an overrepresentation of non-degraded envelope proteins compared to wild type emerged (Knopf et al., 2012). However, this type of analysis did not allow for the identification of RBL11-specific substrates. Thus, we compared the envelope proteome from wild type plants with that of *rbl11* single mutants (Table 1, Suppl. Table 1, Suppl. Table 2, and Suppl. Table 3). Since we were especially interested in RBL11 substrates degraded after onset of cold temperatures, these analyses were carried out on plants grown at 21°C or at 4°C (Table 1).

Protein	Gene locus	Description	Log2FC *rbl11* / Wt	qValue_Interaction Genotype Temperature
TGD5	At1g27695	Encodes a small glycine rich protein that is localized to the chloroplast envelope and is a component of the ER to plastid lipid trafficking pathway.	4.54	0.033
PORA	At5g54190	light-dependent NADPH:protochlorophyllide oxidoreductase A	4.08	0.030
HCF107	At3g17040	Involved in regulating plastidial gene expression and biogenesis	1.56	0.002
unknown	At3g28220	TRAF-like family protein, contains MATH domain	2.73	0.044
KIN14A	At5g10470	Kinesin-like protein required for chloroplast movements and anchor to the plasma membrane	2.18	0.032
DVR	At5g18660	Encodes a protein with 3,8-divinyl protochlorophyllide a 8-vinyl reductase activity	1.85	0.031
unknown	At1g08530	Chitinase-like protein	1.35	0.019
unknown	At5g19850	Alpha/beta-Hydrolases superfamily protein	1.47	0.027
unknown	At1g64850	Calcium-binding EF hand family protein	1.22	0.015
unknown	At5g39410	Saccharopine dehydrogenase	1.06	0.018
unknown	At2g43630	Nucleus envelope protein	0.48	0.004
TGD2	At3g20320	Encodes a permease-like component of an ABC transporter involved in lipid transfer from ER to chloroplast. A phosphatidic acid-binding protein.	0.98	0.022
DEGP2	At2g47940	Encodes DegP2 protease	0.83	0.015
SPPA	At1g73990	Encodes a light-inducible chloroplast protease complex	0.78	0.015
FAX1*	At3g57280	involved in fatty acid and lipid homeostasis and likely functions as a fatty acid transporter that exports fatty acids from the plastid	0.38	0.028

Table 1. The table shows the 15 top hits of proteins that are increased by more than 30% in the mutant at low
temperature. For each protein a two-way ANOVA was performed, which means that only proteins with a significance
level below 0.05 were considered. The qValue_Interaction Genotype Temperature indicates that there is a temperature
effect that differs between mutant and wild type. All proteins were annotated by name, gene locus and description. The
proteins written in bold and in italics are increased under standard conditions as well as after 7 days at 4°C (see Suppl.
Table 1) to the described selected criteria (Abundance > 30%; qValue < 0.05). FAX1 was labeled with an asterisk
because the protein is not within the top hits, but also fulfills the selected criteria.

Putative substrates of RBL11 are expected to be more abundant in *rbl11* mutants when
compared to wild-type plants. In a previous study on proteins accumulating in the inner
envelope of *rbl10rbl11* double mutants, the TRAF-like family protein containing a MATH
domain was also found to accumulate in the double mutant (Knopf et al., 2012). Similarly,
the TRAF-like family protein accumulated in *rbl11* single mutants (Suppl. Table 1)
indicating that this protease is responsible for its degradation. Interestingly, the
abundance of the protein TGD5, which is involved in ER to plastid lipid transfer (Fan et
al., 2015), is also higher in *rbl11* mutants than in wild types, and this effect is independent
of the environmental temperature (Suppl. Table 1 and Table 1). Besides TGD5, the
abundance of another TGD protein, namely TGD2, is also higher in *rbl11* mutants when
exposed for 4 days to 4°C (Table 1). In addition, the chloroplast located protease systems
DegP2 and SPPA are also more abundant in cold treated *rbl11* mutants when compared
to wild type (Table 1). Similarly, the inner envelope associated NADPH:protochloro-
phyllide reductase PORA (Barthélemy et al., 2000) appears, besides the TGD
components and FAX1, as a further RBL11 substrate (Suppl. Table 1), especially after
onset of cold temperature (Table 1). It seems likely that the decrease of some proteins in
rbl11 chloroplasts under warm or cold conditions represents pleiotropic reactions of the
mutant induced by altered FAX1 abundance (Suppl. Table 2 and Suppl. Table 3).
Nevertheless, among these proteins, we also found proteins associated with lipid
metabolism, e.g., LACS9, representing a major long chain acyl-CoA synthetase (Schnurr
et al., 2002), or BASS1, representing a sodium/pyruvate cotransporter (Furumoto et al.,
2011) (Suppl. Table 2 and Suppl. Table 3).

FAX1 overexpressor lines exhibit markedly increased levels of FAX1 protein

The above data indicate a molecular interrelation between increased FAX1 protein abundance (due to an absence of RBL11) and the levels of membrane lipids and Arabidopsis cold tolerance (Figures 1 and 3). This encouraged us to analyze the impact of increased FAX1 abundance on plant cold tolerance. _rbl11_ mutants are not conducive to this, because the absence of this protease does not exclusively affect FAX1 levels, but also the abundance of other envelope associated proteins (Table 1). To avoid the study of pleiotropic effects, we decided to investigate the cold tolerance of Arabidopsis mutants with constitutively increased FAX1 levels (Li et al., 2015).

Our observations that _rbl11_ loss-of-function mutants have increased FAX1 levels under cold conditions (Figure 1E) and that directed synthesis of RBL11 leads to a decrease in FAX1 in the cold (Figure 2) provide evidence that the RBL11 protease is responsible for this process. Nevertheless, for a comprehensive understanding of FAX1's impact on the cold response, it is mandatory to study _FAX1_ transcript levels after transfer to cold conditions. In fact, the observation that _FAX1_ mRNA rises markedly in wild-type plants after exposure to 4°C (Figure 6A) supports our conclusion that RBL11 is responsible for the cold-induced decrease of the level of the FAX1 protein (Figure 1F).

Previously, we generated two independent FAX1 overexpressing lines, FAX1 Ox#2 and FAX1 Ox#4, both containing the _FAX1_ gene under the control of the constitutive cauliflower 35S promotor (Li et al., 2015). So far, it has only been shown that these mutant lines have both, increased _FAX1_ mRNA and increased FAX1 protein levels when grown at ambient temperature (Li et al., 2015). We verified that the _FAX1_ mRNA accumulation is a stable feature when FAX1 overexpressor mutants are cultivated at cold temperature. For this purpose, the _FAX1_ mRNA levels were quantified in plants grown under the standard temperature of 21°C with those grown for one week at 4°C. The two FAX1 overexpressor lines exhibited a substantial accumulation of _FAX1_ mRNA when compared to wild-type plants. Under warm temperature, FAX1 Ox#2 plants contained about 2.1-fold higher _FAX1_ mRNA levels as present in the wild type, while FAX1 Ox#4 plants contained even 1170-fold higher _FAX1_ mRNA (Figure 6B). This marked difference of _FAX1_ mRNA in the two overexpressors concurs with previous observations (Li et al., 2015). After one week of growth at 4°C, FAX1 Ox#2 plants contained about 5.7-fold higher _FAX1_ mRNA

levels, and FAX1 Ox#4 plants contained more than 970-fold higher *FAX1* mRNA levels

than present in the wild type (Figure 6C). Thus, even at 4°C, both FAX1 overexpressor

lines exhibited higher *FAX1* mRNA than present in the wild type.

It was shown that both FAX1 overexpressors contain higher levels of FAX1 protein when

grown at 21°C (Li et al., 2015). To confirm that also under cold conditions, FAX1

overexpressor lines exhibit higher FAX1 protein levels than present in wild types, we

conducted a immunoblot analysis (Li et al., 2015). After one week of growth at 4°C, the

relative FAX1 protein level in both overexpressor plants was higher than observed in wild

type plants (Figure 6D), and a corresponding quantification experiment revealed that

Ox#2 plants contained about 3-fold more FAX1 protein than present in wild type, while

Ox#4 plants contained about 115-fold more FAX1 protein (Figure 6D, please note: loaded

protein extracted from Ox#4 plants was 1:10 diluted when compared to proteins extracted

from wild types and Ox#2 mutants).

Figure 6: Gene expression, immunoblot and phenotypic analysis of two independent Arabidopsis
fatty acid export protein 1 (FAX1) overexpression lines (FAX1 Ox#2 and FAX1 Ox#4) and wild-
type (Wt) plants grown under standard and cold (4°C) conditions. Plants were grown under
standard conditions (21°C day and night, 10h day length and 120µE light intensity) for 3 weeks
and then treated with cold (4°C day and night, 10h day length and 120µE light intensity). A)
Expression of FAX1 by qRT-PCR under standard growth conditions (0 days at 4°C) and several
days during cold treatment (7; 14; 42, and 70 days at 4°C). Data represent relative mean
expression levels of 3 biological replicates and are normalised to standard conditions (0 days at
4°C) using UBQ as an internal control. Relative expression of FAX1 by qRT-PCR under B)
standard growth conditions and C) 7 days after cold treatment. Data represent relative mean
expression levels and are normalised to the wild type using UBQ as an internal control. D)
Immunoblot analysis of FAX1 in crude extract from plants grown at 4°C for 7 days. The image of
the Coomassie stained gel (CBB) shows equal loading of protein samples from Wt and FAX1
Ox#2 with 30µg per lane. The amount of FAX1 Ox#4 protein is reduced to 3µg. E) Rosette
phenotype of representative Wt, FAX1 Ox#2 and FAX1 Ox#4 under standard growth conditions
(0 days at 4°C) and after cold treatment (7; 42, and 70 days at 4°C). F) Number of wilted leaves
per plant after 42 and 70 days of cold treatment. Error bars in A) represent ± SEM. Error bars in
B), C) and F) are ± SD. Statistical differences between wild-type and overexpressor lines in A), B)
and C) were analysed by Student's t-test: p-value ≤0.05: *; p-value ≤0.01: **; p-value ≤0.001: ***.
Letters above error bars in F) indicate significant differences by two-way ANOVA followed by
Tukey's test (p<0.05; (Supplemental file 1)).
490

Similar to *rbl11* mutants, FAX1 overexpressor plants exhibit decreased chilling and
frost tolerance

The data above indicate that *rbl11* mutants exhibit increased cold sensitivity (Figure 1B,
C). Because *rbl11* mutants revealed higher FAX1 levels than present in wild types it was
interesting to analyze whether the two lines constitutively expressing the *FAX1* gene also
show altered response to low temperatures.

Therefore, we cultivated all plants at ambient temperature (21°C) prior to cultivation at
4°C. Subsequently, we inspected morphologic plant appearance and the number of wilted
leaves in the respective growth phase. After 7 days at 4°C none of the three plant lines
developed wilted leaves (Figure 6E). However, after 42 days at 4°C, and even more
pronounced after 70 days at 4°C, leaves from both FAX1 overexpressors gradually wilted
more rapidly than observed in the wild type (Figure 6E).

To quantify this observation we counted the wilted leaves as a proxy for chilling sensitivity
(Trentmann et al., 2020). After 42 days at 4°C wild type plants exhibited on average one
wilted leaf while FAX1 overexpressor Ox#2 exhibited in average three wilted leaves, and

Ox#4 mutants showed in average 4.5 wilted leaves (Figure 6F). After growth for ten weeks
at 4°C, wild type plants displayed on average 2.8 wilted leaves per plant while FAX1
overexpressors Ox#2 and Ox#4 exhibited on average 7.5 and 10.4 wilted leaves,
respectively (Figure 6F).

As done for *rbl11* plants (Figure 2), we compared the frost tolerance of FAX1 over-
expressor plants and wild types. In contrast to both FAX1 overexpressors, wild types
which survived the post-freezing phase restored efficient growth within the next 14 days,
as indicated by a larger rosette size and less wilted leaves when compared to over-
expressor mutants (Figure 7A). FAX1 overexpressor plants able to recover from freezing
exhibited about 80% of wilted leaves, while wild types only exhibited about 46% of wilted
leaves (Figure 7A, B). With about 87% the survival rate of wild types reached a value
similar to previous observations (Trentmann et al., 2020). In contrast, only 57% of the
FAX1 Ox#2 plants survived this stress, and only 43% of FAX1 Ox#4 plants recovered
from frost (Figure 7C).

Figure 7: Recovery from frost is impaired in FAX1 overexpression mutants. Plants were cultivated for 3 weeks under standard growing conditions. Before freezing the temperature was lowered to 4°C for 4 days (day and night temperature) for cold acclimation. Lowering of the temperature for freezing was done stepwise (2°C/ h) and in completely dark. -10°C was kept for 15 h before a stepwise temperature raising to 21°C (2°C/h). A) representative Wt and FAX1 overexpression plants recovered from -10°C freezing. Pictures were taken 7 and 14 days after freezing, from

unacclimated and unfrozen (control) plants. B) Comparison of survival rate between Wt and FAX1
overexpression mutants 7 days after -10°C treatment. Data represent the mean value from four
independent experiments with 11 to 15 plants per line and experiment. C) Quantification of wilted
leaves from Wt and FAX1 overexpressor plants recovered for 7 days from -10°C freezing under
standard growing conditions. Data represent the mean value of 4 independent experiments.
Statistical differences between wildtype and the overexpressor lines in B) and C) was analyzed
by Student's t-test: p-value ≤0.01: **; p-value ≤0.001: *** (Supplemental File 1).

Similar to *rbl11* mutants, lipid biosynthesis in FAX1 overexpressors is shifted to the

eukaryotic biosynthesis pathway under low temperature

Under cold conditions, *rbl11* mutants exhibited a shift towards ER-synthesized membrane

lipids when compared to their corresponding wild type (Figure 5). To check whether this

response is due to high FAX1 protein abundance, we analyzed the lipid composition in

FAX1 overexpressors. For a first overview of cold effects on membrane lipid homeostasis

in FAX1 overexpressor plants, we grew wild types and the representative FAX1 Ox#4 line

(Li et al., 2015) under either control conditions, or for two weeks at 4°C. Subsequently,

leaf lipids were extracted and quantified via mass-spectrometry.

When grown at 21°C, the total levels (nmol mg^{-1} DW) and the relative proportions (mol%)

of the glycolipids MGDG, DGDG and SQDG, or of the phospholipids PA, PS, PI, PG, PE

and PC were similar in leaves from wild types and FAX1 Ox#4 plants (Supplemental

Figure 1A, B and insert). In contrast, after two weeks at 4°C FAX1 overexpressor plants

contained higher levels of MGDG and DGDG when compared to wild types (Supplemental

Figure 1C). In addition, the levels of PS-, PG-, PE- and PC FAX1 Ox#4 plants appeared

to be slightly higher than in corresponding wild types (Supplemental Figure 1C). This

increase led in sum to 158 nmol mg^{-1} DW of membrane lipids in FAX1 Ox#4 plants, while

wild types contained only 125 nmol mg^{-1} DW membrane lipids (insert Supplemental Figure

1C). Similar to the situation at 21°C, the proportions of all lipid species in both plant lines

are nearly identical (Supplemental Figure 1D).

Since FAX1 overexpressor plants revealed impaired cold tolerance after long exposure to

low temperatures, and because the observed changes in lipid composition of FAX1

overexpressors after two weeks of growth at cold temperature appeared in parts to be

moderate (Supplemental Figure 1), we extracted lipids from plants exposed for ten weeks

to 4°C (Figure 8). Similar to plants exposed to only two weeks at 4°C (Supplemental Figure

1), both FAX1 overexpressor lines exhibited increased total levels of the galactolipids

MGDG and DGDG, and also of the phospholipids PI, PG, PE and PC (Figure 8A). The

increase of the two groups of lipids led to an overall higher level of total membrane lipids

in FAX1 overexpressor lines (FAX1 Ox#2, 205 nmol mg^{-1} DW; FAX1 Ox#4, 188 nmol mg^{-}

^{1} DW; wild type, 125 nmol mg^{-1} DW; (insert Figure 8A). The levels of the low abundant

glycolipid SQDG and the phospholipid PS were not altered in FAX1 overexpressors, when

compared to wild type (Figure 8A). Although total amounts of glycolipids in both FAX1

overexpressors were higher when compared to wild types (Figure 8A), the relative

contribution (mol %) of MGDG, DGDG and SQDG in FAX1 Ox#2 and #4 to total

membrane lipids was similar to wild type (Figure 8B), while the relative contribution of the

phospholipids PG, PE and PC in FAX1 Ox#2 and #4 to total membrane lipids was

increased, when compared to the wild type (Figure 8B). The most pronounced alterations

of phospholipids were noted for PC and PE (Figure 8B), which were 40% higher in FAX1

overexpressors than in wild type plants (Figure 8B).

MGDG and DGDG represent the two most abundant glycerolipids in Arabidopsis leaves

(Figure 8A). Interestingly, a closer inspection of the contents of the two MGDG molecular

species 34:6 and 36:6 – which are indicative for either plastid-generated MGDG (34:6) or

ER-born MGDG (36:6) - revealed clear differences between wild types and

overexpressors. Wild-type plants exhibited about 85 mol% of 34:6 MGDG, while leaves

from FAX1 Ox#2 and Ox#4 plants accumulated only 75 and 62 mol% of this MGDG

species, respectively (Figure 8C). In marked contrast to this, wild type contained only 10

mol% of the MGDG molecular species 36:6, while the FAX1 overexpressor plants Ox#2

and Ox#4 accumulated 21 and 30 mol% of 36:6 MGDG, respectively (Figure 8C). The

relative levels of 34:6 and 34:3 DGDG in wild type leaves were higher than in

correspondingly grown FAX1 Ox#2 and Ox#4 plants. 34:6 DGDG in wild type amounted

at 6.5 mol%, while FAX1 Ox#2 and Ox#4 plants contained only 5.6 or 4.6 mol%,

respectively (Figure 8D). Eukaryotic ER-produced 36:6 DGDG in wild types amounted to

a relative abundance of 61 mol%, while, similar to the increase of 36:6 MGDG, the two

FAX1 overexpressors contained 75 and 74 mol% of 36:6 type DGDG, respectively (Figure

8D). In summary, we can conclude that FAX1 overexpressor lines at cold temperatures

accumulate ER-produced phospholipids – namely PC and PE – as well as galactolipids

(MGDG, DGDG) with DAG backbones from the eukaryotic pathway, while the proportions

of prokaryotic 34:x galactolipids are reduced.

Figure 8: Lipid analysis of Wt and FAX1 overexpression lines. Galacto- and phospholipids as well
as galactolipid molecular species were determined in leaves of plants grown for 3 weeks under
standard conditions before lowering the growth temperature to 4°C for 10 weeks. Changes in A)
total contents and B) relative amounts of different galacto- and phospholipids. Boxplot in A)
indicates changes in the sum of measured total lipid contents. Lipid composition of C)
Monogalactosyldiacylglycerol (MGDG) and D) Digalactosyldiacylglycerol (DGDG) molecular
species. Data represent mean values of 5 plants per line. Error bars represent ± SD. Significance
of differences between wildtype and mutant lines was analyzed by Student's t-test: p-value ≤0.05:*
(Supplemental File 1).

FAX1 overexpressor lines show symptoms of impaired photosynthesis at early time

points of cold exposure

One marked phenotype of FAX1 overexpressor plants after transfer to cold conditions is

the appearance of wilted, decayed leaves after some weeks of growth (Figure 6F, H).

However, degradation of the FAX1 protein and first changes of the lipid composition are

already observed after a few days at 4°C (Figure 1F and Supplemental Figure 1). To

search for additional responses, we quantified photosynthetic parameters after short and

longer exposure to 4°C. To this end we grew all plants for 28 days at 21°C prior to transfer

to at 4°C. At the beginning of the transfer, and after one or six weeks of growth at 4°C, we

quantified the photosynthetic performance by measuring the following parameters: the

ratio of variable- to maximal fluorescence (F_v/F_m), PSII efficiency Y(II), non-photochemical

quenching Y(NPQ), non-regulated quenching Y(NO), and the rate of electron transport

(ETR). This comprehensive analyses have been done using the pulse-amplitude-

modulation (PAM) fluorometry method (Schreiber, 2004).

Prior to transfer to the cold, all wild type and the two FAX1 overexpressor plants exhibited

similar F_v/F_m ratios (Figure 9A) which ranged around 0.79 (Figure 9B). After onset of cold

temperatures, the F_v/F_m ratio decreased gradually in all three lines. Already after one

week, the F_v/F_m ratio in both FAX1 overexpressor lines was significantly lower than

displayed by the wild type, and after six weeks in the cold, wild type plants showed a F_v/F_m

ratio of 0.73, while both mutants exhibited a F_v/F_m ratio of about 0.68 (Figure 9A, B). This

altered F_v/F_m ratio in the cold is not reflected by an increased NPQ, although FAX1 Ox#2

plants showed a slightly increased NPQ, after 7 days in the cold. However, this was not

found for FAX1 Ox#4 plants (Figure 9) and even after 5 weeks in the cold, NPQ in all three

lines was similarly decreased (Figure 9C). The photosynthetic quantum yield Y(II) of all

three plant lines was similar when grown under control temperature (Figure 9C). In

contrast, both FAX overexpressors showed after 7 or 42 days in the cold a decreased

Y(II) and an increased energy dissipation via non-regulated quenching Y(NO), which was

especially pronounced after 42 days (Figure 9C). While the chloroplastic electron

transport rate (ETR) of all three plant lines was similar at the beginning of the cold

treatment, the ETR in both FAX1 overexpressor plants was after 7 days at 4°C markedly

lower, when compared to corresponding wild type plants (Figure 9C).

Figure 9: Cold-dependent PSII alterations in FAX1 over expressor lines. After 28 days of growth under short-day (10/14 hours; 110 PAR) conditions at RT, plants (Col-0; FAX1 Oex#2; FAX1 Oex#4) were shifted to a 4°C short-day (110 PAR) chamber. Pulse-Amplitude-Modulation (PAM) induction curve measurements at 110 PAR were performed on day 0 as well as after 7 days and 42 days of the cold treatment. A) Representative PAM images are depicted for PSII capacity (F_v/F_m), yield (Y(II)), the quantum yield of light-induced non-photochemical fluorescence quenching (Y(NPQ)), and quantum yield of nonregulated energy dissipation (Y(NO)). B) F_v/F_m determination of the three genotypes at the start of the cold treatment, after one and five weeks. C) Induction curves were generated until a steady-state phase of Y(II), Y(NPQ); Y(NO), and electron transport rate (ETR [µmol m^{-2} s^{-1}] were reached. The data shown represents this steady-state after 600 seconds. N = 16; Mean; ± SEM; p-value: (one-way ANOVA): * = 0.033; ** = 0.0021; *** = 0.0002; **** = 0.0001 (Supplemental File 1).

FAX1 overexpressor and *rbl11* mutants exhibit signs of reactive oxygen species (ROS) accumulation

Several systemic changes in FAX1 overexpressors and *rbl11* mutants occur quite rapidly after transfer to cold conditions. For example, first changes in the lipid composition of both types of mutants are already present after two weeks at 4°C (Figure 5 and Supplemental Figure 1) and alterations in photosynthetic parameters of FAX1 overexpressors are already present after one week of growth at low temperature (Fig. 7). Given that both types of mutants contain higher FAX1 abundance than present in wild types (Figures 1 and 4) and show similar shifts in lipid biosynthesis (Figure 5 and Supplemental Figure 1) we searched for further similarities in their molecular responses after transfer into cold conditions.

The accumulation of both, anthocyanins and reactive oxygen species (ROS) can be taken as early responses upon onset of abiotic stress stimuli (Chalker-Scott, 1999; Baxter et al., 2014). When grown at 21°C, *rbl11* plants and the two FAX1 overexpressor lines contain similar levels of anthocyanins as wild type plants (Figure 10A). In contrast, when exposed to 4°C for only one week, *rbl11* mutants and FAX1 overexpressors accumulated significantly more anthocyanins than the correspondingly grown wild types (Figure 10A). The levels of the transcript coding for the protein BAP1, which indicates the cold-induced occurrence of ROS (Yang et al., 2007; Zhu et al., 2011), was unchanged between the plant lines when grown at 21°C (Figure 10B). Similar to the anthocyanin accumulation, *BAP1* mRNA accumulated to much higher extents in both types of mutants after one week of growth at 4°C, than in wild type plants (Figure 10B).

To corroborate the differences in cold-induced ROS levels in wild types and the two types of mutants, we analyzed the relative increase in leaf superoxide by nitroblue-tetrazolium (NBT) staining (Doke, 1983; Hoffmann et al., 2013). NBT staining of wild-type tissue at the start of cold treatment and after four days at 4°C does not result in a detectable increase in superoxide (Figure 10C). However, both types of mutants showed a stronger NBT staining after four days at 4°C than at the beginning of cold treatment (Figure 10C).

Figure 10: Anthocyanin and reactive oxygen species (ROS) accumulation in *rbl11* loss of function mutants and FAX1 overexpressors at low temperatures. A) Impact of cold on anthocyanin accumulation under ambient conditions and after 7 days at 4°C. B) Relative transcript levels of *BAP1* during standard conditions and after 7 days at 4°C. C) NBT staining of O_2- accumulation. Plants were grown for 2 weeks under control conditions and subsequently cultivated at 4°C for 4 days. Error bars in A) and B) represent ± SD. Letters displayed over the error bars indicate significant differences analyzed by two-way ANOVA followed by Tukey's multiple comparisons test (p-value <0.05; (Supplemental File 1)).

690 **FAX1 overexpressor plants exhibit altered levels of metabolites when exposed to**

691 **cold temperatures**

692 To gain additional insight into metabolic causes for the impaired cold- and frost tolerance

693 pattern of FAX1 overexpressor lines, we analyzed the levels of 71 primary metabolites via

694 GC-MS. Thus, wild-type plants and both FAX1 overexpressor lines were grown for 28

695 days at 21°C prior to growth at 4°C for two weeks.

696 When cultivated at 4°C for two weeks, the FAX1 overexpressor lines showed the largest

697 differences in their metabolic readjustment compared to correspondingly grown wild-type

698 plants (Figure 11, Supplemental Table 4). The two FAX1 overexpressor lines contained

699 nearly doubled levels of β-alanine, asparagine, serine and ethanolamine, 1.7-fold

700 increased amounts of glycerol-phosphoglycerol and 1.7-fold more malate when compared

701 to wild types (Figure 11). Similar to malate, three further intermediates of the tricarboxylic

702 acid cycle, namely citrate, fumarate and succinate were also significantly higher in cold

703 treated FAX1 overexpressors than in wild type plants (Figure 11). In addition, cold treated

704 FAX1 overexpressors contained about 38% more proline, about twice as much putrescine

705 and 2.3-fold more spermidine when compared to corresponding wild types, whereas the

706 trehalose concentration in FAX1 Ox#2 plants was substantially lower, compared to wild

707 types (Figure 11).

708

Figure 11: Volcano plot of the metabolic differences measured between Wt and FAX1 Ox#2
rosette leaves, cold treated for 14 days at 4°C. Blue dots represent decreased and red dots
increased metabolites with a $\log_2$ fold change ≥ 0.5 and p-value ≤ 0.05. The complete data set is
available in Supplemental Table 4.
Significant differences between the 21°C and 4°C treatments were analyzed using a t-test
(Supplemental File 1).

Discussion

The dynamic modification of plant organelle proteomes is mandatory to achieve new homeostatic levels allowing to cope with challenging environmental conditions (Taylor et al., 2009). As seen in many systems, cellular mRNA level and the respective protein amounts do not necessarily correlate to a high degree (Gygi et al., 1999; Greenbaum et al., 2003; Koussounadis et al., 2015). Thus, other factors are also important for controlling protein abundance. The chloroplast envelope proteome undergoes substantial remodeling in response to changes in light or temperature conditions (Knopf et al., 2012; Nishimura et al., 2016; Adam et al., 2019; Mielke et al., 2020) and it can thus be expected that proteases must play a role for this.

Interestingly, cold-induced changes in the proteome are accompanied by modifications of membrane lipids. Latter process maintains membrane fluidity at low temperatures and stabilizes membrane integrity (preventing rigidification) to ensure proper organelle function (Moellering et al., 2010; Zheng et al., 2011; Barnes et al., 2016; Barrero-Sicilia et al., 2017; Guo et al., 2018). For lipid biosynthesis, especially for the generation of phospholipids, fatty acid export from the chloroplast is mandatory. and the FAX1 protein is the best characterized chloroplast envelope located protein presumed to be involved in this transport process so far (Li et al., 2015; Li et al., 2016; Xiao et al., 2021).

In Arabidopsis FAX1 abundance rapidly decreases after transfer to cold temperatures while other envelope proteins increase (Trentmann et al., 2020). Taking the proteins NTT (the chloroplast ATP importer (Tjaden et al., 1998) or MEX1 (the chloroplast maltose exporter) (Niittylä et al., 2004) as examples, we demonstrated that the relative changes of these carriers are key to proper cold tolerance (Trentmann et al., 2020). However, in the case of FAX1, it is unknown whether the decreased protein abundance is a controlled process required to tolerate low environmental temperatures. In addition, no protease has been discovered that could mechanistically explain the decreased abundance of selected envelope-associated proteins in the cold.

Compared to *RBL10* and *FtsH11* mRNAs, the *RBL11* transcript accumulates early after cold exposure (Figure 1A). This correlation is a first indication of a specific molecular interaction between RBL11 and FAX1, and a physical contact between these two proteins,

which is a prerequisite for FAX1 degradation, was confirmed by BiFC analysis (Figure 2).

The suggestion of a specific effect of RBL11 on FAX1 abundance in the cold is supported

by two observations. First, cold-induced FAX1 degradation did not occur in *rbl11* mutants

(Figure 1E), whereas *rbl10* or *ftsh11* mutants degraded FAX1 in the cold similarly to wild-

type plants (Figure 1E). Second, dexamethasone-induced expression of RBL11 leads to

a decrease in FAX1 protein (Figure 4C).

In general, the ability of RBL11 to degrade intrinsic membrane proteins with multiple

transmembrane domains, such as FAX1 (Li et al., 2015), is consistent with the properties

of other rhomboid proteases (Erez and Bibi, 2009). However, since dexamethasone-

induced expression of RBL11 leads to FAX1 degradation only at cold temperature and

not at 21°C (Figure 4C), we propose a so far unknown post-translational modification of

RBL11 and/or FAX1, which is a prerequisite for catalytic protease activity (Figure 4C). The

observation that RBL11 activity might also be involved in the dynamic change TGD5 and

TGD2 (Table 1), two components of the TGD complex involved in the unidirectional ER

to plastids import of eukaryotic lipids (Xu et al., 2010; Li-Beisson et al., 2017), point to a

central function of this protease in modification of plant lipid homeostasis under

challenging temperature conditions. Notably, RBL10, the closest RBL11 homologue, has

also been shown to affect lipid metabolism as it interacts with the ACYL CARRIER

Protein4 and modulates MGDG biosynthesis (Lavell et al., 2019; Xu et al., 2023). Because

rbl11 mutants exhibit a shift toward eukaryotic lipid biosynthesis (which is due to

decreased FAX1 abundance but not observed in wild types, Figures 3 and 6,

Supplemental Figure 1), we hypothesize that *rbl11* cells attempt to reduce the unintended

stimulation of eukaryotic lipid biosynthesis by downregulating the core plastid lipid

importer TGD (Table 1). This assumption is supported by the observation that Arabi-

dopsis, as a 16:3 plant, normally stimulates the plastid membrane lipid pathway after

exposure to cold conditions (Li et al., 2015; Yu et al., 2023).

Apart from FAX1 and TGD components, which act as substrates for RBL11, it is worth

mentioning that the protochlorophyllide oxidoreductase PORA strongly accumulates in

rbl11 mutants (Table 1). PORA is responsible for the stromal conversion of

protochlorophyllide to chlorophyllide and it was shown, that increased protochlorophyllide

oxidoreductase activity leads to ROS formation (Pattanayak and Tripathy, 2011). Thus,

RBL11 might not only contribute to the regulation of FAX1 activity but also influence
envelope-located mechanisms preventing cold-induced ROS formation.

rbl11 mutants exhibit in cold conditions both, higher FAX1 protein levels than observed in
wild types (Figure 1E) and a shift of their lipid composition towards accumulation of
phospholipids (Figure 5). The latter observation is in accordance with both, the function
of FAX1 as a chloroplast to ER fatty acid (FA) export protein (Li et al., 2015; Li et al., 2016;
Li-Beisson et al., 2017; Takemura et al., 2019; Tian et al., 2019) and the function of the
ER as the cellular site of phospholipid biosynthesis (Li et al., 2015; Li et al., 2016; Li-
Beisson et al., 2017; Takemura et al., 2019; Tian et al., 2019). In fact, apart from a
stimulation of the eukaryotic pathway of lipid biosynthesis in the cold, *rbl11* plants and
FAX1 overexpressors share further similarities, e.g., impaired tolerance to low
temperatures and frost (Figures 1, 2 and 7), and increased levels of anthocyanins, ROS
and *BAP1* transcripts in the cold when compared to wild types (Figure 10A-C). Latter
changes are independent molecular markers for a reinforced stress situation in mutants
(Chalker-Scott, 1999; Choudhury et al., 2017)

Although in sum these similarities are indicative for an important function of the down
regulation of FAX1 during cold tolerance, a detailed analysis of the impact of FAX1 on this
process cannot be made in the *rbl11* mutants, since RBL11 acts on several inner envelope
proteins, which triggers pleiotropic effects in *rbl11* plants (Supplemental Tables 1 and 3,
and Table 1). Thus, the impact of FAX1 on cold and frost tolerance was analyzed using
FAX1 overexpressors. Although the relative abundancies of *FAX1* mRNA and protein
differed substantially between Ox#2 and Ox#4 mutants, it turned out that the physiological
responses of the two plant lines were very similar (Li et al., 2015 and below). Accordingly,
these lines are suitable to search for a potential impact of a cold-induced down regulation
of FAX1 levels for tolerance to low temperatures.

The observation that FAX1 overexpressor leaves exhibited higher levels of ER-derived
membrane lipids (PE, PC) when exposed to cold temperature (Figure 8 and Supplemental
Figure 1B) indicates that the cold-induced decrease of FAX1 in wild types (Trentmann et
al., 2020) is a limiting factor for synthesis of lipid backbones via the eukaryotic pathway at
low temperature. Our assumption that a relative stimulation of lipid biosynthesis in the ER
is causative for increased total levels of galacto- and phospholipids in FAX1

overexpressors gains independent support by observations made on mutants lacking the

envelope proteins TGD2 or TGD3. Corresponding loss-of-function mutants show

decreased levels of MGDG and DGDG with ER-derived DAG-backbones, while ER-borne

phospholipid levels are increased (Awai et al., 2006; Lu et al., 2007). In other words, a

higher substrate availability such as acyl residues at the ER might lead to increased levels

of membrane lipids with ER-derived DAG backbones (Figure 12).

MGDG represents the most abundant chloroplast-located galactolipid (Dorne et al., 1990;

Kobayashi, 2016) and in the 16:3 plant Arabidopsis, two main molecular species occur

i.e., plastid derived 34:6 type MGDG and, as ER derived, 36:6 type MGDG (Boudière et

al., 2014). When compared to wild types, Ox#4 mutants grown at 21°C exhibit a relative

decrease of 34:6 type MGDG, which is nearly balanced by a corresponding increased

level of 36:6 type MGDG (Figure 8C), and the proportion of 34:6 DGDG is decreased,

while 36:6 DGDG is increased when comparing wild type with Ox#4 plants grown at 4°C

(Figure 8D). Thus, the latter observation indicates that a stimulated fatty acid export in

Ox#2 and Ox#4 plants leads to a shift from plastid-derived to ER-derived galactolipids.

Such flexible shift of lipid biosynthesis from plastids to the ER has already been observed

for mutants lacking the plastid glycerol-3-phosphate acyltransferase (ACT) activity

(Falcone et al., 2004; Lusk et al., 2022), and also in *act1* mutants (synonymous: *ats1*) a

stimulation of membrane lipid biosynthesis in the ER largely compensates for the impaired

prokaryotic lipid biosynthesis (Kunst et al., 1988). The observation that FAX1

overexpressors not only exhibit an increased ratio of 36:6 to 34:6 type MGDG (Figure 8C),

but also increased ratios of 36:6 to 34:6-, and 36:6 to 34:3 type DGDGs (Figure 8D) further

underlines the shift from prokaryotic to eukaryotic lipid biosynthesis in these lines. Finally,

the increase of the relative proportions of the two most abundant phospholipids, PC and

PE, supports the conclusion that the eukaryotic lipid biosynthesis pathway is stimulated

in FAX1 overexpressors.

The analyses of the molecular responses of the FAX1 mutants to cold and freezing

conditions showed that these changes are associated with a strongly impaired ability to

resist low environmental temperatures, as evidenced by an increased number of wilted

leaves in the cold and a decreased ability to recover from freezing (Figures 2 and 7). A

similar gradual decay of leaves was also seen for other Arabidopsis mutants exhibiting

decreased amounts of polyunsaturated fatty acids in membrane lipids (Miquel et al., 1993)

and of course for the *rbl11* mutant (Figure 1). It seems likely, that the increased number

of wilted leaves of FAX1 overexpressors when grown at 4°C (Figure 6E and F) is the result

of an impaired photosynthetic performance (Figure 9A,C).

The exact reasons for these marked effects are not clear, but it was shown that changes

in the composition of membrane lipids, induced by modifications of different lipid

biosynthesis genes, affect photosynthetic properties (Botté et al., 2011; Kobayashi, 2016;

Gao et al., 2020). Thus, it seems conceivable that the altered chloroplast membrane

composition observed in FAX1 overexpressors (Supplemental Figure 1 and Figure 8) is

causative for the impaired photosynthetic performance. Indeed, the decreased F_v/F_m ratio

of FAX1 overexpressor plants (Figure 9A,B) is an indicator of impaired PSII function

(Murchie and Lawson, 2013) and a temperature stressed PSII can lead to ROS production

(Pospíšil, 2016), as observed in FAX1 overexpressor plants (Figure 10C-F). However, it

cannot be excluded that the marked effects of RBL11 mutation or FAX1 overexpression

on PC and PA levels (Figures 5C,D and 8A,B) also contribute to ROS accumulation

(Figure 10). This is because the signaling molecule PA has been shown to bind directly

to the plasma membrane NADPH oxidoreductase RBOHD, leading to the activation of this

enzyme. Accordingly, this process stimulates O_2^- production, which ultimately leads to

H_2O_2 accumulation (Zhang et al., 2009).

Wild-type plants and FAX1 overexpressor lines exhibit a quite similar metabolite pattern

when grown at 21°C. In contrast, the metabolite composition of FAX1 Ox plants during

growth at 4°C is to some degree indicative for reinforced cold stress. For example, the

comparably high accumulation of polyamines in form of putrescine and spermidine, of

amino acids like proline and asparagine, or the accumulation of the non-proteinogenic

amino acid β-alanine (Figure 11) represent independent evidences for a pronounced

metabolic response to cold temperatures (Alet et al., 2011; Liang et al., 2013; Zhang et

al., 2016; Marco et al., 2019; Parthasarathy et al., 2019). β-alanine is not only a metabolic

indicator for abiotic stress (Parthasarathy et al., 2019), it also acts as precursor for CoA

synthesis and is thus required for fatty acid and phospholipid biosynthesis (Perrett et al.,

2017). Therefore, β-alanine accumulation is coincident with higher total levels of galacto-

and phospholipids in FAX1 overexpressors in the cold (Figure 11A). Under many stress

conditions, asparagine and proline levels change similarly (Curtis et al., 2018). While the

exact function of asparagine during stress is unclear, the role of proline as a general stress

metabolite is well established (Liang et al., 2013; Ghosh et al., 2022). For example, proline

is able to diminish PS_{II} defects caused by rising ROS levels (Alia and Mohanty, 1997) and

because cold treated FAX1 overexpressors exhibit defects in PS_{II} activity (Figure 9A-C),

the induce proline accumulation might represent a process to tune down such negative

effects.

In summary, we propose a model in which the envelope protease RBL11 is responsible

for the degradation and down-regulation of FAX1 in cold-treated Arabidopsis plants

(Figure 12). This process represents a previously hidden molecular response that is

critical for optimal low temperature acclimation. Most likely, the initial cold response in

wild-type plants is represented by a preference for the prokaryotic lipid synthesis pathway

to protect chloroplast membranes and the photosynthetic machinery from cold damage.

Since the corresponding *FAX1* mRNA is not decreased in cold-treated wild-type plants

(Figure 6A), the decreased FAX1 protein level is most likely due to a specific effect of

RBL11. In near future it will be interesting to search for factors that activate RBL11 and/or

convert selected membrane proteins on the inner envelope proteins into specific

substrates (Figure 12). It seems worth to mention, that the cold-induced degradation of

FAX1 further supports findings on the importance of various chloroplast envelope

associated processes for plant cold and frost tolerance (Moellering et al., 2010; Barnes et

al., 2016; Guan et al., 2019; Schwenkert et al., 2023).

Figure 12: Proposed model for the influence of the fatty acid export protein1 (FAX1) abundance,
controlled by the envelope located rhomboid-like protease11 (RBL11), on cold adaptation via
balancing the glyco- and phospholipid contents in Arabidopsis.
In wild types, the the glyco- and phospholipid levels are balanced to gain ideal conditions for
growth and development under diverse environmental conditions. During low temperature, RBL11
interacts with FAX1, which leads to a decrease of FAX1 abundance. The associated reduced
export of fatty acids, and concomitted slowdown of the eukaryotic pathway for lipid biosynthesis,
seems to be an efficient mechanism for cold acclimation. In FAX1 overexpressing (Ox) plants
however, the permanent increased export of fatty acids from the chloroplast, which stimulates lipid
biosynthesis in the Endoplasmic Reticulum, leads to a shift to phospholipid synthesis. In FAX1 Ox
lines, RBL11 is unable to decrease FAX1 protein abundance, and the disturbed glyco- to
phospholipid ratio impairs the efficient acclimation to cold temperatures, results in cold- and frost
sensitive mutants.
907

Material and Methods

Plant cultivation and growth conditions

Arabidopsis (*Arabidopsis thaliana*) ecotype Columbia (Col-*0*) and transgenic plants were sown on standard soil (type ED73, Einheitserde Patzer; Sinntal-Altengronau, Germany) with 10% (v/v) sand, stratified at 4°C for 48 h and then grown under short-day regime (10 h light/14 h dark) at 60% relative humidity and 120 µmol m^{-2} s^{-1} light intensity at 21°C, representing standard growth conditions. For cold treatment, plants were were grown for 21 days at 21°C first and subsequently transferred to a cultivation chamber (Fitotron SGR223, Weis-Gallenkamp Technik, Heidelberg, Germany) and incubated for several days at 4°C while all other parameters were kept constant.

We described FAX1 Ox#2 and FAX1 Ox#4 lines earlier (Li et al., 2015). *rbl10* mutants (Lavell et al., 2019) were provided by Dr. Christoph Benning (Michigan State University, Wisconsin, USA), *ftsh11* and *rbl11* mutants (Knopf et al., 2012; Adam et al., 2019) were provided by Dr. Zach Adam (Hebrew University, Jerusalem, Israel).

For RNA extraction, metabolite and anthocyanin analysis, Arabidopsis rosette leaves were collected five hours after onset of light, transferred immediately into liquid nitrogen and stored at -80°C until preparation. Leaf material for chloroplast envelope isolation was used directly after harvesting one hour before the start of illumination. For lipid isolation, rosette leaves were collected and directly transferred in a glass tube containing boiling water.

Generation of RBL11 overexpressor lines

The cloning steps to generate the dexamethasone-inducible RBL11 overexpression lines were performed using S7 Fusion Polymerase™ (MD-S7-100, Mobidiag, Espoo, Finland). The sequence of BirA-HA was first amplified from the Vector pcDNA3.1 MCS-BirA(R118G)-HA (Roux et al., 2012) using gene-specific primers and ligated C-terminally of *RBL11* in the pBSK vector (Short et al., 1988) by overlap extension PCR. The *RBL11*-BirA-HA construct was amplified using gene-specific primers with *attB*-sites attached and

inserted first into pDONR and then into the destination vector pTA7001-DEST (Aoyama
and Chua, 1997) via the Gateway Cloning system.

Wildtype Arabidopsis plants were transformed via floral dip using *Agrobacterium*
tumefaciens (GV3101). Positive transformed plants were selected by hygromycin
selection (Harrison et al., 2006) and confirmed by Western Blot analysis using an HA-
antibody after induction of leaves with 30µM dexamethasone. Sequences of gene-specific
primers, which were used for cloning, are provided in Supplemental File 2

Protein extraction from RBL11 overexpressor plants

Prior to protein extraction, approximately 50 leaves of Arabidopsis wildtype, or RBL11-HA
overexpressor plants were cut and placed in 30µM DEX or water, as control. Leaves were
incubated on a shaker for 48h in a plant chamber at either 21°C, or 4°C. Preparation of
soluble and insoluble (membrane) protein fraction and detection of RBL11-HA using HA
antibody via Western Blot was performed as described earlier (Khan et al., 2018).

Arabidopsis chloroplast envelope preparation

Arabidopsis chloroplast envelope membranes were isolated according to an established
protocol (Bouchnak et al., 2018) with few modifications. In a cold room (4°C), before onset
of light, 100 to 200 g of rosette leaves were harvested from six-weeks old plants and
ground in a Waring blender (three cycles, each of 2 seconds, average intensity) in the
presence of grinding medium (Tricine-KOH (20 mM, pH 8.4), sorbitol (0.4 M), EDTA (10
mM, pH 8), and $NaHCO_3$ (10 mM), BSA (0.1% (w/v)). The homogenate was filtered
through one layer of Miracloth and centrifuged for 2 min at 2,070x*g* at 4°C. The
supernatant was discarded, and the sediment was gently resuspended on ice with a soft
natural bristle paint brush in washing medium (1x) (Tricine-KOH (10 mM, pH 7.6), sorbitol
(0.4 M), $MgCl_2$ (2.5 mM), and EDTA (1.25 mM)) with a final volume of the combined
chloroplast solutions = 24 ml). 6 ml of the suspension was equally distributed and loaded
on top of four continuous Percoll (Sigma Aldrich, Heidelberg, Germany) gradients
(containing 50% Percoll / 0.4 M sorbitol, prepared by centrifugation at 38,700x*g*, for 55

min at 4 °C). Loaded gradients were centrifuged for 10 min at 13,300xg, 4°C using a

swinging-bucket rotor. The intact chloroplasts present in the lower phase were retrieved

with a 10 ml pipet. The intact chloroplast suspension was washed twice with 30 ml

washing buffer (1x) and centrifuged for 2 min at 2,070xg at 4°C. After washing, the purified

chloroplasts were lysed by resuspending the sediment in hypotonic medium (MOPS (10

mM, pH 7.8), $MgCl_2$ (4 mM), PMSF (1 mM, dissolved in isopropanol), benzamidine

hydrochloride hydrate (1 mM), and ε-amino caproic acid (0.5 mM). 3 ml of lysed

chloroplasts were loaded on top of two prepared sucrose gradients (4 ml of 0.93 M, 3 ml

of 0.6 M and 2.5 ml of 0.3 M sucrose). Gradients were ultracentrifuged for 1 h at 70,000xg,

4°C in a swinging-bucket rotor. The yellow band of both gradients (containing the

envelope fraction) was retrieved and pooled in one tube. The envelope suspension was

washed in 12 ml membrane washing buffer medium (MOPS (10 mM, pH 7.8), PMSF (1

mM), benzamidine hydrochloride hydrate (1 mM), ε-amino caproic acid (0.5 mM) and

ultracentrifuged again for 1 h at 110,000xg, 4 °C. Supernatants were aspirated by using

a water pump. Approximately 100 µl of membrane washing buffer was used to resuspend

the envelope sediment. Isolated envelopes were stored in liquid nitrogen until use.

982   Bimolecular Fluorescence Complementation for interaction studies

For the cloning of the BiFC constructs, the full-length sequences of *RBL11, FAX1* and

FTSH11 were used. The coding sequences were amplified by PCR using S7 Fusion

Polymerase (MD-S7-100, Mobidiag, Espoo, Finland) and inserted first into pDONR and

then into the pUBC-cYFP and the pUBC-nYFP vectors via the Gateway cloning system

(Grefen et al., 2010). Half of a yellow fluorescent protein (nYFP or cYFP) is thus fused to

the C- terminus of *RBL11*, *FAX1*, or *FTSH11*. The resulting constructs were then

transformed into *Agrobacterium tumefaciens* strain GV3101. Transient expression in

tobacco (*N. benthamiana*) leaves of RBL11, FAX1, and FTSH11, each fused to an nYFP

or cYFP, was performed as described (Walter et al., 2004). *Nicotiana benthamiana* leaves

were infiltrated through the lower epidermis. After 5 days, leaves were analyzed using a

Leica TCS SP5II fluorescence microscope (Leica Instruments, Wetzlar, Germany) (514

nm excitation and 525-582 nm detection of emission through an HCX PL APO 63 × 1.2 W

water immersion objective).

Sequences of gene-specific primers, which were used for cloning, are provided in
Supplemental File 2.

Frost recovery experiment

1000 For detection of the ability to recover from frost, a freezing tolerance test was performed
1001 according to an established approach (Trentmann et al., 2020; Cvetkovic et al., 2021).
1002 Survival rates and the numbers of wilted leaves were documented after 7 days of recovery
1003 under standard growth conditions.

1004

Pulse Amplitude Modulation (PAM) Fluorescence Measurements

1005

Photosystem II parameters at constant light intensities were monitored using an imaging
PAM M-Series IMAG-K7 and the ImagingWinGigE V2.56p (WALZ, Würzburg, Germany)
software. Induction curve settings were on default with 110 PAR as light intensity, 40 s
delay- and 20 s clock-time. Dark adaptation of plants lasted 10 minutes, followed by a 615
s long measurement monitoring PSII capacity (F_v/F_m), PSII effective photochemical
quantum yield (Y(II)), the quantum yield of light-induced non-photochemical fluorescence
quenching (Y(NPQ)) and quantum yield of nonregulated energy dissipation (Y(NO))
(Genty et al., 1989; Kramer et al., 2004). All plants analyzed (Col-*0*, FAX1 Ox#2 and FAX1
Ox#4) were grown for 28 days under short-day conditions (10/14 h; 110 µE) at RT before
being shifted for the duration of 6 weeks to 4°C (10/14 h, 110 µE). PAM measurements
were carried out on the day of the shift to 4°C, after one or six weeks in the cold.

1017

Determination of anthocyanin content

1018

For anthocyanin quantification 1 ml of extraction buffer composed of H_2O, propanol and
HCl (81:18:1) was added to 100 mg of fine grounded rosette plant material and incubated
for 3 min at 95°C, while shaking at 650 rpm and stored over night at RT in full darkness.
After centrifugation for 15 min at 12.500 rpm at RT, the supernatant was used for
photometric quantification at E1=535 nm and at E2=650 nm. The extinction was
determined and corrected ($E_{corr}=[E_{535}-(2.2*E_{650})]$ / mg FW).

1025

1026

Metabolomics

Metabolite profiling was performed according to established protocols (Roessner et al., 2001; Lisec et al., 2006; Erban et al., 2007). In brief, from four plants per genotype and growth condition, 50 mg fresh weight (Fw) of ground rosette material was mixed, in a 1.5 ml reaction tube, with 180 µl of cold (-20°C) methanol containing internal standards (10 µl ribitol, 0.2 mg ml^{-1} in water and 10 µl ^{13}C-sorbitol, 0.2 mg ml^{-1} in water). After 15 min of incubation at 70°C, the extract was cooled down to room temperature and carefully mixed with 100 µl of chloroform and 200 µl of water. To force phase separation, a 15 min centrifugation step at full speed was performed. Fifty µl of the upper (polar) phase was dried *in vacuo* and stored at -80°C. For derivatization, the pellet was resuspended in 10 µl of methoxyamin-hydrochloride (20 mg ml^{-1} in pyridine) and incubated for 90 min at 40°C. After addition of 20 µl of BSTFA (*N,O*-bis[trimethylsilyl]trifluoroacetamide) containing 2.5 µl retention time standard mixture of linear alkanes (n-decane, n-dodecane, n-pentadecane, n-nonadecane, n-docosane, n-octacosane, n-dotriacontane), the preparation was incubated at 40°C for further 45 min.

One µl of each sample was injected into a GC–TOF–MS system (Pegasus HT, Leco, St Joseph, USA). Samples were automatically processed by an autosampler system (Combi PAL, CTC Analytics AG, Zwingen, Switzerland). Helium acted as carrier gas at a constant flow rate of 1 ml min^{-1}. Gas chromatography was performed on an Agilent GC (7890A, Agilent, Santa Clara, CA, USA) using a 30 m VF-5ms column with 10 m EZ-Guard column. The temperature of the split/splitless injector was set to 250°C, as well as the transfer line and the ion source. The initial oven temperature (70°C) was linearly increased to a final temperature of 350 °C by a rate of 9°C/min. Metabolites were ionized and fractionated by an ion pulse of 70 eV. Mass spectra were recorded at 20 scans s^{-1} with an *m/z* 50– 600 scanning range. Chromatograms and mass spectra were evaluated using ChromaTOF 4.72 and TagFinder 4.1 software (Luedemann et al., 2008).

RNA Extraction, cDNA synthesis and qRT-PCR

RNA was extracted from 50 mg of frozen and fine ground rosette leaf material from four biological replicates per genotype and growth condition using the NucleoSpin RNA Plant Kit (Macherey-Nagel, Düren, Germany), according to the manufacturer's protocol. The

1058 synthesis of cDNA from RNA was performed with the qScript cDNA Synthesis Kit

1059 (Quantabio, Beverly, MA, USA). Primers used for gene expression analysis via qRT-PCR

1060 are listed in Supplemental File 2. *AtUBQ* was used as reference gene for normalization.

1061

1062 **Nitroblue tetrazolium (NBT) staining**

1063 For ROS staining, O_2^- was detected by nitroblue tetrazolium staining (Fryer et al., 2002)

1064 in whole rosettes of two weeks old Arabidopsis plants cultivated as described above. For

1065 cold treatments, plants were transferred to 4°C for four days, while control plants were

1066 kept under standard conditions.

1067

Measurement of galacto-, phospho- and sulfolipids

For the analysis of lipids, plants were cultivated as described above and transferred from

standard conditions to 4°C for 2 weeks, 10 weeks, or used directly. From 5 plants per

genotype 100 mg fresh weight of rosette leaf material was harvested and immediately

placed into a glass tube containing boiling water to prevent degradation of phospholipids

through phospholipase D activity. The lipid extraction was performed with

chloroform/methanol after deactivation of lipase activities by boiling the tissue in water as

described earlier (Gasulla et al., 2013). Lipids were measured by tandem mass

spectrometry (Q-TOF 6530 Agilent Technologies) and quantified by MS/MS experiments

with internal standards following the strategy developed earlier (Gasulla et al., 2013; Welti

et al., 2002).

1079

Immunoblotting

Per lane, 8 µg of isolated chloroplast envelope protein or 30 µg of freshly prepared protein

extract from Arabidopsis leaf material were separated via SDS-PAGE (12%). The proteins

in the gel were transferred onto a nitrocellulose membrane by a semi-dry blotting system

(TransBlot® Turbo™ Transfer System, BIO RAD, Göttingen, Germany). The membrane

was blocked in phosphate-buffered saline plus 0.1% (v/v) Tween 20 (PBS-T) with 3% milk

powder for 1 h at room temperature, and then washed three times in PBS-T for 10 min.

The membrane was incubated with a polyclonal rabbit antibody raised against FAX1 (Li

et al., 2015) over night at 4°C at 1:1000 dilution. After three times of washing with PBS-T

for 10 min, the membrane was incubated with a horseradish peroxidase (HPR) conjugated

anti-rabbit antibody (Promega, Walldorf, Germany) diluted 1:10.000 in PBS-T with 3% milk

powder for 1 h. The immunoreaction was visualized by chemiluminescence using ECL

Prime Western blotting reagent (GE Healthcare, Karlsruhe Germany) and a Fusion Solo

S6 (Vilber-Lourmat, Eberhardzell, Germany).

1094

Peptide Mass spectrometry

Peptide Mass spectrometry

Enriched envelope fractions were precipitated in 80% acetone, digested in solution using

Lys-C and trypsin, and resulting peptides were desalted as previously described (Hammel

et al., 2018). Peptide mass spectrometry was performed using a nanoUHPLC-IM-MS

system (nanoElute coupled to timsTOF Pro2, Bruker Daltonics, Bremen, Germany).

Samples were directly loaded on a 25 cm, 75 µm ID, 1.6 µm particle size, C18 column

with integrated emitter (Odyssey/Aurora ionopticks, Melbourne, Australia) set to 50 °C,

and peptides were separated under a flow rate of 0.3 µl/min using buffers A (water, 0.1%

formic acid) and B (acetonitrile, 0.1% formic acid). The gradient employed ramped from

2% B to 25% B within 67 min, then to 37% B within 10 min, followed by washing and

equilibration steps. The MS was operated in positive mode, electrospray voltage was set

to 1.4 kV and spectra were recorded from 100 – 1700 m/z. A total of 10 MS/MS PASEF

ramps ($1/K_0$ 0.6 – 1.43 V*s/cm^2) with 100 ms duration were acquired per cycle, and target

intensity for MS/MS was set to 14500 whereafter the precursors were excluded from

fragmentation for 0.4 min.

1110

Protein Identification and Quantification

Protein Identification and Quantification

Acquired data were searched against the Uniprot protein sequences for *Arabidopsis*

thaliana (UP000006548) using the FragPipe v19.1 processing pipeline choosing the

default LFQ-MBR workflow with minor modifications: peptide length was set to a minimum

of 6 amino acids, missed cleavages to 3, normalization of intensities across runs was

omitted. Mass spectrometry raw and processed data have been deposited at the

ProteomeXChange Consoritum via the PRIDE partner repository (Perez-Riverol et al.,

2021) with the dataset identifier PXD041219.

1119

1120

Data normalization and missing value imputation

Prior to computation of protein-level statistics, replicate groups were normalized using the median-of-ratios method (Anders and Huber, 2010). Subsequently, we computed global variance estimates and local gene-wise mean estimates to impute missing data points as independent draws from normal distributions. When a replicate group did not contain any measurement, the normal distribution was centered at an intensity corresponding to the 5% quantile of all intensities. Proteins were excluded from statistical analysis according to two filter criteria. First, proteins were considered ineligible for downstream analysis if there was no biological replicate group with at least one reading. Second, only proteins reported to be localized to the envelope membrane were considered, either by association with matching Gene Ontology terms retrieved from UniProt or by entry in the manually curated AT_CHLORO database (Ashburner et al., 2000; Bruley et al., 2012; Consortium, 2021).

Statistical Analyses

Statistical analyses regarding the proteomic data were based on log2 transformed imputed values. According to the experimental design which consisted of two factors (genotype and treatment) at each of two levels (wild type/mutant and normal/cold), changes in protein abundance were evaluated using two-way analysis of variance, ANOVA (Fisher, 1925). To account for multiple hypothesis testing,we controlled the False Discovery Rate (FDR) by computing q-values based on the ANOVA p-values as previously described by Storey (Storey and Tibshirani, 2003; Storey et al., 2004). All calculations were carried out using the FSharp.Stats library for statistical computing (Venn et al., 2023). Changes were considered to be significant if a q-value of 0.05 was not exceeded.

For statistical analysis of the numerical data GraphPad Prism 9 and Microsoft Office Excel were used. Significant differences between two groups were analyzed by two-tailed Student's *t*-test. The software Shiny application (https://houssein-assaad.shinyapps.io/TwoWayANOVA/) was used for letter-based representation of all pairwise comparisons using popular statistical tests in two-way ANOVA. Statistical data is provided in Supplemental File 1.

1151

Accession numbers

Sequence data from this article can be found in the ARAMEMNON GenBank data library (http://aramemnon.uni-koeln.de/). *FAX1* (*At3g57280*), *FTSH11* (*At5g53170*), *RBL10* (*At1g25290*), *RBL11* (*At5g25752*), *BAP1* (*At3g61190*).

Acknowledgments

Work in the labs from ML, H-HK, AF MS, TM and HEN was supported by the Deutsche Forschungsgemeinschaft (DFG) within the SFB/Transregio (TRR) 175, The Green Hub.

References

Abdallah F, Salamini F, Leister D (2000) A prediction of the size and evolutionary origin of the proteome of chloroplasts of Arabidopsis. Trends Plant Sci. **5:** 141-142

Adam Z, Aviv-Sharon E, Keren-Paz A, Naveh L, Rozenberg M, Savidor A, Chen J (2019) The chloroplast envelope protease FTSH11 – interaction with CPN60 and identification of potential substrates. Front. Plant Sci. **10**

Alberdi M, Corcuera LJ (1991) Cold-acclimation in plants. Phytochem. **30:** 3177-3184

Alet AI, Sanchez DH, Cuevas JC, Del Valle S, Altabella T, Tiburcio AF, Marco F, Ferrando A, Espasandín FD, González ME, Ruiz OA, Carrasco P (2011) Putrescine accumulation in *Arabidopsis thaliana* transgenic lines enhances tolerance to dehydration and freezing stress. Plant Sig. Behav. **6:** 278-286

Alia PS, Mohanty P (1997) Involvement of proline in protecting thylakoid membranes against free radical-induced photodamage. J. Photochem. Photobiol. **38:** 253-257

Anders S, Huber W (2010) Differential expression analysis for sequence count data. Genome Biol. **11:** R106

Andersson MX, Dörmann P (2009) Chloroplast membrane lipid biosynthesis and transport. *In* AS Sandelius, H Aronsson, eds, The Chloroplast: Interactions with the Environment. Springer, Berlin, Heidelberg, pp 125-158

Aoyama T, Chua NH (1997) A glucocorticoid-mediated transcriptional induction system in transgenic plants. Plant J. **11:** 605-612

Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, Harris MA, Hill DP, Issel-Tarver L, Kasarskis A, Lewis S, Matese JC, Richardson JE, Ringwald M, Rubin GM, Sherlock G (2000) Gene Ontology: tool for the unification of biology. Nature Genetics **25:** 25-29

Awai K, Xu C, Tamot B, Benning C (2006) A phosphatidic acid-binding protein of the chloroplast inner envelope membrane involved in lipid trafficking. Proc. Natl. Acad. Sci. **103:** 10817-10822

Balsera M, Goetze TA, Kovács-Bogdán E, Schürmann P, Wagner R, Buchanan BB, Soll J, Bölter B (2009) Characterization of Tic110, a channel-forming protein at the inner envelope membrane of chloroplasts, unveils a response to Ca^{2+} and a stromal regulatory disulfide bridge. J. Biol. Chem. **284:** 2603-2616

Barnes AC, Benning C, Roston RL (2016) Chloroplast membrane remodeling during freezing stress is accompanied by cytoplasmic acidification activating SENSITIVE TO FREEZING2 Plant Physiol. **171:** 2140-2149

Barrero-Sicilia C, Silvestre S, Haslam RP, Michaelson LV (2017) Lipid remodelling: unravelling the response to cold stress in Arabidopsis and its extremophile relative *Eutrema salsugineum*. Plant Sci. **263:** 194-200

Barthélemy X, Bouvier G, Radunz A, Docquier S, Schmid GH, Franck F (2000) Localization of NADPH-protochlorophyllide reductase in plastids of barley at different greening stages. Photosynth. Res. **64:** 63-76

Baxter A, Mittler R, Suzuki N (2014) ROS as key players in plant stress signalling

3. J Exp Bot **65:** 1229-1240

Botté CY, Deligny M, Roccia A, Bonneau A-L, Saïdani N, Hardré H, Aci S, Yamaryo-Botté Y, Jouhet J, Dubots E, Loizeau K, Bastien O, Bréhélin L, Joyard J, Cintrat J-C, Falconet D, Block MA, Rousseau B, Lopez R, Maréchal E (2011) Chemical inhibitors of monogalactosyldiacylglycerol synthases in *Arabidopsis thaliana*. Nat. Chem. Biol. **7:** 834-842

Bouchnak I, Moyet L, Salvi D, Kuntz M, Rolland N (2018) Preparation of chloroplast sub-compartments from Arabidopsis for the analysis of protein localization by immunoblotting or proteomics. J. Vis. Exp.

Boudière L, Michaud M, Petroutsos D, Rébeillé F, Falconet D, Bastien O, Roy S, Finazzi G, Rolland N, Jouhet J, Block MA, Maréchal E (2014) Glycerolipids in photosynthesis: Composition, synthesis and trafficking. Biochim. Biophys. Acta **1837:** 470-480

Bruley C, Dupierris V, Salvi D, Rolland N, Ferro M (2012) AT_CHLORO: a chloroplast protein database dedicated to sub-plastidial localization. Front. Plant Sci. **3:** 205-205

Chalker-Scott L (1999) Environmental significance of anthocyanins in plant stress responses. Photochemistry and Photobiology **70:** 1-9

Chen J, Burke JJ, Velten J, Xin Z (2006) FtsH11 protease plays a critical role in Arabidopsis thermotolerance. Plant J. **48:** 73-84

Choudhury FK, Rivero RM, Blumwald E, Mittler R (2017) Reactive oxygen species, abiotic stress and stress combination. Plant J. **90:** 856-867

Consortium GO (2021) The Gene Ontology resource: enriching a GOld mine. Nucleic Acids Res **49:** D325-d334

Curtis TY, Bo V, Tucker A, Halford NG (2018) Construction of a network describing asparagine metabolism in plants and its application to the identification of genes affecting asparagine metabolism in wheat under drought and nutritional stress. Food, Energy Sec. **7:** e00126-e00126

Cvetkovic J, Haferkamp I, Rode R, Keller I, Pommerrenig B, Trentmann O, Altensell J, Fischer-Stettler M, Eicke S, Zeeman SC, Neuhaus HE (2021) Ectopic maltase alleviates dwarf phenotype and improves plant frost tolerance of maltose transporter mutants. Plant Physiol. **186:** 315-329

Doke N (1983) Generation of superoxide anion by potato tuber protoplasts during the hypersensitive response to hyphal wall components of *Phytophthora infestans* and specific inhibition of the reaction by suppressors of hypersensitivity. Physiol. Plant Pathol. **23:** 359-367

Dorne AJ, Joyard J, Douce R (1990) Do thylakoids really contain phosphatidylcholine? Proc. Natl. Acad. Sci. USA **87:** 71-74

Erban A, Schauer N, Fernie AR, Kopka J (2007) Nonsupervised construction and application of mass spectral and retention time index libraries from time-of-flight gas chromatography-mass spectrometry metabolite profiles. Methods Mol. Biol. **358:** 19-38

Fan J, Zhai Z, Yan C, Xu C (2015) Arabidopsis TRIGALACTOSYLDIACYLGLYCEROL5 interacts with TGD1, TGD2, and TGD4 to facilitate lipid transfer from the Endoplasmic Reticulum to plastids. Plant Cell **27:** 2941-2955

Fisher RA (1925) Statistical Methods for Research Workers. Oliver and Boyd, Edinburgh, Scotland

Fryer MJ, Oxborough K, Mullineaux PM, Baker NR (2002) Imaging of photo-oxidative stress responses in leaves. J. Exp. Bot. **53:** 1249-1254

Furumoto T, Yamaguchi T, Ohshima-Ichie Y, Nakamura M, Tsuchida-Iwata Y, Shimamura M, Ohnishi J, Hata S, Gowik U, Westhoff P, Brautigam A, Weber AP, Izui K (2011) A plastidial sodium-dependent pyruvate transporter. Nature **476:** 472-475

Gao J, Lunn D, Wallis JG, Browse J (2020) Phosphatidylglycerol composition is central to chilling damage in the Arabidopsis *fab1* mutant. Plant Physiol.**:** pp.01219.02020

Garcia-Molina A, Kleine T, Schneider K, Mühlhaus T, Lehmann M, Leister D (2020) Translational components contribute to acclimation responses to high light, heat and cold in Arabidopsis. iScience **23:** 101331

Gasulla F, vom Dorp K, Dombrink I, Zähringer U, Gisch N, Dörmann P, Bartels D (2013) The role of lipid metabolism in the acquisition of desiccation tolerance in *Craterostigma plantagineum*: a comparative approach. Plant J. **75:** 726-741

Genty B, Briantais J-M, Baker NR (1989) The relationship between the quantum yield of photosynthetic electron transport and quenching of chlorophyll fluorescence. Biochim. Biophys. Acta **990:** 87-92

Ghosh UK, Islam MN, Siddiqui MN, Cao X, Khan MAR (2022) Proline, a multifaceted signalling molecule in plant responses to abiotic stress: understanding the physiological mechanisms. Plant Biol. **24:** 227-239

Greenbaum D, Colangelo C, Williams K, Gerstein M (2003) Comparing protein abundance and mRNA expression levels on a genomic scale. Genome Biol. **4:** 117

Grefen C, Donald N, Hashimoto K, Kudla J, Schumacher K, Blatt MR (2010) A ubiquitin-10 promoter-based vector set for fluorescent protein tagging facilitates temporal stability and native protein distribution in transient and stable expression studies. Plant J. **64:** 355-365

Guan L, Denkert N, Eisa A, Lehmann M, Sjuts I, Weiberg A, Soll J, Meinecke M, Schwenkert S (2019) JASSY, a chloroplast outer membrane protein required for jasmonate biosynthesis. Proc. Natl. Acad. Sci. **116:** 10568-10575

Guo L, Yang H, Zhang X, Yang S (2013) Lipid transfer protein 3 as a target of MYB96 mediates freezing and drought stress in Arabidopsis. J. Exp. Bot. **64:** 1755-1767

Guo X, Liu D, Chong K (2018) Cold signaling in plants: Insights into mechanisms and regulation. J. Integr. Plant Biol. **60:** 745-756

Gygi SP, Rist B, Gerber SA, Turecek F, Gelb MH, Aebersold R (1999) Quantitative analysis of complex protein mixtures using isotope-coded affinity tags. Nat. Biotechnol. **17:** 994-999

Hagio M, Sakurai I, Sato S, Kato T, Tabata S, Wada H (2002) Phosphatidylglycerol is essential for the development of thylakoid membranes in *Arabidopsis thaliana*. Plant, Cell Physiol. **43:** 1456-1464

Hammel A, Zimmer D, Sommer F, Mühlhaus T, Schroda M (2018) Absolute quantification of major photosynthetic protein complexes in *Chlamydomonas reinhardtii* using quantification concatamers (QconCATs). Front. Plant Sci. **9**

Harrison SJ, Mott EK, Parsley K, Aspinall S, Gray JC, Cottage A (2006) A rapid and robust method of identifying transformed Arabidopsis thaliana seedlings following floral dip transformation. Plant Methods **2:** 19

Hoffmann C, Plocharski B, Haferkamp I, Leroch M, Ewald R, Bauwe H, Riemer J, Herrmann JM, Neuhaus HE (2013) From endoplasmic reticulum to mitochondria: absence of the Arabidopsis ATP antiporter Endoplasmic Reticulum Adenylate Transporter1 perturbs photorespiration. Plant Cell **25:** 2647-2660

Hölzl G, Dörmann P (2019) Chloroplast lipids and their biosynthesis. Ann. Rev. Plant Biol. **70:** 51-81

Kerppola TK (2008) Bimolecular fluorescence complementation (BiFC) analysis as a probe of protein interactions in living cells. Ann. Rev. Biophys. **37:** 465-487

Khan M, Youn JY, Gingras AC, Subramaniam R, Desveaux D (2018) In planta proximity dependent biotin
identification (BioID). Sci. Rep. **8:** 9212
Khodakovskaya M, McAvoy R, Peters J, Wu H, Li Y (2006) Enhanced cold tolerance in transgenic tobacco
expressing a chloroplast ω-3 fatty acid desaturase gene under the control of a cold-inducible
promoter. Planta **223:** 1090-1100
Knopf RR, Feder A, Mayer K, Lin A, Rozenberg M, Schaller A, Adam Z (2012) Rhomboid proteins in the
chloroplast envelope affect the level of allene oxide synthase in *Arabidopsis thaliana*. Plant J. **72:**
559-571
Kobayashi K (2016) Role of membrane glycerolipids in photosynthesis, thylakoid biogenesis and
chloroplast development. J. Plant Res. **129:** 565-580
Koevoets IT, Venema JH, Elzenga JTM, Testerink C (2016) Roots withstanding their environment:
exploiting root system architecture responses to abiotic stress to improve crop tolerance. Front.
Plant Sci. **7**
Koussounadis A, Langdon SP, Um IH, Harrison DJ, Smith VA (2015) Relationship between differentially
expressed mRNA and mRNA-protein correlations in a xenograft model system. Scientific Rep. **5:**
10775
Kramer DM, Johnson G, Kiirats O, Edwards GE (2004) New fluorescence parameters for the determination
of QA redox state and excitation energy fluxes. Photosynth. Res. **79:** 209
Kunst L, Browse J, Somerville C (1988) Altered regulation of lipid biosynthesis in a mutant of Arabidopsis
deficient in chloroplast glycerol-3-phosphate acyltransferase activity. Proc. Natl. Acad. Sci. USA **85:**
4143-4147
Lavell A, Froehlich JE, Baylis O, Rotondo AD, Benning C (2019) A predicted plastid rhomboid protease
affects phosphatidic acid metabolism in *Arabidopsis thaliana*. Plant J. **99:** 978-987
Lavell AA, Benning C (2019) Cellular organization and regulation of plant glycerolipid metabolism. Plant
Cell Physiol. **60:** 1176-1183
Li-Beisson Y, Neunzig J, Lee Y, Philippar K (2017) Plant membrane-protein mediated intracellular traffic of
fatty acids and acyl lipids. Curr. Opin. Plant Biol. **40:** 138-146
Li-Beisson Y, Shorrosh B, Beisson F, Andersson MX, Arondel V, Bates PD, Baud S, Bird D, Debono A,
Durrett TP, Franke RB, Graham IA, Katayama K, Kelly AA, Larson T, Markham JE, Miquel M,
Molina I, Nishida I, Rowland O, Samuels L, Schmid KM, Wada H, Welti R, Xu C, Zallot R, Ohlrogge
J (2010) Acyl-lipid metabolism. The Arabidopsis Book **8:** e0133-e0133
Li N, Gügel IL, Giavalisco P, Zeisler V, Schreiber L, Soll J, Philippar K (2015) FAX1, a novel membrane potein
mediating plastid fatty acid export. PLoS Biol. **13:** e1002053
Li N, Xu C, Li-Beisson Y, Philippar K (2016) Fatty acid and lipid transport in plant cells. Trends Plant Sci. **21:**
145-158
Li Q, Zheng Q, Shen W, Cram D, Fowler DB, Wei Y, Zou J (2015) Understanding the biochemical basis of
temperature-induced lipid pathway adjustments in plants. Plant Cell **27:** 86-103
Liang X, Zhang L, Natarajan SK, Becker DF (2013) Proline mechanisms of stress survival. Antioxidants &
Redox Signaling **19:** 998-1011
Lin Y-S, Medlyn BE, Ellsworth DS (2012) Temperature responses of leaf net photosynthesis: the role of
component processes. Tree Physiology **32:** 219-231
Lisec J, Schauer N, Kopka J, Willmitzer L, Fernie AR (2006) Gas chromatography mass spectrometry–based
metabolite profiling in plants. Nature Prot. **1:** 387-396
Lu B, Xu C, Awai K, Jones AD, Benning C (2007) A small ATPase protein of Arabidopsis, TGD3, involved in
chloroplast lipid import. J. Biol. Chem. **282:** 35945-35953
Luedemann A, Strassburg K, Erban A, Kopka J (2008) TagFinder for the quantitative analysis of gas
chromatography-mass spectrometry (GC-MS)-based metabolite profiling experiments.
Bioinformatics **24:** 732-737

Mackey KR, Paytan A, Caldeira K, Grossman AR, Moran D, McIlvin M, Saito MA (2013) Effect of temperature on photosynthesis and growth in marine *Synechococcus spp*. Plant Physiol. **163:** 815-829

Marco F, Busó E, Lafuente T, Carrasco P (2019) Spermine confers stress resilience by modulating abscisic acid biosynthesis and stress responses in Arabidopsis plants. Front. Plant Sci. **10**

Mielke K, Wagner R, Mishra LS, Demir F, Perrar A, Huesgen PF, Funk C (2020) FtsH12 abundance modulates chloroplast development in *Arabidopsis thaliana*. J. Exp. Bot.

Miquel M, James D, Jr., Dooner H, Browse J (1993) Arabidopsis requires polyunsaturated lipids for low-temperature survival. Proc. Natl. Acad. Sci. USA **90:** 6208-6212

Moellering ER, Muthan B, Benning C (2010) Freezing tolerance in plants requires lipid remodeling at the outer chloroplast membrane. Science **330:** 226-228

Murchie EH, Lawson T (2013) Chlorophyll fluorescence analysis: a guide to good practice and understanding some new applications. J. Exp. Bot. **64:** 3983-3998

Nakamura Y (2017) Plant phospholipid diversity: emerging functions in metabolism and protein-lipid interactions. Trends Plant Sci. **22:** 1027-1040

Niittylä T, Messerli G, Trevisan M, Chen J, Smith AM, Zeeman SC (2004) A previously unknown maltose transporter essential for starch degradation in leaves. Science **303:** 87-89

Nishimura K, Kato Y, Sakamoto W (2016) Chloroplast proteases: updates on poteolysis within and across suborganellar compartments. Plant Physiol. **171:** 2280-2293

Obata T, Fernie AR (2012) The use of metabolomics to dissect plant responses to abiotic stresses. Cell. Mol. Life Sci. **69:** 3225-3243

Parthasarathy A, Savka MA, Hudson AO (2019) The synthesis and role of β-alanine in plants. Front. Plant Sci. **10:** 921

Pattanayak GK, Tripathy BC (2011) Overexpression of protochlorophyllide oidoreductase C regulates oxidative stress in Arabidopsis. PLoS ONE **6:** e26532

Patzke K, Prananingrum P, Klemens PAW, Trentmann O, Martins Rodrigues C, Keller I, Fernie AR, Geigenberger P, Bölter B, Lehmann M, Schmitz-Esser S, Pommerrenig B, Haferkamp I, Neuhaus HE (2019) The plastidic sugar transporter pSuT influences flowering and affects cold responses. Plant Physiol. **179:** 569-587

Perez-Riverol Y, Bai J, Bandla C, García-Seisdedos D, Hewapathirana S, Kamatchinathan S, Kundu Deepti J, Prakash A, Frericks-Zipper A, Eisenacher M, Walzer M, Wang S, Brazma A, Vizcaíno Juan A (2021) The PRIDE database resources in 2022: a hub for mass spectrometry-based proteomics evidences. Nucleic Acids Res. **50:** D543-D552

Perrett M, Gothard M, Ludwig A, Rouhier KA (2017) Identifying a source of β-alanine and its broader implications in *Arabidopsis thaliana* by GC/MS. FASEB J. **31:** 626.621-626.621

Pommerrenig B, Ludewig F, Cvetkovic J, Trentmann O, Klemens PAW, Neuhaus HE (2018) In concert: orchestrated changes in carbohydrate homeostasis are critical for plant abiotic stress tolerance. Plant, Cell Physiol. **59:** 1290-1299

Pospíšil P (2016) Production of reactive oxygen species by photosystem II as a response to light and temperature stress. Front. Plant Sci. **7:** 1950

Pottosin I, Shabala S (2016) Transport across chloroplast membranes: optimizing photosynthesis for adverse environmental conditions. Mol. Plant **9:** 356-370

Rawsthorne S (2002) Carbon flux and fatty acid synthesis in plants. Prog. Lipid Res. **41:** 182-196

Roessner U, Lüdemann A, Brust D, Fiehn O, Linke T, Willmitzer L, Fernie A (2001) Metabolic profiling allows comprehensive phenotyping of genetically or environmentally modified plant systems. Plant Cell **13:** 11-29

Roughan PG, Slack CR (1982) Cellular organization of glycerolipid metabolism. Ann. Rev. Plant Physiol. **33:** 97-132

Roux KJ, Kim DI, Raida M, Burke B (2012) A promiscuous biotin ligase fusion protein identifies proximal and interacting proteins in mammalian cells. J.Cell Biol. **196:** 801-810

Schnurr JA, Shockey JM, de Boer GJ, Browse JA (2002) Fatty acid export from the chloroplast. Molecular characterization of a major plastidial acyl-coenzyme A synthetase from Arabidopsis. Plant Physiol. **129:** 1700-1709

Schreiber U (2004) Pulse-Amplitude-Modulation (PAM) fluorometry and saturation pulse method: an overview. *In* GC Papageorgiou, Govindjee, eds, Chlorophyll a Fluorescence: A Signature of Photosynthesis. Springer Netherlands, Dordrecht, pp 279-319

Schwenkert S, Fernie AR, Geigenberger P, Leister D, Möhlmann T, Naranjo B, Neuhaus HE (2022) Chloroplasts are key players to cope with light and temperature stress. Trends Plant Sci. **in press**

Schwenkert S, Lo WT, Szulc B, Yip CK, Pratt AI, Cusack SA, Brandt B, Leister D, Kunz HH (2023) Probing the physiological role of the plastid outer-envelope membrane using the oemiR plasmid collection. G3 (Bethesda)

Short JM, Fernandez JM, Sorge JA, Huse WD (1988) Lambda ZAP: a bacteriophage lambda expression vector with in vivo excision properties. Nucleic Acids Res. **16:** 7583-7600

Smallwood M, Bowles DJ (2002) Plants in a cold climate. Philos. Trans. R Soc. Lond. B Biol. Sci. **357:** 831-847

Song Y, Chen Q, Ci D, Shao X, Zhang D (2014) Effects of high temperature on photosynthesis and related gene expression in poplar. BMC Plant Biol. **14:** 111

Steponkus PL, Garber MP, Myers SP, Lineberger RD (1977) Effects of cold acclimation and freezing on structure and function of chloroplast thylakoids. Cryobiology **14:** 303-321

Storey JD, Taylor JE, Siegmund D (2004) Strong control, conservative point estimation and simultaneous conservative consistency of false discovery rates: A unified approach. J. Royal Stat. Soc. **66:** 187-205

Storey JD, Tibshirani R (2003) Statistical significance for genomewide studies. Proc. Nat. Acad. Sci. **100:** 9440-9445

Takemura T, Imamura S, Tanaka K (2019) Identification of a chloroplast fatty acid exporter protein, *Cm*FAX1, and triacylglycerol accumulation by its overexpression in the unicellular red alga *Cyanidioschyzon merolae*. Algal Res. **38:** 101396

Taylor NL, Tan Y-F, Jacoby RP, Millar AH (2009) Abiotic environmental stress induced changes in the Arabidopsis thaliana chloroplast, mitochondria and peroxisome proteomes. J. Proteom. **72:** 367-378

Tian Y, Lv X, Xie G, Wang L, Dai T, Qin X, Chen F, Xu Y (2019) FAX2 mediates fatty acid export from plastids in developing Arabidopsis seeds. Plant Cell Physiol. **60:** 2231-2242

Tjaden J, Schwöppe C, Möhlmann T, Neuhaus HE (1998) Expression of the plastidic ATP/ADP transporter gene in *Escherichia coli* leads to a functional adenine nucleotide transport system in the bacterial cytoplasmic membrane. J.Biol.Chem. **273:** 9630-9636

Trentmann O, Mühlhaus T, Zimmer D, Sommer F, Schroda M, Haferkamp I, Keller I, Pommerrenig B, Neuhaus HE (2020) Identification of chloroplast envelope proteins with critical importance for cold acclimation. Plant Physiol. **182:** 1239-1255

van Meer G, Voelker DR, Feigenson GW (2008) Membrane lipids: where they are and how they behave. Nat. Rev. Mol. Cell Biol. **9:** 112-124

Venn B, Mühlhaus T, Schneider K, Weil L, Zimmer D (2023) FSharp.Stats: Release 0.4.11. . Zenodo

Wagner R, Aigner H, Pružinská A, Jänkänpää HJ, Jansson S, Funk C (2011) Fitness analyses of *Arabidopsis thaliana* mutants depleted of FtsH metalloproteases and characterization of three FtsH6 deletion mutants exposed to high light stress, senescence and chilling. New Phytol. **191:** 449-458

Wagner R, von Sydow L, Aigner H, Netotea S, Brugière S, Sjögren L, Ferro M, Clarke A, Funk C (2016) Deletion of FtsH11 protease has impact on chloroplast structure and function in *Arabidopsis thaliana* when grown under continuous light. Plant, Cell Environ. **39:** 2530-2544

Walker B, Ariza LS, Kaines S, Badger MR, Cousins AB (2013) Temperature response of in vivo Rubisco kinetics and mesophyll conductance in *Arabidopsis thaliana*: comparisons to *Nicotiana tabacum*. Plant Cell Environ. **36:** 2108-2119

Walter M, Chaban C, Schütze K, Batistic O, Weckermann K, Näke C, Blazevic D, Grefen C, Schumacher K, Oecking C, Harter K, Kudla J (2004) Visualization of protein interactions in living plant cells using bimolecular fluorescence complementation. Plant J. **40:** 428-438

Wang P, Hsu C-C, Du Y, Zhu P, Zhao C, Fu X, Zhang C, Paez JS, Macho AP, Tao WA, Zhu J-K (2020) Mapping proteome-wide targets of protein kinases in plant stress responses. Proc. Natl. Acad. Sci. **117:** 3270–3280

Wang Z, Benning C (2012) Chloroplast lipid synthesis and lipid trafficking through ER–plastid membrane contact sites. Biochem. Soc. Trans. **40:** 457-463

Welti R, Li W, Li M, Sang Y, Biesiada H, Zhou HE, Rajashekar CB, Williams TD, Wang X (2002) Profiling membrane lipids in plant stress responses. Role of phospholipase D alpha in freezing-induced lipid changes in Arabidopsis. J. Biol. Chem. **277:** 31994-32002

Witz S, Panwar P, Schober M, Deppe J, Pasha FA, Lemieux MJ, Möhlmann T (2014) Structure-function relationship of a plant NCS1 member--homology modeling and mutagenesis identified residues critical for substrate specificity of PLUTO, a nucleobase transporter from Arabidopsis. PLoS ONE **9:** e91343

Xiao Z, Tang F, Zhang L, Li S, Wang S, Huo Q, Yang B, Zhang C, Wang D, Li Q, Wei L, Guo T, Qu C, Lu K, Zhang Y, Guo L, Li J, Li N (2021) The *Brassica napus* fatty acid exporter FAX1-1 contributes to biological yield, seed oil content, and oil quality. Biotechnol. Biofuels **14:** 190

Xu C, Moellering ER, Muthan B, Fan J, Benning C (2010) Lipid transport mediated by Arabidopsis TGD proteins is unidirectional from the endoplasmic reticulum to the plastid. Plant Cell Physiol. **51:** 1019-1028

Xu Y, Kambhampati S, Morley SA, Cook R, Froehlich J, Allen DK, Benning C (2023) Arabidopsis ACYL CARRIER PROTEIN4 and RHOMBOID LIKE10 act independently in chloroplast phosphatidate synthesis. Plant Physiol.

Yang H, Yang S, Li Y, Hua J (2007) The Arabidopsis BAP1 and BAP2 genes are general inhibitors of programmed cell death. Plant Physiol. **145:** 135-146

Yu L, Shen W, Fan J, Sah SK, Mavraganis I, Wang L, Gao P, Gao J, Zheng Q, Meesapyodsuk D, Yang H, Li Q, Zou J, Xu C (2023) A chloroplast diacylglycerol lipase modulates glycerolipid pathway balance in Arabidopsis. Plant J. **in press**

Zhang J, Yang D, Li M, Shi L (2016) Metabolic profiles reveal changes in wild and cultivated soybean seedling leaves under salt stress. PloS One **11:** e0159622-e0159622

Zhang Y, Zhu H, Zhang Q, Li M, Yan M, Wang R, Wang L, Welti R, Zhang W, Wang X (2009) Phospholipase Dα1 and phosphatidic acid regulate NADPH oxidase activity and production of reactive oxygen species in ABA-mediated stomatal closure in Arabidopsis. Plant Cell **21:** 2357-2377

Zheng G, Tian B, Zhang F, Tao F, Li W (2011) Plant adaptation to frequent alterations between high and low temperatures: remodelling of membrane lipids and maintenance of unsaturation levels. Plant, Cell Environ. **34:** 1431-1442

Zhu L, He S, Liu Y, Shi J, Xu J (2020) Arabidopsis FAX1 mediated fatty acid export is required for the transcriptional regulation of anther development and pollen wall formation. Plant Mol. Biol. **104:** 187-201

Zhu Y, Yang H, Mang H-G, Hua J (2011) Induction of *BAP1* by a moderate decrease in temperature is mediated by *ICE1* in Arabidopsis. Plant Physiol. **155:** 580-588