Activation of the mitochondrial unfolded protein response regulates the formation of stress

granules

Marta Lopez-Nieto[1,2], Zhaozhi Sun[3], Emily Relton[1,2], Rahme Safakli[4], Brian D. Freibaum[5], J Paul

Taylor[5], Alessia Ruggieri[3], Ioannis Smyrnias[4,*] and Nicolas Locker[1,2,*].

[1] Faculty of Health and Medical Sciences, School of Biosciences and Medicine, University of Surrey,

Guildford, United Kingdom.

[2] The Pirbright Institute, Pirbright, United Kingdom.

[3] Heidelberg University, Medical Faculty, Centre for Integrative Infectious Disease Research (CIID),

Department of Infectious Diseases, Molecular Virology, Heidelberg, Germany.

[4] Faculty of Health and Medical Sciences, School of Veterinary Medicine, University of Surrey,

Guildford, United Kingdom.

[5] Department of Cell and Molecular Biology, St. Jude Children's Research Hospital, Memphis,

Tennessee, USA

* Corresponding authors email: nicolas.locker@pirbright.ac.uk, i.smyrnias@surrey.ac.uk

Running title: the UPR[mt] regulates SG formation

Keywords: UPR[mt], stress granules, integrated stress response, mitochondrial stress response,

GADD34

Summary statement

We describe a novel crosstalk between the mitochondrial unfolded protein response and the

integrated stress response involving stress granules that protects cells from further stress.

Abstract

To rapidly adapt to harmful changes to their environment, cells activate the integrated stress response
(ISR). This results in an adaptive transcriptional and translational rewiring, and the formation of
biomolecular condensates named stress granules (SGs), to resolve stress. In addition to this first line
of defence, the mitochondrial unfolded protein response (UPR^{mt}) activates a specific transcriptional
programme to maintain mitochondrial homeostasis. We present evidence that SGs and UPR^{mt}
pathways are intertwined and communicate. UPR^{mt} induction results in eIF2α phosphorylation and the
initial and transient formation of SGs, which subsequently disassemble. The induction of GADD34
during late UPR^{mt} protects cells from prolonged stress by impairing further assembly of SGs.
Furthermore, mitochondrial functions and cellular survival are enhanced during UPR^{mt} activation when
SGs are absent, suggesting that UPR^{mt}-induced SGs have an adverse effect on mitochondrial
homeostasis. These findings point to a novel crosstalk between SGs and the UPR^{mt} that may
contribute to restoring mitochondrial functions under stressful conditions.

Introduction

Controlling the localisation and function of macromolecules is fundamental for many cellular functions and typically achieved by surrounding them with lipid membranes in organelles such as the nucleus, lysosomes, or mitochondria. In recent years, biomolecular condensates that lack membranes have become increasingly recognised as an alternative way to organise cellular components during stress. The dynamic formation of these condensates is supported by networks of protein-protein, protein-RNA and RNA-RNA interactions, generating high local concentration of RNA and protein components (Tauber et al., 2020). With such remarkable molecular plasticity, biomolecular condensates provide an ideal platform for the regulation of key processes, including mRNA metabolism or intracellular signalling, and are utilised by cells to rapidly adjust and rewire regulatory networks in response to various physiological and pathological triggers (Yoo et al., 2019).

Stress granules (SGs) are among the most characterised cytoplasmic biomolecular condensates and assemble in response to many stresses, including oxidative stress, heat shock, viral infection, proteasomal inhibition, endoplasmic reticulum (ER) stress and UV irradiation (Corbet and Parker, 2019; Hofmann et al., 2020). Capturing mRNAs and proteins, SGs are central to the reorganisation of cellular contents during stress. Stress sensing results in the activation of the integrated stress response (ISR) and a global inhibition of protein synthesis usually driven by phosphorylation of the eukaryotic translation initiation factor 2α (eIF2α), which results in the dissociation of mRNAs from polysomes and their accumulation in cytoplasmic RNP complexes (Hofmann et al., 2020). These RNPs are then recognised by aggregation prone RNA-binding proteins (RBPs), such as Ras-GTPase activating SH3 domain binding protein 1 (G3BP1) and T cell internal antigen-1 (TIA-1), which lead to further recruitment of multiple proteins containing low sequence complexity domains or intrinsically disordered regions. These mediate clustering and fusion events driven by multivalent interactions between their protein and RNA components, at the core of which G3BP1 acts as a key node that promotes RNA-protein, protein-protein and RNA-RNA interactions, ultimately resulting in liquid-liquid phase separation (LLPS) and SG formation (Corbet and Parker, 2019; Wang et al., 2018). SGs are highly dynamic, rapidly assembling to sequester the cytoplasmic stalled mRNAs and dissolving upon stress resolution to release stored mRNAs for future translation (Matheny et al., 2019; Namkoong et al., 2018). The dynamic nature of SGs is also reflected in their proposed stress-selective compositional, structural and functional heterogeneity, with different stresses proposed to induce at least 3 distinct types of SGs (Aulas et al., 2017; Hofmann et al., 2020). Type I and II SGs assemble via eIF2α-dependent and independent pathways, respectively, while type III SGs lack eIFs and are associated with cellular death (Reineke and Neilson, 2019). This suggests that compositionally heterogeneous SGs support specialized functions promoting survival or pro-death outcomes. By concentrating specific protein components and altering their cytoplasmic concentration, SGs have been proposed to alter the course of biochemical reactions and signalling cascades (Aulas et al., 2017; Hofmann et al., 2020). Furthermore, recent evidence hints that metabolic enzymes stored in SGs produce metabolites that can affect their stability (Begovich et al., 2020; Rollins et al., 2017; Saad et al., 2017). Importantly, dysregulation of pathways impacting SG clearance or LLPS is

increasingly associated with neuropathology, in particular amyotrophic lateral sclerosis and related
diseases (Wolozin and Ivanov, 2019). Moreover, SGs can be exploited by cancer cells to adapt to the
tumour microenvironment, with resident SG proteins aberrantly expressed in tumours (Anderson et
al., 2015). Finally, SGs are also proposed to be a crossroad between intracellular signalling and
antiviral responses by controlling the location and function of key sensors and effectors of innate
immunity (Eiermann et al., 2020).

In addition to this first line of defence, cells have evolved tailored quality control processes to protect
the functions of key organelles in times of stress. Mitochondria act as the cell's powerhouse,
generating energy through oxidative phosphorylation but are also critical for essential function as such
as calcium homeostasis, nucleotide metabolism and innate immunity regulation (Banoth and Cassel,
2018; Giorgi et al., 2018; Spinelli and Haigis, 2018). Key to maintaining or adapting mitochondrial
functions, the mitochondrial unfolded protein response (UPR^{mt}) is an evolutionarily conserved
mitochondria-to-nucleus signalling cascade, initially described as a transcriptional response to the
accumulation of misfolded mitochondrial proteins (Martinus et al., 1996; Zhao et al., 2002). Studies
have shown that activation of the UPR^{mt} is primarily regulated by the Activating Transcription Factor 5
(ATF5) and Heat Shock Factor 1 (HSF-1) (Fiorese et al., 2016; Smyrnias et al., 2019; Sutandy et al.,
2023; Wang et al., 2019), and engaged upon a wide range of mitochondrial insults, including
abnormal level of mitochondrial reactive oxygen species (ROS) (Wang et al., 2022b), dissipation of
mitochondrial membrane potential (Rolland et al., 2019), and reduced mitochondrial protein import
efficiency (Nargund et al., 2012). Activated UPR^{mt} serves to restore mitochondrial protein
homeostasis and promote cell survival in various biological processes, such as innate immune
signalling (Pellegrino et al., 2014) and ageing (Houtkooper et al., 2013), as well as adaptation to
pathophysiological conditions, such as neurodegenerative disorders (Sorrentino et al., 2017) and
cardiovascular diseases (Smyrnias et al., 2019; Wang et al., 2019). Notably, growing evidence
implicates the components of the ISR as central elements of the UPR^{mt} (Anderson and Haynes, 2020;
Fu et al., 2023; Quirós et al., 2017).

The ISR is an adaptive gene expression regulation programme activated in response to a range of
pathophysiological conditions, including hypoxia, amino acid deprivation, and viral infection (Brostrom
et al., 1996; Dever et al., 1992; Harding et al., 2003; Ron, 2002; Wek et al., 2006). ISR signalling is
initiated by the phosphorylation of eIF2α by four different kinases, which results in global translational
shut down to dampen cellular workload and accumulation of SGs, while specific transcripts involved in
stress adaptation remain preferentially translated. A key aspect of ISR regulation involves the fine-
tuning of eIF2α phosphorylation by the growth arrest and DNA-damage inducible protein (GADD34).
This stress-induced regulatory subunit of protein phosphatase 1 (PP1) mediates eIF2α
dephosphorylation, acting as a negative feedback loop to prevent prolonged ISR activation to ensure
stress resolution and resuming of protein synthesis (Harding et al., 2000; Lee et al., 2009; Novoa et
al., 2001). Studying the components of the ISR signalling has provided evidence that the ISR acts as
an essential precursor to the UPR^{mt} during mitochondrial stress. Specifically, each of the eIF2α
kinases has been implicated in response to stress associated with mitochondrial dysfunctions (Baker

et al., 2012; D'Souza and Minczuk, 2018; Guo et al., 2020; Samluk et al., 2019; Zhang et al., 2018). In

addition, ATF5, together with ATF4 and C/EBP homologous protein (CHOP), are main regulators of

the ISR that have emerged as essential mediators of a response to mitochondrial dysfunction similar

to the UPR[mt] in *C. elegans* (Fiorese et al., 2016; Jiang et al., 2020; Kaspar et al., 2021; Quirós et al.,

2017; Zhao et al., 2002). Moreover, UPR[mt] activation following mitochondrial dysfunction has been

linked to reduced protein synthesis within the mitochondria (Münch and Harper, 2016), and with

studies demonstrating that SGs impact metabolic processes, act as signalling hubs, and dock to

mitochondria, we propose that SGs may be implicated in the interface between UPR[mt] and ISR

signalling during mitochondrial stress.

Herein, we established a novel crosstalk between the UPR[mt] and SGs. Induction of UPR[mt] during

mitochondrial stress first results in an initial wave of SGs, which then disassemble potentially due to

an upregulation of SG disassembly chaperones. Next, the negative feedback loop mediated by

GADD34 protects cells against further stress insults and the assembly of eIF2α-dependent SGs.

Finally, we demonstrated that the core SG scaffold proteins G3BP1 and G3BP2 (G3BP1/2) negatively

affect mitochondrial respiration, morphology, ROS levels, and translation. Therefore, we propose a

model in which the dynamic assembly of SGs is an important component of the UPR[mt] to maintain

mitochondrial homeostasis, which itself can fine-tune further adaptation to stress by activating

GADD34. This provides further understanding of the cellular adaptation to adverse conditions and the

crosstalk between generalised and specialised stress responses.

Results

UPR[mt] activation by GTPP is associated with a transient SG assembly and translational shut-off

With previous studies suggesting an interplay between the UPR[mt] and the ISR, and given the central

role of SGs in the cellular stress response, we investigated the impact of mitochondrial stressors on

the SG pathway in more details. First, we tested the induction of UPR[mt] signalling in U2OS cells

following mitochondrial stress using the mitochondrial HSP90 inhibitor gamitrinib-

triphenylphosphonium (GTPP), which results in protein misfolding in the mitochondria and induction of

mitochondrial chaperones and proteases (Kang et al., 2009). The levels of typical transcripts induced

upon UPR[mt] and ISR activation were measured using RT-qPCR (Fig 1A). We observed a clear

induction of CHOP and ATF4 after 4 h of GTPP treatment, which are related to ISR activation, as well

as HSPD1 and LONP1, indicating induction of the UPR[mt]. ATF5 acts as a key mediator of the

mammalian UPR[mt] signalling, translocating to the nucleus upon stress to activate a specific

transcriptional cascade (Fiorese et al., 2016). Consistently, siRNA-mediated silencing of ATF5

expression significantly reduced the transcription of all GTPP-induced UPR[mt] markers, but not ISR

markers, further supporting the role of ATF5 as UPR[mt] inducer in our model (Fig 1A).

Next, we tested the impact of UPR[mt] activation on SG assembly by treating U2OS cells with GTPP for

up to 8 hours. Cells were fixed at the indicated time points and stained with anti-G3BP1 and anti-

eIF3η antibodies to detect SG assembly (Fig 1B). Sodium arsenite was used as positive control for

SG assembly as it induces eIF2α-dependent SGs by activating the eIF2α kinase HRI (Taniuchi et al.,

2016). G3BP1 and eIF3η remained diffused in the cytoplasm in DMSO-treated cells and, as expected,

arsenite treatment resulted in the assembly of SGs reflected by the accumulation of G3BP1/eIF3η into

cytoplasmic foci. Interestingly, approximately 25% of cells were positive for SGs at 2 h post GTPP

treatment (Fig 1C). The percentage of SG-positive cells decreased gradually until being undetectable

at 8 h post treatment. To further explore the characteristics of GTPP-induced SGs, we used recently

described small molecules inhibitors of G3BP1 condensation that impair SG assembly and induce

their disassembly (Freibaum et al., 2024). Pre-treatment of cells with the active inhibitor G3Ib

completely blocked GTPP-induced SG formation, while the inactive compound G3Ib' had no impact.

This confirmed that SGs induced following UPR^mt activation, like canonical SGs, are dependent on

G3BP1 condensation for their assembly (Sup Fig 1). Thus, mitochondrial stress following GTPP

treatment in U2OS cells activates the UPR^mt and induces an early and transient phase of SG

assembly.

Several studies have suggested a connection between mitochondrial function and cytosolic protein

translation (Lu and Guo, 2020; Topf et al., 2018). Given that formation of SGs is preceded by an

inhibition of translation, we examined the impact of GTPP on global translational control at a single

cell level using ribopuromycylation assays (David et al., 2012). Briefly, cells were treated with arsenite

as positive control or GTPP for the indicated time and nascent polypeptides were labelled with

puromycin, a tRNA analogue, followed by an emetine pulse to avoid the release of puromycin-nascent

chains from ribosomes. Cells were then fixed and stained with anti-puromycin to detect

puromycylated nascent peptide chains, and with the SG markers G3BP1 and eIF3η (Sup Fig 2).

Overall, GTPP treatment resulted at in strong translation repression at 2, 4, 6 and 8h post treatment

when compared to untreated cells (Fig 1D). In addition, in treated cells displaying SGs, translational

repression was more potent than in cells not forming SGs (Fig 1D). Therefore, these results indicate

that UPR^mt activation by GTPP significantly attenuates cytosolic translation at 2 h post treatment and

this shut off is at least in partial uncoupled from SG formation and maintained over time.

<u>GTPP-mediated mitochondrial stress leads to ISR activation by PERK</u>

Phosphorylation of eIF2α is central event in the ISR signalling and a trigger of SG formation

(Kedersha et al., 2002). Thus, we measured phosphorylation levels of eIF2α as well as protein and

RNA expression levels of ISR markers during early UPR^mt activation. Phosphorylation of eIF2α was

detected within 30 minutes of GTPP treatment, which was followed by an increase of ATF4 and

GADD34 protein levels at 1 h and 3 h post treatment (Fig 2A). Similarly, CHOP and GADD34 mRNA

levels were also highly upregulated following GTPP treatment, for 2 and 3 h, respectively (Fig 2B).

These results indicate that UPR^mt induction by GTPP signals through an early activation of canonical

ISR.

We next tested which of the four eIF2α kinases is activated during UPR^mt induction. We first

hypothesized that the double-stranded RNA (dsRNA)-activated protein kinase (PKR) might be

activated upon GTPP treatment due to mitochondrial dsRNA leakage occurring as a consequence of

mitochondrial stress and membrane damage (Kim et al., 2018). However, mitochondrial dsRNA was

not detected after UPR^{mt} induction (Sup Fig 3). Supporting this result, the expression of the interferon-
stimulated genes such as CCXCL10, CCL5, MX1 and OAS1, reported to be induced upon PKR
activation (Pindel and Sadler, 2011; Radetskyy et al., 2018), was not upregulated during early or late
UPR^{mt} (Fig 2C). Thus, PKR is unlikely to be the kinase responsible for eIF2α phosphorylation upon
GTPP treatment. We next investigated the potential role of PKR-like ER kinase (PERK) and general
control non-derepressible protein 2 (GCN2). PERK is activated in response to ER stress and has
been shown to modulate mitochondrial homeostasis during stress (Almeida et al., 2022), while GCN2
is activated by amino acid depravation or UV light radiation (Deng et al., 2002). We used the
pharmacological inhibitors GSK2606414 and A-92 to inhibit PERK and GCN2 activity in U2OS cells,
respectively. We first confirmed that they reduced eIF2α phosphorylation upon DTT treatment or UV
light exposure (Fig 2D). In response to 2 h GTPP, the GCN2 inhibitor A-92 did not reduce eIF2α
phosphorylation, whereas the PERK inhibitor GSK2606414 decreased it by more than 2-fold,
suggesting that PERK is the main eIF2α kinase responsive to GTPP. To confirm that PERK activation
is responsible for SG assembly during UPR^{mt} activation, cells were treated with PERK inhibitor
followed by GTPP treatment. This resulted in the complete lack of SGs further confirming that GTPP-
induced SGs are dependent on PERK-driven eIF2α phosphorylation (Fig 2E, 2F). Overall, these
results suggest that activation of the UPR^{mt} results is rapidly followed by an early phase characterised
by ISR activation through PERK and SG assembly (0.5-3 h), and a late phase where SGs
disassemble and UPR^{mt} genes are expressed (4-8 h).

<u>Mitochondrial stress impairs the assembly of eIF2α-dependent SGs</u>

As GTPP-mediated mitochondrial stress and UPR^{mt} activation resulted in SG assembly only during
the early and not at the later stages of stress, we hypothesized that the UPR^{mt} may limit the ability of
cells to form SGs during prolonged stress. To test this, cells were stimulated for 2, 4, 6 or 8 h with
GTPP and then challenged with arsenite for 30 min to induce the formation of SGs (Fig 3A and Sup
Fig 4). As expected, in absence of pre-treatment with GTPP, arsenite induced the assembly of SGs in
almost all cells. Interestingly, only some GTPP-treated cells formed SGs after arsenite treatment at
the indicated timepoints. Detailed analysis of U2OS cells revealed that pre-treatment with GTPP
reduced arsenite-induced SG assembly compared to arsenite-only stimulated cells from 92% to 50%,
14%, 28% and 35% after 2, 4, 6 or 8 h treatments, respectively (Fig 3B). These data indicate that
UPR^{mt} induction by GTPP inhibits subsequent arsenite-induced SG formation. SG assembly is a
multistep process by which microscopically invisible "seeds" first coalesce upon stress, and then fuse
into larger aggregates during a growth phase (Wheeler et al., 2016). Thus, we next examined whether
the SG maturation process is also affected when UPR^{mt} is pre-activated. To address this, the average
size of SGs per cell and number of SGs per cell were compared in cells pre-treated with GTPP and
non-treated (Fig 3C). The analysis showed that while a 2 h GTPP treatment does not impact the area
of arsenite-induced SGs, longer pre-treatment results in much smaller ones significantly smaller (in
average 0.47, 0.53, 0.5 μm^2, for 4 h, 6 h and 8 h respectively) compared to non-treated cells (in
average 0.72 μm^2). Likewise, less arsenite-induced SGs were detectable in GTPP pre-treated cells.

These results suggest that SG maturation process and fusion into larger granules may be altered by
GTPP during late UPR^{mt}.

Given that eIF2α phosphorylation drives SG formation, we next assessed how UPR^{mt} activation
impacts on arsenite-induced eIF2α phosphorylation. Cells were treated as mentioned above and
levels of basal and phosphorylated eIF2α measured by immunoblotting. Cells treated with arsenite
displayed high levels of eIF2α phosphorylation, while GTPP treatment slightly increased eIF2α
phosphorylation as shown previously (Fig 3D). Interestingly, when subjected to arsenite treatment,
GTPP pre-treated cells showed lower levels of arsenite-induced eIF2α phosphorylation. Specifically,
phosphorylated levels were reduced approximately 3 to 4-fold compared to arsenite-treated cells in all
timepoints.

To confirm these results, UPR^{mt} was induced with another mitochondrial stressor, paraquat, which
produces high amounts of superoxide and interferes with the electron transport chain (Cocheme and
Murphy, 2008). Treatment of cells with paraquat for 48 h increased by 2 to 3-fold the mRNA levels of
ATF4, CHOP, GADD34 and DNAJA3 compared to non-treated cells, confirming ISR and UPR^{mt}
activation (Sup Fig 5A). Next, we examined how paraquat stimulation impacts on SG assembly and
whether pre-treatment with paraquat affects the ability of cells to form arsenite-induced SGs by
immunofluorescence. Paraquat treatment for up to 48 h did not induce SG formation, and G3BP1 or
eIF3η remain diffused in the cytoplasm (Sup Fig 5B). However, prior induction of UPR^{mt} by treatment
with paraquat for 48 h decreased percentage of cells containing arsenite-induced SGs, from about
96% to 65% (Fig 3E, 3F and Sup Fig 5C). Similar to our previous observation with GTPP, paraquat
pre-treatment resulted in smaller and less numerous arsenite-induced SGs when compared to non-
treated (0.53 μm^2 compared to 0.85 μm^2, and 57 vs 32, respectively) (Fig 3G). These results indicate
that paraquat impairs SG formation and limits their maturation into larger SGs. Paraquat treatment
also resulted in a slight increase in eIF2α phosphorylated levels, supporting ISR induction, and pre-
treatment with paraquat also limited the levels of arsenite-induced eIF2α phosphorylation compared to
non-treated cells (Fig 3H). Overall, activation of UPR^{mt} using two different stressors results in the
inhibition of arsenite-induced eIF2α phosphorylation and consequently prevents SG formation during
prolonged stress. Altogether, these data strongly propose that SG assembly is modulated by prior
induction of the UPR^{mt}.

Given that ATF5 is one of the central regulators in UPR^{mt} activation, we explored its impact on SG
assembly (Fiorese et al., 2016). To study this, cells were transfected with siRNA targeting ATF5 and
SG formation was monitored by immunofluorescence during GTPP treatment for 4 h and with or
without arsenite (Sup Fig 6A). ATF5 silencing did not affect SG assembly nor disassembly after 4 h of
GTPP treatment, around 20% of cells were positive for SGs (Sup Fig 6B). Moreover, assembly of
arsenite-induced SGs in GTPP pre-treated cells was impaired when ATF5 was silenced, with no
differences compared to non-silencing control cells. However, quantitative analyses revealed that
GTPP-induced SGs were significantly smaller when ATF5 was silenced compared to control cells,
while no difference was observed in the number of granules per cell (Sup Fig 6C). In contrast,
whereas ATF5 silencing did not altered the size of arsenite-induced SGs, it resulted in a significant

reduced number of granules compared to non-silencing cells (Sup Fig 6D). SGs differ in their
dynamics, assembly and composition depending on the stress (Aulas et al., 2017), therefore these
findings suggest that the ATF5-mediated UPR^{mt} transcriptional reprogramming alters SG dynamics in
a stress-specific manner.

GADD34 is required for the inhibition of *de novo* eIF2α-dependent SG assembly by preventing eIF2α
phosphorylation

We observed that UPR^{mt} activation by either GTPP or paraquat results in the upregulation of GADD34
(Fig 2A, 2B and Sup Fig 5A), the negative regulator of the ISR, which in complex with PP1
dephosphorylates eIF2α to resume translation. Therefore, we hypothesized that increased levels of
GADD34 could limit eIF2α phosphorylation in response to a consecutive stress and consequently SG
assembly. To test this hypothesis, we generated U2OS GADD34 KO cell clones using a
CRISPR/Cas9 approach (Sup Fig 7A). Control cell clones (Ctrl) were generated for comparison using
a non-targeting crRNA. Ctrl and GADD34 KO cells were stimulated with GTPP for 4 h prior arsenite
treatment and SG assembly was detected by immunofluorescence (Fig 4A and Sup Fig 7B). As
observed previously, prior induction of UPR^{mt} by GTPP impaired the assembly of arsenite-induced
SGs in Ctrl cells. In contrast, we observed no difference in the assembly of SGs in GADD34 KO cells
irrespectively of GTPP pre-treatment. While only 17% of Ctrl cells pre-treated with GTPP were
positive for arsenite-induced SG, 97% of GADD34 KO cells contained SGs in the same conditions
(Fig 4B). Similar results were obtained with paraquat pre-treatment with 66% Ctrl cells and 98%
GADD34 KO cells displaying SGs in response to arsenite treatment (Fig 4C, 4D and Sup Fig 7C).

These results were supported by reduced levels of phosphorylated eIF2α in GTPP and paraquat pre-
treated Ctrl cells compared to GADD34 KO cells (Fig 4E, 4F). In Ctrl cells, either GTPP or paraquat
treatment followed by arsenite stimulation resulted in reduced eIF2α phosphorylation levels compared
to stimulation with arsenite only. In contrast, in GADD34 KO cells, the induction of eIF2α
phosphorylation was comparable between stimulation with arsenite and GTPP or paraquat treatments
followed by arsenite stimulation. Therefore, GADD34 KO rescued the ability of cells to phosphorylate
eIF2α following UPR^{mt} induction. Overall, these results indicate that UPR^{mt} activation results in
increased levels of GADD34 expression, which prevent eIF2α phosphorylation and thereby impair the
ability of cells to assemble SGs during stress adaptation.

GTPP-mediated UPR^{mt} activation differentially modulates the assembly of eIF2α-independent SGs

SG assembly can also be triggered following eIF2α-independent inhibition of translation, for example
by blocking eIF4A or mTOR activity (Aulas et al., 2017). Thus, we examined whether UPR^{mt} activation
alters the assembly of eIF2α-independent SGs following treatment with the eIF4A inhibitor silvestrol
(Bordeleau et al., 2008). To test this, U2OS cells were stimulated with GTPP for 6 h, then treated with
or without silvestrol for 1 h and SG assembly was analysed by immunofluorescence (Fig 5A). As
expected, silvestrol induced formation of SGs in nearly all cells. Interestingly, pre-treatment with
GTPP resulted in a reduction of the percentage of cells displaying silvestrol-induced SGs, from 94%
to 72% (Fig 5B). In contrast, pre-treatment with paraquat did not impair the assembly of silvestrol-

induced SGs. (Fig 5D, 5E). However, features of SGs appeared to be altered by both mitochondrial
stressors, as reported before for arsenite-induced SGs. Detailed analysis revealed that GTPP and
paraquat treatment resulted in a reduction in average size of silvestrol-induced SGs, from 0.98 μm^2 to
0.9 μm^2 and 0.66 μm^2, respectively, and increased average number of SGs, from 18 to 22 and 39,
respectively, when compared to non-treated cells (Fig 5C, 5F). These results indicate that UPR^{mt}
activation by GTPP impairs the ability of cells to assemble eIF2α-independent SGs, although not to
the same extent as eIF2α-dependent SGs, and that both stressors affect their maturation and fusion
process. Taken together, these data suggest that UPR^{mt} activation impact on SG assembly and
dynamics both in an ISR-dependent and -independent manner.

SGs are highly dynamic and disassemble upon stress removal. Recently, the chaperone Hsp90 was
reported to initiate SG disassembly by interacting with and activating the dual specificity tyrosine-
phosphorylation-regulated kinase 3 (DYRK3), leading to the phosphorylation of yet unknown targets.
(Mediani et al., 2021). Given that GTPP-induced SGs disassemble after 4 h treatment (Fig 1C) and to
further examine how UPR^{mt} activation modulates assembly and dynamics of eIF2α-independent SGs,
we measured the mRNA levels of DYRK3 and the stress-inducible and constitutively expressed
Hsp90 forms, HSP90AA1 and HSP90AB1, respectively. Upon UPR^{mt} induction by GTPP, the level of
HSP90AA1 mRNA was significantly increased by approximately 4 to 6-fold in all time points
compared to non-treated cells, while Hsp90AB1 and DYRK3 were slightly upregulated by 2 to 3-fold
at 6 and 8 h GTPP treatment (Fig 5G). In contrast, the mRNA levels of these proteins remained
unchanged upon UPR^{mt} induction by paraquat (Fig 5H). These results are consistent with our
previous results where silvestrol-induced SG assembly was altered by GTPP treatment but not by
paraquat, indicating that GTPP but not paraquat mediated UPR^{mt} activation enhances SG
disassembly process. Altogether, these data suggest that UPR^{mt} activation by GTPP results in an
initial SG assembly which then rapidly disassemble potentially due to an enhancement in the SG
disassembly process.

<u>Cells unable to assemble SGs during stress present improved mitochondrial functions during UPR^{mt}</u>
<u>activation</u>

To investigate the impact of reduced SG formation during mitochondrial stress, we first examined the
effect of GTPP on mitochondrial morphology in U2OS G3BP1/2 double knockout cells (G3BP1/2
dKO) lacking the ability to form SGs compared to control cells (Sup Fig 8) (Yang et al., 2020). Wild-
type (WT) and G3BP1/2 dKO U2OS cells were treated with GTPP for 2 h, the time at which we
previously detected the appearance of SGs, and stained for the mitochondrial marker TOMM20 (Fig
6A). Confocal imaging of DMSO-treated cells showed a typical tubular mitochondrial network while
those treated with GTPP contained round and fragmented mitochondria. Subsequent detailed
analysis indicated that following GTPP treatment, 91.5% of WT cells presented fragmented
mitochondria, compared to 75% of G3BP1/2 dKO cells (P < 0.012) (Fig 6B). These results indicate
that GTPP-induced defective mitochondrial dynamics is partially restored when the SG scaffolding
proteins G3BP1/2 are absent. Alterations in mitochondrial morphology are associated with
mitochondrial ROS (mtROS) and in particular, high mtROS levels promote mitochondria

fragmentation or swelling (Brillo et al., 2021). We then studied whether GTPP results in mtROS

production and how the absence of SGs impacts on mtROS levels. WT and G3B1/2 dKO cells were

stimulated with GTPP for 2 h and incubated with MitoSOX fluorogenic dye which specifically detects

mitochondrial superoxide in live cells. Interestingly, while GTPP treatment induced high production of

mtROS in WT cells, 1.7-fold compared to non-treated, GTPP-induced mtROS levels were significantly

lower in G3BP1/2 dKO cells (Fig 6C and Sup Fig 9). These data demonstrate that loss of G3BP1/2

partially prevents mitochondria fragmentation and reduces mtROS production during mitochondrial

stress.

To further characterise the functional consequences of reduced SG formation during mitochondrial

stress, we compared mitochondrial respiration upon UPR^{mt} activation in WT and G3BP1/2 dKO cells.

Cells were treated with GTPP for 2 h and measured mitochondrial oxygen consumption rate (OCR)

using a Seahorse flux analyser (Fig 6D). As expected GTPP impaired mitochondrial functions in WT

cells resulting in decreased basal, maximal respiration, ATP production-coupled respiration and spare

respiratory capacity compared to untreated cells (Fig 6E, 6F). In contrast, GTPP-stressed G3BP1/2

dKO cells that cannot form SGs presented with significantly improved mitochondrial respiration when

compared to WT cells.

Protein synthesis is regulated during stress conditions to preserve cellular resources. Given that

GTPP stimulation results in arrest of cytosolic translation (Fig 1D), we next investigated how GTPP

impacts on translational control when G3BP1/2 are absent. G3BP1/2 dKO cells were treated with

GTPP, or arsenite as a control, for the indicated times and ribopuromycylation assay was performed

as described above to measure the levels of translation in single cells. As shown in previous studies

(Ohn et al., 2008), arsenite treatment induced a potent translational shut off both in WT and G3BP1/2

dKO cells, confirming that SG formation is not required for translation repression (Fig 7A, 7B).

Interestingly, in G3BP1/2 dKO cells GTPP treatment did not alter the levels of translation which

contrasts with WT cells (Fig 1D and Fig 7A, 7B), suggesting that the translational shut off induced by

GTPP is dependent on the presence of G3BP1/2 and/or the formation of UPR^{mt}-induced SGs. Next,

we hypothesise that the impact of GTPP treatment on protein synthesis could be linked to specific

translational control at the mitochondrial level. To further assess this, we monitored the impact of

GTPP treatment, on specific mitochondrial translation using a SUNSET-based approach (Yousefi et

al., 2021). Cells were incubated in methionine-free media and cytosolic translation was selectively

halted by cycloheximide treatment, before adding the alkyne-containing non-canonical amino acid

L-homopropargylglycine (HPG). This allows HPG to be specifically incorporated into mitochondrial

translation products, instead of methionine, which can be visualized by copper-catalyzed

cycloaddition reaction (click) to azide-containing fluorescent dyes (Fig 7C, 7D, 7E). First, we

confirmed the validity of this approach as the HPG-derived fluorescence signal colocalise with

mitochondrial TOMM20 signal. Second, when cells were left in methionine-containing media or

incubated with the mitochondrial translation inhibitor chloramphenicol (CHL), HPG signal disappeared

showing that HPG is only incorporated into mitochondrial nascent peptides, and its fluorescence

signal correlates with mitochondrial translation (Fig 7C). Quantification of HPG incorporation revealed

that UPR^{mt} activation by GTPP results in decreased mitochondrial translation in WT cells.
Interestingly, G2BP1/2 dKO cells also presented a significant reduction in mitochondrial translation
after stimulation with GTPP (Fig 7D, 7E), indicating that absence of G3BP1/2 does not affect
mitochondrial translation during UPR^{mt} as compared to cytosolic translation. Taken together, these
results demonstrated that the assembly of UPR^{mt}-induced SGs is associated with altered
mitochondrial dynamics, mtROS production without impairing mitochondrial translation during stress.

<u>Loss of SG scaffolding proteins G3BP1/2 results in enhanced cell survival during chronic UPR^{mt}</u>
<u>activation</u>

Mitochondrial homeostasis is crucial for cell survival and prolonged mitochondrial dysfunction can
lead to cell death (Van Houten et al., 2016). We therefore examined how the absence of SGs impacts
on cell survival following chronic UPR^{mt} activation. The kinetics of cell viability after GTPP treatment in
WT and G3BP1/2 dKO U2OS cells was monitored for 28 h using the IncuCyte live-cell imaging
system. Interestingly, we detected increased resistance to GTPP-induced cell death in G3BP1/2 dKO
cells that are unable to form SGs, when compared to WT cells (Fig 8). The number of WT dead cells,
indicated by round-shape orange-fluorescent cells, increased significantly compared with WT DMSO
treated cells at 11 h of GTPP treatment (Fig 8A, 8B). In contrast, we did not detect a significant
increase in G3BP1/2 dKO dead cells until 17 h after GTPP treatment compared to DMSO-treated
cells. These results show that cells unable to form SGs present a prolonged cell death following
UPR^{mt} activation compared to WT cells, suggesting that GTPP-induced SGs are detrimental for
cellular survival.

Discussion

The disruption of mitochondrial homeostasis is associated with several pathologies, such as
neurodegeneration, cancer, aging, and cardiovascular diseases (Chen et al., 2023; Norat et al., 2020;
Quirós et al., 2016; Suomalainen and Battersby, 2018). The UPR^{mt} is an evolutionarily conserved
homeostatic pathway engaged upon a wide range of mitochondrial insults, including mitochondrial
protein misfolding (Zhao et al., 2002), reduced mitochondrial protein import efficiency (Nargund et al.,
2012), dissipation of mitochondrial membrane potential (Donehower et al., 2019), and dysregulated
levels of reactive oxygen species (ROS) (Wang et al., 2022b). UPR^{mt} activation triggers a broad and
protective transcriptional cascade, through enhancing mitochondrial folding capacity and triggering a
mitochondrial recovery program, alleviating neurodegenerative and cardiovascular diseases
symptomatology (Smyrnias et al., 2019; Sorrentino et al., 2017), promoting resistance to pathogen
infection (Mahmud et al., 2020; Pellegrino et al., 2014), and prolonging lifespan (Munkácsy and Rea,
2014). Studies in mammals have shown that the ISR and UPR^{mt} are intertwined; however, how SGs –
which are downstream effectors of the ISR – modulate the outcome of the UPR^{mt} and how their
assembly is regulated upon mitochondrial stress is unknown. Evidence supporting a functional
interplay between SGs and mitochondria include the impaired metabolism linked to altered SG
dynamics in several neurodegenerative diseases, the regulation of SG stability by metabolites

produced by resident SG enzymes such as Sam1 or pyruvate kinase, and the regulation of
mitochondrial permeability and lipid fatty acid oxidation by starvation-induced SGs (Begovich et al.,
2020; Cereghetti et al., 2021; Rollins et al., 2017; Saad et al., 2017; Wang et al., 2022a). SGs can
physically contact different membrane-bound organelles, including lysosomes or the ER, and
starvation-induced SGs contact with mitochondria in human cells (Amen and Kaganovich, 2021; Lee
et al., 2020; Liao et al., 2019). Furthermore, numerous mitochondrial proteins have been associated
with SGs (Amen and Kaganovich, 2021; Aoyama-Ishiwatari et al., 2021; Paget et al., 2023). Our study
identifies SGs as novel signalling components associated with the mammalian UPR^{mt} and SG
dynamics as a key for maintaining mitochondrial function during mitochondrial stress. We show that
UPR^{mt} activation by GTPP or paraquat results in PERK-mediated activation of the ISR, accompanied
with transient SG formation, in the case of GTPP. Moreover, UPR^{mt}-induced upregulation of the
GADD34 protects cells against persistent SG assembly and preventing SG formation improved
mitochondrial functions upon UPR^{mt} induction.

ATF5 has been proposed to drive the protective effect of UPR^{mt} in different diseases models such as
osteoarthritis (Zhou et al., 2022), heart failure (Smyrnias et al., 2019; Wang et al., 2019), and protect
host during enteric infection (Chamseddine et al., 2022). In our experimental system, we show that
ATF5 is essential for the upregulation of the UPR^{mt} markers in cells subjected to mitochondrial stress
(Fig 1), but while it was not involved directly in SG formation, GTPP-induced SGs were smaller when
ATF5 was silenced (Sup Fig 6). Thus, impaired upregulation of UPR^{mt} genes may affect the
composition and maturation of SGs, as well as their function. Recently, a study by Sutandy *et al.*
proposed that the accumulation of mitochondrial protein precursors and ROS in the cytosol results in
nuclear translocation of heat shock factor 1 (HSF1) and activation of UPR^{mt} markers. Importantly, they
demonstrated that ATF5 synergises with HSF1, which affects late activation of the UPR^{mt} (Sutandy et
al., 2023). Thus, we cannot exclude the possibility that other transcription factors such as HSF1 may
contribute to SG dynamics during conditions of mitochondrial stress and induction of the UPR^{mt}.

The disassembly of SGs is crucial for cellular homeostasis since persistent SGs can promote the
formation of aberrant aggregates, which are implicated in several neurodegenerative diseases (Baron
et al., 2013; Repici et al., 2019; Vanderweyde et al., 2012). SGs induced by GTPP begin to
disassemble post 3 h treatment and completely disappear by 8 h (Fig 1C). We first hypothesized that
the increased expression of GADD34 due to ISR activation might be responsible for SG disassembly
in these mid/late phases. However, GADD34 KO cells cleared SGs with a comparable kinetic to
control cells (Fig 5), indicating that a factor other than GADD34 is involved in SG disassembly after
GTPP treatment. In addition, GTPP treatment resulted in increased expression of the kinase DYRK3,
and HSP90, a major chaperone promoting SG disassembly of SGs through binding and stabilising
DYRK3 (Fig 5) (Mediani et al., 2021; Wippich et al., 2013). Upregulation of SG disassembly
chaperones was not detected with paraquat, consistent with this stressor not inducing SG formation.
Interestingly, UPR^{mt} induction by either GTPP or paraquat prevented *de novo* assembly of arsenite-
induced SGs by impairing eIF2α phosphorylation. This effect was rescued in GADD34 KO cells,
indicating that GADD34 upregulation is responsible for the inhibition of SG assembly following a

second stressor (Fig 4). This result is in agreement with recent work demonstrating that sustained

GADD34 levels upon ISR activation reduces cell responsiveness to a second stress pulse, promoting

cell adaptation (Klein et al., 2022; Magg et al., 2024). Our data suggest that UPR^{mt} employs GADD34

as a negative regulator of the ISR to fine-tune cellular stress responses and maintain mitochondrial

function under these conditions.

UPR^{mt} activation by GTPP also inhibited the formation of eIF2α-independent SGs, and, the size and

number of both eIF2α-dependent and independent SGs were altered in cells treated with both GTPP

and paraquat compared to non-treated. This suggests that UPR^{mt}-induced stress could alter SG

biogenesis by limiting the growth of small SGs prior to their fusion into bigger aggregates. We

speculate that the known GTPP and paraquat effects on oxidative phosphorylation could affect SG

biogenesis, whose fusion and growth steps are ATP-dependent (Jain et al., 2016). Further

experiments are needed to fully understand how energy depletion affects SG biogenesis during

mitochondrial stress.

Our results reveal that cells subjected to mitochondrial stress activate the UPR^{mt} and engage

mechanisms to disassemble either SGs induced by mitochondrial stress and/or prevent *de novo*

assembly of SGs, suggesting that clearance of SGs is crucial to the maintenance of mitochondrial

homeostasis. Consistently, we show that GTPP-induced SGs impaired mitochondrial morphology and

respiration, whereas cells unable to form SGs exhibited improved mitochondrial function, including

translation during stress (Fig 6). In addition, absence of SGs improved cell survival during

mitochondrial stress (Fig 8). Furthermore, GTPP treatment blocked cytosolic translation uncoupled

from SG formation and when G3BP1/2 were absent, this translational shut off was rescued (Fig 7).

Besides eIF2α phosphorylation, protein synthesis is also modulated by the mammalian target of

rapamycin (mTOR) signalling pathway. Activation of mTOR leads to hyperphosphorylation of 4E-BP

which allows the correct assembly of eIF4F complex into the 5'-cap and thereby translation initiation.

Insufficient nutrients or energy result in mTOR inhibition, preventing cap-dependent translation

(Haghighat et al., 1995; Zoncu et al., 2011). UPR^{mt} activation by GTPP reduces ATP production-

coupled respiration (Fig 6F) which may inhibit mTOR resulting in translation inhibition independent of

SG formation. G3BP1/2 have been described as anchors of the tuberous sclerosis complex (TSC) to

lysosomes and supress mTOR activity upon starvation, thus, G3BP1 silencing results in mTOR

hyperactivation during those conditions (Prentzell et al., 2021; Rehbein et al., 2021). Therefore, an

hyperactivation of mTOR signalling pathway in G3BP1/2 dKO cells could explain the unaltered

translation levels after UPR^{mt} activation. Further studies are required to unravel the molecular

mechanisms underlying translational control during UPR^{mt} activation and disentangle the effect driven

by SG assembly from those of G3BP1 itself.

Questions remain as to the functional link between inhibition of SG assembly and mitochondrial

homeostasis following UPR^{mt} activation. Several studies support a protective role for SGs, through

impairing pro-apoptotic JNK signalling (Arimoto et al., 2008; Kitajima et al., 2023), sequestration of

caspases (Fujikawa et al., 2023) or suppressing production of ROS (Park and Shin, 2023; Takahashi

et al., 2013) and they have been described as "shock absorbers" dampening excessive innate

immune response to viral infection (Paget et al., 2023). In contrast, persistent SGs are implicated in
pro-death functions during chronic stress conditions as well as neurodegenerative diseases or tumour
chemotherapy (Reineke et al., 2018; Reineke and Neilson, 2019; Schwed-Gross et al., 2022). Future
studies are required to further compositionally characterise SGs, and their functions, during
mitochondrial stress.

Overall, our findings support a model in which mitochondrial stress-induced SGs are formed as a
consequence of the associated ISR activation. Subsequently, cells activate homeostatic mechanisms
to disassemble SGs, *i.e.* the UPR^{mt}, as their persistence may compromise mitochondrial recovery.
This could occur by through various mechanisms, including physically preventing efficient
mitochondrial dynamics, recruiting mitochondrial or related proteins to SGs, preventing translation of
essential mitochondrial proteins, or a combination of these. This underscores a novel functional
interplay between the ISR and UPR^{mt} and further work should aim at revealing the impact of SG
signalling during UPR^{mt}, how UPR^{mt}-induced SGs, and the sequestration of specific resident proteins,
impact on mitochondrial homeostasis. This will open further opportunities for alternative strategies for
the treatment of human diseases with compromised mitochondrial function and associated with an
impaired SG dynamics.

Material and methods

<u>Cell culture and gene silencing</u>
U2OS, U2OS G3BP1/2 double knock out (dKO), U2OS control (Ctrl) and GADD34 KO cell lines were
grown in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum,
100 U/ml penicillin, 100 g/ml streptomycin and 1% L-glutamine in a humidified incubator at 37ºC and
5% CO_2 environment. U2OS G3BP1/2 dKO cells have been described previously (Burke et al., 2020).

Silencing of ATF5 was performed using Lipofectamine-RNAiMax transfection reagent (Thermo
Fisher). U2OS cells were incubated with the transfection mix containing either siRNA negative control
(Invitrogen, #4390843) or siRNA targeting ATF5 (Invitrogen, ID s22425) at a final concentration of 10
nM for 48 h. Next, cells were incubated with subsequent treatments and analysed as described
below.

<u>Generation of U2OS GADD34 KO cell clones</u>
U2OS GADD34 KO cells were generated using a two-crRNA strategy to delete a portion of the
PPP1R15A locus encoding GADD34. crRNAs cutting sequences in *PPP1R15A* exon 2 (IDT,
Hs.Cas9.PPP1R15A.AC: AltR1 / AUC ACU GGG UUG GCA CCA CCG UUU UAG AGC UAU GCU /
AltR2) and exon 3 (IDT, custom design: AltR1 / GAG GCG UGG CUG AGA CCA ACG UUU UAG
AGC UAU GCU / AltR2) were purchased to Integrated DNA technologies (IDT). Individual crRNAs
and tracrRNA (IDT) were resuspended in 1 x TE buffer (10 mM Tris, 0.1 mM EDTA, pH 7.5) to a final
concentration of 200 µM and hybridized at equimolar amounts by heating to 95 °C for 5 min and slow
stepwise cooling (1°C/sec) to room temperature. Pre-assembled crRNA/Cas9 ribonucleoprotein
complexes (RNPs) were obtained by gently mixing 1.2 µl of hybridized RNAs with 17 µg of

recombinant Alt-R Cas9 protein (IDT) and 2.1 µl of PBS, followed by 20 min incubation at room

temperature.

U2OS cells (1×10^6) were washed with PBS, resuspended in 91 µl nucleofection solution from Cell

Line Nucleofector Kit V (Lonza) and mixed with 2.5 µl of each pre-assembled RNPs and 4 µl

electroporation enhancer (Integrated DNA technologies). Nucleofection was performed using an

Amaxa 2b nucleofector (Lonza) and pre-set program X-01. Nucleofected cells were incubated for 48 h

prior to clonal amplification and screening for homozygous clones using target-specific PCR (GoTaq

Hot Start, Promega) on genomic DNA using the following primers: PPP1R15A_forward: 5'-

ATCACACCGGGAGTGTTGTC-3', PPP1R15A_reverse: 5'-AGACGCTGCTCGCTACAAAT-3')

flanking sequences targeted by crRNAs. After genotyping, expression levels of GADD34 in 2 control

and 5 GADD34 KO single cell clones in response to 2 µM thapsigargin (Biotrend) treatment for 3 h

was assessed by Western blotting. After characterization and for further experiments, cell pools were

generated by mixing the clones in equal proportions.

<u>Chemical treatments</u>

For UPR^mt induction, GTPP (kindly provided by Dr Altieri, The Wistar Institute Cancer Center,

Philadelphia USA) was diluted in DMSO and used at concentration of 10 µM for up to 8 h. Paraquat

(Sigma-Aldrich, #36541) was dissolved in H_2O and used at a concentration of 0.5 mM for 48 h. For

SG assembly, sodium arsenite (Sigma Aldrich) was diluted in H_2O and used at a concentration of 0.5

mM for 30 min. Silvestrol (MedChemExpress) was resuspended in DMSO and used at a

concentration of 2 µM for 1 h. Phosphorylation of eIF2α was induced by DTT (Promega, DMSO) at 5

mM for 2 h or by UV irradiation at 20 mJ/cm2 using Stratalinker 1800 (Stratagene) followed by 2 h

incubation at 37 ºC. GCN2 activity was inhibited using 1 µM A-92 (Axon Medchem, dissolved in

DMSO). PERK activity was inhibited using 0.1 µM GSK2606414 (Merck, dissolved in DMSO). SG

formation was inhibited by pre-incubating cells with G3Ib, or the control drug G3Ib' (kindly provided by

Paul Taylor) at 50 µM final concentration for 20 min before further stimulating with other stressors.

<u>Transfection with double-stranded (ds) RNA</u>

Synthesis and purification of 200-bp dsRNA was described previously (Klein et al., 2022). In brief, a

200-bp region from the ampicillin resistance gene was used as template for *in vitro* synthesis of plus

and minus strand transcripts, hybridized and purified. 2×10^5 cells were seeded in a 6-well plate

containing coverslips and transfected with 1 µg dsRNA mixed with Lipofectamine 2000 transfection

reagent (Thermo Fisher Scientific) in a 1:3 ratio. Cells were fixed after 16 h with PBS-PFA 4% and

further processed for immunofluorescence analysis.

<u>Immunofluorescence microscopy</u>

Cells were seeded on sterilized coverslips in a 24-well plate and treated as required. Media was

removed and fixed with 4% paraformaldehyde (PFA) in PBS for 20 min at RT. Fixation solution was

removed, coverslips were rinsed with PB and stored at 4 ºC or processed immediately. Cells were

permeabilized with 0.1% Triton X-100 (Sigma) in PBS for 5 min at RT and blocked with 0.5% bovine

serum albumin (Fisher) in PBS for 1 h at RT. Cells were then incubated with primary antibody solution

for 1 h at RT, washed 3 times with PBS and incubated with secondary antibody solution containing

0.2 μg/ml DAPI for 1 h at RT. Coverslips were then rinsed 3 times with PBS and mounted onto
microscope slides with 5 μl Mowiol 4-88 (Sigma-Aldrich #81381). Cells were visualized and imaged
with a Nikon Ti-Eclipse A1M Confocal Microscope using a 60X objective. Confocal images were
analysed utilizing Image J. SG quantification was either performed manually counting at least 100
cells per condition or automatically using the Analyze Particles function, obtaining further information
such as the average size of SG and number per cell. Quantification of mitochondria morphology was
performed by manually counting cells with distinct mitochondria phenotypes – tubular or fragmented –
based on TOMM20 staining. The primary antibodies used were rabbit anti-G3BP1 (1:600, Sigma),
goat anti-eIF3η (1:600, Santa Cruz), rabbit anti-TOMM20 (1:1000, abcam ab78547), mouse IgG2a
anti-dsRNA (1:250, SCICOBS), mouse IgG1 anti-TOMM20 (1:100, Becton Dickinson), mouse anti-
puromycin (1:5, DSHB #PMY-2A4). Secondary antibodies used were goat anti-rabbit Alexa 488,
donkey anti-goat Alexa 555, chicken anti-mouse 633, goat anti-mouse-IgG2 Alexa 568 and goat anti-
mouse-IgG1 Alexa 488 (1:500, Invitrogen).

<u>Ribopuromycylation assay</u>

Cells were seeded on sterilized coverslips in a 24-well plate and treated as required. Prior fixation
cells were incubated with puromycin (Sigma) at 10 μg/ml at 37° for 5 min to label the nascent
polypeptide chains followed by a 2 min pulse at RT with emetine (Sigma) at 180 μM to block
translation elongation. Media was removed, cells were rinsed with warm DMEM, fixed with 4% PFA in
PBS for 20 min at RT and processed for immunofluorescence as described above. Puromycin
incorporation was quantified using ImageJ. The raw integrated density values were measured from ~
100 cells per condition.

<u>Immunoblotting</u>
Cells were seeded in 6-well plates to be around 80-90% confluent at the indicated times. Cells were
treated as required, then media was removed, and cells were washed twice with cold PBS. Cells were
lysed with protein lysis buffer (50 mM Tris-HCl, 250 mM NaCl, 5 mM EDTA, 1 mM Mg_2Cl, 1% NP40
and supplemented with protease inhibitor cocktail, Roche), incubated on ice for 20 minutes and spun
at 10,000 g for 10 min at 4 ºC. Supernatant was collected and protein concentration was measured
following BCA pierce kit (Fisher) and between 20-60 μg of protein were used depending on the
experiment and quantity obtained. Lysed were mixed with 3x red Loading buffer supplemented with
DTT (Cell Signalling) and boiled at 95 ºC for 5 min. Lysates were separated in 4-20% gradient gel with
1x SDS running buffer and proteins were transferred to a nitrocellulose membrane using the Pierce
Power Station and 1-step Transfer Buffer (Thermo Scientific). Membranes were blocked in 5% BSA or
5% marvel in TBS-Tween (TBS-T) for 1 h at RT and then incubated with primary antibodies overnight
at 4 ºC. Next day, membranes were washed 3 times with TBS-T and subsequently incubated with
secondary antibodies (Dako, 1:5000) diluted in 5% marvel in TBS-T for 1 h at RT. After three washes
with TBS-T, secondary antibodies were labelled with Clarity Western ECL Substrate (Bio-Rad), and
membranes were imaged using the vilber-Lourmat Imaging system. Band intensity was quantified and
analysed using Image Lab Software (Bio-rad). Primary antibodies used were rabbit anti-eIF2α
(1:1000, Cell Signaling), rabbit anti-P-eIF2α (1:1000, Invitrogen), rabbit anti-ATF4 (1:1000, Cell

Signaling), rabbit anti-GADD34 (1:1000, Proteintech), mouse anti-GAPDH (1:5000, Ambion), goat

anti-eIF3η (1:2000, Santa Cruz) and anti-eIF3A (1:2000, Bethyl Laboratories).

<u>Preparation of RNA samples and RT-qPCR</u>

Cells were seeded in 6-well plates to be around 80-90% confluent at the indicated times. Cells were

treated as needed, then were washed twice with cold PBS, and total RNA was extracted using a

quick-RNA micro prep kit (Zymo Research) following the manufacturer's instructions. Purified RNA

was quantified on Nanodrop, and 500 ng of RNA was subject to reverse transcription using Precision

nanoScript2 Reverse Transcription kit from Primer Design. Next, real-time quantitative PCR (qPCR)

was performed in duplicate with 25 ng of template, specific primers listed below used at a final

concentration of 0.3 µM and Precision Plus Master Mix (Primer Design) following manufacturer's

instructions, using the Quant Studio 7 Flex (Applied Biosystems). The comparative Ct method was

used; Ct values from genes of interest were normalized to RpL9 Ct values unless otherwise specified.

The 2-ΔΔCT method was used to calculate the fold change expression relative to non-treated (Livak

and Schmittgen, 2001).

<u>Mitochondrial respiration</u>

Oxygen consumption rate of U2OS WT and G3BP1/2 dKO cells was measured on an extracellular

flux analyzer (Agilent Seahorse) using the XF Cell Mito Stress Test Kit following manufacturer's

protocol. Briefly, cells were seeded one day prior to the experiment on Seahorse XFe24 culture plate

at 80% confluency. Cells were treated as required and 1 h prior running the plate, medium was

removed, and cells were incubated with assay medium supplemented with 1 mM pyruvate, 2 mM

glutamine and 10 mM glucose. Cells were then sequentially stimulated with oligomycin at 1 µM,

FCCP at 2 µM and rotenone and antimycin A at 1 µM and respiratory profile was obtained. Data was

normalised by amount of protein for each well. Following assay, cells were lysed with protein lysis

buffer (50 mM Tris-HCl, 250 mM NaCl, 5 mM EDTA, 1 mM Mg_2Cl, 1% NP40 and supplemented with

protease inhibitor cocktail, Roche), incubated on ice for 20 minutes and spun at 10,000 g for 10 min at

4 ºC. Supernatant was collected and protein concentration was measured following BCA pierce kit

(Fisher).

<u>Mitochondrial ROS measurement</u>

Cells were seeded in 96-well plate to be around 60% confluent for the day of the experiment. After

GTPP treatment for 2 h, cells were washed and incubated with 2.5 µM MitoSOX Red (Invitrogen,

#M36008) diluted in warm PBS for 30 min at 37 ºC. Next, cells were washed three times and

fluorescence was analysed in IncuCyte S3 imaging system. MitoSOX Red mean intensity was

extracted from 4 images taken at 20X per condition and plotted relative to DMSO-treated cells.

<u>Mitochondrial translation assay</u>

To label newly synthesized mitochondrial-encoded proteins, a click chemistry-based approach was

applied (Yousefi et al., 2021). Briefly, WT and G3BP1/2 dKO U2OS cells were treated with 10µM

GTPP or DMSO for 2 h, before replacing existing media with methionine-free media (D0422, Sigma)

supplemented with 2nM L-glutamine. Cytoplasmic translation was inhibited with cycloheximide

(50µg/ml, 20 mins), before addition of methionine analogue L-Homopropargylglycine (HPG) which is

preferentially incorporated into nascent mitochondrial peptides, for 15 mins. To validate this approach,

cells were left in methionine-containing media, whereby methionine was preferentially incorporated

into mitochondrial ribosomes instead of HPG, resulting in loss of HPG fluorescence signal. For control

samples, mitochondrial translation was blocked using Chloramphenicol (CHL, 150 µg/ml, 50 mins)

before HPG labelling. Cells were then incubated in buffer A (10 mM HEPES, 10 mM NaCl, 5 mM KCl,

300 mM sucrose, and 0.015% digitonin) for 2 min on ice, followed by 15 s in buffer A without digitonin

before fixation with 4% PFA in PBS for 30 min at RT. Next, click chemistry and immunofluorescence

was performed. After fixation, coverslips were rinsed in PBS for 5 min then quenched with 100 mM

NH4Cl in PBS for 15 min and blocked with 5% BSA in PBS for 10 min. Click chemistry was performed

as per the manufacturer guidelines. Briefly, coverslips were then incubated with 500µl Picolyl azide

AF-555 (CLK-090-1, Jena Bioscience) in CuAAC BTTAA Cell Reaction Buffer (CLK-073, Jena

Bioscience) in the dark for 60 min at RT. Coverslips were washed twice with 3% BSA in PBS for 5 min

before IF staining. The probes used in this experiment were TOMM20 (AB78547, Abcam) and

chicken anti-mouse Alexa fluor 488 (A21200, life technologies).

<u>Cell viability assay</u>

Cells were seeded in 96-well plate, incubated with GTPP and Cytotox dye (Sartorius, #4632) at a final

concentration of 250 nM and monitored overtime using IncuCyte S3 live cell analysis system. When

plasma membrane integrity is disrupting in dying cells, IncuCyte Cytotox Dye enters cells and

massively increases its fluorescence upon DNA binding. About 9 pictures were taken at 10X per

condition in 96-wells every 2-3 h over a 28 h period.

<u>Statistical analyses</u>

Statistical analyses were performed using Graphpad Prism software and three biological replicates

were analysed in all experiments. Statistical tests performed are indicated in the figure legends (****,

$P < 0.0001$; ***, $P < 0.001$; **, $P < 0.01$; *, $P < 0.05$; ns, not significant).

<u>Primer list</u>

GENE	FORWARD PRIMER (5'→3')	REVERSE PRIMER (5'→3')
ATF4	GCTAAGGCGGGCTCCTCCGA	ACCCAACAGGGCATCCAAGTCG
ATF5	TTTGCAGTGCGGGAAGAT	AAAATGAACACCCAGTCACCA
CHOP	GGAGCATCAGTCCCCCACTT	TGTGGGATTGAGGGTCACATC
DNAJA3	GAGGACGAGACAGATGTGGAG	TCCTGCGGAGCTATCCAT
DYRK3	TCCTTCTGAACCACCTCCAC	CCTTCATCTCACCTCCATCC
GADD34	CAGAAACCCCTACTCATGATCC	AAATGGACAGTGACCTTCTCG
HSPD1	GATGGAGAAGCTCTAAGTACACTCG	GCTGGTTCTTTCTATTGTCACCA
HSP90AA1	ATGGCAGCAAAGAAACAC	GTATCATCAGCAGTAGGGTCA
HSP90BB1	TGGCAGTCAAGCACTTTTCTGT	GCCCGACGAGGAATAAATAGC
LONP1	AACGTGTCGGGCTACGTG	CTCTCATCCAAGCCACACAG
RPL9	GGTGGGGTAACAGAAAGGAAC	CGTTGATGGGGAAGTGAGC
RPS18	CCAAGAGGGCGGGAGAAC	TATTTTCCATCCTTTACATCCTTCTG

CCL5	ACGTCAAGGAGTATTTCTACAC	GATGTATTCTTGAACCCACT
CXCL10	TGGCATTCAAGGAGTACCTC	TTGTAGCAATGATCTCAACACG
MXA	ATCCTGGGATTTTGGGGC TT	CCGCTTGTCGCTGGTGTCG
OAS1	AGGAAAGGTGCTTCCGAGGTAG	GGACTGAGGAAGACAACCAGGT

Figure legends

Figure 1: **ATF5-dependent UPR^{mt} induction by GTPP is associated with early SG formation and**
translational shut off. (**A**) U2OS cells were transfected with non-targeting siRNA (siCont) or ATF5-
specifc siRNA (siATF5). After 48 h cells were treated either with DMSO or 10 µM GTPP for 4h and
mRNA levels of UPR^{mt} markers were measured by RT-qPCR. Results shown as mean ± SD, n=3,
normalised to RPL9 mRNA and shown relative to siCont DMSO expression level, analysed by two-
way ANOVA (****, $P < 0.0001$; ***, $P < 0.001$; **, $P < 0.01$; *, $P < 0.05$). (**B**) Cells were treated with
GTPP up to 8 h and analysed by immunofluorescence for the SG markers G3BP1 (cyan) and eIF3η
(magenta). Nuclei were stained with DAPI. Scale bars indicate 10 µm. (**C**) Quantification by manual
counting of cells with SGs using ImageJ, at least 100 cells were counted per replicate. Results are
shown as mean ± SD, n=3 and were analysed by one-way ANOVA compared to DMSO treated cells
(****, $P < 0.0001$; *, $P < 0.05$). (**D**) Representative violin plots from three independent experiments
showing *de novo* protein synthesis upon GTPP treatment over time. Arsenite (Ars)-treated cells were
included as positive control, shown in yellow. Cells positive for SGs are shown in light purple. Figure
shows the median and quartiles of puromycin signal intensity measured by immunofluorescence,
number of cells analysed from this replicate is shown above per each condition and were analysed by
one-way ANOVA compared to DMSO treated cells (****, $P < 0.0001$; **, $P < 0.01$; *, $P < 0.05$).
Representative images are shown in Sup Fig 2.

Figure 2: UPR^{mt} activation by GTPP results in activation of the integrated stress response
(ISR) through PERK. (**A**) Representative western blot of three independent experiments showing
protein levels of GADD34, ATF4, and p-eIF2α in U2OS cells treated with GTPP for up to 6. Band
intensities were normalised to that of GAPDH and to DMSO for each time point. (**B**) RT-qPCR
analysis of ATF4, ATF5, CHOP and GADD34 mRNA levels in U2OS cells treated with GTPP for up to
6 h. Results shown as mean ± SD, n=3, normalised to RPL9 mRNA and shown relative to DMSO
expression level, analysed by one-way ANOVA (****, $P < 0.0001$; *, $P < 0.05$). (**C**) RT-qPCR analysis
of the interferon-stimulated genes CXCL10, CCL5, MX1 and OAS1 after 1, 4 and 8 h GTPP exposure.
Results shown as mean ± SD, n=3, normalised to RPL9 mRNA and shown relative to DMSO
expression level. (**D**) Representative immunoblot from three independent experiments showing levels
of p-eIF2α in U2OS cells treated with either DMSO, 0.1 µM PERK inhibitor GSK2606414 or 1 µM
GCN2 inhibitor A-92 for 1 h, followed by treatment with 10 µM GTPP for 2 h, or 5 mM DTT for 2 h, or
2 h-post UV light exposure at 20mJ/cm³. Band quantifications were normalised to GAPDH and
relative to DMSO for each inhibitor used. (**E**) Cells were pre-incubated with either DMSO or 0.1 µM
PERK inhibitor GSK2606414, further stimulated with GTPP for 2 h and analysed by

immunofluorescence for the SG markers G3BP1 (cyan) and eIF3η (magenta). Nuclei were stained
with DAPI. Scale bars indicate 10 µm. (**F**) Quantification by manual counting of cells with SGs using
ImageJ, at least 100 cells were counted per replicate. Results are shown as mean ± SD, n=3 and
were analysed by two-way ANOVA compared to DMSO treated cells (**, $P < 0.001$).

Figure 3: Pre-treatment with UPR^{mt} inducers impairs assembly of eIF2α-dependent SGs and
prevents eIF2α phosphorylation. Representative immunofluorescence images (**A, E**) and quantified
data (**B, F**) of U2OS cells pre-treated with 10 µM GTPP (**A, E**) or 0.5 Paraquat (**B, F**). Sodium
arsenite (0.5 mM) was added 30 min before harvesting of cells to induce SG formation (schematic
diagram of the experimental setup is shown on the top right). Cells were analysed by
immunofluorescence for the SG markers G3BP1 (cyan) and eIF3η (magenta). Nuclei were stained
with DAPI. Scale bars indicate 10 µm. Quantification was done by manual counting of cells with SGs
using ImageJ and at least 100 cells were counted per replicate. Results are shown as mean ± SD,
n=3 and were analysed by two-way ANOVA (****, $P < 0.0001$; ***, $P < 0.001$). The average size of
SGs per cell and the number of SGs per cell in U2OS cells treated with GTPP (C) or Paraquat (G)
were quantified using the G3BP1 channel and Analyze Particle plugin from ImageJ,, as displayed as
violin plot displaying showing the median and quartiles of at least 60 cells per condition from three
independent experiments analysed by (C) Krustal-Wallis test (*, $P < 0.05$; **, $P < 0.01$; (****, $P < $
0.0001) and (G) by Mann-Whitney test (****, $P < 0.0001$). (**D, H**) Representative blot from three
independent experiments showing p-eIF2α protein levels in U2OS cells treated as in (A)and (E)..
eIF2α phosphorylation levels were normalized to GAPDH expression levels and are shown as relative
to DMSO/NT cells; NT, non-treated.

Figure 4: GADD34 upregulation is responsible for preventing eIF2α-phosphorylation and SG
assembly. U2OS control (Ctrl) or GADD34 KO cells were treated with GTPP 4 h (**A**) or paraquat (Pq)
48 h (**C**) and 0.5 mM sodium arsenite (Ars) was added 30 min before harvest and cells were analysed
by immunofluorescence for the SG markers G3BP1 (cyan) and eIF3η (magenta). Nuclei were stained
with DAPI. Scale bars indicate 10 µm. (**B, D**) Panels show quantification by manual counting of cells
with SGs using ImageJ, at least 100 cells were counted per replicate. Results are shown as mean ±
SD, n=3 and were analysed by two-way ANOVA (****, $P < 0.0001$; **, $P < 0.01$). (**E, F**) Representative
western blot analysis of three independent experiments of same conditions as above showing
GADD34 protein levels and arsenite-induced eIF2α phosphorylation levels in Ctrl and GADD34 U2OS
cells pre-treated with GTPP (**E**) and Pq (**F**). eIF2α phosphorylation levels were normalized to eIF3A or
GAPDH expression levels and are shown as relative to DMSO/NT cells; NT, non-treated.

Figure 5: UPR^{mt} activation also alters eIF2α-independent SG formation associated with an
upregulation of SG disassembly chaperones. U2OS were pre-treated with GTPP (**A**) or paraquat
(Pq) (**D**) and 2 µM silvestrol (Silv) was added 1 h before harvest, analysed by immunofluorescence for
the SG markers G3BP1 (cyan) and eIF3η (magenta). Nuclei were stained with DAPI. Scale bars
indicate 10 µm. (**B, E**) Panels show quantification by manual counting of cells with SG using ImageJ,
at least 100 cells were counted per replicate. Results are shown as mean ± SD, n=3, and were
analysed by two-way ANOVA (****, $P < 0.0001$; ns, non-significant). (**C, F**) Violin plot displaying the

average size of SGs per cell and the number of SGs per cell of data from A and D, respectively, using

G3BP1 channel and Analyze Particle plugin from ImageJ. Figures show the median and quartiles of

at least 100 cells per condition from three independent experiments and were analysed by Mann-

Whitney test (****, $P < 0.0001$; **, $P < 0.01$). (**G, H**) RT-qPCR analysis of SG disassembly chaperones

HSP90AA1, HSP90AB1 and DYRK3 mRNA expression levels in response to GTPP (**G**) and Pq (**H**)

treatment. Results shown as mean ± SD, n=3, normalised to RpL9 mRNA and shown relative to

DMSO/NT expression level, analysed by two-way ANOVA (****, $P < 0.0001$; **, $P < 0.01$; *, $P < 0.05$).

Figure 6: Presence of SGs alters mitochondrial functions during UPR^mt. (**A**) WT and G3BP1/2

dKO cells were treated with either DMSO or GTPP for 2 h and analysed by immunofluorence for the

mitochondrial marker TOMM20 (cyan). Nuclei were stained with DAPI. Scale bars indicate 10 μm. T:

tubular mitochondria; F: fragmented mitochondria. (**B**) Quantification of the mitochondrial phenotypes

observed in **A**, at least 100 cells were counted per biological replicate. Figure shows mean ± SD, n=3

and analysed by two-way ANOVA (**, $P < 0.01$). (**C**) Following GTPP treatment for 2 h, U2OS WT

and G3BP1/2 cells were incubated with 2.5 uM MitoSOX dye for 30 mins, and mitochondrial ROS

were quantitatively measured using IncuCyte S3 live cell analysis software. Figure shows mean ± SD,

n=3, relative to DMSO-treated cells, and analysed by two-way ANOVA (****, $P < 0.0001$). (**D**)

Schematic diagram of the respiratory profile of live cells using Seahorse Cell Mito Stress Test,

showing the key parameters of mitochondrial function and the drugs being used. Briefly, oxygen

consumption rate is measured when cells are sequentially stimulated with oligomycin (ATP synthase

inhibitor), FCCP (protonophore, uncoupling) and antimycin A/rotenone (inhibitors of complex III and I).

(**E-F**) Representative traces (**E**) and averaged OCR data (**F**) from three independent experiments in

WT (solid line) or G3BP1/2 dKO (dashed line) U2OS cells treated with DMSO (black) or GTPP

(purple) for 2 h. Results shown as mean ± SD, n=3 and as fold change to DMSO-treated cells for

each cell line, analysed by two-way ANOVA (****, $P < 0.0001$; ***, $P < 0.001$; **, $P < 0.01$; *, $P < 0.05$).

Figure 7: G3BP1/2 are required for effective cytosolic translational shut off but not

mitochondrial during UPR^mt activation. (**A**) G3BP1/2 dKO cells were treated with GTPP or arsenite

for the indicated times and stimulated with puromycin followed by emetine before fixation. Then, cells

were analysed by immunofluorescence with anti-puromycin (grey). Nuclei were stained with DAPI.

Scale bars indicate 10 μm. (**B**) Representative violin plots from three independent experiments

showing *de novo* protein synthesis upon GTPP treatment over time. Arsenite (Ars)-treated cells were

included as positive control, shown in yellow. Figure shows the median and quartiles of puromycin

signal intensity measured by immunofluorescence, number of cells analysed from this replicate is

shown above per each condition and were analysed by one-way ANOVA compared to DMSO treated

cells (****, $P < 0.0001$). (**C**) U2OS WT cells were either incubated in normal media (Met, methionine)

as a control, or methionine-free media in the presence and/or absence of cycloheximide (CHX) and/or

chloramphenicol (CHL), before adding HPG for 15 minutes. After fixation, AF555-azide (cyan) was

used for click-chemistry reaction to visualise HPG incorporation and cells were additionally labelled

with TOMM20 as mitochondrial marker (magenta). Nuclei stained with DAPI. Scale bars indicate 10

μm. (**D**) WT and G3BP1/2 dKO cells were treated with either DMSO or GTPP for 2h, before inhibiting

cytosolic translation with cycloheximide and performing HPG labelling. After fixation, cells were fixed,
labelled and imaged as in (**C**). Scale bars indicate 10 µm. (**E**) Representative violin plots from three
independent experiments showing HPG incorporation upon 2h GTPP treatment. For quantification,
images were subjected to thresholding before AF555 signal intensity was measured using measure
function in ImageJ. Results were analysed by unpaired t-test for each cell line compared to DMSO
treated cells (***, $P < 0.001$; ****, $P < 0.0001$).

Figure 8: Loss of SG scaffolding proteins G3BP1/2 delays cell death following prolonged
UPR^{mt} activation. GTPP and Cytotox Dye were added to WT and G3BP1/2 U2OS cells and
monitored for 28 h using IncuCyte S3 live cell analysis software. (**A**) Selected representative time
frames of DMSO and GTPP-treated cells. Scale bar indicates 200 µm. (**B**) Representative cell viability
kinetics of three independent experiments of WT and G3BP1/2 dKO cells when exposed to GTPP.
The figure shows mean ± SEM of cell death count (orange cytotox dye) from 9 images per condition.
Data were analysed by two-way ANOVA (****, $P < 0.0001$; **, $P < 0.01$; *, $P < 0.05$; when comparing
WT GTPP vs G3BP1/2 dKO GTPP. #, $P < 0,001$; when comparing WT DMSO vs WT GTPP. ^, $P <$
0.0001; when comparing G3BP1/2 dKO DMSO vs G3BP1/2 dKO GTPP).

Supplementary Figure 1: Inhibition of SGs formation by G3Ib blocks GTPP-induced SG
assembly. U2OS cells were preincubated with either DMSO or G3Ib/G3Ib' at 50 µM for 20 min and
GTPP was added to the cells at final concentration of 10 µM. SG formation was analysed by
immunofluorescence using SG marker G3BP1 (cyan). Nuclei were stained with DAPI. Scale bars
indicate 10 µm.

Supplementary Figure 2: Translational shut off upon UPR^{mt} activation by GTPP. U2OS WT cells
were stimulated with GTPP for the indicated times or arsenite for 1 h and treated with puromycin
followed by emetine before fixation. Then, cells were analysed by immunofluorescence with anti-
puromycin (grey) and the SG markers anti-G3BP1 (cyan) and anti-eIF3η (magenta). Nuclei were
stained with DAPI. Scale bars indicate 10 µm.

Supplementary Figure 3: GTPP treatment does not result in mitochondrial dsRNA leakage.
Cells were treated with GTPP for 2 h or transfected with dsRNA (positive control) and analysed by
immunofluorescence with anti-dsRNA (magenta) and the mitochondrial marker anti-TOMM20 (cyan).
Nuclei were stained with DAPI. Scale bars indicate 10 µm.

Supplementary Figure 4: **GTPP treatment impairs subsequent SG formation by arsenite.**
Individual panels are shown from Fig 3A.

Supplementary Figure 5: Paraquat treatment results in UPR^{mt} activation but does not induce
SG formation in U2OS cells. (A) RT-qPCR analysis of ISR and UPR^{mt} markers in U2OS cells
treated with 0.5 mM paraquat for 48 h. Results shown as mean ± SD, n=3, normalised to RPL9
mRNA and shown relative to NT expression level, analysed by two-way ANOVA (****, $P < 0.0001$; *, P
< 0.05); NT, non-treated. (**B**) Treatment of U2OS cells with paraquat for 1, 2, 6, 24 and 48 h does not
result in SG assembly. Representative images from three independent experiments are shown. Cells

were analysed by immunofluorescence for the SG markers G3BP1 (cyan) and eIF3η (magenta);
nuclei were stained with DAPI. Scale bars indicate 10 µm. (**C**) Individual panels are shown from Fig
3E.

Supplementary Figure 6: ATF5 silencing modulates SG dynamics in a stress-specific manner.
(**A**) U2OS cells were transfected with non-targeting siRNA (siCont) or ATF5-specifc siRNA (siATF5),
after 48 h cells were treated with GTPP 4 h and sodium arsenite (Ars) was added 30 min before
harvesting and cells were analysed by immunofluorescence for the SG markers G3BP1 (cyan) and
eIF3η (magenta). Nuclei were stained with DAPI. Scale bars indicate 10 µm. (**B**) Panel shows
quantification by manual counting of cells with SGs using ImageJ, at least 100 cells were counted per
replicate. Results are shown as mean ± SD, n=3 and were analysed by two-way ANOVA (ns, non-
significant). (**C, D**) Violin plot displaying the average size of SGs per cell and the number of SGs per
cell from GTPP treated cells (**C**) and GTPP+Ars treated cells (**D**) quantification was performed using
G3BP1 channel and Analyze Particle plugin from ImageJ. Figure shows the median and quartiles of
at least 60 cells per condition from three independent experiments and were analysed by Mann-
Whitney test (**, $P < 0.01$).

Supplementary Figure 7: Validation of U2OS GADD34 KO single cell clones by Western blot.
(**A**) Clones were treated with 2 µM thapsigargin (Thap) or with DMSO (-) for 3 h and GADD34 protein
levels were analysed by western blot. eIF3A was used as loading control. (**B, C**) Individual panels are
shown from Fig 4A and 4C, respectively.

Supplementary Figure 8: U2OS G3BP1/2 dKO cells do not form SGs. U2OS WT and G3BP1/2
dKO cells were treated with arsenite (Ars) for 30 min or GTPP for 2 h and analysed by
immunofluorescence using the SG markers G3BP1 (cyan) and eIF3η (magenta). Nuclei were stained
with DAPI. Scale bars indicate 10 µm.

Supplementary Figure 9: U2OS G3BP1/2 dKO cells presented reduced MitoSOX Red signal
after GTPP treatment compared to WT cells. Representative IncuCyte images from Figure 6C
showing MitoSOX Red and phase channels.

Data availability Statement

All the data are included in the article. All unique reagents generated in this study are
available from the corresponding authors with a completed Materials Transfer Agreement.

884
Acknowledgments

We thank P. J. McCormick (Queen Mary University London) for helpful discussions during
the study.

Author contributions

N. L., A. R. and I. S. conceptualization; N. L., M. L-N., A. R., B.D.F., J.P.T., Z. S., R. S. and I. S.
methodology; N. L., M. L-N., A. R., and Z. S. formal analysis; N. L., M. L-N., E.R., A. R., Z. S., R. S.
and I. S. investigation; N. L. and M. L-N. writing–original draft; N. L., M. L-N., A. R., E.R., Z. S., R. S.,
B.D.F., J.P.T. and I. S. writing–review & editing; N.L. supervision. N. L., A. R., and I. S. funding
acquisition.

Funding and additional information

Work in N.L.'s laboratory is supported by the Biotechnology and Biological Sciences Research
Council research and strategic grants BB/V014528/1 and BBS/E/PI/230002A. Bioimaging studies are
supported by BBSRC Core Capability Grant BBS/E/I/00007039. M. L-N. and R. S. are supported by
DCSA4 Ph.D studentships from the University of Surrey. Work in A.R.'s laboratory was supported by
the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) project numbers
240245660 SFB1129 TP13 and 278001972 TRR186 A14.

Conflict of interest

The authors declare that they have no conflicts of interest with the contents of this article.

References

Almeida, L.M., B.R. Pinho, M.R. Duchen, and J.M.A. Oliveira. 2022. The PERKs of
mitochondria protection during stress: insights for PERK modulation in neurodegenerative and
metabolic diseases. *Biol Rev Camb Philos Soc*. 97:1737-1748.
Amen, T., and D. Kaganovich. 2021. Stress granules inhibit fatty acid oxidation by modulating
mitochondrial permeability. *Cell Rep*. 35:109237.
Anderson, N.S., and C.M. Haynes. 2020. Folding the Mitochondrial UPR into the Integrated
Stress Response. *Trends Cell Biol*. 30:428-439.
Anderson, P., N. Kedersha, and P. Ivanov. 2015. Stress granules, P-bodies and cancer.
Biochim Biophys Acta. 1849:861-870.
Aoyama-Ishiwatari, S., T. Okazaki, S.I. Iemura, T. Natsume, Y. Okada, and Y. Gotoh. 2021.
NUDT21 Links Mitochondrial IPS-1 to RLR-Containing Stress Granules and Activates Host Antiviral
Defense. *J Immunol*. 206:154-163.
Arimoto, K., H. Fukuda, S. Imajoh-Ohmi, H. Saito, and M. Takekawa. 2008. Formation of stress
granules inhibits apoptosis by suppressing stress-responsive MAPK pathways. *Nat Cell Biol*.
10:1324-1332.
Aulas, A., M.M. Fay, S.M. Lyons, C.A. Achorn, N. Kedersha, P. Anderson, and P. Ivanov. 2017.
Stress-specific differences in assembly and composition of stress granules and related foci. *J Cell
Sci*. 130:927-937.
Baker, B.M., A.M. Nargund, T. Sun, and C.M. Haynes. 2012. Protective coupling of
mitochondrial function and protein synthesis via the eIF2α kinase GCN-2. *PLoS Genet*. 8:e1002760.

Banoth, B., and S.L. Cassel. 2018. Mitochondria in innate immune signaling. *Transl Res.*
202:52-68.

Baron, D.M., L.J. Kaushansky, C.L. Ward, R.R. Sama, R.J. Chian, K.J. Boggio, A.J. Quaresma,
J.A. Nickerson, and D.A. Bosco. 2013. Amyotrophic lateral sclerosis-linked FUS/TLS alters stress
granule assembly and dynamics. *Mol Neurodegener.* 8:30.

Begovich, K., A.Q. Vu, G. Yeo, and J.E. Wilhelm. 2020. Conserved metabolite regulation of
stress granule assembly via AdoMet. *J Cell Biol.* 219:e201904141.

Bordeleau, M.E., F. Robert, B. Gerard, L. Lindqvist, S.M. Chen, H.G. Wendel, B. Brem, H.
Greger, S.W. Lowe, J.A. Porco, Jr., and J. Pelletier. 2008. Therapeutic suppression of translation
initiation modulates chemosensitivity in a mouse lymphoma model. *J Clin Invest.* 118:2651-2660.

Brillo, V., L. Chieregato, L. Leanza, S. Muccioli, and R. Costa. 2021. Mitochondrial Dynamics,
ROS, and Cell Signaling: A Blended Overview. *Life (Basel).* 11:332.

Brostrom, C.O., C.R. Prostko, R.J. Kaufman, and M.A. Brostrom. 1996. Inhibition of
translational initiation by activators of the glucose-regulated stress protein and heat shock protein
stress response systems. Role of the interferon-inducible double-stranded RNA-activated eukaryotic
initiation factor 2alpha kinase. *J Biol Chem.* 271:24995-25002.

Burke, J.M., E.T. Lester, D. Tauber, and R. Parker. 2020. RNase L promotes the formation of
unique ribonucleoprotein granules distinct from stress granules. *J Biol Chem.* 295:1426-1438.

Cereghetti, G., C. Wilson-Zbinden, V.M. Kissling, M. Diether, A. Arm, H. Yoo, I. Piazza, S.
Saad, P. Picotti, D.A. Drummond, U. Sauer, R. Dechant, and M. Peter. 2021. Reversible amyloids of
pyruvate kinase couple cell metabolism and stress granule disassembly. *Nat Cell Biol.* 23:1085-
1094.

Chamseddine, D., S.A. Mahmud, A.K. Westfall, T.A. Castoe, R.E. Berg, and M.W. Pellegrino.
2022. The mitochondrial UPR regulator ATF5 promotes intestinal barrier function via control of the
satiety response. *Cell Rep.* 41:111789.

Chen, W., H. Zhao, and Y. Li. 2023. Mitochondrial dynamics in health and disease:
mechanisms and potential targets. *Signal Transduction and Targeted Therapy.* 8:333.

Cocheme, H.M., and M.P. Murphy. 2008. Complex I is the major site of mitochondrial
superoxide production by paraquat. *J Biol Chem.* 283:1786-1798.

Corbet, G.A., and R. Parker. 2019. RNP Granule Formation: Lessons from P-Bodies and Stress
Granules. *Cold Spring Harb Symp Quant Biol.* 84:203-215.

D'Souza, A.R., and M. Minczuk. 2018. Mitochondrial transcription and translation: overview.
Essays Biochem. 62:309-320.

David, A., B.P. Dolan, H.D. Hickman, J.J. Knowlton, G. Clavarino, P. Pierre, J.R. Bennink, and
J.W. Yewdell. 2012. Nuclear translation visualized by ribosome-bound nascent chain
puromycylation. *J Cell Biol.* 197:45-57.

Deng, J., H.P. Harding, B. Raught, A.C. Gingras, J.J. Berlanga, D. Scheuner, R.J. Kaufman, D.
Ron, and N. Sonenberg. 2002. Activation of GCN2 in UV-irradiated cells inhibits translation. *Curr
Biol.* 12:1279-1286.

Dever, T.E., L. Feng, R.C. Wek, A.M. Cigan, T.F. Donahue, and A.G. Hinnebusch. 1992.
Phosphorylation of initiation factor 2 alpha by protein kinase GCN2 mediates gene-specific
translational control of GCN4 in yeast. *Cell*. 68:585-596.

Donehower, L.A., T. Soussi, A. Korkut, Y. Liu, A. Schultz, M. Cardenas, X. Li, O. Babur, T.K.
Hsu, O. Lichtarge, J.N. Weinstein, R. Akbani, and D.A. Wheeler. 2019. Integrated Analysis of TP53
Gene and Pathway Alterations in The Cancer Genome Atlas. *Cell Rep*. 28:1370-1384.e1375.

Eiermann, N., K. Haneke, Z. Sun, G. Stoecklin, and A. Ruggieri. 2020. Dance with the Devil:
Stress Granules and Signaling in Antiviral Responses. *Viruses*. 12:984.

Fiorese, C.J., A.M. Schulz, Y.-F. Lin, N. Rosin, M.W. Pellegrino, and C.M. Haynes. 2016. The
Transcription Factor ATF5 Mediates a Mammalian Mitochondrial UPR. *Curr Biol*. 26:2037-2043.

Freibaum, B.D., J. Messing, H. Nakamura, U. Yurtsever, J. Wu, H.J. Kim, J. Hixon, R.M.
Lemieux, J. Duffner, W. Huynh, K. Wong, M. White, C. Lee, R.E. Meyers, R. Parker, and J.P. Taylor.
2024. Identification of small molecule inhibitors of G3BP-driven stress granule formation. *J Cell Biol*.
223:e202308083.

Fu, Y., O. Sacco, E. DeBitetto, E. Kanshin, B. Ueberheide, and A. Sfeir. 2023. Mitochondrial
DNA breaks activate an integrated stress response to reestablish homeostasis. *Mol Cell*. 83:3740-
3753.

Fujikawa, D., T. Nakamura, D. Yoshioka, Z. Li, H. Moriizumi, M. Taguchi, N. Tokai-Nishizumi, H.
Kozuka-Hata, M. Oyama, and M. Takekawa. 2023. Stress granule formation inhibits stress-induced
apoptosis by selectively sequestering executioner caspases. *Curr Biol*. 33:1967-1981.e1968.

Giorgi, C., S. Marchi, and P. Pinton. 2018. The machineries, regulation and cellular functions of
mitochondrial calcium. *Nat Rev Mol Cell Biol*. 19:713-730.

Guo, X., G. Aviles, Y. Liu, R. Tian, B.A. Unger, Y.T. Lin, A.P. Wiita, K. Xu, M.A. Correia, and M.
Kampmann. 2020. Mitochondrial stress is relayed to the cytosol by an OMA1-DELE1-HRI pathway.
Nature. 579:427-432.

Haghighat, A., S. Mader, A. Pause, and N. Sonenberg. 1995. Repression of cap-dependent
translation by 4E-binding protein 1: competition with p220 for binding to eukaryotic initiation factor-
4E. *Embo j*. 14:5701-5709.

Harding, H.P., I. Novoa, Y. Zhang, H. Zeng, R. Wek, M. Schapira, and D. Ron. 2000. Regulated
translation initiation controls stress-induced gene expression in mammalian cells. *Mol Cell*. 6:1099-
1108.

Harding, H.P., Y. Zhang, H. Zeng, I. Novoa, P.D. Lu, M. Calfon, N. Sadri, C. Yun, B. Popko, R.
Paules, D.F. Stojdl, J.C. Bell, T. Hettmann, J.M. Leiden, and D. Ron. 2003. An integrated stress
response regulates amino acid metabolism and resistance to oxidative stress. *Mol Cell*. 11:619-633.

Hofmann, S., N. Kedersha, P. Anderson, and P. Ivanov. 2020. Molecular mechanisms of stress
granule assembly and disassembly. *Biochim Biophys Acta Mol Cell Res*:118876.

Houtkooper, R.H., L. Mouchiroud, D. Ryu, N. Moullan, E. Katsyuba, G. Knott, R.W. Williams,
and J. Auwerx. 2013. Mitonuclear protein imbalance as a conserved longevity mechanism. *Nature*.
497:451-457.

Jain, S., J.R. Wheeler, R.W. Walters, A. Agrawal, A. Barsic, and R. Parker. 2016. ATPase-

Modulated Stress Granules Contain a Diverse Proteome and Substructure. *Cell*. 164:487-498.

Jiang, D., H. Cui, N. Xie, S. Banerjee, R.M. Liu, H. Dai, V.J. Thannickal, and G. Liu. 2020. ATF4

Mediates Mitochondrial Unfolded Protein Response in Alveolar Epithelial Cells. *Am J Respir Cell Mol*

Biol. 63:478-489.

Kang, B.H., J. Plescia, H.Y. Song, M. Meli, G. Colombo, K. Beebe, B. Scroggins, L. Neckers,

and D.C. Altieri. 2009. Combinatorial drug design targeting multiple cancer signaling networks

controlled by mitochondrial Hsp90. *J Clin Invest*. 119:454-464.

Kaspar, S., C. Oertlin, K. Szczepanowska, A. Kukat, K. Senft, C. Lucas, S. Brodesser, M.

Hatzoglou, O. Larsson, I. Topisirovic, and A. Trifunovic. 2021. Adaptation to mitochondrial stress

requires CHOP-directed tuning of ISR. *Sci Adv*. 7:eabf0971.

Kedersha, N., S. Chen, N. Gilks, W. Li, I.J. Miller, J. Stahl, and P. Anderson. 2002. Evidence

that ternary complex (eIF2-GTP-tRNA(i)(Met))-deficient preinitiation complexes are core constituents

of mammalian stress granules. *Mol Biol Cell*. 13:195-210.

Kim, Y., J. Park, S. Kim, M. Kim, M.G. Kang, C. Kwak, M. Kang, B. Kim, H.W. Rhee, and V.N.

Kim. 2018. PKR Senses Nuclear and Mitochondrial Signals by Interacting with Endogenous Double-

Stranded RNAs. *Mol Cell*. 71:1051-1063.e1056.

Kitajima, H., R. Maruyama, T. Niinuma, E. Yamamoto, A. Takasawa, K. Takasawa, K. Ishiguro,

A. Tsuyada, R. Suzuki, G. Sudo, T. Kubo, K. Mitsuhashi, M. Idogawa, S. Tange, M. Toyota, A.

Yoshido, K. Kumegawa, M. Kai, K. Yanagihara, T. Tokino, M. Osanai, H. Nakase, and H. Suzuki.

2023. TM4SF1-AS1 inhibits apoptosis by promoting stress granule formation in cancer cells. *Cell*

Death Dis. 14:424.

Klein, P., S.M. Kallenberger, H. Roth, K. Roth, T.B.N. Ly-Hartig, V. Magg, J. Ales, S.R. Talemi,

Y. Qiang, S. Wolf, O. Oleksiuk, R. Kurilov, B. Di Ventura, R. Bartenschlager, R. Eils, K. Rohr, F.A.

Hamprecht, T. Hofer, O.T. Fackler, G. Stoecklin, and A. Ruggieri. 2022. Temporal control of the

integrated stress response by a stochastic molecular switch. *Sci Adv*. 8:eabk2022.

Lee, J.E., P.I. Cathey, H. Wu, R. Parker, and G.K. Voeltz. 2020. Endoplasmic reticulum contact

sites regulate the dynamics of membraneless organelles. *Science*. 367:eaay7108.

Lee, Y.Y., R.C. Cevallos, and E. Jan. 2009. An Upstream Open Reading Frame Regulates

Translation of GADD34 during Cellular Stresses That Induce eIF2α Phosphorylation*. *Journal of*

Biological Chemistry. 284:6661 - 6673.

Liao, Y.C., M.S. Fernandopulle, G. Wang, H. Choi, L. Hao, C.M. Drerup, R. Patel, S. Qamar, J.

Nixon-Abell, Y. Shen, W. Meadows, M. Vendruscolo, T.P.J. Knowles, M. Nelson, M.A. Czekalska, G.

Musteikyte, M.A. Gachechiladze, C.A. Stephens, H.A. Pasolli, L.R. Forrest, P. St George-Hyslop, J.

Lippincott-Schwartz, and M.E. Ward. 2019. RNA Granules Hitchhike on Lysosomes for Long-

Distance Transport, Using Annexin A11 as a Molecular Tether. *Cell*. 179:147-164.e120.

Livak, K.J., and T.D. Schmittgen. 2001. Analysis of relative gene expression data using real-

time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods*. 25:402-408.

Lu, B., and S. Guo. 2020. Mechanisms Linking Mitochondrial Dysfunction and Proteostasis

Failure. *Trends Cell Biol*. 30:317-328.

Magg, V., A. Manetto, K. Kopp, C.C. Wu, M. Naghizadeh, D. Lindner, L. Eke, J. Welsch, S.M.

Kallenberger, J. Schott, V. Haucke, N. Locker, G. Stoecklin, and A. Ruggieri. 2024. Turnover of

PPP1R15A mRNA encoding GADD34 controls responsiveness and adaptation to cellular stress. *Cell*

Rep. 43:114069.

Mahmud, S.A., M.A. Qureshi, M. Sapkota, and M.W. Pellegrino. 2020. A pathogen branched-

chain amino acid catabolic pathway subverts host survival by impairing energy metabolism and the

mitochondrial UPR. *PLoS Pathog*. 16:e1008918.

Martinus, R.D., G.P. Garth, T.L. Webster, P. Cartwright, D.J. Naylor, P.B. Høj, and N.J.

Hoogenraad. 1996. Selective induction of mitochondrial chaperones in response to loss of the

mitochondrial genome. *Eur J Biochem*. 240:98-103.

Matheny, T., B.S. Rao, and R. Parker. 2019. Transcriptome-Wide Comparison of Stress

Granules and P-Bodies Reveals that Translation Plays a Major Role in RNA Partitioning. *Mol Cell*

Biol. 39:e00313-00319.

Mediani, L., F. Antoniani, V. Galli, J. Vinet, A.D. Carrà, I. Bigi, V. Tripathy, T. Tiago, M. Cimino,

G. Leo, T. Amen, D. Kaganovich, C. Cereda, O. Pansarasa, J. Mandrioli, P. Tripathi, D. Troost, E.

Aronica, J. Buchner, A. Goswami, J. Sterneckert, S. Alberti, and S. Carra. 2021. Hsp90-mediated

regulation of DYRK3 couples stress granule disassembly and growth via mTORC1 signaling. *EMBO*

reports. 22:e51740.

Münch, C., and J.W. Harper. 2016. Mitochondrial unfolded protein response controls matrix pre-

RNA processing and translation. *Nature*. 534:710-713.

Munkácsy, E., and S.L. Rea. 2014. The paradox of mitochondrial dysfunction and extended

longevity. *Exp Gerontol*. 56:221-233.

Namkoong, S., A. Ho, Y.M. Woo, H. Kwak, and J.H. Lee. 2018. Systematic Characterization of

Stress-Induced RNA Granulation. *Mol Cell*. 70:175-187 e178.

Nargund, A.M., M.W. Pellegrino, C.J. Fiorese, B.M. Baker, and C.M. Haynes. 2012.

Mitochondrial import efficiency of ATFS-1 regulates mitochondrial UPR activation. *Science*. 337:587-

590.

Norat, P., S. Soldozy, J.D. Sokolowski, C.M. Gorick, J.S. Kumar, Y. Chae, K. Yağmurlu, F.

Prada, M. Walker, M.R. Levitt, R.J. Price, P. Tvrdik, and M.Y.S. Kalani. 2020. Mitochondrial

dysfunction in neurological disorders: Exploring mitochondrial transplantation. *npj Regenerative*

Medicine. 5:22.

Novoa, I., H. Zeng, H.P. Harding, and D. Ron. 2001. Feedback inhibition of the unfolded protein

response by GADD34-mediated dephosphorylation of eIF2alpha. *J Cell Biol*. 153:1011-1022.

Ohn, T., N. Kedersha, T. Hickman, S. Tisdale, and P. Anderson. 2008. A functional RNAi

screen links O-GlcNAc modification of ribosomal proteins to stress granule and processing body

assembly. *Nat Cell Biol*. 10:1224-1231.

Paget, M., C. Cadena, S. Ahmad, H.T. Wang, T.X. Jordan, E. Kim, B. Koo, S.M. Lyons, P.

Ivanov, B. tenOever, X. Mu, and S. Hur. 2023. Stress granules are shock absorbers that prevent

excessive innate immune responses to dsRNA. *Mol Cell*. 83:1180-1196.e1188.

Park, J.Y., and O.S. Shin. 2023. Stress Granules Inhibit Coxsackievirus B3-Mediated Cell
Death via Reduction of Mitochondrial Reactive Oxygen Species and Viral Extracellular Release. *J*
Microbiol Biotechnol. 33:582-590.

Pellegrino, M.W., A.M. Nargund, N.V. Kirienko, R. Gillis, C.J. Fiorese, and C.M. Haynes. 2014.
Mitochondrial UPR-regulated innate immunity provides resistance to pathogen infection. *Nature*.
516:414-417.

Pindel, A., and A. Sadler. 2011. The role of protein kinase R in the interferon response. *J*
Interferon Cytokine Res. 31:59-70.

Prentzell, M.T., U. Rehbein, M. Cadena Sandoval, A.S. De Meulemeester, R. Baumeister, L.
Brohée, B. Berdel, M. Bockwoldt, B. Carroll, S.R. Chowdhury, A. von Deimling, C. Demetriades, G.
Figlia, M.E.G. de Araujo, A.M. Heberle, I. Heiland, B. Holzwarth, L.A. Huber, J. Jaworski, M. Kedra,
K. Kern, A. Kopach, V.I. Korolchuk, I. van 't Land-Kuper, M. Macias, M. Nellist, W. Palm, S. Pusch,
J.M. Ramos Pittol, M. Reil, A. Reintjes, F. Reuter, J.R. Sampson, C. Scheldeman, A. Siekierska, E.
Stefan, A.A. Teleman, L.E. Thomas, O. Torres-Quesada, S. Trump, H.D. West, P. de Witte, S.
Woltering, T.E. Yordanov, J. Zmorzynska, C.A. Opitz, and K. Thedieck. 2021. G3BPs tether the TSC
complex to lysosomes and suppress mTORC1 signaling. *Cell*. 184:655-674.e627.

Quirós, P.M., A. Mottis, and J. Auwerx. 2016. Mitonuclear communication in homeostasis and
stress. *Nat Rev Mol Cell Biol*. 17:213-226.

Quirós, P.M., M.A. Prado, N. Zamboni, D. D'Amico, R.W. Williams, D. Finley, S.P. Gygi, and J.
Auwerx. 2017. Multi-omics analysis identifies ATF4 as a key regulator of the mitochondrial stress
response in mammals. *J Cell Biol*. 216:2027-2045.

Radetskyy, R., A. Daher, and A. Gatignol. 2018. ADAR1 and PKR, interferon stimulated genes
with clashing effects on HIV-1 replication. *Cytokine Growth Factor Rev*. 40:48-58.

Rehbein, U., M.T. Prentzell, M. Cadena Sandoval, A.M. Heberle, E.P. Henske, C.A. Opitz, and
K. Thedieck. 2021. The TSC Complex-mTORC1 Axis: From Lysosomes to Stress Granules and
Back. *Front Cell Dev Biol*. 9:751892.

Reineke, L.C., S.A. Cheema, J. Dubrulle, and J.R. Neilson. 2018. Chronic starvation induces
noncanonical pro-death stress granules. *J Cell Sci*. 131:jcs220244.

Reineke, L.C., and J.R. Neilson. 2019. Differences between acute and chronic stress granules,
and how these differences may impact function in human disease. *Biochem Pharmacol*. 162:123-
131.

Repici, M., M. Hassanjani, D.C. Maddison, P. Garcao, S. Cimini, B. Patel, E.M. Szego, K.R.
Straatman, K.S. Lilley, T. Borsello, T.F. Outeiro, L. Panman, and F. Giorgini. 2019. The Parkinson's
Disease-Linked Protein DJ-1 Associates with Cytoplasmic mRNP Granules During Stress and
Neurodegeneration. *Mol Neurobiol*. 56:61-77.

Rolland, S.G., S. Schneid, M. Schwarz, E. Rackles, C. Fischer, S. Haeussler, S.G. Regmi, A.
Yeroslaviz, B. Habermann, D. Mokranjac, E. Lambie, and B. Conradt. 2019. Compromised
Mitochondrial Protein Import Acts as a Signal for UPR(mt). *Cell Rep*. 28:1659-1669.e1655.

Rollins, M., S. Huard, A. Morettin, J. Takuski, T.T. Pham, M.D. Fullerton, J. Côté, and K. Baetz. 2017. Lysine acetyltransferase NuA4 and acetyl-CoA regulate glucose-deprived stress granule formation in Saccharomyces cerevisiae. *PLoS Genet*. 13:e1006626.

Ron, D. 2002. Translational control in the endoplasmic reticulum stress response. *J Clin Invest*. 110:1383-1388.

Saad, S., G. Cereghetti, Y. Feng, P. Picotti, M. Peter, and R. Dechant. 2017. Reversible protein aggregation is a protective mechanism to ensure cell cycle restart after stress. *Nat Cell Biol*. 19:1202-1213.

Samluk, L., M. Urbanska, K. Kisielewska, K. Mohanraj, M.J. Kim, K. Machnicka, E. Liszewska, J. Jaworski, and A. Chacinska. 2019. Cytosolic translational responses differ under conditions of severe short-term and long-term mitochondrial stress. *Mol Biol Cell*. 30:1864-1877.

Schwed-Gross, A., H. Hamiel, G.P. Faber, M. Angel, R. Ben-Yishay, J.I.C. Benichou, D. Ishay-Ronen, and Y. Shav-Tal. 2022. Glucocorticoids enhance chemotherapy-driven stress granule assembly and impair granule dynamics, leading to cell death. *J Cell Sci*. 135:jcs259629.

Smyrnias, I., S.P. Gray, D.O. Okonko, G. Sawyer, A. Zoccarato, N. Catibog, B. López, A. González, S. Ravassa, J. Díez, and A.M. Shah. 2019. Cardioprotective Effect of the Mitochondrial Unfolded Protein Response During Chronic Pressure Overload. *J Am Coll Cardiol*. 73:1795-1806.

Sorrentino, V., M. Romani, L. Mouchiroud, J.S. Beck, H. Zhang, D. D'Amico, N. Moullan, F. Potenza, A.W. Schmid, S. Rietsch, S.E. Counts, and J. Auwerx. 2017. Enhancing mitochondrial proteostasis reduces amyloid-β proteotoxicity. *Nature*. 552:187-193.

Spinelli, J.B., and M.C. Haigis. 2018. The multifaceted contributions of mitochondria to cellular metabolism. *Nat Cell Biol*. 20:745-754.

Suomalainen, A., and B.J. Battersby. 2018. Mitochondrial diseases: the contribution of organelle stress responses to pathology. *Nat Rev Mol Cell Biol*. 19:77-92.

Sutandy, F.X.R., I. Gößner, G. Tascher, and C. Münch. 2023. A cytosolic surveillance mechanism activates the mitochondrial UPR. *Nature*. 618:849-854.

Takahashi, M., M. Higuchi, H. Matsuki, M. Yoshita, T. Ohsawa, M. Oie, and M. Fujii. 2013. Stress granules inhibit apoptosis by reducing reactive oxygen species production. *Mol Cell Biol*. 33:815-829.

Tauber, D., G. Tauber, A. Khong, B. Van Treeck, J. Pelletier, and R. Parker. 2020. Modulation of RNA Condensation by the DEAD-Box Protein eIF4A. *Cell*. 180:411-426 e416.

Topf, U., I. Suppanz, L. Samluk, L. Wrobel, A. Böser, P. Sakowska, B. Knapp, M.K. Pietrzyk, A. Chacinska, and B. Warscheid. 2018. Quantitative proteomics identifies redox switches for global translation modulation by mitochondrially produced reactive oxygen species. *Nat Commun*. 9:324.

Van Houten, B., S.E. Hunter, and J.N. Meyer. 2016. Mitochondrial DNA damage induced autophagy, cell death, and disease. *Front Biosci (Landmark Ed)*. 21:42-54.

Vanderweyde, T., H. Yu, M. Varnum, L. Liu-Yesucevitz, A. Citro, T. Ikezu, K. Duff, and B. Wolozin. 2012. Contrasting pathology of the stress granule proteins TIA-1 and G3BP in tauopathies. *J Neurosci*. 32:8270-8283.

Wang, J., J.M. Choi, A.S. Holehouse, H.O. Lee, X. Zhang, M. Jahnel, S. Maharana, R.

Lemaitre, A. Pozniakovsky, D. Drechsel, I. Poser, R.V. Pappu, S. Alberti, and A.A. Hyman. 2018. A

Molecular Grammar Governing the Driving Forces for Phase Separation of Prion-like RNA Binding

Proteins. *Cell*. 174:688-699 e616.

Wang, T., X. Tian, H.B. Kim, Y. Jang, Z. Huang, C.H. Na, and J. Wang. 2022a. Intracellular

energy controls dynamics of stress-induced ribonucleoprotein granules. *Nat Commun*. 13:5584.

Wang, Y.T., Y. Lim, M.N. McCall, K.T. Huang, C.M. Haynes, K. Nehrke, and P.S. Brookes.

2019. Cardioprotection by the mitochondrial unfolded protein response requires ATF5. *Am J Physiol

Heart Circ Physiol*. 317:H472-H478.

Wang, Z., H. Bo, Y. Song, C. Li, and Y. Zhang. 2022b. Mitochondrial ROS Produced by Skeletal

Muscle Mitochondria Promote the Decisive Signal for UPRmt Activation. *Biomed Res Int*.

2022:7436577.

Wek, R.C., H.Y. Jiang, and T.G. Anthony. 2006. Coping with stress: eIF2 kinases and

translational control. *Biochem Soc Trans*. 34:7-11.

Wheeler, J.R., T. Matheny, S. Jain, R. Abrisch, and R. Parker. 2016. Distinct stages in stress

granule assembly and disassembly. *Elife*. 5:e18413.

Wippich, F., B. Bodenmiller, M.G. Trajkovska, S. Wanka, R. Aebersold, and L. Pelkmans. 2013.

Dual specificity kinase DYRK3 couples stress granule condensation/dissolution to mTORC1

signaling. *Cell*. 152:791-805.

Wolozin, B., and P. Ivanov. 2019. Stress granules and neurodegeneration. *Nat Rev Neurosci*.

20:649-666.

Yang, P., C. Mathieu, R.M. Kolaitis, P. Zhang, J. Messing, U. Yurtsever, Z. Yang, J. Wu, Y. Li,

Q. Pan, J. Yu, E.W. Martin, T. Mittag, H.J. Kim, and J.P. Taylor. 2020. G3BP1 Is a Tunable Switch

that Triggers Phase Separation to Assemble Stress Granules. *Cell*. 181:325-345.e328.

Yoo, H., C. Triandafillou, and D.A. Drummond. 2019. Cellular sensing by phase separation:

Using the process, not just the products. *J Biol Chem*. 294:7151-7159.

Yousefi, R., E.F. Fornasiero, L. Cyganek, J. Montoya, S. Jakobs, S.O. Rizzoli, P. Rehling, and

D. Pacheu-Grau. 2021. Monitoring mitochondrial translation in living cells. *EMBO Rep*. 22:e51635.

Zhang, S., A. Macias-Garcia, J. Velazquez, E. Paltrinieri, R.J. Kaufman, and J.J. Chen. 2018.

HRI coordinates translation by eIF2αP and mTORC1 to mitigate ineffective erythropoiesis in mice

during iron deficiency. *Blood*. 131:450-461.

Zhao, Q., J. Wang, I.V. Levichkin, S. Stasinopoulos, M.T. Ryan, and N.J. Hoogenraad. 2002. A

mitochondrial specific stress response in mammalian cells. *Embo j*. 21:4411-4419.

Zhou, Z., J. Lu, M. Yang, J. Cai, Q. Fu, J. Ma, and L. Zhu. 2022. The mitochondrial unfolded

protein response (UPR(mt)) protects against osteoarthritis. *Exp Mol Med*. 54:1979-1990.

Zoncu, R., A. Efeyan, and D.M. Sabatini. 2011. mTOR: from growth signal integration to cancer,

diabetes and ageing. *Nat Rev Mol Cell Biol*. 12:21-35.

1199

Figure 1

Figure 2

Figure 3

Figure 4

A

	DMSO	Ars	GTPP 4 h	GTPP + Ars
Ctrl				
GADD34 KO				

G3BP1 eIF3η DAPI

B

Cells displaying SG (%)

** ****

○ Ctrl
■ GADD34 KO

DMSO   Ars   GTPP   GTPP+Ars

C

	NT	Ars	Pq 48 h	Pq + Ars
Ctrl				
GADD34 KO				

G3BP1 eIF3η DAPI

D

Cells displaying SG (%)

****

○ Ctrl
■ GADD34 KO

NT   Ars   Pq   Pq+Ars

E

Ctrl | GADD34 KO
GTPP: - - + + - - + +
Ars: - + - + - + - +
kDa
100 — GADD34
37 — p-eIF2α
1 305 4 135 1 208 33 187
37 — eIF2α total
150 — eIF3A

F

Ctrl | GADD34 KO
Pq: - - + + - - + +
Ars: - + - + - + - +
kDa
100 — GADD34
37 — p-eIF2α
1 269 6 123 1 616 7 519
37 — eIF2α total
37 — GAPDH

Figure 5

Figure 6

A

WT   G3BP1/2 dKO
DMSO   GTPP   DMSO   GTPP

TOMM20

B

mitochondria morphology %

■ Tubular
⋯ Fragmented

**

GTPP   -   +   -   +
        WT   G3BP1/2
                  dKO

C

mitoSOX mean intensity (relative to DMSO)

****
****   ****

□ DMSO
■ GTPP

WT   G3BP1/2 dKO

D

OCR (pmol/min/ug)

Oligomycin   FCCP   Rotenone & antimycin A

Spare capacity

Basal respiration

ATP-linked respiration

Maximal respiration

H⁺ leak

Non-mitochondrial oxygen consumption

Time (minutes)

E

OCR (pmol/min/ug)

—■— WT DMSO
--○-- G3BP1/2 dKO DMSO
—■— WT GTPP
--○-- G3BP1/2 dKO GTPP

Time (minutes)

F

OCR fold change

Basal respiration

*

WT   G3BP1/2 dKO

Maximal respiration

* * *

WT   G3BP1/2 dKO

ATP production-coupled respiration

* *

WT   G3BP1/2 dKO

% Spare respiratory capacity

* * * *   * *

WT   G3BP1/2 dKO

% Coupling efficiency

* * * *

WT   G3BP1/2 dKO

□ DMSO
■ GTPP

Figure 7

Figure 8

A

B