Identification and characterization of intermediate states in mammalian neural crest cell

epithelial to mesenchymal transition and delamination

Ruonan Zhao[1,2], Emma L. Moore[1], Madelaine M Gogol[1], Jay R. Uhruh[1], Zulin Yu[1], Allison Scott[1], Yan Wang [1], Naresh Kumar Rajendran[1] and Paul A. Trainor[1,2*]

[1]Stowers Institute for Medical Research, Kansas City, MO, USA

[2]Department of Anatomy and Cell Biology, University of Kansas Medical Center, Kansas City, KS, USA

*Author for Correspondence: pat@stowers.org

Keywords: neural crest cells; epithelial to mesenchymal transition; EMT; epithelial to mesenchymal plasticity; EMP; mouse; cell cycle; Dlc1

Abstract

Epithelial to mesenchymal transition (EMT) is a cellular process that converts epithelial cells to
mesenchymal cells with migratory potential in both developmental and pathological processes.
Although originally considered a binary event, EMT in cancer progression involves intermediate
states between a fully epithelial and a fully mesenchymal phenotype, which are characterized
by distinct combinations of epithelial and mesenchymal markers. This phenomenon has been
termed epithelial to mesenchymal plasticity (EMP), however, the intermediate states remain
poorly described and it's unclear whether they exist during developmental EMT. Neural crest
cells (NCC) are an embryonic progenitor cell population that gives rise to numerous cell types
and tissues in vertebrates, and their formation is a classic example of developmental EMT. An
important feature of NCC development is their delamination from the neuroepithelium via
EMT, following which NCC migrate throughout the embryo and undergo differentiation. NCC
delamination shares similar changes in cellular state and structure with cancer cell invasion.
However, whether intermediate states also exist during NCC EMT and delamination remains
unknown. Through single cell RNA sequencing, we identified intermediate NCC states based on
their transcriptional signature and then spatially defined their locations in situ in the
dorsolateral neuroepithelium. Our results illustrate the progressive transcriptional and spatial
transitions from premigratory to migratory cranial NCC during EMT and delamination. Of note
gene expression and trajectory analysis indicate that distinct intermediate populations of NCC
delaminate in either S phase or G2/M phase of the cell cycle, and the importance of cell cycle
regulation in facilitating mammalian cranial NCC delamination was confirmed through cell cycle
inhibition studies. Additionally, transcriptional knockdown revealed a functional role for the
intermediate stage marker *Dlc1* in regulating NCC delamination and migration. Overall, our
work identifying and characterizing the intermediate cellular states, processes, and molecular
signals that regulate mammalian NCC EMT and delamination furthers our understanding of
developmental EMP and may provide new insights into mechanisms regulating pathological
EMP.

Introduction

Epithelial to mesenchymal transition (EMT) is a cellular process that converts epithelial cells to mesenchymal cells with migratory potential (Hay 1995). EMT plays an essential role in various developmental and pathological processes such as embryonic morphogenesis, wound healing, tissue fibrosis and cancer progression (Zhao and Trainor 2023). Studies of EMT, particularly in the field of cancer biology, have increased exponentially in the past 5 years due to the implied role of EMT in numerous aspects of malignancy such as cancer cell invasion, survival, stemness, metastasis, therapeutic resistance and tumor heterogeneity (Yang et al. 2020a).

EMT has traditionally been considered a binary process comprising either full epithelial cell or mesenchymal cell states. However, studies of cancer have uncovered an alternative scenario termed epithelial to mesenchymal plasticity (EMP), in which multiple intermediate states exist along the EMT spectrum (Dong et al. 2018; Gonzalez et al. 2018; Huang et al. 2013; Karacosta et al. 2019; Kumar et al. 2019; Pastushenko et al. 2018). More specifically, in some cases of head and neck cancer, primary tumors and matching lymph nodes contain a subpopulation of tumor cells in a partial EMT state, as defined by their expression of both epithelial marker genes and mesenchymal marker genes (Puram et al. 2017). Moreover, cells in this partial EMT state are highly invasive and located at the leading edge of tumors *in vivo*. Similarly, in mouse xenograft models of skin cancer and breast cancer, the absence of epithelial cellular adhesion molecules in several cell populations was discovered to represent early EMT hybrid states as they also expressed both Vimentin and Cytokeratin 14 at intermediate levels and consequently had high metastatic potential (Pastushenko et al. 2018). Partial EMT states have also been recognized in lung cancer and ovarian cancer (Gonzalez et al. 2018; Karacosta et al. 2019), with tumor cells expressing both the epithelial marker E-cadherin, and the mesenchymal marker Vimentin. Besides primary tumors, EMT intermediate states have also been identified in circulating tumor cells from patient samples (Yu et al. 2013). Even though EMT intermediate states have been discovered in numerous studies, few have focused on describing and understanding the molecular and cellular mechanisms governing or defining each intermediate state due to the challenges of studying cancer initiation and progression *in vivo*.

Since EMT during embryogenesis and cancer progression have been shown to share analogous phenotypic changes that involve similar core transcription factors and molecular mechanisms, it was proposed that the initiation and development of carcinoma could be attributed to an unusual activation of EMT factors involved in normal developmental processes (Hay 1995). However, compared to tumorigenesis, it remains largely unknown whether intermediate or transition states exist or play a role in classic developmental EMT. Therefore, identifying and characterizing intermediate states during developmental EMT can further our understanding of the cellular processes, and molecular signaling networks that regulate EMP.

NCC formation is a classic example of developmental EMT (Lee et al. 2013; Zhao and Trainor 2023). NCC are a migratory progenitor cell population unique to vertebrates. Formed during neurulation in the dorsolateral domain of the neural plate, EMT facilitates their delamination from the neuroepithelium and migration throughout the body, where they differentiate into neurons and glia of the peripheral nervous system, pigment cells in the skin,

craniofacial bone and cartilage, as well as many other cell types (Bhatt et al. 2013; Dash and
Trainor 2020; Lièvre and Douarin 1975; Trainor 2005; Weston 1983). Disruption of NCC
delamination and migration can result in developmental abnormalities, referred to as
neurocristopathies (Achilleos and Trainor 2015; Watt and Trainor 2014), hence it is important
to study the mechanisms that regulate mammalian NCC development.
We performed single cell RNA sequencing (scRNA-seq) to identify and define
intermediate transcriptional and cellular states during mouse cranial NCC EMT and
delamination. We identified two NCC EMT intermediate populations distinguished by their S or
G2/M cell cycle phase state during delamination. Interestingly, trajectory analyses reveal that
these distinct intermediate populations are formed simultaneously, and independently, but
then converge into a single or common pool, suggesting they do not have distinct fates
following migration. This is consistent with the known plasticity and potency of early migrating
NCC (Golding et al. 2000; Sandell and Trainor 2006; Trainor and Krumlauf 2000a; Trainor and
Krumlauf 2000b, 2001). Transcriptional profiling revealed that the intermediate NCC
populations could also be defined by unique transcriptional signatures, including differential
expression of genes involved in cell protrusion, such as *Dlc1*, *Pak3* and *Sp5*. Further
interrogation using SABER-FISH, revealed that these intermediate NCC populations were
spatially localized in the dorsolateral region of the neural plate. In addition, knocking down the
NCC EMT intermediate marker *Dlc1* led to a significant reduction in the number of migratory
NCC, which revealed a critical role for *Dlc1* in the regulation of mouse cranial NCC delamination.
Overall, our findings provide novel, detailed, high-resolution descriptions of the intermediate
cell populations, and transcriptional states that occur during cranial NCC EMT and delamination
in mouse embryos. Our work further illustrates that molecular characterization of NCC EMT
intermediate states can reveal essential regulatory components of mouse NCC EMT and
delamination. These results shed light on similar mechanisms of NCC EMT and delamination in
other mammalian species as well and will also serve as a resource for the community. In
addition to furthering our understanding of the cellular processes, and molecular signaling
networks that regulate NCC and EMT delamination, our work may also help to inform the
phenotypic changes and corresponding gene regulatory control of EMP in other developmental
EMT events as well as pathological conditions such as tissue fibrosis and cancer progression.
118
Results

Identification of intermediate stages during mouse cranial NCC EMT and delamination

To investigate the biological process and mechanisms governing mouse cranial NCC EMT
and delamination, we performed scRNA-seq on dissociated cranial tissues isolated from E8.5
mouse embryos with 7-9 somites (Figure 1A). More specifically, embryos were collected from
two transgenic mouse lines: *Wnt1-Cre;RosaeYFP* (Chai et al. 2000) and *Mef2c-F10N-LacZ* (Aoto
et al. 2015) (Figure 1A). In E8.5 *Wnt1-Cre;RosaeYFP* embryos, YFP is expressed by *Wnt1+*
neuroepithelial cells located in the dorsolateral neural plate, which encompasses premigratory
NCC (Figure 1B). Consequently, *Wnt1-Cre;RosaeYFP* labels premigratory and migratory NCC and
other lineage labeled cells derived from the *Wnt1+* cell population (Figure 1B) (Chai et al. 2000).
In contrast, *Mef2c-F10N-LacZ* predominantly labels migratory NCC, with lacZ activity driven by

the F10N enhancer of the *Mef2C* gene (Figure 1B) (Aoto et al. 2015). The two different
transgenic lines allowed us to distinguish premigratory from migratory NCC spatially within an
embryo, but also later bioinformatically following single cell dissociation and RNA-sequencing.
133   The scRNA-seq data was processed and analyzed as previously described (Falcon et al.
2022)*.* We initially identified 6 major cell or tissue types present in E8.5 mouse embryonic
cranial tissues based on the differential expression of classic cell or tissue type specific markers,
and we clustered the data accordingly (Figure 1C and 1D; Figure 1-figure supplement 1). For
example, *Sox1* and *Sox2* were used to delineate neural ectoderm, whereas *Cdh1* (E-cadherin)
was used to define non-neural ectoderm. *eYFP*, *lacZ*, *Sox10* and *Twist* delineated migrating
neural crest cells. *Tbx1* was primarily used as a marker of mesoderm cells and *Kdr* (*Vegfr2*)
demarcated mesoderm-derived endothelial cells. We then bioinformatically segregated the
cranial NCC cluster, which includes both premigratory and migratory NCC, and divided it into 5
subclusters at a resolution of 0.26 (Figure 2A).
143   To characterize these NCC subclusters, we then interrogated the expression of known
neuroepithelial and neural plate border markers, as well as genes expressed by NCC during
their specification and migration (Supplementary Table 1) (Echelard et al. 1994; Hafemeister
and Satija 2019; Lee et al. 2013; Murdoch et al. 2012; Parr et al. 1993; Sauka-Spengler and
Bronner-Fraser 2008; Wood and Episkopou 1999). A high percentage of cells in subclusters 0-3
express elevated levels of markers of migratory NCC (*Vim*, *Sox10*, *Twist1*), while only
subclusters 0 and 1 exhibit high levels of expression of NCC specifier genes (*Zeb2*, *Pax3*, *Nr6a1*,
Sox9, *Foxd3*, *Snai1)* (Figure 2B) (Cheung et al. 2005; Dottori et al. 2001; Hari et al. 2012;
Kobayashi et al. 2020; Lee et al. 2013; Li et al. 2000; Murdoch et al. 2012; Schorle et al. 1996;
Soo et al. 2002; Van de Putte et al. 2003). Since NCC specifiers are downregulated as NCC
migrate and later differentiate into specific lineages, the combinatorial expression of genes
suggests that subclusters 0 and 1 likely represent an earlier stage of NCC delamination and
migration than subclusters 2 and 3. This conclusion was further verified by integrating
previously published networks of genes that represent an early migratory NCC program versus
a late migratory NCC program (Soldatov et al. 2019). Early migratory NCC program genes are
expressed by the majority of migratory NCC whereas late migratory NCC program genes are
only expressed in a subset of migratory NCC as they have already begun to mature.
160   Subclusters 0-3 each exhibits significant expression of early migratory NCC program
genes (Figure 2-figure supplement 1). Furthermore, subclusters 2 and 3 express a significantly
higher level of late migratory NCC program genes than subclusters 0 and 1, which demonstrates
that subclusters 0 and 1 contain early migratory NCC, whereas subclusters 2 and 3 comprise
late migratory NCC (Figure 2-figure supplement 1)*.* Interestingly, we also observed that a small
percentage of subcluster 0 cells express early migratory NCC genes, but at a lower intensity
compared to subcluster 1. This implies that subcluster 0 might also contain premigratory NCC
that do not yet express any migratory NCC genes (Figure 2-figure supplement 1). Despite similar
expression profiles, NCC subclusters 2 and 3 possibly represent undifferentiated NCC derived
mesenchyme tissue in different parts of the head. More specifically, subcluster 2 displays a high
level of expression of pharyngeal arch NCC mesenchyme marker *Dlx2* (Figure 2-figure
supplement 1) (Bulfone et al. 1993). In contrast, subcluster 3 expresses a frontonasal
mesenchyme marker *Alx1* (Figure 2-figure supplement 1) (Iyyanar et al. 2022). Consistent with
these observations, neither population expresses a high level of neurogenic lineage markers

such as *Nrp1* and Nrp2, which is indicative of their undifferentiated state (Figure 2-figure
supplement 1) (Lumb et al. 2014).
Subcluster 4 exhibits a unique transcriptional profile distinct from subclusters 0-3
(Figure 2B). Higher expression of *Sox1* and *Sox2* is indicative of a neuroepithelial identity for
subcluster 4 (Figure 2B). During NCC formation, Sox1 and Sox2 activity are downregulated in
the dorsal neural plate border domain but remain strongly expressed more ventrally
throughout the neural plate (Supplementary Table 1). Concomitantly, Sox9 and Sox10 are
activated in what is known as the SoxB (1/2) to SoxE (9/10) switch (Mandalos et al. 2014;
Remboutsika et al. 2011; Wakamatsu et al. 2004). Consistent with this model, overexpressing
Sox2 in the dorsal neural tube has been shown to repress NCC specification, whereas
overexpressing *Sox9* and *Sox10* have been shown to precociously promote NCC formation
(Aybar et al. 2003; Cheung and Briscoe 2003; Mandalos et al. 2014; McKeown et al. 2005;
Remboutsika et al. 2011).
To determine if mouse cranial NCC EMT is non-binary, and occurs through intermediate
or transition states, we further subdivided the 5 cranial NCC clusters into smaller subclusters
(resolution=2.0; subcluster 1'-21') and extracted 15 subclusters out of the original early
migratory NCC subclusters 0, 1 and 4 (Figure 2C; Figure 2-figure supplement 1). Through
heatmap analysis, expression of the same NCC marker genes as described above was examined
within these new subclusters, and the order of the subclusters was arranged according to their
combinatorial expression patterns (Figure 2D)*. For example, *Wnt1* and *Sox10* were used to
identify NCC transitioning from premigratory to migratory states since *Wnt1* is only expressed
in premigratory NCC and is immediately downregulated as NCC delaminate and start to
migrate. In contrast, *Sox10* is activated only after NCC have delaminated and begun to migrate.
The heatmap shows that a significant number of cells in subcluster 17' express a much higher
level of *Wnt1* than any other subcluster, suggesting that subcluster 17' comprises premigratory
NCC (Figure 2D). Consistent with this observation, subcluster 17' cells also express other neural
plate border and NCC specifier genes such as *Zic2*, *Pax7*, *Nr6a1*, *Pax3*, *Sox9*, and *Foxd3,* but
does not express migratory NCC markers such as *Sox10* and *Vim* (Figure 2D). Subclusters 2' and
10' share a similar expression profile to subcluster 17'. However, subcluster 2' and 10' express
less *Wnt1* and less neural plate border specifiers such as *Zic2* and *Pax7* than subcluster 17'
(Figure 2D; Figure 2-figure supplement 1). This data suggests that subclusters 2' and 10' could
represent EMT intermediate states as premigratory NCC transition to migratory NCC during
delamination. Lastly, the remaining early NCC subclusters express *Sox10* and *Vim* indicating that
they comprise or represent migratory NCC (Figure 2D).
To further validate the identity of subclusters 2' and 10' as representing intermediate
cellular stages of EMT, we assessed the expression of genes associated with adherens junctions,
tight junctions, and apical-basal polarity, which are required to maintain epithelial integrity, and
cytoskeleton rearrangement that is typically associated with EMT (Dongre and Weinberg 2019;
Matsuuchi and Naus 2013; Radisky and Radisky 2010) (Zhao and Trainor 2023). Intermediate
NCC exhibit a decrease in *Nectin*, *Cadherin* and *Tight Junction Protein* gene expression
consistent with intercellular tight junction breakdown and degradation of apicobasal polarity
(Figure 2E). At the same time, subcluster 2 and 10 cells exhibit an increase in *Rac*, *Rack1* and
Cdc42 gene expression which is indicative of cytoskeletal rearrangement and the formation of

cell protrusions (Figure 2E). These alterations in gene expression are indicative of the molecular
cellular mechanisms that underpin EMT (Figure 2E).
219
220 **Mouse cranial NCC undergo EMT and delamination in S or G2/M phase cell cycle**
221
Since intermediate NCC stages represent a transitional or intermediate phase between
premigratory and migratory NCC, the molecular and signaling pathway signatures uniquely
expressed by these intermediate NCC can reveal essential regulatory mechanisms governing
NCC EMT and delamination. Analysis of cell cycle gene expression, for example, indicated that
EMT intermediate NCC populations exhibit distinct cell cycle phase properties or characteristics.
NCC in subcluster 2' primarily express S phase cell cycle genes such as *Pcna*, *Pol1* subunits, *Plk1*,
Ccnd and *Mcm* family members. In contrast, NCC in subcluster 10' predominantly express G2/M
phase cell cycle genes including *Mik67*, *Aurka/b*, *Cenp* and *Kif* family members (Figure 3A and
3B).
To understand the potential for any temporal or lineage relationships between the two
intermediate stage NCC populations based on their different cell cycle states, we performed
pseudotime trajectory analysis on the entire NCC population using Monocle 3 (Figure 3C and
3D). Premigratory and EMT intermediate NCC were identified as the earliest discrete
populations to form among the entire cranial NCC population (Figure 3C). In contrast, late
migratory NCC represent a more mature stage of NCC development (Figure 3C). Results from
the pseudotime analysis indirectly support the identities previously assigned to the cranial NCC
subclusters under both resolutions. The trajectory analysis also demonstrates that NCC can
arise through two independent paths and initially become two distinct intermediate
populations (subcluster 2' and 10') during EMT (Figure 3D). Later the trajectories or lineages of
the intermediate NCC subclusters then merge back together into a single or common
population of early migratory NCC, before ultimately maturing into still as yet undifferentiated
late migratory NCC as they colonize the frontonasal or pharyngeal arch mesenchyme (Figure
3D). These results imply that NCC representing two distinct intermediate stages form
simultaneously and independently during development. Moreover, the different cell cycle
status of subclusters 2' and 10' suggest that premigratory NCC can undergo EMT and
delamination in either S phase or G2/M phase of the cell cycle. The trajectory analysis further
depicts that EMT intermediate NCC and their immediate lineages are not fate restricted to any
specific cranial NCC derivative at this timepoint.
250
251 **Cell cycle regulation is critical for mouse cranial NCC EMT and delamination**
252
Since EMT intermediate NCC are either in S phase or G2/M phase of the cell cycle, we
then investigated whether cell cycle regulation plays a significant role in driving mouse cranial
NCC delamination. We dissected E8.5 *Wnt1-Cre;RosaeYFP* mouse embryos and examined the
expression of cell cycle markers to compare the cell cycle status of delaminating NCC at the
neural plate border in cranial tissues versus premigratory non-delaminating NCC in in the neural
plate border of the trunk. EdU and phospho-histone H3 (pHH3) were used to label S phase and
G2/M phase of the cell cycle respectively (Figure 4A). A majority of delaminating cranial NCC
express either EdU or pHH3 or both. Only a very small percentage of cells do not express either

of these cell cycle markers (Figure 4C). In contrast, almost 50% of premigratory, non-
delaminating trunk NCC in the dorsolateral neural tube, do not express either cell cycle marker
(Figure 4D). These observations imply that specific cell cycle phases are intimately connected to
mouse cranial NCC EMT and delamination.
To further validate the association between cell cycle status and cranial NCC EMT and
delamination, we inhibited S phase during early NCC development by incubating E8.0 CD1
mouse embryos in whole embryo roller culture with Aphidicolin. After 12 hours of treatment,
we quantified the number of migratory NCC via Sox10 immunostaining to determine the
number of premigratory NCC that delaminated. Cell cycle status was also evaluated in DMSO
(control) and Aphidicolin treated samples via EdU and pHH3 staining. Based on our trajectory
analyses, we hypothesized that inhibiting S phase progression would block S phase
delamination but not G2/M phase delamination. As we expected, the EdU signal was
completely absent in Aphidicolin treated embryos demonstrating that cells cannot enter S
phase post treatment (Figure 4-figure supplement 1). The Aphidicolin treatment did not induce
cell death as the level of TUNEL staining in treated embryos was similar to that observed in
controls (Figure 4-figure supplement 1). Quantification of Sox10 positive cells revealed
significantly fewer migratory neural crest cells in the craniofacial region of Aphidicolin treated
embryos compared to DMSO treated control embryos (Figure 4E). Similarly, we also treated
E8.0 *Mef2c-F10N-LacZ* embryos with Aphidicolin for 12 hours in roller culture and observed
fewer migratory NCC compared to control embryos as evidenced by LacZ staining (Figure 4-
figure supplement 1). Interestingly, pHH3 is expressed by a major proportion of the remaining
migratory NCC after Aphidicolin treatment (Figure 4B). This data is consistent with the
pseudotime trajectory analysis that cranial NCC delamination in G2/M phase of the cell cycle
(subcluster 10' NCC) is independent of delamination in S phase of the cell cycle (subcluster 2'
NCC). Disrupting S phase of the cell cycle didn't prohibit EMT intermediate NCC in G2/M phase
of the cell cycle from delaminating and forming migratory NCC that express G2/M phase cell
cycle markers. Thus, our data shows that cranial NCC delamination is disrupted upon S phase
cell cycle inhibition, which supports the hypothesis that cell cycle regulation is critical for cranial
NCC delamination in mouse embryos.
290
291 **Spatiotemporal localization of intermediate stage NCC *in vivo***
292
To identify and define EMT intermediate stage NCC *in vivo* during mouse cranial NCC
development, we used the scRNA-seq data to extract genes that were differentially expressed
(threshold based on average logFC ≥ 0.25) in the intermediate NCC populations (Figure 5-figure
supplement 1). Among the potential marker genes, we then selected *Dlc1*, *Sp5* and *Pak3* based
on their relatively high expression levels and specificity in cranial NCC, and more importantly,
intermediate NCC populations (Figure 5A; Figure 5-figure supplement 1). During cranial NCC
development, *Sp5* and *Pak3* are expressed at high levels in both premigratory and intermediate
stage NCC. In contrast, *Dlc1* is highly expressed in intermediate and migratory NCC (Figure 5A).
To further distinguish between premigratory, intermediate and migratory NCC, *Wnt1* (a
premigratory NCC marker) and *Sox10* (a migratory NCC marker) were also included in the *in situ*
identification analyses alongside *Dlc1*, *Sp5* and *Pak3* (Figure 5A).

To confirm that the intermediate stage markers *Dlc1, Sp5* and *Pak3* are expressed
during NCC delamination, and determine the spatial location of the intermediate stage NCC, we
performed signal amplification by exchange reaction for multiplexed fluorescent in situ
hybridization (SABER-FISH). SABER-FISH was chosen for our multiplexed analyses of gene
expression because SABER-FISH probes lack secondary structure which facilitates increased
sensitivity and depth of tissue penetration. SABER-FISH oligo pools were designed for *Wnt1*,
Sox10, Dlc1, Sp5 and *Pak3,* using stringent parameters (Kishi et al. 2019). Intermediate stage
marker probes were validated by comparing the expression patterns of the SABER-FISH staining
for *Dlc1, Sp5* and *Pak3* with traditional in situ hybridization staining in transverse histological
sections (Figure 5-figure supplement 2). The expression of *Dlc1, Sp5* and *Pak3* matched
between SABER-FISH and traditional in situ hybridization methods, validating our probe design
(Figure 5-figure supplement 2*).*
Following individual validation, we then performed combined staining to visualize the
spatial distribution of all the genes in the same tissue section (Figure 5B). The intermediate
stage markers *Dlc1, Sp5* and *Pak3* appeared to overlap in expression in the dorsal most region
of the neural fold, where EMT takes place as evidenced by the presence of *Sox10* labelled
migratory NCC adjacent to the neuroepithelium (Figure 5B). This overlap in expression was
notably not observed in older embryos in areas where EMT had concluded. To better visualize
and confirm co-localized expression of these genes in the same dorsolateral region of the
neural plate border, we generated polyline kymographs depicting the average intensity of each
gene's fluorescent signal along the dorsal most region of the neural fold and into the migratory
NCC population (Figure 5C). We observed a consistent pattern of activity in which *Wnt1* is
highly expressed in the dorsal neuroepithelium (Figure 5C). However, in the most dorsolateral
domain, where *Wnt1* expression is slightly diminished, the intermediate stage NCC markers
Dlc1, Sp5 and *Pak3* are highly expressed (Figure 5C). In contrast, minimal *Sox10* expression is
detected in this transition region at the edge of the neuroepithelium, but high levels of *Sox10* in
the clear absence of *Wnt1, Dlc1, Sp5* and *Pak3,* is observed in migratory NCC located more
ventrally (Figure 5C). Therefore, our data indicates that EMT intermediate stages can not only
be transcriptionally defined, but also spatially resolved to the dorsal most region of the
neuroepithelium.
334
335   **EMT intermediate stage marker gene Dlc1 regulates NCC delamination**
336
Having transcriptionally defined intermediate stage NCC and determined their special
location during delamination, it was important to test whether any of the intermediate stage
signature genes, *Dlc1*, *Pak3* or *Sp5*, play functional or essential roles in NCC development. We
prioritized *Dlc1* over *Pak3* and *Sp5* because *Dlc1* is not expressed by premigratory NCC but is
expressed at high levels in all EMT intermediate stage NCC. Furthermore, *Dlc1* null mutant mice
are embryonically lethal and exhibit craniofacial malformation phenotypes, which is suggestive
of a perturbation of NCC development. In contrast, *Pak3* and *Sp5* are expressed in premigratory
NCC, but in only 50-60% of intermediate stage NCC (Figure 5A). Moreover, *Pak3* and *Sp5* null
mutant mice are healthy and fertile with no obvious abnormalities. Therefore, we hypothesized
that alone, *Dlc1* loss-of-function would more likely perturb cranial NCC delamination.

To test this hypothesis, we knocked down *Dlc1* by injecting *Dlc1* shRNA-based
lentiviruses and control scrambled shRNA lentiviruses into the amniotic cavity of E7.5 CD1
mouse embryos. Since the neural plate remains open at this developmental stage, all
neuroepithelial cells are exposed or in contact with amniotic fluid containing virus. The embryos
were then cultured for 24 hours after which we assessed the number of migratory NCC via
Sox10 staining, to determine how many premigratory NCC underwent EMT and delamination
(Figure 6A). We subsequently observed that the number of migratory NCC was significantly
reduced in all *Dlc1* knockdown embryos (Figure 6B). In fact, each of the *Dlc1* shRNA constructs
led to a significant reduction in the number of migratory NCC compared to their respective
control (Figure 6-figure supplement 1*).* Importantly, we did not observe any difference in cell
death in *Dlc1* knockdown embryos comparable to controls (Figure 6D). On average, a 30%
reduction of *Dlc1* expression was achieved by each *Dlc1* shRNA lentivirus knockdown (Figure 6C;
Figure 6-figure supplement 1), and notably, the *Dlc1* isoforms targeted by the different *Dlc1*
shRNA constructs used in this study suggest a correlation with *Dlc1* null mouse embryos and
their phenotypes.
Dlc1a shRNA construct targets exon 1 specifically in *Dlc1* mRNA variant 2. Consequently,
Dlc1a shRNA is capable of exclusively eliminating the expression of *Dlc1* mRNA variant 2
(isoform 2) since the same exon region is not present in variant 1 or 3. Interestingly, one of the
Dlc1 null mouse models, *Dlc1*$^{gt/gt}$, carries a gene trap vector inserted into intron 1, which results
in reduction of the 6.1 kb transcript (Dlc1 isoform 2) alone (Sabbir et al. 2010). Therefore, it is
possible that the craniofacial phenotypes observed in *Dlc1*$^{gt/gt}$ null mice are caused by abnormal
cranial NCC EMT and delamination as shown by *Dlc1a* knockdown in our data. In contrast, *Dlc1b*
and *Dlc1c* shRNA constructs both target exon 5 of *Dlc1* mRNA variant 2 and 3, which is the same
as exon 9 of variant 1. Consequently, all three *Dlc1* variants should be diminished by *Dlc1b* and
Dlc1c shRNAs. In another *Dlc1* null mouse model, exon 5 was deleted by replacing it with a
neomycin resistance gene, which caused a reading frame shift and subsequently premature
translation termination (Durkin et al. 2005). This led to the synthesis of truncated polypeptides
containing only the first 77 amino acids, which encode the sterile alpha motif (SAM) protein
interaction domain and 23 novel residues. Since *Dlc1*$^{-/-}$ null, *Dlc1b* and *Dlc1c* shRNAs all cause
disruptions in exon 5/9 of *Dlc1* transcripts, there is a strong correlation between the *Dlc1*$^{-/-}$ null
craniofacial phenotypes and cranial NCC EMT defects observed in *Dlc1b* and *Dlc1c* knockdown
mouse embryos. Collectively, these results therefore demonstrate an important functional role
for Dlc1 in mammalian NCC EMT and delamination. Moreover, our results suggest that genes
that are differentially expressed in intermediate NCC could play a regulatory role during EMT
and delamination.
382
Discussion

EMT is a cellular process that converts epithelial cells to mesenchymal cells. EMT is
essential for normal development and is a key driver of disease pathogenesis, particularly
cancer metastasis. Although classically considered to be a binary event, studies of EMT in
cancer identified multiple intermediate states within the EMT spectrum, a phenomenon termed
epithelial to mesenchymal plasticity (EMP). However, it remained to be determined whether
developmental EMT is also a developmental EMP process. Our goal therefore was to determine

whether intermediate stages of NCC development during EMT could be transcriptionally and
spatially defined, and then test whether transitional stage associated genes are functionally
required for NCC EMT and delamination.
Through scRNA-seq analysis of mouse cranial tissues at E8.5, which coincided with the
onset of NCC EMT and delamination, we identified two populations of NCC, whose gene
expression profiles or signatures were representative of intermediate stages between
premigratory and migratory NCC. Furthermore, we determined that the two intermediate
populations could be defined by their distinct transcriptional states which were consistent with
being in either S phase or G2/M phase of the cell cycle. Pseudotime trajectory analysis suggests
that these intermediate stage cranial NCC populations can undergo EMT and delaminate in
either S phase or G2/M phase, simultaneously, and independently of each other, but then later
merge into a single or common pool of early migratory NCC, suggesting they do not have
distinct fates following migration. This is consistent with the known plasticity and potency of
migrating NCC (Sandell and Trainor 2006; Trainor and Krumlauf 2000a; Trainor and Krumlauf
2000b, 2001).
These results correlate with observations that cell cycle status is also a critical factor
regulating NCC delamination in avian and zebrafish embryos. For example, BrdU incorporation,
which demarcates proliferating cells in S phase of the cell cycle, was previously used to evaluate
the cell cycle status of emigrating trunk NCC, dorsal midline neuroepithelial cells and
surrounding cells at the segmental plate, epithelial somite and dissociating somite axial levels in
chicken embryos (Burstyn-Cohen and Kalcheim 2002). Most emigrating trunk NCC (about 80%)
at the epithelial somite and dissociating somite axial levels were in S phase of the cell cycle,
while less than 50% of dorsal neuroepithelial cells were BrdU+. Similarly, slice culture of the
trunk of chicken embryos also revealed that most premigratory NCC contained basally
positioned nuclei indicative of S phase. Interestingly, however, a small proportion of
premigratory NCC presented as round mitotic cells, whose daughter cells later became
migratory (Ahlstrom and Erickson 2009). In comparison, *in vivo* time-lapse imaging of chicken
embryos revealed that half of the delaminating trunk NCC that were tracked, displayed signs of
cell division (McKinney et al. 2013). However, in most cases, only one progeny of a mitotic
premigratory NCC was observed to exit the neural tube and become a migratory NCC. In
contrast to delaminating trunk NCC, only around 30% of delaminating cranial NCC in chicken
embryos were found to be in S phase (Théveneau et al. 2007) illustrating considerable
differences between cranial and trunk neural crest cells and in the correlation between cell
cycle phase and delamination. Interestingly, time-lapse imaging of zebrafish embryos also
revealed active cell division in dorsal neuroepithelial cells prior to NCC delamination and EMT.
The daughter cells of those divisions then translocate into the basal side of the
neuroepithelium, where EMT subsequently occurs (Berndt et al. 2008). Whether delaminating
NCC undergo proliferation and cell division in zebrafish embryos remains to be investigated,
however, active cell division in the dorsal neuroepithelium is a shared feature of NCC
delamination in avian and aquatic species. Even though our findings primarily illustrate the
association of distinct cell cycle phases with intermediate stage NCC during EMT and
delamination, this work has emphasized the importance of further examining the cell division
and cell cycle activities of delaminating NCC in mouse embryos as critical contributors to normal
development and the pathogenesis of neurocristopathies.

Blocking G1/S transition in chicken embryos via *in ovo* electroporation, or via small
molecule inhibitors in explanted neural primordia, prevents the onset of NCC delamination
(Burstyn-Cohen and Kalcheim 2002). Furthermore, BMP and Wnt canonical signaling regulates
the G1/S transition and promotes trunk NCC delamination (Burstyn-Cohen et al. 2004).
Although it remains to be determined which signaling pathways regulate cell cycle phase
progression in the intermediate or transitional populations of NCC in mouse embryos, our
results illustrate an evolutionarily conserved mechanistic role for cell cycle progression in NCC
delamination in vertebrate embryos.
The intermediate populations of cranial NCC in mouse embryos exhibited
transcriptional profiles that were characterized by the downregulation of tight junction and
polarity genes. This is consistent with the breakdown of intercellular tight junctions and
degradation of apicobasal polarity, which are hallmarks of EMT (Zhao and Trainor 2023).
Further interrogation of genes that were differentially expressed in the intermediate NCC
populations revealed *Dlc1*, *Sp5* and *Pak3* based on their relatively high expression levels as
potentially specific markers, and regulators of intermediate NCC populations. Through SABER-
FISH staining of *Dlc1*, *Sp5* and *Pak3* in combination with *Wnt1* as a marker of premigratory NCC,
and *Sox10* as a marker of migratory NCC, we spatially resolved the location of intermediate NCC
to the most dorsolateral domain of the cranial neural plate in E8.5 mouse embryos. We then
prioritized *Dlc1* for functional analyses because it is expressed at high levels in all EMT
intermediate stage NCC, but not in premigratory NCC. Lentiviral shRNA knockdown of *Dlc1* in
cultured mouse embryos resulted in a significant reduction in the number of migratory neural
crest cells, which may account for the craniofacial and cardiac malformation phenotypes
observed in *Dlc1* null mutant mice (Sabbir et al. 2010). Thus *Dlc1*, which is primarily expressed
in intermediate stage NCC, plays an important role in NCC EMT and delamination. Dlc1 is a Rho
GTPase Activating Protein (GAP) that regulates the activity of Rho family GTPases Rho and
Cdc42 (Kim et al. 2008). RhoGTPases are known to regulate cell morphology and motility
through modulating the activity of the actin cytoskeleton. More specifically, Rho has been
shown to facilitate the formation of stress fibers, while Cdc42 is involved in filopodium
formation (Kim et al. 2008). RhoGTPases also regulate the organization of tight junctions, which
breakdown during EMT (Popoff and Geny 2009; Terry et al. 2010). Therefore, the effect of *Dlc1*
knockdown on cranial NCC EMT and delamination may be mediated via disrupted RhoGTPase
activity and subsequent downstream cellular changes.
A similar regulatory role for Dlc1 and other GAP family members has also been observed
during NCC EMT and delamination in chicken and zebrafish embryos. In chicken embryos, *Dlc1*
overexpression results in ectopic trunk NCC delamination, including apically into the neural
tube lumen, due to a disruption in apical-basal polarity of dorsal neuroepithelial cells (Liu et al.
2017). Furthermore, NCC overexpressing *Dlc1* exhibit a loss of directionality during migration.
Conversely, *Dlc1* inhibition and depletion results in less NCC emigration and thus fewer
migratory NCC. In addition, the downregulation of *Dlc1* in migrating NCC restricts their motility.
Thus, *Dlc1* regulates trunk NCC delamination and migration in chicken embryos. Interestingly, in
zebrafish embryos, another GAP family member Arhgap has been shown to modulate NCC EMT
and delamination via the localization of Rho activation to designated subcellular compartments
and the promotion of localized actomyosin contraction to trigger directional cell motility (Clay
and Halloran 2013). More specifically, the knockdown of Arhgap in NCC results in Rho

activation, which in turn inhibits NCC EMT and delamination. These data suggest that the role
of Dlc1 during NCC delamination might be evolutionarily conserved in vertebrate embryos.
However, whether Dlc1 regulates mouse cranial NCC delamination through localized activation
of Rho remains to be determined. Nevertheless, consistent with similarities in EMT,
delamination and cell migration between neural crest cells and cancer cells, Dlc1 may also play
an important role in promoting cell migration during cancer progression. High levels of DLC1
expression are detected in most melanoma tissues, and functional studies have revealed that
DLC1 is both sufficient and required for melanoma growth and metastasis (Yang et al. 2020b).
In conclusion, through scRNA-sequencing we transcriptionally identified two distinct
intermediate stages of NCC during EMT and delamination based primarily on cell cycle status.
Delamination in S phase or G2/M phase seems to occur simultaneously but also independently,
resulting in a single or common pool of early migratory NCC. Further interrogation of our
transcriptomic dataset revealed *Dlc1* to be a key molecular marker of intermediate stage NCC,
and their location in situ in the dorsolateral neural plate, which we spatially resolved in E8.5
mouse embryos. Lastly, we tested and functionally validated that *Dlc1* plays an important role
in NCC delamination in mouse embryos. Taken together, our identification and characterization
of intermediate stage cranial NCC during their delamination are consistent with NCC EMT being
a developmental EMP event. Similar to intermediate EMT states in cancer metastasis, NCC
downregulate certain epithelial cell features but maintain co-expression of epithelial cell
markers and mesenchymal cell markers during EMT and delamination. Additionally,
intermediate stage NCC are localized at the dorsolateral edge of the neural plate border, which
is reminiscent of the localization of intermediate EMT states at the leading edge of invasion in
several types of primary tumors. However, unlike certain intermediate EMT cells present in the
lymph nodes or circulating tumor cells, mouse EMT intermediate NCC represent a transient
state and eventually form migratory NCC with mesenchymal character. This suggests that EMP
may be a more common developmental phenomenon. Our transcriptional data and signatures
of intermediate stage NCC during EMT and delamination can serve as a useful resource for the
community. This also now sets the stage for uncovering the gene regulatory networks that
govern intermediate stage NCC development and function, and for exploring whether EMP is a
feature of other developmental and pathological EMT events such as in gastrulation, wound
healing, and fibrosis.
509
Materials and Methods

Mice and animal husbandry

All mice were kept in a 16h-light and 8h-dark light cycle in the Laboratory Animal
Services Facility at the Stowers Institute for Medical Research. All animal experiments were
conducted in accordance with Stowers Institute for Medical Research Institutional Animal Care
and Use Committee (IACUC)-approved protocol (IACUC no. 2022-143). *Wnt1-Cre* mice
(*H2afv*^Tg(Wnt1-cre)11Rth Tg(Wnt1-GAL4)11Rth/J, Jax stock #003829) and *RosaeYFP* mice were
obtained from the Jackson Laboratory and maintained and genotyped as previously described
(Chai et al. 2000; Jiang et al. 2000). *Wnt1-Cre* was maintained as a heterozygous allele and
crossed to homozygous *RosaeYFP* to generate *Wnt1-Cre;RosaeYFP*. *Mef2c-F10N-LacZ* mice, in

which LacZ expression is regulated by a neural crest cell specific enhancer of *Mef2c*, were
maintained and genotyped as previously described (Aoto et al. 2015).
522
Immunohistochemistry

Wholemount

Embryos were dissected and fixed in 4% PFA (in PBS) overnight. The next day the
embryos were rinsed 3 times for 5 minutes in PBS, dehydrated through an increasing methanol
series from 25% methanol/PBS to 100% methanol and stored at -20°C until needed. Embryos
were incubated in Dent's bleach (4:1:1 Methanol:DMSO:Hydrogen peroxide) for 2 hours in the
dark at room temperature, rinsed in 100% methanol for 15 minutes, and rehydrated through a
decreasing methanol series from 100% methanol to PBT (0.1% Triton in 1X PBS). Embryos were
blocked in 2% goat serum + 2% Bovine Serum Albumin for 2 hours at room temperature before
primary antibody (GFP, 1:500, Invitrogen #A6455) was added and the embryos were incubated
overnight at 4°C with rocking. Embryos were washed three times for 5 minutes in PBT and then
6 times for 1 hour at room temperature with rocking. Embryos were incubated in secondary
antibody (Alexa Fluor 488 Goat anti-Rabbit, 1:500, Invitrogen #11008; DAPI, 1:1200, Sigma-
Aldrich #D9564) at 4°C in the dark overnight with rocking, then washed 3 times for 5 minutes in
PBT and 6 times for 1 hour at room temperature with rocking. Embryos were then mounted in
VECTASHIELD® Antifade Mounting Medium with DAPI (Vector Laboratories #H-1200) for
imaging.
540
Cryosection

Embryos were dissected and fixed in 4% PFA (in PBS) overnight. Embryos were then
transferred to 1X PBS and rocked for 10 minutes at 4°C, before being incubated in 30% sucrose
in 1X PBS, at 4°C for 30 minutes-5 hours with rocking (or until embryos were equilibrated as
indicated by their sinking to the bottom of the tube). Embryos were embedded in Tissue-Tek
O.C.T. Compound (VWR #25608-930), sectioned transversely at 10um thickness on a NX70
Cryostar cryostat, permeabilized 3 times for 5 minutes in PBT, and blocked in 2% goat serum +
2% Bovine Serum Albumin for 2 hours at room temperature. Sections were then incubated in
primary antibody overnight at 4°C and the following primary antibodies were used: Sox10
(1:500, Abcam #ab155279), phospho-histone H3 (1:500, Millipore #05-806), GFP (1:500,
Invitrogen #G10362). Sections were then washed 3 times for 5 minutes in PBT at room
temperature, before diluted secondary antibody and/or conjugated primary antibody solutions
were added and the sections were incubated for 2 hours in the dark at room temperature.
Sections were washed 3 times for 5 minutes in PBT before mounting in VECTASHIELD®.
Secondary antibodies included Alexa Fluor 488 Goat anti-Rabbit (1:500, Invitrogen, #11008),
546 Goat anti-Mouse (1:500, Invitrogen, #A21045), 647 Goat anti-Rat (1:500, Invitrogen,
#A21267), 647 Donkey anti-Mouse (1:500, Invitrogen, #A32787), DAPI (1:1200, Sigma-Aldrich,
#D9564).
559
TUNEL staining

Cryosections were washed 3 times for 5 minutes in PBT and then permeabilized in 0.1%
sodium citrate/PBT for 5 minutes. Cryosections were then washed 3 times for 5 minutes in PBT
before being incubated with 1:10 TUNEL enzyme buffer (Roche #12156792910) at 37°C in the
dark for 2 hours. Sections were washed 3 times for 5 minutes in PBS prior to counterstaining
with DAPI and/or mounting in VECTASHIELD®.
566
EdU labeling

EdU staining was performed on cultured whole mouse embryos. Briefly, complete media
containing 50% DMEM-F12-Glutamax, 50% rat serum, and 1X penicillin/streptomycin was pre-
warmed at 37°C in roller culture bottles in a 5% CO2, 5% O2 and 90% N2 atmosphere (Muñoz
and Trainor 2019; Sakai and Trainor 2014). After E8.5 CD1 embryos were dissected in Tyrode's
buffer with an intact yolk sac, and equilibrated in culture media for 30 minutes, EdU (Invitrogen
#C10638) was added to the media according to the manufacturer's protocol at a working
concentration of 500uM. DMSO was added to control embryos. Embryos were then incubated
for 15 minutes, after which they were briefly rinsed in warm equilibrated culture media
followed by Tyrode's buffer before fixing in 4% PFA (in PBS) overnight at 4°C. Fixed embryos
were embedded and sectioned at 10um, and then stained following the manufacturer's
instructions, with fluorescent immunostaining performed as needed after EdU staining.
579

Single cell RNA sequencing

Tissue collection, cell processing and sequencing, and data processing were performed
as previously described (Falcon et al. 2022). Briefly, 6 *Mef2c-F10N-LacZ* (Aoto et al. 2015) and 6
*Wnt1-Cre;*RosaeYFP (Chai et al. 2000) E8.5 mouse embryos were collected, dissociated into
single cells, and 12,000-15,000 cells per sample were loaded on a Chromium Single Cell
Controller (10x Genomics). Libraries were prepared using the Chromium Next GEM Single Cell 3'
Library & Gel Bead Kit v3.1 (10x Genomics), quality control checked, and then pooled at equal
molar concentrations and sequenced on an Illumina NovaSeq 6000 S1 flow cell. Raw
sequencing data was processed using Cell Ranger (v3.0.0, 10x Genomics) and after
mitochondria and other feature thresholding, the final dataset used for analysis consisted of
21,190 cells (12,498 cells for Wnt1-Cre;RosaeYFP and 8,692 for Mef2c-F10N-LacZ and 29,041
genes, and is available at the Gene Expression Omnibus (accession no. GSE168351). R (v3.6.1)
was used for downstream analysis. The Seurat package (v3.1.5.9003) (Stuart et al. 2019) was
used to normalize data via the SCTransform method (Hafemeister and Satija 2019). For
clustering, 3000 highly variable genes were selected, and the first 46 principal components
based on those genes were used to identify 6 initial clusters at a resolution of 0.05 using the
shared nearest neighbor method. Cranial NCC were identified as one of the 6 initial clusters
based on the expression of tissue specific marker genes. The cranial NCC cluster was then
subdivided at resolution=0.26 into 5 subclusters, three of which were characterized as the early
migratory NCC based on NCC development gene expression. Early migratory NCC were further
subdivided into 15 subclusters at resolution=2.0 to explore the presence of EMT intermediate
NCC.
602

Pseudotime trajectory analysis

Trajectory analysis was performed using Monocle3 (0.2.2) with closed loop set to true and use partition set to false on cranial neural crest cells, which were subset from the single cell data set after doublet identification with DoubletFinder (2.0.2). The resulting pseudotime estimates were added back to the Seurat metadata table for visualization and the trajectory was plotted on the original clusters identified in the Seurat analysis.

SABER-FISH staining and imaging

All oligo pools, concatemer hairpins and fluorophore probes were designed and ordered through IDT using stringent settings as previously described (Kishi et al. 2019). Only fluorophores 488, 594 and 647 were used to prevent overlap in spectral emission and channel bleed-through. SABER-FISH probes for each gene were made following the PER concatamerization protocol. The probes were allowed to elongate for 2 hours at 37°C before heat inactivation of the polymerase. A sample of the probes was run on a gel to confirm elongation length and ensure that no secondary products had formed in the process. Probes were then cleaned using the Qiagen PCR Purification kit and the final concentration of the generated probes were measured by Nanodrop.

CD1 embryos were collected at the 5-6ss and fixed overnight in 4% PFA (in PBS) at 4°C. Embryos were then washed 5 times in PBS with DEPC-PBTW (1X DEPC-PBS + 0.1% Tween-20) and transferred into 30% sucrose PBTW and rocked overnight at 4°C. Next, embryos were embedded in OCT, frozen and cryosectioned at 20μm thickness. Sections were placed on Histogrip treated slides and warmed to encourage adherence, washed 3 times for 5 minutes in PBTW, incubated in Hybridization wash buffer and allowed to equilibrate to 37°C (the lowest melting temperature limit out of our probe set). Sections were then incubated with pre-equilibrated probes overnight (roughly 16 hours) at 37°C. The next day the sections were washed with Hybridization wash buffer, 2X SSCT (2×SSC + 0.1% Tween-20) and PBTW and then the first set of fluorophores were applied to the sections [set one: Sox10, Wnt1 and Sp5; set two: Pak3 and Dlc1] and left to hybridize at 37°C for 30 minutes. Following fluorophore probe hybridization, sections were washed in PBTW with 1:1000 DAPI for 10 minutes followed by PBTW 2 times for 5 minutes. The slides were then mounted in VECTASHIELD® containing DAPI with a 1.5 glass coverslip.

Sections were imaged with a Nikon CSU-W1 inverted spinning disk equipped with a sCMOS camera. Each laser was set to 400μs acquisition speed and a z-stack of 32 slices (1μm per slice) was acquired. Following imaging of the first set of probes, the coverglass was gently removed from each slide. The first set of fluorophore probes were removed by washing 3 times for 5 minutes in PBTW washes, followed by 3 washes with Displacement buffer and another 3 washes in PBTW. Slides were remounted with VECTASHIELD®, covered with a coverslip and imaged to ensure all previous fluorescent probes had been removed. After imaging, the coverslip was removed again, and the sections were rinsed in PBTW. 3 times for 5 minutes. The second fluorophore probes were then applied to the sections and allowed to hybridize for 30 minutes at 37 °C. Slides were then washed 3 times for 5minutes in PBTW and mounted a final

644   time in VECTASHIELD® containing DAPI. The second set of probes were imaged using the exact
645   same parameters as the first.

647   **SABER-FISH image processing**

648   All image processing was performed in FIJI/ImageJ (Schindelin et al. 2012) and plugin
649   source code can be accessed through https://github.com/jayunruh/Jay_Plugins. Plugins can be
650   used by following the Stowers Fiji update site. Convenience macros that combine these plugin
651   functionalities are included in the supplemental materials.
652   After acquisition, images were scaled by 0.5 with averaging (i.e. binning). Background
subtraction was run on the scaled images using the "roi average subtract jru v1" plugin. This
plugin takes the average intensity signal in a selected region of interest or chosen area of
background in the image and removes that average across the image. The two series of z-stack
images acquired for each set of probes were then aligned and combined through a
"registration_macro.ijm". In summary, this macro registers the two z-stacks according to DAPI
signal from a selected representative z-slice. Alignment is achieved through using an
implementation of the TurboReg tool set to a rigid body transformation (Thevenaz et al. 1998).
660   Once the two sets of probes have been aligned, the signals are measured and tracked
with the macro "all_combined_aftersubreg.ijm". Images were first copied, and then sum
projected in z before undergoing a Gaussian blur with sigma value 4. Nuclei are then identified
in the DAPI channel by a maximum finding approach with a minimum spot distance of 70 pixels
and a threshold of 10% of the maximum intensity (Varberg et al. 2022). Those spots provided a
locational value for later mapping of transcript signals in two dimensions. Individual transcripts
are found in three dimensions in the original combined images with a Gaussian blur with sigma
value 1 and a rolling ball background subtraction with a radius of 10 pixels. Because the signal is
very punctate and slides can accumulate auto-fluorescent debris over the course of staining, we
next removed the 20 brightest spots in 3D with a spheroid of xy diameter 15 and z diameter 5
to ensure we were evaluating true signal. Spots were found using the same maximum finding
approach as above but in 3 dimensions with a minimum separation of 12 pixels in xy and 4
slices in z and a threshold at 7% of the maximum intensity in each channel. The positions of
those found maxima were then sum projected in z to estimate the number of transcripts in the
vicinity of each nuclear maximum (see above).
675   An image showing the number of transcripts per cell was generated using
"make_nuclear_image.py" macro based on the nuclei and transcript locations identified in the
previous step. To generate the polyline kymograph showcasing spatial expression through the
tissue, a zoomed in region of the transcript mapped images was generated around the dorsal
neural fold tips. A line of 100 pixel width was drawn starting from the middle of the
neuroepithelium towards the dorsal most tip of the neural fold and then ventrally into the
underlying mesoderm and migratory neural crest cell population. The polyline kymograph was
generate based on this line using the plugin "polyline kymograph jru v1".

RNA in situ hybridization

Dlc1, Pak3 and *Sp5 in situ* plasmids were previously published (Dunty et al. 2014;
Piccand et al. 2014). RNA in situ hybridization was performed as follows: Embryos were
rehydrated through a descending methanol series from 100% methanol to DEPC-PBTW, then
washed 2 times for 5 minutes in DEPC-PBTW. Embryos were bleached in 6% hydrogen peroxide
in the dark for 15 minutes with rocking and then incubated in 10µg/ml Proteinase K in DEPC-
PBTW for 4-5 minutes at room temperature, without rocking. Embryos were wash with 2 mg/ml
glycine in DEPC-PBTW for 5 minutes, then 2 times for 5 minutes in DEPC-PBTW before being
refixed in 4% DEPC-PFA + 0.2% glutaraldehyde for 20 minutes. Embryos were then washed 3
times for 5 minutes in DEPC-PBTW, rinsed in prewarmed (68°C) hybridization buffer (50%
formamide, 5X SSC pH 4.5, 0.05% EDTA pH 8, 0.2% Tween20, 0.1% CHAPS, 20mg/ml Boehringer
blocking powder, 1 mg/ml torula RNA, 0.05 mg/ml heparin), and incubated in hybridization
buffer with rocking for 1 hour at 68°C. Embryos were then incubated in digoxygenin-labeled
riboprobes (~2 ng/ul) in hybridization buffer overnight at 68°C. Day 2: Embryos were washed 2
times for 30 minutes with hybridization buffer at 68°C. followed by prewarmed Solution I (50%
formamide, 1X SSC pH 4.5, 0.1% Tween20), 3 times for 30 minutes at 65°C. Embryos were then
washed in 50% Solution I/50% MABT (1X MAB – Maleic acid, 0.1% Tween-20) for 30 minutes at
room temperature, followed by MABT, 3 times for 5 minutes. Embryos were then blocked in
MABT + 2% Boehringer blocking powder for 1 hour at room temperature, followed by MABT +
2% Boehringer blocking powder + 20% heat-treated goat serum for 2 hours. Embryos were then
incubated overnight in anti-digoxygenin-AP diluted 1:2000 in MABT + 2% Boehringer blocking
powder + 20% heat-treated goat serum, at 4°C. The embryos were then rinsed in MABT,
washed 2 times for 15 minutes in MABT, followed by 5-8 further washes in MABT for 1-1.5
hours each, and an overnight wash in MABT at room temperature. The embryos were then
washed 3 times for 10 minutes in NTMT (100mM NaCl, 100mM Tris pH 9.5, 50mM MgCl2, 0.1%
Tween-20), and incubated in NTMT + BCIP/NBT at room temperature in the dark. Color
development at room temperature was allowed to continue until the desired darkness of the
substrate was achieved. The color reaction was stopped by washing in PBTW, after which the
embryos were stored. long-term in 4% PFA (in PBS)/0.1% glutaraldehyde at 4 ºC. Stained
embryos were imaged, and sectioned transversely at 10um thickness prior to imaging.
714
Aphidicolin treatment

E7.5-E8.0 CD1 embryos were dissected in Tyrode's buffer with their yolk sac intact and
incubated in pre-warmed culture media for 30 minutes-1 hour as previously described (Muñoz
and Trainor 2019; Sakai and Trainor 2014, 2016). After 30 minutes equilibration, 0.5ug/ml
aphidicolin (Sigma-Aldrich, #A0781) was added to the culture media to inhibit S phase of the
cell cycle. The same quantity of DMSO was added to the control embryos. Following 12-13
hours of roller culture, EdU as added during the final 15 minutes. Embryos were then fixed in
4% PFA (in PBS) overnight at 4°C. overnight. Fixed embryos were embedded and sectioned at
10um and stained using the EdU kit following the manufacturer's instructions. Sox10 and
phospho-histone H3 fluorescent immunostaining was performed after EdU staining as
described above. TUNEL staining (described above) was performed after fluorescent
immunostaining if applicable.
727

ShRNA-based lentiviral plasmids and lentivirus production

A shRNA plasmid clone set targeting *Dlc1* was obtained from GeneCopoeia
(MSH100727-LVRU6MP). Each set contains 3 shRNA expression constructs and 1 scrambled
shRNA control. Each shRNA hairpin consists of a 7 base loop and 19-29 base stem optimized for
specific gene sequences as detailed by the manufacturer. Glycerol stocks of shRNA-based
lentiviral plasmids were cultured in LB buffer with 100 µg/ml of ampicillin (Amresco # 0339).
Plasmids were purified using a HiSpeed Plasmid midi kit (Qiagen #12643).
A total of $4e^6$-$5e^6$ of 293T cells were seeded in one 10cm plate with 12ml of media
without antibiotics the day before transfection. On the following day, when the cells reached
70-80% confluence, transfection was performed as follows: shRNA-based lentiviral plasmids
(7ug), Pax2 packaging plasmid (7ug) and VSVG envelop plasmid (1ug) were mixed with 45ul of
Fugene HD transfection reagent (Promega, cat#E5911) in 1.5ml of Opti-MEM (Gibco,
cat#31985070). The mixture was incubated at room temperature for 15 minutes and then
added dropwise to the plate of 293T cells. Virus-containing old culture media was harvested
after 48 hours and 72 hours upon transfection. We then mixed the virus-containing media and
added HEPES to reach 10mM final concentration. The media was spun down at 500g for 5
minutes at 4°C to remove cell debris and the supernatant was then collected and filtered
through 0.45um filter into a falcon tube. We added 1 volume of 4x lentivirus concentrator
solution (40% PEG-8000 in 1.2M NaCl) to 3 volumes of the filtered virus, mixed well and placed
the tube on a shaker (60rpm/minute) at 4°C for overnight. The filtered virus was centrifuged at
1600g for 1hour at 4°C and the supernatant was carefully removed without disturbing the
pellet. The pellet was then thoroughly resuspended with 1ml of cold PBS by gently pipetting up
and down. The solution was transferred to 1.5ml tube and placed at room temperature for 10
minutes. We gently pipetted the virus again about 20 times and spun it down in a
microcentrifuge at full speed for 3 minutes to pellet the protein debris. The supernatant was
then aliquoted and stored at -80°C for future use.

Lentivirus injection

E7.5-E8.0 CD1 embryos were dissected in Tyrode's buffer with an intact yolk sac and
equilibrated in pre-warmed culture media for 30 minutes as previously described (Muñoz and
Trainor 2019; Sakai and Trainor 2014, 2016). Viruses in 5ul aliquots were thawed and kept on
ice, then pipetted onto a piece of parafilm in a small dish with the lid on to prevent
evaporation. Using the Eppendorf CellTram Vario system, 1-2ul of each virus was injected into
the amnionic cavity of each embryo (or until the amnion had visibly expanded) after which
time, the embryos were returned to roller culture. After 24 hours of incubation, embryos were
rinsed in PBS and fixed in 4% PFA (in PBS) at 4°C overnight. Fixed embryos were imaged,
embedded, and sectioned at 10um thickness before fluorescent immunostaining with Sox10.

RNA isolation, cDNA preparation and quantitative reverse transcription-PCR (qRT-PCR)

Individual E8.5 mouse embryo head and tail tissues post culturing were collected into
1.5mL tubes and flash froze on dry ice with minimal DEPC-Tyrode's buffer remained in the tube.
RNA was extracted using the Qiagen miRNeasy Micro Kit (Qiagen #217084) with on-column

DNase treatment. RNA concentration was determined by Nanodrop. The SuperScript™ III First-
Strand Synthesis System (Invitrogen #18080051) was used to synthesize cDNA for qRT-PCR with
random hexamer primers, and qRT-PCR was performed on an ABI7000 (Thermo QuantStudio 7)
using Perfecta SYBR Green (Quantbio #95072-250). Primers are listed in Supplemental Table 2.
No template and no reverse transcription controls were run as negative controls. ΔΔCt method
was used to calculate fold change. One-way ANOVA was used for statistical analysis and
significance was determined based on $p < 0.05$.
777
β-galactosidase staining

E8.5 *Mef2c-F10N-LacZ* embryos were collected and fixed in 2% formalin, 0.2%
glutaraldehyde in 1X PBS for 15-20 minutes. Embryos were rinsed with PBS and stained
according to manufacturer's protocol (Millipore #BG-6-B, #BG-7-B, #BG-8-C). Embryos were
then fixed again in 4% PFA (in PBS) at 4°C with rocking overnight followed by washing in PBS for
whole embryo brightfield imaging.
784
Fluorescent imaging

Fluorescently stained section images were captured on an upright Zeiss LSM-700 laser
scanning confocal microscope using 405 nm, 488 nm, 555 nm and 639 nm excitation lasers.
Emissions filters used to acquire images were Far-red: LP 640nm, Red: BP 505-600 nm, GFP: BP
490-555 nm, DAPI: SP 490 nm. Images were acquired with a Zeiss Fluar 10x objective lens. For
each specimen, a z-stack of images was collected and processed as a maximum intensity
projection.
All the images used for quantification were acquired with an Orca Flash 4.0 sCMOS
100fps at full resolution on a Nikon Eclipse Ti2 microscope equipped with a Yokagawa CSU W1
10,000 rpm Spinning Disk Confocal system. The spinning disk confocal is equipped with a quad
filter for excitation with 405/488/561/640. Emissions filters used to acquire images were Far-
red: 669-741nm, Red: 579-631 nm, GFP: 507-543 nm, DAPI: 430-480 nm. A Nikon Plan
Apochromat Lambda LWD 40x objective was used to acquire the images with 50-100 ms
exposure times.
799
Image processing

All analyses of fluorescent intensity were performed using Fiji and custom-written
ImageJ Macro and python notebooks. Prior to the analyses, raw images were processed by
subtracting background and were then projected for the max intensity to form single multiple-
color images. Individual cells were segmented based on DAPI channel with a pre-trained
cellpose model (Stringer et al. 2021), and then the mean intensity of the individual channel was
measured by FIJI. The segmented cells were classified as either positive or negative based on
their intensity in the corresponding channel. For images from the shRNA lentivirus injection
experiments, due to the large variety of background signals, we manually labeled positive cells
and then trained a cellpose model based on the manually labeled cells. Positive cells were
classified based on the trained model, and their coordinates were recorded and saved to
ImageJ ROI files for future verification.

For the cell cycle staining analysis, cells in the most dorsal lateral domain of the cranial and trunk neural plate were selected and saved to ImageJ ROI files. In cranial neural plate border cells, the following quantifications were performed: EdU+%=the percentage of EdU positive cells within eYFP positive delaminating premigratory NCC; pHH3+%= the percentage of pHH3 positive cells within eYFP positive delaminating premigratory NCC; EdU+pHH3+%= the percentage of EdU and pHH3 double positive cells within eYFP positive delaminating premigratory NCC; EdU-pHH3-%= the percentage of EdU and pHH3 double negative cells within eYFP positive delaminating premigratory NCC. In trunk neural plate border cells, the following quantifications were performed: EdU+%=the percentage of EdU positive cells within DAPI positive neural plate border cells at the trunk axial level; pHH3+%= the percentage of pHH3 positive cells within DAPI positive trunk neural plate border cells; EdU+pHH3+%= the percentage of EdU and pHH3 double positive cells within DAPI positive trunk neural plate border cells; EdU-pHH3-%= the percentage of EdU and pHH3 double negative cells within DAPI positive trunk neural plate border cells. For the Aphidicolin treatment experiments, neuroepithelial cells and Sox10 positive migratory NCC were selected and saved to ImageJ ROI files. The ratio of Sox10 positive migratory NCC over DAPI positive neural plate/neuroepithelial cells was calculated. For the shRNA lentivirus injection experiments, Sox10 positive migratory NCC were selected and saved to ImageJ ROI files. The number of Sox10 positive cells was quantified.

Brightfield imaging

Embryos were imaged on a Leica MZ16 microscope equipped with a Nikon DS-Ri1 camera and NIS Elements imaging software. Manual Z stacks were taken and then assembled using Helicon Focus software. Sections from embryos stained by in situ hybridization were imaged on a ZEISS Axio Vert and stitched using ImageJ if needed.

Acknowledgements

We thank members of the Trainor laboratory for their insights and suggestions on the project. We also acknowledge our amazing animal technician, Melissa Childers, and the Laboratory Animal Services facility at Stowers Institute for Medical Research for animal husbandry and care of the mice used in this work. The *Dlc1*, *Sp5* and *Pak3* ISH plasmids were generously provided by Dr. Marian Durkin, Dr. Gerard Gradwohl, and Dr. Terry Yamaguchi, respectively. We also thank Mark Miller, who created the beautiful illustration for the experimental workflow included in Figure 1. Funding for this research was provided by the Stowers Institute for Medical Research (P.A.T) and the National Institute for Dental and Craniofacial Research F31 DE032256 (E.L.M.). The scRNA-seq data was generated through the University of Kansas Medical Center Genomics Core. Funding for the Genomics Core services was provided by the Kansas Intellectual and Developmental Disabilities Research Center (NIH U54 HD 090216), the Molecular Regulation of Cell Development and Differentiation – COBRE (P30 GM122731-03)

852   and the NIH S10 High-End Instrumentation Grant (NIH S10OD021743) at the University of
853   Kansas Medical Center, Kansas City, KS 66160.

References

Achilleos A, Trainor PA. 2015. Mouse models of rare craniofacial disorders. Review of. *Curr Top Dev Biol*
115:413-458. doi: 10.1016/bs.ctdb.2015.07.011
Ahlstrom JD, Erickson CA. 2009. The neural crest epithelial-mesenchymal transition in 4d: A 'tail' of
multiple non-obligatory cellular mechanisms. Review of. *Development (Cambridge, England)*
136:1801-1812. doi: 10.1242/dev.034785
Aoto K, Sandell LL, Butler Tjaden NE, Yuen KC, Watt KEN, Black BL, Durnin M, Trainor PA. 2015. Mef2c-
f10n enhancer driven β-galactosidase (lacz) and cre recombinase mice facilitate analyses of gene
function and lineage fate in neural crest cells. Review of. *Developmental Biology* **402**:3-16. doi:
10.1016/j.ydbio.2015.02.022
Aybar MJ, Nieto MA, Mayor R. 2003. Snail precedes slug in the genetic cascade required for the
specification and migration of the xenopus neural crest. Review of. *Development (Cambridge,*
England) **130**:483-494. doi: 10.1242/dev.00238
Berndt JD, Clay MR, Langenberg T, Halloran MC. 2008. Rho-kinase and myosin ii affect dynamic neural
crest cell behaviors during epithelial to mesenchymal transition in vivo. Review of.
Developmental Biology **324**:236-244. doi: 10.1016/j.ydbio.2008.09.013
Bhatt S, Diaz R, Trainor PA. 2013. Signals and switches in mammalian neural crest cell differentiation.
Review of. *Cold Spring Harbor Perspectives in Biology* **5**:a008326. doi:
10.1101/cshperspect.a008326
Bulfone A, Kim H-J, Puelles L, Porteus MH, Grippo JF, Rubenstein JLR. 1993. The mouse dlx-2 (tes-1) gene
is expressed in spatially restricted domains of the forebrain, face and limbs in midgestation
mouse embryos. Review of. *Mechanisms of Development* **40**:129-140. doi: 10.1016/0925-
4773(93)90071-5
Burstyn-Cohen T, Kalcheim C. 2002. Association between the cell cycle and neural crest delamination
through specific regulation of g1/s transition. Review of. *Developmental Cell* **3**:383-395. doi:
10.1016/s1534-5807(02)00221-6
Burstyn-Cohen T, Stanleigh J, Sela-Donenfeld D, Kalcheim C. 2004. Canonical wnt activity regulates trunk
neural crest delamination linking bmp/noggin signaling with g1/s transition. Review of.
Development **131**:5327-5339. doi: 10.1242/dev.01424
Chai Y, Jiang X, Ito Y, Bringas P, Han J, Rowitch DH, Soriano P, McMahon AP, Sucov HM. 2000. Fate of the
mammalian cranial neural crest during tooth and mandibular morphogenesis. Review of.
Development (Cambridge, England) **127**:1671-1679. doi: 10.1242/dev.127.8.1671
Cheung M, Briscoe J. 2003. Neural crest development is regulated by the transcription factor sox9.
Review of. *Development* **130**:5681-5693. doi: 10.1242/dev.00808
Cheung M, Chaboissier M-C, Mynett A, Hirst E, Schedl A, Briscoe J. 2005. The transcriptional control of
trunk neural crest induction, survival, and delamination. Review of. *Developmental Cell* **8**:179-
192. doi: 10.1016/j.devcel.2004.12.010
Clay MR, Halloran MC. 2013. Rho activation is apically restricted by arhgap1 in neural crest cells and
drives epithelial-to-mesenchymal transition. Review of. *Development* **140**:3198-3209. doi:
10.1242/dev.095448
Dash S, Trainor PA. 2020. The development, patterning and evolution of neural crest cell differentiation
into cartilage and bone. Review of. *Bone* **137**:115409. doi: 10.1016/j.bone.2020.115409
Dong J, Hu Y, Fan X, Wu X, Mao Y, Hu B, Guo H, Wen L, Tang F. 2018. Single-cell rna-seq analysis unveils a
prevalent epithelial/mesenchymal hybrid state during mouse organogenesis. Review of.
Genome Biology **19**:31. doi: 10.1186/s13059-018-1416-2

Dongre A, Weinberg RA. 2019. New insights into the mechanisms of epithelial-mesenchymal transition
and implications for cancer. Review of. *Nature Reviews. Molecular Cell Biology* **20**:69-84. doi:
10.1038/s41580-018-0080-4
Dottori M, Gross MK, Labosky P, Goulding M. 2001. The winged-helix transcription factor foxd3
suppresses interneuron differentiation and promotes neural crest cell fate. Review of.
Development **128**:4127-4138. doi: 10.1242/dev.128.21.4127
Dunty WC, Jr., Kennedy MW, Chalamalasetty RB, Campbell K, Yamaguchi TP. 2014. Transcriptional
profiling of wnt3a mutants identifies sp transcription factors as essential effectors of the
wnt/beta-catenin pathway in neuromesodermal stem cells. Review of. *PLOS ONE* **9**:e87018. doi:
10.1371/journal.pone.0087018
Durkin ME, Avner MR, Huh C-G, Yuan B-Z, Thorgeirsson SS, Popescu NC. 2005. Dlc-1, a rho gtpase-
activating protein with tumor suppressor function, is essential for embryonic development.
Review of. *FEBS Letters* **579**:1191-1196. doi: 10.1016/j.febslet.2004.12.090
Echelard Y, Vassileva G, McMahon AP. 1994. Cis-acting regulatory sequences governing wnt-1
expression in the developing mouse cns. Review of. *Development* **120**:2213-2224. doi:
10.1242/dev.120.8.2213
Falcon KT, Watt KEN, Dash S, Zhao R, Sakai D, Moore EL, Fitriasari S, Childers M, Sardiu ME, Swanson S,
Tsuchiya D, Unruh J, Bugarinovic G, Li L, Shiang R, Achilleos A, Dixon J, Dixon MJ, Trainor PA.
2022. Dynamic regulation and requirement for ribosomal rna transcription during mammalian
development. Review of. *Proceedings of the National Academy of Sciences* **119**:e2116974119.
doi: 10.1073/pnas.2116974119
Golding JP, Trainor P, Krumlauf R, Gassmann M. 2000. Defects in pathfinding by cranial neural crest cells
in mice lacking the neuregulin receptor erbb4. Review of. *Nat Cell Biol* **2**:103-109. doi:
10.1038/35000058
Gonzalez VD, Samusik N, Chen TJ, Savig ES, Aghaeepour N, Quigley DA, Huang Y-W, Giangarrà V,
Borowsky AD, Hubbard NE, Chen S-Y, Han G, Ashworth A, Kipps TJ, Berek JS, Nolan GP, Fantl WJ.
2018. Commonly occurring cell subsets in high-grade serous ovarian tumors identified by single-
cell mass cytometry. Review of. *Cell Reports* **22**:1875-1888. doi: 10.1016/j.celrep.2018.01.053
Hafemeister C, Satija R. 2019. Normalization and variance stabilization of single-cell rna-seq data using
regularized negative binomial regression. Review of. *Genome Biology* **20**:296. doi:
10.1186/s13059-019-1874-1
Hari L, Miescher I, Shakhova O, Suter U, Chin L, Taketo M, Richardson WD, Kessaris N, Sommer L. 2012.
Temporal control of neural crest lineage generation by wnt/β-catenin signaling. Review of.
Development (Cambridge, England) **139**:2107-2117. doi: 10.1242/dev.073064
Hay ED. 1995. An overview of epithelio-mesenchymal transformation. Review of. *Acta Anatomica* **154**:8-
20. doi: 10.1159/000147748
Huang RY-J, Wong MK, Tan TZ, Kuay KT, Ng AHC, Chung VY, Chu Y-S, Matsumura N, Lai H-C, Lee YF, Sim
W-J, Chai C, Pietschmann E, Mori S, Low JJH, Choolani M, Thiery JP. 2013. An emt spectrum
defines an anoikis-resistant and spheroidogenic intermediate mesenchymal state that is
sensitive to e-cadherin restoration by a src-kinase inhibitor, saracatinib (azd0530). Review of.
Cell Death & Disease **4**:e915. doi: 10.1038/cddis.2013.442
Iyyanar PPR, Wu Z, Lan Y, Hu Y-C, Jiang R. 2022. Alx1 deficient mice recapitulate craniofacial phenotype
and reveal developmental basis of alx1-related frontonasal dysplasia. Review of. *Frontiers in Cell
and Developmental Biology* **10**.
Jiang X, Rowitch DH, Soriano P, McMahon AP, Sucov HM. 2000. Fate of the mammalian cardiac neural
crest. Review of. *Development* **127**:1607-1616. doi: 10.1242/dev.127.8.1607
Karacosta LG, Anchang B, Ignatiadis N, Kimmey SC, Benson JA, Shrager JB, Tibshirani R, Bendall SC,
Plevritis SK. 2019. Mapping lung cancer epithelial-mesenchymal transition states and

trajectories with single-cell resolution. Review of. *Nature Communications* **10**:5587. doi:
10.1038/s41467-019-13441-6
Kim TY, Healy KD, Der CJ, Sciaky N, Bang Y-J, Juliano RL. 2008. Effects of structure of rho gtpase-
activating protein dlc-1 on cell morphology and migration*. Review of. *Journal of Biological*
Chemistry **283**:32762-32770. doi: 10.1074/jbc.M800617200
Kishi JY, Lapan SW, Beliveau BJ, West ER, Zhu A, Sasaki HM, Saka SK, Wang Y, Cepko CL, Yin P. 2019.
Saber amplifies fish: Enhanced multiplexed imaging of rna and DNA in cells and tissues. Review
of. *Nature Methods* **16**:533-544. doi: 10.1038/s41592-019-0404-0
Kobayashi GS, Musso CM, Moreira DdP, Pontillo-Guimarães G, Hsia GSP, Caires-Júnior LC, Goulart E,
Passos-Bueno MR. 2020. Recapitulation of neural crest specification and emt via induction from
neural plate border-like cells. Review of. *Stem Cell Reports* **15**:776-788. doi:
10.1016/j.stemcr.2020.07.023
Kumar SU, Kumar DT, Siva R, Doss CGP, Zayed H. 2019. Integrative bioinformatics approaches to map
potential novel genes and pathways involved in ovarian cancer. Review of. *Frontiers in*
Bioengineering and Biotechnology **7**:391. doi: 10.3389/fbioe.2019.00391
Lee RTH, Nagai H, Nakaya Y, Sheng G, Trainor PA, Weston JA, Thiery JP. 2013. Cell delamination in the
mesencephalic neural fold and its implication for the origin of ectomesenchyme. Review of.
Development (Cambridge, England) **140**:4890-4902. doi: 10.1242/dev.094680
Li J, Chen F, Epstein JA. 2000. Neural crest expression of cre recombinase directed by the proximal pax3
promoter in transgenic mice. Review of. *genesis* **26**:162-164. doi: 10.1002/(SICI)1526-
968X(200002)26:2<162::AID-GENE21>3.0.CO;2-R
Lièvre CSL, Douarin NML. 1975. Mesenchymal derivatives of the neural crest: Analysis of chimaeric quail
and chick embryos. Review of. *Development* **34**:125-154. doi: 10.1242/dev.34.1.125
Liu JA, Rao Y, Cheung MPL, Hui M-N, Wu M-H, Chan L-K, Ng IO-L, Niu B, Cheah KSE, Sharma R, Hodgson L,
Cheung M. 2017. Asymmetric localization of dlc1 defines avian trunk neural crest polarity for
directional delamination and migration. Review of. *Nature Communications* **8**:1185. doi:
10.1038/s41467-017-01107-0
Lumb R, Wiszniak S, Kabbara S, Scherer M, Harvey N, Schwarz Q. 2014. Neuropilins define distinct
populations of neural crest cells. Review of. *Neural Development* **9**:24. doi: 10.1186/1749-8104-
9-24
Mandalos N, Rhinn M, Granchi Z, Karampelas I, Mitsiadis T, Economides AN, Dollé P, Remboutsika E.
2014. Sox2 acts as a rheostat of epithelial to mesenchymal transition during neural crest
development. Review of. *Frontiers in Physiology* **5**:345. doi: 10.3389/fphys.2014.00345
Matsuuchi L, Naus CC. 2013. Gap junction proteins on the move: Connexins, the cytoskeleton and
migration. Review of. *Biochimica et Biophysica Acta (BBA) - Biomembranes* **1828**:94-108. doi:
10.1016/j.bbamem.2012.05.014
McKeown SJ, Lee VM, Bronner-Fraser M, Newgreen DF, Farlie PG. 2005. Sox10 overexpression induces
neural crest-like cells from all dorsoventral levels of the neural tube but inhibits differentiation.
Review of. *Developmental Dynamics* **233**:430-444. doi: 10.1002/dvdy.20341
McKinney MC, Fukatsu K, Morrison J, McLennan R, Bronner ME, Kulesa PM. 2013. Evidence for dynamic
rearrangements but lack of fate or position restrictions in premigratory avian trunk neural crest.
Review of. *Development (Cambridge, England)* **140**:820-830. doi: 10.1242/dev.083725
Muñoz WA, Trainor PA. 2019. Mouse embryo culture for the study of neural crest cells. Review of.
Methods in Molecular Biology (Clifton, N.J.) **1976**:107-119. doi: 10.1007/978-1-4939-9412-0_9
Murdoch B, DelConte C, García-Castro MI. 2012. Pax7 lineage contributions to the mammalian neural
crest. Review of. *PLOS ONE* **7**:e41089. doi: 10.1371/journal.pone.0041089

Parr BA, Shea MJ, Vassileva G, McMahon AP. 1993. Mouse wnt genes exhibit discrete domains of
expression in the early embryonic cns and limb buds. Review of. *Development* **119**:247-261. doi:
10.1242/dev.119.1.247
Pastushenko I, Brisebarre A, Sifrim A, Fioramonti M, Revenco T, Boumahdi S, Van Keymeulen A, Brown
D, Moers V, Lemaire S, De Clercq S, Minguijón E, Balsat C, Sokolow Y, Dubois C, De Cock F,
Scozzaro S, Sopena F, Lanas A, D'Haene N, Salmon I, Marine J-C, Voet T, Sotiropoulou PA,
Blanpain C. 2018. Identification of the tumour transition states occurring during emt. Review of.
Nature **556**:463-468. doi: 10.1038/s41586-018-0040-3
Piccand J, Meunier A, Merle C, Jia Z, Barnier J-V, Gradwohl G. 2014. Pak3 promotes cell cycle exit and
differentiation of β-cells in the embryonic pancreas and is necessary to maintain glucose
homeostasis in adult mice. Review of. *Diabetes* **63**:203-215. doi: 10.2337/db13-0384
Popoff MR, Geny B. 2009. Multifaceted role of rho, rac, cdc42 and ras in intercellular junctions, lessons
from toxins. Review of. *Biochimica et Biophysica Acta (BBA) - Biomembranes* **1788**:797-812. doi:
10.1016/j.bbamem.2009.01.011
Puram SV, Tirosh I, Parikh AS, Patel AP, Yizhak K, Gillespie S, Rodman C, Luo CL, Mroz EA, Emerick KS,
Deschler DG, Varvares MA, Mylvaganam R, Rozenblatt-Rosen O, Rocco JW, Faquin WC, Lin DT,
Regev A, Bernstein BE. 2017. Single-cell transcriptomic analysis of primary and metastatic tumor
ecosystems in head and neck cancer. Review of. *Cell* **171**:1611-1624.e1624. doi:
10.1016/j.cell.2017.10.044
Radisky ES, Radisky DC. 2010. Matrix metalloproteinase-induced epithelial-mesenchymal transition in
breast cancer. Review of. *Journal of Mammary Gland Biology and Neoplasia* **15**:201-212. doi:
10.1007/s10911-010-9177-x
Remboutsika E, Elkouris M, Iulianella A, Andoniadou CL, Poulou M, Mitsiadis TA, Trainor PA, Lovell-
Badge R. 2011. Flexibility of neural stem cells. Review of. *Frontiers in Physiology* **2**:16. doi:
10.3389/fphys.2011.00016
Sabbir MG, Wigle N, Loewen S, Gu Y, Buse C, Hicks GG, Mowat MR. 2010. Identification and
characterization of dlc1 isoforms in the mouse and study of the biological function of a single
gene trapped isoform. Review of. *BMC Biology* **8**:17. doi: 10.1186/1741-7007-8-17
Sakai D, Trainor PA. 2014. Gene transfer techniques in whole embryo cultured post-implantation mouse
embryos. Review of. *Methods Mol Biol* **1092**:227-234. doi: 10.1007/978-1-60327-292-6_15
Sakai D, Trainor PA. 2016. Face off against ros: Tcof1/treacle safeguards neuroepithelial cells and
progenitor neural crest cells from oxidative stress during craniofacial development. Review of.
Dev Growth Differ **58**:577-585. doi: 10.1111/dgd.12305
Sandell LL, Trainor PA. 2006. Neural crest cell plasticity. In *Neural crest induction and differentiation*,
edited by Saint-Jeannet J-P, 78-95. Boston, MA: Springer US.
Sauka-Spengler T, Bronner-Fraser M. 2008. A gene regulatory network orchestrates neural crest
formation. Review of. *Nature Reviews. Molecular Cell Biology* **9**:557-568. doi: 10.1038/nrm2428
Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld
S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K, Tomancak P, Cardona A. 2012. Fiji:
An open-source platform for biological-image analysis. Review of. *Nature Methods* **9**:676-682.
doi: 10.1038/nmeth.2019
Schorle H, Meier P, Buchert M, Jaenisch R, Mitchell PJ. 1996. Transcription factor ap-2 essential for
cranial closure and craniofacial development. Review of. *Nature* **381**:235-238. doi:
10.1038/381235a0
Soldatov R, Kaucka M, Kastriti ME, Petersen J, Chontorotzea T, Englmaier L, Akkuratova N, Yang Y, Häring
M, Dyachuk V, Bock C, Farlik M, Piacentino ML, Boismoreau F, Hilscher MM, Yokota C, Qian X,
Nilsson M, Bronner ME, Croci L, Hsiao W-Y, Guertin DA, Brunet J-F, Consalez GG, Ernfors P, Fried

K, Kharchenko PV, Adameyko I. 2019. Spatiotemporal structure of cell fate decisions in murine
neural crest. Review of. *Science (New York, N.Y.)* **364**. doi: 10.1126/science.aas9536
Soo K, O'Rourke MP, Khoo P-L, Steiner KA, Wong N, Behringer RR, Tam PPL. 2002. Twist function is
required for the morphogenesis of the cephalic neural tube and the differentiation of the cranial
neural crest cells in the mouse embryo. Review of. *Developmental Biology* **247**:251-270. doi:
10.1006/dbio.2002.0699
Stringer C, Wang T, Michaelos M, Pachitariu M. 2021. Cellpose: A generalist algorithm for cellular
segmentation. Review of. *Nature Methods* **18**:100-106. doi: 10.1038/s41592-020-01018-x
Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, 3rd, Hao Y, Stoeckius M, Smibert P,
Satija R. 2019. Comprehensive integration of single-cell data. Review of. *Cell* **177**:1888-1902
e1821. doi: 10.1016/j.cell.2019.05.031
Terry S, Nie M, Matter K, Balda MS. 2010. Rho signaling and tight junction functions. Review of.
Physiology **25**:16-26. doi: 10.1152/physiol.00034.2009
Thevenaz P, Ruttimann UE, Unser M. 1998. A pyramid approach to subpixel registration based on
intensity. Review of. *IEEE Transactions on Image Processing* **7**:27-41. doi: 10.1109/83.650848
Théveneau E, Duband J-L, Altabef M. 2007. Ets-1 confers cranial features on neural crest delamination.
Review of. *PLOS ONE* **2**:e1142. doi: 10.1371/journal.pone.0001142
Trainor P, Krumlauf R. 2000a. Plasticity in mouse neural crest cells reveals a new patterning role for
cranial mesoderm. Review of. *Nat Cell Biol* **2**:96-102. doi: 10.1038/35000051
Trainor PA. 2005. Specification of neural crest cell formation and migration in mouse embryos. Review
of. *Seminars in Cell & Developmental Biology* **16**:683-693. doi: 10.1016/j.semcdb.2005.06.007
Trainor PA, Krumlauf R. 2000b. Patterning the cranial neural crest: Hindbrain segmentation and hox
gene plasticity. Review of. *Nat Rev Neurosci* **1**:116-124. doi: 10.1038/35039056
Trainor PA, Krumlauf R. 2001. Hox genes, neural crest cells and branchial arch patterning. Review of.
Curr Opin Cell Biol **13**:698-705. doi: 10.1016/s0955-0674(00)00273-8
Van de Putte T, Maruhashi M, Francis A, Nelles L, Kondoh H, Huylebroeck D, Higashi Y. 2003. Mice
lacking zfhx1b, the gene that codes for smad-interacting protein-1, reveal a role for multiple
neural crest cell defects in the etiology of hirschsprung disease-mental retardation syndrome.
Review of. *American Journal of Human Genetics* **72**:465-470. doi: 10.1086/346092
Varberg JM, Unruh JR, Bestul AJ, Khan AA, Jaspersen SL. 2022. Quantitative analysis of nuclear pore
complex organization in schizosaccharomyces pombe. Review of. *Life Science Alliance*
5:e202201423. doi: 10.26508/lsa.202201423
Wakamatsu Y, Endo Y, Osumi N, Weston JA. 2004. Multiple roles of sox2, an hmg-box transcription
factor in avian neural crest development. Review of. *Developmental Dynamics* **229**:74-86. doi:
10.1002/dvdy.10498
Watt KEN, Trainor PA. 2014. Chapter 17 - neurocristopathies: The etiology and pathogenesis of disorders
arising from defects in neural crest cell development. In *Neural crest cells*, edited by Trainor PA,
361-394. Boston: Academic Press.
Weston JA. 1983. Embryonic development: The neural crest. Nicole le douarin. Cambridge university
press, new york, 1983. Xii, 260 pp., illus. $65. Developmental and cell biology series, 12. Review
of. *Science* **220**:1372-1372. doi: 10.1126/science.220.4604.1372
Wood HB, Episkopou V. 1999. Comparative expression of the mouse sox1, sox2 and sox3 genes from
pre-gastrulation to early somite stages. Review of. *Mechanisms of Development* **86**:197-201. doi:
10.1016/s0925-4773(99)00116-1
Yang J, Antin P, Berx G, Blanpain C, Brabletz T, Bronner M, Campbell K, Cano A, Casanova J, Christofori G,
Dedhar S, Derynck R, Ford HL, Fuxe J, Garcia de Herreros A, Goodall GJ, Hadjantonakis AK, Huang
RYJ, Kalcheim C, Kalluri R, Kang Y, Khew-Goodall Y, Levine H, Liu J, Longmore GD, Mani SA,
Massague J, Mayor R, McClay D, Mostov KE, Newgreen DF, Nieto MA, Puisieux A, Runyan R,

Savagner P, Stanger B, Stemmler MP, Takahashi Y, Takeichi M, Theveneau E, Thiery JP,
Thompson EW, Weinberg RA, Williams ED, Xing J, Zhou BP, Sheng G, Association EMTI. 2020a.
Guidelines and definitions for research on epithelial-mesenchymal transition. Review of. *Nat Rev*
Mol Cell Biol **21**:341-352. doi: 10.1038/s41580-020-0237-9

Yang X, Hu F, Liu JA, Yu S, Cheung MPL, Liu X, Ng IO, Guan XY, Wong KKW, Sharma R, Lung HL, Jiao Y, Lee
LTO, Cheung M. 2020b. Nuclear dlc1 exerts oncogenic function through association with foxk1
for cooperative activation of mmp9 expression in melanoma. Review of. *Oncogene* **39**:4061-
4076. doi: 10.1038/s41388-020-1274-8

Yu M, Bardia A, Wittner BS, Stott SL, Smas ME, Ting DT, Isakoff SJ, Ciciliano JC, Wells MN, Shah AM,
Concannon KF, Donaldson MC, Sequist LV, Brachtel E, Sgroi D, Baselga J, Ramaswamy S, Toner
M, Haber DA, Maheswaran S. 2013. Circulating breast tumor cells exhibit dynamic changes in
epithelial and mesenchymal composition. Review of. *Science (New York, N.Y.)* **339**:580-584. doi:
10.1126/science.1228522

Zhao R, Trainor PA. 2023. Epithelial to mesenchymal transition during mammalian neural crest cell
delamination. Review of. *Seminars in Cell & Developmental Biology* **138**:54-67. doi:
10.1016/j.semcdb.2022.02.018

1104

1105

Figures and figure legends

Figure 1. Single-cell RNA seq analysis of mouse early E8.5 cranial tissues. (A) Schematic of
experimental design. *Wnt1-Cre;RosaeYFP* and *Mef2c-F10N-LacZ* embryos with between 7-9
somites (6 each) were dissected and cranial tissues anterior to rhombomere 3 were collected.
Tissues were dissociated into single cell suspensions before being processed through the 10X
Genomics pipeline. The final dataset used for analysis consisted of 21,190 cells (12,498 cells
from *Wnt1-Cre;RosaeYFP* and 8,692 from *Mef2c-F10N-LacZ*) and 29,041 genes. (B) YFP and LacZ
staining of E8.5 *Wnt1-Cre;RosaeYFP* and *Mef2c-F10N-LacZ* embryos and 10um cranial
transverse sections. YFP (green) labels cells located in the dorsal neuroepithelium and their
lineages. As a result, both premigratory and migratory NCC are marked by YFP expression. LacZ
(blue) labels migratory NCC. (C) Uniform Manifold Approximation and Projection (UMAP) and
clustering of 6 major tissue types in the cranial region of E8.5 mouse embryos: cranial NCC,
neuroectoderm, non-neural ectoderm, mesoderm, endothelial cells, and embryonic blood cells.
(D) Dotplot showing the expression of tissue specific markers used for cluster identification. Dot

size indicates the percentage of cells in each corresponding cluster (y-axis) that expresses a

specific gene (x-axis). Dot color intensity indicates the average expression level of a specific

gene in a cell cluster.

1124

Figure 2. Expression of NCC development related genes and EMT functional genes identifies
NCC EMT intermediate populations. (A) UMAP and re-clustering of the cranial NCC cluster into
5 smaller subclusters at a resolution of 0.26. (B) Dotplot showing the expression of NCC

development related genes in 5 cranial NCC subclusters. (C) UMAP and re-clustering of the early
migratory NCC subclusters 0, 1 and 4 into smaller subclusters at a resolution of 2.0. (D)
Heatmap showing the expression of NCC development related genes in the smaller early
migratory NCC subclusters at a resolution of 2.0 shown in (C). High levels of expression are
indicated in yellow, and low levels of expression are indicated in pink. Based on the gene
expression profile of each subcluster, subcluster 17' was determined to be premigratory NCC;
subcluster 2' and 10' are EMT intermediate NCC; the remaining subclusters are migratory NCC.
(E) Dotplot showing the expression of EMT functional genes in premigratory NCC subcluster 17'
and intermediate NCC subclusters 2' and 10'. EMT intermediate NCC display reduced
expression of adherens junction, tight junction and apical basal polarity genes compared to
premigratory NCC, whereas protrusion related genes are upregulated in intermediate NCC.

1139

Figure 3. Mouse cranial NCC delaminate in S phase or G2/M phase cell cycle independently. (A)
Dotplot showing the expression of G2/M phase cell cycle genes in premigratory NCC (PM) and
EMT intermediate NCC. G2/M phase cell cycle genes are expressed in PM and intermediate
subcluster 10' cells. (B) Dotplot showing the expression of S phase cell cycle genes in PM and
EMT intermediate NCC. S phase cell cycle genes are expressed in PM and intermediate
subcluster 2' cells. (C) Pseudotime analysis of the cranial NCC cluster reveals the temporal
relationship between intermediate NCC subcluster 2' and 10'. Dark color indicates early NCC
development, and light color indicates later NCC development. PM and intermediate NCC
subclusters represent the earliest developmental timepoints among all cranial NCC. (D)
Trajectory analysis of the cranial NCC cluster reveals lineage/fate relationship between PM and
intermediate NCC subcluster 2' and 10'. Two intermediate NCC subclusters develop
simultaneously and independently from premigratory NCC. Apart from their cell cycle status,
early migratory NCC formed from the different intermediate subclusters are transcriptionally
indistinguishable.

1154

Figure 4. Cell cycle regulation plays an important role in mouse cranial NCC delamination and
EMT. (A) Cell cycle marker staining of early E8.5 mouse embryonic cranial and trunk tissues
reveals differences in cell cycle status between cranial delaminating premigratory NCC and
trunk neural plate border cells. EdU (magenta) and pHH3 (cyan) staining were performed on

10um transverse sections of early E8.5 (5-7 somites) *Wnt1-Cre;RosaeYFP* mouse embryo cranial
and trunk tissues. (B) E8.0 CD1 mouse embryos treated with Aphidicolin exhibit reduced
migratory NCC that primarily express pHH3. Cranial sections of treated embryos were stained
with Sox10, EdU and pHH3 (magenta) and arrowheads indicate migratory NCC expressing
pHH3. Most remaining migratory NCC in Aphidicolin treated samples express pHH3. In contrast,
a small proportion of migratory NCC in control DMSO treated samples express pHH3. (C) Cell
cycle staining quantification of delaminating premigratory NCC in the cranial neural plate
border shows that most cells express cell cycle markers. Staining and quantification were
performed on delaminating premigratory NCC in the cranial neural plate border of 5-7 somite
Wnt1-Cre;RosaeYFP mouse embryos (n=3). The neural plate border region was manually
selected in the most dorsolateral domain of the neural plate. EdU+%=the percentage of EdU
positive cells within eYFP positive delaminating premigratory NCC in the selected neural plate
border domain. pHH3+%= the percentage of pHH3 positive cells within eYFP positive
delaminating premigratory NCC. EdU+pHH3+%= the percentage of EdU and pHH3 double
positive cells within eYFP positive delaminating premigratory NCC. EdU-pHH3-%= the
percentage of EdU and pHH3 double negative cells within eYFP positive delaminating
premigratory NCC. (D) Cell cycle staining quantification of trunk neural plate border cells shows
that a significant proportion of cells do not express any cell cycle markers. Staining and
quantification were performed on trunk neural plate border cells of 5-7 somite *Wnt1-*
Cre;RosaeYFP mouse embryos (n=3). The neural plate border region was manually selected in
the most dorsolateral domain of the neural plate. EdU+%=the percentage of EdU positive cells
within DAPI positive neural plate border cells at the trunk axial level. pHH3+%= the percentage
of pHH3 positive cells within DAPI positive trunk neural plate border cells. EdU+pHH3+%= the
percentage of EdU and pHH3 double positive cells within DAPI positive trunk neural plate
border cells. EdU-pHH3-%= the percentage of EdU and pHH3 double negative cells within DAPI
positive trunk neural plate border cells. (E) Quantification of Sox10 expressing migratory NCC
upon Aphidicolin and control treatment reveals fewer cranial migratory NCC in Aphidicolin
treated embryos. Sox10 staining and quantification were performed on cranial sections of 4-6
somite CD1 mouse embryos post treatment (n=3 per treatment; ****$p<0.0001$). For
quantification, we calculated the ratio of Sox10 positive migratory NCC over DAPI positive
neural plate/neuroepithelial cells.

1190

Figure 5. SABER-FISH of EMT intermediate stage markers pinpoints the location of EMT
intermediate NCC within the dorsal most region of the neural fold. (A) Dotplot showing the
expression of selected EMT intermediate NCC markers in early migratory NCC subclusters
(resolution 2.0). (B) SABER-FISH staining of premigratory, EMT intermediate stage and
migratory NCC marker genes on the same section. Higher magnification insets of the left side
neural fold (box) showing that *Wnt1* is expressed in the neuroepithelium and *Sox10* is
expressed in migratory NCC populating the underlying mesenchyme. *Dlc1*, *Sp5* and *Pak3* are
expressed in the dorsolateral most region of the neuroepithelium. (C) 2D map showing the
number of transcripts per cell, calculated from the SABER-FISH staining. To evaluate the
expression of each gene within and across tissues, a polyline kymograph was generated along

the track indicated by the arrows at a width of 100 pixels. The polyline kymograph can be seen
to the right of each neural fold map it depicts. At the beginning of the track, *Wnt1* expression is
highest, demarcating the dorsal lateral domain of the neuroepithelium. Towards the middle of
the track, at the location of the most dorsolateral region of the neuroepithelium, *Wnt1* is
expressed along with the intermediate stage markers *Dlc1, Sp5* and *Pak3*. As the track
progresses to just outside of the neuroepithelium, *Sox10* expression appears and increases as
the track continues through the migratory NCC population.

1208

Figure 6. Dlc1 plays a regulatory role in mouse cranial NCC EMT and delamination. (A) Sox10
immunostaining was performed on cranial sections of E8.5 control and *Dlc1* knockdown mouse
embryos. (B) *Dlc1* knockdown significantly reduced the number of migratory NCC compared to
the control. The number of Sox10+ migratory NCC was quantified in control (n=4) and all *Dlc1*
knockdown (n=12) embryos. All datapoints in *Dlc1* knockdown samples were normalized to the
control samples. ****p<0.0001. (C) *Dlc1* shRNA-based lentiviruses achieved an average of 30%
reduction of *Dlc1* expression in all *Dlc1* knockdown embryos based on qRT-PCR analysis.
*p<0.05. (D) TUNEL staining showed minimal cell death in *Dlc1* knockdown samples.

1217

Figure 1-figure supplement 1. Expression of tissue specific marker genes that identify 6 major
cell type clusters in early E8.5 mouse embryonic cranial tissues. FeaturePlots that show the
expression of (A) *Sox2*, neuroectoderm marker; (B) *Sox1*, neuroectoderm marker; (C) *eYFP*,
premigratory and migratory NCC marker; (D) *LacZ*, migratory NCC marker; (E) *Sox10*, migratory
NCC marker; (F) *Twist1*, NCC and mesoderm marker; (G) *Tbx1*, mesoderm marker; (H) *Kdr*,
endothelial cell marker; (I) *Cdh1*, non-neural ectoderm marker; (J) *Hba-x*, embryonic blood cell
marker.

1225

Figure 2-figure supplement 1. Additional gene expressions that support the identity of cranial NCC subclusters at 0.26 and 2.0 resolutions. (A) Expression of NCC development program genes (Soldatov et al., 2019) confirms NCC subcluster 0 and 1 as early migratory NCC and subcluster 2 and 3 as late migratory NCC. Dotplots showing the expression of early and late NCC program genes in 5 NCC subclusters at resolution 0.26. Subcluster 0-3 all express early migratory NCC program genes, but only subcluster 2 and 3 express a significant level of late migratory NCC program genes. (B) UMAP and re-clustering of the cranial NCC cluster into 21 smaller

subclusters at a resolution of 2.0. (C-F) FeaturePlots show the expression of mesenchyme and
neurogenic lineage markers in cranial NCC. Subcluster 2 displays a high expression of
pharyngeal arch mesenchyme marker *Dlx2* (C). The frontonasal mesenchyme marker *Alx1* is
specifically expressed by subcluster 3 (D). Neurogenic lineage markers *Nrp1* and *Nrp2* are
expressed in late migratory NCC (E and F). (G-I) FeaturePlots showing expression of the
premigratory NCC marker *Wnt1* (G) and neural plate border specifiers *Pax7 and Zic2* (H and I) in
early migratory NCC. Subcluster 2' and 10' cells express reduced levels of *Wnt1*, *Zic2* and *Pax7*
than subcluster 17' cells.

1241

Figure 4-figure supplement 1. Aphidicolin treatment on *Mef2c-F10N-LacZ* embryos show
consistent results as CD1 embryos. (A) Aphidicolin treatment in early E8.5 CD1 embryos for 12-
13 hours prevented cells from entering S phase cell cycle and induced minimal cell death.
Aphidicolin treated embryos exhibit a lack of EdU incorporation and a similar level of TUNEL
signal compared to the control. (B) *Mef2c-F10N-LacZ* embryos treated with Aphidicolin for 12-
13 hours exhibit reduced migratory NCC as evidenced by β-galactosidase staining.

1248

Figure 5-figure supplement 1. Expression of EMT intermediate NCC markers in scRNA-seq data.
(A) Dotplot showing the expression of all EMT intermediate NCC markers in early migratory NCC
subclusters (resolution 2.0). (B-D) FeaturePlots showing the expression of *Dlc1*, *Sp5* and *Pak3* in
all early E8.5 cranial mouse embryo tissues. The expression of *Dlc1*, *Sp5* and *Pak3* is largely
restricted to NCC, neuroectoderm and mesoderm cells. Arrowheads indicate *Dlc1*, *Sp5* and *Pak3*
expression specifically in premigratory and intermediate NCC.

1255

Figure 5-figure supplement 2. Expression of intermediate NCC markers *Dlc1*, *Sp5* and *Pak3* by
SABER-FISH and traditional in situ hybridization in E8.5 mouse embryos and cranial sections.
Arrowheads indicate positive signals in the dorsolateral neuroepithelium. (A) *Dlc1* signal was
observed in the dorsolateral region of the neuroepithelium as well as in cells just outside of the
neuroepithelium in the underlying mesenchyme. (B) *Sp5* expression is localized within the
neuroepithelium and underlying dorsal mesenchyme. (C) *Pak3* expression is observed is
ectodermal tissues including the neuroectoderm or neuroepithelium.

A **B**

1263

Figure 6-figure supplement 1. *Dlc1* plays a regulatory role in mouse cranial NCC EMT and

delamination. (A) *Dlc1a*, *Dlc1b* and *Dlc1c* shRNA-based lentiviruses consistently achieved 30%

reduction of *Dlc1* expression based on qRT-PCR analysis. *p<0.05. (B) Embryos injected with

Dlc1a, *Dlc1b* and *Dlc1c* shRNA-based lentiviruses consistently showed significantly fewer

migratory NCC compared to the control. The number of Sox10+ migratory NCC was quantified

in control (n=4), *Dlc1a* (n=4), *Dlc1b* (n=4) and *Dlc1c* (n=4) embryos. All datapoints in *Dlc1a*,

Dlc1b and *Dlc1c* samples were normalized to the control samples. *p<0.05. ****p<0.0001.

1271

1272 Supplementary Table 1. NCC development related genes.

Gene	Expression pattern in relation to NCC development	References
Sox2	Neural epithelium (reduced in neural plate border)	Hafemeister & Satija, 2019; Lee et al., 2013; Wood & Episkopou, 1999
Sox1	Neural epithelium (reduced in neural plate border)	Hafemeister & Satija, 2019; Lee et al., 2013; Wood & Episkopou, 1999
Zic1/2	Neuroepithelium	Sauka-Spengler & Bronner-Fraser, 2008
Wnt1	Neural plate border	Echelard et al., 1994; Parr et al., 1993
Pax7	Neural plate border	Murdoch et al., 2012
GCNF (Nr6a1)	Neural epithelium and early MNCC	
Zeb2	Neural epithelium and early MNCC	Van de Putte et al., 2003
Pax3	Neural plate border and early MNCC	Li et al., 2000
(Wnt1-Cre) EYFP	Neural plate border and early MNCC	Hari et al., 2012
Sox9	Neural plate border and early MNCC	Lee et al., 2013
Foxd3	Neural plate border and early MNCC	Dottori et al., 2001

Snail1	Neural plate border and early MNCC	Cheung et al., 2005
Mef2c-F10N-LacZ	Predominantly MNCC	Aoto et al., 2015
Vimentin	MNCC (marker of mesenchymal cells)	Kobayashi et al., 2020
Sox10	MNCC	Hari et al., 2012
Twist1	MNCC	Soo et al., 2002

1273

1274   Supplementary Table 2. Primers for qRT-PCR

	Forward (5'-3')	Reverse (5'-3')
Dlc1	AGCGGCTGTGAAAGAAA	GCATTACCCTTGGAGAAGA
B2M	CACTGACCGGCCTGTATGC	GGTGGCGTGAGTATACTTGAATTTG
CANX	CCAGACCCTGATGCAGAGAAG	CCTCCCATTCTCCGTCCATA

1275

1276

1277