RpoS activates *Salmonella* Typhi biofilms and drives persistence in a Zebrafish model

Stuti K. Desai[1*], Yiyang Zhou[1], Rahul Dilawari[1], Andrew L. Routh[1,2,3,5],

Vsevolod Popov[3,4], and Linda J. Kenney[1,2,3*]

[1]Department of Biochemistry and Molecular Biology, [2]Sealy Center for Structural Biology and Molecular

Biophysics, [3]Institute for Human Infections and Immunity, and [4]Department of Pathology,

University of Texas Medical Branch, Galveston, TX 77555, and [5]Department of Immunology and

Microbiology, Scripps Research, 10550 N. Torrey Pines Rd., La Jolla, CA 92037, USA

[*]Address correspondence to: stdesai@utmb.edu or likenney@utmb.edu

Keywords: Static immersions, Transposon directed insertion site sequencing, ClickSeq, Asymptomatic Typhoid carriage, IraP, Fimbriae, Vi-antigen, heterologous host, Zebrafish

Abbreviations: red, dry and rough (rdar); colony forming unit (cfu); room temperature (RT); hour (h); Polymerase chain reaction (PCR); Reverse transcription quantitative real-time PCR (RT-qPCR); Transposon (Tn); next-generation sequencing (NGS); two-component regulatory system (TCRS)

Abstract

The development of strategies for targeting the asymptomatic carriage of *Salmonella* Typhi in chronic

typhoid patients has suffered owing to our basic lack of understanding of the molecular mechanisms that

enable the formation of *S.* Typhi biofilms. Traditionally, studies have relied on cholesterol-attached

biofilms formed by a closely related serovar, Typhimurium, to mimic multicellular Typhi communities

formed on human gallstones. In long-term infections, *S.* Typhi adopts the biofilm lifestyle to persist in vivo

and survive in the carrier state, ultimately leading to the spread of infections via the fecal-oral route of

transmission. In the present work, we studied *S.* Typhi biofilms directly, applied targeted as well as genome-

wide genetic approaches to uncover unique biofilm components that do not conform to the CsgD-dependent

pathway as established in *S.* Typhimurium. We adopted a genome-wide *Tn5* mutation screen in *S.* Typhi in

gallstone-mimicking conditions and generated New Generation Sequencing libraries based on the ClickSeq

technology to identify the key regulators, IraP and RpoS, and the matrix components as Sth fimbriae, Vi

capsule and lipopolysaccharide. We discovered that the starvation sigma factor, RpoS, was required for the

transcriptional activation of matrix-encoding genes in vitro, and for *S.* Typhi colonization in persistent

infections in vivo, using a heterologous fish larval model. Overall, our work established a novel RpoS-

driven paradigm for the formation of cholesterol-attached Typhi biofilms and emphasized the role(s) of

stress signaling pathways for adaptation in chronic infections.

Introduction

Salmonella enterica is a rod-shaped enteric bacterium that easily spreads through contaminated

food or water via the fecal-oral route in poor hygiene conditions. The human-restricted serovar of

Salmonella enterica, serovar Typhi (STy), causes typhoid fever and continues to be a dangerous pathogen

throughout the world[1]. The global incidence of typhoid fever in mostly children, adolescents and older

adults is between 12 to 27 million and in total 116,815 succumbed to the disease in 2017[2]. In contrast, its

closely related serovar Typhimurium (STm) infects diverse hosts such as humans, cattle, poultry and

reptiles to cause gastroenteritis that is mostly self-limiting in healthy adults.

Upon successful invasion of intestinal epithelial cells, *Salmonella* is phagocytosed by

macrophages, where it resides in a modified vacuole in a self-nourishing niche called a *Salmonella-*

Containing Vacuole (SCV) to ultimately reach the systemic sites of liver, spleen and the gall bladder. The

SsrA/B Two-Component Regulatory System (TCRS) is essential for the activation of the *Salmonella*

Pathogenicity Island-2 (SPI-2) regulon genes encoding a type-three secretory system and effectors that are

involved in formation of the SCV[3-5]. *Salmonella* also resides encased in extracellular matrix as

multicellular communities or biofilms, on intestinal epithelial cells[6], gallstones[7], tumors[8] and in the large

intestine[9].

Interestingly, SsrB, a response regulator of the SsrA/B TCRS, is essential for switching on the

multicellular lifestyle of *S.* Typhimurium by relieving H-NS silencing at the *csgD* promoter[10, 11]. CsgD is

the master regulator of STm biofilms, and it activates the transcription of extracellular matrix components

including curli fimbriae, cellulose, BapA and the O-antigen capsule[12-16]. Moreover, the SsrB-CsgD

regulatory pathway drives STm persistence in the heterologous host *Caenorhabditis elegans* by enabling

the formation of biofilms, which eventually promotes host life span through the p38-*Mitogen-activated*

Protein Kinase (p38-MAPK) innate immunity pathway[17].

Biofilms in the gall bladder are important for maintaining the carrier state of *Salmonella* Typhi,

allowing it to persist in 2 to 4% of chronic typhoid patients[18-20]. STy reservoirs in human carriers, who are

characteristically asymptomatic, play a crucial role in the spread of typhoid in epidemic regions, as well as

in its introduction to non-epidemic regions. Indeed, such long-term colonization of STy in human carriers,

coupled with the rise of multi-drug resistant strains, for example, those belonging to the H58 haplotype,

prevents effective control of typhoid fever [21-23]. Unfortunately, all of our understanding of the regulation

of STy biofilms on gallstones has been based on the assumption that it shares a high degree of conservation

with canonical biofilms formed by the closely related serovar, *S.* Typhimurium.

Herein, we establish that STy biofilms are fundamentally distinct from STm biofilms. In particular,

CsgD and the STm lifestyle regulator, SsrB, are not required for formation of multicellular aggregates on

surfaces coated with cholesterol that forms the major component of human gallstones. In order to identify

unique components of STy biofilms, we employed a whole genome transposon mutagenesis approach, Tn-

ClickSeq, and identified the starvation sigma factor RpoS as a crucial determinant of STy biofilms. We

discovered that the formation of large STy aggregates was defective in the absence of *rpoS* owing to the

down-regulation of extracellular matrix components comprised of Sth fimbriae, the Vi-antigen and the

lipopolysaccharide core.

Finally, we developed a heterologous host model, *Danio rerio*, to investigate STy lifestyles in

persistent infections for visualizing the colonization in real-time by confocal microscopy and measuring

the ensuing effects on host physiology. Zebrafish is a powerful vertebrate model for many human diseases

owing to a high degree of conservation of immuno-signaling pathways and colonization characteristics of

bacterial infections[24-27]. Previous studies have established that exposure of zebrafish larvae to *S.*

Typhimurium, *Mycobacterium marinum*, *Shigella flexneri*, and *Pseudomonas aeruginosa* leads to

successful pathogenesis[28-32]. *Shigella sonnei* and *S.* Typhimurium have also been recently shown to

persistently colonize macrophages in zebrafish[33, 34]. We infected zebrafish larvae with STy using static

immersions. Intestinal colonization was diminished in the *rpoS* null and larval survival at 6 days post

infection was greater, corroborating a crucial role of RpoS in prolonged STy survival in vivo.

Results

***S*. Typhi biofilms employ unique components that differ from *S*. Typhimurium biofilm**

Our previous analysis of *S.* Typhimurium (STm) biofilms established that unphosphorylated SsrB

was required for the expression of the master biofilm regulator CsgD to drive biofilm formation in vitro

and in vivo in the intestinal lumen of *C. elegans*[10, 35]. *S.* Typhi (STy) forms biofilms on gallstones in the

gall bladder and this ability is an important aspect of maintaining its carrier state[18-20]. It was therefore of

interest to examine whether STy employed similar pathways as STm for biofilm formation. We grew the

STy strain H58 under conditions that were proposed to mimic the gall bladder environment in humans[36],

hereinafter referred to as gallstone-mimicking conditions, and observed robust surface-attached

communities after two days in cholesterol-coated tubes, as measured by a crystal violet staining assay (Fig.

1A). The requirement of such distinct physico-chemical factors to develop biofilms was not specific to H58,

as three other isolates including Ty2b, CT18 and ct118 formed comparable cholesterol-attached biomass

after two days of growth (Fig. 1A). Not surprisingly, H58 also failed to form the characteristic rough, dry

and red ('rdar') morphotype at 30°C and low osmolality, as classically observed for the STm wild type

14028s (Supplementary Fig. 1A). The loss of the rdar morphotype, as well as the inability to form biofilms

in STm-specific environmental conditions has also been recently described for six other STy wild type

strains[37]. We followed the developmental course of H58 biofilm formation at two, four and six days in

cholesterol-coated tubes and observed only a marginal increase in the amount of biofilms at days 4 and 6

compared to day 2 (Supplementary Fig. 1B). We therefore focused on understanding STy biofilms at day

2 in our further investigations.

The response regulator SsrB plays a dual role in activating the virulent lifestyle in *S*. Typhimurium

(SsrB~P-dependent activation of SPI-2 genes), while driving the formation of biofilms in its

unphosphorylated form by de-repressing *csgD*[10, 11 for a review]. Hence, we examined whether complete

deletions of *ssrB* and *csgD* affected the ability to form cholesterol-attached *S*. Typhi biofilms. Interestingly,

biofilm formation was not affected by the loss of STy homologs encoding either the STm lifestyle regulator

SsrB, or the master regulator of STm biofilms CsgD, emphasizing the fundamental differences in

mechanisms of biofilm formation in these two closely related serovars (Fig. 1B). We also investigated

whether adhesion of STy aggregates to cholesterol-coated surfaces required curli fibers and O-Antigen, as

observed in the STm model, by inactivating *csgA* and *yihO/P,* respectively[13, 38]. We discovered that curli

fibers and O-Antigen did not play any role in the formation of cholesterol-attached STy biofilms, strongly

establishing that STy biofilms are drastically different from the model STm biofilms (Fig. 1B).

Tn-ClickSeq reveals the unique genetic signature of STy biofilms

Transposon-directed insertion site sequencing (TraDIS) has been immensely useful to study the

genetic repertoires essential for growth in STy and *Escherichia coli*, evolution of the invasive STm lineage

ST313, and biofilm formation in *E. coli* and *Pseudomonas aeruginosa*[39-43]. We applied a similar approach,

Tn-ClickSeq, by combining genome-wide transposon (Tn) mutagenesis with ClickSeq. ClickSeq is

advantageous here since it is a fragmentation-free next-generation sequencing (NGS) library synthesis

technique and is capable of generating focused NGS read data upstream of chosen target sites[44, 45], thus

greatly simplifying the digital transposon display protocol and removing artefactual recombination events

inherent to common NGS library preparation techniques. Using primers targeting the 3' or 5' ends of

inserted transposons, Tn-ClickSeq can sensitively and specifically sequence the junctions of a transposon

and the adjacent genomic loci of the integration site and thus identify genetic loci involved in forming

cholesterol-attached biofilms in vitro (Fig. 2A). A Tn-library in the *S*. Typhi strain H58 was kindly provided

by Stephen Baker, Cambridge University, UK[46]. We grew *S*. Typhi biofilms for two days using the H58-

Tn library in gallstone-mimicking conditions and isolated planktonic and biofilm fractions from a pool of

thirty cholesterol-coated tubes. NSG libraries were generated from genomic DNA isolated from each of

these sub-populations using the Tn-Click-Seq approach[45] (Fig. 2A).

Since Tn-ClickSeq returns short sequence reads containing a short fragment of the 3' or 5' end of

the inserted (known) Tn as well as a fragment of the adjacent genomic DNA, we developed a simple

computational pipeline that identified and trimmed the Tn-derived sequences from individual reads, and

then mapped the remaining fragment to the H58 genome (SAMEA3110714) using HISAT2. The 3' end of

these Tn-mapping reads represent the exact nucleotide junction of the insertion site of the Tn in the genome.

With this approach, we identified transposon insertions as 47%, 52% and 20% per million raw reads in the

input, planktonic and biofilm sub-populations, respectively (Supplementary Fig. 2A and B). Mapping of

insertion sites returned genomic locations of the inserted Tn in each dataset as well as the frequency of

these inserts within each original sample. After controlling for PCR duplication using UMIs included in the

'Click-Adaptor'[44], we assigned insertion indices by dividing the number of Tn-insertions per gene with 1

Kbp of gene length (TnClickSeq insertion indices.xlsx). With this approach, gene insertion frequencies in

each condition revealed clear differences in the planktonic and biofilm fractions, indicating unique genetic

components driving H58 lifestyles in gallstone-mimicking conditions (Fig. 2B). Principal Component

Analysis (Fig. 2C) and hierarchical clustering analysis (Fig. 2D) returned distinct clusters of genome

insertions sites of the three replicates of Tn-ClickSeq libraries obtained from the input, planktonic and

biofilm sub-populations. Overall, we successfully adopted the Tn-ClickSeq approach in *S*. Typhi to

generate a comprehensive view of genetic systems that determine the development of cholesterol-attached

biofilms from unattached planktonic cells.

Identifying matrix components and regulators of STy biofilms

To determine the exact mechanism(s) by which STy forms multicellular communities on

cholesterol surfaces in gallstone-mimicking conditions, we compared insertion indices between the

planktonic and biofilm fractions. Transposon insertions in biofilm genes likely disrupt functions resulting

in the inability of such mutants to form biofilms and would be enriched in the planktonic sub-population.

Interestingly, the majority of insertion indices had drastically lower values in the biofilm fraction compared

to the planktonic group, and these were of immediate interest for validation as STy biofilm targets

(TnClickSeq insertion indices.xlsx). We focused on loci having mean insertion indices greater than 100 in

the planktonic sub-population and performed Gene Ontology analysis on 1,515 such genes and identified a

significant enrichment in cell membrane components, transmembrane ion transport pathways, and other

membrane-related activities (Supplementary Fig. 2C). However, it was not possible to perform essentiality

analysis using a standard bioinformatics pipeline, because most of these genes had insertion indices equal

to zero in the biofilm fraction (see Methods and Discussion). In the absence of any *a priori* list of 'essential'

biofilm components, we narrowed our list of 1,515 genes to the top 200 and sought to investigate some for

their roles in STy biofilms. It was noteworthy that genes/operons encoding STm 'biofilm' homologs; *ssrB*,

csgDEFG, *csgBAC* or *yihPO* did not appear on our list of selected Tn-ClickSeq targets. This established a

strong correlation of our targeted genetic approach as described in Figure 1B, to our whole genome

transposon mutagenesis Tn-ClickSeq approach (Fig. 2A).

We next generated precise deletions of selected loci identified by Tn-ClickSeq that might encode

the structural components of STy biofilms in our gallstone-mimicking conditions: *sthC*, part of the *sth*

fimbrial operon[47, 48], *waaZ*, encoding an enzyme involved in the biosynthesis of the LPS core forming the

outer membrane[49] and *tviD*, a part of the *tviBCDE* operon encoding for the surface-exposed Vi-Antigen;

TviA is the regulator[50, 51]. We performed crystal violet staining assays at day 2 and determined that the

loss of *sthC*, *waaZ, tviA* and *tviD* led to a decrease in *S.* Typhi biofilms (Fig. 3A).

In parallel, while unraveling the regulatory circuits that drive the formation of surface-attached STy

communities, we were intrigued by the gene *iraP*, as it encodes an anti-adapter that inhibits the ClpXP-

dependent proteolysis of the starvation master regulator RpoS[52]. We tested the biofilm formation

capabilities of an *iraP* null H58 strain and observed a substantial reduction in biofilms compared to the wild

type parent at 2 days (Fig. 3B). Since the only known role of IraP is an indirect positive role in activating

the starvation stress response in enteric bacteria[53], we examined whether deletion of *rpoS* affected *S.* Typhi

biofilms. Indeed, we discovered a positive effect of RpoS in regulating the formation of cholesterol-attached

STy biofilms (Fig. 3B). Over-expression of RpoS *in trans* (a kind gift from Roy Curtiss III, University of

Florida), complemented the defect in biofilm formation of the *rpoS* null strain (Fig. 3B) and increased

biofilms compared to the wildtype. Our discovery of RpoS as an activator of *S.* Typhi biofilms is significant,

as biofilms were not mediated through CsgD (unlike in *S.* Typhimurium, see Fig. 1B). Finally, we ruled out

an effect of mere growth differences on biofilm formation in the STy mutants by monitoring the planktonic

growth of *iraP*, *rpoS*, *sthC*, *waaZ* and *tviD* null mutants compared to the wild type H58 parent. We observed

similar growth rates among all the strains indicating that IraP, RpoS, SthC, WaaZ and TviD were necessary

for the formation of *S.* Typhi surface-attached communities (Supplementary Fig. 3).

Cholesterol-attached aggregates of STy biofilm mutants exhibit a poor ultrastructure

Our understanding of the CsgD-dependent mechanisms of biofilm formation in *S.* Typhimurium has been

achieved by screening mutant strains for their inability to exhibit the rdar morphotype and high-resolution

imaging of biofilm components by fluorescence confocal microscopy[10, 35, 37, 54-56]. Unfortunately, neither

of these approaches can be applied to unequivocally establish the molecular players that drive the formation

of STy biofilms on cholesterol-coated surfaces because of a high level of autofluorescence in the gallstone-

mimicking conditions (see above and Discussion). We therefore grew STy biofilms in cholesterol-coated

tubes for two days and obtained direct visualizations of surface-attached communities by scanning electron

microscopy. We observed dense aggregates of the wild type parent having a rich network of extracellular

matrix that connected the cells, while the strains defective in *rpoS*, *sthC*, *waaZ* and *tviD* formed smaller

STy aggregates with poor network connections (Fig. 4). Our high-resolution observations of cholesterol-

attached STy communities by scanning electron microscopy consolidated our in vitro measurements of the

sessile biomass formed by the STy strains by the standard crystal violet staining assay (Fig. 3, above).

Overall, we demonstrated that the Tn-ClickSeq approach enabled the identification of previously

unidentified players that drive the formation of atypical STy biofilms.

RpoS activates transcription of STy biofilm matrix genes

We next wanted to determine the possible role of RpoS in regulating the expression of STy biofilm

components, including: SthC, WaaZ and the Vi capsule. We first investigated if the expression of *rpoS*,

sthC, *waaZ* and *tviB* (the *tviBCDE* operon mediates biosynthesis of Vi-polysaccharide[57]), was up-regulated

in biofilm-inducing conditions. We isolated planktonic and biofilm sub-populations from wild type strain

grown in cholesterol-coated tubes and measured steady state transcript levels by reverse transcription

quantitative real-time PCR (RT-qPCR). Indeed, we detected an increase in the transcription of *rpoS* (~20-

fold), *sthC* (~4-fold), *waaZ* (~200-fold) and *tviB* (~10-fold) in biofilms formed by the H58 parent in

gallstone-mimicking conditions (Fig. 5A).

In order to investigate the exact role of RpoS in favoring the biofilm lifestyle, we grew the wild

type parent H58 and its *rpoS* null derivative in cholesterol-coated tubes in vitro and isolated total RNA from

the biofilm fractions after two days. We then compared the transcription of matrix encoding genes *waaZ*,

sthC and *tviB* by RT-qPCR and observed a positive effect of RpoS at each of these loci (Fig. 5B).

Transcription of the matrix-encoding genes was clearly down-regulated when RpoS was absent (Fig. 5B).

This result established a molecular link between the stress sigma factor, RpoS, and development of STy

cholesterol-attached biofilms.

RpoS regulates infection outcomes in zebrafish larvae and drives persistence in vivo

Since the formation of STy biofilms plays a crucial role in enabling persistence in human carriers,

we investigated the role of RpoS in persistent infections in a zebrafish model. We replicated the natural

mode of Typhi entry by employing static immersions of zebrafish larvae in tank water contaminated with

specific doses of *S.* Typhi strains. We exposed 5 days post-fertilization (dpf) zebrafish larvae to 10^8 cfu/ml

of mCherry-expressing wild type and the isogenic *rpoS* null mutant in system water for 24 h. After 24 h,

the larvae were shifted to plain system water and monitored for six days under standard conditions of

zebrafish husbandry[adapted from 29]. Control larvae were exposed to an equal volume of PBS. We discovered

that *S*. Typhi colonization of infected larvae was reduced when RpoS was absent, as evident from both

bacterial load measurements (Fig. 6A), as well as real-time visualizations at 2, 4 and 6 dpi using confocal

fluorescence microscopy (Fig. 6B). At longer times of 4 and 6 dpi, wild type *S*. Typhi clearly persisted in

the intestine (Fig. 6A, B), as also observed previously in a mouse model of STm persistence[58]. Finally,

persistence was strongly correlated with pathogenesis outcomes as larvae infected with the *rpoS* null strain

survived in significantly greater numbers than the wild type infections at 6 dpi (Fig. 6C). In fact, the survival

rate of *rpoS* infections was similar to the uninfected PBS control from 1 to 6 dpi, strongly emphasizing the

requirement of RpoS for enabling STy survival in long term infections. Monitoring infected larvae beyond

6 dpi is challenging, owing to a greater requirement of live food (Paramecia) which masks pathogen-driven

physiological effects[59 and see Discussion]. In conclusion, we revealed a positive role for RpoS in chronic STy

infections and established that RpoS was necessary for persistence in vivo.

Discussion

The dangerous transmission of *S*. Typhi from seemingly healthy but chronically infected

individuals is well documented and the first cases in the United States of Mary Mallon, 'Typhoid Mary',

and Mr. N in the United Kingdom, were reported in the early 1900s[60, 61]. Since then several epidemiological

studies have established that persistent colonization of STy in the gall bladders of asymptomatic patients

forms the basis for typhoid carriage[reviewed in 18, 62]. Despite clear evidence of the role of STy biofilms in

spreading the disease, previous studies have failed to delineate any genetic mechanisms that regulate the

development of gallstone biofilms in STy. This is chiefly because biofilms formed by *S*. Typhimurium have

been employed as a surrogate for understanding the multicellular behavior of *S*. Typhi[63-65]. In the present

work, we studied STy biofilms directly and discovered that the pathways for biofilm formation in STy and

STm were completely different. More importantly CsgD, the master regulator of STm surface-attached

communities, was entirely dispensable for the formation of STy sessile aggregates on cholesterol-coated

surfaces. As a result, matrix production in STy biofilms did not require curli fibers or biosynthesis of O-

Antigen via the *yihO/P* system.

We grew a transposon library in the STy parent strain H58 as biofilms in vitro in gallstone-

mimicking conditions and developed Tn-ClickSeq analysis to map the transposon genome junctions

enriched in the planktonic and biofilm sub-populations. We found Tn-genome junctions in 47% and 20%

of total sequencing reads from the planktonic and biofilm fractions, respectively. These differences

indicated that presumably inactivating transposon insertions were tolerated in only a low number of genes

in biofilms, which correlated with their enrichment in the planktonic fraction. Also the technical challenges

of isolating genomic DNA with high efficiency from STy cells attached to cholesterol-coated surfaces may

affect the total sequencing yields, leading to very low insertion indices in biofilms. Nevertheless, our Tn-

ClickSeq analysis generated novel insights by identifying Sth fimbriae, Vi capsule and the

lipopolysaccharide core as structural components and IraP as a regulator of STy cholesterol-attached

biofilms. Our comprehensive analysis of the STy biofilms also opens up exciting new directions for

mapping the complete genetic signature of STy carriage.

The discovery of IraP is significant because it motivated our subsequent investigations on the

significance of the downstream starvation-induced sigma factor, RpoS, during STy biofilm development.

We established that the formation of STy biofilms required RpoS and determined its role in activating the

transcription of the extracellular matrix components: Sth fimbriae, Vi capsule and lipopolysaccharide.

Interestingly, the binding sites for RpoS have been identified in the *waaZ* promoter in *E. coli*[66]. A previous

study had also described a regulatory interplay between RpoS and the TCRS RcsC/B to regulate Vi capsule

synthesis under different osmotic conditions[67].

A high degree of autofluorescence and the non-specific binding of fluorescent dyes and antibodies

on cholesterol surfaces prevents the visualization of mature STy biofilms using high-resolution

fluorescence microscopy techniques. Although some protocols have been developed to improve the

fluorescence imaging of STy aggregates in vitro and ex vivo, crucial controls that included STy strains

defective in forming biofilms were absent[68-70]. Therefore, in order to clearly visualize STy biofilms

attached to cholesterol surfaces, we employed scanning electron microscopy and observed distinct

aggregates of the wild type H58 parent and substantially reduced aggregate formation in the STy biofilm

mutants validated from our Tn-ClickSeq analysis.

Our use of zebrafish larvae as a heterologous host for *S.* Typhi validated the important role of RpoS

in enabling *S.* Typhi colonization in long-term infections. Reducing STy persistence by inactivating *rpoS*

significantly decreased the pathogen load and prolonged host survival. Following STy infections beyond

6 dpi may be possible by rearing germ-free zebrafish larvae using established methods[71].

Finally, a strong correlation has been observed between the ability to form biofilms and the duration

of STy shedding and carriage in typhoid patients from Pakistan[72]. Most recent phylogenomic analysis has

proposed that the ancestral H58 haplotype originated in a chronic carrier from India and evolved to give

rise to the three sub-lineages that cause a majority of typhoid infections in Asia and Africa[23]. In the light

of these results, it is tempting to propose that RpoS contributes to niche adaptation in *S.* Typhi by activating

the formation of biofilms in chronic carriers.

(A) (B)

Figure 1: The master regulator CsgD, and the STm lifestyle regulator SsrB, are not required for the formation of cholesterol-attached STy biofilms. Crystal violet staining assays: (A) Wild type strains Ty2b, CT18, ct117 and H58 formed robust biofilms at day 2 in gallstone-inducing conditions and (B) STy null strains deleted for the indicated STm-biofilm homologs formed similar biofilms as the H58 wild type strain at day 2. Growth medium added to cholesterol-coated Eppendorf tubes was used as the control and subtracted from all measurements. N = 3 in at least triplicates, error bars represent Mean ± SD, ns = not significant by One-way ANOVA.

(A) (B)

Supplementary Figure 1: H58 forms 'atypical' biofilms. (A) Examination of a macrocolony of *S.* Typhimurium (STm) displaying a <u>r</u>ed, <u>d</u>ry and <u>r</u>ough (rdar) morphotype on low salt Luria-Bertani agar medium containing the congo red dye, while *S.* Typhi (STy) are 'smooth' and lack the rdar morphology. (B) The cholesterol-attached biomass formed by wild type H58 did not increase significantly at days 4 and 6 compared to day 2, as determined by crystal violet staining assay. Growth medium added to cholesterol-coated Eppendorf tubes was used as the control and subtracted from all measurements. N = 3, in at least triplicates, error bars represent Mean ± SD, ns = not significant by one-way ANOVA.

Figure 2: Tn-ClickSeq analysis investigated novel pathways that drive the formation of STy biofilms. (A) A general scheme depicting Tn-ClickSeq analysis, from the isolation of planktonic and biofilm fractions, DNA extraction and the generation of Tn-ClickSeq libraries using a reverse transcription reaction (RT) with a Tn-specific primer and azido-nucleotides/dNTP mixtures. The azido-terminated cDNAs were click-ligated to an alkyne-adapted illumina P5 adapter to generate Illumina libraries. After PCR and barcoding, the reads that contained partial Tn sequence and genome insertion site were pair-end sequenced and subjected to downstream analysis. (B) A linearized map of the average Tn-insertion sites from three biological replicates showed genome-wide differences in the planktonic sub-population (orange) and biofilms (green). The H58-Tn library inoculum (blue) and the H58 parent without any transposons (black, control) were used as positive and negative controls, respectively. (C) Principal component analysis showed distinct clustering of respective replicates of planktonic (orange) and biofilm (green) fractions as compared to the input library (blue). (D) The number of transposon insertions of each gene was normalized to per 1000 bp of gene length and averaged among replicates for each group to depict a hierarchical clustering of Tn-ClickSeq targets.

Supplementary Figure 2: (A) Percentages of reads that matched the primer, transposon sequence (IR sequence) and the host genome from left to right: input, planktonic and biofilm sub-populations. (B) Graphical representation of Tn insertion sites per one million of raw reads as obtained by Tn-ClickSeq showing an average of 470,000 reads for the input fraction, 520,000 for the planktonic fraction and 197,000 for the biofilm fraction. A similar analysis using an H58 strain without any transposon insertions resulted in only 100 false positives. (C) A Gene Ontology analysis of transposon insertion sites in 1515 genes that were enriched in the planktonic sub-population as compared to biofilms showed a significant enrichment of cell membrane components, transmembrane ion transport pathways and other membrane related activities.

Figure 3: Biofilms are defective in STy null mutants identified by Tn-ClickSeq. (A) Defective biofilms formed by *tviA/tviD*, *sthC* and *waaZ* null strains suggested a role of Vi capsule, Sth fimbriae and the lipopolysaccharide core in extracellular matrix production, respectively. (B) H58 strains deleted for *iraP*, which regulates the protein stability of RpoS, and *rpoS,* which encodes the starvation sigma factor, formed less biofilms than the wild type parent. The defect in biofilm formation of the *rpoS* null strain was complemented by overexpression of *rpoS* from a plasmid *in trans*. N = 3, in at least triplicates, error bars represent Mean ± SD, in a crystal violet staining assay. Growth medium added to cholesterol-coated Eppendorf tubes was used as the control and subtracted from all measurements, ns = not significant, *p ≤ 0.05, **p ≤ 0.01 and ***p ≤ 0.001 by one-way ANOVA.

Supplementary Figure 3: Biofilm mutants are not compromised in planktonic growth.

Strains deleted of *iraP*, *rpoS*, *waaZ*, *sthC* or *tviD* and the wild type H58 strain were grown

for 12 hours in Luria-Bertani broth at 37°C/250 rpm and the absorbance at 600 nm was

measured every 2 hours.

Figure 4: Ultra-structure visualization reveals the loss of dense aggregate formation in STy biofilm mutants. Representative scanning electron microscopy images of rich *S*. Typhi biofilms formed by the H58 (wt) and strikingly smaller sized aggregates formed by the strains deleted of *rpoS*, *sthC*, *waaZ* or *tviD* when grown in cholesterol-coated Eppendorf tubes in gallstone-mimicking conditions.

Scale bar = 5 μm.

Figure 5: Steady-state levels of biofilm components are increased in gallstone-mimicking conditions and RpoS is required for the expression of biofilm matrix genes. Real time RT-qPCR analysis showed (A) a significant increase in transcription of *rpoS*, *sthC*, *waaZ* and *tviB* in the cholesterol-attached fraction compared to the planktonic fraction in the wild type H58 parent and (B) In the *rpoS* null strain, there was a significant decrease of biofilm matrix components *waaZ*, *tviD* and *sthC*. *rrsA* transcript levels were used as an internal control. N = 3, in triplicates, error bars represent Mean ± SD, *p ≤ 0.05, **p ≤ 0.01, ***p ≤ 0.001 and ****p ≤ 0.0001 by one-way ANOVA.

(A)

(C)

(B)

Figure 6: STy colonizes the intestines of chronically infected zebrafish, and RpoS is required for persistent colonization in vivo. (A) The number of STy colonies recovered from persistently infected whole larvae was drastically reduced in the *rpoS* null strain (blue bars) compared to the wild type (green bars) at days 2, 4 and 6 post infection. Control larvae exposed to PBS were void of *Salmonella*. N = 3, five infected larvae from each group, error bars represent Mean ± SD and ****p ≤ 0.0001 by two-way ANOVA with Sidak's multiple comparison tests. (B) Representative merged images (red channel and bright field) of live zebrafish larvae showing the presence of wild type mCherry-expressing *S.* Typhi in the gut. There was a stark reduction of bacteria in the infections using the mCherry-tagged *rpoS* null strain at 2, 4 and 6 dpi. No fluorescence was detected for the PBS control at all time points. The cartoon on top depicts an infected zebrafish larva with the highlighted intestinal region was adapted from BioRender. 20x magnification, Scale bar = 10 µm. (C) Larvae infected with the *rpoS* null strain (blue line) survived longer and showed a significant increase in the percentage of survival at 6 dpi compared to the wild type H58 (green line). The survival rate of the PBS control is shown as a dotted black line. N = 3 with 30 to 60 larvae in each group, error bars represent Mean ± SD, *p ≤ 0.05 by two-way ANOVA with Sidak's multiple comparison tests.

Methods

Bacterial strains and growth

The bacterial strains and plasmids used in this study are listed in Supplementary Table 1. *S.* Typhi strains

were routinely grown in Luria-Bertani broth (LB) or Nutrient broth (NB) (BD Difco) medium with shaking

at 275 rpm at 37°C in the presence of 100 μg/ml Ampicillin, 12.5 μg/ml Tetracycline, 25 μg/ml

Chloramphenicol or 50 μg/ml Kanamycin (Millipore Sigma) when necessary. For the growth of STy

cholesterol-attached biofilms, a modified NB medium (NB*) containing 3% w/v NB, 1.75% w/v sodium

chloride, 0.25% w/v potassium chloride, 1% w/v sodium choleate and 2% w/v glucose (Millipore Sigma)

was used [36]. For observing the rdar morphotype, 40 μl of overnight LB broth cultures of STm wild type

strain 14028s and STy wild type strain H58 were spotted on agar plates containing 1% w/v Tryptone and

0.5% w/v Yeast Extract supplemented with Congo Red (40 μg/ml) (Millipore Sigma) and kept at 30°C for

two days.

Molecular Biology techniques

All DNA manipulation procedures were carried out according to[73] using reagents procured from Qiagen,

Millipore Sigma or Invitrogen. All transformations in STy wild type strain H58 were performed by standard

electroporation protocols[73]. Polymerase chain reactions (PCR) were carried out using oligonucleotides as

listed in the Supplementary Table 2 following standard protocols[73].

Strain construction

The *ssrB* null mutation in H58 strain was generated by transducing the *ssrB::kan* allele from the STm strain

DW85 using standard P22 transduction protocols[74]. Other gene deletions in the STy wild type strain H58,

as listed in Supplementary Table 1, were generated by the lambda *red* homologous recombination technique

as described in[75]. Briefly, plasmids pKD3, pKD4 or the *TetRA* DNA were used to generate linear DNA

fragments by PCR using gene-specific hybrid primers as listed in Supplementary Table 2. A 10 ml LB broth

culture of H58 transformed with the plasmid pKD46 (containing 100 μg/ml Ampicillin and 20 mM

Arabinose) was used to electroporate 600 ng to 1 μg of purified PCR product following the protocol as

described in[75]. The cells were recovered for at least 6 h at 275 rpm/30 °C after which the cells were

harvested and plated on respective selective plates. Chromosomal deletions were confirmed by PCR using

flanking primer pairs as listed in Supplementary Table 2.

STy cholesterol-attached biofilms in gallstone-mimicking conditions

STy biofilms were routinely grown following protocols adapted from[36, 38]. Sterile 1.5 ml Eppendorf tubes

containing 200 µl of 10 mg/ml cholesterol (Millipore Sigma) in diethyl ether were air dried aseptically (2-

3 h) and 20 µl of STy strains grown overnight in LB/NB was added to 180 µl NB* medium. The tubes were

incubated for two days at 275 rpm at 37 °C. Cholesterol-coated tubes containing only 200 µl NB* medium

served as controls in all the experiments. For monitoring the time course of biofilm formation, the

cholesterol-coated tubes were further incubated for four and six days, with fresh 200 µl NB* medium

replaced every two days.

Crystal violet staining assay

The amount of STy cholesterol-attached biofilms in gallstone-mimicking conditions was estimated using

crystal violet staining assays adapted from[10, 38]. The supernatant/growth medium was removed, and each

tube was washed once with 400 µl of Phosphate-buffered Saline (PBS). The attached biofilms were then

stained with 200 µl of 0.1 % w/v crystal violet solution (filtered using Whatman Grade 1 filter paper) for

five minutes at room temperature (RT). This was followed by washing once with 400 µl PBS and addition

of 200 µl absolute ethanol. Appropriate dilutions were then measured for absorbance at 595 nm using an

iMark™ Microplate Absorbance Reader. Each experiment was performed in triplicates or pentuplicates.

Isolation of fractions for Tn-ClickSeq libraries

A *TnTMDH5deloriR6K* genome library in the *S*. Typhi wild type strain H58 (H58-Tn) was kindly provided

by Dr. Stephen Baker, University of Cambridge, UK. Details on the construction of the library have been

previously described in[46]. A 1 µl loop of frozen stock of the H58-Tn library was inoculated in 10 ml NB*

cultures and incubated overnight at 275 rpm at 37 °C. This was the 'input' fraction, which was used to

prepare thirty tubes of cholesterol-attached biofilms in gallstone-mimicking conditions. After two days, the

culture supernatants were removed and pooled to obtain the 'planktonic' fraction. To isolate the 'biofilm'

fraction, 400 µl of PBS was added to each tube followed by sonication in a XUB Digital Ultrasonic Bath

(Grant Instruments) for 20 minutes, maximum power, no leap, at RT. The harvested biomass was pooled,

and the sonication step was repeated thrice. The final pool at the end of four sonication cycles was the

'biofilm' fraction.

Genomic DNA isolation for generating Tn-ClickSeq libraries

Planktonic and biofilm fractions were centrifuged at 24,000 g at 4 °C for 1.5 h in a Beckman Coulter Avanti

J-26XP centrifuge. Respective supernatants were discarded, and the pellets were stored on ice. 1 ml of the

input fraction was also centrifuged at 15,500 g at 4 °C for 10 min and the pellet was stored on ice. Each of

the pellet fractions were then resuspended in 600 µl TE buffer (pH 8.0) with 40 µl 10% w/v sodium dodecyl

sulphate, 4 µl Proteinase-K and 2 µl RNaseE and mixed well by vortexing. Samples were incubated at 37

°C for 1 h, after which an equal volume of Phenol:Chloroform mixture (pH 6.7/8.0) was added, mixed and

centrifuged at 15,500 g at 4 °C for 15 min. The upper aqueous phase was added to a fresh tube and an equal

volume of chloroform was added, mixed, and centrifuged at 13,000 rpm at 4 °C for 15 min. The supernatants

were removed, 2.5 to 3 volumes cold absolute ethanol was added, and stored overnight at -20 °C. DNA

pellets were obtained by centrifuging the samples at 15,500 g at 4 °C for 15 min, followed by a 70% ethanol

wash. The pellets were air dried, and DNA was resuspended in 40 µl of nuclease-free water.

Preparation of Tn-ClickSeq libraries

For Tn-ClickSeq, genomic DNA from the input, planktonic and biofilm fractions was reverse transcribed

using a reverse transcriptase (SSIII, Invitrogen) and Azido-NTPs. A Tn-specific reverse primer (3'21-39,

Supplementary Table 2) was designed to the 3' proximal end, 21 to 39 bp of transposon, with an overhang

of the reverse complementary sequence of Illumina adapter (Supplementary Figure 4, Supplementary Table

2). 500 ng DNA was mixed with 1 µL of 5 µM primer, and 1 µL of 10 mM AzNTP/dTNP mixture

(AzNTP:dNTP = 1:35). This initial reaction mix was heated at 95 °C for 5 min and then cooled to 50 °C in

gradual steps of 0.1 °C per second. Other reaction components for the reverse transcription reaction were

then added: buffer, DTT, SSIII (as per manufacturer's protocol), and kept in a thermocycler at 50 °C for 50

min, after which the reaction was terminated at 95 °C for 5 min. This was followed by the standard ClickSeq

protocol as previously described[44, 76, 77]. Briefly, immediately after denaturing, the DNA products were

purified with Solid Phase Reversible Immobilization (SPRI) beads and click-ligated with a 5'-alkyne-

modified adapter including a 12-nucleotide unique molecular identifier (UMI). The click-ligated product

was then purified, barcoded, amplified with 18 to 20 cycles of PCR, and analyzed by agarose gel

electrophoresis. The final Tn-ClickSeq libraries were subjected to pair-end sequencing on a NexSeq 550

platform at the Genomics core, UTMB.

Bioinformatics analysis

We designed a bioinformatics pipeline to process the hybrid reads that originate from the inserted

transposon and extend into the host genome (Supplementary Figure 5). The raw paired-end FASTQ reads

were first pre-processed to trim the Illumina adapter, filter low-quality reads and extract UMIs using

fastp[78]: -a AGATCGGAAGAGC -U --umi_loc read1 --umi_len 14 --umi_prefix umi -l 30. We then used

FASTX toolkit (http://hannonlab.cshl.edu/fastx_toolkit/index.html) to reverse complement the R2 reads for

ease of downstream analyses. We filtered the reads that contained the 19 bp primer sequence (targeting the

Tn) with 1 nucleotide mismatch allowance with *cutadapt*[79]: -a cctatagtgagtcgtatta -e 0.1 -O 19 -m 30 --

discard-untrimmed. To obtain the reads containing primer sequences, we further filtered reads that

contained the last 10 nucleotides of the 3' invert repeat (IR) with 0% error rate allowance with *cutadapt*: -

a ctgtctctta -e 0 -O 10 -m 30 --discard-untrimmed. After trimming the IR sequence, the rest of IR-containing

reads   were   mapped   to   the   H58   genome

(https://www.ncbi.nlm.nih.gov/datasets/genome/GCF_001051385.1/) with *hisat2*[80], and then processed

with *SAMtools*[81]: view/sort/index. We de-duplexed the data to minimize PCR bias with umi_tools[82]:

dedup --method=unique. Finally, the locations of the transposon insertion sites were extracted with

BEDTools[83]: genomeCoverageBed: -3 -bg. Raw sequencing data is available at the Sequence Read

Archive, Project ID, PRJNA1029173.

Gene mapping and target analysis

The number of insertion reads at each insertion site revealed by Tn-ClickSeq were ratiometrically

normalized across different samples. This was followed by the assignment of gene names and annotations

within H58 (GCF_001051385.1). We normalized the number of insertions to the length of each annotated

gene (per 1 Kbp length of gene) to obtain the insertion indices for each samples. Matrices of all the

normalized insertion dataset were then processed with DESeq2 to identify genes enriched or depleted in

each conditions and perform Principal component analysis[84]. Hierarchical clustering was conducted with

Cluster 3.0 (http://bonsai.hgc.jp/~mdehoon/software/cluster/), followed by TreeView

(http://jtreeview.sourceforge.net/) to build the graphic map. Gene ontology analysis was conducted with

GeneOntology web server (http://geneontology.org/) with *Salmonella* Typhimurium as a reference.

RNA isolation

Planktonic and biofilm fractions were centrifuged at 24,000 g at 4 °C for 1.5 h in a Beckman Coulter Avanti

J-26XP centrifuge. Respective supernatants were discarded, and the pellets were either stored at -80 °C, or

immediately processed for total RNA isolation by the Trizol method[85]. Briefly, the pellets were

resuspended in 1 ml TRIzol reagent (Life Technologies) and incubated at RT for 5 min. 200 µl chloroform

was added, mixed well, and incubated for 3 min at RT. The mixtures were centrifuged at 15,500 g for 15

min in cold and 500 µl isopropanol was added to the supernatants. The samples were transferred to -20 °C

overnight, after which the RNA was pelleted by centrifugation at 15,500 g for 15 min in cold. This was

followed by a 75% ethanol wash. The pellets were air dried, and the RNA was resuspended in 20 µl of

nuclease-free water.

RT-qPCR

1 µg of total RNA extracted from the biofilm fraction of two days old cholesterol-attached biofilms was

used for a reverse transcription reaction with iScript Supermix (Bio-Rad) according to the manufacturer's

protocol. This was followed by amplifying 50 ng cDNA by real-time qPCR (RT-qPCR) using SsoFast

EvaGreen Supermix (Bio-Rad) and internal primers specific for *rpoS*, *sthC*, *waaZ* and *tviB*; *rrsA* was used

as a normalization control (Supplementary Table 2). The annealing temperature for all the primer pairs was

56 °C. All experiments were performed in triplicates with at least three independently isolated RNA

preparations. Relative expression was determined using the $2^{-\Delta\Delta C_T}$ (Livak) method as described in[85] and

plotted using the GraphPad Prism 10 software.

Scanning electron microscopy

STy cholesterol-attached biofilms were grown in gallstone-mimicking conditions for two days and after

removal of the growth medium were fixed in a primary fixative containing 2.5% w/v formaldehyde (made

from paraformaldehyde), 0.1% v/v glutaraldehyde, 0.01% v/v trinitrophenol and 0.03% w/v $CaCl_2$ in 0.05M

sodium cacodylate buffer (pH 7.3). Samples were post-fixed in 1% w/v OsO_4 in cacodylate buffer,

dehydrated in ethanol and infiltrated with hexamethyldisilazane (HMDS) to prevent cracking during drying.

After air drying, the conical parts of the Eppendorf tubes were cut into strips, mounted on SEM specimen

holders (metal stubs) and sputter-coated with iridium in an Emitech K575X (Emitech, Houston, TX) sputter

coater for 30 seconds at 20 mA, at the Electron Microscopy Laboratory, Department of Pathology, UTMB.

The samples were examined in a JEOL JSM-6330F Scanning electron microscope at the Texas Center for

Superconductivity, University of Houston, at 4 µA and 2 kV.

Zebrafish husbandry

All the protocols used for zebrafish experiments were approved by the University of Texas Medical Branch

Institutional Care and Use Committee. Adult and larval zebrafish were maintained using standard

husbandry procedures[86] at our in-house satellite facility. The wild type AB line was used for lifespan

analysis and Typhi load measurements and the *casper* line was used for visualizing mCherry-tagged STy

strains in infected larvae by confocal fluorescence imaging. Eggs were obtained by the natural spawning

method[86] and kept at 28 °C in embryo medium containing methylene blue for two days. At 2 days post-

fertilization (dpf), the larvae were shifted to an embryo medium containing 25 µg/ml gentamicin for 6 h,

after which they were transferred to a sterile embryo medium at 28 °C. Fresh embryo medium was replaced

daily.

Static immersions of zebrafish larvae

All infections were performed using mCherry-tagged STy strains, as listed in Supplementary table 2. For

static immersions, 1 ml of an overnight LB broth culture of mCherry-tagged STy strains was inoculated in

10 ml LB broth containing 100 µg/ml ampicillin and grown for 4.5 hours at 275 rpm at 37 °C. Growth was

normalized by measuring absorbance at 600 nm. The strains were harvested by centrifugation at 4,200 g

for 15 min at RT and resuspended in 500 µl sterile embryo medium. Ten larvae 5 dpf were added to 8 ml

embryo medium in a 6-well polystyrene plate, followed by the addition of 80 µl of harvested STy cultures

(10^9 cfu/ml). Equal infection doses across strains were verified using aliquots of the harvested STy cultures

by the total viable counting method. Larvae exposed to equal volume of PBS served as controls. After

24 h, the larvae were transferred to fresh embryo medium in a 6-well plate. Fresh embryo medium was

replaced daily. The larvae were fed once daily from 9 dpf, as recommended[59, 86]. Typically, infections were

performed with thirty to sixty larvae in each group.

Bacterial load estimates from infected larvae

For enumerating the STy load at 2, 4 and 6 dpi, infected larvae were isolated and added to embryo medium

containing 60 to 100 µg/ml buffered Tricaine. The anesthetized larva was transferred to a 1.5 ml Eppendorf

tube containing 200 µl PBS and homogenized well with a motorized micropestle. The homogenate was

serially diluted and colony forming units (cfu) were estimated by the total viable counting method on LB

agar plates containing 100 µg/ml ampicillin. For each time point, five larvae were used from each group.

PBS control larvae remained sterile. The experiment was repeated at least three times.

Lifespan analysis

Infected larvae were checked daily under a dissection microscope, and the percentage survival was scored.

Larvae that did not show any heart beats were considered dead.

Confocal fluorescence imaging

Five larvae from each group were withdrawn at 2, 4 and 6 dpi and anesthetized in an embryo medium

containing 60 to 100 µg/ml buffered Tricaine. These were then mounted on 35 mm glass-bottomed dishes

with a drop of 0.8% low melting point agarose (supplemented with 60 to 100 µg/ml buffered tricaine).

Images were acquired on an Olympus SpinSR-10 Yokogawa spinning disk confocal microscope fitted with

an ORCA Fusion sCMOS camera (Hamamatsu) using a 20x objective (NA 0.8, Olympus), 561 CSU and a

step size of 1.5 µ in the Z-dimension. Images were analyzed using Image J software.

Statistics

The Prism 10 software was used to plot all graphs and perform statistical analyses.

Acknowledgements:

S.D. and L.J.K. thank the following for their help; Stephen Baker (University of Cambridge) and Tran Vu

Thieu Nga for the H58 Tn library, Swine Chen, Kurosh Mehershahi and Varnica Khetrapal for initial

attempts at TraDIS at the Genome Institute of Singapore, Zhixia Ding (Department of Pathology, UTMB)

and Dezhi Wang (The Texas Center for Superconductivity, University of Houston) for SEM sample

preparations and image acquisitions and Roy Curtiss III and Soo-Young Wanda, University of Florida, for

the *rpoS* construct. S.D. and R.D. thank the Zebrafish International Resource Center (ZIRC) for fish lines

and April Freeman (ZIRC) for advice on fish husbandry. Y.Z. and A.L.R. were supported by a subaward

from U54 Center Grant, AI170855, NIH/NIAID. L.J.K. acknowledges grant support from UTMB Start up,

Texas Star Award and Cancer prevention and Research Institute of Texas (CPRIT) Grant, RP2000650.

1006

1007

1008

1009

1010

1011

1012

1013

1014

1015

1016

Supplementary Table 1: List of bacterial strains and plasmids

Strain or plasmid	Description	Source or Reference
14028s	*Salmonella enterica* serovar Typhimurium wild type strain	Lab collection
DW85	14028s (*ssrB*::*kan*)	Lab collection
H58 ('wild type' in this study)	*Salmonella enterica* serovar Typhi wild type strain	Stephen Baker, Cambridge University, UK
Ty2b	*Salmonella enterica* serovar Typhi wild type strain	Salmonella Genetic Stock Center, University of Calgary, CA
CT18	*Salmonella enterica* serovar Typhi wild type strain	Salmonella Genetic Stock Center, University of Calgary, CA
ct117	*Salmonella enterica* serovar Typhi wild type strain	Salmonella Genetic Stock Center, University of Calgary, CA
ssrB	H58 (*ssrB*::*kan*)	This study
csgD	H58 (*csgD*::*cat*)	This study
csgA	H58 (*csgA*::*kan*)	This study
yihO	H58 (*yihO*::*kan*)	This study
yihP	H58 (*yihP*::*kan*)	This study
yihO yihP	H58 (Δ*yihO yihP*::*kan*)	This study
iraP	H58 (*iraP*::*kan*)	This study
rpoS	H58 (*rpoS*::*kan*)	This study
sthC	H58 (*sthC*::*kan*)	This study
waaZ	H58 (*waaZ*::*kan*)	This study
tviA	H58 (*tviA*::*cat*)	This study
tviB	H58 (*tviD*::*kan*)	This study
rpoSc	H58 (*rpoS*::*kan*, pBR322::*rpoS*)	This study
pBR322::*rpoS* plasmid	*rpoS* cloned between EcoRI and ScaI sites in pBR322	Roy Curtiss III, University of Florida, USA
pFPV::mCherry plasmid	mCherry cloned between XbaI and SphI sites in pFPV	Olivia Steele-Mortimer (Addgene plasmid # 20956 http://n2t.net/addgene:20956; RRID:Addgene_20956

Supplementary Table 2: List of oligonucleotides

Primers	Sequence
Tn-ClickSeq	
3'21-39	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTTAATACGACTCACTATAGG [Black = Illumina p7 adapter, red = Tn-specific]
Gene deletions	

csgD knockout forward	GCAGCTGTCAGATGTGCGATTAAAAAAAGTGGAGTTTCATCATGTTTAATGTGTAGGCTGTAGCTGCTTC
csgD knockout reverse	CTCTGCTGCTACAATCCAGGTCAGATAGCGTTTCATGGCCTTACCGCCTGCCTCCTTAGTTCCTATTCCG
csgA knockout forward	CACCCAACGCTAATACCGTTACGACTTTTAAATCAATCAATCCGGTGTAGGCTGGAGCTGCTTC
csgA knockout reverse	AGGGCTTATGCCCTGTTTTTTTATTAGCGCAGACGCTAAACATATGAATATCCTCCTTAG
yihO knockout forward	ATGTCTAATCATGATCCGCTAACGCTAAAGTTGAGCCTGCGTGTAGGCTGGAGCTGCTTC
yihO knockout reverse	TTAATTATTTACAGTAGAAATACTTTGTTTATTATTAGTTCATATGAATATCCTCCTTAG
yihP knockout forward	GAGAAGAATAATGAGTCAAACATCTGTGTAGGCTGGAGCTGCTTC
yihP knockout reverse	GTTATATTTTATTGTTGTTAAACCGTATGCATATGAATATCCTCCTTA
iraP knockout forward	ATGAAAAATCTCATAGCAGAGTTGTTGCTTAAGCTAGCCCGTGTAGGCTGGAGCTGCTTC
iraP knockout reverse	TTAGTGCCGGGGGGTGTCTCAGCAACTTTTTTACATATTGGCATATGAATATCCTCCTTAG
rpoS knockout forward	ATGAGTCAGAATACGCTGAAAGTTCATGATTTAAATGAAGGTGTAGGCTGGAGCTGCTTC
rpoS knockout reverse	TTACTCGCGGAACAGCGCTTCGATATTCAGCCCCTGCGTCCATATGAATATCCTCCTTAG
tviA knockout forward	ATGAGGTTTCATCATTTCTGGCCTCCGAATGATATCTATTGTGTAGGCTGGAGCTGCTTC
tviA knockout reverse	TTACAGTAAAGTAACTGAATCCGGCAATAACAGATAGCGCCATATGAATATCCTCCTTAG
tviD knockout forward	ATGAATTTAATGAAATCGTCAGGGATGTTTACGCTTACAGGTGTAGGCTGGAGCTGCTTC
tviD knockout reverse	TTACGACTTCCCTGATGTATTTTTTTGTAATGCGGTTATGCATATGAATATCCTCCTTAG

waaZ knockout forward	ATGGGCAGCGTTAACTTCATAACTCACGCCGATGTTCTGC GTGTAGGCTGGAGCTGCTTC
waaZ knockout reverse	CTAGACAATTTTATCGTAATATTTCATCTTCAAGTTCCGA CATATGAATATCCTCCTTAG
sthC knockout forward	TTGATCGCCGACCCGCGAGAATTCGTCCCTACCAGCAAAT GTGTAGGCTGGAGCTGCTTC
sthC knockout reverse	TCACTGGCATTGCTCGTGTAAGATTTCTACGCCAGAAGCG CATATGAATATCCTCCTTAG
csgA forward	CACCCAACGCTAATACCGTT
csgA reverse	AGGGCTTATGCCCTGTTTTT
csgD forward	CAGCTGTCAGATGTGCGATT
csgD reverse	TCTGCTGCTACAATCCAGGT
yihO forward	TGCGGGGCGTTTTGAGAGGCGA
yihO reverse	TGAATAAAGCGGCAAGCGTCG
yihP forward	GTTGCGATTGGACGCTGTACCTG
yihP reverse	GTAATATGCAGGCATCCCGAGTTC
iraP forward	AGTGATAACGTCACCCTGGAAC
iraP reverse	AGTAACGTTATAACAACTGTGT
rpoS forward	CAGTCTGTCGACTGGCCTTT
rpoS reverse	CTAGTTCCGTCAAGGGATCA
tviA forward	AGGTTATTTCAGCATAAGGA
tviA reverse	TGTCCGTGTTTTACTCAATA
tviD forward	CTCGGTATAACTACTCACTT
tviD reverse	TTCCTAGTGCAGCTAACT
waaZ forward	GACATACTTGAGAGAATTTG
waaZ reverse	TGCGTGCCGAAGCAACGCAA
sthC forward	ACCGGCTGGATTTAGCGATC

sthC reverse	CATGATGCCAGACCCGTGAA	
RT-qPCR		
rrsA internal forward	GCACCGGCTAACTCCGTGCC	
rrsA internal reverse	GCAGTTCCCAGGTTGAGCCCG	
rpoS internal forward	CCTGCGTCTGGTGGTAAA	
rpoS internal reverse	TTCTCGACTGCACGGATAAG	
waaZ internal forward	CCGCTATTCAGGTTGCCTATTC	
waaZ internal reverse	AGGGCTGGTAGATTCGTCATAG	
sthC internal forward	GATGAAGACGACGATACGGAAG	
sthC internal reverse	CGTCTTTCGCGGAGTTCATA	
tviB internal forward	TGTGGTAAAGGAACTCGGTAAA	
tviB internal reverse	GACTTCCGATACCGGGATAATG	

1024
1025

1026

1027

1028

1029

1030

1031

1032

References

1. Dougan, G. and S. Baker, *Salmonella enterica serovar Typhi and the pathogenesis of typhoid fever.* Annu Rev Microbiol, 2014. **68**: p. 317-36.
2. Typhoid, G.B.D. and C. Paratyphoid, *The global burden of typhoid and paratyphoid fevers: a systematic analysis for the Global Burden of Disease Study 2017.* Lancet Infect Dis, 2019. **19**(4): p. 369-381.
3. Cirillo, D.M., et al., *Macrophage-dependent induction of the Salmonella pathogenicity island 2 type III secretion system and its role in intracellular survival.* Mol Microbiol, 1998. **30**(1): p. 175-88.
4. Feng, X., et al., *The response regulator SsrB activates transcription and binds to a region overlapping OmpR binding sites at Salmonella pathogenicity island 2.* Mol Microbiol, 2004. **54**(3): p. 823-35.
5. Walthers, D., et al., *The response regulator SsrB activates expression of diverse Salmonella pathogenicity island 2 promoters and counters silencing by the nucleoid-associated protein H-NS.* Mol Microbiol, 2007. **65**(2): p. 477-93.
6. Boddicker, J.D., et al., *Differential binding to and biofilm formation on, HEp-2 cells by Salmonella enterica serovar Typhimurium is dependent upon allelic variation in the fimH gene of the fim gene cluster.* Mol Microbiol, 2002. **45**(5): p. 1255-65.
7. Prouty, A.M., W.H. Schwesinger, and J.S. Gunn, *Biofilm formation and interaction with the surfaces of gallstones by Salmonella spp.* Infect Immun, 2002. **70**(5): p. 2640-9.
8. Crull, K., et al., *Biofilm formation by Salmonella enterica serovar Typhimurium colonizing solid tumours.* Cell Microbiol, 2011. **13**(8): p. 1223-33.
9. Miller, A.L., et al., *In vivo synthesis of bacterial amyloid curli contributes to joint inflammation during S. Typhimurium infection.* PLoS Pathog, 2020. **16**(7): p. e1008591.
10. Desai, S.K., et al., *The horizontally-acquired response regulator SsrB drives a Salmonella lifestyle switch by relieving biofilm silencing.* Elife, 2016. **5**.
11. Desai, S.K. and L.J. Kenney, *To ~P or Not to ~P? Non-canonical activation by two-component response regulators.* Mol Microbiol, 2017. **103**(2): p. 203-213.
12. Gerstel, U., C. Park, and U. Romling, *Complex regulation of csgD promoter activity by global regulatory proteins.* Mol Microbiol, 2003. **49**(3): p. 639-54.
13. Gerstel, U. and U. Romling, *The csgD promoter, a control unit for biofilm formation in Salmonella typhimurium.* Res Microbiol, 2003. **154**(10): p. 659-67.
14. Romling, U., et al., *Curli fibers are highly conserved between Salmonella typhimurium and Escherichia coli with respect to operon structure and regulation.* J Bacteriol, 1998. **180**(3): p. 722-31.
15. Zogaj, X., et al., *The multicellular morphotypes of Salmonella typhimurium and Escherichia coli produce cellulose as the second component of the extracellular matrix.* Mol Microbiol, 2001. **39**(6): p. 1452-63.
16. Gibson, D.L., et al., *Salmonella produces an O-antigen capsule regulated by AgfD and important for environmental persistence.* J Bacteriol, 2006. **188**(22): p. 7722-30.
17. Desai, S.K. and L.J. Kenney, *Switching Lifestyles Is an in vivo Adaptive Strategy of Bacterial Pathogens.* Front Cell Infect Microbiol, 2019. **9**: p. 421.
18. Gunn, J.S., et al., *Salmonella chronic carriage: epidemiology, diagnosis, and gallbladder persistence.* Trends Microbiol, 2014. **22**(11): p. 648-55.
19. Crawford, R.W., et al., *Gallstones play a significant role in Salmonella spp. gallbladder colonization and carriage.* Proc Natl Acad Sci U S A, 2010. **107**(9): p. 4353-8.
20. Basnyat, B. and S. Baker, *Typhoid carriage in the gallbladder.* Lancet, 2015. **386**(9998): p. 1074.
21. Wong, V.K., et al., *Phylogeographical analysis of the dominant multidrug-resistant H58 clade of Salmonella Typhi identifies inter- and intracontinental transmission events.* Nat Genet, 2015. **47**(6): p. 632-9.

22. Rowe, B., L.R. Ward, and E.J. Threlfall, *Multidrug-resistant Salmonella typhi: a worldwide*
epidemic. Clin Infect Dis, 1997. **24 Suppl 1**: p. S106-9.
23. Carey, M.E., et al., *The origins of haplotype 58 (H58) Salmonella enterica serovar*
Typhi. bioRxiv, 2022: p. 2022.10.03.510628.
24. Gomes, M.C. and S. Mostowy, *The Case for Modeling Human Infection in Zebrafish.* Trends
Microbiol, 2020. **28**(1): p. 10-18.
25. Torraca, V. and S. Mostowy, *Zebrafish Infection: From Pathogenesis to Cell Biology.* Trends Cell
Biol, 2018. **28**(2): p. 143-156.
26. Berg, R.D. and L. Ramakrishnan, *Insights into tuberculosis from the zebrafish model.* Trends Mol
Med, 2012. **18**(12): p. 689-90.
27. Ramakrishnan, L., *The zebrafish guide to tuberculosis immunity and treatment.* Cold Spring Harb
Symp Quant Biol, 2013. **78**: p. 179-92.
28. Varas, M., et al., *Live-cell imaging of Salmonella Typhimurium interaction with zebrafish larvae*
after injection and immersion delivery methods. J Microbiol Methods, 2017. **135**: p. 20-25.
29. Varas, M., et al., *Salmonella Typhimurium induces cloacitis-like symptomsin zebrafish larvae.*
Microb Pathog, 2017. **107**: p. 317-320.
30. Davis, J.M., et al., *Real-time visualization of mycobacterium-macrophage interactions leading to*
initiation of granuloma formation in zebrafish embryos. Immunity, 2002. **17**(6): p. 693-702.
31. Mostowy, S., et al., *The zebrafish as a new model for the in vivo study of Shigella flexneri*
interaction with phagocytes and bacterial autophagy. PLoS Pathog, 2013. **9**(9): p. e1003588.
32. Brannon, M.K., et al., *Pseudomonas aeruginosa Type III secretion system interacts with*
phagocytes to modulate systemic infection of zebrafish embryos. Cell Microbiol, 2009. **11**(5): p.
755-68.
33. Torraca, V., et al., *Shigella serotypes associated with carriage in humans establish persistent*
infection in zebrafish. J Infect Dis, 2023.
34. Leiba, J., et al., *Dynamics of macrophage polarization support Salmonella persistence*
in a whole living organism. bioRxiv, 2023: p. 2023.05.09.539693.
35. Desai, S.K., et al., *Salmonella biofilms program innate immunity for persistence in Caenorhabditis*
elegans. Proc Natl Acad Sci U S A, 2019. **116**(25): p. 12462-12467.
36. Ganjali Dashti, M., et al., *Optimization of Salmonella Typhi biofilm assay on polypropylene*
microtiter plates using response surface methodology. Biofouling, 2016. **32**(4): p. 477-87.
37. MacKenzie, K.D., et al., *Parallel evolution leading to impaired biofilm formation in invasive*
Salmonella strains. PLoS Genet, 2019. **15**(6): p. e1008233.
38. Crawford, R.W., et al., *Identification of a bile-induced exopolysaccharide required for Salmonella*
biofilm formation on gallstone surfaces. Infect Immun, 2008. **76**(11): p. 5341-9.
39. Goodall, E.C.A., et al., *The Essential Genome of Escherichia coli K-12.* mBio, 2018. **9**(1).
40. Canals, R., et al., *The fitness landscape of the African Salmonella Typhimurium ST313 strain*
D23580 reveals unique properties of the pBT1 plasmid. PLoS Pathog, 2019. **15**(9): p. e1007948.
41. Langridge, G.C., et al., *Simultaneous assay of every Salmonella Typhi gene using one million*
transposon mutants. Genome Res, 2009. **19**(12): p. 2308-16.
42. Holden, E.R., et al., *Massively parallel transposon mutagenesis identifies temporally essential*
genes for biofilm formation in Escherichia coli. Microb Genom, 2021. **7**(11).
43. Nolan, L.M., et al., *A global genomic approach uncovers novel components for twitching motility-*
mediated biofilm expansion in Pseudomonas aeruginosa. Microb Genom, 2018. **4**(11).
44. Jaworski, E., et al., *Tiled-ClickSeq for targeted sequencing of complete coronavirus genomes with*
simultaneous capture of RNA recombination and minority variants. Elife, 2021. **10**: p. e68479.
45. Routh, A., et al., *ClickSeq: Fragmentation-Free Next-Generation Sequencing via Click Ligation of*
Adaptors to Stochastically Terminated 3'-Azido cDNAs. J Mol Biol, 2015. **427**(16): p. 2610-6.
46. Tran, V.T.N., *The humoral response against Salmonella Typhi protein antigens during acute,*
convalescent, and chronic typhoid fever. 2019. p. xiv, 288 p.

47. Baumler, A.J., et al., *Contribution of horizontal gene transfer and deletion events to development*
of distinctive patterns of fimbrial operons during evolution of Salmonella serotypes. J Bacteriol,
1997. **179**(2): p. 317-22.
48. Baumler, A.J., R.M. Tsolis, and F. Heffron, *Fimbrial adhesins of Salmonella typhimurium. Role in*
bacterial interactions with epithelial cells. Adv Exp Med Biol, 1997. **412**: p. 149-58.
49. Frirdich, E., et al., *Overexpression of the waaZ gene leads to modification of the structure of the*
inner core region of Escherichia coli lipopolysaccharide, truncation of the outer core, and
reduction of the amount of O polysaccharide on the cell surface. J Bacteriol, 2003. **185**(5): p. 1659-
71.
50. Waxin, H., et al., *Identification of six open reading frames in the Salmonella enterica subsp.*
enterica ser. Typhi viaB locus involved in Vi antigen production. Res Microbiol, 1993. **144**(5): p.
363-71.
51. Pickard, D., et al., *Composition, acquisition, and distribution of the Vi exopolysaccharide-encoding*
Salmonella enterica pathogenicity island SPI-7. J Bacteriol, 2003. **185**(17): p. 5055-65.
52. Bougdour, A., S. Wickner, and S. Gottesman, *Modulating RssB activity: IraP, a novel regulator of*
sigma(S) stability in Escherichia coli. Genes Dev, 2006. **20**(7): p. 884-97.
53. Gottesman, S., *Trouble is coming: Signaling pathways that regulate general stress responses in*
bacteria. J Biol Chem, 2019. **294**(31): p. 11685-11700.
54. MacKenzie, K.D., et al., *Bistable Expression of CsgD in Salmonella Connects Virulence to*
Persistence. Infect Immun, 2015.
55. Anwar, N., et al., *Modulation of biofilm-formation in Salmonella enterica serovar Typhimurium by*
the periplasmic DsbA/DsbB oxidoreductase system requires the GGDEF-EAL domain protein
STM3615. PLoS One, 2014. **9**(8): p. e106095.
56. Gallo, P.M., et al., *Amyloid-DNA Composites of Bacterial Biofilms Stimulate Autoimmunity.*
Immunity, 2015. **42**(6): p. 1171-84.
57. Santander, J., K.L. Roland, and R. Curtiss, 3rd, *Regulation of Vi capsular polysaccharide synthesis*
in Salmonella enterica serotype Typhi. J Infect Dev Ctries, 2008. **2**(6): p. 412-20.
58. Monack, D.M., D.M. Bouley, and S. Falkow, *Salmonella typhimurium persists within macrophages*
in the mesenteric lymph nodes of chronically infected Nramp1+/+ mice and can be reactivated by
IFNgamma neutralization. J Exp Med, 2004. **199**(2): p. 231-41.
59. Hernandez, R.E., et al., *Delay of Initial Feeding of Zebrafish Larvae Until 8 Days Postfertilization*
Has No Impact on Survival or Growth Through the Juvenile Stage. Zebrafish, 2018. **15**(5): p. 515-
518.
60. Marineli, F., et al., *Mary Mallon (1869-1938) and the history of typhoid fever.* Ann Gastroenterol,
2013. **26**(2): p. 132-134.
61. Mortimer, P.P., *Mr N the milker, and Dr Koch's concept of the healthy carrier.* Lancet, 1999.
353(9161): p. 1354-6.
62. Gopinath, S., S. Carden, and D. Monack, *Shedding light on Salmonella carriers.* Trends Microbiol,
2012. **20**(7): p. 320-7.
63. Crawford, R.W., K.E. Reeve, and J.S. Gunn, *Flagellated but not hyperfimbriated Salmonella*
enterica serovar Typhimurium attaches to and forms biofilms on cholesterol-coated surfaces. J
Bacteriol, 2010. **192**(12): p. 2981-90.
64. Adcox, H.E., et al., *Salmonella Extracellular Matrix Components Influence Biofilm Formation and*
Gallbladder Colonization. Infect Immun, 2016. **84**(11): p. 3243-3251.
65. Gonzalez, J.F., et al., *Human Bile-Mediated Regulation of Salmonella Curli Fimbriae.* J Bacteriol,
2019. **201**(18).
66. Shimada, T., K. Tanaka, and A. Ishihama, *The whole set of the constitutive promoters recognized*
by four minor sigma subunits of Escherichia coli RNA polymerase. PLoS One, 2017. **12**(6): p.
e0179181.
67. Santander, J., et al., *Role of RpoS in fine-tuning the synthesis of Vi capsular polysaccharide in*
Salmonella enterica serotype Typhi. Infect Immun, 2007. **75**(3): p. 1382-92.

68. Marshall, J.M., et al., *Visualization of extracellular matrix components within sectioned Salmonella*
biofilms on the surface of human gallstones. PLoS One, 2014. **9**(2): p. e89243.
69. Devaraj, A., et al., *Enhanced biofilm and extracellular matrix production by chronic carriage*
versus acute isolates of Salmonella Typhi. PLoS Pathog, 2021. **17**(1): p. e1009209.
70. Hahn, M.M., et al., *The Abundance and Organization of Salmonella Extracellular Polymeric*
Substances in Gallbladder-Mimicking Environments and In Vivo. Infect Immun, 2021. **89**(11): p.
e0031021.
71. Milligan-Myhre, K., et al., *Study of host-microbe interactions in zebrafish.* Methods Cell Biol,
2011. **105**: p. 87-116.
72. Raza, A., et al., *Effect of biofilm formation on the excretion of Salmonella enterica serovar Typhi*
in feces. Int J Infect Dis, 2011. **15**(11): p. e747-52.
73. Sambrook J, F.E., Maniatis T, *Molecular cloning: a laboratory manual*. 1989: Cold Spring Harbor
Laboratory Press, Cold spring Harbor, NY.
74. Davis R, B.D., Roth J, *Advanced Bacterial Genetics*. 1980: Cold spring Harbor Laboratory, NY.
16-18.
75. Karlinsey, J.E., *lambda-Red genetic engineering in Salmonella enterica serovar Typhimurium.*
Methods Enzymol, 2007. **421**: p. 199-209.
76. Jaworski, E. and A. Routh, *ClickSeq: Replacing Fragmentation and Enzymatic Ligation with Click-*
Chemistry to Prevent Sequence Chimeras, in *Next Generation Sequencing: Methods and Protocols*,
S.R. Head, P. Ordoukhanian, and D.R. Salomon, Editors. 2018, Springer New York: New York,
NY. p. 71-85.
77. Jaworski, E. and A. Routh, *Parallel ClickSeq and Nanopore sequencing elucidates the rapid*
evolution of defective-interfering RNAs in Flock House virus. PLoS Pathog, 2017. **13**(5): p.
e1006365.
78. Chen, S., et al., *fastp: an ultra-fast all-in-one FASTQ preprocessor.* Bioinformatics, 2018. **34**(17):
p. i884-i890.
79. Martin, M., *Cutadapt removes adapter sequences from high-throughput sequencing reads.*
EMBnet.journal; Vol 17, No 1: Next Generation Sequencing Data AnalysisDO -
10.14806/ej.17.1.200, 2011.
80. Kim, D., et al., *Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype.*
Nat Biotechnol, 2019. **37**(8): p. 907-915.
81. Li, H., et al., *The Sequence Alignment/Map format and SAMtools.* Bioinformatics (Oxford,
England), 2009. **25**(16): p. 2078-2079.
82. Smith, T., A. Heger, and I. Sudbery, *UMI-tools: modeling sequencing errors in Unique Molecular*
Identifiers to improve quantification accuracy. Genome Res, 2017. **27**(3): p. 491-499.
83. Quinlan, A.R. and I.M. Hall, *BEDTools: a flexible suite of utilities for comparing genomic features.*
Bioinformatics, 2010. **26**(6): p. 841-2.
84. Love, M.I., W. Huber, and S. Anders, *Moderated estimation of fold change and dispersion for*
RNA-seq data with DESeq2. Genome Biol, 2014. **15**(12): p. 550.
85. Schmittgen, T.D. and K.J. Livak, *Analyzing real-time PCR data by the comparative C(T) method.*
Nat Protoc, 2008. **3**(6): p. 1101-8.
86. Westerfield, M., *The zebrafish book. A guide for the laboratory use of zebrafish (Danio rerio)*. 4th
ed ed. 2000, Eugene: Univ. of Oregon Press.
1227