RpoS activates formation of *Salmonella* Typhi biofilms and drives persistence in the gall bladder

Stuti K. Desai[1*], Yiyang Zhou[1], Rahul Dilawari[1], Andrew L. Routh[1,2,3,5],

Vsevolod Popov[3,4], and Linda J. Kenney[1,2,3*]

[1]Department of Biochemistry and Molecular Biology, [2]Sealy Center for Structural Biology and Molecular Biophysics, [3]Institute for Human Infections and Immunity, and [4]Department of Pathology, University of Texas Medical Branch, Galveston, TX 77555, and [5]Department of Immunology and Microbiology, Scripps Research, 10550 N. Torrey Pines Rd., La Jolla, CA 92037, USA

[*]Address correspondence to: stdesai@utmb.edu or likenney@utmb.edu

Keywords: Static immersions, Transposon directed insertion site sequencing, ClickSeq, Asymptomatic Typhoid carriage, IraP, Fimbriae, Vi polysaccharide, CsgD

Abbreviations: red, dry and rough (rdar); colony forming unit (CFU); room temperature (RT); hour (h); Polymerase chain reaction (PCR); Reverse transcription quantitative real-time PCR (RT-qPCR); Transposon (Tn); next-generation sequencing (NGS); two-component regulatory system (TCRS)

Abstract

The development of strategies for targeting the asymptomatic carriage of *Salmonella* Typhi in chronic typhoid patients has suffered owing to our basic lack of understanding of the molecular mechanisms that enable the formation of *S.* Typhi biofilms. Traditionally, studies have relied on cholesterol-attached biofilms formed by a closely related serovar, Typhimurium, to mimic multicellular Typhi communities formed on human gallstones. In long-term infections, *S.* Typhi adopts the biofilm lifestyle to persist in vivo and survive in the carrier state, ultimately leading to the spread of infections via the fecal-oral route of transmission. In the present work, we studied *S.* Typhi biofilms directly, applied targeted as well as genome-wide genetic approaches to uncover unique biofilm components that do not conform to the CsgD-dependent pathway established in *S.* Typhimurium. We undertook a genome-wide *Tn5* mutation screen in H58, a clinically relevant multidrug resistance strain of *S.* Typhi, in gallstone-mimicking conditions. We generated New Generation Sequencing libraries based on the ClickSeq technology to identify the key regulators, IraP and RpoS, and the matrix components Sth fimbriae, Vi capsule and lipopolysaccharide. We discovered that the starvation sigma factor, RpoS, was required for the transcriptional activation of matrix-encoding genes in vitro, and for *S.* Typhi colonization in persistent infections in vivo, using a heterologous fish larval model. An *rpoS* null mutant failed to colonize the gall bladder in chronic zebrafish infections. Overall, our work uncovered a novel RpoS-driven, CsgD-independent paradigm for the formation of cholesterol-attached Typhi biofilms, and emphasized the role(s) of stress signaling pathways for adaptation in chronic infections. Our identification of the biofilm regulators in *S.* Typhi paves the way for the development of drugs against typhoid carriage, which will ultimately control the increased incidence of gall bladder cancer in typhoid carriers.

Introduction

Salmonella enterica is a rod-shaped enteric bacterium that easily spreads through contaminated food or water via the fecal-oral route in poor hygiene conditions. The human-restricted serovar of *Salmonella enterica* serovar Typhi (STy), causes typhoid fever and continues to be a dangerous pathogen throughout the world[1]. The global incidence of typhoid fever in mostly children, adolescents and older

adults is between 12 to 27 million and in total 116,815 succumbed to the disease in 2017[2]. In contrast, its

closely related serovar Typhimurium (STm) infects diverse hosts such as humans, cattle, poultry and

reptiles to cause gastroenteritis that is mostly self-limiting in healthy adults.

Upon successful invasion of intestinal epithelial cells, *Salmonella* is phagocytosed by

macrophages, where it resides in a modified vacuole in a self-nourishing niche called a *Salmonella-*

<u>C</u>ontaining <u>V</u>acuole (SCV) to ultimately reach the systemic sites of liver, spleen and the gall bladder. The

SsrA/B <u>T</u>wo-<u>C</u>omponent <u>R</u>egulatory <u>S</u>ystem (TCRS) is essential for activation of the *Salmonella*

<u>P</u>athogenicity <u>I</u>sland-2 (SPI-2) regulon genes encoding a type-three secretory system and effectors that are

involved in formation of the SCV[3-5]. *Salmonella* also resides encased in extracellular matrix as

multicellular communities or biofilms, on intestinal epithelial cells[6], gallstones[7], tumors[8] and in the large

intestine[9].

Interestingly, SsrB, a response regulator of the SsrA/B TCRS, is essential for switching on the

multicellular lifestyle of *S.* Typhimurium by relieving H-NS silencing at the *csgD* promoter[10, 11]. CsgD is

the master regulator of STm biofilms, and it activates the transcription of extracellular matrix components

including curli fimbriae, cellulose*,* BapA and O-Antigen[12-16]. Moreover, the SsrB-CsgD regulatory

pathway drives STm persistence in the heterologous host *Caenorhabditis elegans,* by enabling the

formation of biofilms, which eventually promotes host life span through the p38-<u>M</u>itogen-<u>a</u>ctivated Protein

<u>K</u>inase (p38-MAPK) innate immunity pathway[17].

Biofilms in the gall bladder are important for maintaining the carrier state of *Salmonella* Typhi,

allowing it to persist in 2 to 4% of chronic typhoid patients[18-20]. STy reservoirs in human carriers, who are

characteristically asymptomatic, play a crucial role in the spread of typhoid in endemic regions, as well as

in its introduction to non-endemic regions. Indeed, such long-term colonization of STy in human carriers,

coupled with the rise of multi-drug resistant strains, for example, those belonging to the H58 haplotype,

prevents effective control of typhoid fever [21-23]. Unfortunately, all of our understanding of the regulation

of STy biofilms on gallstones has been based on the assumption that it shares a high degree of conservation

with canonical biofilms formed by the closely related serovar, *S.* Typhimurium[18, 24].

Herein, we establish that the components of STy biofilms are fundamentally distinct from STm biofilms. In particular, CsgD and the STm lifestyle regulator, SsrB, are not required for formation of multicellular aggregates on surfaces coated with cholesterol that forms the major component of human gallstones. In order to identify unique components of STy biofilms, we employed a whole genome transposon mutagenesis approach, Tn-ClickSeq, and identified the starvation sigma factor RpoS as a crucial determinant of STy biofilms. We discovered that the formation of large STy aggregates was defective in the absence of *rpoS*, owing to the down-regulation of extracellular matrix components comprised of Sth fimbriae, the Vi polysaccharide and the lipopolysaccharide core.

Finally, we developed a heterologous host model, *Danio rerio*, to investigate STy lifestyles in persistent infections for visualizing the colonization in real-time by confocal microscopy and measuring the ensuing effects on host physiology. Zebrafish is a powerful vertebrate model for many human diseases owing to a high degree of conservation of immuno-signaling pathways and colonization characteristics of bacterial infections[25-28]. Previous studies have established that exposure of zebrafish larvae to *S.* Typhimurium, *Mycobacterium marinum*, *Shigella flexneri*, and *Pseudomonas aeruginosa* leads to successful pathogenesis[29-33]. *Shigella sonnei* and *S.* Typhimurium have also been recently shown to persistently colonize macrophages in zebrafish[34, 35]. We infected zebrafish larvae with STy using static immersions. Intestinal colonization was diminished, and larval survival was greater in the *rpoS* null, corroborating a crucial role of RpoS in prolonged STy survival in vivo. Ultimately, we harnessed the immense imaging potential of the zebrafish model to reveal the absolute requirement of RpoS in enabling chronic gall bladder colonization.

Results

***S.* Typhi biofilms employ unique components that differ from *S.* Typhimurium biofilms**

S. Typhi (STy) forms biofilms on gallstones in the gall bladder and this ability is an important aspect of maintaining its carrier state[18-20]. It was therefore of interest to examine whether STy employed similar pathways as STm for biofilm formation. We grew the STy strain H58 under conditions that were proposed to mimic the gall bladder environment in humans[36], hereafter referred to as gallstone-mimicking

conditions, and observed robust surface-attached communities after two days in cholesterol-coated tubes,

as measured by a crystal violet staining assay (Fig. 1A). The requirement of such distinct physico-chemical

factors to develop biofilms was not specific to H58, as three other isolates including Ty2-b, CT18 and

CT117 formed comparable cholesterol-attached biomass after two days of growth (Fig. 1A). Not

surprisingly, H58 failed to form the characteristic rough, dry and red ('rdar') morphotype as classically

observed for the STm wild type strain 14028s (Supplementary Fig. 1A). This failure to develop rdar

colonies has also been reported for other STy strains[37, 38]. Similarly, crystal violet staining of static biofilms

formed in the standard '*Salmonella*-conditions', of 30°C and low osmolality, demonstrated that H58 formed

extremely poor biofilms when compared to the STm strain 14028s (Supplementary Fig. 1B). We followed

the developmental course of H58 biofilm formation at two, four and six days in cholesterol-coated tubes

and observed only a marginal increase in the amount of biofilms at days 4 and 6 compared to day 2

(Supplementary Fig. 1C). We therefore focused on understanding STy biofilms at day 2 in our further

investigations.

The response regulator SsrB plays a dual role in regulating *S.* Typhimurium lifestyles. For

virulence, SsrB~P activates SPI-2 genes, while the unphosphorylated form drives the formation of biofilms

by de-repressing *csgD*[10, 11 for a review]. Hence, we examined whether complete deletions of *ssrB* and *csgD*

affected the ability of *S.* Typhi to form cholesterol-attached biofilms. Interestingly, biofilm formation was

not affected by the loss of STy homologs encoding either SsrB, the STm lifestyle regulator, or CsgD, the

master regulator of STm biofilms, emphasizing the fundamental differences in mechanisms of biofilm

formation in these two closely related serovars (Fig. 1B). We also investigated whether other requirements

for STm biofilms: *csgA* (thin aggregative curli fimbriae), and *yihO*/*P* adhesion (O-Antigen) were required

for STy aggregates to cholesterol-coated surfaces[13, 14, 39, 40]. We discovered that curli fibers and O-Antigen

did not play any role in the formation of cholesterol-attached STy biofilms, strongly establishing that STy

biofilms are distinct from the model STm biofilms (Fig. 1B and see Discussion).

Tn-ClickSeq reveals the unique genetic signature of STy biofilms

Transposon-directed insertion site sequencing (TraDIS) has been immensely useful to study the

genetic repertoires essential for growth in STy and *Escherichia coli*, evolution of the invasive STm lineage

ST313, and biofilm formation in *E. coli* and *Pseudomonas aeruginosa*[41-45]. We applied a similar approach,

Tn-ClickSeq, by combining genome-wide transposon (Tn) mutagenesis with ClickSeq. ClickSeq is

advantageous here, since it is a fragmentation-free next-generation sequencing (NGS) library synthesis

technique and is capable of generating focused NGS read data upstream of chosen target sites[46, 47]. These

features greatly simplify the digital transposon display protocol and remove artifactual recombination

events inherent to common NGS library preparation techniques. Using primers targeting the 3' or 5' ends

of inserted transposons, Tn-ClickSeq can sensitively and specifically sequence the junctions of a transposon

and the adjacent genomic loci of the integration site and thus identify genetic loci involved in forming

cholesterol-attached biofilms in vitro (Fig. 2A and Supplementary Fig. 2A). A Tn-library in the *S.* Typhi

strain H58 was kindly provided by Stephen Baker, Cambridge University, UK[48 and see Methods]. We grew *S.*

Typhi biofilms for two days using the H58-Tn library in gallstone-mimicking conditions and isolated

planktonic and biofilm fractions from a pool of thirty cholesterol-coated tubes. NGS libraries were

generated from genomic DNA isolated from each of these sub-populations using the Tn-ClickSeq

approach[47] (Fig. 2A).

Since Tn-ClickSeq returns short sequence reads containing a short fragment of the 3' or 5' end of

the inserted (known) Tn as well as a fragment of the adjacent genomic DNA, we developed a simple

computational pipeline (Supplementary Fig. 2A) that identified and trimmed the Tn-derived sequences from

individual reads, and then mapped the remaining fragment to the H58 genome (BioSample ID -

SAMEA3110714) using a HISAT2 program[49]. The 3' end of these Tn-mapping reads represents the exact

nucleotide junction of the insertion site of the Tn in the genome. With this approach, we identified

transposon insertions as 47%, 52% and 20% per million raw reads in the input, planktonic and biofilm sub-

populations, respectively (Supplementary Fig. 2B, C and D). Mapping of insertion sites returned genomic

locations of the inserted Tn in each dataset, as well as the frequency of these inserts within each original

sample. After controlling for PCR duplication using unique molecular identifiers (UMIs) included in the

'Click-Adaptor'[46], we assigned insertion indices by dividing the number of Tn-insertions per gene with 1

Kbp of gene length (TnClickSeq insertion indices.xlsx). With this approach, gene insertion frequencies in

each condition revealed clear differences in the planktonic and biofilm fractions, indicating unique genetic

components driving H58 lifestyles in gallstone-mimicking conditions (Fig. 2B). Principal Component

Analysis (Fig. 2C) and hierarchical clustering analysis (Fig. 2D) returned distinct clusters of genome

insertion sites of the three replicates of Tn-ClickSeq libraries obtained from the input, planktonic and

biofilm sub-populations. Overall, we successfully adopted the Tn-ClickSeq approach in *S.* Typhi to

generate a comprehensive view of genetic systems that determine the development of cholesterol-attached

biofilms from unattached planktonic cells.

Identifying matrix components and regulators of STy biofilms

To determine the exact mechanism(s) by which STy forms multicellular communities on

cholesterol surfaces in gallstone-mimicking conditions, we compared insertion indices between the

planktonic and biofilm fractions. Transposon insertions in biofilm genes likely disrupt functions resulting

in the inability of such mutants to form biofilms and would be enriched in the planktonic sub-population.

Interestingly, the majority of insertion indices had drastically lower values in the biofilm fraction compared

to the planktonic group, these were of immediate interest for validation as STy biofilm targets (TnClickSeq

insertion indices.xlsx). We focused on loci having mean insertion indices greater than 100 in the planktonic

sub-population and performed Gene Ontology analysis on 1,515 such genes. We identified a significant

enrichment in cell membrane components, transmembrane ion transport pathways, and other membrane-

related activities (Supplementary Fig. 3). However, it was not possible to perform essentiality analysis using

a standard bioinformatics pipeline, because most of these genes had insertion indices equal to zero in the

biofilm fraction (see Methods and Discussion). In the absence of any *a priori* list of 'essential' biofilm

components, we narrowed our list of 1,515 genes to the top 300 and sought to investigate some for their

roles in STy biofilms. It was noteworthy that genes/operons encoding STm 'biofilm' homologs; *ssrB*,

csgDEFG, *csgBAC* or *yihPO* did not appear on our list of selected Tn-ClickSeq targets. This established a

strong correlation of our targeted genetic approach (Figure 1B), to our whole genome transposon mutagenesis Tn-ClickSeq approach (Fig. 2A).

We next generated precise deletions of selected loci identified by Tn-ClickSeq that might encode the structural components of STy biofilms in our gallstone-mimicking conditions: *sthC*, part of the Typhi-specific *sth* fimbrial operon[50, 51], *waaZ*, encoding an enzyme involved in the biosynthesis of the LPS core forming the outer membrane[52] and *tviD*, a part of the *tviBCDE* operon encoding the surface-exposed Vi polysaccharide (which is not encoded in STm); TviA is the regulator of the pathway[53, 54]. We performed crystal violet staining assays at day 2 and determined that the loss of *sthC*, *waaZ, tviA* and *tviD* led to a decrease in *S*. Typhi biofilms (Fig. 3A). The double mutant strains, *sthC tviD* and *waaZ sthC*, showed a similar reduction, of around 50%, as compared to the wild type parent in forming cholesterol-attached biofilms. This result indicated a possible functional redundancy or an inter-dependency in biosynthesis in the mature STy biofilms (Supplementary Fig. 4A). For example, an interplay between O-Antigen and K2 capsule synthesis has been recently described in extra-intestinal pathogenic *E. coli* for conferring serum resistance[55].

Intriguingly, the gene *iraP* rated highly on our insertion index list (insertion index = 670). It encodes an anti-adapter that inhibits the ClpXP-dependent proteolysis of the starvation master regulator RpoS[56]. We tested the biofilm-forming capabilities of an *iraP* null H58 strain and observed a substantial reduction in biofilms compared to the wild type parent at 2 days (Fig. 3B). The only known role of IraP is an indirect positive role in activating the RpoS-dependent starvation stress response in enteric bacteria[57], yet the *rpoS* insertion index was substantially lower than *iraP* (25). This was below our arbitrary threshold for selecting Tn-ClickSeq targets (insertion index > 100), thus we had excluded *rpoS* from our Gene Ontology analysis (TnClickSeq insertion indices.xlsx). It was therefore of interest to examine whether deletion of *rpoS* affected *S*. Typhi biofilms. Indeed, the *rpoS* null strain was inhibited in the formation of cholesterol-attached STy biofilms by over 50% (Fig. 3B). Over-expression of RpoS *in trans* (a kind gift from Roy Curtiss III, University of Florida), complemented the defect in biofilm formation of the *rpoS* null strain (Fig. 3B) and increased biofilms to a level similar to the wildtype. Our discovery of RpoS as an

activator of *S.* Typhi biofilms was significant and highlighted the divergence between *S.* Typhi and *S.*

Typhimurium biofilms. In in *S.* Typhimurium, RpoS impacts biofilms by effects on the master regulator

CsgD, whereas *S.* Typhi biofilms are CsgD-independent. (Fig. 1B). Finally, we ruled out an effect of mere

growth differences on biofilm formation in the STy mutants by monitoring the planktonic growth of *iraP*,

rpoS, *sthC*, *waaZ* and *tviD* null mutants compared to the wild type H58 parent. We observed similar growth

rates among all the strains, indicating that IraP, RpoS, SthC, WaaZ and TviD were necessary for the

formation of *S.* Typhi surface-attached communities (Supplementary Fig. 4B).

Cholesterol-attached aggregates of STy biofilm mutants exhibit a poor ultrastructure

Our understanding of the CsgD-dependent mechanisms of biofilm formation in *S.* Typhimurium

has been achieved by screening mutant strains for their inability to exhibit the rdar morphotype and high-

resolution imaging of biofilm components by fluorescence confocal microscopy[10, 37, 58-61]. Unfortunately,

neither of these approaches can be applied to unequivocally establish the molecular players that drive the

formation of STy biofilms on cholesterol-coated surfaces because of a high level of autofluorescence in the

gallstone-mimicking conditions (see above and Discussion). We therefore grew STy biofilms in

cholesterol-coated tubes for two days and obtained direct visualization of surface-attached communities by

scanning electron microscopy. We observed dense aggregates of the wild type parent having a rich network

of extracellular matrix that connected the cells, while the strains defective in *rpoS*, *sthC*, *waaZ* and *tviD*

formed smaller STy aggregates with poor network connections (Fig. 4). Our high-resolution observations

of cholesterol-attached STy communities by scanning electron microscopy consolidated our in vitro

measurements of the sessile biomass formed by the STy strains by the standard crystal violet staining assay

(Fig. 3). Overall, we demonstrated that the Tn-ClickSeq approach enabled the identification of previously

unidentified players that drive the formation of atypical STy biofilms.

RpoS activates transcription of STy biofilm matrix genes

We next wanted to determine the possible role of RpoS in regulating the expression of STy biofilm

components, including: SthC, WaaZ and the Vi capsule. We first investigated if the expression of *rpoS*,

sthC, *waaZ* and *tviB* (the *tviBCDE* operon mediates biosynthesis of Vi-polysaccharide[62]), were up-

regulated in biofilm-inducing conditions. We isolated planktonic and biofilm sub-populations from the wild

type strain grown in cholesterol-coated tubes and measured steady state transcript levels by reverse

transcription quantitative real-time PCR (RT-qPCR). Indeed, we detected an increase in the transcription

of *rpoS* (~20-fold), *sthC* (~4-fold), *waaZ* (~200-fold) and *tviB* (~10-fold) in biofilms formed by the H58

parent in gallstone-mimicking conditions (Fig. 5A).

In order to investigate the exact role of RpoS in driving the biofilm lifestyle, we grew the wild type

parent H58 and its *rpoS* null derivative in cholesterol-coated tubes in vitro and isolated total RNA from the

biofilm fractions after two days. We then compared the transcription of matrix-encoding genes *waaZ*, *sthC*

and *tviB* by RT-qPCR and observed a positive effect of RpoS at each of these loci (Fig. 5B). Transcription

of the matrix-encoding genes was clearly down-regulated when RpoS was absent, ~ 60% for *waaZ*, ~40%

for *tviB* and ~20% for *sthC* (Fig. 5B). This result established a molecular link between the stress sigma

factor, RpoS, and development of STy cholesterol-attached biofilms.

Further, since the binding sites for RpoS have been previously mapped for several stress responsive

genes in *E. coli*[63, 64], we analyzed the regulatory regions of *waaZ*, *sthC* and *tviA* (the regulator of the

tviBCDE operon) in silico and found a high degree of conservation of classical RpoS recognition sequences

for *waaZ* and *tviA* and a partial conservation for *sthC* (Supplementary Fig. 5A). Interestingly, binding of

RpoS to the *waaZ* promoter has also been reported in *E. coli*[65]. To investigate direct binding, we amplified

biotinylated fragments of these regulatory regions and tested the formation of DNA-protein complexes by

electrophoretic mobility shift assays using purified His-tagged RpoS[66]. We detected RpoS binding by

visualizing slower migrating DNA-protein complexes of *waaZ*, *sthC*, *tviA* and *tviB* (Fig. 5C). Binding was

specific, as confirmed by a decrease in the amount of DNA-protein complexes formed in the presence of

excess non-biotinylated promoter fragments as competitor DNA. Thus, RpoS directly binds and activates

the formation of extracellular matrix in gallstone-mimicking conditions. RpoS binding affinity may be

enhanced by the adapter protein, Crl (insertion index = 154), which was also present in our total of 1,515

Tn-ClickSeq targets (Supplementary Fig. 3; TnClickSeq insertion indices.xlsx)(see Discussion). A

proposed global role of RpoS in enabling lifestyle transitions in STy is also supported by the enrichment of

Tn insertions in several RpoS-regulated genes such as *gadC*, *osmB*, *bolA*, *dps*, *mscL, uspA, uspB* and *bfr*[63,

64] in the planktonic fraction as compared to the biofilm fraction (Supplementary Fig. 5B and Discussion).

RpoS regulates long-term infection outcomes in zebrafish larvae

Since the formation of STy biofilms plays a crucial role in enabling persistence in human carriers,

we investigated the role of RpoS in persistent infections in a zebrafish model. We replicated the natural

mode of Typhi entry by employing static immersions of zebrafish larvae in tank water contaminated with

specific doses of *S.* Typhi strains. We exposed zebrafish larvae 5 days post-fertilization (dpf) to 10^8 cfu/ml

of mCherry-expressing wild type and the isogenic *rpoS* null mutant in system water for 24 h. After 24 h,

the larvae were shifted to plain system water and monitored for six days under standard conditions of

zebrafish husbandry[adapted from 30]. Control larvae were exposed to an equal volume of PBS. We discovered

that *S.* Typhi colonization of infected larvae was reduced when RpoS was absent, as evident from both

bacterial load measurements (Fig. 6A), as well as real-time visualizations at 2, 4 and 6 dpi using confocal

fluorescence microscopy (Fig. 6B). At longer times of 4 and 6 dpi, wild type *S.* Typhi clearly persisted in

the intestine (Fig. 6A, B), as also observed in a mouse model of STm persistence[67]. Finally, persistence

was strongly correlated with pathogenesis outcomes, as larvae infected with the *rpoS* null strain survived

infections at 6 dpi in significantly greater numbers than the wild type (Fig. 6C). The percentage survival of

larvae infected with an *rpoS* mutant was similar to the uninfected PBS control from 1 to 6 dpi, strongly

emphasizing the requirement of RpoS for enabling STy survival in long term infections. Monitoring

infected larvae beyond 6 dpi is challenging, owing to a greater requirement of live food (Paramecia), which

masks pathogen-driven physiological effects[68 and see Discussion]. In conclusion, zebrafish infections revealed a

role for RpoS in chronic STy infections and for persistence in vivo.

RpoS is essential for STy to persist in the gall bladder

We were then able to address the ultimate question: was RpoS necessary to colonize the

hepatobiliary system? This is of importance because adaptation of STy to the gall bladder niche forms the

basis for long-term transmission in humans, and increases the risks of developing hepatobiliary carcinomas.

We infected zebrafish larvae with mCherry-tagged wild type H58 and *rpoS* null strains via static

immersions at 5 dpf and performed whole-mount immunohistochemistry against mCherry at 6 dpi. We

observed the gall bladder region by confocal fluorescence microscopy and detected anti-mCherry antibody

staining in chronic infections of the H58 parent (Fig. 7, middle right zoomed-in image). Uninfected larvae

were clear of any fluorescent signals in the hepatobiliary region and served as negative controls.

Remarkably, we did not detect the presence of STy in the hepatobiliary system in infections using the *rpoS*

null strain, emphasizing an essential role of RpoS in enabling persistent gall bladder colonization. The *rpoS*

null strain was poorly visible in the intestine at 6 dpi, in complete agreement with results described above

(Fig. 6A and B), further emphasizing the essentiality of RpoS signaling in hepatobiliary persistence. Since

the Vi-polysaccharide capsule is also a component of the STy biofilm matrix (Fig. 4), we also performed

whole-mount immunohistochemistry against the Vi polysaccharide at 6 dpi and detected clusters of

bacteria, presumably in vivo STy communities, in the hepatobiliary region of zebrafish larvae infected with

the wild type parent (Supplementary Fig. 6). Overall, this work provides an excellent tool, the zebrafish

persistent infection model, for future investigations to unravel the exact nature of STy lifestyles in the gall

bladder.

Discussion

The dangerous transmission of *S.* Typhi from seemingly healthy, but chronically infected

individuals is well documented, and the first cases in the United States of Mary Mallon, 'Typhoid Mary',

and Mr. N in the United Kingdom, were reported in the early 1900s[69, 70]. Since then several epidemiological

studies have established that persistent colonization of STy in the gall bladders of asymptomatic patients

forms the basis for typhoid carriage[reviewed in 18, 24, 71]. Despite clear evidence of the role of STy biofilms in

spreading the disease, previous studies have failed to delineate any genetic mechanisms that regulate the

development of gallstone biofilms in STy. This is chiefly because biofilms formed by *S.* Typhimurium have

been employed as a surrogate for understanding the multicellular behavior of *S.* Typhi[72-74]. In the present

work, we studied STy biofilms directly, and discovered that the pathways for biofilm formation in STy and

STm were distinct. Our observations that STy failed to form biofilms in the standard laboratory conditions

of low osmolality and lower temperatures (RT or 30°C) routinely employed for studying STm biofilms

were in clear agreement with previous studies that observed an inability of typhoidal strains to exhibit the

rdar morphotype, a hallmark of biofilm capability in the non-typhoidal *Salmonella* strains[37, 38]. More

importantly, CsgD, the master regulator of STm surface-attached communities, was entirely dispensable

for the formation of STy sessile aggregates on cholesterol-coated surfaces. As a result, matrix production

in STy biofilms did not require curli fibers or biosynthesis of O-Antigen via the *yihO/P* system (Fig. 1B).

The YihO/P permeases were previously known to transport the O-Antigen subunits for LPS biosynthesis

in *E. coli* and *Salmonellae*[16, 39], but are now classified in sulfoquinovose catabolism in *E. coli*[75, 76].

Elucidating the mechanism of STy biofilm formation

In order to identify the players that drive STy biofilms, we grew a transposon library in the STy

parent strain H58 as biofilms in vitro in gallstone-mimicking conditions and developed Tn-ClickSeq

analysis to map the transposon genome junctions enriched in the planktonic and biofilm sub-populations.

We found Tn-genome junctions in 47% and 20% of total sequencing reads from the planktonic and biofilm

fractions, respectively. These differences indicated that inactivating transposon insertions were presumably

tolerated in only a low number of genes in biofilms, which correlated with their enrichment in the planktonic

fraction. Also the technical challenges of isolating genomic DNA with high efficiency from STy cells

attached to cholesterol-coated surfaces may affect the total sequencing yields, leading to very low insertion

indices in biofilms. Nevertheless, our Tn-ClickSeq analysis generated novel insights by identifying Sth

fimbriae, Vi capsule and the lipopolysaccharide core as structural components and IraP as a regulator of

STy cholesterol-attached biofilms. Notably, Sth fimbriae belongs to a different fimbrial class than curli

fimbriae and are not present in STm strains[50, 51]. Our comprehensive analysis of the STy biofilms also

opens up exciting new directions for mapping the complete genetic signature of STy carriage.

RpoS is a key regulator of STy biofilms

The discovery of IraP was significant, because it motivated our subsequent investigations of the

downstream starvation-induced sigma factor RpoS during STy biofilm development. The reduction in

biofilm formation was similar in the *iraP* and *rpoS* deletion strains (Fig. 3B), which suggested that epistatic

interactions between IraP and RpoS regulated the development of STy surface-attached communities. The

activation of IraP in low phosphate environments[77] provides a regulatory insight on the activation of RpoS

in bi. 3Bofilm favoring gallstone-mimicking conditions. The observation that elimination of *iraP* or *rpoS*

only reduced biofilms by 50% (Fig. 3B) suggests other inputs are also involved and need to be

characterized. Strong candidates include the global regulators OxyR, BolA and LeuO and the TCRS

RcsC/B.

Possible experimental biases in transposon library preparations may have resulted in the exclusion

of *rpoS* from our list of Tn-ClickSeq targets due to its low insertion index (25). For example, the *rpoS*

insertions were deterimental to survival in gallstone-mimicking conditions as shown by a higher insertion

index of 65 in our input Tn-ClickSeq library compared to the planktonic sub-population. An alternative

approach of serial passage of a transposon library in biofilm favoring conditions to specfically identify

genetic systems regulating the development of community behavior might be informative. Nevertheless,

we established that the formation of STy biofilms required RpoS and determined its role in activating the

transcription of the extracellular matrix components: Sth fimbriae, Vi capsule and lipopolysaccharide (Fig.

5B). Other studies have shown that RpoS and the TCRS RcsC/B regulate Vi capsule synthesis under

osmotic stress[78], and RpoS was required for the SPI-9-mediated adhesion of STy to epithelial cells[79]. The

mechanisms of biofilm formation in STy are unique, in STm, the starvation response regulator RpoS

activates biofilms via CsgD[80, 81], but we specifically eliminated a role for CsgD and curli in the formation

of STy biofilms (Fig. 1B). We propose that RpoS directly binds to the upstream regulatory sequences of

STy genes encoding the extracellular biofilm matrix components (Fig. 5C) that are AT-rich and harbor a

possible extended -10 promoter element, as previously characterized for the RpoS regulon in *E. coli*[63, 64].

It is important to note here that the protein sequences of RpoS are highly conserved between *E. coli* and

H58 (99% identical). A complete understanding of RpoS signaling pathways that enable the switch from

the planktonic lifestyle to biofilms will require validating the precise σ^S-recognition sequences, and hence

the transcriptional start sites, for *waaZ*, *sthC*, *tviA* and *tviB*. Further understanding will also require

investigating a possible co-regulatory function of Crl in stimulating the RpoS activity to enable

transcriptional activation, as determined for other RpoS-responsive genes, including the *csg* operon, *adrA*

and *ssrA*[66, 82-84]. Overall, the presence of other quintessential RpoS-regulated genes, such as *dps*, *bolA*,

uspA and *osmB* in our Tn-ClickSeq dataset (Supplementary Fig. 5B) raises an interesting question as to

how RpoS coordinates stress signaling and matrix production in gallstone-mimicking conditions.

Visualization of STy biofilms by electron microscopy

A high degree of autofluorescence and the non-specific binding of fluorescent dyes and antibodies

to cholesterol surfaces prevents the visualization of mature STy biofilms using high-resolution fluorescence

microscopy techniques. Although some protocols have been developed to improve the fluorescence

imaging of STy aggregates in vitro and ex vivo, crucial controls that included STy strains defective in

forming biofilms were lacking[85-87]. Therefore, in order to clearly visualize STy biofilms attached to

cholesterol surfaces, we employed scanning electron microscopy and observed distinct aggregates of the

wild type H58 parent and substantially reduced aggregate formation in the STy biofilm mutants validated

from our Tn-ClickSeq analysis (Fig. 4).

Significance of the STy-Zebrafish infection model

Our use of zebrafish larvae as a heterologous host for *S.* Typhi validated the important role of RpoS

in enabling *S.* Typhi colonization in long-term infections. Reducing STy persistence by inactivating *rpoS*

significantly decreased the pathogen load, prolonged host survival and most importantly, abolished

hepatobiliary colonization at 6 dpi. Following STy infections beyond 6 dpi may be possible by rearing

germ-free zebrafish larvae using established methods[88].

Possible role of RpoS in the evolution of STy carriage

Finally, a strong correlation has been observed between the ability to form biofilms and the duration

of STy shedding and carriage in typhoid patients from Pakistan[89]. Most recent phylogenomic analysis has

proposed that the ancestral H58 haplotype originated in a chronic carrier from India and evolved to give

rise to the three sub-lineages that cause a majority of typhoid infections in Asia and Africa[23]. In the light

of these results, it is tempting to propose that RpoS contributes to niche adaptation in *S.* Typhi by activating

the formation of biofilms in chronic carriers.

Figure 1: The master regulator CsgD, and the STm lifestyle regulator SsrB, are not required for the formation of cholesterol-attached STy biofilms. Crystal violet staining assays: (A) Wild type strains Ty2-b, CT18, CT117 and H58 formed robust biofilms at day 2 in gallstone-inducing conditions and (B) STy null strains deleted for the indicated STm-biofilm homologs formed similar biofilms as the H58 wild type strain at day 2. Growth medium added to cholesterol-coated Eppendorf tubes was used as the control and subtracted from all measurements. N = 3 in at least triplicates, error bars represent Mean ± SD, ns = not significant by One-way ANOVA.

Supplementary Figure 1: H58 forms 'atypical' biofilms. (A) Examination of a macrocolony of

S. Typhimurium (STm) displaying a <u>r</u>ed, <u>d</u>ry and <u>r</u>ough (rdar) morphotype on Luria-Bertani (LB)

agar without salt medium containing the congo red dye, while *S.* Typhi (STy) are 'smooth' and lack

the rdar morphology. (B) Wild type STy strain H58 was unable to form biofilms compared to the

wild type STm strain 14028s, when grown in LB broth without salt medium in polystyrene plates at

two days as determined by a crystal violet staining assay and (C) The cholesterol-attached biomass

formed by wild type H58 did not increase significantly at days 4 and 6 compared to day 2, as

determined by a crystal violet staining assay. Growth medium added to cholesterol-coated

Eppendorf tubes was used as the control and subtracted from all measurements. N = 3, in at least

triplicates, error bars represent Mean ± SD, ns = not significant by one-way ANOVA.

443
444
445

Figure 2: Tn-ClickSeq analysis investigates novel pathways that drive the formation of STy biofilms. (A) A general scheme depicting Tn-ClickSeq analysis, from the isolation of planktonic and biofilm fractions, DNA extraction and the generation of Tn-ClickSeq libraries using a reverse transcription reaction (RT) with a Tn-specific primer and azido-nucleotides/dNTP mixtures. The azido-terminated cDNAs were click-ligated to an alkyne-adapted illumina P5 adapter to generate Illumina libraries. After PCR and barcoding, the reads that contained partial Tn sequence and genome insertion site were pair-end sequenced and subjected to downstream analysis. (B) A linearized map of the average Tn-insertion sites from three biological replicates showed genome-wide differences in the planktonic sub-population (orange) and biofilms (green). The H58-Tn library inoculum (blue) and the H58 parent without any transposons (black, control) were used as positive and negative controls, respectively. (C) Principal component analysis showed distinct clustering of respective replicates of planktonic (orange) and biofilm (green) fractions as compared to the input library (blue) and (D) The number of transposon insertions of each gene was normalized per 1000 bp of gene length and averaged among replicates for each group to depict a hierarchical clustering of Tn-ClickSeq targets.

(A)

(B)

(C)

(D)

Sample	Raw reads (after QC)	Reads with primer	Reads with Tn	Reads mapping to H58 genome
Input#1	14, 256, 570	13, 914, 412	7, 099, 879	6, 891, 853
Input#2	11, 418, 976	10, 962, 217	5, 572, 442	5, 365, 704
Input#3	8, 352, 374	8, 093, 450	3, 974, 593	3, 825, 148
Planktonic#1	15, 228, 533	15, 000, 105	8, 234, 051	8, 013, 378
Planktonic#2	13, 131, 165	12, 881, 673	7, 440, 776	7, 260, 709
Planktonic#3	16, 164, 559	15, 905, 926	7, 942, 465	7, 759, 788
Biofilms#1	406, 289	23, 158	20, 738	20, 543
Biofilms#2	540, 650	82, 179	57, 380	55, 349
Biofilms#3	459, 357	299, 041	207, 996	202, 089
Negative control	18, 888, 763	18, 359, 878	2, 433	2, 331

Supplementary Figure 2: (A) A flowchart of the bioinformatics pipeline employed for Tn-ClickSeq analysis depicting the filtering of raw reads containing both primer and partial Tn sequences, the subsequent identification of genome insertion sites using the reference H58 genome and lastly de-duplexing to remove PCR biases. (B) Percentages of reads that matched the primer, transposon sequence (IR sequence) and the host genome from left to right: input, planktonic and biofilm sub-populations. (C) Sequencing yields of the respective input, planktonic and biofilm Tn-ClickSeq libraries in triplicates and of an H58 strain without any transposon insertions as a negative control and (D) Graphical representation of Tn insertion sites per one million of raw reads showing an average of 470,000 reads for the input fraction, 520,000 for the planktonic fraction and 197,000 for the biofilm fraction. A similar analysis using an H58 strain without any transposon insertions resulted in only 100 false positives.

Supplementary Figure 3: A Gene Ontology analysis of transposon insertion sites in 1515 genes that were enriched in the planktonic sub-population as compared to biofilms showed a significant enrichment of cell membrane components, transmembrane ion transport pathways and other membrane related activities.

Figure 3: Biofilms are defective in STy null mutants identified by Tn-ClickSeq. (A) Defective biofilms formed by *tviA/tviD*, *sthC* and *waaZ* null strains suggested a role of Vi capsule, Sth fimbriae and the lipopolysaccharide core in extracellular matrix production, respectively and (B) H58 strains deleted for *iraP*, which regulates the protein stability of RpoS, and *rpoS,* which encodes the starvation sigma factor, formed less biofilms than the wild type parent. The defect in biofilm formation of the *rpoS* null strain was complemented by overexpression of *rpoS* from a plasmid *in trans*. N = 3, in at least triplicates, error bars represent Mean ± SD, in a crystal violet staining assay. Growth medium added to cholesterol-coated Eppendorf tubes was used as the control and subtracted from all measurements, ns = not significant, *p ≤ 0.05, **p ≤ 0.01 and ***p ≤ 0.001 by one-way ANOVA.

Supplementary Figure 4: (A) Double null mutant derivatives of an H58 parent defective for Sth fimbriae and Vi-polysaccharide (*sthC tviD*), and for LPS biosynthesis and Sth fimbriae (*waaZ sthC*) showed a reduced ability to form cholesterol-attached biofilms, by around 50%, compared to the wild type parent. N = 3, in at least triplicates, error bars represent Mean ± SD, in a crystal violet staining assay. Growth medium added to cholesterol-coated Eppendorf tubes was used as the control and subtracted from all measurements, ****$p \leq 0.0001$ by one-way ANOVA and (B) Biofilm mutants are not compromised in planktonic growth - Strains deleted of *iraP*, *rpoS*, *waaZ*, *sthC* or *tviD* and the wild type H58 strain were grown for 12 hours in Luria-Bertani broth at 37°C/250 rpm and the absorbance at 600 nm was measured every 2 hours.

Figure 4: Ultra-structure visualization reveals the loss of dense aggregate formation in STy biofilm mutants. Representative scanning electron microscopy images of rich *S.* Typhi biofilms formed by the H58 (wt) and strikingly smaller-sized aggregates formed by the strains deleted of *rpoS*, *sthC*, *waaZ* or *tviD* when grown in cholesterol-coated Eppendorf tubes in gallstone-mimicking conditions. Scale bar = 5 μm.

Figure 5: Steady-state levels of biofilm components are increased in gallstone-mimicking conditions and RpoS is required for direct transcriptional activation of biofilm matrix genes. Real time RT-qPCR analysis showed (A) a significant increase in transcription of *rpoS*, *sthC*, *waaZ* and *tviB* in the cholesterol-attached fraction compared to the planktonic fraction in the wild type H58 parent. (B) In the *rpoS* null strain, there was a significant decrease of biofilm matrix components *waaZ*, *tviD* and *sthC*. *rrsA* transcript levels were used as an internal control. N = 3, in triplicates, error bars represent Mean ± SD, ***p ≤ 0.001 and ****p ≤ 0.0001 by one-way ANOVA and (C) Electrophoretic mobility shift assays showing free biotinylated DNA, biotinylated DNA-protein complexes in the presence of RpoS (σ^S) (**) and a decrease in the biotinylated DNA-protein complexes in the presence of excess amounts of non-biotinylated DNA (*) for *waaZ*, *sthC*, *tviA and tviB,* respectively.

(A)

(B)

Supplementary Figure 5: (A) Sequences of upstream regulatory regions of *waaZ*, *sthC* and *tviA* are shown as snapshots from the IGV genome browser with the RpoS binding sites, an extended -10 element followed by an AT-rich sequence, highlighted as dark blue lines (total conservation) and cyan lines (partial conservation) and (B) Hierarchical clustering of a subset of Tn-ClickSeq targets showing an enrichment of RpoS-regulated genes[59,60] in the planktonic library. For example, these include genes that confer acid resistance (*gadC*), adaptation to osmotic (*osmB*) and other environmental stresses (*psiF*, *uspA*, *uspB*), encode stress responsive regulators (*bolA* and *dps*), an iron storage protein (*bfr*), a toxin (*tomB*) and a mechanosensitive channel (*mscL*).

Figure 6: STy colonizes the intestines of chronically infected zebrafish, and RpoS is required for persistent colonization in vivo. (A) The number of STy colonies recovered from persistently infected whole larvae was drastically reduced in the *rpoS* null strain (magenta bars) compared to the wild type (green bars) at days 2, 4 and 6 post infection. Control larvae exposed to PBS were void of *Salmonella*. N = 3, five infected larvae from each group, error bars represent Mean ± SD and ns = not significant, *p ≤ 0.05 and ***p ≤ 0.001 by two-way ANOVA with Sidak's multiple comparison tests. (B) Representative merged images (red channel and bright field) of live zebrafish larvae showing the presence of wild type mCherry-expressing *S.* Typhi in the gut. There was a stark reduction of bacteria in the infections using the mCherry-tagged *rpoS* null strain at 2, 4 and 6 dpi. No fluorescence was detected for the PBS control at all time points. The cartoon on top depicts an infected zebrafish larva with the highlighted intestinal region and was adapted from BioRender. 20x magnification, Scale bar = 100 µm and (C) Larvae infected with the *rpoS* null strain (magenta line) survived longer and showed a significant increase in the percentage of survival at 6 dpi compared to the wild type H58 (green line). The survival rate of the PBS control is shown as a dotted black line. N = 3 with 30 to 60 larvae in each group, error bars represent Mean ± SD, *p ≤ 0.05 by two-way ANOVA with Sidak's multiple comparison tests.

1073

1074

1075

1076

1077

1078

1079

1080

1081

1082

1083

1084

1085

1086

1087

1088

1089

1090

1091

1092

1093

1094

1095

1096

1097

1098

Figure 7: Chronic colonization in the gall bladder requires RpoS. Representative whole-mount
immunohistochemistry images showing the successful detection of anti-mCherry antibody signal (magenta) in
the gall bladder region of a larva infected with an mCherry-tagged wild type strain at 6 dpi (middle panel). The
hepatobiliary system is highlighted by yellow rectangles in merged images and zoomed-in images show mCherry-
positive fluorescent signals, validating the hepatobiliary colonization of STy in wild type infections at 6 dpi
(middle right panel). In contrast, there was a drastic reduction in fluorescent signals from the gall bladder region
of a larva infected with an mCherry-tagged *rpoS* null strain, with only a weak fluorescent signal as observed in
the anterior intestine at 6 dpi (lower right panel, zoomed-in image). A low level of background fluorescence was
also detected in the uninfected PBS control (top right panel, zoomed-in image). 20X magnification, Scale bar =
100 μm for all images except for zoomed-in images with 10 μm scale bars. N = 3 with 5 to 10 larvae analyzed in
each group.

Supplementary Figure 6: Representative whole-mount immunohistochemistry images showing the successful detection of anti-Vi-polysaccharide antibody signal (magenta) in the gall bladder region, as marked by a yellow rectangle in the merged image of a wild type infected larva at 6 dpi (middle right panel). Liver (L) and intestine (I) are marked in the bright field image on the left to pinpoint the gall bladder (G) position. The presence of Vi-antigen-positive STy clusters in the gall bladder is highlighted in the respective zoomed-in image (middle right). The gall bladder region, as marked by a yellow rectangle, of an *rpoS* null infected larva remained negative for anti-Vi-antigen antibody staining (lower panel, zoomed-in image on the right). No fluorescence was detected in the uninfected PBS control (top panel). 20X magnification, Scale bar = 100 μm for all images except for zoomed-in images with 10 μm scale bars. N = 3 with 5 to 10 larvae analyzed in each group.

Methods

Bacterial strains and growth

The bacterial strains and plasmids used in this study are listed in Supplementary Table 1. *S.* Typhi strains

were routinely grown in Luria-Bertani broth (LB) or Nutrient broth (NB) (BD Difco) medium with shaking

at 275 rpm at 37°C in the presence of 100 μg/mL Ampicillin, 12.5 μg/mL Tetracycline, 25 μg/mL

Chloramphenicol or 50 μg/mL Kanamycin (Millipore Sigma) when necessary. For the growth of STy

cholesterol-attached biofilms, a modified NB medium (NB*) containing 3% w/v NB, 1.75% w/v sodium

chloride, 0.25% w/v potassium chloride, 1% w/v sodium choleate and 2% w/v glucose (Millipore Sigma)

was used [36]. For observing the rdar morphotype, 40 μL of overnight LB broth cultures of STm wild type

strain 14028s and STy wild type strain H58 were spotted on agar plates containing 1% w/v Tryptone and

0.5% w/v Yeast Extract (LB broth without salt) supplemented with Congo Red (40 μg/mL) (Millipore

Sigma) and kept at 30°C for two days. Biofilms in 96-well polystyrene plates were grown in LB broth

without salt as described previously[10].

Molecular Biology techniques

All DNA manipulation procedures were carried out according to[90] using reagents procured from Qiagen,

Millipore Sigma or Invitrogen. All transformations in STy wild type strain H58 were performed by standard

electroporation protocols[90]. Polymerase chain reactions (PCR) were carried out using oligonucleotides as

listed in the Supplementary Table 2 following standard protocols[90].

Strain construction

The *ssrB* null mutation in H58 strain was generated by transducing the *ssrB::kan* allele from the STm strain

DW85 using standard P22 transduction protocols[91]. Other gene deletions in the STy wild type strain H58,

as listed in Supplementary Table 1, were generated by the lambda *red* homologous recombination technique

as described in[92, 93]. Briefly, plasmids pKD3, pKD4 or the *TetRA* DNA were used to generate linear DNA

fragments by PCR using gene-specific hybrid primers as listed in Supplementary Table 2. A 10 ml LB broth

culture of H58 transformed with the plasmid pKD46 (containing 100 μg/mL Ampicillin and 20 mM

Arabinose) was used to electroporate 600 ng to 1 μg of purified PCR product following the protocol as

described in[92]. The cells were recovered for at least 6 h at 275 rpm/30 °C after which the cells were

harvested and plated on respective selective plates. The *sthC tviD* and *waaZ sthC* double null strains were

generated similarly except the Kan^R cassettes were first removed from the *sthC* and *waaZ* null mutant strains

by transforming with the temperature-sensitive plasmid, pCP20[93]. Chromosomal deletions were confirmed

by PCR using flanking primer pairs as listed in Supplementary Table 2.

1208 STy cholesterol-attached biofilms in gallstone-mimicking conditions

STy biofilms were routinely grown following protocols adapted from[36, 39]. Sterile 1.5 ml Eppendorf tubes

containing 200 μL of 10 mg/mL cholesterol (Millipore Sigma) in diethyl ether were air dried aseptically

(2-3 h) and 20 μL of STy strains grown overnight in LB/NB* was added to 180 μL NB* medium. The tubes

were incubated for two days at 275 rpm at 37 °C. Cholesterol-coated tubes containing only 200 μL NB*

medium served as controls in all the experiments. For monitoring the time course of biofilm formation, the

cholesterol-coated tubes were further incubated for four and six days, with fresh 200 μL NB* medium

replaced every two days.

1216 Crystal violet staining assay

The amount of STy cholesterol-attached biofilms in gallstone-mimicking conditions was estimated using

crystal violet staining assays adapted from[10, 39]. The supernatant/growth medium was removed, and each

tube was washed once with 400 μL of Phosphate-buffered Saline (PBS). The attached biofilms were then

stained with 200 μL of 0.1 % w/v crystal violet solution (filtered using Whatman Grade 1 filter paper) for

five minutes at room temperature (RT). This was followed by washing once with 400 μL PBS and addition

of 200 μL absolute ethanol. Appropriate dilutions were then measured for absorbance at 595 nm using an

iMark™ Microplate Absorbance Reader. Each experiment was performed in triplicates or pentuplicates.

1224 Generation of H58-Tn library

1225 The *TnTMDH5deloriR6K* genome library in the *S.* Typhi wild type strain H58 (H58-Tn) was kindly

1226 provided by Dr. Stephen Baker, University of Cambridge, UK[48]. The Tn5 transposon was derived from

the plasmid EZ-Tn5<R6Kgori/KAN-2> (Epicentre Biotechnologies) as described in the original reference[43]. Briefly, the plasmid was digested in a 10 μL reaction containing 2.5 μL TnTMDH5deloriR6K plasmid DNA(10ng/ mL), 1μL 10X NEB4 buffer, 5.5 μL water, 0.5 μL MspAI1 restriction enzyme and 1 μL BSA, for 2 h at 37 °C. The transposon was PCR amplified using the oligonucleotides (5'-CTGTCTCTTATACACATCTC CCT and 5'-CTGTCTCTTATACACATCTCTTC) and Pfu DNA polymerase Ultra Fusion II, (Stratagene). The PCR steps were; 95 °C/ 90 seconds, followed by 30 cycles of denaturation at 95 °C/10 sec, annealing at 58 °C/20 sec and extension at 72 °C/ 20 sec, and a final extension at 72 °C/3 min.

The Tn5 amplicons were PCR purified and phosphorylated at their 5' ends in a 80 μL phosphorylation reaction including 70 μL purified Tn5 amplicons (approximately 50 μg/ml), 8 μL 10X T4 buffer, 1 μl ATP (75 mM, Roche) and 1 μL T4 Polynucleotide Kinase (NEB). The reaction was incubated at 37 °C/45 min followed by inactivation at 65 °C/20 min. The phosphorylated Tn5 transposons were purified with a 0.5 volume of a 1:1 mixture of phenol:choloroform and centrifuged at 14,000 g/10 min/ 4 °C. The supernatant was isolated and DNA was precipitated in a 30 μL reaction including 10 μL supernatant, 1μL 3M sodium acetate (pH7.5) and 19 μL absolute ethanol. The mixture was centrifuged at 14,000 g/10 min/4 °C, the supernatant step was discarded and the pellet was rinsed twice with 200 μL   of 70 % (w/v) ethanol. The pellet was eluted in 10 μL 1X Tris-EDTA (TE) buffer (pH7.5) and stored at -20 °C.

Next, transposomes were prepared using 2 μL of above purified phosphorylated Tn5 DNA (~70 μg/mL), 4 μL EZ-Tn5 Transposase (Epicentre Biotechnologies), 2 μL 100% glycerol and 4μL 50% v/v glycerol. The reaction was carried out for 30 min/ RT. 0.2 μL of transposomes were then mixed with 60 μL electrocompetent cells of STy H58 strain for electroporation. Electrocompetent cell preparations and electrotransformations were performed exactly as described previously[43]. Finally, the kanamycin resistant colonies were resuspended in 10 % v/v glycerol for storage at -80 °C and each batch of the H58-Tn mutant library included 10 electrotransformations used to generate approximately 3×10^5 mutants.

Isolation of fractions for Tn-ClickSeq libraries

A 1 μL loop of frozen stock of the H58-Tn library was inoculated in 10 mL NB* cultures and incubated overnight at 275 rpm at 37 °C. This was the 'input' fraction, which was used to prepare thirty tubes of cholesterol-attached biofilms in gallstone-mimicking conditions. After two days, the culture supernatants were removed and pooled to obtain the 'planktonic' fraction. To isolate the 'biofilm' fraction, 400 μL of PBS was added to each tube followed by sonication in a XUB Digital Ultrasonic Bath (Grant Instruments) for 20 min, maximum power, no leap, at RT. The harvested biomass was pooled, and the sonication step was repeated thrice. The final pool at the end of four sonication cycles was the 'biofilm' fraction.

Genomic DNA isolation for generating Tn-ClickSeq libraries

Planktonic and biofilm fractions were centrifuged at 24,000 g at 4 °C for 1.5 h in a Beckman Coulter Avanti J-26XP centrifuge. Respective supernatants were discarded, and the pellets were stored on ice. 1 ml of the input fraction was also centrifuged at 15,500 g at 4 °C for 10 min and the pellet was stored on ice. Each of the pellet fractions were then resuspended in 600 μL TE buffer (pH 8.0) with 40 μL 10% w/v sodium dodecyl sulphate, 4 μl of 20mg/mL Proteinase-K (Invitrogen) and 2 μl of 10mg/mL RNaseE (Invitrogen) and mixed well by vortexing. Samples were incubated at 37 °C for 1 h, after which an equal volume of Phenol:Chloroform mixture (pH 6.7/8.0) was added, mixed and centrifuged at 15,500 g at 4 °C for 15 min. The upper aqueous phase was added to a fresh tube and an equal volume of chloroform was added, mixed, and centrifuged at 13,000 rpm at 4 °C for 15 min. The supernatants were removed, 2.5 to 3 volumes cold absolute ethanol was added, and stored overnight at -20 °C. DNA pellets were obtained by centrifuging the samples at 15,500 g at 4 °C for 15 min, followed by a 70% ethanol wash. The pellets were air dried, and DNA was resuspended in 40 μL of nuclease-free water.

Preparation of Tn-ClickSeq libraries

For Tn-ClickSeq, genomic DNA from the input, planktonic and biofilm fractions was reverse transcribed using a reverse transcriptase (SSIII, Invitrogen) and Azido-NTPs. A Tn-specific reverse primer (3'21-39, Supplementary Table 2) was designed to the 3' proximal end, 21 to 39 bp of transposon, with an overhang

of the reverse complementary sequence of Illumina adapter (Supplementary Figure 4, Supplementary Table

2). 500 ng DNA was mixed with 1 µL of 5 µM primer, and 1 µL of 10 mM AzNTP/dTNP mixture

(AzNTP:dNTP = 1:35). This initial reaction mix was heated at 95 °C for 5 min and then cooled to 50 °C in

gradual steps of 0.1 °C per second. Other reaction components for the reverse transcription reaction were

then added: buffer, DTT, SSIII (as per manufacturer's protocol), and kept in a thermocycler at 50 °C for 50

min, after which the reaction was terminated at 95 °C for 5 min. This was followed by the standard ClickSeq

protocol as previously described[46, 94, 95]. Briefly, immediately after denaturing, the DNA products were

purified with Solid Phase Reversible Immobilization (SPRI) beads and click-ligated with a 5'-alkyne-

modified adapter including a 12-nucleotide unique molecular identifier (UMI). The click-ligated product

was then purified, barcoded, amplified with 18 to 20 cycles of PCR, and analyzed by agarose gel

electrophoresis. The final Tn-ClickSeq libraries were subjected to pair-end sequencing on a NexSeq 550

platform at the Genomics core, UTMB.

Bioinformatics analysis

We designed a bioinformatics pipeline to process the hybrid reads that originate from the inserted

transposon and extend into the host genome (Supplementary Figure 5). The raw paired-end FASTQ reads

were first pre-processed to trim the Illumina adapter, filter low-quality reads and extract UMIs using

fastp[96]: -a AGATCGGAAGAGC -U --umi_loc read1 --umi_len 14 --umi_prefix umi -l 30. We then used

FASTX toolkit (http://hannonlab.cshl.edu/fastx toolkit/index.html) to reverse complement the R2 reads for

ease of downstream analyses. We filtered the reads that contained the 19 bp primer sequence (targeting the

Tn) with 1 nucleotide mismatch allowance with *cutadapt*[97]*:* -a cctatagtgagtcgtatta -e 0.1 -O 19 -m 30 --

discard-untrimmed. To obtain the reads containing primer sequences, we further filtered reads that

contained the last 10 nucleotides of the 3' invert repeat (IR) with 0% error rate allowance with *cutadapt*: -

a ctgtctctta -e 0 -O 10 -m 30 --discard-untrimmed. After trimming the IR sequence, the rest of IR-containing

reads   were   mapped   to   the   H58   genome

(https://www.ncbi.nlm.nih.gov/datasets/genome/GCF_001051385.1/) with *hisat2*[49], and then processed

with *SAMtools*[98]: view/sort/index. We de-duplexed the data to minimize PCR bias with umi_tools[99]:

dedup --method=unique. Finally, the locations of the transposon insertion sites were extracted with

BEDTools[100]: genomeCoverageBed: -3 -bg. Raw sequencing data is available at the Sequence Read

Archive, Project ID, PRJNA1029173.

Gene mapping and target analysis

The number of insertion reads at each insertion site revealed by Tn-ClickSeq were ratiometrically

normalized across different samples. This was followed by the assignment of gene names and annotations

within H58 (GCF_001051385.1). We normalized the number of insertions to the length of each annotated

gene (per 1 Kbp length of gene) to obtain the insertion indices for each samples. Matrices of all the

normalized insertion dataset were then processed with DESeq2 to identify genes enriched or depleted in

each conditions and perform Principal component analysis[101]. Hierarchical clustering was conducted with

Cluster   3.0   (http://bonsai.hgc.jp/~mdehoon/software/cluster/),   followed   by   TreeView

(http://jtreeview.sourceforge.net/) to build the graphic map. Gene ontology analysis was conducted with

GeneOntology web server (http://geneontology.org/) with *Salmonella* Typhimurium as a reference.

RNA isolation

Planktonic and biofilm fractions were centrifuged at 24,000 g at 4 °C for 1.5 h in a Beckman Coulter Avanti

J-26XP centrifuge. Respective supernatants were discarded, and the pellets were either stored at -80 °C, or

immediately processed for total RNA isolation by the Trizol method[102]. Briefly, the pellets were

resuspended in 1 mL TRIzol reagent (Life Technologies) and incubated at RT for 5 min. 200 μL chloroform

was added, mixed well, and incubated for 3 min at RT. The mixtures were centrifuged at 15,500 g for 15

min in cold and 500 μL isopropanol was added to the supernatants. The samples were transferred to -20 °C

overnight, after which the RNA was pelleted by centrifugation at 15,500 g for 15 min in cold. This was

followed by a 75% ethanol wash. The pellets were air dried, and the RNA was resuspended in 20 μL of

nuclease-free water.

RT-qPCR

1 μg of total RNA extracted from the biofilm fraction of two days old cholesterol-attached biofilms was

used for a reverse transcription reaction with iScript Supermix (Bio-Rad) according to the manufacturer's

protocol. This was followed by amplifying 50 ng cDNA by real-time qPCR (RT-qPCR) using SsoFast

EvaGreen Supermix (Bio-Rad) and internal primers specific for *rpoS*, *sthC*, *waaZ* and *tviB*; *rrsA* was used

as a normalization control (Supplementary Table 2). The annealing temperature for all the primer pairs was

56 °C. All experiments were performed in triplicates with at least three independently isolated RNA

preparations. Relative expression was determined using the $2^{-\Delta\Delta C_T}$ (Livak) method as described in[102] and

plotted using the GraphPad Prism 10 software.

Scanning electron microscopy

STy cholesterol-attached biofilms were grown in gallstone-mimicking conditions for two days and after

removal of the growth medium were fixed in a primary fixative containing 2.5% w/v formaldehyde (made

from paraformaldehyde), 0.1% v/v glutaraldehyde, 0.01% v/v trinitrophenol and 0.03% w/v $CaCl_2$ in 0.05M

sodium cacodylate buffer (pH 7.3). Samples were post-fixed in 1% w/v OsO_4 in cacodylate buffer,

dehydrated in ethanol and infiltrated with hexamethyldisilazane (HMDS) to prevent cracking during drying.

After air drying, the conical parts of the Eppendorf tubes were cut into strips, mounted on SEM specimen

holders (metal stubs) and sputter-coated with iridium in an Emitech K575X (Emitech, Houston, TX) sputter

coater for 30 seconds at 20 mA, at the Electron Microscopy Laboratory, Department of Pathology, UTMB.

The samples were examined in a JEOL JSM-6330F Scanning electron microscope at the Texas Center for

Superconductivity, University of Houston, at 4 μA and 2 kV.

Zebrafish husbandry

All the protocols used for zebrafish experiments were approved by the University of Texas Medical Branch

Institutional Care and Use Committee. Adult and larval zebrafish were maintained using standard

husbandry procedures[103] at our in-house satellite facility. The wild type AB line was used for lifespan

analysis and Typhi load measurements and the *casper* line was used for visualizing mCherry-tagged STy

strains in infected larvae by confocal fluorescence imaging. Eggs were obtained by the natural spawning

method[103] and kept at 28 °C in embryo medium containing methylene blue for two days. At 2 days post-

fertilization (dpf), the larvae were shifted to an embryo medium containing 25 µg/mL gentamicin for 6 h,

after which they were transferred to a sterile embryo medium at 28 °C. Fresh embryo medium was replaced

daily.

Static immersions of zebrafish larvae

All infections were performed using mCherry-tagged STy strains, as listed in Supplementary table 2. For

static immersions, 1 ml of an overnight LB broth culture of mCherry-tagged STy strains was inoculated in

10 ml LB broth containing 100 µg/mL ampicillin and grown for 4.5 hours at 275 rpm at 37 °C. Growth was

normalized by measuring absorbance at 600 nm. The strains were harvested by centrifugation at 4,200 g

for 15 min at RT and resuspended in 500 µl sterile embryo medium. Ten larvae 5 dpf were added to 8 ml

embryo medium in a 6-well polystyrene plate, followed by the addition of 80 µl of harvested STy cultures

(10^9 cfu/mL). Equal infection doses across strains were verified using aliquots of the harvested STy cultures

by the total viable counting method. Larvae exposed to equal volume of PBS served as controls. After

24 h, the larvae were transferred to fresh embryo medium in a 6-well plate. Fresh embryo medium was

replaced daily. The larvae were fed Sparos Zebrafeed (<100 µ) once daily from 9 dpf, as adopted from a

delayed initial feeding model previously described[68]. Typically, infections were performed with thirty to

sixty larvae in each group.

Bacterial load estimates from infected larvae

For enumerating the STy load at 2, 4 and 6 dpi, infected larvae were isolated and added to embryo medium

containing 60 to 100 µg/mL buffered Tricaine. The anesthetized larva was transferred to a 1.5 ml Eppendorf

tube containing 200 µl PBS and homogenized well with a motorized micropestle. The homogenate was

serially diluted and colony forming units (cfu) were estimated by the total viable counting method on LB

agar plates containing 100 µg/ml ampicillin. For each time point, five larvae were used from each group.

PBS control larvae remained sterile. The experiment was repeated at least three times.

Lifespan analysis

1377 Infected larvae were checked daily under a dissection microscope, and the percentage survival was scored.

1378 Larvae that did not show any heart beats were considered dead.

1379 Confocal fluorescence imaging

Five to ten larvae from each group were withdrawn at 2, 4 and 6 dpi and anesthetized in an embryo medium

containing 60 to 100 μg/mL buffered Tricaine. These were then mounted on 35 mm glass-bottomed dishes

with a drop of 0.8% low melting point agarose (supplemented with 60 to 100 μg/ml buffered tricaine).

Images were acquired on an Olympus SpinSR-10 Yokogawa spinning disk confocal microscope fitted with

an ORCA Fusion sCMOS camera (Hamamatsu) using a 20x objective (NA 0.8, Olympus), 561 CSU (or

using 640 CSU for anti-mCherry IHC, below) and a step size of 1.5 μ in the Z-dimension. Images were

analyzed using Image J software.

1387 Whole-mount immunohistochemistry (IHC) of zebrafish larvae

To detect STy colonization in the gall bladder we performed immunofluorescence of whole zebrafish larvae

at 6 dpi[104]. Briefly, AB larvae infected with wild type H58 and *rpoS* null strains were fixed in 4%

paraformaldehyde in PBS (pH 7.4) overnight at 4°C, washed thrice with PBS for 5 minutes and treated with

a mixture of 3% hydrogen peroxide and 0.5% potassium hydroxide for 30 mins at RT to remove

pigmentation. After three PBS washes, dehydration was performed using a graded methanol/PBS series of

25%, 50%, 75% and 100% with each step of 5 mins. Larvae were then kept in 100% methanol for overnight

at 4°C. Rehydration steps involved a graded methanol/PBS series of 75%, 50% and 25%, with each step of

5 mins, followed by three washes with PBS containing 0.1% Tween-20 (PBS-T). For antigen retrieval, the

larvae were transferred to a Tris buffer (150 mM Tris-HCl, pH 9.0) for 15 min at 70°C, washed twice with

PBS-T and thrice with milli-Q water on ice. Specimens were then incubated in 100% acetone stored at

-20°C for 20 mins, followed by washes with milli-Q water and PBS-T. In case of IHC using the anti-

mCherry primary antibody (abcam AB213511), an additional step of treatment with 0.3% v/v Triton-X 100

for 10min/RT was included at this step. Samples were washed twice with PBS-T for 5 min before blocking

in 10% BSA in PBS-T for 2h at RT, washed with PBS-T and kept overnight in 1% BSA solution in

PBS-T containing a 1:100 dilution of anti-mCherry antibody or 1:500 dilution of anti-rabbit-Vi-

polysaccharide antibody (BD BioSciences) overnight at 4°C. Subsequently, four PBS-T washes of 15 min

each were performed, and the larvae were incubated in a secondary antibody solution containing 1% BSA

in PBS-T and 1:1000 donkey-anti-rabbit 647 (for α-mCherry primary antibody) or 1:1000 goat-anti-rabbit-

Alexa Fluor 555 antibody (for α-Vi-capsule primary antibody, Invitrogen) for overnight at 4°C. Final

washes included four PBS-T washes and one PBS wash at RT. The larvae were then mounted on agarose

and imaged on an Olympus SpinSR-10 Yokogawa spinning disk confocal microscope as described in the

section on confocal imaging.

Purification of His-RpoS protein

His$_6$-tagged RpoS protein was purified from an *E. coli* DH5α strain transformed with the plasmid pUHE21-

lacI^q::*rpoS*[66] (expresses RpoS from STm 14028s strain which is 100% identical to RpoS from STy H58)

following the protocol as described in Kim et al., Front Microbiol, 2021, with the following modifications:

A 30 ml culture was grown in LB broth containing 100 μg/mL ampicillin till 0.6 OD$_{600}$ and induced with

0.05 mM isopropyl-β-D-1-galactopyranoside (IPTG) for 7 hours. After sonication cells were resuspended

in a lysis buffer (Buffer A) containing 50 mM NaH$_2$PO4 and 300 mM NaCl, pH 8.0. A Nickel-

Nitrilotriacetic acid (Ni-NTA, GOLDBIO) column was washed with water and equilibrated in Buffer A

containing 30 mM imidazole. The cell lysates were incubated in Ni-NTA resin for an hour, pelleted and

washed thrice with Buffer A containing 30 mM imidazole. Protein fractions were eluted in Buffer A

containing 300 mM imidazole and dialyzed using a 6 to 8K MWCO dialysis tubing at 4 °C in a buffer

containing 20mM Tris-HCl, pH 8.0, 150 mM NaCl, 0.1mM EDTA, 5mM dithiothreitol and 20% v/v

glycerol for 2 to 3 hours. The dialyzed protein was quantified by measuring the absorbance at 280 nm using

a NanoDrop (Thermo Fisher Scientific) and aliquots were stored at -20 °C for immediate use or at -80 °C

for future use.

Electrophoretic mobility shift assay (EMSA)

Electrophoretic mobility shift assays were performed using the LightShift™ Chemiluminescent EMSA Kit

(Thermo Fisher Scientific) according to the manufacturer's protocol. Primers used for amplifying the

upstream regulatory regions of *waaZ*, *sthC*, *tviA* and *tviB* are described in Supplementary Table 2.

Biotinylated forward primers were used for generating respective biotinylated DNA fragments by PCR.

Each 20 μl binding reaction contained 1 ng DNA, 2 μL binding buffer, 1 μl 1μg/μL poly dI.dC, 1 μL 50%

glycerol, 1 μL 1M KCl and 1 μL 1% w/v NP-40. 1.4 μM purified RpoS, 125 ng non-biotinylated *waaZ* or

sthC DNA and 160 ng non-biotinylated *tviA* and *tviB* were added when appropriate.

Statistics

The Prism 10 software was used to plot all graphs and perform statistical analyses.

Acknowledgements:

S.D. and L.J.K. thank the following for their assistance; Stephen Baker (University of Cambridge) and Tran

Vu Thieu Nga for the H58 Tn library, Swaine Chen, Kurosh Mehershahi and Varnica Khetrapal for initial

attempts at TraDIS at the Genome Institute of Singapore, Zhixia Ding (Department of Pathology, UTMB)

and Dezhi Wang (The Texas Center for Superconductivity, University of Houston) for SEM sample

preparations and image acquisitions, Hyunjin Yoon and Eunsuk Kim, Ajou University, South Korea, for

the His-RpoS construct, Leslie Morgan in the LJK group for His-RpoS purification and Roy Curtiss III and

Soo-Young Wanda, University of Florida, for the *rpoS* construct. S.K.D. and R.D. thank the Zebrafish

International Resource Center (ZIRC) for fish lines and April Freeman (ZIRC) for advice on fish husbandry.

Z.Y. and A.L.R. were supported by a subaward from U54 Center Grant, AI170855, NIH/NIAID. L.J.K.

acknowledges support from UTMB Start up Funds, Texas Star Award, and Cancer prevention and Research

Institute of Texas (CPRIT) Grant RP2000650.

1447

1448

1449

1450

1451

1452 **Supplementary Table 1: List of bacterial strains and plasmids**
1453

Strain or plasmid	Description	Source or Reference
14028s	*Salmonella enterica* serovar Typhimurium	Lab collection
H58 ('wild type' in this study)	*Salmonella enterica* serovar Typhi	Stephen Baker, Cambridge University, UK
Ty2-b	*Salmonella enterica* serovar Typhi strain SGSC 2408	Salmonella Genetic Stock Center, University of Calgary, CA
CT18	*Salmonella enterica* serovar Typhi strain SGSC 4072	Salmonella Genetic Stock Center, University of Calgary, CA
CT117	*Salmonella enterica* serovar Typhi	Stephen Baker, Cambridge University, UK
ssrB	H58 (*ssrB*::*kan*)	This study
csgD	H58 (*csgD*::*cat*)	This study
csgA	H58 (*csgA*::*kan*)	This study
yihO	H58 (*yihO*::*kan*)	This study
yihP	H58 (*yihP*::*kan*)	This study
yihO yihP	H58 (Δ*yihO yihP*::*kan*)	This study
iraP	H58 (*iraP*::*kan*)	This study
rpoS	H58 (*rpoS*::*kan*)	This study
sthC	H58 (*sthC*::*kan*)	This study
waaZ	H58 (*waaZ*::*kan*)	This study
tviA	H58 (*tviA*::*cat*)	This study
tviB	H58 (*tviD*::*kan*)	This study
sthC tviD	H58 Δ*sthC* (*tviD*::*kan*)	This study
waaZ sthC	H58 Δ*waaZ* (*sthC*::*cat*)	This study
rpoSc	H58 (*rpoS*::*kan*, pBR322::*rpoS*)	This study
pUHE21-lacIq::*rpoS*	pUHE21-lacIq construct expressing His6-RpoS from 14028s	Kim et al., 2021[62]
pBR322::*rpoS* plasmid	*rpoS* cloned between EcoRI and ScaI sites in pBR322	Roy Curtiss III, University of Florida, USA
pFPV::mCherry plasmid	mCherry cloned between XbaI and SphI sites in pFPV	Olivia Steele-Mortimer (Addgene plasmid # 20956 http://n2t.net/addgene:20956; RRID:Addgene_20956

1454
1455
1456 **Supplementary Table 2: List of oligonucleotides**
1457

Primers	Sequence
Tn-ClickSeq	
3'21-39	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTTAATACGACTCACTATAGG [Black = Illumina p7 adapter, red = Tn-specific]
Gene deletions	

csgD knockout forward	GCAGCTGTCAGATGTGCGATTAAAAAAAGTGGAGTTTCATCATGTTTAATGTGTAGGCTGTAGCTGCTTC
csgD knockout reverse	CTCTGCTGCTACAATCCAGGTCAGATAGCGTTTCATGGCCTTACCGCCTGCCTCCTTAGTTCCTATTCCG
csgA knockout forward	CACCCAACGCTAATACCGTTACGACTTTTAAATCAATCAATCCGGTGTAGGCTGGAGCTGCTTC
csgA knockout reverse	AGGGCTTATGCCCTGTTTTTTTATTAGCGCAGACGCTAAACATATGAATATCCTCCTTAG
yihO knockout forward	ATGTCTAATCATGATCCGCTAACGCTAAAGTTGAGCCTGCGTGTAGGCTGGAGCTGCTTC
yihO knockout reverse	TTAATTATTTACAGTAGAAATACTTTGTTTATTATTAGTTCATATGAATATCCTCCTTAG
yihP knockout forward	GAGAAGAATAATGAGTCAAACATCTGTGTAGGCTGGAGCTGCTTC
yihP knockout reverse	GTTATATTTTATTGTTGTTAAACCGTATGCATATGAATATCCTCCTTA
iraP knockout forward	ATGAAAAATCTCATAGCAGAGTTGTTGCTTAAGCTAGCCCGTGTAGGCTGGAGCTGCTTC
iraP knockout reverse	TTAGTGCCGGGGGGTGTCTCAGCAACTTTTTTACATATTGGCATATGAATATCCTCCTTAG
rpoS knockout forward	ATGAGTCAGAATACGCTGAAAGTTCATGATTTAAATGAAGGTGTAGGCTGGAGCTGCTTC
rpoS knockout reverse	TTACTCGCGGAACAGCGCTTCGATATTCAGCCCCTGCGTCCATATGAATATCCTCCTTAG
tviA knockout forward	ATGAGGTTTCATCATTTCTGGCCTCCGAATGATATCTATTGTGTAGGCTGGAGCTGCTTC
tviA knockout reverse	TTACAGTAAAGTAACTGAATCCGGCAATAACAGATAGCGCCATATGAATATCCTCCTTAG
tviD knockout forward	ATGAATTTAATGAAATCGTCAGGGATGTTTACGCTTACAGGTGTAGGCTGGAGCTGCTTC
tviD knockout reverse	TTACGACTTCCCTGATGTATTTTTTTGTAATGCGGTTATGCATATGAATATCCTCCTTAG
waaZ knockout forward	ATGGGCAGCGTTAACTTCATAACTCACGCCGATGTTCTGCGTGTAGGCTGGAGCTGCTTC
waaZ knockout reverse	CTAGACAATTTTATCGTAATATTTCATCTTCAAGTTCCGACATATGAATATCCTCCTTAG
sthC knockout forward	TTGATCGCCGACCCGCGAGAATTCGTCCCTACCAGCAAATGTGTAGGCTGGAGCTGCTTC
sthC knockout reverse	TCACTGGCATTGCTCGTGTAAGATTTCTACGCCAGAAGCGCATATGAATATCCTCCTTAG
csgA forward	CACCCAACGCTAATACCGTT
csgA reverse	AGGGCTTATGCCCTGTTTTT
csgD forward	CAGCTGTCAGATGTGCGATT
csgD reverse	TCTGCTGCTACAATCCAGGT
yihO forward	TGCGGGGCGTTTTGAGAGGCGA
yihO reverse	TGAATAAAGCGGCAAGCGTCG

yihP forward	GTTGCGATTGGACGCTGTACCTG
yihP reverse	GTAATATGCAGGCATCCCGAGTTC
iraP forward	AGTGATAACGTCACCCTGGAAC
iraP reverse	AGTAACGTTATAACAACTGTGT
rpoS forward	CAGTCTGTCGACTGGCCTTT
rpoS reverse	CTAGTTCCGTCAAGGGATCA
tviA forward	AGGTTATTTCAGCATAAGGA
tviA reverse	TGTCCGTGTTTTACTCAATA
tviD forward	CTCGGTATAACTACTCACTT
tviD reverse	TTCCTAGTGCAGCTAACT
waaZ forward	GACATACTTGAGAGAATTTG
waaZ reverse	TGCGTGCCGAAGCAACGCAA
sthC forward	ACCGGCTGGATTTAGCGATC
sthC reverse	CATGATGCCAGACCCGTGAA
RT-qPCR	
rrsA internal forward	GCACCGGCTAACTCCGTGCC
rrsA internal reverse	GCAGTTCCCAGGTTGAGCCCG
rpoS internal forward	CCTGCGTCTGGTGGTAAA
rpoS internal reverse	TTCTCGACTGCACGGATAAG
waaZ internal forward	CCGCTATTCAGGTTGCCTATTC
waaZ internal reverse	AGGGCTGGTAGATTCGTCATAG
sthC internal forward	GATGAAGACGACGATACGGAAG
sthC internal reverse	CGTCTTTCGCGGAGTTCATA
tviB internal forward	TGTGGTAAAGGAACTCGGTAAA
tviB internal reverse	GACTTCCGATACCGGGATAATG
EMSA	
Biotin-waaZpro forward	Biotin - GCTGACTGACTTTTATTTGC
waaZpro reverse	ACTCGTATGTTTATCATGCA
waaZpro forward	GCTGACTGACTTTTATTTGC
Biotin-sthCpro forward	Biotin - ACTTCCCCGAGCTTAAAAAT

sthCpro reverse	CGCGATTTGAGGGATCGCTA
sthCpro forward	ACTTCCCCGAGCTTAAAAAT
Biotin-tviApro forward	Biotin - CGTTAGTACTATTAAAATTAGG
tviApro reverse	CTCCTTATGCTGAAATAACCTAA
tviApro forward	CGTTAGTACTATTAAAATTAGG
Biotin-tviBpro	Biotin-GTACGGTTATACGTTTTCAT
tviBpro reverse	CACTCTTATTAATCCTTTACT

1458

1459

1460

1461

1462

1463

1464

1465

1466

1467

1468

1469

1470

1471

1472

1473

1474

1475

1476

1477

1478

References

References

1.   Dougan, G. and S. Baker, *Salmonella enterica serovar Typhi and the pathogenesis of typhoid fever.*
Annu Rev Microbiol, 2014. **68**: p. 317-36.
2.   Typhoid, G.B.D. and C. Paratyphoid, *The global burden of typhoid and paratyphoid fevers: a*
systematic analysis for the Global Burden of Disease Study 2017. Lancet Infect Dis, 2019. **19**(4):
p. 369-381.
3.   Cirillo, D.M., et al., *Macrophage-dependent induction of the Salmonella pathogenicity island 2*
type III secretion system and its role in intracellular survival. Mol Microbiol, 1998. **30**(1): p. 175-
88.
4.   Feng, X., et al., *The response regulator SsrB activates transcription and binds to a region*
overlapping OmpR binding sites at Salmonella pathogenicity island 2. Mol Microbiol, 2004. **54**(3):
p. 823-35.
5.   Walthers, D., et al., *The response regulator SsrB activates expression of diverse Salmonella*
pathogenicity island 2 promoters and counters silencing by the nucleoid-associated protein H-NS.
Mol Microbiol, 2007. **65**(2): p. 477-93.
6.   Boddicker, J.D., et al., *Differential binding to and biofilm formation on, HEp-2 cells by Salmonella*
enterica serovar Typhimurium is dependent upon allelic variation in the fimH gene of the fim gene
cluster. Mol Microbiol, 2002. **45**(5): p. 1255-65.
7.   Prouty, A.M., W.H. Schwesinger, and J.S. Gunn, *Biofilm formation and interaction with the*
surfaces of gallstones by Salmonella spp. Infect Immun, 2002. **70**(5): p. 2640-9.
8.   Crull, K., et al., *Biofilm formation by Salmonella enterica serovar Typhimurium colonizing solid*
tumours. Cell Microbiol, 2011. **13**(8): p. 1223-33.
9.   Miller, A.L., et al., *In vivo synthesis of bacterial amyloid curli contributes to joint inflammation*
during S. Typhimurium infection. PLoS Pathog, 2020. **16**(7): p. e1008591.
10.   Desai, S.K., et al., *The horizontally-acquired response regulator SsrB drives a Salmonella lifestyle*
switch by relieving biofilm silencing. Elife, 2016. **5**.
11.   Desai, S.K. and L.J. Kenney, *To ~P or Not to ~P? Non-canonical activation by two-component*
response regulators. Mol Microbiol, 2017. **103**(2): p. 203-213.
12.   Gerstel, U., C. Park, and U. Romling, *Complex regulation of csgD promoter activity by global*
regulatory proteins. Mol Microbiol, 2003. **49**(3): p. 639-54.
13.   Gerstel, U. and U. Romling, *The csgD promoter, a control unit for biofilm formation in Salmonella*
typhimurium. Res Microbiol, 2003. **154**(10): p. 659-67.
14.   Romling, U., et al., *Curli fibers are highly conserved between Salmonella typhimurium and*
Escherichia coli with respect to operon structure and regulation. J Bacteriol, 1998. **180**(3): p. 722-
31.
15.   Zogaj, X., et al., *The multicellular morphotypes of Salmonella typhimurium and Escherichia coli*
produce cellulose as the second component of the extracellular matrix. Mol Microbiol, 2001. **39**(6):
p. 1452-63.
16.   Gibson, D.L., et al., *Salmonella produces an O-antigen capsule regulated by AgfD and important*
for environmental persistence. J Bacteriol, 2006. **188**(22): p. 7722-30.
17.   Desai, S.K. and L.J. Kenney, *Switching Lifestyles Is an in vivo Adaptive Strategy of Bacterial*
Pathogens. Front Cell Infect Microbiol, 2019. **9**: p. 421.
18.   Gunn, J.S., et al., *Salmonella chronic carriage: epidemiology, diagnosis, and gallbladder*
persistence. Trends Microbiol, 2014. **22**(11): p. 648-55.
19.   Crawford, R.W., et al., *Gallstones play a significant role in Salmonella spp. gallbladder*
colonization and carriage. Proc Natl Acad Sci U S A, 2010. **107**(9): p. 4353-8.
20.   Basnyat, B. and S. Baker, *Typhoid carriage in the gallbladder.* Lancet, 2015. **386**(9998): p. 1074.
21.   Wong, V.K., et al., *Phylogeographical analysis of the dominant multidrug-resistant H58 clade of*
Salmonella Typhi identifies inter- and intracontinental transmission events. Nat Genet, 2015.
47(6): p. 632-9.

22. Rowe, B., L.R. Ward, and E.J. Threlfall, *Multidrug-resistant Salmonella typhi: a worldwide*
epidemic. Clin Infect Dis, 1997. **24 Suppl 1**: p. S106-9.
23. Carey, M.E., et al., *The origins of haplotype 58 (H58) Salmonella enterica serovar*
Typhi. bioRxiv, 2022: p. 2022.10.03.510628.
24. Harrell, J.E., et al., *Salmonella Biofilm Formation, Chronic Infection, and Immunity Within the*
Intestine and Hepatobiliary Tract. Front Cell Infect Microbiol, 2020. **10**: p. 624622.
25. Gomes, M.C. and S. Mostowy, *The Case for Modeling Human Infection in Zebrafish.* Trends
Microbiol, 2020. **28**(1): p. 10-18.
26. Torraca, V. and S. Mostowy, *Zebrafish Infection: From Pathogenesis to Cell Biology.* Trends Cell
Biol, 2018. **28**(2): p. 143-156.
27. Berg, R.D. and L. Ramakrishnan, *Insights into tuberculosis from the zebrafish model.* Trends Mol
Med, 2012. **18**(12): p. 689-90.
28. Ramakrishnan, L., *The zebrafish guide to tuberculosis immunity and treatment.* Cold Spring Harb
Symp Quant Biol, 2013. **78**: p. 179-92.
29. Varas, M., et al., *Live-cell imaging of Salmonella Typhimurium interaction with zebrafish larvae*
after injection and immersion delivery methods. J Microbiol Methods, 2017. **135**: p. 20-25.
30. Varas, M., et al., *Salmonella Typhimurium induces cloacitis-like symptomsin zebrafish larvae.*
Microb Pathog, 2017. **107**: p. 317-320.
31. Davis, J.M., et al., *Real-time visualization of mycobacterium-macrophage interactions leading to*
initiation of granuloma formation in zebrafish embryos. Immunity, 2002. **17**(6): p. 693-702.
32. Mostowy, S., et al., *The zebrafish as a new model for the in vivo study of Shigella flexneri*
interaction with phagocytes and bacterial autophagy. PLoS Pathog, 2013. **9**(9): p. e1003588.
33. Brannon, M.K., et al., *Pseudomonas aeruginosa Type III secretion system interacts with*
phagocytes to modulate systemic infection of zebrafish embryos. Cell Microbiol, 2009. **11**(5): p.
755-68.
34. Torraca, V., et al., *Shigella serotypes associated with carriage in humans establish persistent*
infection in zebrafish. J Infect Dis, 2023.
35. Leiba, J., et al., *Dynamics of macrophage polarization support Salmonella persistence*
in a whole living organism. bioRxiv, 2023: p. 2023.05.09.539693.
36. Ganjali Dashti, M., et al., *Optimization of Salmonella Typhi biofilm assay on polypropylene*
microtiter plates using response surface methodology. Biofouling, 2016. **32**(4): p. 477-87.
37. MacKenzie, K.D., et al., *Parallel evolution leading to impaired biofilm formation in invasive*
Salmonella strains. PLoS Genet, 2019. **15**(6): p. e1008233.
38. Ou, C., C.M. Dozois, and F. Daigle, *Differential regulatory control of curli (csg) gene expression*
in Salmonella enterica serovar Typhi requires more than a functional CsgD regulator. Sci Rep,
2023. **13**(1): p. 14905.
39. Crawford, R.W., et al., *Identification of a bile-induced exopolysaccharide required for Salmonella*
biofilm formation on gallstone surfaces. Infect Immun, 2008. **76**(11): p. 5341-9.
40. Hammar, M., Z. Bian, and S. Normark, *Nucleator-dependent intercellular assembly of adhesive*
curli organelles in Escherichia coli. Proc Natl Acad Sci U S A, 1996. **93**(13): p. 6562-6.
41. Goodall, E.C.A., et al., *The Essential Genome of Escherichia coli K-12.* mBio, 2018. **9**(1).
42. Canals, R., et al., *The fitness landscape of the African Salmonella Typhimurium ST313 strain*
D23580 reveals unique properties of the pBT1 plasmid. PLoS Pathog, 2019. **15**(9): p. e1007948.
43. Langridge, G.C., et al., *Simultaneous assay of every Salmonella Typhi gene using one million*
transposon mutants. Genome Res, 2009. **19**(12): p. 2308-16.
44. Holden, E.R., et al., *Massively parallel transposon mutagenesis identifies temporally essential*
genes for biofilm formation in Escherichia coli. Microb Genom, 2021. **7**(11).
45. Nolan, L.M., et al., *A global genomic approach uncovers novel components for twitching motility-*
mediated biofilm expansion in Pseudomonas aeruginosa. Microb Genom, 2018. **4**(11).
46. Jaworski, E., et al., *Tiled-ClickSeq for targeted sequencing of complete coronavirus genomes with*
simultaneous capture of RNA recombination and minority variants. Elife, 2021. **10**: p. e68479.

47.   Routh, A., et al., *ClickSeq: Fragmentation-Free Next-Generation Sequencing via Click Ligation of*
1581   *Adaptors to Stochastically Terminated 3'-Azido cDNAs.* J Mol Biol, 2015. **427**(16): p. 2610-6.
1582   48.   Tran, V.T.N., *The humoral response against Salmonella Typhi protein antigens during acute,*
1583   *convalescent, and chronic typhoid fever.* 2019. p. xiv, 288 p.
1584   49.   Kim, D., et al., *Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype.*
1585   Nat Biotechnol, 2019. **37**(8): p. 907-915.
1586   50.   Townsend, S.M., et al., *Salmonella enterica serovar Typhi possesses a unique repertoire of fimbrial*
1587   *gene sequences.* Infect Immun, 2001. **69**(5): p. 2894-901.
1588   51.   Dufresne, K., J. Saulnier-Bellemare, and F. Daigle, *Functional Analysis of the Chaperone-Usher*
1589   *Fimbrial Gene Clusters of Salmonella enterica serovar Typhi.* Front Cell Infect Microbiol, 2018.
1590   **8**: p. 26.
1591   52.   Frirdich, E., et al., *Overexpression of the waaZ gene leads to modification of the structure of the*
1592   *inner core region of Escherichia coli lipopolysaccharide, truncation of the outer core, and*
1593   *reduction of the amount of O polysaccharide on the cell surface.* J Bacteriol, 2003. **185**(5): p. 1659-
1594   71.
1595   53.   Waxin, H., et al., *Identification of six open reading frames in the Salmonella enterica subsp.*
1596   *enterica ser. Typhi viaB locus involved in Vi antigen production.* Res Microbiol, 1993. **144**(5): p.
1597   363-71.
1598   54.   Pickard, D., et al., *Composition, acquisition, and distribution of the Vi exopolysaccharide-encoding*
1599   *Salmonella enterica pathogenicity island SPI-7.* J Bacteriol, 2003. **185**(17): p. 5055-65.
1600   55.   McGarry, N., D. Roe, and S.G.J. Smith, *Synergy between Group 2 capsules and lipopolysaccharide*
1601   *underpins serum resistance in extra-intestinal pathogenic Escherichia coli.* Microbiology
1602   (Reading), 2024. **170**(8).
1603   56.   Bougdour, A., S. Wickner, and S. Gottesman, *Modulating RssB activity: IraP, a novel regulator of*
1604   *sigma(S) stability in Escherichia coli.* Genes Dev, 2006. **20**(7): p. 884-97.
1605   57.   Gottesman, S., *Trouble is coming: Signaling pathways that regulate general stress responses in*
1606   *bacteria.* J Biol Chem, 2019. **294**(31): p. 11685-11700.
1607   58.   Desai, S.K., et al., *Salmonella biofilms program innate immunity for persistence in Caenorhabditis*
1608   *elegans.* Proc Natl Acad Sci U S A, 2019. **116**(25): p. 12462-12467.
1609   59.   MacKenzie, K.D., et al., *Bistable Expression of CsgD in Salmonella Connects Virulence to*
1610   *Persistence.* Infect Immun, 2015.
1611   60.   Anwar, N., et al., *Modulation of biofilm-formation in Salmonella enterica serovar Typhimurium by*
1612   *the periplasmic DsbA/DsbB oxidoreductase system requires the GGDEF-EAL domain protein*
1613   *STM3615.* PLoS One, 2014. **9**(8): p. e106095.
1614   61.   Gallo, P.M., et al., *Amyloid-DNA Composites of Bacterial Biofilms Stimulate Autoimmunity.*
1615   Immunity, 2015. **42**(6): p. 1171-84.
1616   62.   Santander, J., K.L. Roland, and R. Curtiss, 3rd, *Regulation of Vi capsular polysaccharide synthesis*
1617   *in Salmonella enterica serotype Typhi.* J Infect Dev Ctries, 2008. **2**(6): p. 412-20.
1618   63.   Peano, C., et al., *Characterization of the Escherichia coli sigma(S) core regulon by Chromatin*
1619   *Immunoprecipitation-sequencing (ChIP-seq) analysis.* Sci Rep, 2015. **5**: p. 10469.
1620   64.   Weber, H., et al., *Genome-wide analysis of the general stress response network in Escherichia coli:*
1621   *sigmaS-dependent genes, promoters, and sigma factor selectivity.* J Bacteriol, 2005. **187**(5): p.
1622   1591-603.
1623   65.   Shimada, T., K. Tanaka, and A. Ishihama, *The whole set of the constitutive promoters recognized*
1624   *by four minor sigma subunits of Escherichia coli RNA polymerase.* PLoS One, 2017. **12**(6): p.
1625   e0179181.
1626   66.   Kim, S.I., E. Kim, and H. Yoon, *sigma(S)-Mediated Stress Response Induced by Outer Membrane*
1627   *Perturbation Dampens Virulence in Salmonella enterica serovar Typhimurium.* Front Microbiol,
1628   2021. **12**: p. 750940.

1629	67.	Monack, D.M., D.M. Bouley, and S. Falkow, *Salmonella typhimurium persists within macrophages in the mesenteric lymph nodes of chronically infected Nramp1+/+ mice and can be reactivated by IFNgamma neutralization.* J Exp Med, 2004. **199**(2): p. 231-41.

1629   67.   Monack, D.M., D.M. Bouley, and S. Falkow, *Salmonella typhimurium persists within macrophages*
1630   *in the mesenteric lymph nodes of chronically infected Nramp1+/+ mice and can be reactivated by*
1631   *IFNgamma neutralization.* J Exp Med, 2004. **199**(2): p. 231-41.
1632   68.   Hernandez, R.E., et al., *Delay of Initial Feeding of Zebrafish Larvae Until 8 Days Postfertilization*
1633   *Has No Impact on Survival or Growth Through the Juvenile Stage.* Zebrafish, 2018. **15**(5): p. 515-
1634   518.
1635   69.   Marineli, F., et al., *Mary Mallon (1869-1938) and the history of typhoid fever.* Ann Gastroenterol,
1636   2013. **26**(2): p. 132-134.
1637   70.   Mortimer, P.P., *Mr N the milker, and Dr Koch's concept of the healthy carrier.* Lancet, 1999.
1638   **353**(9161): p. 1354-6.
1639   71.   Gopinath, S., S. Carden, and D. Monack, *Shedding light on Salmonella carriers.* Trends Microbiol,
1640   2012. **20**(7): p. 320-7.
1641   72.   Crawford, R.W., K.E. Reeve, and J.S. Gunn, *Flagellated but not hyperfimbriated Salmonella*
1642   *enterica serovar Typhimurium attaches to and forms biofilms on cholesterol-coated surfaces.* J
1643   Bacteriol, 2010. **192**(12): p. 2981-90.
1644   73.   Adcox, H.E., et al., *Salmonella Extracellular Matrix Components Influence Biofilm Formation and*
1645   *Gallbladder Colonization.* Infect Immun, 2016. **84**(11): p. 3243-3251.
1646   74.   Gonzalez, J.F., et al., *Human Bile-Mediated Regulation of Salmonella Curli Fimbriae.* J Bacteriol,
1647   2019. **201**(18).
1648   75.   Denger, K., et al., *Sulphoglycolysis in Escherichia coli K-12 closes a gap in the biogeochemical*
1649   *sulphur cycle.* Nature, 2014. **507**(7490): p. 114-7.
1650   76.   Speciale, G., et al., *YihQ is a sulfoquinovosidase that cleaves sulfoquinovosyl diacylglyceride*
1651   *sulfolipids.* Nat Chem Biol, 2016. **12**(4): p. 215-7.
1652   77.   Bougdour, A. and S. Gottesman, *ppGpp regulation of RpoS degradation via anti-adaptor protein*
1653   *IraP.* Proc Natl Acad Sci U S A, 2007. **104**(31): p. 12896-901.
1654   78.   Santander, J., et al., *Role of RpoS in fine-tuning the synthesis of Vi capsular polysaccharide in*
1655   *Salmonella enterica serotype Typhi.* Infect Immun, 2007. **75**(3): p. 1382-92.
1656   79.   Velasquez, J.C., et al., *SPI-9 of Salmonella enterica serovar Typhi is constituted by an operon*
1657   *positively regulated by RpoS and contributes to adherence to epithelial cells in culture.*
1658   Microbiology (Reading), 2016. **162**(8): p. 1367-1378.
1659   80.   Romling, U., et al., *Occurrence and regulation of the multicellular morphotype in Salmonella*
1660   *serovars important in human disease.* Int J Med Microbiol, 2003. **293**(4): p. 273-85.
1661   81.   Newman, S.L., et al., *The curli regulator CsgD mediates stationary phase counter-silencing of*
1662   *csgBA in Salmonella Typhimurium.* Mol Microbiol, 2018. **108**(1): p. 101-114.
1663   82.   Arnqvist, A., et al., *The Crl protein activates cryptic genes for curli formation and fibronectin*
1664   *binding in Escherichia coli HB101.* Mol Microbiol, 1992. **6**(17): p. 2443-52.
1665   83.   Pratt, L.A. and T.J. Silhavy, *Crl stimulates RpoS activity during stationary phase.* Mol Microbiol,
1666   1998. **29**(5): p. 1225-36.
1667   84.   Typas, A., et al., *Stationary phase reorganisation of the Escherichia coli transcription machinery*
1668   *by Crl protein, a fine-tuner of sigmas activity and levels.* EMBO J, 2007. **26**(6): p. 1569-78.
1669   85.   Marshall, J.M., et al., *Visualization of extracellular matrix components within sectioned Salmonella*
1670   *biofilms on the surface of human gallstones.* PLoS One, 2014. **9**(2): p. e89243.
1671   86.   Devaraj, A., et al., *Enhanced biofilm and extracellular matrix production by chronic carriage*
1672   *versus acute isolates of Salmonella Typhi.* PLoS Pathog, 2021. **17**(1): p. e1009209.
1673   87.   Hahn, M.M., et al., *The Abundance and Organization of Salmonella Extracellular Polymeric*
1674   *Substances in Gallbladder-Mimicking Environments and In Vivo.* Infect Immun, 2021. **89**(11): p.
1675   e0031021.
1676   88.   Milligan-Myhre, K., et al., *Study of host-microbe interactions in zebrafish.* Methods Cell Biol,
1677   2011. **105**: p. 87-116.
1678   89.   Raza, A., et al., *Effect of biofilm formation on the excretion of Salmonella enterica serovar Typhi*
1679   *in feces.* Int J Infect Dis, 2011. **15**(11): p. e747-52.

1680   90.   Sambrook J, F.E., Maniatis T, *Molecular cloning: a laboratory manual*. 1989: Cold Spring Harbor
1681   Laboratory Press, Cold spring Harbor, NY.
1682   91.   Davis R, B.D., Roth J, *Advanced Bacterial Genetics*. 1980: Cold spring Harbor Laboratory, NY.
1683   16-18.
1684   92.   Karlinsey, J.E., *lambda-Red genetic engineering in Salmonella enterica serovar Typhimurium.*
1685   Methods Enzymol, 2007. **421**: p. 199-209.
1686   93.   Datsenko, K.A. and B.L. Wanner, *One-step inactivation of chromosomal genes in Escherichia coli*
1687   *K-12 using PCR products.* Proc Natl Acad Sci U S A, 2000. **97**(12): p. 6640-5.
1688   94.   Jaworski, E. and A. Routh, *ClickSeq: Replacing Fragmentation and Enzymatic Ligation with Click-*
1689   *Chemistry to Prevent Sequence Chimeras*, in *Next Generation Sequencing: Methods and Protocols*,
1690   S.R. Head, P. Ordoukhanian, and D.R. Salomon, Editors. 2018, Springer New York: New York,
1691   NY. p. 71-85.
1692   95.   Jaworski, E. and A. Routh, *Parallel ClickSeq and Nanopore sequencing elucidates the rapid*
1693   *evolution of defective-interfering RNAs in Flock House virus.* PLoS Pathog, 2017. **13**(5): p.
1694   e1006365.
1695   96.   Chen, S., et al., *fastp: an ultra-fast all-in-one FASTQ preprocessor.* Bioinformatics, 2018. **34**(17):
1696   p. i884-i890.
1697   97.   Martin, M., *Cutadapt removes adapter sequences from high-throughput sequencing reads.*
1698   EMBnet.journal; Vol 17, No 1: Next Generation Sequencing Data AnalysisDO -
1699   10.14806/ej.17.1.200, 2011.
1700   98.   Li, H., et al., *The Sequence Alignment/Map format and SAMtools.* Bioinformatics (Oxford,
1701   England), 2009. **25**(16): p. 2078-2079.
1702   99.   Smith, T., A. Heger, and I. Sudbery, *UMI-tools: modeling sequencing errors in Unique Molecular*
1703   *Identifiers to improve quantification accuracy.* Genome Res, 2017. **27**(3): p. 491-499.
1704   100.   Quinlan, A.R. and I.M. Hall, *BEDTools: a flexible suite of utilities for comparing genomic features.*
1705   Bioinformatics, 2010. **26**(6): p. 841-2.
1706   101.   Love, M.I., W. Huber, and S. Anders, *Moderated estimation of fold change and dispersion for*
1707   *RNA-seq data with DESeq2.* Genome Biol, 2014. **15**(12): p. 550.
1708   102.   Schmittgen, T.D. and K.J. Livak, *Analyzing real-time PCR data by the comparative C(T) method.*
1709   Nat Protoc, 2008. **3**(6): p. 1101-8.
1710   103.   Westerfield, M., *The zebrafish book. A guide for the laboratory use of zebrafish (Danio rerio)*. 4th
1711   ed ed. 2000, Eugene: Univ. of Oregon Press.
1712   104.   Santos, D., S.M. Monteiro, and A. Luzio, *General Whole-Mount Immunohistochemistry of*
1713   *Zebrafish (Danio rerio) Embryos and Larvae Protocol.* Methods Mol Biol, 2018. **1797**: p. 365-
1714   371.
1715