Reproductive character displacement explains strengthening of mechanical barriers in damselflies

ABSTRACT

Reinforcing natural selection against maladaptive hybrids can favor the strengthening of premating reproductive isolation driving a pattern of Reproductive Character Displacement (RCD). In a recent study conducted in North-West (NW) (older) Spanish hybrid zone, was detected an asymmetric reinforcement of the mechanical isolation in the reciprocal cross direction between *I. graellsii* males and *I. elegans* females. Furthermore, in the North-Central and Mediterranean (NCM) (younger) hybrid zone was also detected a similar strengthening of the mechanical isolation, consistent with a pattern of asymmetric reinforcement in this hybrid zone as well. In this study, we did geometric morphometrics analyses, shape, and Centroid Size (CS), on male and female secondary sexual traits to investigate whether reinforcement has generated a pattern of RCD of these traits in both hybrid zones. We detected, in the NW hybrid zone, unidirectional RCD (CS) of the male caudal appendages of *I. graellsii*, and bidirectional RCD (shape) of the female prothorax. Consistently with the prediction that the signal of reinforcement may diminish rapidly once reinforcement ceases to operate, a stronger signal of RCD was detected in the NCM than in the NW hybrid region. In this region, was detected unidirectional RCD (CS) of the male caudal appendages which was consistent with the lock-and-key mechanism of genital coevolution, as well as RCD (shape) of the female prothorax of *I. elegans*. Interestingly, our study highlights the importance of using geometric morphometrics to deal with the complexity of reproductive structures and controlling for environmental and geographic factors to investigate RCD.

Keywords: Reproductive Character Displacement, Reinforcement, Hybridization, damselflies, Mechanical Barriers, Geometric Morphometrics

INTRODUCTION

Character displacement was defined by Brown and Wilson (1956) as a pattern in which two sympatric species present greater differences in certain ecological or reproductive characters than they do in allopatry. When this pattern is produced by selection to avoid the excessive cost of maladaptive hybridization it is named "Reproductive Character Displacement" (RCD) (Servedio and Noor, 2003; Coyne and Orr, 2004). This process has been earlier named by Dobzhansky (1940) as "reinforcing reproductive isolation". Available empirical evidence suggests that RCD plays an important role in the evolution of diversity (Schluter, 2000; Pfennig and Pfennig, 2005). This mechanism has been reported in different organisms like insects (Brown and Wilson, 1956; Walker 1974; Kawano 2002), fishes (Schluter, 2000; Crampton et al., 2011; Roth-Monzón et al., 2020), birds (Brown and Wilson, 1956; Diamond et al., 1989; Seddon and Tobias, 2010), reptiles (Melville 2002; Dayan and Simberloff, 2005) and amphibians (Brown and Wilson, 1956; Johanet et al., 2009). Commonly, the presence of RCD in genital divergence between closely related species has been used as support for the lock-and-key hypothesis of genital evolution, i.e., as evidence of natural selection acting on genital morphology to prevent hybridization. Although the number of studies testing RCD on genital divergence has increased and are mainly focused on insects (e.g., Kawano, 2002; 2003; Usami et al., 2006; Augustijnen et al., 2022) and arachnids (Kuntner et al., 2009; Costa-Schmidt and Mellender de Araújo, 2010; Muster and Michalik, 2019), these studies have predominantly focused on male divergence, which have prevented a comprehensive investigation of lock-and-key hypothesis of genital evolution.

In damselflies, mechanical isolation due to the incompatibility to form a tandem position is the strongest component of prezygotic isolation (Sánchez-Guillén et al., 2014a; Nava-Bolaños et al., 2017; Wellenreuther and Sánchez-Guillén 2015; Barnard et al., 2017). In damselflies copula involves two contact points. Firstly, the male grasps the female by her prothorax using his caudal appendages, thus achieving the first contact point known as "tandem position". The second contact point implies female recognition, through tactile isolation: the female bends her abdomen bringing both genitalia into contact, thus achieving the second contact point known as "wheel position". There is great variation in the caudal male appendages among odonates. McPeek et al. (2011) observed significant differences in the shape of male cerci among six *Enallagma* species, but not among populations within species. Moreover, McPeek studies in *Enallagma* damselflies (McPeek et al. 2008; 2009) have provided empirical support for "lock-and-key" coevolutionary divergence in damselflies. In their research, McPeek et al. (2008, 2009) identified a correlated evolution between male and female secondary genitalia, specifically, male cerci and female mesostigmal plates. They quantified both tempo and mode of male and female genital evolution and they found a similar pattern of punctuated evolution in both structures, proposing that this pattern might be indicative of a lock-and-key mechanism in the tandem position (see also Paulson, 1974, Masly 2012).

The damselflies *Ischnura elegans* and *I. graellsii* overlap their distributions in north and central Spain, where they are in sympatry (see Figure 1). Currently, *I. elegans* has a patchy distribution in Spain, consistent with a mottled hybrid zone (Sánchez-Guillén et al. 2023). This sympatric distribution comprises two unconnected hybrid zones: the North-West (NW) hybrid zone, which extends along the north-west Spanish coast. *Ischnura elegans* was first discovered in this zone during the early 1980s (Ocharan, 1987). The second hybrid zone, a larger zone known as the North-Central and Mediterranean (NCM) hybrid zone, extends from norther and central Spain and along the Mediterranean coast. The earlier records of *I. elegans* in the Mediterranean zone are before 1900, although certain parts of the central zone, records are more recent, specifically from the early 2000s (Sánchez-Guillén et al., 2021b). In previous genomic studies, on-going hybridization and bilateral introgression were detected in the NW hybrid zone (Sánchez-Guillén et al. 2011; Wellenreuther et al. 2018; Sánchez-Guillén et al. 2023), whereas on-going hybridization but unilateral introgression (thorough *I. graellsii*) was detected in the NCM hybrid region. Recently, Arce-Valdés et al. (2023) tested the reinforcement of reproductive isolation in the NW hybrid zone by comparing the intensity of five prezygotic barriers in crosses between *I. elegans* and *I. graellsii* from allopatry *versus* sympatry, and by measuring the fitness of their hybrids. In the reciprocal cross involving *I. graellsii* males and *I. elegans* females, it was detected that the mechanical barrier implicated in tandem formation was more intense in sympatry than in allopatry, and the hybrids resulting from this cross direction exhibited reduced fitness compared to the parental species. However, in the other reciprocal cross direction, the one between *I. elegans* males and *I. graellsii* females, none of the estimated five prezygotic barriers were more intense in sympatry than in allopatry, and hybrids were not formed in allopatry by the accumulated action of these five reproductive barriers, but were formed in sympatry, possibly due to the purge of BDM incompatibilities. Therefore, Arce-Valdés et al. (2023) detected asymmetric reinforcement occurring only in the reciprocal cross involving *I. graellsii* males and *I. elegans* females. Although the reinforcement has not yet been tested in the NCM hybrid zone, the strength of the mechanical barrier reinforced in the NW hybrid zone was similar in strength to that in the NCM zone (unpublished data). This is consistent with the reinforcement of this mechanical barrier occurring in the NW hybrid zone.

Taking advantage of these features, we investigated whether the reinforcement of mechanical isolation in the reciprocal cross between *I. graellsii* males and *I. elegans* females in the NW and the NCM hybrid zones have generated a pattern of RCD in the secondary-genital divergence of both, *I. graellsii* males and *I. elegans* females. To this end, we performed geometric morphometrics analyses on both shape and Centroid Size (CS) of male and female secondary sexual structures of individuals of *I. elegans* and *I. graellsii* from allopatry and both hybrid zones (NW and NCM). Moreover, because of phenotypic characters may vary along environmental gradients (Goldberg and Lande, 2006), to discriminate between phenotypic differences caused by sympatry from those caused by

environmental factors we have controlled for environmental factors (Johanet et al., 2009).
Our first prediction is that we will detect a pattern of RCD in *I. graellsii* males and *I.*
elegans females within both hybrid zones. This prediction in based on the enhanced
mechanical isolation detected in previous studies (Arce-Valdés et al. 2023; unpublished-
data) but only in the reciprocal cross between *I. graellsii* males and *I. elegans* females
within both hybrid zones. Our second prediction is that RCD could also be detected in *I.*
graellsii females and *I. elegans* males within both hybrid zones. This prediction is based on
the lock-and-key coevolution of the genitalia, which will drive a genital coevolution as a
mechanism to maintain the integrity of species boundaries and to prevent hybridization (see
Sloan and Simmons, 2019). Our third prediction is that RCD will be unilateral or
asymmetric between species. This prediction is based on that the strength of RCD depends
on the interplay of several non-mutually exclusive factors: 1) the symmetry/asymmetry of
reinforcement of the reproductive isolation (cf. Cooley, 2007; Yikweon, 2008); 2)
proximity to parental populations, since hybrid populations closer to parental populations
experience higher levels of gene flow (Liou and Price 1994; Pfennig and Ryan 2006); the
3) the abundance of both parental species, since rarer species have a greater chance of
encountering/mating with the common species than *vice versa* (Peterson et al., 2005;
Hochkirch et al., 2007). Our fourth prediction is that the signal of RCD will be stronger in
the NCM than in the NW hybrid region because the signal of reinforcement may diminish
rapidly once reinforcement ceases to operate. This prediction is based on higher levels of
interspecific gene flow detected in the NW compared to NCM hybrid region (see Sánchez-
Guillén et al. 2023). Our fifth prediction is that RCD will also vary between populations
because of differences in local species proportions (see Sánchez-Guillén et al. 2023).

METHODOLOGY

Population samplings

During summers from 2003 to 2021, adult males and females were collected from 15
localities belonging to six allopatric localities, four localities from the NW hybrid zone, and
five localities from the NCM hybrid zone (Table 1; Figure 1). Populations from both hybrid
zones were selected for this study based on our knowledge of reproductive isolation and
hybridization patterns from previous studies. Arce-Valdés et al. (2023) detected the
strengthening of the mechanical isolation due to the mismatch between male and female
secondary genital structures to form the tandem in both conspecific crosses of both species,
and heterospecific crosses in sympatry with respect to allopatry. From each locality, we
sampled between 14-31 adult males and 9-28 adult females of both species depending on
the dominant species in each locality (Table 1). Sampled specimens were preserved in
ethanol (96%) and stored at -20°C.

Geometric morphometrics analyses

We used a two-dimensional geometric morphometric approach to investigate the RCD of
genital structures because it allows to break down the complexity in shape and CS of the
studied reproductive structures (Klingenberg, 2016). Thus, our analyses investigated RCD
in two views (lateral and posterior) of both shape and CS of male caudal appendages and
female prothorax because these are structures that can influence the ability to achieve
copulation and contribute to defining species-specificity (Simmons and García-Gonzalez,
2011; Song, 2009). In fact, morphological differences in the male's caudal appendages and
the female's prothorax are used in morphological identification of these species (Askew,
1988; Monetti et al., 2002). In addition, both structures are involved in the first point of
contact during damselfly reproduction which is known as the tandem position. During the
tandem male grasps the female by her thorax by using his caudal appendages to achieve
this position (Supplementary Figure S1A-B), thereafter, the second point of contact implies
a prior recognition of the female, which bends her abdomen putting into contact with both
genitalia, thus achieving the copula.

Photographs of the posterior and lateral view of the male's caudal appendages and the
female's prothorax were obtained under a stereoscopic microscope (Zeiss Stemi 305) with
an integrated camera (Axiocam ERc5s). The posterior view of the male's caudal
appendages shape was sampled with a combination of eight landmarks and 80 semi-
landmarks in four curves (left and right sides, Supplementary Figure S2A-B), which were
assigned above and below each of the cerci. The lateral view was sampled with seven
landmarks and 36 semi-landmarks in two curves which were assigned on the cercus and the
paraproct of the left side (Supplementary Figure S2C-D). Likewise, the posterior view of
the female's prothorax shape was sampled with a combination of four landmarks and 21
semi-landmarks in one curve (Supplementary Figure S3A-B), which was assigned
surrounding the pronotum. The left lateral view was sampled with 4 landmarks and 19
semi-landmarks in one curve (Supplementary Figure S3C-D), also surrounding the
pronotum Cartesian coordinates of each posterior and lateral view of males and females
were registered with tpsDIG2 version 2.04 (Rohlf, 2006): 88 landmarks in the posterior
view of the caudal appendages, and 43 landmarks in lateral view; 25 landmarks in posterior
view of pronotum and 23 landmarks in lateral view; in which x and y coordinates of the
geometric configurations were obtained for each image. The configuration of the landmarks
on each of the male and female reproductive structures of both species is described in table
S1.

From these, a Generalized Procrustes Analysis (GPA; Rohlf and Slice, 1990) was generated
to superimpose landmark configurations and variation due to differences in translation,
orientation, and size was removed followed by a thin-plate spline analysis (Bookstein 1991)
using the geomorph package version 4.0.4 (Adams et al. 2022). The GPA translates all
samples to an origin, rescales them to a CS, and rotates them by least squares to the
coordinates of the corresponding points, aligning them as closely as possible (Adams et al.,
2017). Semilandmarks were aligned using the Bending Energy minimization criteria, which

optimize the semilandmarks position with the lowest deformation energy from a consensus curve of reference (Gunz and Mitteroecker 2013). Additionally, shape variation of both reproductive structures (male's caudal appendages and female's prothorax) was then evaluated graphically along the first two axes of a Principal Components Analysis (PCA) and through deformation grids to facilitate anatomical description of any implied morphological changes using the same geomorph package version 4.0.4 (Adams et al. 2022).

Morphological variation: environmental and geographic factors and/or reinforcement of RI?

Ecological changes in habitat or resource use may generate changes in the species that promote RCD (Pfennig and Pfennig, 2009). Earlier studies in odonates, for instance, have shown that phenotypic variables, such as body size or wing size and shape show a latitudinal variation (Johansson, 2003; Hassall et al., 2008). Since our allopatric and sympatric studied populations are distributed in a latitudinal and longitudinal gradient, environmental factors may also be responsible for explaining their phenotypic characters (Goldberg and Lande, 2006). Thus, to discriminate between phenotypic differences caused by sympatry from those caused by environmental/geographic we tested whether environmental and geographic factors and sympatry might be correlated with the morphological characters measured.

To this end, we analyzed five environmental (precipitation, annual mean temperature, maximum temperature, minimum temperature, and elevation) and two geographic (latitude and longitude) factors. Environmental variables were chosen from those identified in previous studies as variables that contribute to the geographical distribution of *I. elegans* and that are involved in odonate selection processes (Lancaster et al., 2015; Dudaniec et al., 2018; 2021; Wellenreuther et al., 2012). Climate data were generated using bioclimatic parameters from the Worldclim database version 1.4 (Fick and Hijmans, 2017) (at 1km cell resolution). Rasters from each variable were obtained using the function 'getData' from the 'raster' package. Furthermore, we explored a correlation between environmental variables using the 'PerformanceAnalytics' package version 2.0.4 (Peterson et al., 2022). Thus, for both environmental and geographic data, when two variables presented an absolute value of correlation coefficients ($|r|$)>0.7, one of them was not included in the same model because the relationship between these is considered strong (see Dormann et al., 2013).

Finally, we fitted separate gaussian generalized linear models (GLMs) with an identity link function for the shape (described by the first principal component of shape variance to each geometric configuration), and for the CS of male and female reproductive structures in each species. All models included the geographic variables (latitude and longitude) because in all data groups, they showed correlation coefficient values lower than 0.7 (Table S2), then we added those environmental variables with less than 0.7 absolute correlation coefficient

with both geographic variables. When two environmental variables had low correlation

coefficients with the geographic variables but were highly correlated between them, we

selected the variable with the overall lowest correlation coefficients with the geographic

variables. For males and females of *I. elegans* and for males of *I. graellsii*, the explanatory

variables of the GLMs included one environmental variable (annual mean temperature), the

two geographical variables (latitude and longitude), and the study zones (allopatry, NW and

NCM hybrid zones) as a fixed effect to test for zone variation in individual phenotypic

responses to these environmental factors. While for *I. graellsii* females, the explanatory

variables used in the model were the elevation, the two geographical variables, and the

study zones (allopatry, NW and NCM hybrid zones) as a fixed effect. During model

selection, Corrected Akaike Information Criterion (AICc) and the significance of fixed

effects were used to distinguish among competing models. All analyses were conducted

using the statistical software R-project (version 3.6.2, R Core Team, 2019).

Reproductive character displacement by regions

To tested our first four predictions: 1) RCD will be detected in *I. graellsii* males and *I.*

elegans females by reinforcement of RI; 2) RCD could also be detected in *I. graellsii*

females and *I. elegans* females by lock-and-key coevolution; 3) RCD could be unilateral or

asymmetric between species because it depends on the interplay of several non-mutually

exclusive factors; and 4) RCD will be more pronounced in the younger NCM hybrid zone

because of the signal of reinforcement may diminish rapidly once reinforcement ceases to

operate, in the following manner.

Because RCD is a pattern whereby reproductive characters can already differ or not in

allopatry (Slatkin, 1980; Liou and Price, 1994), but the variance in distance between these

reproductive characters should be larger in sympatry than in allopatry, first we measured

the distance variance between *I. elegans* and *I. graellsii* from allopatry and from each

hybrid zone. When the variance in distance was larger in the hybrid zone compared to the

allopatric zone (distance in sympatry - distance in allopatry $\square$0) this provided evidence of

RCD (see Table 2). Second, in cases where RCD was detected given its potential for being

unilateral or bilateral, occurring in either one or both species we conducted intraspecific

comparisons (distance in sympatry *versus* distance in allopatry). Therefore, when the

variance in distance was significantly different only in one intraspecific comparison, we

inferred that RCD was asymmetric (occurring in one species), and when the variance in

distance was significantly different in both intraspecific comparisons, we inferred that RCD

was symmetric.

Statistical analyses were done in the following manner. Firstly, to assess the statistical

hypothesis investigating RCD patterns of shape variation and covariation for the set of

(Procrustes-aligned) coordinates, explained by CS, geographic group (allopatry, NW and

NCM hybrid zones) and their interaction we performed a procrustes ANOVA model using

the 'procD.lm' function from the geomorph package version 4.0.4 (Adams et al., 2022),

which calculates the Procrustes distances variance explained by each factor of the model.
Secondly, when the group factor showed statistical significance, we complemented this
analysis with a pairwise evaluation using the 'permudist' function from the 'morpho'
package version 2.8 (Schlager, 2017). This function compares the distance between two
group means with the distances obtained by random assignment (here 1,000 permutations)
of observations to the compared groups (Schlager, 2017). Thirdly, to assess differences in
CS we used the function 'lm.rrpp' from the RRPP package (Collyer and Adams et al.,
2018; 2021) to perform the linear models. When the group factor showed statistical
significance, we conducted posthoc pairwise comparisons to detect which groups were
statistically different using the 'pairwise' function.

Reproductive character displacement by localities

To test our fifth prediction, that RCD will vary between populations because of differences
in local species proportions (see Sánchez-Guillén et al. 2023) we separately analyzed the
shape and CS of male caudal appendages and female prothorax in each population.

To assess the statistical hypothesis investigating RCD patterns of shape variation and
covariation for the set of (Procrustes-aligned) coordinates, explained by CS, locality, and
their interaction we performed a procrustes ANOVA model using the 'procD.lm' function
from the geomorph package version 4.0.4 (Adams et al., 2022). When the locality showed
statistical significance, we complemented this analysis with a pairwise evaluation using the
'permudist' function from the 'morpho' package version 2.8 (Schlager, 2017). To assess
the statistical hypothesis investigating RCD patterns of CS variation we used the function
'lm.rrpp' from the RRPP package (Collyer and Adams et al., 2018; 2021) to perform the
linear models. When the locality factor showed statistical significance, we conducted
posthoc pairwise comparisons to detect which localities were statistically different using
the 'pairwise' function.

Lock-and key coevolution of male-female genitalia

We evaluated the role of the lock-and-key on the coevolution of male and female genitalia
of *I. elegans* and *I. graellsii* males and females given RCD was detected in both males and
females (see Table 2 for detailed results). To this end, we followed two approximations.

Our first approximation was based on that the lock-and-key mechanism, which predicts that
genital coevolution results from reinforcement selection, as the formation of new species
requires reproductive isolation (Masly 2012). Reinforcing natural selection represents a
stabilizing force that acts to strengthen RI between species. As a result, it is expected to
promote allometric relationships between body size and genital size. This occurs because it
pushes to maintain an equilibrium between these two traits, ensuring optimal reproductive
compatibility. To test this prediction, we examined a subset of the data, which included one
allopatric population of each species (Cyprien for *I. elegans* and Gamillazo for *I. graellsii*)
and two populations from each hybrid zone (NW hybrid zone: Laxe for *I. elegans* and

Cachadas for *I. graellsii*, and the NCM hybrid zone: Perdiguero for *I. elegans* and Valpierre for *I. graellsii*). We investigated whether male and female genital size changed allometrically in relation to body length by using wing length as a proxy for body size (Nava et al., 2011; 2014; Córdoba et al., 2015). This approach was necessary due to abdominal deformation during preservation in ethanol. We included a set of 10 males and 10 females from each population, zone, and species. The wings were dissected using fine scissors and examined under a stereoscopic microscope (Zeiss Stemi 305), and we measured the hindwing length between two points: the arculus and the proximal corner of the pterostigma (Carchini et al., 2000) using IMAGEJ software (version 1.53k, National Institutes of Health, MD, USA). To keep a constant measurement error, the wing length was measured by the same person. We performed a Pearson correlation between CS genital size and hindwing length. Both the CS data and the wing length data were log-transformed to linearize the relationship between genital traits and body size indicators. We used R software (R Development Core Team 2009) for statistical analysis.

Secondly, we directly examined the lock-and-key genital coevolution (shape and CS, and both views) of the complementary male caudal appendages and female prothorax in allopatry as well as in both hybrid zones. To test this prediction, we examined a subset of data, in which we select the same number of males as females. Using randomization in R, we adjusted the number of pairs according to the minimum number of individuals of either sex in each population. To assess the genital coevolution of the shape of male and female genital structures, we conducted a two-block partial least squares analysis on Procrustes shape variables using paired samples with 100 iterations, utilizing the 'two.b.pls' function from the geomorph package version 4.0.4 (Adams et al. 2022). To assess the genital coevolution of the CS of male and female genital structures, we performed a Pearson correlation with 100 iterations. All analyses were done with the statistical software R-project (version 3.6.2, R Core Team, 2019).

RESULTS

Morphological variation: environmental and geographic factors and/or reinforcement of RI?

A total of 192 *I. elegans* males and 137 *I. graellsii* males were included in the morphometric analyses of male caudal appendages, and a total of 134 *I. elegans* females and 135 *I. graellsii* females were included in the morphometric analyses of the prothorax. Table 3 includes all statistical GLM results explaining which part of the morphological variation detected on male and female secondary genitalia was explained by the annual mean temperature or elevation (as they were the only uncorrelated and statistically significant environmental factors), the two geographic factors (latitude and longitude) and the distribution type (allopatry, NW and NCM hybrid zones).

In *I. elegans* males GLM modeling and scoring using the AICc suggested that the morphological variation of the shape, at both views, of their caudal appendages was influenced by the annual mean temperature, the longitude, and the distribution type, while the morphological variation of the CS differed between views: the lateral view was influenced by the annual mean temperature, the longitude, and the distribution type; while for the lateral view only the longitude and the distribution type were included in the best model. According to the post-hoc comparisons, based on the shape (both views) the allopatric zone was different from both hybrid zones (Table 3 and S3), while based on the CS (both views), the allopatric zone was only different from the NCM hybrid zone (Table 3 y S3). Otherwise, in *I. elegans* females GLM modeling and scoring using the AIC suggested that the morphological variation of the shape, at both views, of their prothorax was not influenced by any environmental or geographical variable, only the distribution type was included in the best model, while the morphological variation of the CS, at both views, was influenced by the longitude and the distribution type. Similarly, to that detected in males, post-hoc comparisons based on the shape (both views) the allopatric zone was different from the hybrid zones (Table 3 y S3), while based on the CS (both views), the allopatric zone was only different from the NCM hybrid zone (Table 3 y S3).

In *I. graellsii* males GLM modeling and scoring using the AICc suggested that the morphological variation of the lateral view of the shape of their caudal appendages was influenced by the distribution type, while the morphological variation of the CS differed between views: the lateral view was influenced by the longitude and the distribution type, while for the posterior view only the distribution type was included in the best model. According to the post-hoc comparisons, based on the shape (lateral view) the allopatric zone was different from both hybrid zones (Table 3 and S3), while based on the CS the allopatric zone was only different from the NCM hybrid zone for the posterior view, and was only different from the NW hybrid zone for the lateral view (Table 3 y S3). Otherwise, in *I. graellsii* females GLM modeling and scoring using the AIC suggested that the morphological variation of the shape, at both views, of their prothorax was not influenced by any environmental variable or the distribution type, only by the geographical variables were included in the best model: the longitude at both views and the latitude only the lateral view (Table 3), similarly, the morphological variation of the CS differed between views: for the posterior view, the best model included only the latitude, while for the lateral view, the best model included only the distribution type. Post-hoc comparisons based on the CS (lateral vie) indicated differences between the allopatric zone and the hybrid zones (Table 3 y S3).

Shape variation of male caudal appendages

PCA plots of the two views (posterior and lateral) showed a clear separation between species in the three zones (Table S4, Figure 2A-F). Procrustes ANOVA showed statistically significant differences between *I. elegans* and *I. graellsii*: 1) from the allopatric zone

(posterior view: F=326.66, *p*<0.001, d=0.31; lateral view: F=144.25, *p*<0.001, d=0.22); 2)
from the NCM hybrid zone (posterior view: F=226.17, *p*<0.001, d=0.26; lateral view:
F=90.29, *p*<0.001, d=0.18); and 3) from the NW hybrid zone (posterior view: F=318.78,
p<0.001, d=0.28; lateral view: F=89.06, *p*<0.001, d=0.19) (Table S5-S6). However,
distance variance (both views) did not support the RCD, as the distance variance in both
hybrid zones the NW and the NCM hybrid zones was not larger than that in the allopatric
zone (distance sympatry - distance allopatry □0) (Table 2).

CS variation of male caudal appendages

Procrustes ANOVA showed statistically significant differences between *I. elegans* and *I.*
graellsii from allopatry, particularly in the posterior view (posterior view F=18.603,
p<0.001, d=41.634; lateral view F=0.017, *p*=0.894, d=1.231), and from the NCM (posterior
view: F=29.581, *p*<0.001, d=59.347; lateral view: F=5.151, *p*=0.018, d=23.529) and NW
(posterior view: F=6.465, *p*=0.013, d=25.324; lateral view: F=13.629, *p*<0.001, d=31.454)
hybrid zones (Tables S7-S8, Figure 3A-B). The CS had significant but low effects in all the
models (6 to 16% of shape variance, *p*<0.01) and the lowest variances were explained by
the interaction between CS and groups (2 to 4%). Distance variance analysis confirmed
RCD in both views in the NCM but only in the lateral view in the NW hybrid zone, where
the difference between sympatry and allopatry distances was greater than zero (Table 2).

To investigate whether the detected RCD was unidirectional or bidirectional, i.e., present in
one or both species, we conducted comparisons within each species. In *I. elegans*, the CS
distance was significantly different among the three zones (posterior view: F=31.345,
p<0.001; lateral view F=15.194, *p*<0.001; respectively, Table S9; Figure 3A-B). Post hoc
analysis showed that the caudal appendages of males from the NCM hybrid zone were
significantly larger than those from allopatry (posterior view: z=4.146, p<0.001, d=64.323;
lateral view: z=3.444, p<0.001, d=45.350; Table S10), while the males from the NW hybrid
zone were not significantly different compared to males from allopatry (Table S10).
Similarly, in *I. graellsii*, the CS distance was significantly different among the three zones
(posterior view: F=14.533, *p*<0.001; lateral view F=4.080, *p*=0.011; Table S9; Figure 3A-
B). Post hoc analysis showed that the caudal appendages of males from the NCM hybrid
zone were significantly larger than those from allopatry (posterior view: z=3.265, *p*<0.001,
d=46.610; lateral view: z=1.809, *p*=0.023, d=20.588, Table S10), while the males from the
NW hybrid zone were only significantly larger than those from allopatry for the lateral
view (z=2.219, *p*=0.003, d=25.891). Therefore, RCD (both views) was unilateral, as it was
only detected in *I. elegans* from the NCM hybrid zone, while although RCD (lateral view)
was also unilateral in the NW hybrid zone, it was only detected in the *I. graellsii* males.
(Table 2).

Shape variation of female Prothorax

413 PCA plots of the two views (posterior and lateral) showed a clear separation between
414 species in the three zones (Table S11, Figure 2G-L). Procrustes ANOVA showed

statistically significant differences between *I. elegans* and *I. graellsii*: from the allopatric zone (posterior view: F=45.47, *p*<0.001, d=0.12 and lateral view: F=21.21, *p*<0.001, d=0.19); from the NCM hybrid zone (posterior view: F =106.19, p<0.001, d=0.22 and lateral view: F=15.02, p<0.001, d=0.30); and from the NW hybrid zone (posterior view: F=144.23, *p*<0.001, d=0.19 and lateral view: F=42.90, *p*<0.001, d=0.27) (Tables S12-S13). Distance variance analysis confirmed RCD in both views and hybrid regions the NW and the NCM, as distance variance was larger than that in the allopatric zone (distance sympatry - distance allopatry □0) (Table 2).

Within species in *I. elegans,* PC axis1 explained 62.9% (posterior view) and 43.6% (lateral view), and PC axis 2 the 12.4% (posterior view) and 21.7% (lateral view) of the shape variance. They were mainly discriminated in the first axis (Supplementary Figure S4). Procrustes ANOVA showed statistically significant differences between zones (posterior view: F=15.83, *p*<0.001); lateral view: F=13.80, *p*<0.001)] (Table S14). Post hoc paired tests showed statistically significant differences between *I. elegans* individuals from the allopatric zone and the NCM hybrid zone (posterior view: *p*<0.001; d=0.08, and lateral view: *p*<0.001; d=0.11), and between *I. elegans* individuals from the allopatric and the NW hybrid zone at the lateral view (posterior view: *p*=0.486; d=0.03; lateral view: *p*=0.036; d=0.06) (Table S15). Similarly, within species in *I. graellsii*, PC axis1 explained 34.1% (posterior view) and 43.7% (lateral view), and PC axis 2 the 29.3% (posterior view) and 25.1% (lateral view) of the shape variance. They were mainly discriminated in the first axis (Supplementary Figure S4). Procrustes ANOVA showed statistically significant differences between the three zones (Posterior view: F=2.27, *p*=0.008 and lateral view: F=2.18, *p*=0.026) (Table S14). Post hoc paired tests only detected statistically significant differences between individuals of *I. graellsii* from the allopatric and NW hybrid zone in the lateral view (posterior view: *p*=0.470; d=0.02 lateral view: *p*=0.011; d=0.05, Table S15). Therefore, RCD (both views) was unilateral, as it was only detected in *I. elegans* from the NCM hybrid zone, while RCD (lateral view) was bilateral in the NW hybrid zone (Table 2).

CS variation of female Prothorax

Procrustes ANOVA showed statistically significant differences between *I. elegans* and *I. graellsii* from allopatry (posterior view: F=24.228, *p*<0.001, d=33.346, and lateral view: F=122.45, *p*<0.001, d=55.109), from the NCM hybrid zone (posterior view: F=54.319, *p*<0.001, d=55.194, and lateral view: F=415.07, *p*<0.001, d=93.158), and from the NW hybrid zone (posterior view: F=0.052, *p*<0.811, d=2.083, lateral view: F=100.34, *p*<0.001, d=52.052) (Tables S16-S17, Figure 3C-D). Distance variance analysis confirmed RCD in both views only in the NCM hybrid zone, where the difference between sympatry and allopatry distances was greater than zero. In contrast, in the NW hybrid zone, the difference between sympatry and allopatry distances was less than zero (Table 2).

Within species, in *I. elegans,* the CS distance of the of female prothoraxes from the NCM
hybrid zone was significantly larger than those from allopatry (posterior view: z=3.389,
p<0.001, d=38.990; lateral view: z=4.357, *p*<0.001, d=51.163; Table S18). Within species,
in *I. graellsii,* although the CS distance in the lateral view of females from the NCM hybrid
zone (lateral view: z=2.607, *p*<0.001, d=13.115) was significantly larger than those of
females from allopatry, they shifted in the same direction than *I. elegans* did, i.e., so that I.
graellsii females did not contribute to RCD. In summary, RCD (both views) was unilateral,
only detected in *I. elegans* females from the NCM hybrid zone (Table 2).

Reproductive character displacement at the local scale

Consistently with our fourth prediction that RCD will vary in its strength between
populations because of differences in local species proportions (see Sánchez-Guillén et al.
2023). Results are given by shape or CS, sexes, and views (Figure 4).

Shape variation of the male caudal appendages

Procrustes ANOVA applied to assess the shape of the male caudal appendages showed
statistically significant differences between localities of both species and in both views: *I.*
elegans posterior view: F=5.840, *p*<0.001, and lateral view: F=5.704, *p*<0.001; and *I.*
graellsii posterior view: F=4.166, *p*<0.001, and lateral view: F=3.176, *p*<0.001 (Table
S19). Posthoc pairwise comparisons between sympatric and allopatric populations of *I.*
elegans, showed statistically significant differences for two localities in the NW zone
(Doniños and Laxe) and for three localities in the NCM zone (Perdiguero, Valpierre, and
Villar) in both the posterior and lateral views, except for Valpierre in the lateral view. In *I.*
graellsii, posthoc pairwise comparisons showed statistically significant differences for two
localities in the NW zone (Cachadas and Xuño) and for three localities in the NCM zone
(Mateo, Plaiaundi, and Valpierre) in both the posterior and lateral views (Table S20).

Shape variation of female prothorax

Procrustes ANOVA for the shape of the female prothorax showed statistically significant
differences between localities of both species, *I. elegans* (posterior view: F=6.471, *p*<0.001,
and lateral view: F=7.248, *p*<0.001) and *I. graellsii* (posterior view: F=1.786, *p*=0.004, and
lateral view: F=1.798, *p*=0.016) (Table S21). In *I. elegans* posthoc pairwise comparisons
(sympatric populations *versus* allopatric populations) showed statistically significant
differences for three out of the four localities from the NCM hybrid zone (posterior view:
Perdiguero and Villar; and lateral view: Valpierre, Perdiguero and Villar), and the two
localities from the NW hybrid zone (lateral view: Doniños and Laxe). In *I. graellsii*,
posthoc pairwise comparisons showed statistically significant differences for two localities
from the NW hybrid zone (posterior view: Cachadas and Xuño; lateral view: Cachadas) and
two localities from NCM hybrid zone (posterior view: Valpierre and Mateo; lateral view:
Mateo) (Table S22).

CS variation of male caudal appendages

Procrustes ANOVA for the CS of the male caudal appendages showed statistically
significant differences between localities of both species *I. elegans* (posterior view:
F=25.352, *p*<0.001, and lateral view: F=14.13, *p*<0.001) (Table S23) and *I. graellsii*
(posterior view: F=6.930, *p*<0.001, and lateral view: F=3.083, *p*=0.004) (Table S23). In *I.*
elegans, posthoc pairwise comparisons (sympatric populations *versus* allopatric
populations) showed statistically significant differences for three out of the four localities
from the NCM hybrid zone (both views: Villar, Perdiguero, and Plaiaundi). In *I. graellsii*,
posthoc pairwise comparisons showed statistically significant differences for the three
populations from the NCM hybrid zone (posterior view: Valpierre, Mateo and Plaiaundi),
(Table S24, Figure 4).

CS variation of female prothorax

Procrustes ANOVA for the CS of the female prothorax showed statistically significant
differences between localities of both species *I. elegans* (posterior view: F=15.355,
p<0.001, and lateral view: F=16.460, *p*<0.001) (Table S25) and *I. graellsii* (posterior view:
F=0.933, *p*=0.469, and lateral view: F=2.213, *p*=0.044) (Table S25). In *I. elegans*, posthoc
pairwise comparisons (sympatric populations *versus* allopatric populations) showed
statistically significant differences for two localities from the NCM hybrid zone (both
views: Villar, Perdiguero), (Table S26, Figure 4). In *I. graellsii*, posthoc pairwise
comparisons (sympatric populations *versus* allopatric populations) showed non-statistically
significant differences for any population from the NCM hybrid zone.

Evidence of the lock-and-key coevolution

Consistently with our prediction that reinforcement selection which is stabilizing force, will
promote allometric relationships between body size and genital size, we detected a positive
allometry between the CS (posterior view) of the caudal appendages of the *I. graellsii*
males and the proxy to body size in the NW (Cachadas: r =0.851; *p* =0.001) and in the
NCM hybrid zone (Valpierre: r = 0.636; *p* = 0.047), and between the CS (posterior view) of
the *I. elegans* females prothorax and the proxy to body size in the NW (Laxe: r=0.657;
p=0.039), and in the NCM (Perdiguero: r=0.749; *p*=0.012) hybrid zone. However, in the
allopatric populations that correlations were not detected (Table S27).

Consistently with our prediction of the lock-and-key coevolution between the male and
female genital, we detected moderate coevolution between male caudal appendages and
female prothorax in allopatry, but this correlation was slightly lower in both hybrid zones,
probably because the different force of the reinforcement and the lock-and key. In detail,
for the CS we detected a moderate percentage of coevolution in allopatry: 19% and 20%,
posterior and lateral views, respectively in *I. elegans*, and 19% and 4% posterior and lateral
views, respectively in *I. graellsii*. However, in both hybrid zones, especially in the NCM,

that correlations were considerably reduced: in the NW hybrid zone, 13% and 2%, posterior and lateral views, respectively in *I. elegans*, and 6% and 5% posterior and lateral views, respectively in *I. graellsii;* in the NCM hybrid zone, 7% and 2%, posterior and lateral views, respectively in *I. elegans*, and 1% and 6% posterior and lateral views, respectively in *I. graellsii.* However, for the shape we detected a low percentage of coevolution in allopatry: 3% and 5%, posterior and lateral views, respectively in *I. elegans*, and 7% and 5% posterior and lateral views, respectively in *I. graellsii.* However, in both hybrid zones, especially in the NW, that correlations were slightly increased: in the NW hybrid zone, 6% and 10%, posterior and lateral views, respectively in *I. elegans*, and 6% and 4% posterior and lateral views, respectively in *I. graellsii;* in the NCM hybrid zone, 5% and 5%, posterior and lateral views, respectively in *I. elegans*, and 1% and 6% posterior and lateral views, respectively in *I. graellsii* (Table S28).

DISCUSSION

The main criterion for RCD is that divergence in reproductive structures is greater in sympatry than in allopatry (see Brown and Wilson 1956).

Divergence in reproductive traits via sexual selection may play a key role in determining the crossability between incipient species (Panhuis et al. 2001; Ritchie 2007). While male genitalia have traditionally been recognized as the most variable and divergent morphological structure, a growing body of empirical evidence supports coevolutionary processes, a pattern of divergent evolution of female's genitalia matching males' variation (Simmons and Garcia-Gonzalez 2011, Macagno et al., 2011; Simmons and Fitzpatrick, 2019). Mechanical isolation hypothesis predicts that physical incompatibility between primary genitalia of males and females hinders hybridization (so-called ''lock-and-key'' or ''structural mechanical isolation'', Dufour, 1848; Coyne and Orr 2004). Contact and grasping structures that are not intromissive have been included in the category of genital traits (Brennan 2016). In damselflies, species recognition takes place during tandem via tactile interactions, and the female can then accept or refuse to cooperate with the male (Wellenreuther and Sánchez-Guillén, 2016). Moreover, male caudal appendages and female prothorax must match to form a tandem, thus any change not only in size but also in shape will affect the mismatch of these structures. Considering the complexity of these structures we analyzed both the shape and the CS in two complementary views (posterior and lateral) to capture most of the variation.

We directly measured the lock-and-key genital coevolution (shape and CS, and both views) of the complementary male caudal appendages and female prothorax in allopatry as well as both hybrid zones, although coevolution should not be limited to morphological correspondence (Brennan and Prum 2015). We found that while the proportion of interactions showing a positive correlation between the CS of male and female structures was moderate in allopatry (19-20%), it was higher than in each hybrid zone. Note that only the CS analysis (not shape analysis) was useful to investigate the mechanism of lock-and

key RI because RCD of the shape was only detected in females. Moreover, consistent with our prediction that genital coevolution results from reinforcing natural selection which represents a stabilizing force that acts to strengthen RI between species, and as a result, it is expected to promote allometric relationships between body size and genital size, we detected a positive allometry between the CS of the *I. graellsii* males and the *I. elegans* females from both hybrid zones.

Our results showed that the more variation was analyzed, the more likely RCD was detected: while the CS was useful to find RCD in both males' caudal appendages and females' prothorax, the shape was only useful to find RCD on the female's prothorax. In a similar manner, the analyses of the two different views of the males' and females' genital structures were useful to test the prediction of RCD: while both views (lateral and posterior) were useful to find RCD on the shape of the females' prothorax, only the lateral view was useful to find RCD on the CS of the males' caudal appendages and the females' prothorax. By other hand, the use of geometric morphometrics was opportune since odonate males usually tend to show a pattern of lower allometry than females, i.e., the genital size changes little relative to body size (Nava-Bolaños et al. 2014), which was recovered from our statistical analyses.

Reinforcing selection in response to maladaptive hybridization in hybrid zones can lead to a pattern of RCD of traits involved in premating isolation in sympatry (Howard 1993; Coyne and Orr 2004). However, ecological changes in habitat or resource use may generate changes in the species that also promote RCD (Pfennig and Pfennig, 2009). We detected that both environmental and geographic factors can in part explain several detected patterns of variation in the CS and shape in both hybrid zones and species. Interestingly, the RCD detected of the shape of the *I. graellsii* females was only explained by both geographic factors, latitude, and longitude. Thus, our results highlight the importance of investigating environmental and geographic factors when testing RCD.

RCD pattern can be asymmetric (occurring only in one species or being more pronounced in one of the involved species/sexes) or symmetric (in both involved species). The direction of this pattern depends on the strength of selection to avoid hybridization over each species. As we predicted, based on the reinforcement of the mechanical isolation to form the tandem position in the reciprocal cross direction between *I. graellsii* males and *I. elegans* females (see Arce-Valdés et al., 2023), controlling for environmental and geographic factors we detected that in the NW hybrid zone, CS of the caudal appendages of *I. elegans* and *I. graellsii* males differed more than in allopatry, mainly due to an increase in these structures among the *I. graellsii* males indicating unidirectional RCD, and that shape of the prothorax of *I. elegans* and *I. graellsii* females differed more than in allopatry, resulting from changes in females of both species indicating bidirectional RCD.

Consistently with the prediction that the signal of reinforcement may diminish rapidly once reinforcement ceases to operate, a stronger signal of RCD was detected in the younger

NCM hybrid zone than in the older NW hybrid region: CS of the male caudal appendages as well as CS and shape of the female prothorax of *I. elegans* and *I. graellsii* differed more than in allopatry. These differences were due to the increase of these structures in both males and females, which was consistent with the lock-and-key coevolution, but only of *I. elegans* indicating unidirectional RCD. For instance, the beetles *Odontolabis mouhoti* and *O. cuvera* show more divergence in body size, genitalia length and coloration in sympatry than in allopatry; in this system, both species have shifted their relative sizes in opposite directions, i.e., in a bilateral pattern of RCD (Kawano, 2003). However, the size of genital traits of two closely related *Ohomopterus* species differed between populations in contact, however, although both male and female of *C. maiyasanus* increased in sympatry with *C. iwawakianus*, those of *C. iwawakianus* did not differ between allopatric and sympatric populations, indicating a unilateral RCD and providing novel evidence for the lock-and-key hypothesis of genital evolution in beetles (Nishimura et al., 2022).

Asymmetric patterns of RCD arise from different factors such as the asymmetric reinforcement of the reproductive isolation (cf. Cooley, 2007; Yikweon, 2008), the abundance of both parental species, since rarer species have a greater chance of encountering/mating with the common species than *vice versa* (Peterson et al., 2005; Hochkirch et al., 2007), and the level of gene flow with the parental populations (homogenization by gene flow) (Liou and Price 1994; Pfennig and Ryan 2006). When *I. elegans* and *I. graellsii* came into secondary contact, natural selection to avoid hybridization may be stronger for the reciprocal cross between *I. graellsii* males and *I. elegans* females. This is because prezygotic barriers allow hybrid formation, and hybrids experience more disadvantages than the parental species (Arce-Valdés et al, 2023). In contrast, the formation of hybrids from the reciprocal cross between *I. elegans* males and *I. graellsii* females from allopatry is almost completely precluded by gametic isolation (Arce-Valdés et al, 2023). In fact, in the NW hybrid zone (see Arce-Valdés et al, 2023) and the NCM hybrid zone (personal observation) after at least 50 years of sympatry, the strength of the reciprocal crosses between *I. graellsii* males and *I. elegans* females has been enhanced, to the point to be almost completely prevented by the increase of the incompatibility between the male caudal appendages and the female prothorax to form the tandem position. On the contrary, this has not occurred in the reciprocal cross between *I. elegans* males and *I. graellsii* females, among whom the intensity of mechanical isolation is similar between allopatry and sympatry, but the intensity of the postzygotic isolation is lower in sympatry than in allopatry, possibly due to the purge of Bateson–Dobzhansky–Muller (BDM) incompatibilities in sympatry (see Arce-Valdés et al, 2023).

Additionally, *I. elegans* is probably the rarer species in each new Spanish hybrid locality because the secondary contact between *I. elegans* and *I. graellsii* is the result of the range expansion of *I. elegans* over the *I. graellsii* distribution in Spain, so that, the rarer *I. elegans* females have a greater chance of encountering with the common *I. graellsii* males than *vice versa,* and thus be involved in heterospecific matings (Sánchez-Guillén et al. 2023; Arce-

Valdés et al, 2023). For instance, *Pterostichus thunbergi* and *P. habui* beetles show unilateral RCD pattern of genital morphology. *Pterostichus thunbergi* differed more from *P. habui* at sympatry than at allopatry, likely because of *P. habui* arrives at the contact zone in small numbers, and thus, would experience stronger selection pressures than *P. habui* (Kosuda et al., 2016). Additionally, interspecific gene flow between *I. elegans* and *I. graellsii* can also explain the increase in the CS of male caudal appendages in *I. graellsii* from the NW hybrid zone, and thus, explain the reduction of the distance variance between *I. elegans* and *I. graellsii* males, because of caudal appendages of hybrids are intermediate between those of the two species (Monetti et al. 2002). In fact, the measured genetic divergence between *I. elegans* and *I. graellsii* in NW hybrid zone was lower than in the NCM hybrid zones and allopatry indicating higher level of interspecific gene flow in the NW than in the NCM hybrid zone (Sánchez-Guillén et al. 2023). A similar finding was detected in the *Carabus, C. maiyasanus* and *C. iwawakianus*, in which hybrid genital traits were intermediate to those of the two species, thus explaining elongation of the genital traits of *C. iwawakianus* (Sasabe et al. 2007).

At the local scale, RCD in the CS and shape of male caudal appendages and female prothorax was detected in all localities (except to in the CS of *I. graellsii* females). Both hybrid zones have introgressed populations of each parental species, i.e., bimodal distribution, and hybrid populations with F_1, F_2, and backcrosses, i.e., unimodal distribution (Sánchez-Guillén et al. 2022). According to this, we would expect a higher relative cost of the hybridization, and thus, RCD will more probably be detected in *I. elegans* females than in *I. graellsii* females (see Kyogoku and Wheatcroft, 2020), but because of the mottled pattern of the hybrid zones, the strength and/or direction of the RCD can lack of consistency between hybrid zones and even between localities (Harrison, 1993). Moreover, the outcome of reproductive interference in sympatric localities is density-dependent (Westman et al., 2002; Hochkirch et al., 2007) thus the species at lower density will experience stronger selective pressure as a result of reproductive interference (Okuzaki, 2013).

Conclusions

Our study presents novel results in nonterritorial damselflies with few visual displays, but species recognition through mechanic-tactile stimuli, and consequently almost random mating attempts among congeneric species in sympatry. We found evidence of RCD and evidence of the lock-and-key hypothesis of genital evolution in *Ischnura elegans and I. graellsii*. RCD was in part explained by environmental and geographic factors, and in part explained by enhanced reproductive isolation between *I. elegans* females and *I. graellsii* males. Our study adds to previous empirical studies from which has emerged a consensus on the taxonomically widespread role of RCD to lessen reproductive interactions between species (see Goldberg and Lande, 2006). Moreover, our study highlights the importance of

discarding environmental and geographic factors when testing RCD and using geometric morphometrics to deal with the complexity of reproductive structures.

REFERENCES

Abràmoff, M. D., Magalhães, P. J., and Ram, S. J. (2004). Image processing with ImageJ. *Biophotonics international*, 11(7), 36-42.

Adams, D. C. (2004). Character displacement via aggressive interference in Appalachian salamanders. *Ecology*, 85(10), 2664-2670.

Adams, D., Collyer, M., Kaliontzopoulou, A., Baken, E. (2022). "Geomorph: Software for geometric morphometric analyses. R package version 4.0.4." https://cran.r-project.org/package=geomorph

Arce-Valdés, L. R., Ballén-Guapacha, V. A., Chávez-Ríos, J. R., Wellenreuther, M., Hansson, B., and Sánchez-Guillén, R. A. (under review). Testing the predictions of reinforcement: long-term empirical data from a damselfly mosaic hybrid zone Journal: *Evolution*.

Askew, A. R. (1988). The dragonflies of Europe. England.

Augustijnen, H., Patsiou, T., and Lucek, K. (2022). Secondary contact rather than coexistence—Erebia butterflies in the Alps. *Evolution*, 76(11), 2669-2686.

Barnard, A. A., Fincke, O. M., McPeek, M. A., and Masly, J. P. (2017). Mechanical and tactile incompatibilities cause reproductive isolation between two young damselfly species. *Evolution*, 71(10), 2410-2427.

Barton, N. H. and Hewitt, G. M. (1989). Adaptation, speciation, and hybrid zones. *Nature*, 341(6242), 497.

Bates, D., Maechler, M., Bolker, B., Walker, S., Christensen, R. H. B., Singmann, H., ... and Scheipl, F. (2012). Package 'lme4'. CRAN. R Foundation for Statistical Computing, Vienna, Austria.

Bookstein, F. L. 1991. Morphometric tools for landmark data: geometry and biology. *Cambridge University Press*, Cambridge, UK.

Boudot JP, Kalkman VJ, Azpilicueta-Amorín M, Bogdanovic T, Dommanget L, and Ferreira S, (2009) Odonata of the Mediterranean and North Africa. *Libellula Supplement*. 9: 1–256.

Brennan, P. L., and Prum, R. O. (2015). Mechanisms and evidence of genital coevolution:
the roles of natural selection, mate choice, and sexual conflict. *Cold Spring Harbor*
Perspectives in Biology, 7(7), a017749.

Brennan, P. L. (2016). Studying genital coevolution to understand intromittent organ
morphology. Integrative and comparative biology, 56(4), 669-681.Brown, W. L., and
Wilson, E. O. (1956). Character displacement. *Systematic Zoology*, 5(2), 49-64.

Carchini, G., Chiarotti, F., Di Domenico, M., and Paganotti, G. (2000). Fluctuating
asymmetry, size and mating success in males of *Ischnura elegans* (Vander
Linden)(Odonata: Coenagrionidae). *Animal Behaviour*, 59(1), 177-182.

Comeault, A. A., and Matute, D. R. (2016). Reinforcement's incidental effects on
reproductive isolation between conspecifics. *Current Zoology*, 62(2), 135-143.

Cooley, J. R. (2007). Decoding Asymmetries in Reproductive Character Displacement.
Proceedings of the Academy of Natural Sciences of Philadelphia, 156(1), 89–96.

Costa-Schmidt, L. E., and de Araújo, A. M. (2010). Genitalic variation and taxonomic
discrimination in the semi-aquatic spider genus Paratrechalea (Araneae: Trechaleidae). *The*
Journal of Arachnology, 38(2), 242-249.

Collyer, M. L., Adams, D. C. (2021). "RRPP: Linear Model Evaluation with Randomized
Residuals in a Permutation Procedure, R package version 1.1.2." https://cran.r-
project.org/package=RRPP.

Collyer, M. L, Adams, D. C. (2018). "RRPP: An R package for fitting linear models to
high□dimensional data using residual randomization."

Coyne, J. A., and Orr, H. A. (1989). Patterns of speciation in Drosophila. *Evolution*, 43(2),
362-381.

Coyne, J. A., and Orr, H. A. (1997). "Patterns of speciation in Drosophila" revisited.
Evolution, 51(1), 295-303.

Coyne JA, and Orr HA (2004) Speciation. Book. Sinauer, Sunderland, MA de Wilde, J.,
and de Loof, A. (1973). Reproduction. In *The physiology of Insecta* (pp. 11-95). Academic
Press.

Crampton, W. G., Lovejoy, N. R., and Waddell, J. C. (2011). Reproductive character
displacement and signal ontogeny in a sympatric assemblage of electric fish. *Evolution:*
International Journal of Organic Evolution, 65(6), 1650-1666.

Dayan, T., and Simberloff, D. (2005). Ecological and community□wide character
displacement: the next generation. *Ecology letters*, 8(8), 875-894.

Diamond, J., Pimm, S. L., Gilpin, M. E., and LeCroy, M. (1989). Rapid evolution of
character displacement in myzomelid honeyeaters. *The American Naturalist*, 134(5), 675-
708.

Dobzhansky, T. (1940). Speciation as a stage in evolutionary divergence. *The American
Naturalist*, 74(753), 312-321.

Dormann, C. F., Elith, J., Bacher, S., Buchmann, C., Carl, G., Carré, G., ... and Lautenbach,
S. (2013). Collinearity: a review of methods to deal with it and a simulation study
evaluating their performance. *Ecography*, 36(1), 27-46.

Dudaniec, R. Y., Yong, C. J., Lancaster, L. T., Svensson, E. I., and Hansson, B. (2018).
Signatures of local adaptation along environmental gradients in a range□expanding
damselfly (*Ischnura elegans*). *Molecular Ecology*, 27(11), 2576-2593.

Dudaniec, R. Y., Carey, A. R., Svensson, E. I., Hansson, B., Yong, C. J., and Lancaster, L.
T. (2021). Latitudinal clines in sexual selection, sexual size dimorphism and sex□specific
genetic dispersal during a poleward range expansion. *Journal of Animal Ecology*, 91(6),
1104-1118.

Dufour, L. (1844). Anatomie générale des Dipteres. *Ann. Sci. Nat*, 1, 244-264.

Eberhard, W. G. (1992). Species isolation, genital mechanics, and the evolution of
species□specific genitalia in three species of Macrodactylus beetles (Coleoptera,
Scarabaeidae, Melolonthinae). *Evolution*, 46(6), 1774-1783.

Fick, S.E. and R.J. Hijmans, 2017. WorldClim 2: new 1km spatial resolution climate
surfaces for global land areas. *International Journal of Climatology* 37 (12): 4302-4315.

Goldberg, E., and Lande, R. (2006). Ecological and reproductive character displacement of
an environmental gradient. *Evolution*, 60(7), 1344-1357.

González□Tokman, D., Córdoba□Aguilar, A., Dáttilo, W., Lira□Noriega, A.,
Sánchez□Guillén, R. A., and Villalobos, F. (2020). Insect responses to heat: physiological
mechanisms, evolution, and ecological implications in a warming world. *Biological
Reviews*, 95(3), 802-821.

Gunz, P., and Mitteroecker, P. (2013). Semilandmarks: a method for quantifying curves and
surfaces. *Hystrix, the Italian journal of mammalogy*, 24(1), 103-109.

Hassall, C., Thompson, D. J., and Harvey, I. F. (2008). Latitudinal variation in morphology
in two sympatric damselfly species with contrasting range dynamics (Odonata:
Coenagrionidae). *European Journal of Entomology*, 105(5), 939–944.

Harrison, R. G., and Harrison, R. G. (Eds.). (1993). Hybrid zones and the evolutionary
process. *Oxford University Press on Demand*.

Harrison, R. G., and Larson, E. L. (2014). Hybridization, introgression, and the nature of
species boundaries. *Journal of Heredity*, 105(S1), 795-809.

Hendry, A. P., Wenburg, J. K., Bentzen, P., Volk, E. C., and Quinn, T. P. (2000). Rapid
evolution of reproductive isolation in the wild: evidence from introduced salmon. *Science*,
290(5491), 516-518.

Hochkirch, A., Gröning, J., and Bücker, A. (2007). Sympatry with the devil: reproductive
interference could hamper species coexistence. *Journal of Animal Ecology*, 76(4), 633-642.

Hollander J, Montaño-Rendón M, Bianco G, Yang X, Westram AM, Duvaux L, Reid DG,
Butlin RK (2018). Are assortative mating and genital divergence driven by reinforcement?
Evolution Letters 16;2(6):557-566.

Hoskin, C. J., Higgie, M., McDonald, K. R., and Moritz, C. (2005). Reinforcement drives
rapid allopatric speciation. *Nature*, 437(7063), 1353-1356.

Hoskin, C. J., and Higgie, M. (2013). Hybridization: its varied forms and consequences.
Journal of Evolutionary Biology, 26, 276-278.

Howard, D. J. (1986). A zone of overlap and hybridization between two ground cricket
species. *Evolution*, 40(1), 34-43.

Johanet, A., Secondi, J., Pays, O., Pagano, A., Lodé, T., and Lemaire, C. (2009). A case of
reproductive character displacement in female palmate newts (Lissotriton helveticus).
Comptes Rendus Biologies, 332(6), 548-557.

Johansson, F. (2003). Latitudinal shifts in body size of *Enallagma cyathigerum* (Odonata).
Journal of Biogeography, 30(1), 29-34.

Jombart, T. (2008). adegenet: a R package for the multivariate analysis of genetic markers.
Bioinformatics, 24(11), 1403-1405.

Kamimura, Y., and Mitsumoto, H. (2012). Lock□and□key structural isolation between
sibling Drosophila species. *Entomological Science*, 15(2), 197-201.

Kawano, K. (2002). Character displacement in giant rhinoceros beetles. *The American
Naturalist*, 159(3), 255-271.

Kawano, K. (2003). Character displacement in stag beetles (Coleoptera: Lucanidae). *Annals
of the Entomological Society of America*, 96(4), 503-511.

Kosuda, S., Sasakawa, K., and Ikeda, H. (2016). Directional mitochondrial introgression
and character displacement due to reproductive interference in two closely related
Pterostichus ground beetle species. *Journal of evolutionary biology*, 29(6), 1121-1130.

Kuntner, M., Coddington, J. A., and Schneider, J. M. (2009). Intersexual arms race? Genital
coevolution in nephilid spiders (Araneae, Nephilidae). *Evolution*, 63(6), 1451-1463.

Klingenberg, C. P. (2016). Size, shape, and form: concepts of allometry in geometric
morphometrics. Development genes and evolution, 226(3), 113-137.

Kyogoku, D., and Wheatcroft, D. (2020). Heterospecific mating interactions as an interface
between ecology and evolution. *Journal of Evolutionary Biology*, 33(10), 1330-1344.

Leong, J. M., and Hafernik Jr, J. E. (1992). Hybridization between two damselfly species
(Odonata: Coenagrionidae): Morphometric and genitalic differentiation. *Annals of the*
Entomological Society of America, 85(6), 662-670.

Liou, L. W., and Price, T. D. (1994). Speciation by reinforcement of premating isolation.
Evolution, 48(5), 1451-1459.

Lorenzo-Carballa, M. O., Watts, P. C., and Cordero-Rivera, A. (2014). Hybridization
between *Calopteryx splendens and C. haemorrhoidalis* confirmed by morphological and
genetic analyses. *International Journal of Odonatology*, 17(2-3), 149-160.

Macagno, A. L., Pizzo, A., Parzer, H. F., Palestrini, C., Rolando, A., and Moczek, A. P.
(2011). Shape-but not size-codivergence between male and female copulatory structures in
Onthophagus beetles. *PLoS One*, 6(12), e28893.

Mallet, J. (1986). Hybrid zones of Heliconius butterflies in Panama and the stability and
movement of warning colour clines. *Heredity*, 56(2), 191-202.

Mallet, J., Wynne, I. R., and Thomas, C. D. (2011). Hybridisation and climate change:
brown argus butterflies in Britain (Polyommatus subgenus Aricia). *Insect Conservation and*
Diversity, 4(3), 192-199.

Masly, J. P. (2012). 170 years of "lock-and-key": genital morphology and reproductive
isolation. *International Journal of Evolutionary Biology*, 2012.

McPeek, M. A., Shen, L., Torrey, J. Z., and Farid, H. (2008). The tempo and mode of three-
dimensional morphological evolution in male reproductive structures. *The American*
Naturalist, 171(5), E158-E178.

McPeek, M. A., Shen, L., and Farid, H. (2009). The correlated evolution of
three☐dimensional reproductive structures between male and female damselflies.
Evolution: International Journal of Organic Evolution, 63(1), 73-83.

McPeek, M. A., Symes, L. B., Zong, D. M., and McPeek, C. L. (2011). Species recognition
and patterns of population variation in the reproductive structures of a damselfly genus.
Evolution, 65(2), 419-428.

Melville, J. (2002). Competition and character displacement in two species of scincid
lizards. *Ecology letters*, 5(3), 386-393.

Mendelson, T. C. (2003). Sexual isolation evolves faster than hybrid inviability in a diverse
and sexually dimorphic genus of fish (Percidae: Etheostoma). *Evolution*, 57(2), 317-327.

Monetti, L., Sánchez-Guillén, R. A., and Rivera, A. C. (2002). Hybridization between
Ischnura graellsii (Vander Linder) and 2. elegans (Rambur) (Odonata: Coenagrionidae): are
they different species?. *Biological Journal of the Linnean Society*, 76(2), 225-235.

Muster, C., and Michalik, P. (2019). Cryptic diversity in ant☐mimic Micaria spiders
(Araneae, Gnaphosidae) and a tribute to early naturalists. *Zoologica Scripta*, 49(2), 197-
209.

Nava☐Bolaños, A., Córdoba☐Aguilar, A., and Munguía☐Steyer, R. (2012). A test of
genital allometry using two damselfly species does not produce hypoallometric patterns.
Ethology, 118(2), 203-213.

Nava-Bolaños, A., Sánchez-Guillén, R. A., Munguía-Steyer, R., and Córdoba-Aguilar, A.
(2014). Isometric patterns for male genital allometry in four damselfly species. *Acta*
Ethologica, 17(1), 47–52.

Nava-Bolaños, A., Sánchez-Guillén, R. A., Munguía-Steyer, R., and Córdoba-Aguilar, A.
(2017). Isolation barriers and genetic divergence in non-territorial Argia damselflies.
Biological Journal of the Linnean Society, 120(4), 804-817.

Nava☐Bolaños, A., Sánchez☐Guillén, R. A., Wellenreuther, M., Muñoz, J.,
Torres☐Pachón, M., Novelo☐Gutiérrez, R., and Córdoba☐Aguilar, A. (2019). Predicting
hybridisation as a consequence of climate change in damselflies. *Insect Conservation and*
Diversity, 12(5), 427-436.

Nishimura, T., Nagata, N., Terada, K., Xia, T., Kubota, K., Sota, T., and Takami, Y.
(2022). Reproductive character displacement in genital morphology in Ohomopterus
ground beetles. *The American Naturalist*, 199(3), E76-E90.

Nishimura, T., Terada, K., Xia, T. and Takami, Y. (2023). Relationships between
reproductive character displacement in genital morphology and the population-level cost of
interspecific mating: implications for the Templeton effect. *Biological Journal of the*
Linnean Society, 138(1), 14–26.

Noor, M. A. (1995). Speciation driven by natural selection in Drosophila. *Nature*,
375(6533), 674-675.

Nosil, P., Crespi, B. J., and Sandoval, C. P. (2003). Reproductive isolation driven by the
combined effects of ecological adaptation and reinforcement. *Proceedings of the Royal*
Society of London. Series B: Biological Sciences, 270(1527), 1911-1918.

Okuzaki, Y., Takami, Y., and Sota, T. (2010). Resource partitioning or reproductive isolation: the ecological role of body size differences among closely related species in sympatry. *Journal of Animal Ecology*, 79(2), 383-392.

Paulson, D. R. (1974). Reproductive isolation in damselflies. *Systematic Biology*, 23(1), 40-49.

Peterson, M. A., Honchak, B. M., Locke, S. E., Beeman, T. E., Mendoza, J., Green, J., ... and Monsen, K. J. (2005). Relative abundance and the species□specific reinforcement of male mating preference in the Chrysochus (Coleoptera: Chrysomelidae) hybrid zone. *Evolution*, 59(12), 2639-2655.

Pfennig, K. S., and Simovich, M. A. (2002). Differential selection to avoid hybridization in two toad species. *Evolution*, 56(9), 1840-1848.

Pfennig, K. S., and Pfennig, D. W. (2005). Character displacement as the "best of a bad situation": fitness trade□offs resulting from selection to minimize resource and mate competition. *Evolution*, 59(10), 2200-2208.

Pfennig, K. S., and Ryan, M. J. (2006). Reproductive character displacement generates reproductive isolation among conspecific populations: an artificial neural network study. *Proceedings of the Royal Society B: Biological Sciences*, 273(1592), 1361-1368.

Pfennig, K., and Pfennig, D. (2009). Character displacement: ecological and reproductive responses to a common evolutionary problem. *The Quarterly Review of Biology*, 84(3), 253-276.

Pilgrim, E. M., Roush, S. A., and Krane, D. E. (2002). Combining DNA sequences and morphology in systematics: testing the validity of the dragonfly species *Cordulegaster bilineata*. *Heredity*, 89(3), 184-190.

R Core Team. (2019). A language and environment for statistical computing. R Foundation for Statistical Computing. Vienna, Austria. Retrieved from http://www.R-project.org/

Robertson, H. M., and Paterson, H. E. (1982). Mate recognition and mechanical isolation in Enallagma damselflies (Odonata: Coenagrionidae). *Evolution*, 243-250.

Robinson, J. V., and Allgeyer, R. (1996). Covariation in life-history traits, demographics and behaviour in ischnuran damselflies: the evolution of monandry. *Biological Journal of the Linnean Society*, 58(1), 85-98.

Rohlf, F. J. (2006). TpsDig, program for Digitizing landmarks and outline for Geometric morphometric analyses, Version 2.04. Depto Ecol. and Evol., Stony Brook, New York.

Rohlf, F. J. (2008). TpsUtil, ver. 1.40. Department of Ecology and Evolution, State University of New York at Stony Brook, Stony Brook.

Roth-Monzón, A. J., Belk, M. C., Zúñiga-Vega, J. J., and Johnson, J. B. (2020). Beyond
pairwise interactions: Multispecies character displacement in Mexican freshwater fish
communities. *The American Naturalist*, 195(6), 983-996.

Sánchez-Guillén, R. A., Van Gossum, H., and Cordero Rivera, A. (2005). Hybridization
and the inheritance of female colour polymorphism in two ischnurid damselflies (Odonata:
Coenagrionidae). *Biological Journal of the Linnean Society*, 85(4), 471-481.

Sánchez-Guillén, R. A., Wellenreuther, M., Cordero-Rivera, A., and Hansson, B. (2011).
Introgression and rapid species turnover in sympatric damselflies. *BMC evolutionary
biology*, 11(1), 210.

Sánchez☐Guillén, R. A., Wellenreuther, M., and Cordero Rivera, A. (2012). Strong
asymmetry in the relative strengths of prezygotic and postzygotic barriers between two
damselfly sister species. *Evolution: International Journal of Organic Evolution*, 66(3), 690-
707.

Sánchez-Guillén, R. A., Córdoba-Aguilar, A., Cordero-Rivera, A., and Wellenreuther, M.
(2014a). Rapid evolution of prezygotic barriers in non-territorial damselflies. *Biological
Journal of the Linnean Society*, 113(2), 485-496.

Sánchez☐Guillén, R. A., Córdoba☐Aguilar, A., Cordero☐Rivera, A., and Wellenreuther,
M. (2014b). Genetic divergence predicts reproductive isolation in damselflies. *Journal of
evolutionary biology*, 27(1), 76-87.

Sánchez-Guillén, R. A., Sánchez-Herrera, M., and Vega-Sánchez, Y. M. (2023). Genetic
structure, cryptic species, and hybridization: causes and evolutionary consequences in
Odonata. In *Dragonflies and Damselflies. Second Edition. Edited by Alex Córdoba-
Aguilar, Christopher D. Beatty and Jason T. Bried, Oxford University Press.*

Sánchez-Guillén, R.A., Arce-Valdés, R., Swaegers, J., Chauhan, P., Chávez-Ríos, J. R.,
Rivas–Torres, A., … Hansson, B. (under review). Variable genomic patterns of
hybridization in an expanding hybrid region of damselflies. *Molecular Ecology*.
10.22541/au.168259144.47256890/v1.

Schlager, S. (2017). Morpho and Rvcg–Shape Analysis in R: R-Packages for geometric
morphometrics, shape analysis and surface manipulations. In Statistical shape and
deformation analysis (pp. 217-256). Academic Press.

Schluter, D. (2000). Ecological character displacement in adaptive radiation. *The American
Naturalist*, 156(S4), S4-S16.

Seddon, N., and Tobias, J. A. (2010). Character displacement from the receiver's
perspective: species and mate recognition despite convergent signals in suboscine birds.
Proceedings of the Royal Society B: Biological Sciences, 277(1693), 2475-2483.

Servedio, M. R. (2000). Reinforcement and the genetics of nonrandom mating. *Evolution*,
54(1), 21-29.

Servedio, M. R., and Noor, M. A. (2003). The role of reinforcement in speciation: theory
and data. *Annual Review of Ecology, Evolution, and Systematics*, 34(1), 339-364.

Simmons, L. W., and Garcia-Gonzalez, F. (2011). Experimental coevolution of male and
female genital morphology. *Nature communications*, 2(1), 374.

Simmons, L. W. (2014). Sexual selection and genital evolution. *Austral Entomology*, 53(1),
1-17.

Simmons, L. W., and Fitzpatrick, J. L. (2019). Female genitalia can evolve more rapidly
and divergently than male genitalia. *Nature communications*, 10(1), 1-8.

Slatkin, M. (1980). Ecological character displacement. *Ecology*, 61(1), 163-177.

Sloan, N. S., and Simmons, L. W. (2019). The evolution of female genitalia. *Journal of
Evolutionary Biology*, 32(9), 882-899.

Solano, E., Hardersen, S., Audisio, P., Amorosi, V., Senczuk, G., and Antonini, G. (2018).
Asymmetric hybridization in Cordulegaster (Odonata: Cordulegastridae): Secondary
postglacial contact and the possible role of mechanical constraints. *Ecology and Evolution*,
8(19), 9657-9671.

Song, H. (2009). Species-specificity of male genitalia is characterized by shape, size, and
complexity. Insect Systematics and Evolution, 40(2), 159-170.

Sota, T., and Kubota, K. (1998). Genital lock and key as a selective agent against
hybridization. *Evolution*, 52(5), 1507-1513.

Stre, G. P., Moum, T., Bureš, S., Král, M., Adamjan, M., and Moreno, J. (1997). A sexually
selected character displacement in flycatchers reinforces premating isolation. *Nature*,
387(6633), 589-592.

Taylor, E. B., Boughman, J. W., Groenenboom, M., Sniatynski, M., Schluter, D., and Gow,
J. L. (2006). Speciation in reverse: morphological and genetic evidence of the collapse of a
three spined stickleback (Gasterosteus aculeatus) species pair. *Molecular Ecology*, 15(2),
343-355.

Tregenza, T., Pritchard, V. L., and Butlin, R. K. (2000). The origins of premating
reproductive isolation: testing hypotheses in the grasshopper Chorthippus parallelus.
Evolution, 54(5), 1687-1698.

Tynkkynen, K., Grapputo, A., Kotiaho, J. S., Rantala, M. J., Väänänen, S., and Suhonen, J.
(2008). Hybridization in Calopteryx damselflies: the role of males. *Animal Behaviour*,
75(4), 1431-1439.

Usami, T., Yokoyama, J., Kubota, K., and Kawata, M. (2006). Genital lock-and-key system
and premating isolation by mate preference in carabid beetles (Carabus subgenus
Ohomopterus). *Biological Journal of the Linnean Society*, 87(1), 145-154.

Walker, T. J. (1974). Character displacement and acoustic insects. *American Zoologist*,
14(4), 1137-1150.

Wellenreuther, M., and Sánchez☐Guillén, R. A. (2016). Nonadaptive radiation in
damselflies. *Evolutionary Applications*, 9(1), 103-118.

Wellenreuther, M., Muñoz, J., Chávez☐Ríos, J. R., Hansson, B., Cordero☐Rivera, A., and
Sánchez☐Guillén, R. A. (2018). Molecular and ecological signatures of an expanding
hybrid zone. *Ecology and Evolution*, 8(10), 4793-4806.
993
994 Westman, K., Savolainen, R., and Julkunen, M. (2002). Replacement of the native crayfish
995 Astacus astacus by the introduced species *Pacifastacus leniusculus* in a small, enclosed
996 Finnish lake: a 30☐year study. *Echography*, 25(1), 53-73.
997
998 Yikweon, J. (2008). Asymmetry in reproductive character displacement. *Journal of*
999 *Ecology and Field Biology,* 31(4), 255–260.
1000

 Tables and Figure captions

1002 **Table 1**. Sample details of *I. elegans* and *I. graellsii* for the morphometric analysis. Species proportions indicate species proportion in
1003 each sampled population. N males and N females indicated the number of samples included in the morphometric analyses. The
1004 distribution indicates allopatric and sympatric populations: NW (North-west) and NCM (North-central and Mediterranean) hybrid
1005 zones.

Species	Locality	Distribution	Species proportion	Males	Females	Year	Latitude	Longitude
I. elegans	Marais D'Orx, France	Allopatry	100%	25	7	2015	43°35'58"N	1°23'30"W
I. elegans	Sablière, France	Allopatry	100%	26	8	2016	46°49'53"N	1°51'53"W
I. elegans	St. Cyprien, France	Allopatry	100%	26	25	2016	42°37'07"N	3°00'16"W
I. graellsii	Gamillazo, Salamanca	Allopatry	100%	14	20	2021	40°57'22"N	5°17'21"W
I. graellsii	Saïdia, Morocco	Allopatry	100%	-	16	2009	35°05'27"N	2°15'57"W
I. graellsii	Alba, Galicia	Allopatry	100%	16	9	2003	42°26'30"N	8°38'05"W
I. elegans	Doniños, Galicia	Sympatry (NW)	100%	24	23	2016	43°29'27"N	8°18'29"W
I. elegans	Laxe, Galicia	Sympatry (NW)	99-100%	27	22	2018	43°12'44"N	8°59'43"W
I. graellsii	Cachadas, Galicia	Sympatry (NW)	99-100%	31	28	2018-2019	42°27'12"N	8°50'52"W
I. graellsii	Xuño, Galicia	Sympatry (NW)	90-100%	30	13	2005, 2013, 2016	42°37'59"N	9°02'19"W
I. elegans	Perdiguero, Rioja	Sympatry (NCM)	71-90%	19	20	2018	42°17'06"N	1°58'55"W
I. elegans	Plaiaundi, Rioja	Sympatry (NCM)	99-100%	16	-	2016	43°20'44"N	1°47'40"W
I. elegans	Villar, Rioja	Sympatry (NCM)	74-100%	25	22	2018	42°24'03"N	2°42'15"W
I. graellsii	Valpierre, Rioja	Sympatry (NCM)	95-100%	30	22	2018	42°27'11"N	2°51'48"W
I. graellsii	Mateo, Rioja	Sympatry (NCM)	95-100%	21	33	2018	42°30'34"N	2°44'05"W

1006

1007 **Table 2.** Reproductive character displacement (RCD) in *I. elegans* and *I. graellsii* from both hybrid zones. First and second columns
1008 indicate whether analyses correspond to the shape or CS, sexes (male or females) and views (lateral or posterior). Third and fourth
1009 columns include interspecific and intraspecific comparisons. Fifth column indicate when genital variance was consistent with a pattern
1010 of RCD in one or both species. Sixth column indicates the selective force explaining the RCD pattern.

Dimension	View	Genital Divergence (Interspecific comparisons)			Genital Divergence (Intraspecific comparisons)		RCD		Selective force	
		Dist Symp	Dist Allop	Diff Symp-Allop	*I. elegans* Symp vs. Allop	*I. graellsii* Symp vs. Allop	*I. elegans*	*I. graellsii*	*I. elegans*	*I. graellsii*
Nort-west (NW) hybrid zone										
Shape	♂ Post view	0.28	0.31	-0.03	=	=	No	No		
	♀ Post view	0.19	0.12	**0.07**	=	=	No	No		
	♂ Lat view	0.19	0.22	-0.03	=	=	No	No		
	♀ Lat view	0.27	0.19	**0.08**	Symp≠Allop	Symp≠Allop	**Yes**	**Yes**	**Reinforcement**	
CS	♂ Post view	25.32	41.63	-16.31	=	=	No	No		
	♀ Post view	2.08	33.35	-31.27	=	=	No	No		
	♂ Lat view	31.45	1.23	**30.22**	=	Symp≠Allop	No	**Yes**		**Reinforcement**
	♀ Lat view	52.05	55.11	**3.06**	Symp≠Allop	Symp≠Allop	**Yes**	**Yes**	**Reinforcement**	**Lock-and-key**
North-central and Mediterranean (NCM) hybrid zone										
Shape	♂ Post view	0.26	0.31	-0.05	=	=	No	No		
	♀ Post view	0.22	0.12	**0.10**	Symp≠Allop	=	**Yes**	No	**Reinforcement**	
	♂ Lat view	0.18	0.22	-0.04	=	=	No	No		
	♀ Lat view	0.30	0.19	**0.11**	Symp≠Allop	=	**Yes**	No	**Reinforcement**	
CS	♂ Post view	59.35	41.63	**17.62**	Symp≠Allop	Symp≠Allop	**Yes**	**Yes**	**Lock-and-key**	**Reinforcement**
	♀ Post view	54.19	33.35	**20.84**	Symp≠Allop	=	**Yes**	No	**Reinforcement**	
	♂ Lat view	23.53	1.23	**22.3**	Symp≠Allop	Symp≠Allop	**Yes**	**Yes**	**Lock-and-key**	**Reinforcement**
	♀ Lat view	93.16	55.11	**38.05**	Symp≠Allop	Symp≠Allop	**Yes**	**Yes**	**Reinforcement**	**Lock-and-key**

1011

Table 3. Influence of environmental and geographical factors on the male and female reproductive characters of *Ischnura elegans* and *I. graellsii* damselflies from allopatric and hybrid zones. Estimated regression parameters, estándar errors, t-values and P-values for the Gaussian GLMs applied on the shape and the CS of reproductive structures.

Morphological character	Character view		Environmental/geographical factors	*I. elegans*				*I. graellsii*			
				Estimate	Std. Error	t value	P-value	Estimate	Std. Error	t value	P-value
Caudal appendage	Posterior	CS	Annual mean temperature	-28.018	7.780	-3.601	**<0.001**	NS	NS	NS	NS
		PC1	Annual mean temperature	-0.018	0.006	-2.958	**0.003**	NS	NS	NS	NS
		CS	Longitude	-8.715	3.568	-2.443	**0.015**	9.674	4.797	2.017	**0.045**
		PC1	Longitude	0.007	0.003	2.479	**0.014**	NS	NS	NS	NS
		CS	NW Zone	-94.936	31.718	-2.993	**0.003**	NS	NS	NS	NS
		PC1	NW Zone	0.081	0.025	3.297	**0.001**	0.039	0.011	3.623	**<0.001**
		PC1	NCM Zone	NS	NS	NS	NS	0.051	0.011	4.638	**<0.001**
		PC1	Allopatry Zone	-0.028	0.010	-2.903	**0.004**	NS	NS	NS	NS
Caudal appendage	Lateral	PC1	Annual mean temperature	-0.025	0.011	-2.317	**0.021**	NS	NS	NS	NS
		CS	Longitude	-11.331	3.216	-3.523	**<0.001**	18.650	8.255	2.259	**0.025**
		PC1	Longitude	0.011	0.005	2.141	**0.033**	NS	NS	NS	NS
		CS	NW Zone	-119.91	28.588	-4.194	**<0.001**	41.805	12.375	3.378	**<0.001**
		PC1	NW Zone	0.129	0.043	2.988	**0.003**	NS	NS	NS	NS
Prothorax	Posterior	CS	Latitude	NS	NS	NS	NS	3.388	1.58	2.144	**0.033**
		CS	Longitude	-13.664	5.216	-2.62	**0.009**	NS	NS	NS	NS
		PC1	Longitude	NS	NS	NS	NS	-0.003	0.001	-2.405	**0.017**
		CS	NW Zone	-149.918	45.423	-3.3	**0.001**	NS	NS	NS	NS
		PC1	NCM Zone	-0.058	0.019	-3.142	**0.002**	NS	NS	NS	NS
		PC1	Allopatry Zone	0.077	0.014	5.448	**<0.001**	NS	NS	NS	NS
Prothorax	Lateral	PC1	Latitude	NS	NS	NS	NS	0.009869	0.002746	3.593	**<0.001**
		CS	Longitude	-4.148	1.869	-2.219	**0.028**	NS	NS	NS	NS
		PC1	Longitude	NS	NS	NS	NS	0.005	0.002	2.263	**0.025**
		CS	NW Zone	NS	NS	NS	NS	10.454	3.944	2.651	**0.009**

		CS	NCM Zone	36.018	8.897	4.048	**<0.001**	13.116	3.771	3.478	**<0.001**
		PC1	NCM Zone	0.062	0.021	2.989	**0.003**	*NS*	*NS*	*NS*	*NS*
		PC1	Allopatry Zone	-0.136	0.016	-8.573	**<0.001**	*NS*	*NS*	*NS*	*NS*

1015

Figure Legends

Figure 1. Geographical distribution of *Ischnura elegans* and *I. graellsii* in allopatry and sympatry
(contact zone). Yellow dots show sampled localities in both hybrid zones (NW: northwest zone and
NCM: north-central and Mediterranean zone), blue dots indicate allopatric *I. graellsii* localities, and
purple dots indicate allopatric *I. elegans* localities.

Figure 2. PCA plots showing the variation of males' caudal appendage in both species (*Ischnura*
elegans and *I. graellsii*) in the allopatry regions (A, posterior view and D, lateral view), in the
northwest hybrid region (B, posterior view and E, lateral view) and the north-central region (C,
posterior view and F, lateral view), and of female's prothorax (G, posterior view and J, lateral
view), in the northwest hybrid region (H, posterior view and K, lateral view) and the north-central
region (I, posterior view and L, lateral view).

Figure 3. Media values of CS of the male caudal appendages of *I. elegans* and *I. graellsii* in A)
posterior and B) lateral views and female prothorax of *I. elegans* and *I. graellsii* in C) posterior and
B) lateral views from allopatry and both hybrid zones. Solid lines indicate statistically significant
differences at a significant level of $p < 0.05$ between conspecific males between zones.

Figure 4. Boxplot representing the morphological variation of CS among studied populations. A)
posterior view of male caudal appendage, B) lateral view of male caudal appendage, C) posterior
view of female prothorax, D) lateral view of female prothorax. Colors indicate species: purple= *I.*
elegans, blue= *I. graellsii*.

Zone