Pre-Cambrian origin of *envelope*-carrying endogenous retroviruses in metazoans

Shashank Chary[1] and Rippei Hayashi[1 #]

1. The Shine-Dalgarno Centre for RNA Innovation, The John Curtin School of Medical Research, The
Australian National University, 131 Garran Road, Acton, ACT 2601, Australia

correspondence: rippei.hayashi@anu.edu.au.

Abstract

Retrotransposons or endogenous retroviruses (ERVs) essentially carry open reading frames of *gag* and *pol*,
which are utilized to selfishly replicate themselves in the host's germline genome. One rare example of
ERVs that additionally carry *envelope* genes, are Ty3/*gypsy* errantiviruses in Drosophila. Though they are
structurally analogous to vertebrate retroviruses, their evolutionary relationship as well as the origin of
envelope genes remain elusive. We systematically searched for intact *envelope*-containing ERVs that are
homologous to Ty3/*gypsy* and vertebrate retroviruses in metazoan genomes and found that they are
widespread across taxa including ancient animals such as Cnidarians, Ctenophores and Tunicates. Most
elements are multiplied in their respective genomes, implying recent germline expansion. *Envelope* genes
are classified into those that resemble glycoprotein F from paramyxoviruses (F-type), glycoprotein B from
herpesviruses (HSV/gB-type), and GP2 glycoprotein from retroviruses and Ebolavirus (GP2-type). Both F-
type and HSV/gB-type ERVs are ubiquitously found in invertebrate genomes while vertebrate genomes
carry F-type and GP2-type ERVs. Interestingly, vertebrate ERVs share a distinct and structurally conserved
domain of POL with Ty3/*gypsy* elements from insects and ancient animals, suggesting that they share
common origins. Furthermore, phylogenetic and structural analyses revealed that *envelope* genes have
largely diverged with *pol* genes as well as with the host organisms throughout their evolutionary history
and recombined infrequently, suggesting that the *envelope* acquisition to ERVs is ancient and likely dates
to before the split of bilaterian and non-bilaterian animals in Pre-Cambrian era.

Introduction

Retroviruses have two mechanisms through which they survive; most commonly, they exogenously infect other hosts, transmitting horizontally [1]. However, they can occasionally infect the germline cells of the host; where genetic information is stored for the next generation and integrate into the genome [1,2]. They become fixed in the genome, vertically transferring to the next generation and become known as endogenous retroviruses (ERVs) [1-4]. They are well characterised in vertebrates, making up 5-10% of their genomes. The best-known example of this are Koala retroviruses (KoRVs), in which nearly identical copies of an exogenous lymphoma-causing retrovirus are found in the genomes of geographically distinct Koala populations [5,6]. Though, it should be noted that such germline infection events have rarely been identified, and the exact exogenous origins of many ERVs is not known.

Once a virus becomes endogenized, they have two possible fates. Most commonly, proviral copies in the genome acquire deletions and frameshifts, degrading over time, until only fragments remain [7]. Some retroviral genes are even co-opted by the host to assist in biological functions [8]. It is for this reason that ERVs tend to be thought of as ancient relics of past retroviral infection [9,10]. However, some ERVs retain functionality of their retroviral proteins, and can endogenously create new copies in the genome, keeping their code intact. This requires the activity of the retroviral genes *gag* and *pol*, which are usually flanked by long terminal repeats (LTRs). These active intact ERVs are more broadly classified as class I LTR retrotransposons [11].

Retrotransposons usually exist in the genome as 'selfish' genes, endogenously creating new copies in the germline to survive long term [2,12]. Sometimes, they can be lost from the genome, if they do not remain active enough, or if they acquire random mutations that affects their ability to transpose. The study of intact retrotransposons therefore provides an interesting opportunity, as they can provide great insights into the evolution of retroviruses. It has been postulated that intact ERVs and retrotransposons are the result of more recent retroviral infection [13].

The Ty3/*gypsy* family of retrotransposons is one of the largest and best studied families of LTR retrotransposons, present ubiquitously in the genomes of most eukaryotes. Their origin, likely to be highly ancient [14], has been suggested to be linked to that of vertebrate retroviruses as well. In fact, the *gag* and *pol* genes from vertebrate retroviruses are most closely linked to those of Ty3 elements [15-17]. A recent study identified a new class of ERV in vertebrates called Lokiviruses [18], which has been suggested to be a link between the two, further supporting the idea that the *gag* and *pol* genes of Ty3 retrotransposons and vertebrate retroviruses have a common origin.

One specific branch of Ty3 elements in Drosophila species carry an open reading frame (ORF) for the retroviral *envelope* or *env* gene in addition to *gag* and *pol*, often referred to as 'errantiviruses' [19,20]. Genetic evidence in Drosophila shows that these errantiviruses behave as bona-fide retrotransposons, creating new germline copies that are vertically inherited [21,22]. Interestingly, errantiviruses are expressed in the somatic cells of the female ovaries, forming virus like particles (VLPs) that infect the adjacent germline cells [21,22].

Essentially, errantiviruses can actively infect other cells, rather than maintain copies in the same cell as
typical retrotransposons. This means that if the errantivirus remains active in the somatic cells, it can still
maintain the generation of new copies that can be vertically transmitted to the next generation,
endogenously via cell-to-cell transmission.

Errantiviruses are an interesting branch of retrotransposons, as they are the most 'intact' elements
compared to retroviruses, given the presence of *env*. But although they share structural similarities, the
exact origins are poorly understood. It has been suggested that Ty3 elements acquired the *env* gene from
infecting baculoviruses in an ancestral Drosophila species via recombination events [23,24]. Errantiviruses are
only known to be found in dipteran genomes, identified recently in mosquito species [25]. However,
Baculoviruses are also known to infect non-dipteran insect orders, most famously Lepidoptera species [26].
Additionally, the origins for the *env* gene in baculoviruses is also poorly understood, leading to ambiguity
in the true origins of these elements. Furthermore, LTR retrotransposons with *env* genes have been
sporadically documented in other animals, such as in nematodes and rotifers [27,28]. However, their origins
are even less clear, due to a lack of systematic studies to mine genomes for such elements.

Errantiviruses may therefore be more widespread than previously thought, and the search for intact
errantivirus elements in metazoan orders allows us a unique opportunity to study the acquisition of
retroviral *env* genes by retrotransposons over time.

Results

We systematically searched for homologous elements of Dipteran Ty3/*gypsy* errantiviruses in the
referenced genomes across metazoan phyla using the annotated protein coding sequences of *gag*, *pol* and
env for tBLASTn searches (see methods and Fig S1). We additionally used the protein coding sequences of
known vertebrate retroviruses to search for homologous retroelements in vertebrate genomes. We took
genomic copies that carry all three open reading frames (ORFs). Where possible, we focused our search on
elements that are multiplied in the genome to understand relationships of currently active elements. The
first tBLASTn search identified elements outside Diptera and Vertebrata. Therefore, we extended the
search using the representative of newly identified elements as baits. We repeated this process and
examined whether the additional baits uncovered elements that were not represented in the previous
tBLASTn searches by identifying new clades in the phylogenetic tree of POL (see methods and Fig S1).
After a five iteration of tBLASTn searches, we no longer discovered elements that formed a new POL
phylogenetic clade (Fig S1B). These steps allowed us to obtain a comprehensive set of elements in
metazoan genomes that are homologous to Dipteran Ty3/*gypsy* errantiviruses and vertebrate retroviruses
(Table S1, summarised in Fig 1). Most of the newly identified genomic copies carry putative LTRs and
tRNA primer binding sequences (1308 out of 1765 elements, Table S1), suggesting their recent
integrations. Additionally, most of the annotated elements sit within contigs larger than 0.1 mega bases
(1649 elements, Table S1), indicating their true origins in their respective genomes. We ran multiple
sequence alignments using the conserved Integrase (INT) and Reverse Transcriptase (RT) domains of Pol
and assembled maximum probability trees to infer phylogenetic relationships of the identified elements.

ORFs of selected annotated Drosophila Ty3 elements and vertebrate retroviruses were also included in the

trees to aid classifications. We refer to elements found in non-vertebrate metazoan phyla as

"errantiviruses", and elements found in vertebrate genomes as "vERVs".

<u>Multiple intact elements of *env*-carrying Ty3 retrotransposons are found widespread across metazoan</u>

<u>species</u>

We found that intact errantiviruses are found not only outside of insect orders but are ubiquitously present

across most major metazoan phyla. This includes species from ancient phyla such as Cnidaria, Tunicates

and Ctenophora, but excludes Echinodermata and Porifera (Fig 1B and Table S1). This range of hosts,

which includes ancient phyla, suggests that the elements identified in this study are highly ancient. This is

also supported by the phylogenetic tree assembled on *pol* RT and INT proteins. Firstly, the phylogenetic

tree assembled on *pol* RT reflects that of *pol* INT, with errantiviruses within most clades in one tree

mapping to the same or closely related clades in the other (Fig S2B). Importantly, errantiviruses found in

phyla outside of insects, cluster separately to those found within, which we call "ancient errantiviruses"

and "insect errantiviruses", respectively (Fig 1C and S2A). Notably however, few elements found in

insects sit within clades of ancient errantiviruses, rather than other insect ones (Fig 1C). Annotated

Drosophila *gypsy* elements included in the tree, namely, Gypsy-ZAM, Gypsy-Gypsy/Gypsy-5_DGri and

Gypsy-DM176/Idefix present themselves in different clades while they cluster together with other insect

errantiviruses, in an arrangement with previous studies that have classified Drosophila *gypsy* elements [29]

(Fig 1C). Though, one element from a cnidarian species *Paleozoanthus reticulatus* clusters together with

insect elements, in a clade adjacent to DM176 and Idefix (Fig 1A), its origin is unclear as the element is

found on a short genomic contig (Table S1). While we have potentially identified new classifications of

insect errantiviruses, given new clades (tree positions I3 - I6 in Fig S2A) that are formed between known

insect *gypsy* and "ancient errantiviruses", the separation between "insect errantiviruses" and "ancient

errantiviruses" is still highly indicative of their ancient age.

Within "insect errantiviruses", the clades are strongly supported by bootstrap values greater than 95 and

are additionally supported by the annotations of the tRNA primer binding site (PBS) found in each

element. For example, Tree position I1 within "insect errantiviruses" mainly carry tRNA^{Ser}, whereas Tree

position I2, consisting of only dipteran elements, predominantly use tRNA^{Lys}, though few use tRNA^{Ser} or

tRNA^{Pro} (Fig S2A and S2C). Elements in insects within positions I4, I5 and I6 that predominantly carry

tRNA^{Leu} primer binding sites, in addition to Serine and other uncommon tRNAs (Fig S2C). This is mostly

consistent with past works, that have identified Drosophila errantiviruses to carry either Serine or Lysine

tRNA PBS [30,31].

The "ancient errantivirus" group also consists of clades strongly supported by bootstrap values greater than

95, though the tRNA PBS content within the clades is much more divergent (Fig S3C). Notably however,

errantiviruses found in insects that sit within this group, in tree positions A1, A2 and A7 continue to carry

predominantly tRNA^{Ser}, tRNA^{Lys} and tRNA^{Leu} (Fig S3B). The clades that are formed in this group also

reflect their ancient nature, as they consistently are made up of elements in hosts from the same, or closely

related phyla (Fig S3B). For example, tree position A5 consists of elements found in Thecostraca, Annelida and Mollusca, each of which form their own discrete clade (Fig S3B). Even within larger clades, the separation is evident. For example, tree position A9 consists of elements in a wide variety of hosts, but strongly supported clades only consist of elements from the same order of host (Fig S3B). This congruence between clades and species is highly indicative of long-term vertical transfer, as known in non *env*-containing retrotransposons [32,33].

Most elements that were identified have multiple intact copies in their respective genomes (as defined as copies with >98% nucleotide identity across >98% of the three ORFs, see methods, Fig 1C, Table S1), a hallmark feature of recent transposition [34,35], indicating that most elements that we have annotated likely behave as bona-fide retrotransposons. Activity of these elements are highly varied, as the copy number can be as little as one intact copy, to hundreds of intact copies in the genome (Fig 1C, Table S1). Importantly, multi-copy ERVs are present in nearly all animal phyla where we see intact elements, suggesting that these elements are actively transposing in a wide range of metazoan species today.

The *env* ORF downstream of *pol* in many of the elements resembles glycoprotein F1, found more commonly in paramyxoviruses, such as Nipah, Hendra and Respiratory Syncytial Viruses [36,37], and is also carried in insect baculoviruses [23,24]. However, some errantiviruses instead carry *env* that resembles glycoprotein B (gB), famously found as part of the glycoprotein complex in Herpes Simplex Viruses (HSV). Both types of *env* glycoproteins are carried by errantiviruses found in ancient hosts such as rotifers, tunicates, cnidarians and ctenophores, as well as in hosts from insect and arthropod orders (Fig 1B and S3B). Though, errantiviruses found in Nematoda, Bryozoa and Platyhelminthes only carry HSV/gB-type *env*, and the ORFs are arranged as GAG-ENV-POL rather than GAG-POL-ENV. Furthermore, clades formed on the trees using *pol* proteins tend to carry the same type of *env* (Fig S2A and S3A). For example, within A6 in the "ancient errantivirus" group, elements that carry HSV/gB are in a separate clade to those carrying F (Fig S3B). Though some clades appear to be a mix of both, the HSV/gB elements still tend to cluster together, indicating that they have stayed with *pol* for long evolutionary periods of time.

<u>The divergence of *env* glycoproteins carried by errantiviruses follows that of *pol*</u>

We found that errantiviruses can carry HSV/gB-type *env* in addition to glycoprotein F, in both "insect errantiviruses" and "ancient errantiviruses" (Fig 2A). One clade that exclusively consists of hymenopteran errantiviruses carry both HSV/gB-type and F-type *env* genes (Fig 2A). Interestingly, elements with HSV/gB-type *env* and F-type *env* separate at the base of this clade while they both are found in a wide range of species superfamily within Hymenoptera, suggesting that both HSV/gB-type *env* and F-type *env* have been together with their *pol* genes at least throughout the evolution of hymenopteran species.

To further characterise the ENV proteins carried by errantiviruses, we identified the ectodomains within the *env* ORF, which is flanked by signal peptides and a transmembrane domain, using DeepTMHMM (https://doi.org/10.1101/2022.04.08.487609). We then analysed trimeric configurations of these ectodomains from representative errantiviruses from the phylogenetic tree using Alphafold 2 [38].

F glycoproteins analysed are highly structured, and consistently form 4 domains that are highly conserved

across all elements (Fig S4A). They also predicted a highly conserved Furin cleavage site and hydrophobic

fusion peptides in the ectodomain (Fig S4A and S5). The 3D fold of the ectodomain structures forms a

"head" (domains 1-3) and a "stalk" (domain 4), strongly resembling 3D structures of F ectodomains

characterised in previous studies, such as those from RSVs [36] (Fig S4D). Most errantiviruses also have an

alpha helix at the C terminus of the ectodomain, referred to as "Domain A" (Fig S4A and S4D). The

predicted 3D trimeric structure also reveals two potential configurations of the glycoprotein, based on

domain 4: either a long alpha helix, or two shorter helices (Fig S4D). These long and short barrel structures

resemble pre- and post-fusion structures of glycoprotein F described in prior studies [36]. These highly

structured domains in the intact *env* ORFs in these errantiviruses demonstrates that *env* has remained intact

and could suggest that their function also remains intact as well.

To explore the phylogenetic relationships between *pol* and *env*, we generated multiple sequence alignments

of the ectodomains and RT domains from representative elements and assembled phylogenetic trees. We

found that the ENV ectodomains cluster similarly to RT in *pol*, and ENV proteins tend to form clades

within the same or closely related species as they do for RT (Fig S4B and S4C). Importantly, the ENV

ectodomains found in "ancient errantiviruses" cluster separately to those found in "insect errantiviruses".

Additionally, insect elements that present themselves with ancient errantiviruses in the RT tree, are

similarly positioned in clades within the ENV tree (Fig S4C). This suggests that *env* genes have stayed

with *pol*, and the acquisition of *env* likely occurred anciently.

The ectodomain of errantiviruses that carry HSV/gB-type *env* are either flanked by signal peptides and a

transmembrane domain, or by two transmembrane domains and the Furin cleavage site is not always

conserved (Fig 2B). Like F, the trimeric structures of HSV/gB found in errantiviruses are also highly

structured. They carry 5 highly conserved and distinct domains (designated as domain I – V) arranged in a

hairpin configuration, that fold into a trimeric structure with a base, stalk and crown (Fig 2B). Additionally,

domain I carries highly conserved "fusion loops" consisting of hydrophobic amino acid residues. These

domains have also been characterized in Herpes Simplex Viruses in prior studies [39,40]. Notably,

errantiviruses found in insect orders that carry HSV/gB, only carry one of these fusion loops (Fig S6).

The structure of the HSV/gB carried by these errantiviruses are stabilized by disulphide bonds between

Cysteine residues, which we refer to as Cysteine bridges [41,42]. These bridges are highly conserved but

appear to vary the most within domains I and II of the HSV/gB ectodomain (Fig 2C and S6). Additionally,

in some cases, a Cysteine bridge within domain I is located near the fusion loops (Fig 2C and S6). The

configuration of Cysteine bridges appears to be specific to phyla, though this does not seem to be reflected

by their positions in the tree based on *pol* RT. For example, Bryozoa elements in tree position A3, and

Bryozoa elements in tree position A8, both carry configuration C1c, though they each cluster themselves

separately from other elements (Fig S3B, 2D and S6). Errantiviruses carrying HSV/gB found in insect

phyla mostly have the same configuration regardless of their position in the *pol* RT tree. For example,

insect elements within the "ancient errantivirus" group, and insect elements within the "insect errantivirus"

group, all have the same Cysteine bridge configuration in HSV/gB (Fig 2A, 2D and S6). There are,
however, cases where more than one type of configuration can be found within the same phyla, namely
Nematoda and Platyhelminthes (Fig 2D). Though, in those cases, they appear to continue to separate based
on *pol* RT (Fig S6). For example, Nematoda elements in Position A2 have configurations C1a or C1d,
while those in Position A8 have configuration C2a or C2c (Fig S6).

As mentioned above, the clades that form on RT mostly separate elements that carry HSV/gB from F
within discrete clades (Fig S3B). This, with the fact that the same substructures can be found in different
orders, for example in Nematoda, Bryozoa and Platyhelminthes, suggests that their origin is likely ancient.
The overall specificity of certain residues to host orders further suggests that recombination events would
have only occurred between elements within the same animal order. Furthermore, the overall separation of
distinct Cysteine bridge configurations of HSV/gB, that are represented in the *pol* RT tree within ancient
phyla, suggests that those recombination events occurred anciently. While insect elements are separated on
the tree, ERVs with HSV/gB-type ENV tend to cluster separately from those with F-type ENV. For
example, in the "insect errantivirus" group, the clade that almost exclusively carries HSV/gB-type ENV
sits sister to another clade that only carries glycoprotein F like ENV (Fig 2A'). Not only do they separate
from each other, but HSV/gB-type ENV can be found in a variety of superfamilies within Hymenoptera,
which suggests their ancient origin (Fig 2A). Some species, such as a hymenopteran *Lasioglossum*
lativentre, carry elements with HSV/gB, as well as those with F (Table S1). Despite being found in the
genome of the same species, the *pol* of errantiviruses with F and HSV/gB do not stay together, further
supporting their more ancient origin (Fig 2A'). It should be noted that some clades, such as ones in tree
position 1 within "ancient errantiviruses", have elements that carry HSV/gB and F sitting together within
the same discrete clade, suggestive of more recent recombination (Fig S3B).

<u>The RNase H Domains within the bridge region of *pol* separates "insect errantiviruses" from "ancient</u>
<u>errantiviruses"</u>

To further characterize the errantiviruses identified in this study, we analysed the region between the INT
and RT domains of Pol, referred to as the "bridge region", for representative elements. This region is
known to carry a ribonuclease H (RNase H) domain, which in vertebrate retroviruses has duplicated, and
the N-terminal copy, while partially degraded, maintains the same structure and 3D fold arrangement,
referred to as "tether" [43]. We therefore characterised structures within the bridge region using the 3D fold
arrangement of the core β-sheet structure using Alphafold 2.

We found that almost all errantiviruses that sit within the "insect errantivirus" group have similarly
duplicated the RNase H domain, with the C terminal copy presenting itself as a "pseudo" RNase H domain
(Fig 3A-B). The "pseudo" RNase H domain retains the barrel-like structure made of five core β-sheets, but
the order of the second and the third β-sheets is flipped (Fig 3B).

On the other hand, "ancient errantiviruses" carry an RNase H domain, and a C-terminal "mini" domain.
This consists of three small core β-sheets and α-helix residues at its N terminus. This "mini" domain is also

carried by insect elements in this group, further suggesting that those insect elements are equally ancient as

the other errantiviruses in this group. Furthermore, this "mini" domain is also found in the bridge region of

yeast S. *pombe* Ty3 *pol* (Fig 3), indicating it as a feature of highly ancient transposons.

We also identified a few uncommon structures, that are found in some phylogenetic groups in the tree, all

of which sit at the C terminus of the bridge region (Fig 3B). One notable outlier is an insect element within

the "ancient errantivirus" group that carries a variation of a "pseudo" RNase H domain (position 6, noted

as "pseudo" domain 2 in Fig 3). This same domain can be found in a phylogenetic group in the "insect

errantivirus" group (position 3 in Fig 3A). Furthermore, some elements in this "insect errantivirus" group

carry a "mini" domain as well (position 3 in Fig 3A). Another is that some elements within the "insect

errantivirus" group only carry a solitary RNase H domain (positions 4 and 5 in Fig 3A). It should be noted

that these outliers are mostly specific to small phylogenetic groups/clades, and likely also arose somewhat

anciently.

<u>Vertebrate genomes carry intact copies of *env*-carrying ERVs</u>

After finding that *env*-carrying retrotransposons are widespread in invertebrate metazoan genomes, and

that ancient errantiviruses present themselves as distinct to insect errantiviruses, we extended the search for

intact retroelements to vertebrate genomes. We identified intact elements with the same approach as

errantiviruses, and assembled a phylogenetic tree using the sequences of *pol* RT and INT. We also verified

that the assembled tree on RT reflects the tree of INT; elements in one clade are found in the same or

similar clades on the other tree (Fig S7A).We included the sequences of Epsilon, Gamma, Foamy, Delta,

Lenti, Alpha, Beta and Loki viruses, as well as the sequences of SNRV and ERV-L to aid the classification

of the genomic copies that we identify.

Like errantiviruses, every class within vertebrates carry intact copies of *env*-containing retroviruses

endogenously within their genomes, referred to herein as "vERVs" (Fig 4). We classified the vERVs based

on where annotated vertebrate retroviruses sit and found that each retrovirus strongly forms its own clade

with corresponding vERVs, supported by bootstrap values greater than 95 (Fig 4A). The exceptions are

Foamy, Delta and Lentiviruses, which do not form clades with vERVs, and rather form solitary clades on

their own. Regardless, the classification and phylogeny of each vertebrate retroviral clade agrees with

previous analysis of vertebrate retroviruses [18,44] (Fig 4B). In addition to existing retrovirus classification,

we identify new clades of ERVs that sit basal to known retroviral clades (labelled as vERV 1-4) (Fig 4A).

Furthermore, many vERVs that sit in the same large clade as gamma retroviruses are highly widespread

and divergent, forming their own clades without known exogenous virus relatives, namely, the clades in

vertebrate subgroups V5, V6, V7 and V8 (Fig 4A and S7D). Importantly, most vERV clades are found in a

wide range of hosts, though they mostly cluster according to the host phylogeny. For example, gamma

vERVs in subgroups V4 to V7, and V8 exclusively consist of mammalian and fish elements, respectively

(Fig S7D). When multiple vertebrate classes are represented in the same subgroups of gamma vERVs,

elements from evolutionarily related hosts tend to cluster together. For example, subgroup V6 consists of

bird and reptile vERVs, while Amphibian and retile elements cluster together in subgroups V5 and V7 (Fig

S7D). Like errantiviruses, this is indicative of long-term vertical transfer. Furthermore, many of the
elements identified carry intact LTRs, as well as tRNA PBS, further supporting the intactness of the
elements. Though the tRNAs are highly varied and divergent, like those found in "ancient errantiviruses"
(Fig S7E). Additionally, many of these elements have multiple intact copies in their respective genome
suggesting recent transpositions (Fig 4D).

We further characterised the vERVs that we identified in this study, by analysing the bridge region of *pol*.
As previously mentioned, vertebrate retroviruses have duplicated the RNase H domain, and the N-terminal
copy, while partially degraded, maintains the same structure and 3D fold arrangement, referred to as
"tether" [43]. Our analysis similarly finds that all vERVs carry both RNase H and tether, however, many of
the elements additionally carry the "mini" domain identified in "ancient errantiviruses" and yeast Ty3
elements (summarised in Fig 4A-B, and an example of vERV "mini" RNase H domains shown in Fig 4F).
This strongly supports the idea that many of these vERVs are highly ancient, and potentially share a
common origin with Ty3 elements. One area of note, however, is that some vERVs in gamma retroviruses
that do not carry the "mini" domain and form their own discrete clade with KoRV-A, potentially indicating
that they have emerged more recently (Fig S7D).

We finally characterised the *env* ORF carried by these vERVs. Most carry Glycoprotein 1 (GP1) and
Glycoprotein 2 (GP2), resembling *env* structures from other retroviruses as well as Ebolaviruses [45-47]. All
elements analysed carry intact GP2, however we found structural variations within the GP1 domain in
vERVs, which we analysed using Alphafold 2. Mainly, we find that GP1 from vERVs either carry a
receptor binding domain (RBD) and a C domain, or they lack the RBD entirely (Fig 4A-C). All have a
highly conserved Furin cleavage site upstream of GP2, and most carry two Cysteine residues upstream of
the C domain (Fig 4C and S8). Additionally, we found other highly conserved Cysteine residues that likely
form Cysteine bridges which are shared between groups of elements (Fig S8). Interestingly, some of the
vERVs instead carry GP1 that resembles the structure of Syncytin-2 (Fig S7C-D). Notably, Syncytin-2 like
GP1 domains are not restricted to a single order, being found in reptiles and mammals (Fig S7D). These
highly consistent structures suggest that *env* in vERVs that we identified has not undergone degradation,
and its function has remained intact as well. The presence or absence of RBD in these elements is mostly
consistent with the formations of the clades, though some large clades are mixed in representation (Fig 4A-
B and S7D). There is some ambiguity in these mixed clades, as it may suggest that the RBD has been lost
frequently in different clades over time. However, even within those mixed clades, elements with only the
C domain and elements with both RBD and C domains tend to form their own distinct clades. For
example, within vertebrate group V5, elements with RBD and C form their own clade separate to elements
with C only, suggesting their relatively ancient origin (Fig S7D). Vertebrate group V8 shows a greater
degree of mixing suggestive of more recent recombination, though the outliers have larger branch lengths,
and are not supported by bootstrap values greater than 95, indicating greater divergence within *pol* (Fig
S7D).

Additionally, Cysteine bridges that are formed within GP1 are conserved between elements from the same
vertebrate classes. For example, of the three distinct Cysteine bridges within the C domain of GP1
structures that only have the C domain, one of them (marked as "1" in Fig S8) are predominantly present
in Amphibian and fish vERVs while no bird or reptile elements carry it. Similarly, one Cysteine bridge in
the C domain within the RBD + C group (marked as "4" in Fig S8) is mainly carried by mammalian, bird
and reptile vERVs while no fish or Amphibian vERVs carry it. These observations further suggest that the
env genes in vERVs have stayed with *pol*, and that recombination events only occurred between closely
related species or occurred anciently.

Lokivirus and SNRV vERVs do not carry GP1 and GP2 but carry *env* that resembles glycoprotein F
instead. They are found in more ancient vertebrate classes, including reptiles, amphibians, fish and jawless
fish, sit distantly in a sister clade to ERV-L, and carry RNase H, tether and a "mini" domain in their bridge
region, highly suggestive of an ancient origin (Fig 4A-B). The ectodomain of the F glycoprotein carried by
these vERVs retains the highly structured domains found in errantivirus F glycoproteins, as well as the
highly conserved Furin cleavage site (Fig 4F and S5), suggesting that its function is intact.

We assembled a *pol* RT tree to include both errantiviruses and vERVs, and find that vERVs, "ancient
errantiviruses" and "insect errantiviruses" are clearly distinct from one another. However, Lokivirus
vERVs sit as a sister clade to ancient errantiviruses (Fig 4E-F). The commonality in features, such as the
classification of *env* and the "mini" domain present in the bridge region of *pol*, is highly suggestive of a
common origin between the two. Furthermore, phylogenetic analysis of the *env* F ectodomain clusters
vertebrate *env* F with that of ancient errantiviruses, specifically that of tunicate *env* F (Fig S4C), strongly
suggesting that the origin of *env* in these retroelements is highly ancient, and potentially shared between
ancient animals and vertebrates.

Discussion

Endogenous Ty3 retroelements carrying *env* genes were only previously identified in Dipteran genomes.
However, we found that genomes from nearly every major order carry such elements, and they are fully
intact with uninterrupted ORFs and LTRs. Additionally, many of the elements annotated have multiple
intact copies in their respective genomes, indicating that they have been transposing recently. The
intactness of these elements is further supported by structural analysis of the *env* glycoproteins, where we
found that highly conserved structural domains are still retained, despite divergences in the primary
sequence.

We found that these errantiviruses are likely to be highly ancient, being found in ancient animals such as
Cnidaria, Ctenophora and Tunicata. Phylogenetically, they are found within an "ancient errantivirus" group
that sits separately to elements found in insects. These ancient elements are also structurally distinct, with
the bridge region between INT and RT of *pol* resembling that of yeast Ty3 retrotransposons, rather than
that of insects. Furthermore, the divergence of *env* genes appear to follow that of *pol*, with few exceptions,
indicating that the *env* genes are similarly ancient, and that recombination events are likely rarely

occurring. Clades on the *pol* tree also tend to consist of elements from the same, or closely related hosts,
rather than consist of elements within highly divergent host species, further supporting their ancient nature,
and indicating that they have mainly been vertically transmitting over time.

Finally, we found that *env*-containing endogenous retroviruses are similarly found in vertebrate genomes,
and the same observations made in errantiviruses mostly apply to these elements as well. The recently
identified vertebrate Lokiviruses sit as a sister clade to "ancient errantiviruses" and shares structural
features with them, suggesting their common origins.

It was originally speculated that errantiviruses acquired *env* from baculoviruses in insects, some of which
carry an F-type ENV protein Ld130, potentially via recombination events [23,24]. While Ty3 elements may
have obtained the *env* gene via recombination, it is unlikely that baculoviruses are the source for the *env*
gene found in errantiviruses, as they are far more widespread than insects. The congruence of phylogeny
between *env* and *pol* indicate that they have likely co-evolved (Fig S4). This, alongside the presence of
errantiviruses in ancient host orders such as Cnidaria, Ctenophora and Tunicata, suggests that *env*-carrying
Ty3 elements are highly ancient transposons, rather than the result of a recent recombination event with an
exogenous virus. HSV/gB-type *env* in errantiviruses appears to be equally as ancient, as it is similarly
found in ancient animal orders (Fig 1). The arrangement of Cysteine bridges that stabilize the structure of
the ectodomain, are consistent between elements within the same order, even though *pol* presents itself
separately on the tree (Fig 2 and S6). This implies that the *env* has diverged with the host, but likely
undergoes recombination events with other elements within the same species. Though, *pol* with HSV/gB-
type *env* mostly cluster within their own clades, and are not mixed with *pol* carrying glycoprotein F,
indicating that recombination events are not happening frequently enough to randomise their spread, and
likely occurred anciently. The origins and acquisition, therefore, of either type of *env* protein, likely
occurred in the last common ancestor of metazoans.

Finding intact, multicopy elements in such a widespread manner brings into question the origins of these
unique retrotransposons. One explanation could be that errantiviruses frequently undergo horizontal
transfer, with shorter periods of vertical transmission. Drosophila errantiviruses have been previously
suggested to horizontally transfer between species [29,48]. It has also been suggested that some LTR
retrotransposons in bivalves can horizontally transfer between different species and can even jump across
phyla as well [49]. While horizontal transfer of errantiviruses between distantly related species cannot be
excluded, it is likely to be a rare occurrence, given that *pol* from the same order tend to form their own
discrete clades (Fig S3 and S7). In instances where clades are formed from different phyla of hosts, they
are magnitudes of order apart from each other, and it is unlikely that they can easily transfer. Additionally,
many of these clades tend to have much longer branches, and it is therefore possible that they are still
distantly related and diverged a long time ago. For example, within tree position A10 (Fig S3), a clade
consisting of elements from Hexanauplia and Annelida are separated by longer branch lengths. The reason
for these types of clades, and as to why we do not see elements in species between those distantly related
orders, is likely due to the frequent loss of these retrotransposons from the genome. In such cases, distantly

related species have seen such retrotransposon copies being active, which are vertically inherited, and

diverge along with the genome of the host species, while other species do not see this activity, and those

inactive copies would be degraded over time.

In Drosophila, errantiviruses occupy the somatic cell niche in the ovaries that surround the germline, and

the expression of *env* allows the Ty3 virus-like particles to infect the germline to create new copies [21]. As

an adaptation, Drosophila species have duplicated the germline transposon defence mechanism to target

errantiviruses in the somatic cells of the gonad [50]. It is possible that this is more universal, and that *env*-

containing retrotransposons occupy the cell niches surrounding the germline, and that expression of *env*

allows them to continue to infect the germline. For example, murine IAP retrotransposons that carry *env*

genes also remain infectious, like Drosophila errantiviruses [51]. However, there are evolutionary costs

associated with keeping additional ORFs, and genes in endogenous retroviral elements, as they are

constantly subject to evolutionary pressure from their hosts [52,53]. It is therefore significant that the *env* gene

is as highly conserved as we have observed, and that it does not seem to have been subject to frequent loss

and acquisition. Though, the mechanisms of transmission via the action of this *env* gene remain elusive for

most of the elements we identified, except for those in Drosophila species.

However, the structures formed by *env*, for both HSV/gB and glycoprotein F, form the same functional

structures as those found in exogenous viruses, required for cell entry [36,39,40]. While entry of HSV also

requires three other glycoproteins, gD, gH and gL, gB is specifically required for membrane fusion [54,55].

The mechanism of cell entry, therefore, is unknown for these errantiviruses that carry gB, and necessitates

further investigation regarding their expression. Though, given their highly structured configuration, it is

likely that they retain membrane fusion capabilities. Additionally, given the widespread nature of these

env-containing elements, it may also be possible that somatic transposon defence mechanisms against them

are more common.

We found that vERVs are similarly as widespread as errantiviruses, found ubiquitously in vertebrate

classes. Many of the vERVs that we have identified in this study tend to form clades with known

exogenous and endogenous retroviruses, and their phylogeny is consistent with previous studies [18,44].

However, many vERVs form their own clades, without known/annotated retroviruses. Interestingly, most

of the vERVs have multiple intact copies in their respective genomes, meaning that they have been

recently transposing as well (Fig 4).

Of the known classifications, only vERVs resembling Epsilon viruses or SNRVs are found specific to one

vertebrate order, while the rest are found across multiple. This may be an indication that these vERVs have

existed for long periods of time, remaining in the germline, and may not be derived from their exogenous

counterparts. This is supported by the observation that vERVs found within the same vertebrate classes

tend to cluster together, even within larger clades that have a mix of representation from different classes

(Fig S7). Additionally, structural features of the vERVs tend to be different than those of the known

counterparts. For example, MuERV-L, is not known to carry *env* genes [56,57], but vERVs in the same

discrete clade do (Fig 4). Furthermore, these elements are found in reptiles, mammals and birds (Fig S7).

This magnitude of difference between the orders indicate that it is unlikely that they are sourced from the
same exogenous virus. Additionally, the bridge region of most vERVs carries an RNase H domain, a tether
domain, as well as a "mini" domain. The "mini" domain is not shared by many of the annotated retrovirus
in this study (Fig 4). Though, the same "mini" domain is found in yeast Ty3, as well as "ancient
errantiviruses", potentially supporting their common origin. Some vERVs do not have the "mini" domain
either, but they are restricted to discrete clades. An example of this is a discrete clade of vERVs that
resemble the Koala endogenous retrovirus; KoRV-A (Fig S7).

Evidence suggests that KoRV-A is the result of a recent endogenization event of the exogenous retrovirus
KoRV-B [5,6]. Though, we find that KoRV-A forms a clade with vERVs found in Rodentia, Pilosa,
Chiroptera and Perissodactyla, and not marsupials, it is still possible that they are the result of recent
exogenous integrations. However, recent integrations of exogenous viruses cannot solely be responsible for
all the genomic copies identified in this study, given the multicopy nature of these elements, and there is no
evidence of exogenous counterparts to many of the clades of vERVs (Fig 4). It is possible that elements
can be shared via horizontal transfer, potentially mediated by exogenous viruses. However, given the
separation of *pol* clades between different classes of vertebrates, such horizontal transfer is either limited
between closely related species, or does not happen frequently enough to randomize the distribution of *pol*.
The relative frequencies of endogenization and germline expansion, and how they contribute to the
evolution of vERVs remains as an open question. Identification of more exogenous retroviruses and better
understanding of how vERVs make new germline copies will help address this fundamental question in the
future.

As seen in errantiviruses with HSV/gB-type ENV, structural features of GP1 vary within large clades of the
vERV *pol* RT tree (Fig 4 and S7). But they are still consistent in smaller distinct clades, indicating that *pol*
and *env* are co-evolving in vERVs. For example, some discrete clades of gamma retroviruses carry GP1
that keeps both RBD and the C domain, and others only carry the C domain (Fig S7). Additionally, some
vERVs carry GP1 that resembles syncytin-2. This is the first known characterisation of such intact
endogenous viral copies, and most do not cluster with known exogenous retroviruses either. Notably,
vERVs carrying Syncytin-2 are found in different vertebrate orders, potentially indicating its ancient
origin. Importantly, GP1 and GP2 are structurally intact in most vERVs, and likely continue to be
functional as membrane fusion proteins.

The clade of newly identified vERVs resembling SNRVs and Lokiviruses do not carry GP1/GP2, but
instead carry glycoprotein F, consistent with previous studies [18] (Fig 4). Additionally, this clade carries the
"mini" domain, which appears to be a hallmark of an ancient endogenous element (Fig 4). Lokivirus-like
vERVs are also found in more ancient orders of vertebrates, Fish, Amphibians and Jawless fish. The
structural features of Lokiviruses are shared with "ancient errantiviruses", and they indeed appear to be
closely related to them, based on the phylogeny of their ENV as well as POL proteins (Fig S4). These
findings strongly indicate that vERVs and Errantiviruses have a common origin which is highly ancient.

Our study sheds light on the potential origins of *env* containing endogenous retroelements. We propose that

these *env* containing retrotransposons are highly ancient in nature, and in cases where they remain active,

they diverge with their hosts, infrequently recombining with other elements within the same hosts. While

the origin of the *env* genes is still not entirely known, we suggest that both HSV/gB-type and F-type *env*

were anciently acquired before the split of bilaterian and non-bilaterian animals in Pre-Cambrian era, while

GP1+GP2 *env* were later acquired in the common ancestor of vertebrates.

Materials and methods

<u>Genome assemblies used in the study</u>

Reference genome sequences were retrieved from NCBI and grouped into the following insect orders,

vertebrate classes and other metazoan phyla: Diptera, Lepidoptera, Coleoptera, Hymenoptera, Hemiptera,

Polyneoptera, and Palaeoptera for insects; jawless-fish, fish, Amphibian, reptile, bird, and mammal for

vertebrates; Arachnida, Branchiopoda, Chilopoda, Collembola, Diplopoda, Hexanauplia, Malacostraca,

Merostomata, Ostracoda, and Thecostraca for Arthropoda; Acanthocephala, Annelida, Brachiopoda,

Bryozoa, Dicyemida, Mollusca, Nemertea, Onychophora, Orthonectida, Phoronida, Platyhelminthes,

Priapulida, Rotifera, and Tardigrada for Protostomia; Cephalochordata, Echinodermata, Hemichordata, and

Tunicata for Deuterostomia; Cnidaria, Ctenophora, and Porifera. We obtained lists of reference genomes in

each category in May 2023 using the Entrez Direct command line tool except for the vertebrate genomes

which we used the list from the previous work by Wang et al. [18]. The list of the genome assemblies used

can be found in Supplementary table S4.

<u>Identification of active genomic copies of vertebrate endogenous retroviruses (vERVs) and Ty3/*gypsy*</u>

<u>errantiviruses</u>

We ran a total of five iterations of tBLASTn (ncbi-blast-2.9.0) to search for the genomic copies of vERVs

and Ty3/*gypsy* errantiviruses, firstly using the baits of known elements, and subsequently using the

elements that were identified after each iteration. We chose new baits that were not represented in the

phylogenetic tree of the POL (see below for the detail) from the previous iteration and repeated the process

until we no longer found new elements. We used two sets of sequences for the first tBLAST search. The

first set included GAG, POL and ENV protein sequences of the Dipteran *gypsy* errantiviruses from the

RepBase (https://www.girinst.org/repbase/). The second set included GAG, POL and ENV protein

sequences of selected vertebrate retroviruses that were used in a previous study [18]. The bait sequences for

all   tBLASTn   iterations   can   be   found   at

https://github.com/RippeiHayashi/errantiviruses_ERVs_2024/tree/main. We applied different criteria to

identify ERVs from species outside insects and vertebrates as we did not have prior knowledge of whether

or what kind of ERVs are found in those species. We briefly summarise our approach below.

We first used open reading frames (ORFs) of vertebrate retroviruses as baits for the vertebrate genomes

and those of the Dipteran errantiviruses for the other genomes to run tBLASTn with default parameters (-

evalue 10). Genomic regions that showed homology to the POL baits with a bit score of greater than 50 or

to the GAG and ENV baits with a bit score of greater than 30 were individually pooled. Genomic regions

that had hits from multiple baits were merged for their coordinates using bedtools/2.28.0. Regions that

were larger than 1500 nucleotides for POL and 300 nucleotides for GAG or ENV were collected.

Following, for insect genomes we looked for regions that carry GAG, POL, ENV sequences that are next

to each other in this order allowing up to 500 nucleotides of gaps between them as well as excluding

regions of smaller than 4000 nucleotides or larger than 10000 nucleotides. For the other genomes we

allowed up to 1000 nucleotides of gaps between POL and ENV and collected regions that had both ORFs

that are not smaller than 3000 nucleotides or larger than 10000 nucleotides.

We subsequently checked whether the regions contain uninterrupted ORFs by making translated sequences

using transeq from EMBOSS-6.6.0 and excluding ORFs with more than five stop codons. We clustered all

the identified GAG, POL and ENV from each genome by Clustal Omega

(https://www.ebi.ac.uk/Tools/msa/clustalo/) and identified genomic regions that harbour all three ORFs

that are multiplied in the genome. We also searched for ORFs regardless of the copy number when there

were no multiplied elements in the genome. When genomic copies were abundantly found in multiple

closely related species, we only annotated the elements from representative genomes. Genomes that were

annotated and the number of predicted ORFs per genome are summarised in Supplementary table S4.

The above process yielded abundant putative intact copies of ERVs from insect genomes whereas we took

additional steps for the other genomes. For vertebrate genomes, we extended the genomic regions of

predicted POL-ENV to upstream by 2000 nucleotides and manually annotated putative GAG ORFs using

the online sequence viewer Benchling (https://www.benchling.com/). The identity of GAG ORFs were

further confirmed by HHpred (https://toolkit.tuebingen.mpg.de/tools/hhpred). For Arthropoda genomes

outside insects as well as other Protostomia genomes, we extended the genomic regions of predicted GAG-

POL to downstream by 3000 nucleotides and manually annotated ENV ORFs using Benchling. For

Cnidaria, Ctenophora and Deuterostomia genomes outside vertebrates we used transeq to annotate putative

GAG and ENV ORFs from 5000 nucleotides downstream and 3000 nucleotides upstream to the predicted

POL ORFs, respectively.

We took confident 1128 POL ORFs from the first tBLASTn search, including both invertebrate and

vertebrate elements, and made a multiple sequence alignment of the conserved regions of the Integrase

domain using mafft-7.505 with the option '--auto', and inferred the phylogenetic relationships using iqtree-

1.6.12 with the option '-m rtREV+R4 -bb 1000' (see more details in the section below). We took 178

representative POL from the phylogenetic tree and used the POL and the corresponding GAG and ENV

ORFs from the same genomic regions for a second tBLASTn search. We repeated the above steps after

combining the results of the two tBLASTn searches. We did not find any ERVs in Nematoda genomes that

have GAG, POL and ENV in this order while the previous study reported non-canonical Nemetoda ERVs

that have flipped the order of ENV and POL [27]. Thus, we searched for regions that have ENV and POL in

this order allowing gaps of up to 2000 nucleotides between them. GAG was subsequently predicted from

5000 nucleotides upstream to the predicted ENV-POL. We extended the search for potential GAG-ENV-

POL insertions to other non-insect genomes and found similar elements in Bryozoa and Platyhelminthes.

We also found ENV-GAG-POL elements in Arachnida genomes (*Medioppia subpectinata* and *Oppia*

nitens). These elements were recovered through the search of ENV-POL elements because GAG ORF was

short. We did not extend our search to find other arrangements of ORFs.

Additionally, we found that the glycoprotein gene present in mosquito *BEL/pao* retrotransposons [25] are

also found next to *gypsy* POL genes. The glycoprotein gene resembles the *envelope* genes from Chuviridae

and is structurally similar to glycoprotein B from Herpesviruses. We identified ten mosquito *BEL/pao*

retrotransposons that carry this glycoprotein from RepBase (BEL-199_AA, BEL-279_AA, BEL-652_AA,

BEL-801_AA, BEL-804_AA, BEL-879_AA, BEL-51_AnFu, BEL-56_AnFu, BEL-81_AnFu, and BEL-

71_CQ) and used them as baits for a tBLASTn search against Arthropoda genomes. The tBLASTn search

identified 114 *gypsy* elements with homologous glycoprotein genes from Coleoptera, Hymenoptera,

Polyneoptera, and Hexaneuplia species. Combined together, the second iteration of tBLASTn search

identified 491 new elements. The phylogenetic analysis of POL from the elements identified in the first

and second tBLASTn runs revealed clades that only contained new elements from the second tBLASTn

run, suggesting that the first two tBLASTn runs were not comprehensive. Therefore, we took 42 additional

baits to cover all the major clades in the phylogenetic tree for a third tBLASTn search, which identified 87

new elements across metazoan phyla. We repeated tBLASTn runs five times in total; the fourth and fifth

runs used 21 and 7 new baits and identified 53 and 6 new elements, respectively. We stopped the

tBLASTn search after five iterations because the phylogenetic analysis of POL did not reveal new clades.

Following, we further curated newly identified ERVs. Firstly, we used hmmer 3.3.1 (http://hmmer.org/) to

make sure that the ERVs that we identified are classified as Ty3/*gypsy*. We compared POL ORFs from the

newly identified ERVs against the POL ORFs of known retroelements that are available from the Gypsy

Database 2.0 (https://gydb.org/index.php/Main_Page). The Gypsy Database 2.0 collects hmm profiles of

Ty1/Copia, Bel/Pao, Ty3/*gypsy*, and Retrovirus POL proteins. The hmmer search classified all the newly

identified ERVs into Ty3/*gypsy* or Retroviruses. Secondly, we ran HHpred from hh-suite 3.3.0 [58] or the

online tool of HHpred (https://toolkit.tuebingen.mpg.de/tools/hhpred) to predict GAG, ENV, and POL

ORFs using the default search parameters. For the command line HHpred, we first made multiple sequence

alignments using hhblits and the Uniclust database UniRef30/2022_02 (https://uniclust.mmseqs.com/).

This was followed by hhsearch to predict domain structures using the collection of pdb70 from March

2022 (https://wwwuser.gwdg.de/~compbiol/data/hhsuite/databases/hhsuite_dbs/). We used default settings

to run hhblits and hhsearch. ENV and POL ORFs with large truncations were removed after the domain

structure prediction. ERVs that predicted all GAG, ENV and POL ORFs were taken for further analyses.

We additionally ran BLASTn using the genomic regions that cover all three ORFs as baits to interrogate

the copy number of ERVs in the genome where they are found. The results are summarised in

Supplementary table S1 where we counted the number of BLASTn hits with a more than 98% identity in

more than 98% of their length to the bait sequences. A majority of newly identified ERVs (1586 out of

1765 elements) carry ORFs that are found at least twice in the genome, further validating our approach of

finding active or recently active insertions.

<u>Identification of putative long terminal repeats and transfer RNA (tRNA) primer binding sequences</u>

Long terminal repeats (LTRs) of ERVs were identified as follows. Firstly, we fetched the genomic

coordinates of ERV insertions and extended 3000 nucleotides at either ends from the Start codon of GAG

and the Stop codon of ENV ORFs (POL was taken for the GAG-ENV-POL ERVs) for all species

categories except for Cnidaria, Ctenophora and Deuterostomia where we extended 5000 nucleotides at

either ends from the Start and Stop codons of POL. We retrieved the genomic sequence of the extended

ERVs and ran BLASTn against the same sequence to identify repeated sequences. We considered repeated

regions of greater than 99% identity and the tBLASTn bit score between 200 and 5000 as putative LTRs.

We excluded LTRs whose distance between them is shorter than 5000 nucleotides or longer than 14000

nucleotides. Following, we searched for putative tRNA primer binding sequences (PBS). tRNA sequences

were retrieved from genomic annotations from representative genomes in respective species categories

except for Ctenophora where we took tRNA sequences from Protostomia, Cnidaria and Deuterostomia

because there were no annotated genomes available at the time of the study. Representative genomes that

we used to retrieve tRNA sequences are: GCF_000001215.4 and GCF_013141755.1 for Diptera,

GCF_014905235.1 and GCF_905147365.1 for Lepidoptera, GCF_000002335.3 and GCF_917563875.1

for Coleoptera, GCF_003254395.2 and GCF_016802725.1 for Hymenoptera, GCF_014356525.2 and

GCF_020184175.1 for Hemiptera, GCF_002891405.2 and GCF_021461395.2 for Polyneoptera,

GCF_921293095.1 for Palaeoptera, GCF_000671375.1, GCF_020631705.1, GCF_002217175.1,

GCF_016086655.3, GCF_024679095.1, GCF_019059575.1 for Arthropoda, GCF_000326865.2,

GCF_020536995.1, GCF_000002985.6, GCF_000715545.1 for Protostomia, GCF_000222465.1 and

GCF_022113875.1 for Cnidaria, GCF_013122585.1, GCF_000003605.2, GCF_000003815.2, and

GCF_000002235.5 for Deuterostomia, and GCF_015237465.2, GCF_000001405.40, GCF_017654675.1,

GCF_900496995.4, GCF_900700375.1, and GCF_010993605.1 for Vertebrata. The minimum

requirements of the priming events are incompletely understood, but the length of PBS for

retrotransposons is generally between 8 and 18 nucleotides and the PBS is positioned near the 3' end of the

5' LTR [59]. We chose the 12 nucleotides at the 3' end of tRNAs as putative primer sequences and searched

for PBS in the 100 nucleotide-window centred at the 3' end of the 5' LTR using bowtie-1.2.3 allowing up

to two mismatches. We added "CCA" to the 3' end of the tRNA sequence when it is not present in the

genomic sequences. We successfully identified putative PBSs in 1308 out of 1765 ERVs that we identified

in this study. We grouped ERVs based on the type of tRNA used for the PBS and summarised the results

in Figures S2, S3 and S7 as well as Supplementary table S1.

<u>Phylogenetic analysis of POL</u>

We obtained multiple sequence alignments of the RT and Integrase domains of POL as follows. For the

RT domain, we first took the HHpred alignments of POL against the RT domains of Moloney Murine

Leukaemia Virus (Protein Data bank entry 4MH8_A) and Human Immunodeficiency Virus type 1 (Protein

Data bank entry 4G1Q_B), spanning regions that correspond to their fingers, palm, thumb and connection subdomains [60], namely the amino acid positions of 28 to 426 and 7 to 405, respectively. However, this process caused small truncations at either ends of the domain because of poor alignments at the individual substructures. To overcome this issue and to facilitate sequence alignments, we extended the N- and C-termini by 10 and 15 aa, respectively, after merging the two predicted regions of RT domains. We then ran mafft-7.505 to generate a multiple sequence alignment, from which we manually trimmed the either ends to have intact fingers, palm, thumb and connection subdomains for all POL ORFs. Similarly, we took HHpred alignments of POL against the Integrase domains of Human Spumaretrovirus (Protein Data bank entry 3OYM_A) and Simian T-lymphotropic Virus type 1 (Protein Data bank entry 7OUH_B), spanning regions that correspond to their N-terminal domain (NTD) and catalytic core domain (CCD) [61], namely the amino acid positions of 54 to 273 and 8 to 197, respectively. We used the predicted Integrase domains with 15 and 30 aa extensions at the N- and C-termini, respectively, for generating a multiple sequence alignment before manually trimming the ends. We inspected the multiple sequence alignments using Mview (https://www.ebi.ac.uk/Tools/msa/mview/) and removed ERVs that contained truncated RT or Integrase domains from the analyses. We included ORFs from 5 Drosophila errantiviruses (DM176, IDEFIX, ZAM, GYPSY, and Gypsy-5_DGri) and 27 representative vertebrate retroviruses (sequences used in the previous study [18]: BLV, FLV, HTLV-2, KoRV, LDV, MMTV, MuLV, REV, RSV, SRV-1, WDSV, MuERV-L, BFV, EFV, FFV, SFV, HIV-1, EIAV, FIV, SnRV, Loki-Ofa, Loki-Pma, Loki-Sri, Loki-Str, and Loki-Xtr; these two sequences were additionally used in this study: epsilonretrovirus, WEHV1 (AAD30048.1) and avian gammaretrovirus, DIAV (AGV92859.1)) in the multiple sequence alignments.

We ran iqtree-1.6.12 with the option '-m rtREV+R4 -bb 1000' to generate the phylogenetic trees of all Integrase and RT domains, and those from ERVs found in vertebrate and invertebrate genomes, respectively. We then selected representative elements from clades, taking into consideration the relative distances between nodes in the same clade. For example, if the distance between two nodes is less than 0.1, meaning that there is little – no variations between their sequences, only one node in that position is taken. Additionally, we made sure that the representatives captured the diversity of *env* found within the clade. For example, if a clade contains both F and HSV elements, at least one for both is taken, regardless of distance. Furthermore, the phylogeny of the hosts was represented as fully as possible. For example, if a clade has elements from Diptera and Coleoptera, at least one from both is taken regardless of distance. Once representatives were collected, the tree was then accordingly pruned. It should be noted that the final pruned tree of representatives was only used as a figure aid for clarity (Fig 1-4), and all analyses were done on the full INT and RT phylogenetic trees. We used a similar approach as detailed to select representatives for structural analysis of *env* ectodomains, as well as the *pol* bridge region (Table S2). If any outliers were found, remaining elements from the same clade were also analysed, to check whether structural inconsistencies were consistent across the clade.

For congruence plots in figures S2, S4 and S7, we grouped elements based on which node they sit in the

tree, ensuring that the node is well supported by bootstrap values greater than 95. We used the networkD3

package in R to generate Sankey plots to visualise the representation of elements from one tree in another.

The node definitions are summarized in Table S3.

Further, we took 88 representatives of the F-type ENV based on their positions in the corresponding POL

phylogenetic tree for the analysis. We identified the signal peptides and transmembrane regions by

DeepTMHMM (https://dtu.biolib.com/DeepTMHMM) and obtained the ectodomains that are in between

them. We subsequently ran mafft-7.505 to generate a multiple sequence alignment.

<u>Structure prediction of ENV proteins and the RNase H domains using Alphafold 2</u>

We took 88, 32, 130, and 243 representatives of the F-type, HSV/gB-type, and GP2-type ENV proteins

and the region of POL between INT and RT, that we termed the bridge region, using branch definitions

from the phylogenetic trees of *pol* RT. We used the colabfold wrap v1.4.0 of Alphafold 2

(https://github.com/YoshitakaMo/localcolabfold/releases/tag/v1.4.0), and used MMSeqs2 for the database

search and clustering of protein sequences before performing the structural prediction [62]. We used the

whole ectodomains (see above) of the F-type and HSV/gB-type ENV proteins and predicted the trimer

structures using the multimer mode of Alphafold 2 (AlphaFold2-multimer-v2). For the GP2-type ENV

proteins, we first defined the ectodomain that is in between the predicted signal peptides and

transmembrane regions and identified the GP1 domain between the signal peptides and the putative Furin

cleavage site. We predicted monomer structures of the GP1 domains and the bridge regions. We then

visualised and characterised visible structures and disulfide bonds or Cysteine bridges using Maestro

(Schrödinger Release 2024-3) and grouped them by common substructures (Table S2).

Figure Legends

Figure 1 – Phylogeny and copy number of invertebrate metazoan errantiviruses. (A) Open reading

frame arrangement of intact errantivirus copies. **(B)** Summary of Errantiviruses found in Metazoan phyla

with type of *env* protein indicated. **(C)** Maximum likelihood tree constructed on representative *pol* RT

proteins of intact errantiviruses, with "ancient errantiviruses" and "insect errantiviruses" phylogenetic

groups highlighted in green and white respectively. Positions of known dipteran errantiviruses,

Gypsy_ZAM (1), Gypsy_Gypsy5_Dgri (2), Gypsy_Gypsy (3), Gypsy_Idefix (4), and Gypsy_DM176 (5)

are indicated. Host classifications of errantiviruses are annotated in colour, non-arthropod Protostomia

(yellow), Deuterostomia (red), Diploblasta (orange), non-insect Arthropoda (green), Insecta (blue). One

cnidarian element within the group of insect errantiviruses is marked by an asterisk. **(D)** Copy number of

intact errantivirus copies in metazoans. Column represents a genome assembly, each tile representative of

annotated errantivirus, colour of tile represents copy number.

Figure 2 – Representation of *env* in errantiviruses, and structures of errantivirus glycoprotein B. (A)

Shown is a *pol* RT maximum likelihood tree of the representative errantiviruses (see methods) with host

(*) and type of *env* protein (**) indicated. Colours represent non-insect animals (*) (yellow), insects (*)

(blue), glycoprotein F (F) (**) (orange), and glycoprotein B (HSV/gB) (**) (blue). Clades containing
hymenopteran elements in "Insect errantiviruses" that carry either F or HSV/gB *env* are annotated as 1 and
2. **(A')** Close up of clades 1 and 2 in "Insect errantiviruses". F-type and HSV/gB-type errantiviruses that
are both found in *Lasioglossum lativentre* are marked by asterisks. **(B)** Structural arrangements of
Ectodomains, Transmembrane domains and signal peptides found in HSV/gB-type errantiviruses. **(B')**
Shown are the arrangement of domains of the HSV/gB ectodomain, and Alphafold 2-predicted 3D
monomer and trimer structures of the ectodomain. The structure is taken from
Hemiptera_Pemphigidae_Elan_errantivirus_1. **(C)** Fusion loops and conserved Cysteine bridges are
indicated on structure of the errantivirus HSV/gB ectodomain. **(C')** Shown are variable Cysteine bridges in
the errantivirus HSV/gB ectodomains. Cysteine bridge configurations of domains I and II, with proximity
of Cysteine bridge to fusion loops of domain I are indicated. Structures are taken from
Cnidaria_Hydrozoa_Chem_errantivirus_5, Bryozoa_Gymnolaemata_Cpal_errantivirus_2, and
Tunicata_Ascidiacea_Clep_errantivirus_1, for C1, C2 and C3 configurations, respectively. **(D)** Shown is
the summary of the Cysteine bridge configurations found in HSV/gB of different metazoan orders. Exact
positions of the Cysteine bridges can be found in Supplementary Figure S6.

Figure 3 – RNase H domains found in the bridge region of errantiviruses. (A) Shown is the *pol* RT
maximum likelihood tree of representative errantiviruses, with RNase H domain architecture in bridge
region of *pol* highlighted. "Insect errantiviruses" with RNase H + pseudo RNase H (blue), "insect
errantiviruses" with bridge region outliers (dark blue), "ancient errantiviruses" with RNase H + mini
domain (green), "ancient errantiviruses" with bridge region outliers (yellow). Insect and non-insect
elements are annotated as blue and yellow, respectively. **(A')** RNase H domain configurations found in
bridge region of "Insect errantiviruses" and "Ancient errantiviruses", with outliers indicated, and the
structure of yeast Schizosaccharomyces *pombe* Ty3 bridge region predicted by Alphafold 2 is shown. **(B)**
Shown are Alphafold 2-predicted structures of domains found in the bridge regions of errantiviruses, with
topology maps of α helices (grey) and β sheets (red). Structures are taken from
Annelida_Polychaeta_Apac_errantivirus_7, Bryozoa_Gymnolaemata_Cpal_errantivirus_3,
Coleoptera_Coccinellidae_Npum_errantivirus_1, Annelida_Polychaeta_Slim_errantivirus_9,
Diptera_Cecidomyiidae_Smos_errantivirus_1 and Hymenoptera_Apoidea_Obic_errantivirus_2 for RNase
H/Tether, 'mini' RNase H, 'pseudo' RNase H domains, "Mini" domain 2, "Pseudo" domain 2 and
"Pseudo" domain 3, respectively.

Figure 4 – Widespread presence of *env*-containing endogenous retroviruses in Vertebrate classes.

(A) Maximum likelihood tree of *pol* RT from intact *env*-containing endogenous retroviruses (vERVs) in
vertebrate classes. Classification of clades highlighted on tree, and detailed guide to clades in panel B.
Host phylogeny annotated (*); mammals (orange), amphibians (blue), birds (pink), fish (green), reptiles
(yellow), jawless fish (purple), known exogenous retroviruses (white). Bridge region structures annotated
(**); RNase H + RNase H/Tether (orange), RNase H + RNase H/Tether + mini domain (blue). ENV
structure and type annotated (***); GP1 with receptor binding domain (RBD) and C domain (yellow), GP1

with C domain only (blue), Mix of GP1 with RBD+C and GP1 with C only (teal), Glycoprotein F (grey).

(B) Summary of vERV classification, with colours annotated to match tree classification in panel A. The

bridge region architecture, *env* structure, and host phylogeny of clades are shown. **(C)** Structure of

Glycoprotein 1 (GP1) domain of *env* with Alphafold 2-predicted structure of the receptor binding domain

(RBD), C domain, and conserved Cysteine residues highlighted. Shown are the GP1 structures of

Bird_Passeriformes_Nchr_vERV_2 (C only) and Amphibian_Anura_Rmar_vERV_3 (RBD + C). **(D)**

Genomic copy number of vERVs. Columns represent individual genome assemblies grouped by vertebrate

classes, each tile representative of annotated vERV, and colour of tile represents copy number. **(E)** *pol* RT

maximum likelihood tree of representative errantiviruses and vERVs, with "insect errantiviruses" (white),

"ancient errantiviruses" (green), "lokiviruses" (magenta) and "Vertebrate ERVs" (indigo) highlighted, with

env types and bridge region structures (red – RNase H, Blue – mini domain, green – pseudo RNase H)

indicated below. Host phylogeny indicated on tree, Insects (blue), non-insect Arthropoda (green), non-

arthropod Protostomia (yellow), Diploblasta (orange), non-vertebrate Deuterostomia (red), vertebrates

(purple). **(F)** Positional relationship of Lokiviruses to Errantiviruses, with Alphafold 2-predicted 3D

structures of the bridge region and *env* ectodomain. Both structures are taken from

fish_Cyprinodontiformes_Fhet_vERV_1. **(G)** Alphafold 2-predicted 3D structures of bridge region

domains in known vertebrate retroviruses. Shown are structures from MMTV, FLV and SNRV.

Supplementary Figure S1 – Identification of errantiviruses and vERVs in metazoan genomes. **(A)**

Computational approach to identify and annotate *env*-containing retroelements in metazoan genomes. **(B)**

A pruned maximum likelihood tree of *pol* RT from both errantiviruses and vERVs with protein baits for

consecutive tBLASTn searches annotated in colour, showing a comprehensive coverage of the baits in the

tree. **(C)** *pol* RT maximum likelihood trees with representative elements taken for Alphafold 2 analysis

highlighted in red for bridge domains (All RT), F *env* ectodomains (errantiviruses RT only), HSV/gB *env*

ectodomains (HSV/gB-type RT only) and GP1 (vERVs RT only).

Supplementary Figure S2 – Complete errantivirus tree and annotations of tRNA primer binding

sites of insect errantiviruses. **(A)** Complete Maximum likelihood tree constructed on *pol* RT proteins of

intact errantiviruses, with discrete phylogenetic groups within "insect errantiviruses" indicated (I1 - I6).

"ancient errantivirus" group highlighted in green, "insect errantivirus group highlighted in white. Host

phylogeny (*) annotated for insects (blue), non-insect arthropods (green), non-arthropod Protostomia

(yellow), Deuterostomia (Red) and Diploblasta (Orange). ENV type (**) annotated for Glycoprotein F (F)

(orange), and glycoprotein B (HSV/gB) (blue). **(B)** A Sankey plot of *pol* RT and *pol* INT maximum

likelihood trees for errantiviruses, showing an overall phylogenetic congruence between them. **(C)** Shown

are annotations of tRNA primer binding sites of "insect errantiviruses" within phylogenetic groups of *pol*

RT, demonstrating that the types of tRNAs used reflects the separation of *pol* RT. Number of elements are

indicated on Y axis. Sequence information of the tRNA PBS can be found in Table S1.

Supplementary Figure S3 – Ancient errantivirus clades. **(A)** Maximum likelihood tree of *pol* RT for the

"ancient errantivirus" group in isolation, with host phylogeny and *env* type, and discrete phylogenetic

groups (A1-A11) indicated. Host phylogeny (*) is annotated for insects (blue), non-insect arthropods
(green), non-arthropod Protostomia (yellow), Deuterostomia (Red) and Diploblasta (Orange). ENV type
(**) annotated for Glycoprotein F (F) (orange), and glycoprotein B (HSV) (blue). **(B)** Detailed view of
phylogenetic groupings in "ancient errantiviruses" with host (*) and *env* (**) annotated. Bryozoa elements
that are found in distinct clades of the tree are highlighted. Each element is named for the host species and
simplified to Phylum/Class/Order, Superfamily, Genus and Species. For example, elements found in
Diptera Drosophilidae *Drosophila melanogaster* are named as Diptera Drosophilidae Dmel. **(C)** tRNA
annotations of "ancient errantiviruses" within phylogenetic groups of *pol* RT. Insect orders in the groups
A1, A2 and A7 are marked by hash. Number of elements indicated on Y axis.

Supplementary Figure S4 – Glycoprotein F structures and Phylogeny. (A) Architecture of glycoprotein
F ectodomain in Errantiviruses. **(B)** A Sankey plot of *pol* RT and glycoprotein F *env* ectodomain maximum
likelihood trees for representative errantiviruses, showing a phylogenetic congruence between them.
"ancient errantivirus" group is highlighted in green, and vERVs are highlighted in purple. **(C)** Detailed
view of maximum likelihood trees of *pol* RT and *env* F ectodomains of representative elements for
errantiviruses and vERVs. Host species annotated for insects (blue), vertebrates (purple), and other animals
(yellow). **(D)** Alphafold 2-prediction of glycoprotein F ectodomains. Monomeric and trimeric structures of
the "pre-fusion configuration" and the "post fusion configuration" are shown in **(D')** and **(D'')**,
respectively. Schematic cartoons of *env* F ectodomain domain 4 are shown in **(D''')**. Structures are taken
from Diptera_Sciomyzidae_Cmar_errantivirus_19 and Branchipoda_Diplostraca_Dcar_errantivirus_3 for
"pre fusion" and "post fusion" structures in **(D')** and **(D'')**, respectively.

Supplementary Figure S5 – Conserved predicted Furin cleavage sites and hydrophobic fusion
peptides of glycoprotein F. Shown is a multiple sequence alignment of the middle part of the
representative ectodomains of glycoprotein F. Arginine (R) residues in the putative Furin cleavage site and
hydrophobic residues (Leu, Ile, Val, Phe, and Trp) in the fusion peptides are highlighted in red and green,
respectively.   Furin   cleavage   sites   were   predicted   using   ProP-1.0
(https://services.healthtech.dtu.dk/services/ProP-1.0/). Conserved alpha helix structures are shown
upstream and downstream of the Furin cleavage site. Sequences from vertebrate elements are marked by
asterisks.

Supplementary Figure S6 – Conservation of Cysteine bridges in errantivirus HSV/gB ectodomains.
Shown are alignments of domain I and II of all errantivirus HSV/gB ectodomains with Cysteine residues
highlighted in red and hydrophobic residues in the fusion loop peptides highlighted in green. Sequences
from different metazoan orders are separately aligned. Conserved Cysteine bridges within and across the
alignments are numbered, and the tree positions of ERVs are shown where "insect errantiviruses" and
"ancient errantiviruses" are highlighted in white and green, respectively, as in Fig 2A. ERVs that were used
for   the   structural   analysis   are   marked   by   asterisks.   The   structure   of
Hymenoptera_Formicoidea_Tbic_errantivirus_1 is shown to highlight the position of the Cysteine bridge
in domain I of insect errantivirus HSV/gB ectodomain and the absence of the fusion loop II. Note that the

single HSV/gB-type ENV found in Annelida errantiviruses did not predict a trimer structure, therefore, it is not included in the structure classification.

Supplementary Figure S7 – Structural characterisation of vertebrate vERVs. (A) Phylogenetic congruence of *pol* RT and *pol* INT maximum likelihood trees for vertebrate endogenous retroviruses (vERVs). **(B)** *pol* RT maximum likelihood tree of vERVs with discrete phylogenetic groups (V1 – V18) annotated. **(C)** Alphafold 2-predicted 3D structures of GP1 outlier domains; Syncytin-2, expanded C domain, and extra domain. Structures are taken from Mammal_Diprotodontia_Neug_vERV_2, Bird_Galliformes_Csqu_vERV_2 and Amphibian_Gymnophonia_Rbiv_vERV_1 for Syncytin-2 like, expanded C domain, and extra domain GP1 structures, respectively. **(D)** A detailed view of phylogenetic groupings within vERVs is shown. Bridge region structures (*) annotated for RNase H + Tether + Mini (blue) and RNase H + Tether (orange). Host species (**) annotated for mammals (orange), birds (pink), reptiles (yellow), amphibians (blue), fish (green) and jawless fish (purple) as well as known exogenous retroviruses (white). ENV annotated for GP1 with RBD and C domains (yellow), GP1 with C domain only (blue), GP1 with Syncytin-2 (dark blue), and glycoprotein F (grey). Expanded C domain and extra domain annotated with green and red markers respectively. **(E)** tRNA annotations of vERVs within phylogenetic groups of *pol* RT. Number of elements indicated on Y axis.

Supplementary Figure S8 – Conservation of Cysteine residues in GP1 C domains. Shown are alignments of the representative C domains within GP1 structures analysed with Alphafold 2. Only the C domain sequences are shown for the RBD + C group. Cysteine residues (red), Cysteine bridges marked for conserved groups, and Furin cleavage site (cyan) are highlighted. Furin cleavage sites were predicted using ProP-1.0 (https://services.healthtech.dtu.dk/services/ProP-1.0/). Positions of numbered Cysteine bridges within each GP1 structure are consistent across different alignments. The conserved "CxxC" motif is also indicated. Tree positions of elements annotated with the vERV *pol* RT tree are shown. vERV names are coloured according to classes of their origin; fish (green), Amphibians (blue), reptiles (yellow), birds (magenta) and mammals (orange).

Supplementary Table 1 – Master ERV table. Table of all 1765 newly identified *env*-containing retroelements identified in this study, with host phylogeny, genome assembly name and coordinates, contig size, type of ENV, copy number, tRNA PBS type, 3' and 5' LTR coordinates and PBS mismatches. The second tab summarises the tRNA PBS types identified in this study and their 12 nucleotides sequences.

Supplementary Table 2 – Structural Analysis of errantiviruses and vERVs. RNase H domains found in the bridge region of *pol* for representative elements taken for Alphafold 2 analysis (RNase H). List of representative elements taken for Alphafold 2 prediction of *env* F ectodomain, and further analysis (F). List of representative elements taken for Alphafold 2 prediction of GP1 with domain architecture indicated (GP1). List of representative elements taken for Alphafold 2 prediction of HSV/gB ectodomain, with Cysteine bridge configuration indicated (HSV).

Supplementary Table 3 – Node information for errantiviruses and vERVs. List of errantiviruses and

phylogenetic node where they are found for errantivirus RT and INT (INT-RT Nodes Errantivirus FigS2),

vERV RT and INT (INT-RT Nodes vERVs FigS7), as well as F representatives INT and ENV (INT-ENV

Nodes F-type FigS4). Data was then used for congruence plots in Fig S2, S4, and S7.

Supplementary Table 4 – Genome assemblies analysed in this study. List of genome assemblies with

species name, size of assembly, class, family and number of tBLASTn hits for vERV/errantivirus ORFs.

Genomes that we used to annotate vERVs and errantiviruses are indicated. Tabs are individually detailed

for Diptera, Lepidoptera, Coleoptera, Hymenoptera, Hemiptera, Polyneoptera, Palaeoptera, non-insect

Arthropoda, non-arthropod Protostomia, Cnidaria, Ctenophora, Porifera, non-vertebrate Deuterostomia,

and Vertebrata.

Acknowledgements

We acknowledge Julius Brennecke, Kirsten Senti and Alex Greenwood for valuable comments in

preparation of the manuscript. This work is supported by the Australian Research Council (DP210102385).

Author contributions

S.C. and R.H. analysed and interpreted the data and wrote the manuscript. The idea of this work was

conceived by R.H., and S.C. and R.H. together made key findings.

Data Availability

The following data are available from the git repository

(https://github.com/RippeiHayashi/errantiviruses_ERVs_2024/tree/main): the code used to annotate and

analyse genomic copies of errantiviruses and vERVs, genomic sequences and open reading frames of the

newly identified ERVs, pdb files for the predicted structures of ENV proteins and the POL bridge region,

sequences of GAG, POL and ENV open reading frames that are used for tBLASTn search, multiple

sequence alignments and the tree files that are generated in this study for the structural and phylogenetic

analyses.

Conflicts of interests

The authors declare no conflicts of interests.

References

1   Johnson, W. E. Endogenous Retroviruses in the Genomics Era. *Annu Rev Virol* **2**,
862   135-159 (2015). https://doi.org/10.1146/annurev-virology-100114-054945
863   2   Feschotte, C. & Gilbert, C. Endogenous viruses: insights into viral evolution and
864   impact on host biology. *Nat Rev Genet* **13**, 283-296 (2012).
865   https://doi.org/10.1038/nrg3199
866   3   Zhuo, X., Rho, M. & Feschotte, C. Genome-wide characterization of endogenous
867   retroviruses in the bat Myotis lucifugus reveals recent and diverse infections. *J Virol*
868   **87**, 8493-8501 (2013). https://doi.org/10.1128/JVI.00892-13
869   4   Dewannieux, M. & Heidmann, T. Endogenous retroviruses: acquisition, amplification
870   and taming of genome invaders. *Curr Opin Virol* **3**, 646-656 (2013).
871   https://doi.org/10.1016/j.coviro.2013.08.005
872   5   Tarlinton, R. E., Meers, J. & Young, P. R. Retroviral invasion of the koala genome.
873   *Nature* **442**, 79-81 (2006). https://doi.org/10.1038/nature04841
874   6   Zheng, H. *et al.* Koala retrovirus diversity, transmissibility, and disease associations.
875   *Retrovirology* **17**, 34 (2020). https://doi.org/10.1186/s12977-020-00541-1
876   7   Zheng, J., Wei, Y. & Han, G. Z. The diversity and evolution of retroviruses:
877   Perspectives from viral "fossils". *Virol Sin* **37**, 11-18 (2022).
878   https://doi.org/10.1016/j.virs.2022.01.019
879   8   Gasparotto, E. *et al.* Transposable Elements Co-Option in Genome Evolution and
880   Gene Regulation. *Int J Mol Sci* **24** (2023). https://doi.org/10.3390/ijms24032610
881   9   Naville, M. & Volff, J. N. Endogenous Retroviruses in Fish Genomes: From Relics of
882   Past Infections to Evolutionary Innovations? *Front Microbiol* **7**, 1197 (2016).
883   https://doi.org/10.3389/fmicb.2016.01197
884   10   Grandi, N. & Tramontano, E. Human Endogenous Retroviruses Are Ancient Acquired
885   Elements Still Shaping Innate Immune Responses. *Front Immunol* **9**, 2039 (2018).
886   https://doi.org/10.3389/fimmu.2018.02039
887   11   Coffin, J. M., Hughes, S. H. & Varmus, H. E.   (1997).
888   12   Platt, R. N., Vandewege, M. W. & Ray, D. A. Mammalian transposable elements and
889   their impacts on genome evolution. *Chromosome Res* **26**, 25-43 (2018).
890   https://doi.org/10.1007/s10577-017-9570-z
891   13   Mason, A. S., Fulton, J. E., Hocking, P. M. & Burt, D. W. A new look at the LTR
892   retrotransposon content of the chicken genome. *BMC Genomics* **17**, 688 (2016).
893   https://doi.org/10.1186/s12864-016-3043-1
894   14   Sandmeyer, S., Patterson, K. & Bilanchone, V. Ty3, a Position-specific
895   Retrotransposon in Budding Yeast. *Microbiol Spectr* **3**, MDNA3-0057-2014 (2015).
896   https://doi.org/10.1128/microbiolspec.MDNA3-0057-2014
897   15   Doolittle, R. F., Feng, D. F., Johnson, M. S. & McClure, M. A. Origins and evolutionary
898   relationships of retroviruses. *Q Rev Biol* **64**, 1-30 (1989).
899   https://doi.org/10.1086/416128
900   16   Xiong, Y. & Eickbush, T. H. Origin and evolution of retroelements based upon their
901   reverse transcriptase sequences. *EMBO J* **9**, 3353-3362 (1990).
902   https://doi.org/10.1002/j.1460-2075.1990.tb07536.x
903   17   Hayward, A. Origin of the retroviruses: when, where, and how? *Curr Opin Virol* **25**,
904   23-27 (2017). https://doi.org/10.1016/j.coviro.2017.06.006
905   18   Wang, J. & Han, G. Z. A Sister Lineage of Sampled Retroviruses Corroborates the
906   Complex Evolution of Retroviruses. *Mol Biol Evol* **38**, 1031-1039 (2021).
907   https://doi.org/10.1093/molbev/msaa272
908   19   Pelisson, A., Mejlumian, L., Robert, V., Terzian, C. & Bucheton, A. Drosophila
909   germline invasion by the endogenous retrovirus gypsy: involvement of the viral env
910   gene. *Insect Biochem Mol Biol* **32**, 1249-1256 (2002). https://doi.org/10.1016/s0965-
911   1748(02)00088-7

20 Keegan, R. M., Talbot, L. R., Chang, Y. H., Metzger, M. J. & Dubnau, J. Intercellular
viral spread and intracellular transposition of Drosophila gypsy. *PLoS Genet* **17**,
e1009535 (2021). https://doi.org:10.1371/journal.pgen.1009535

21 Song, S. U., Gerasimova, T., Kurkulos, M., Boeke, J. D. & Corces, V. G. An env-like
protein encoded by a Drosophila retroelement: evidence that gypsy is an infectious
retrovirus. *Genes Dev* **8**, 2046-2057 (1994). https://doi.org:10.1101/gad.8.17.2046

22 Brasset, E. *et al.* Viral particles of the endogenous retrovirus ZAM from Drosophila
melanogaster use a pre-existing endosome/exosome pathway for transfer to the
oocyte. *Retrovirology* **3**, 25 (2006). https://doi.org:10.1186/1742-4690-3-25

23 Malik, H. S., Henikoff, S. & Eickbush, T. H. Poised for contagion: evolutionary origins
of the infectious abilities of invertebrate retroviruses. *Genome Res* **10**, 1307-1318
(2000). https://doi.org:10.1101/gr.145000

24 Rohrmann, G. F. & Karplus, P. A. Relatedness of baculovirus and gypsy
retrotransposon envelope proteins. *BMC Evol Biol* **1**, 1 (2001).
https://doi.org:10.1186/1471-2148-1-1

25 Dezordi, F. Z., Vasconcelos, C. R. D. S., Rezende, A. M. & Wallau, G. L. In and Outs
of Chuviridae Endogenous Viral Elements: Origin of a Potentially New Retrovirus and
Signature of Ancient and Ongoing Arms Race in Mosquito Genomes. *Front Genet* **11**,
542437 (2020). https://doi.org:10.3389/fgene.2020.542437

26 Jehle, J. A. *et al.* Molecular identification and phylogenetic analysis of baculoviruses
from Lepidoptera. *Virology* **346**, 180-193 (2006).
https://doi.org:10.1016/j.virol.2005.10.032

27 Sacco, M. A., Lau, J., Godinez-Vidal, D. & Kaloshian, I. Non-canonical nematode
endogenous retroviruses resulting from RNA virus glycoprotein gene capture by a
metavirus. *J Gen Virol* **103** (2022). https://doi.org:10.1099/jgv.0.001739

28 Rodriguez, F., Kenefick, A. W. & Arkhipova, I. R. LTR-Retrotransposons from Bdelloid
Rotifers Capture Additional ORFs Shared between Highly Diverse Retroelement
Types. *Viruses* **9** (2017). https://doi.org:10.3390/v9040078

29 Bargues, N. & Lerat, E. Evolutionary history of LTR-retrotransposons among 20
Drosophila species. *Mob DNA* **8**, 7 (2017). https://doi.org:10.1186/s13100-017-0090-
3

30 Stefanov, Y., Salenko, V. & Glukhov, I. Drosophila errantiviruses. *Mob Genet*
Elements **2**, 36-45 (2012). https://doi.org:10.4161/mge.19234

31 Terzian, C., Pélisson, A. & Bucheton, A. Evolution and phylogeny of insect
endogenous retroviruses. *BMC Evol Biol* **1**, 3 (2001). https://doi.org:10.1186/1471-
2148-1-3

32 Sormacheva, I. *et al.* Vertical evolution and horizontal transfer of CR1 non-LTR
retrotransposons and Tc1/mariner DNA transposons in Lepidoptera species. *Mol Biol*
Evol **29**, 3685-3702 (2012). https://doi.org:10.1093/molbev/mss181

33 Wanner, N. M. & Faulk, C. Suggested Absence of Horizontal Transfer of
Retrotransposons between Humans and Domestic Mammal Species. *Genes (Basel)*
12 (2021). https://doi.org:10.3390/genes12081223

34 Cordaux, R. & Batzer, M. A. The impact of retrotransposons on human genome
evolution. *Nat Rev Genet* **10**, 691-703 (2009). https://doi.org:10.1038/nrg2640

35 Gagnier, L., Belancio, V. P. & Mager, D. L. Mouse germ line mutations due to
retrotransposon insertions. *Mob DNA* **10**, 15 (2019). https://doi.org:10.1186/s13100-
019-0157-4

36 McLellan, J. S., Ray, W. C. & Peeples, M. E. Structure and function of respiratory
syncytial virus surface glycoproteins. *Curr Top Microbiol Immunol* **372**, 83-104
(2013). https://doi.org:10.1007/978-3-642-38919-1_4

37 Dang, H. V. *et al.* An antibody against the F glycoprotein inhibits Nipah and Hendra
virus infections. *Nat Struct Mol Biol* **26**, 980-987 (2019).
https://doi.org:10.1038/s41594-019-0308-9

38 Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature*
596, 583-589 (2021). https://doi.org:10.1038/s41586-021-03819-2

39   Heldwein, E. E. *et al.* Crystal structure of glycoprotein B from herpes simplex virus 1.
Science **313**, 217-220 (2006). https://doi.org:10.1126/science.1126548
40   Connolly, S. A., Jackson, J. O., Jardetzky, T. S. & Longnecker, R. Fusing structure
and function: a structural view of the herpesvirus entry machinery. *Nat Rev Microbiol*
9, 369-381 (2011). https://doi.org:10.1038/nrmicro2548
41   Lopper, M. & Compton, T. Disulfide bond configuration of human cytomegalovirus
glycoprotein B. *J Virol* **76**, 6073-6082 (2002). https://doi.org:10.1128/jvi.76.12.6073-
6082.2002
42   Vollmer, B. *et al.* The prefusion structure of herpes simplex virus glycoprotein B. *Sci*
Adv **6** (2020). https://doi.org:10.1126/sciadv.abc1726
43   Malik, H. S. & Eickbush, T. H. Phylogenetic analysis of ribonuclease H domains
suggests a late, chimeric origin of LTR retrotransposable elements and retroviruses.
Genome Res **11**, 1187-1197 (2001). https://doi.org:10.1101/gr.185101
44   Gifford, R., Kabat, P., Martin, J., Lynch, C. & Tristem, M. Evolution and distribution of
class II-related endogenous retroviruses. *J Virol* **79**, 6478-6486 (2005).
https://doi.org:10.1128/JVI.79.10.6478-6486.2005
45   Lee, J. E. *et al.* Structure of the Ebola virus glycoprotein bound to an antibody from a
human survivor. *Nature* **454**, 177-182 (2008). https://doi.org:10.1038/nature07082
46   Hötzel, I. Deep-Time Structural Evolution of Retroviral and Filoviral Surface Envelope
Proteins. *J Virol* **96**, e0006322 (2022). https://doi.org:10.1128/jvi.00063-22
47   Hogan, V. & Johnson, W. E. Unique Structure and Distinctive Properties of the
Ancient and Ubiquitous Gamma-Type Envelope Glycoprotein. *Viruses* **15** (2023).
https://doi.org:10.3390/v15020274
48   Loreto, E. L., Carareto, C. M. & Capy, P. Revisiting horizontal transfer of transposable
elements in Drosophila. *Heredity (Edinb)* **100**, 545-554 (2008).
https://doi.org:10.1038/sj.hdy.6801094
49   Metzger, M. J., Paynter, A. N., Siddall, M. E. & Goff, S. P. Horizontal transfer of
retrotransposons between bivalves and other aquatic species of multiple phyla. *Proc*
Natl Acad Sci U S A **115**, E4227-E4235 (2018).
https://doi.org:10.1073/pnas.1717227115
50   Brennecke, J. *et al.* Discrete small RNA-generating loci as master regulators of
transposon activity in Drosophila. *Cell* **128**, 1089-1103 (2007).
https://doi.org:10.1016/j.cell.2007.01.043
51 Ribet, D. *et al.* An infectious progenitor for the murine IAP retrotransposon:
emergence of an intracellular genetic parasite from an ancient retrovirus. *Genome*
Res **18**, 597-609 (2008). https://doi.org:10.1101/gr.073486.107
52 Wang, J. & Han, G. Z. Frequent Retroviral Gene Co-option during the Evolution of
Vertebrates. *Mol Biol Evol* **37**, 3232-3242 (2020).
https://doi.org:10.1093/molbev/msaa180
53 Wang, W., Zhao, H. & Han, G. Z. Host-Virus Arms Races Drive Elevated Adaptive
Evolution in Viral Receptors. *J Virol* **94** (2020). https://doi.org:10.1128/JVI.00684-20
54 Johnson, D. C., Wisner, T. W. & Wright, C. C. Herpes simplex virus glycoproteins gB
and gD function in a redundant fashion to promote secondary envelopment. *J Virol*
85, 4910-4926 (2011). https://doi.org:10.1128/JVI.00011-11
55 Jambunathan, N. *et al.* Two Sides to Every Story: Herpes Simplex Type-1 Viral
Glycoproteins gB, gD, gH/gL, gK, and Cellular Receptors Function as Key Players in
Membrane Fusion. *Viruses* **13** (2021). https://doi.org:10.3390/v13091849
56 Kigami, D., Minami, N., Takayama, H. & Imai, H. MuERV-L is one of the earliest
transcribed genes in mouse one-cell embryos. *Biol Reprod* **68**, 651-654 (2003).
https://doi.org:10.1095/biolreprod.102.007906
57 Ribet, D. *et al.* Murine endogenous retrovirus MuERV-L is the progenitor of the
"orphan" epsilon viruslike particles of the early mouse embryo. *J Virol* **82**, 1622-1625
(2008). https://doi.org:10.1128/JVI.02097-07

58   Steinegger, M. *et al.* HH-suite3 for fast remote homology detection and deep protein
annotation. *BMC Bioinformatics* **20**, 473 (2019). https://doi.org:10.1186/s12859-019-
3019-7
59   Wilhelm, M. & Wilhelm, F. X. Reverse transcription of retroviruses and LTR
retrotransposons.   *Cell   Mol   Life   Sci*   **58**,   1246-1262   (2001).
https://doi.org:10.1007/PL00000937
60   Das, D. & Georgiadis, M. M. The crystal structure of the monomeric reverse
transcriptase from Moloney murine leukemia virus. *Structure* **12**, 819-829 (2004).
https://doi.org:10.1016/j.str.2004.02.032
61   Engelman, A. & Cherepanov, P. Retroviral Integrase Structure and DNA
Recombination   Mechanism.   *Microbiol   Spectr*   **2**,   1-22   (2014).
https://doi.org:10.1128/microbiolspec.MDNA3-0024-2014.
62   Mirdita, M. *et al.* ColabFold: making protein folding accessible to all. *Nat Methods* **19**,
679-682 (2022). https://doi.org:10.1038/s41592-022-01488-1

1034

Figure 1

Figure 2

Figure 3

A

A'

pol

Reverse Transcriptase (RT)	Bridge Region	Integrase (Int)

Insect Errantiviruses

RNase H/Tether + "Pseudo" RNase H

RNase H/Tether + "Pseudo" RNase H + "Pseudo" RNase H[2]

RNase H/Tether + "Mini" domain 2[1]

RNase H/Tether + "Pseudo" domain 2[3]

RNase H/Tether + "Mini" domain[3]

RNase H/Tether + "mini" domain 3[1,4]

RNase H/Tether[4,5]

"Ancient" Errantiviruses

RNase H/Tether + "Mini" domain

RNase H/Tether + "Pseudo" domain 2[6]

RNase H/Tether + "Mini" domain 2[7]

RNaseH/Tether

"Mini" Domain

Schizosaccharomyces *pombe* tf1 Bridge region

Insects

Other Animals

pol

RT	INT

Bootstrap values >95 ●

B

RNaseH/Tether   "Mini" Domain   "Pseudo" RNaseH   "Mini" Domain 2   "Pseudo" Domain 2   "Mini" Domain 3

Figure 4

Supplementary Figure S1

A

Step 1. identify GAG, POL, ENV open reading frames (ORFs) by tBLASTn
using GAG, POL and ENV from RepBase Dipteran errantiviruses as baits against **1901 invertebrate genome assemblies**
using GAG, POL and ENV from RefSeq vertebrate retroviruses as baits against **747 vertebrate genome assemblies**

Step 2. identify loci where GAG, POL and ENV ORFs are next to one another

- Only the GAG-POL-ENV elements were annotated in the insect and vertebrate genomes.
- For non-insect Arthropoda and other Protostomia genomes, ENV was predicted from the downstream region of GAG-POL.
- For non-vertebrate Deuterostomia, Cnidaria and Ctenophora genomes, ENV was predicted from the downstream region of POL.
- GAG-ENV-POL elements were found in Nematoda, Bryozoa and Platyhenminthes genome.
- ENV-GAG-POL elements were uniquely found in Arachnida genome.

Step 3. cluster ERVs per genome using clustal omega and pick representative genomic copies

Step 4. curate the identified ERVs by domain structure prediction using hmmscan and HHpred
***All** three ORFs are required to predict homology to known viral proteins listed below*
GAG and POL: homologous to Ty3/gypsy or retroviral GAG and POL
ENV: homologous to paramyxovirus glycoprotein F (F-type), herpesvirus glycoprotein B (HSV/gB-type), or retrovirus glycoprotein (GP2-type)

Step 5. repeat the process until all clades in the phylogenetic tree of POL are represented by tBLASTn baits

tBLASTn interations	identified ERVs
1st	1128
2nd	491
3rd	87
4th	53
5th	6

new clade

baits used in previous runs

B

pol
RT | INT

Second Set
Third Set
Fourth Set
Fifth Set

C

Bridge Regions

HSV/gB Ectodomains

F Ectodomains

GP1 Domains

Supplementary Figure S2

Supplementary Figure S3

Supplementary Figure S4

A

Furin Cleavage Site

| Signal Peptides | Ectodomain | Transmembrane Domain |

N | 3 | 1 | 2 | αα | 4 | 2 | 1 | 3 | A | C

Furin Cleavage Site

B

env — RT

C

env

SS | Ectodomain | TM

pol

RT | INT

D' "Pre-fusion"

Domain A
Domain 3
N
Domain 1
Domain 2
Furin Cleavage Site
αα
Domain 4
Ectodomain Trimer

D'' "Post-Fusion"

Domain 3
N
Domain 1
Domain 2
αα
Furin Cleavage Site
Domain 4
Domain A
C
Ectodomain Trimer

D'''

pre-fusion

Domain 2
R
αα
Domain 4

post-fusion

Domain 2
R
αα
Domain 4

Ancient Errantiviruses | Insects | Bootstrap values >95 ●

Insect Errantiviruses | Other Animals

vERVs | Vertebrates

Supplementary Figure S5

Supplementary Figure S6

Supplementary Figure S7

Supplementary Figure S8

C only

RBD + C

Syncytin-2

Fish
Amphibians
Reptiles
Birds
Mammals