<u>Intact genomic copies of envelope carrying retroelements are found widespread across metazoans</u>

Shashank Chary[1] and Rippei Hayashi[1, 2 #]

1. The John Curtin School of Medical Research, The Australian National University, 131 Garran Road,
Acton, ACT 2601, Australia

2. The Shine-Dalgarno Centre for RNA Innovation, The Australian National University, 131 Garran Road,
Acton, ACT 2601, Australia

correspondence: rippei.hayashi@anu.edu.au.

Abstract

Endogenous retroviruses (ERVs) are generally seen as remnants of past retroviral infections, but some can
make new germline insertions that are transmitted across generations. ERVs can carry membrane fusion
glycoproteins, or *env* genes, which allow them to remain infectious. We systematically searched for such
active ERVs containing *env* in metazoan genomes and found that multiple intact copies are ubiquitously
present across taxa including ancient animals such as Cnidarians, Ctenophores and Tunicates. Phylogenetic
analyses revealed that the divergence of ERVs, including those found in vertebrate and insect genomes,
largely follow the divergence of host organisms, and likely share a common and highly ancient origin.

Introduction

Retroviruses have two mechanisms through which they survive; most commonly, they exogenously infect other hosts, transmitting horizontally [1]. However, they can occasionally infect the germline cells of the host; where genetic information is stored for the next generation, and integrate into the genome [1,2]. In this way, they survive by transmitting vertically down into the next generation. These proviral copies in the germline genome are referred to as endogenous retroviruses (ERVs) [1,3-5]. ERVs are best characterised in vertebrates and make up 5 to 10% of their genomes. An intact copy of an ERV would contain viral *gag*, *pol* and *env* genes, and be flanked by long terminal repeats (LTRs), broadly classified under class I LTR retrotransposons [6].

LTR retrotransposons are ubiquitously found in higher eukaryotes. One of the best studied examples is Ty3/Gypsy. Interestingly, a branch of *gypsy* transposons in Drosophila species carry *env* genes and are referred to as errantiviruses [7,8]. When the defence mechanism against transposable elements is affected, errantiviruses are expressed in the somatic cells of the female ovaries, forming virus like molecules that infect the adjacent germline cells [9,10]. This means that if the errantivirus remains active in the somatic cells, it can maintain the generation of new copies that can be vertically transmitted to the next generation.

Though Drosophila errantiviruses share structural similarities to vertebrate ERVs, their exact origin is poorly understood. It has been suggested that Ty3 elements acquired the *env* gene from infecting baculoviruses in an ancestral Drosophila species [11,12]. However, a preliminary search revealed that errantiviruses may be more widespread than previously thought. This combined with their unique cell niche in the Drosophila gonad prompted us to further explore their origins, as well as how widespread they are.

Results

We systematically searched for homologous elements of vertebrate ERVs and errantiviruses in the referenced genomes across metazoan phyla using the annotated protein coding sequences of *gag*, *pol* and *env* for tBLASTn searches (see methods). We focused our search on the genomic copies that are recently acquired to understand relationships of currently active elements. We identified 1610 elements that carry nearly intact open reading frames of all three genes (Fig 1A, Table S1). A majority of the newly identified genomic copies carry putative LTRs and tRNA primer binding sequences (1205 elements, summarised in Fig S1B and Table S2), suggesting their recent integrations. We ran multiple sequence alignments using the conserved Integrase and Reverse Transcriptase (RT) domains of Pol as well as the Env proteins and assembled maximum probability trees to infer phylogenetic relationships of the identified elements. ORFs of selected vertebrate retroviruses and Drosophila errantiviruses were also included in the tree to aid classifications. We refer to elements found within and outside Vertebrata as vERVs and *gypsy* errantiviruses, respectively.

We found that intact copies of errantiviruses are ubiquitously present across most major metazoan phyla. This includes species from ancient phyla such as Cnidaria, Tunicates and Ctenophora, but excludes

Echinodermata and Porifera (Fig 1B). We similarly find that intact copies of *env*-containing vERVs are

found in all vertebrate orders including jawless fish, which is consistent with previous studies[13,14]. In both

cases, each element has multiple intact copies in their respective host genomes (Table S1), indicating that

they have likely been actively transposing recently.

We assembled phylogenetic trees using the sequences of the *pol* Integrase and RT domain, and identified

discrete clades based on distance and supported by bootstrap values greater than 95 (Fig S1A, S2A) When

we investigated where the elements in the Integrase tree are represented in RT, we found that the trees

assembled are congruent with each other (Fig S2B, Table S3). For example, RT clade A only consists of

elements found in Integrase clade A. In almost all cases, the clades in one tree are represented by similar

clades in the other. RT clade N, for example, mostly consists of elements from Integrase clade O, but also

contains elements from clade P, as well as elements that are not present in a clade but sit between Integrase

clades N, O, P and Q. This congruence suggests that the Pol sequences analysed in this study have not

undergone recent degradation, further confirming the 'intactness' of these elements.

We then characterised the type of tRNA primer binding sites (PBS) that are found in the errantiviruses that

we identified in this study. Drosophila errantiviruses have been identified to have Serine or Lysine tRNAs

priming the elements [15,16]. In our investigation we find that insect elements in Integrase clades A to N

predominantly carry the same tRNA^{Lys} and tRNA^{Ser} PBS (Fig S1B). Clades A and H, which contain

Diptera and Hemiptera elements respectively, mostly have tRNA^{Lys} PBS, while the rest predominantly use

Serine. The exception, however, is clade N, which consists of elements carrying tRNA^{Leu}, tRNA^{Arg},

tRNA^{Glu}, tRNA^{His} and tRNA^{Pro} in addition to tRNA^{Ser} PBS. The tRNA primer binding site for Leucine is

the most prominent in this clade and is found in elements from insect orders (Table S2). The tRNA^{Ser} PBS

in this clade is found in Hymenoptera and Diptera, while the rest outside this clade are only found within

Hymenoptera. Outside of insects, in Integrase clades O and P, the tRNA PBS becomes more diverse,

appearing to reflect the phylogeny of the elements. However, there appears to be some consistency in the

primers. For example, Branchiopoda elements in clade O all use tRNA^{Ala}, Coleoptera elements in the same

clade continue to use mainly tRNA^{Ser} and tRNA^{Leu}, and Mollusca elements mainly use tRNA^{His} (Table S2,

Fig S1B). Some non-insect Arthropoda elements as well as two elements from Mollusca within these

clades also use tRNA^{Leu}.

Elements in Vertebrate clades Q-V also have a large variety of tRNA PBS, but follow a similar pattern to

clade O. For example, tRNA^{Pro}, tRNA^{Gly} and tRNA^{Gln} are mainly used by mammalian elements with a few

elements from the other orders also using the same, while tRNA^{Phe} and tRNA^{Ser} are mainly used by

elements in fish. While the type of tRNA PBS carried by an element tends to overall be roughly respective

of their host species and their branch, the pattern is not consistent enough to draw a conclusion as to

whether they reflect the overall divergence of the errantiviruses and vERVs on the tree.

The phylogenetic tree assembled using the sequences of the *pol* Integrase domains appears to separate into

clades that reflect the hosts that they are found in (Fig 1A). For example, Integrase clades A, B and C

exclusively contain dipteran elements, whereas clades F and J are exclusive to Lepidoptera. Some clades

contain a mix of hosts, though they are normally related. The Integrase clade E consists of mostly

coleopteran elements, but also contains some elements found in other holometabolous insect orders

Hymenoptera and Diptera as well. The notable exceptions are Integrase clades O and P, where elements

found in non-insect Arthropoda, non-Arthropoda Protostomia and ancient orders are found. Though, within

each respective clade, they still appear to separate based on their host (Fig S3C-D). Gamma-retroviruses

are found in Integrase clade S, while other known retroviruses, including alpha-, beta-, delta-, lenti- and

formy-retroviruses, are all found in Q. The more recently identified loki-retroviruses[17] are found in clade R

(Fig 1A, Table S3). Known dipteran errantiviruses Gypsy-Gypsy, and Gypsy-5-Dgri sit within clade A,

Gypsy-17.6 and Idefix sit within clade B, and Gypsy-ZAM sits within clade C (Fig 1A, Table S3). This

arrangement is consistent with previous studies that have classified retroviruses and Drosophila *gypsy*

elements [17-19].

To further explore the congruence between host phylogeny, we specifically identified discrete clades for

Arthropoda (A1-A24) and Vertebrata (V1-V12) in the *pol* Integrase tree and investigated the host makeup

of those clades (Fig S3A-B). The balloon plot is presented as a proportion of representation. A data point

of 0.5 would indicate that 50% of the elements from that host are represented in that clade. We found that

most elements within Arthropoda and Vertebrata clades are formed by elements either from single or

multiple related orders or classes (Fig S3B). For example, clades V3 and V5 only contain elements found

in mammals, and V1 contains elements from birds, but also contains elements from Mammals and

Reptiles. Within Arthropoda, the pattern is similar, though it should be noted that some of the elements

found in insects that sit within clades that contain non-insect Arthropoda carry a different type of *env* than

the others, namely, an *env* that resembles that from herpesviruses.

We further examined the Env protein found in these intact copies of errantiviruses and vERVs, as well as

the divergence between Env and Integrase. We firstly found that the errantiviruses identified in this study

carry two types of Env; those resembling fusion glycoproteins from Nipah viruses (referred to as

glycoprotein F), and those resembling glycoprotein B from herpesviruses (referred to as HSV) (Fig 2,

S4A-B). The vERVs identified also carry glycoprotein F, but most carry glycoprotein GP2, a typical

vertebrate retrovirus Env. While individual clades contain a mixture of Env representation, the congruence

between Integrase and Env appears to hold (Fig 2, Fig S4, Table S5). For example, Integrase clade O

contains both elements that carry HSV and F (Fig 2C), meaning that similar Pol proteins carry different

Env. However, when individual trees are made using only HSV or F, elements appear to form similar

clades between Integrase and Env (Fig S4), meaning that similar Pol proteins carry similar Env genes

individually for HSV or F. For example, clade H in the Integrase tree for HSV consists of elements only

from clade H in the Env HSV tree (Fig S4B). Similar congruence can be seen between Integrase and Env

for glycoprotein F (Fig S4A), though elements that carry GP2 are less congruent than either F or HSV (Fig

S4C).

We subsequently characterised the region between the Integrase and RT domains of Pol, referred to as the

"bridge region", for representative elements within the Integrase/RT phylogenetic trees. The bridge region

is known to carry a ribonuclease H (RNase H) domain, which in vertebrate retroviruses has duplicated, and
the N-terminal copy, while partially degraded, maintains the same structure and 3D fold arrangement,
referred to as "tether" [20]. We therefore characterised structures within the bridge region using the 3D fold
arrangement of the core β-sheet structure. Aside from a few elements in insects, nearly all elements have
duplicated the RNase H domain, which have undergone degradation and structural changes. We mainly
found three types of structures widespread in the bridge region of Pol; the typical RNase H/Tether
arrangement, a "pseudo" RNase H domain, and a "mini" domain (Fig 3C). The "pseudo" RNase H-like
domain retains the barrel-like structure made of five core β-sheets, but the order of the second and the third
β-sheets is flipped. The "mini" domain consists of three small core β-sheets. In addition, we found three
other structures that are less commonly found in the bridge region (Fig S5A). The separation of clades in
the Integrase tree appears to agree with the domain configurations of the bridge region of the elements,
further supporting our clade definition. For example, all representatives taken from Integrase clade A (Fig
1) carry the "pseudo" domain (Fig 3A). Furthermore, Integrase clades O and P, which contains a wide
variety of host species, mostly carry the "mini" domain. The exception is Integrase clade N, whose
elements carry either the "pseudo" domain, or "pseudo" domain 3 (Fig 3A, Fig S5), though within the
clade, the elements are still separated (Fig 3A).

The "pseudo" RNase H domain is mostly specific to insects only, whereas vertebrate elements are the only
ones to have two RNase H-like/Tether domains within their bridge region (Fig 3). However, most
vertebrate elements additionally carry the "mini" domain, which almost all elements in ancient animals,
some elements in insects, and even yeast Ty3 elements carry in the bridge region of their Pol (Fig S5B).
While insect elements mainly carry a "Pseudo" RNase H, some can also carry other uncommon domains,
and they overall have greater variability in the structures present in the "bridge region". It should be noted
that one Cnidaria element that sits in a clade with insects on the Integrase and RT trees carries the
"pseudo" domain, and that the annotated Loki-retrovirus "loki-xtr" only carries one RNase H domain and
the "mini" domain (Table S6).

Our findings show that multiple intact copies of errantiviruses and *env*-containing vERVs are ubiquitously
present in metazoan genomes, indicating recent activity. Additionally, the errantiviruses identified in this
study adhere to their hosts, and the phylogeny of ORFs for *env* and *pol* resemble each other. Finally,
vERVs and errantiviruses share common features such as structures in the bridge domain of Pol, the
structure of Env as well as their host-adhering nature.

Discussion

We found that *env*-containing errantiviruses are more widespread than previously thought. Genomes from
nearly every major order contain multiple intact copies, with uninterrupted ORFs and intact flanking
LTRs, which is suggestive of recent transposition. Though they are recent, they still appear to reflect the
phylogeny of their hosts, meaning that errantiviruses found from one order are more like elements found
either within the same order, or closely related ones, though there are exceptions which will be discussed
shortly. Non-insect errantiviruses are structurally distinct from insect errantiviruses, which is reflected in

their own separate clades. For example, they do not carry a "pseudo" RNase H domain within the bridge

region of Pol, but rather carry a "mini" domain. However, there are still insect errantiviruses present in

these distinct clades, carrying same structural features as non-insect errantiviruses, with their Pol

resembling those from ancient animals more than they do other insects. These findings suggest that the

errantiviruses found in insects may have more ancient origins than previously thought, in addition to being

highly widespread. It was originally speculated that errantiviruses acquired *env* from baculoviruses in

insects, some of which carry an F-type ENV protein Ld130, potentially via recombination events [11,12].

While Ty3 elements may have obtained the *env* gene via recombination, it is unlikely that baculoviruses

are the source for the *env* gene found in errantiviruses, as they are more widespread than insects, and can

carry different types of *env* genes as well. Additionally, whether errantiviruses originally acquired HSV or

F type *env* is also unclear, as non-insect clades which include elements found in ancient animals, carry

both types of *env*. Insect errantivirus clade N (Fig 2) also contains both types of *env*, but the elements that

carry HSV type *env* are distinct from the other errantiviruses within this clade, further indicating that

recombination events may have happened, but they are not happening frequently enough to randomise

their spread.

There are a few ways to explain the widespread and multicopy nature of errantiviruses in metazoans, one

of which is that these errantiviruses frequently undergo horizontal transfer. Drosophila errantiviruses have

been previously suggested to horizontally transfer between species [19,21] and would explain their widespread

presence across metazoans. It has also been suggested that some LTR retrotransposons in bivalves can

horizontally transfer between different species and can even jump across phyla as well [22]. While horizontal

transfer of errantiviruses between distantly related species cannot be excluded, it is likely to be a rare

occurrence, given the congruence of the phylogeny of *int* with that of their hosts. Additionally, some of the

different elements that cluster together are found in hosts that are magnitudes of order apart from each

other, and it is unlikely that they can easily transfer. Furthermore, it is unlikely that horizontal transfer

followed by re-entry into the germline can happen frequently enough to explain their widespread nature.

Alternatively, the widespread nature of these errantiviruses can be explained via an "endogenized

retrovirus model", in which exogenous retroviruses enter the germline by chance, and become fixed in the

population, where they remain as evolutionary relics of past infection, degrading over time [23,24]. Those

exogenous viruses would likely be species specific, given that they are mostly unlikely to freely move

away from their natural host species, often referred to as a species barrier [25,26]. This model is supported by

studies that have identified exogenous virus counterparts of endogenous retroviral copies, for gibbon ape

leukaemia viruses and koala retroviruses [27,28]. For example, the koala gamma retrovirus KoRV-A is the

endogenous form of the exogenous KoRV-B retrovirus [29]. This model would explain the phylogeny of the

errantiviruses following that of their hosts. However, exogenous retroviruses are widely considered to be

vertebrate-specific[30]. We find it unlikely that elements that are as abundant and ubiquitous as we observed

in invertebrate hosts remain undetected to date. Furthermore, some insect elements resemble those from

ancient animals such as non-insect arthropods, Nematoda, Platyhelminthes and Thecostraca (Fig S3C-D),

which would indicate that the potential exogenous viruses are not limited to one host species, contrary to
the "endogenized retrovirus model".

Another possibility that explains both the phylogeny as well as the widespread and multicopy nature of
these elements, is that they are highly ancient germline transposons that continue to remain active. In this
scenario, the last common ancestor of metazoans would have had this element in their genome, which
remained active as species diverged. As long as one of the transposon copies remains active, it would be
vertically inherited, and diverge along with the genome of the host species, whereas older and inactive
copies would be degraded over time. Additionally, it is possible that *env* was present in the original LTR
retrotransposon, or an ancient recombination event is responsible for the presence of the *env* gene.

One question that arises, is why these elements continue to retain *env* over such long periods of time,
remaining potentially infectious when expressed. In Drosophila, errantiviruses occupy the somatic cell
niche in the ovaries that surround the germline, and the expression of *env* allows the Ty3 virus-like
particles to infect the germline to create new copies [9]. As an adaptation, Drosophila species have
duplicated the germline transposon defence mechanism to target errantiviruses in the somatic cells of the
gonad [31] . It is possible that this is more universal, and that *env*-containing retrotransposons occupy the cell
niches surrounding the germline, and that expression of *env* allows them to continue to infect the germline.
For example, murine IAP retrotransposons that carry *env* genes also remain infectious, similar to
Drosophila errantiviruses [32]. Given the widespread nature of intact copies of *env* containing retroelements
identified in this study, it may, therefore, also be possible that a defence mechanism against them is more
common.

We finally found that vERVs are similarly as widespread as errantiviruses, and like them, also carry
multiple intact copies in their respective genomes. While retroviruses are known to have duplicated the
RNase H domain and degrade the catalytic residues to form a 'tether' domain [20], duplications of RNase H
have frequently occurred in errantiviruses as well, and have diverged into a "pseudo" domain. Although
the "tether" domain is strictly found in vERVs, the duplication of the RNase H domain alone may not
separate retroviruses from other errantiviruses. We also identified a "mini" domain towards the C-terminus
of the bridge region of vERVs, which is a feature shared with errantiviruses in ancient animals, and even
yeast Ty3 elements. More strikingly, some vERVs carry F-type *env* proteins, which are carried by a
majority of the errantiviruses identified in this study. These elements form a separate clade, along with the
recently identified loki-retroviruses, which were also found to carry the same *env* type [17]. One possible
explanation for this is that vERVs undergo recombination events with other viruses. However, given that
the F-type vERVs form their own clade, it is unlikely that this happens frequently or recently, else they
would be more widespread across the phylogenetic tree. Another possibility is that the divergence of
vERVs and errantiviruses occurred after the acquisition of *env*, which would imply that they are equally
ancient, and may have a common evolutionary ancestor and origin.

Acknowledgements

We acknowledge Julius Brennecke, Kirsten Senti and Alex Greenwood for valuable comments in

preparation of the manuscript. This work is supported by the Australian Research Council (DP210102385).

Author contributions

S.C. and R.H. analysed and interpreted the data and wrote the manuscript. The idea of this work was

conceived by R.H., and S.C. and R.H. together made key findings.

Data Availability

The following data are available from the git repository

(https://github.com/RippeiHayashi/errantiviruses_ERVs): the code used to annotate and analyse genomic

copies of errantiviruses and vERVs, genomic sequences and open reading frames of the newly identified

ERVs, pdb files for the predicted structures of the POL bridge region, sequences of GAG, POL and ENV

open reading frames that are used for tBLASTn search, multiple sequence alignments and the tree files that

are generated in this study for the phylogenetic analyses.

Conflicts of interests

The authors declare no conflicts of interests.

Materials and methods

Genome assemblies used in the study

Reference genome sequences were retrieved from NCBI and grouped into the following insect orders, vertebrate classes and other metazoan phyla: Diptera, Lepidoptera, Coleoptera, Hymenoptera, Hemiptera, Polyneoptera, and Palaeoptera for insects; jawless-fish, fish, Amphibian, reptile, bird, and mammal for vertebrates; Arachnida, Branchiopoda, Chilopoda, Collembola, Diplopoda, Hexanauplia, Malacostraca, Merostomata, Ostracoda, and Thecostraca for Arthropoda; Acanthocephala, Annelida, Brachiopoda, Bryozoa, Dicyemida, Mollusca, Nemertea, Onychophora, Orthonectida, Phoronida, Platyhelminthes, Priapulida, Rotifera, and Tardigrada for Protostomia; Cephalochordata, Echinodermata, Hemichordata, and Tunicata for Deuterostomia; Cnidaria, Ctenophora, and Porifera. We obtained lists of reference genomes in each category in May 2023 using the Entrez Direct command line tool except for the vertebrate genomes which we used the list from the previous work by Wang et al. [17]. The list of the genome assemblies used can be found in Supplementary table S7.

Identification of active genomic copies of vertebrate endogenous retroviruses (vERVs) and *gypsy* errantiviruses

We ran two iterations of tBLASTn (ncbi-blast-2.9.0) to search for the genomic copies of vERVs and *gypsy* errantiviruses, firstly using the baits of known elements, and secondly using the baits identified from the first iteration. We used two sets of sequences for the first tBLAST search. The first set included GAG, POL and ENV protein sequences of the Dipteran *gypsy* errantiviruses from the RepBase (https://www.girinst.org/repbase/). The second set included GAG, POL and ENV protein sequences of selected vertebrate retroviruses that were used in the previous study [17]. We applied different criteria to identify ERVs from species outside insects and vertebrates as we did not have prior knowledge of whether or what kind of ERVs are found in those species. We briefly summarise our approach below.

We first used open reading frames (ORFs) of vertebrate retroviruses as baits for the vertebrate genomes and those of the Dipteran errantiviruses for the other genomes to run tBLASTn with default parameters. Genomic regions that showed homology to the POL baits with a bit score of greater than 50 or to the GAG and ENV baits with a bit score of greater than 30 were individually pooled. Genomic regions that had hits from multiple baits were merged for their coordinates using bedtools/2.28.0. Regions that were larger than 1500 nucleotides for POL and 300 nucleotides for GAG or ENV were collected. We then looked for regions that carry GAG, POL and ENV sequences that are next to each other in this order for insect genomes, and POL and ENV for the other genomes. We allowed up to 500 nucleotides of gaps between them for insect and vertebrate elements while we allowed up to 1000 nucleotides for the other genomes. For GAG-POL-ENV elements, we excluded regions of smaller than 4000 nucleotides or larger than 10000 nucleotides. For POL-ENV elements, we excluded regions of smaller than 3000 nucleotides or larger than 10000 nucleotides.

We subsequently checked whether the regions contain uninterrupted ORFs by making translated sequences

using transeq from EMBOSS-6.6.0 and excluding ORFs with more than five stop codons. We clustered all

the identified GAG, POL and ENV from each genome by Clustal Omega

(https://www.ebi.ac.uk/Tools/msa/clustalo/) and identified genomic regions that harbour all three ORFs

that are duplicated in the genome. These processes eliminated genomic copies of ERVs that are being

degraded.

The above process yielded abundant putative intact copies of ERVs from insect and vertebrate genomes

whereas we took additional steps for the other genomes. We extended the genomic regions of predicted

POL-ENV to upstream by 2000 nucleotides and manually annotated putative GAG ORFs using the online

sequence viewer Benchling (https://www.benchling.com/) for the Arthropoda genomes outside insects as

well as other Protostomia genomes. The identity of GAG ORFs were further confirmed by HHpred

(https://toolkit.tuebingen.mpg.de/tools/hhpred). For Cnidaria, Ctenophora and Deuterostomia genomes

outside vertebrates we used transeq to annotate putative GAG and ENV ORFs from 5000 nucleotides

downstream and 3000 nucleotides upstream to the predicted POL ORFs, respectively. Where possible only

the duplicated copies of ERVs were taken for further analyses outside insects and vertebrates as well.

We took confident 1128 POL ORFs from the first tBLASTn search and made a multiple sequence

alignment of the conserved regions of the Integrase domain using mafft-7.505 with the option '—auto',

and inferred the phylogenetic relationships using iqtree-1.6.12 with the option '-m rtREV+R4 -bb 1000'

(see more details in the section below). We took 178 representative POL from the phylogenetic tree and

used the POL and the corresponding GAG and ENV ORFs from the same genomic regions for a second

tBLASTn search. We repeated the above steps after combining the results of the two tBLASTn searches.

We did not find any ERVs in Nematoda genomes that have GAG, POL and ENV in this order. A previous

study reported non-canonical Nemetoda ERVs that have flipped the order of ENV and POL [33]. Thus, we

searched for regions that have ENV and POL in this order allowing gaps of up to 2000 nucleotides

between them. GAG was subsequently predicted from 5000 nucleotides upstream to the predicted ENV-

POL. We extended the search for potential GAG-ENV-POL insertions to other Protostomia genomes

outside Arthropoda and found similar elements in Bryozoa and Platyhelminthes. Furthermore, we found

that the glycoprotein gene present in mosquito *BEL/pao* retrotransposons [34] are also found next to *gypsy*

POL genes. The glycoprotein gene resembles the *envelope* genes from Chuviridae and is structurally

similar to glycoprotein B from Herpesviruses. We identified ten mosquito *BEL/pao* retrotransposons that

carry this glycoprotein from RepBase (BEL-199_AA, BEL-279_AA, BEL-652_AA, BEL-801_AA, BEL-

804_AA, BEL-879_AA, BEL-51_AnFu, BEL-56_AnFu, BEL-81_AnFu, and BEL-71_CQ) and used them

as baits for a tBLASTn search against Arthropoda genomes. The tBLASTn search identified 114 *gypsy*

elements with homologous glycoprotein genes from Coleoptera, Hymenoptera, Polyneoptera, and

Hexaneuplia species.

We ran Clustal Omega to select ORFs that are found multiple times in the genome for insects and

vertebrates while we focused on finding intact ORFs for the other genomes (see below) regardless of the

copy number. We additionally ran tBLASTn using the fully annotated ORFs as baits to interrogate the

copy number of ERVs in the genome where they are found. The results are summarised in Supplementary

table x where we counted the number of tBLASTn hits with a more than 90% identity to the bait

sequences. A majority of newly identified ERVs (1516 out of 1610 elements) carry ORFs that are found at

least twice in the genome (Table S1), further validating our approach of finding active or recently active

insertions.

Following, we further curated newly identified ERVs. Firstly, we used hmmer 3.3.1 (http://hmmer.org/) to

make sure that the ERVs that we identified are classified as Ty3/gypsy. We compared POL ORFs from the

newly identified ERVs against the POL ORFs of known retroelements that are available from the Gypsy

Database 2.0 (https://gydb.org/index.php/Main_Page). The Gypsy Database 2.0 collects hmm profiles of

Ty1/Copia, Bel/Pao, Ty3/Gypsy, and Retrovirus POL proteins. The hmmer search classified all the newly

identified ERVs into Ty3/Gypsy or Retroviruses. Secondly, we ran HHpred from hh-suite 3.3.0 [35] or the

online tool of HHpred (https://toolkit.tuebingen.mpg.de/tools/hhpred) to predict GAG, ENV, and POL

ORFs using the default search parameters. For the command line HHpred, we first made multiple sequence

alignments using hhblits and the Uniclust database UniRef30/2022_02 (https://uniclust.mmseqs.com/).

This was followed by hhsearch to predict domain structures using the collection of pdb70 from March

2022 (https://wwwuser.gwdg.de/~compbiol/data/hhsuite/databases/hhsuite_dbs/). We used default settings

to run hhblits and hhsearch. Furthermore, we reannotated ORFs in ERVs except for those from insects

when they are only partially predicted homology to baits in the tBLASTn search. We extended the

predicted ORFs in the genomic region to upstream until a start codon is found, and to downstream until a

stop codon is found. We used benchling to facilitate the identification of continuous and intact ORFs.

ERVs that predicted all GAG, ENV and POL ORFs were taken for further analyses.

Phylogenetic analysis of POL and ENV domains

We obtained multiple sequence alignments of the RT and Integrase domains of POL as follows. For the

RT domain, we first took the HHpred alignments of POL against the RT domains of Moloney Murine

Leukaemia Virus (Protein Data bank entry 4MH8_A) and Human Immunodeficiency Virus type 1 (Protein

Data bank entry 4G1Q_B), spanning regions that correspond to their fingers, palm, thumb and connection

subdomains [36], namely the amino acid positions of 28 to 426 and 7 to 405, respectively. However, this

process caused small truncations at either ends of the domain because of poor alignments at the individual

substructures. To overcome this issue and to facilitate sequence alignments, we extended the N- and C-

termini by 10 and 15 aa, respectively, after merging the two predicted regions of RT domains. We then ran

mafft-7.505 to generate a multiple sequence alignment, from which we manually trimmed the either ends

to have intact fingers, palm, thumb and connection subdomains for all POL ORFs. Similarly, we took

HHpred alignments of POL against the Integrase domains of Human Spumaretrovirus (Protein Data bank

entry 3OYM_A) and Simian T-lymphotropic Virus type 1 (Protein Data bank entry 7OUH_B), spanning

regions that correspond to their N-terminal domain (NTD) and catalytic core domain (CCD) [37], namely the

amino acid positions of 54 to 273 and 8 to 197, respectively. We used the predicted Integrase domains with

15 and 30 aa extensions at the N- and C-termini, respectively, for generating a multiple sequence

alignment before manually trimming the ends. We inspected the multiple sequence alignments using

Mview (https://www.ebi.ac.uk/Tools/msa/mview/) and removed ERVs that contained truncated RT or

Integrase domains from the analyses. We included ORFs from 5 Drosophila errantiviruses (DM176,

IDEFIX, ZAM, GYPSY, and Gypsy-5_DGri) and 23 representative vertebrate retroviruses (BLV, FLV,

HTLV-2, KoRV, LDV, MMTV, MuLV, REV, RSV, SRV-1, BFV, EFV, FFV, SFV, HIV-1, EIAV, FIV,

SnRV, Loki-Ofa, Loki-Pma, Loki-Sri, Loki-Str, and Loki-Xtr [17]) in the multiple sequence alignments.

We identified ERVs that carry three distinct types of ENV or Fusogen proteins, namely glycoprotein

F/Nipahvirus type (F-type), glycoprotein B from Herpes simplex viruses (HSV-type), and glycoprotein 2

type (GP2-type), and separately generated multiple sequence alignments. Briefly, we took HHpred

alignments of F-type ENV against metapneumovirus fusion protein (Protein Data bank entries 7SEM_F

and 5WB0_F), RSV F glycoprotein (6VKE_F and 6Q0S_F), parainfluenza virus fusion protein (4GIP_F

and 2B9B_B), Newcastle disease virus (1G5G_E and 3MAW_A), paramyxovirus (1ZTM_C), canine

Distemper virus (6XYE_B), and Nipah virus fusion glycoprotein (5EVM_D) and merged the predicted

regions. We subsequently ran mafft-7.505 to generate a multiple sequence alignment, from which we

manually removed ENVs with large truncations and trimmed sequences at either ends that are less

conserved. We then ran a second multiple sequence alignment only including the trimmed and conserved

regions of ENVs. We included F-type ENVs from SnRV and the above mentioned five Loki retroviruses in

the alignment. We took a similar approach for the HSV-type ENV. We took HHpred alignments of Human

herpesvirus 5 prefusion glycoprotein B (7KDP_B and 5C6T_A), Herpes simplex virus type I glycoprotein

B (5V2S_A), Human alphaherpesvirus 3 glycoprotein B (6VN1_C and 2GUM_C), and Suid

alphaherpesvirus 1 glycoprotein B, and merged the predicted regions. We then ran mafft-7.505 once to

generate a multiple sequence alignment, which we further trimmed either ends to obtain alignments from a

highly conserved region. HHpred found homology to Syncytin-2 (7OIX_A) for all the GP2-type ENV

ORFs that we identified. Some GP2-type ENV ORFs additionally showed homology to the N-terminal

receptor binding domain of Feline Leukemia Virus glycoprotein (1LCS_B), Murine Leukemia virus

glycoprotein 70 (1AOL_A), and the Human endogenous retrovirus EnvP(b)1 (6W5Y_B). We first aligned

the full length ORFs of the GP2-type ENVs, and trimmed either ends to have C-terminal regions that are

homologous to Syncytin-2. ENVs with large truncations are removed. We then performed a second

alignment using a set of intact and trimmed ENV ORFs. Among the above mentioned ORFs from the

representative vertebrate retroviruses, we included BLV, FLV, HTLV-2, KoRV, LDV, MMTV, MuLV,

REV, RSV and SRV-1 in the alignment of the GP2-type ENVs.

1610 ERVs from all phyla combined retained intact Integrase and RT domains in POL and one of the three

types of ERVs, and thus were used for further analyses. We also included the above mentioned 5

Drosophila errantiviruses and 23 vertebrate retroviruses for phylogenetic analyses. We ran iqtree-1.6.12

with the option '-m rtREV+R4 -bb 1000' to generate the phylogenetic trees of all Integrase and RT

domains, those from ERVs with each of the three types of ENVs, and the trees of each ENV type.

Clades are then defined mainly based on how closely they cluster together on the tree, as well as the

relative distance from neighbouring clades. In some cases, multiple clades are combined based on their

relative position to each other from the root. Clade definitions are summarised in Supplementary Table S3

and represented in Figure 1. We then utilize these clade definitions to analyse congruency between

Integrase and host phylogeny, Integrase and Env, and Integrase and RT, (Table S3-5). We also use these

clade definitions to select random representative ERVs for structural prediction of the RNase H domain

using Alphafold 2, discussed later (Table S6).

For our analysis on host phylogeny, we separately define clades that contain ERVs found in Arthropoda or

Vertebrata using a similar approach, but this time excluding clades that do not contain either. We labelled

these clades A1-A24 for Arthropoda, and V1-V2 for vertebrates (Table S4). We then constructed a balloon

plot of clade against host species using the ggballoon plot package in R. The size of each data point was

made proportional to species representation. For example, a balloon size of 0.5 for an order in one clade,

would mean that 50% of the elements found in that order are found in that given clade.

For our analysis on Integrase and RT, as well as Integrase and Env, we constructed balloon plots in a

similar manner, but the size of the balloon is simply proportional to the total number of elements

represented in the point, and not as a proportion.

<u>Identification of putative long terminal repeats and transfer RNA (tRNA) primer binding sequences</u>

Long terminal repeats (LTRs) of ERVs were identified as follows. Firstly, we fetched the genomic

coordinates of ERV insertions and extended 3000 nucleotides at either ends from the Start codon of GAG

and the Stop codon of ENV ORFs (POL was taken for the GAG-ENV-POL ERVs) for all species

categories except for Cnidaria, Ctenophora and Deuterostomia where we extended 5000 nucleotides at

either ends from the Start and Stop codons of POL. We retrieved the genomic sequence of the extended

ERVs and ran BLASTn against the same sequence to identify repeated sequences. We considered repeated

regions of greater than 99% identity and the BLASTn bit score between 200 and 5000 as putative LTRs.

We excluded LTRs whose distance between them is shorter than 5000 nucleotides or longer than 14000

nucleotides. Following, we searched for putative tRNA primer binding sequences (PBS). tRNA sequences

were retrieved from genomic annotations from representative genomes in respective species categories

except for Ctenophora where we took tRNA sequences from Protostomia, Cnidaria and Deuterostomia

because there were no annotated genomes available at the time of the study. Representative genomes that

we used to retrieve tRNA sequences are: GCF_000001215.4 and GCF_013141755.1 for Diptera,

GCF_014905235.1 and GCF_905147365.1 for Lepidoptera, GCF_000002335.3 and GCF_917563875.1

for Coleoptera, GCF_003254395.2 and GCF_016802725.1 for Hymenoptera, GCF_014356525.2 and

GCF_020184175.1 for Hemiptera, GCF_002891405.2 and GCF_021461395.2 for Polyneoptera,

GCF_921293095.1 for Palaeoptera, GCF_000671375.1, GCF_020631705.1, GCF_002217175.1,

GCF_016086655.3, GCF_024679095.1, GCF_019059575.1 for Arthropoda, GCF_000326865.2,

GCF_020536995.1, GCF_000002985.6, GCF_000715545.1 for Protostomia, GCF_000222465.1 and

GCF_022113875.1 for Cnidaria, GCF_013122585.1, GCF_000003605.2, GCF_000003815.2, and

GCF_000002235.5 for Deuterostomia, and GCF_015237465.2, GCF_000001405.40, GCF_017654675.1,

GCF_900496995.4, GCF_900700375.1, and GCF_010993605.1 for Vertebrata. The minimum

requirements of the priming events are incompletely understood, but the length of PBS for

retrotransposons is generally between 8 and 18 nucleotides and the PBS is positioned near the 3' end of the

5' LTR [38]. We chose the 12 nucleotides at the 3' end of tRNAs as putative primer sequences and searched

for PBS in the 100 nucleotide-window centred at the 3' end of the 5' LTR using bowtie-1.2.3 allowing up

to two mismatches. We added "CCA" to the 3' end of the tRNA sequence when it is not present in the

genomic sequences. We successfully identified putative PBSs in 1205 out of 1610 ERVs that we identified

in this study. We grouped ERVs based on the type of tRNA used for the PBS and summarised the results

in Supplementary figure S1 and Supplementary table S2.

<u>Structure prediction of the RNase H domains using Alphafold 2</u>

Using clade definitions from the phylogenetic trees of Integrase and RT, we selected representative ERVs

from each clade for the structural prediction of the RNase H domain. We used the colabfold wrap v1.4.0 of

Alphafold 2 (https://github.com/YoshitakaMo/localcolabfold/releases/tag/v1.4.0), and used MMSeqs2 for

the database search and clustering of protein sequences before performing the structural prediction [39]. We

then visualised the RNase H domains using Maestro from Schrödinger

(https://newsite.schrodinger.com/platform/products/maestro/), and characterised visible structures by the

orientation and order of β sheets present, grouping them by letter (Table S6). Once the first set of proteins

were analysed, elements with unique structures were further analysed by characterising RNase H domains

from the remaining ERVs in that clade.

Figure Legends

Figure 1 – Phylogenetic relationships between *gypsy* errantiviruses and vertebrate endogenous retroviruses (vERVs). (A) Maximum likelihood tree constructed based on *pol* Integrase proteins of *gyspy* errantiviruses and vERVs, with host phyla annotated in colour. (B) Distribution of vERVs/*gypsy* errantiviruses in metazoa with *env* type indicated.

Figure 2 – Representation of *env* type in phylogenetic tree of Integrase in *gypsy* errantiviruses and vERVs. (A) Maximum likelihood tree constructed based on *pol* Integrase proteins of *gyspy* errantiviruses and vERVs, with Env type annotated in colour. (B-D) Close up view of Integrase clades N, O and P respectively. Host species for sub-branches within clade annotated, with *env* type annotated for mixed sub-branches.

Figure 3 – Domain architecture and 3D structure of Ribonuclease H (RNase H) in *gypsy* errantiviruses and vERVs. (A) Maximum likelihood tree constructed based on *pol* Integrase proteins of *gyspy* errantiviruses and vERVs with RNase H domain architecture annotated in colour. V – Vertebrates, I – Insects, AA – Ancient Animals. Integrase clades N, O and P are also shown in grey. (B) Summary of bridge region RNase H domains found in representative elements in Insects, Ancient Animals and Vertebrates. (C) Alphafold-predicted 3D structure of canonical RNase H/Tether, 'mini' RNase H and 'pseudo' RNase H domains, with topology maps of α helices (grey) and β sheets (red). (D) Distribution of shared RNase H domain architectures in the bridge region of elements within insects, vertebrates, and ancient animals. Structures for Alphafold taken from Annelida_Polychaeta_Apac_errantivirus_7, Bryozoa_Gymnolaemata_Cpal_errantivirus_3 and Coleoptera_Coccinellidae_Npum_errantivirus_1 for RNase H/Tether, 'mini' RNase H and 'pseudo' RNase H domains respectively.

Supplementary Figure S1 – Bootstrap values and tRNA primer binding sites of *gypsy* errantiviruses and vERVs. (A) Maximum likelihood tree constructed using *pol* Integrase proteins. Red triangles represent branches/nodes with bootstrap values >95. The tree and the clade definition are the same as in Fig 1A. (B) Type of tRNAs carried by *gypsy* errantiviruses and vERVs in discrete clades from the tree.

Supplementary Figure S2 – Phylogenetic congruence between Integrase and RT proteins of vERVs and *gypsy* errantiviruses. (A) Maximum likelihood tree constructed based on Pol RT proteins of *gypsy* errantiviruses and vERVs. Bootstrap values >95 annotated on nodes with red triangles. (B) Balloon plot of number of elements shared between clades in Integrase and RT. Size of balloon is proportional to number of elements shared between discrete clades of Integrase and RT.

Supplementary Figure S3 – Phylogenetic relationships between *gyspy* errantiviruses/vERVs and their hosts. (A) Arthropoda (A1-A24) and Vertebrate (V1-V12) specific clades on maximum likelihood tree constructed based on Integrase proteins. The same tree was used in Figure 1A. (B) Balloon plot representing Arthropoda/Vertebrate clade to order congruence. Size of balloon represented in proportion of

total elements from an order, where 1 represents a balloon that contains all the elements from that order.

(C-D) Close ups of clades O and P from Figure 1A. (E) Expanded and detailed colour code for host

phylum annotation in maximum likelihood trees.

Supplementary Figure S4 – Phylogenetic congruence between *pol* Integrase and Env for *gypsy*

errantiviruses and vERVs. Trees for Integrase and Env with clade definitions matching the annotation in

the balloon plot of number of elements shared between clades in Integrase and Env for elements containing

type F (A), GP2 (B) and HSV (C). Size of balloon is proportional to number of elements shared between

discrete clades of Integrase and Env. (A') Maximum likelihood tree constructed on *int* proteins of elements

carrying EnvF. (A") Maximum likelihood tree constructed on Env type F. (A'") Balloon plot of Integrase

vs Env for F type elements. (B') Maximum likelihood tree constructed on Integrase proteins of elements

carrying Env HSV. (B") Maximum likelihood tree constructed on Env type HSV. (B") Balloon plot of

Integrase vs Env for HSV type elements. (C') Maximum likelihood tree constructed on *int* proteins of

elements carrying Env GP2. (C") Maximum likelihood tree constructed on Env type GP2. (C'") Balloon

plot of Integrase vs Env for GP2 type elements.

Supplementary Figure S5 – 3D structure of uncommon and yeast RNase H-like domains in *gypsy*

errantiviruses. (A) Alphafold-predicted 3D structure of uncommonly found RNase H-like domains with

topology maps of α helices (grey) and β sheets (red). (B) 3D structures and topology maps of 'mini' RNase

H domains in yeast Ty3/*gypsy* elements. Structures for Alphafold taken from

Annelida_Polychaeta_Slim_errantivirus_9, Diptera_Cecidomyiidae_Smos_errantivirus_1 and

Hymenoptera_Apoidea_Obic_errantivirus_2 for "Mini" domain 2, "Pseudo" domain 2 and "Pseudo"

domain 3 respectively.

Supplementary table S1 – Master *gypsy* errantivirus and vERV table. Information on name, host

classification, genome assembly, genomic coordinates, *env* type, type of tRNA used as primer, LTR

coordinates and the number of mismatches between annotated tRNA sequences and the tRNA primer

binding sequence in tab 1. Copy number of *gypsy* errantivirus and vERV open reading frames in tab 2,

tRNA classification guide in tab 3.

Supplementary table S2 – Raw data of Integrase clade definition against tRNA primer classification for

supplementary figure S1B.

Supplementary table S3 – Clade definitions of Integrase and RT maximum likelihood trees. Clade

definitions for Integrase in tab 1. Clade definitions for RT in tab 2. Raw balloon plot data for

supplementary figure S2B of Integrase vs RT in tab 3.

Supplementary table S4 – Raw data for supplementary figure S3. Arthropoda clade definitions for

Integrase in tab 1. Raw balloon plot table of host order/class against Arthropoda clade in tab 2, with total

number of elements from order/class in tree given. Vertebrate clade definitions for Integrase in tab 3. Raw

balloon plot table of host order/class against Vertebrate clade in tab 4, with total number of elements from
order/class in tree given.

Supplementary table S5 – Raw data for supplementary figure 4. Clade definitions for Integrase carrying F
type Env in tab 1. Clade definitions for Env F in tab 2. Raw balloon plot of Integrase vs Env for F carrying
ERVs in tab 3. Clade definitions for Integrase carrying HSV type Env in tab 4. Clade definitions for Env
HSV in tab 5. Raw balloon plot of Integrase vs Env for HSV carrying ERVs in tab 6. Clade definitions for
Integrase carrying GP2 type Env in tab 7. Clade definitions for Env GP2 in tab 8. Raw balloon plot of
Integrase vs Env for GP2 carrying ERVs in tab 9.

Supplementary table S6 – RNase H classification based on arrangement of β sheets for representative
elements.

Supplementary table S7 – Total list and details of analysed genomes. Diptera genomes in tab 1.
Lepidoptera genomes in tab 2. Coleoptera genomes in tab 3. Hymenoptera genomes in tab 4. Hemiptera
genomes in tab 5. Polyneoptera genomes in tab 6. Palaeoptera genomes in tab 7. Non-insect Arthropoda
genomes in tab 8. Protostomia genomes excluding Arthropoda and Nematoda in tab 9. Nematoda genomes
in tab 10. Cnidaria genomes in tab 11. Ctenophora genomes in tab 12. Porifera genomes in tab 13.
Deuterostomia genomes excluding vertebrates in tab 14. Vertebrate genomes in tab 15. Numbers of
predicted ORFs are shown. ORFs that are found downstream of predicted POL were also counted for
Protostomia, Cnidaria, Ctenophora, Porifera and Deuterostomia. Genomes that we used to annotate vERVs
and errantiviruses are indicated.

References

1   Johnson, W. E. Endogenous Retroviruses in the Genomics Era. *Annu Rev Virol* **2**, 135-159
(2015). https://doi.org:10.1146/annurev-virology-100114-054945
2   Feschotte, C. & Gilbert, C. Endogenous viruses: insights into viral evolution and impact on
host biology. *Nat Rev Genet* **13**, 283-296 (2012). https://doi.org:10.1038/nrg3199
3   Zhuo, X., Rho, M. & Feschotte, C. Genome-wide characterization of endogenous retroviruses
in the bat Myotis lucifugus reveals recent and diverse infections. *J Virol* **87**, 8493-8501
(2013). https://doi.org:10.1128/JVI.00892-13
4   Metegnier, G. *et al.* Comparative paleovirological analysis of crustaceans identifies multiple
widespread viral groups. *Mob DNA* **6**, 16 (2015). https://doi.org:10.1186/s13100-015-0047-3
5   Dewannieux, M. & Heidmann, T. Endogenous retroviruses: acquisition, amplification and
taming of genome invaders. *Curr Opin Virol* **3**, 646-656 (2013).
https://doi.org:10.1016/j.coviro.2013.08.005
6   Coffin, J. M., Hughes, S. H. & Varmus, H. E.   (1997).
7   Pelisson, A., Mejlumian, L., Robert, V., Terzian, C. & Bucheton, A. Drosophila germline
invasion by the endogenous retrovirus gypsy: involvement of the viral env gene. *Insect*
Biochem Mol Biol **32**, 1249-1256 (2002). https://doi.org:10.1016/s0965-1748(02)00088-7
8   Keegan, R. M., Talbot, L. R., Chang, Y. H., Metzger, M. J. & Dubnau, J. Intercellular viral
spread and intracellular transposition of Drosophila gypsy. *PLoS Genet* **17**, e1009535 (2021).
https://doi.org:10.1371/journal.pgen.1009535
9   Song, S. U., Gerasimova, T., Kurkulos, M., Boeke, J. D. & Corces, V. G. An env-like protein
encoded by a Drosophila retroelement: evidence that gypsy is an infectious retrovirus.
Genes Dev **8**, 2046-2057 (1994). https://doi.org:10.1101/gad.8.17.2046
10   Brasset, E. *et al.* Viral particles of the endogenous retrovirus ZAM from Drosophila
melanogaster use a pre-existing endosome/exosome pathway for transfer to the oocyte.
Retrovirology **3**, 25 (2006). https://doi.org:10.1186/1742-4690-3-25
11   Malik, H. S., Henikoff, S. & Eickbush, T. H. Poised for contagion: evolutionary origins of the
infectious abilities of invertebrate retroviruses. *Genome Res* **10**, 1307-1318 (2000).
https://doi.org:10.1101/gr.145000
12   Rohrmann, G. F. & Karplus, P. A. Relatedness of baculovirus and gypsy retrotransposon
envelope proteins. *BMC Evol Biol* **1**, 1 (2001). https://doi.org:10.1186/1471-2148-1-1
13   Hayward, A., Cornwallis, C. K. & Jern, P. Pan-vertebrate comparative genomics unmasks
retrovirus macroevolution. *Proc Natl Acad Sci U S A* **112**, 464-469 (2015).
https://doi.org:10.1073/pnas.1414980112
14   Wang, J. & Han, G. Z. Genome mining shows that retroviruses are pervasively invading
vertebrate genomes. *Nat Commun* **14**, 4968 (2023). https://doi.org:10.1038/s41467-023-
40732-w
15   Stefanov, Y., Salenko, V. & Glukhov, I. Drosophila errantiviruses. *Mob Genet Elements* **2**, 36-
45 (2012). https://doi.org:10.4161/mge.19234
16   Terzian, C., Pélisson, A. & Bucheton, A. Evolution and phylogeny of insect endogenous
retroviruses. *BMC Evol Biol* **1**, 3 (2001). https://doi.org:10.1186/1471-2148-1-3
17   Wang, J. & Han, G. Z. A Sister Lineage of Sampled Retroviruses Corroborates the Complex
Evolution of Retroviruses. *Mol Biol Evol* **38**, 1031-1039 (2021).
https://doi.org:10.1093/molbev/msaa272
18   Gifford, R., Kabat, P., Martin, J., Lynch, C. & Tristem, M. Evolution and distribution of class II-
related endogenous retroviruses. *J Virol* **79**, 6478-6486 (2005).
https://doi.org:10.1128/JVI.79.10.6478-6486.2005
19   Bargues, N. & Lerat, E. Evolutionary history of LTR-retrotransposons among 20 Drosophila
species. *Mob DNA* **8**, 7 (2017). https://doi.org:10.1186/s13100-017-0090-3

20 Malik, H. S. & Eickbush, T. H. Phylogenetic analysis of ribonuclease H domains suggests a
late, chimeric origin of LTR retrotransposable elements and retroviruses. *Genome Res* **11**,
1187-1197 (2001). https://doi.org:10.1101/gr.185101
21 Loreto, E. L., Carareto, C. M. & Capy, P. Revisiting horizontal transfer of transposable
elements in Drosophila. *Heredity (Edinb)* **100**, 545-554 (2008).
https://doi.org:10.1038/sj.hdy.6801094
22 Metzger, M. J., Paynter, A. N., Siddall, M. E. & Goff, S. P. Horizontal transfer of
retrotransposons between bivalves and other aquatic species of multiple phyla. *Proc Natl
Acad Sci U S A* **115**, E4227-E4235 (2018). https://doi.org:10.1073/pnas.1717227115
23 Grandi, N. & Tramontano, E. Human Endogenous Retroviruses Are Ancient Acquired
Elements Still Shaping Innate Immune Responses. *Front Immunol* **9**, 2039 (2018).
https://doi.org:10.3389/fimmu.2018.02039
24 Johnson, W. E. Origins and evolutionary consequences of ancient endogenous retroviruses.
Nat Rev Microbiol **17**, 355-370 (2019). https://doi.org:10.1038/s41579-019-0189-2
25 Louz, D., Bergmans, H. E., Loos, B. P. & Hoeben, R. C. Cross-species transfer of viruses:
implications for the use of viral vectors in biomedical research, gene therapy and as live-
virus vaccines. *J Gene Med* **7**, 1263-1274 (2005). https://doi.org:10.1002/jgm.794
26 Subbarao, K. The Critical Interspecies Transmission Barrier at the Animal⁻Human Interface.
Trop Med Infect Dis **4** (2019). https://doi.org:10.3390/tropicalmed4020072
27 Zheng, H. *et al.* Koala retrovirus diversity, transmissibility, and disease associations.
Retrovirology **17**, 34 (2020). https://doi.org:10.1186/s12977-020-00541-1
28 Mottaghinia, S. *et al.* A recent gibbon ape leukemia virus germline integration in a rodent
from New Guinea. *Proc Natl Acad Sci U S A* **121**, e2220392121 (2024).
https://doi.org:10.1073/pnas.2220392121
29 Tarlinton, R. E., Meers, J. & Young, P. R. Retroviral invasion of the koala genome. *Nature* **442**,
79-81 (2006). https://doi.org:10.1038/nature04841
30 Zheng, J., Wei, Y. & Han, G. Z. The diversity and evolution of retroviruses: Perspectives from
viral "fossils". *Virol Sin* **37**, 11-18 (2022). https://doi.org:10.1016/j.virs.2022.01.019
31 Brennecke, J. *et al.* Discrete small RNA-generating loci as master regulators of transposon
activity in Drosophila. *Cell* **128**, 1089-1103 (2007). https://doi.org:10.1016/j.cell.2007.01.043
32 Ribet, D. *et al.* An infectious progenitor for the murine IAP retrotransposon: emergence of
an intracellular genetic parasite from an ancient retrovirus. *Genome Res* **18**, 597-609 (2008).
https://doi.org:10.1101/gr.073486.107
33 Sacco, M. A., Lau, J., Godinez-Vidal, D. & Kaloshian, I. Non-canonical nematode endogenous
retroviruses resulting from RNA virus glycoprotein gene capture by a metavirus. *J Gen Virol*
103 (2022). https://doi.org:10.1099/jgv.0.001739
34 Dezordi, F. Z., Vasconcelos, C. R. D. S., Rezende, A. M. & Wallau, G. L. In and Outs of
Chuviridae Endogenous Viral Elements: Origin of a Potentially New Retrovirus and Signature
of Ancient and Ongoing Arms Race in Mosquito Genomes. *Front Genet* **11**, 542437 (2020).
https://doi.org:10.3389/fgene.2020.542437
35 Steinegger, M. *et al.* HH-suite3 for fast remote homology detection and deep protein
annotation. *BMC Bioinformatics* **20**, 473 (2019). https://doi.org:10.1186/s12859-019-3019-7
36 Das, D. & Georgiadis, M. M. The crystal structure of the monomeric reverse transcriptase
from Moloney murine leukemia virus. *Structure* **12**, 819-829 (2004).
https://doi.org:10.1016/j.str.2004.02.032
37 Engelman, A. & Cherepanov, P. Retroviral Integrase Structure and DNA Recombination
Mechanism. *Microbiol Spectr* **2**, 1-22 (2014). https://doi.org:10.1128/microbiolspec.MDNA3-
0024-2014.
38 Wilhelm, M. & Wilhelm, F. X. Reverse transcription of retroviruses and LTR retrotransposons.
Cell Mol Life Sci **58**, 1246-1262 (2001). https://doi.org:10.1007/PL00000937

39   Mirdita, M. *et al.* ColabFold: making protein folding accessible to all. *Nat Methods* **19**, 679-
682 (2022). https://doi.org:10.1038/s41592-022-01488-1

Figure 1

Figure 1A. Unrooted phylogenetic tree with clades labeled A–V. Inset legend showing color coding for INSECTS (Diptera, Lepidoptera, Coleoptera, Hymenoptera, Hemiptera, Polyneoptera, Palaeoptera), ARTHROPODS (Collembola, Hexanauplia, Malacostraca, Thecostraca, Branchiopoda, Arachnida), PROTOSTOMIA (Nematoda, Mollusca, Bryozoa, Annelida, Nemertea, Platyhelminthes), ANCIENT ANIMALS (Hemichordata, Tunicata, Ctenophora, Cnidaria), VERTEBRATES (Jawless Fish, Fish, Amphibian, Reptile, Bird, Mammal). Tree scale: 1.

B

	GAG-POL-ENV	Env Type
Platyhelminthes	Yes	HSV[1]
Nemertea	Yes	F
Annelida	Yes	F
Bryozoa	Yes	HSV[1]
Mollusca	Yes	F
Nematoda	Yes	HSV[1]
Arthropoda*	Yes	F, HSV
Vertebrata**	Yes	GP2, F
Tunicata	Yes	F, HSV
Hemichordata	Yes	F
Echinodermata	No	-
Cnidaria	Yes	F, HSV
Ctenophora	Yes	F, HSV
Porifera	No	-

Protostomia (Lophotrochozoa, Ecdysozoa); Deuterostomia (Chordata); Coelenterata, Diploblasta.

1 - Orientation of ORFs is GAG-ENV-POL

	GAG-POL-ENV	Env Type
Insecta***	Yes	F, HSV
Collembola	Yes	F
Hexanauplia	Yes	F, HSV
Malacostraca	Yes	F
Thecostraca	Yes	F
Branchiopoda	Yes	F
Arachnida	Yes	F
Mammals	Yes	GP2
Birds	Yes	GP2
Reptiles	Yes	GP2, F
Amphibians	Yes	GP2, F
Fish	Yes	GP2, F
Jawless Fish	Yes	F
Diptera	Yes	F
Lepidoptera	Yes	F
Coleoptera	Yes	F, HSV
Hymenoptera	Yes	F, HSV
Hemiptera	Yes	F, HSV
Polyneoptera	Yes	F, HSV
Palaeoptera	Yes	F

Arthropoda (*); Vertebrata (**); Insecta (***).

Figure 2

A

P

O

N

F
HSV
GP2

Tree scale: 1

B

Lepidoptera, Coleoptera,
Hymenoptera, Hemiptera,
Polyneoptera,
Palaeoptera

Hemiptera   Hymenoptera

Hymenoptera

Clade N   Tree scale: 1

C

Nematoda*, Platyhelminthes*,
Arachnida**, Branchiopoda**,
Coleoptera*, Polyneoptera***
Palaeoptera*

Coleoptera,
Collembola,
Branchiopoda

Cnidaria

Bryozoa

*   HSV type env only
**   F type env only
***   F and HSV type env

Clade O   Tree scale: 1

D

Hemiptera

Nematoda

Diptera**, Collembola**,
Annelida**, Nematoda*,
Ctenophora**

Bryozoa*, Thecostraca**,
Nematoda*

*   HSV type env only
**   F type env only

Clade P   Tree scale: 1

Figure 3

A

Integrase clade A

I

I+AA

I

V

V

AA

Integrase clade P

Integrase clade O

I+AA

Integrase clade N

AA

Tree scale: 1

B

Vertebrates

RNase H/Tether + RNase H/Tether + "Mini" domain
RNase H/Tether + "Mini" domain
RNase H/Tether + RNase H/Tether + "Mini" domain 2
RNase H/Tether + RNase H/Tether

Insects

RNase H/Tether + "Pseudo" RNase H
RNase H/Tether + "Pseudo" RNase H + "Pseudo" RNase H
RNase H/Tether + "Mini" domain 2
RNase H/Tether + "Pseudo" domain 2
RNase H/Tether + "Mini" domain
RNase H/Tether + "Pseudo" domain 3
RNase H/Tether

Ancient Animals

RNase H/Tether + "Mini" domain
RNase H/Tether + "Mini" domain 2
RNase H/Tether + "Pseudo" RNase H

C

B'   B''   B'''

α
5   α 4   α 1   2   3

1   2   3
α
α
α

5   α 4   α 1   3   2

RNaseH/Tether   "Mini" Domain   "Pseudo" RNaseH

D

Vertebrates

Ancient Animals

RNaseH
+
Tether
+
"Mini"

RNaseH
+
Tether

Insects

RNaseH
+
"Pseudo"

RNaseH
+
"Mini"

Supplementary Figure S1

Supplementary Figure S2

Supplementary Figure S3

Supplementary Figure S4

Supplementary Figure S5

A

A' "Mini" Domain 2

A'' "Pseudo" Domain 2

A''' "Pseudo" Domain 3

B

B' 7Q5B

B'' Pombe tf1

B''' Pombe tf2