Activity in the reticulospinal tract scales with handgrip force in humans

Tyler L. Danielson[1], Layla Gould[2], Jason M. DeFreitas[1], Rob J. MacLennan[3,4], Chelsea Ekstrand[5], Ron
Borowsky[6], Jonathan P. Farthing[7], Justin W. Andrushko[8]

1. Applied Neuromuscular Physiology Laboratory, College of Education and Human Sciences,
Oklahoma State University, Stillwater, Oklahoma, USA
2. Department of Surgery, Division of Neurosurgery, College of Medicine, University of
Saskatchewan, Saskatoon, Saskatchewan, Canada
3. Department of Neurology, College of Medicine, University of Florida, Gainesville, Florida,
USA
4. Brain Rehabilitation Research Center, Malcom Randall VA Medical Center, Gainesville,
Florida, USA
5. Department of Neuroscience, Faculty of Arts and Science, University of Lethbridge,
Lethbridge, Alberta, Canada
6. Department of Psychology and Health Studies, College of Arts and Science, University of
Saskatchewan, Saskatchewan, Canada
7. College of Kinesiology, University of Saskatchewan, Saskatoon, Saskatchewan, Canada
8. Department of Sport, Exercise and Rehabilitation, Faculty of Health and Life Sciences,
Northumbria University, Newcastle upon Tyne, Tyne and Wear, United Kingdom

Corresponding Author

Justin W. Andrushko, Ph.D.
Justin.andrushko@northumbria.ac.uk
ORCiD: https://orcid.org/0000-0003-2258-1689

Co-corresponding Author

Jonathan P. Farthing, Ph.D.
jon.farthing@usask.ca
ORCiD: https://orcid.org/0000-0002-2549-1227

Keywords

Descending motor pathways, reticular formation, Pons, motor control, force modulation, strength,
fMRI, brain imaging

Abstract

Introduction: The neural pathways that contribute to force production in humans are currently poorly understood, as the relative roles of the corticospinal tract and brainstem pathways, such as the reticulospinal tract (RST), vary substantially across species. Using functional magnetic resonance imaging (fMRI) we aimed to measure activation in the pontine reticular nuclei during different submaximal handgrip contractions to determine the potential role of the RST in force modulation. **Methods:** Thirteen neurologically intact participants (age: 28 ± 6 years) performed unilateral handgrip contractions at 25%, 50%, 75% of maximum voluntary contraction during brain scans. We quantified the magnitude of RST activation from the contralateral and ipsilateral sides in addition to the contralateral primary motor cortex during each of the three contraction intensities. **Results:** A repeated measures ANOVA demonstrated a significant main effect of force ($p = 0.012$, $\eta_p^2 = 0.307$) for RST activation, independent of side (i.e., activation increased with force for both contralateral and ipsilateral nuclei). Further analyses of these data involved calculating the linear slope between the magnitude of activation and handgrip force for each ROI at the individual-level. One-sample t-tests on the slopes revealed significant group-level scaling for the RST bilaterally, but only the ipsilateral RST remained significant after correcting for multiple comparisons. **Conclusions:** Here, we show evidence of task dependent activation in the RST pontine nuclei that was positively related to handgrip force. These data build on a growing body of literature that highlights the RST as a functionally relevant motor pathway for force modulation in humans.

New & Noteworthy

In this short report, we used a task-based fMRI paradigm to show that activity in the reticulospinal tract, but not the contralateral motor cortex, scales linearly with increasing force during a handgrip task. These findings directly support recently proposed hypotheses that the reticulospinal tract may play an important role in modulating force production in humans.

Introduction

Within the central nervous system, there are multiple descending pathways that control motor

function, and the roles and relative contributions from each pathway may vary substantially across

species. In non-mammals, such as reptiles and birds, the main motor pathways descending onto

motoneurons originate in the brainstem (1). However, in humans, it is believed that upper limb fine

motor control and force modulation is primarily controlled through the corticospinal tract [CST; (2,

3)]; a descending pathway originating from the primary motor cortex (M1). An added difficulty in

quantifying the role of the CST is how much its size and functionality differs between species. Within

humans, the cortico-motoneuronal connections are substantially more developed than in non-

human primates and other mammals (4), suggesting that the role of CST in humans likely cannot be

inferred accurately from animal-model research. Increased development of the CST correlates

positively with indices of dexterity (5), giving further evidence for the role of the CST in fine, distal

motor production. Due to this specialization of the CST within humans, most research studies

investigating the mechanisms of human motor control and muscle force have primarily focused on

the CST, with equivocal and heterogeneous findings (6, 7).

Recent evidence, however, now suggests the reticulospinal tract (RST); a brainstem pathway thought

to be limited in humans to postural and proximal muscle control, may be more involved in upper

limb motor control than previously thought (8–10). In support of this, there has been evidence of

distal RST innervation in non-human primate models (11) and in humans (8, 9). In addition to its

contribution to gross movement control, recent papers have suggested that the RST may also have a

role in force modulation (12–14). Some studies have used animal models to study the RST's role in

voluntary force production, but these models require the implantation of electrodes on the RST

which is not possible in humans (10, 11, 15–17). For example, Glover and Baker (16) found that the

RST appears to influence contraction force in macaque monkeys. Others have examined this control

in humans with stroke and spinal cord injury, allowing for models analogous to animal lesion studies

(18, 19).

In humans, these studies primarily involve non-invasive stimulation to produce evoked contractions

(19–21), while animal models have allowed for studies during voluntary movement (11, 15–17, 22).

Examining the role of the RST during voluntary force production is important to elucidate how it

contributes to the control of human movement. To our knowledge, no study to date has used

functional magnetic resonance imaging (fMRI) to corroborate the role of the RST for controlling force

in humans despite the benefits of its non-invasive nature. The purpose of this study was to quantify

whether the RST is more active during handgrip contractions at higher compared to lower intensities.

Materials and Methods

As a secondary analysis on a data set that has previously been published (23) we aimed to determine

the relationship between fMRI BOLD signal modulation in the pontine reticular formation nuclei and

parametric increases in handgrip force during three separate fMRI runs (Figure 1).

Figure 1. A) Handgrip contractions at three different force levels were performed during
three separate functional magnetic resonance imaging (fMRI) runs of the cerebrum and
brainstem. The monitor in the top right shows our primary region of interest (ROI) in MNI-
space, the pontine nuclei of the reticulospinal tract (RST). Section **B)** depicts a simplified
diagram of the hypothesized pathways involved in voluntary handgrip force production. For
comparison, activity levels in the contralateral motor cortex (M1) were also assessed. Not
shown, but also potential contributors, are the contralateral supplemental motor area (SMA)
and its bilateral projections to the RST on each side, potential connections to the RST from
the premotor cortices, and uncrossed pathways from the ipsilateral M1.

Participants

Thirteen neurologically intact individuals volunteered to participate in this study (11 right-handed, 2

left-handed; sex: 7 female, 6 male; age: 28 ± 6 years; height: 170.9 ± 9.8 cm; weight: 75.1 ± 16.7 kg).

Participants were eligible if they were 18 years or older and were MRI safe. The presence of

intracranial metal clips, cardiac pacemaker, or any factor that would prevent a safe MRI scan

disqualified individuals to participate.

Ethical approval

112 All participants provided written informed consent before participation. The study adhered to the

113 Declaration of Helsinki and received approval from the University of Saskatchewan Biomedical

114 Research Ethics Board (Ethics # Bio 01-125).

Experimental outline

During a single fMRI session, participants engaged in three experimental tasks. These tasks were

executed in a random sequence and consisted of submaximal unimanual right isometric handgrip

contractions at 25%, 50%, and 75% of their maximum voluntary contraction (MVC). The session

began with the acquisition of a structural MRI brain scan, followed by having the participants

perform three MVC's. To prevent fatigue, a one-minute rest interval was allotted between each MVC

attempt. Based on the MVC results, target force levels were set for the three submaximal

experimental tasks. Participants then proceeded to complete the three experimental tasks during

separate fMRI runs in random order.

Behavioural motor task

Each of the submaximal handgrip conditions were acquired in separate block design fMRI runs. For

each run, participants performed 5 sets × 5 repetitions of either a 25%, 50%, or 75% MVC handgrip

visuomotor target matching task with an MRI-compatible hand clench dynamometer (Biopac

Systems Inc. Aero Camino Goleta, CA). Task blocks involved 1,650 ms [i.e., corresponding to the

repetition time (TR) of the T2* fMRI scan] contractions alternating with 1,650 ms of rest (16,500 ms

total task block). Each task block was separated by blocks of complete rest (16,500 ms total rest

block). During each fMRI run, the participants wore MRI compatible goggles and viewed a computer

screen projection, where they were able to view a custom-built LabView (version 8.6) interface.

Participants saw clear target lines and go/no-go on screen flashing lights and were cued when to

contract or relax. The LabView interface was triggered by the MRI through a fibre optic cable that

sent a signal at the beginning of each TR. This signal went through a Siemens fMRI trigger converter

and passed to the LabView interface via a BNC connection. This allowed for precise continuous

synchronization between the MRI and the experimental paradigm computer throughout the scan

(24). Target lines were presented relative to the individual's peak MVC and force feedback was

presented as a vertical force bar that was responsive to each participant's handgrip contraction (i.e.,

harder contraction resulted in the bar rising vertically). Participants were visually instructed when to

contract and when to relax with a green light that would turn on when the participant was to

contract. A second red light was programmed to turn on during rest blocks to indicate a sustained

rest. Participants were instructed to relax their non-active arm throughout each of the fMRI runs.

Handgrip motor task processing

The handgrip force data were processed in Matlab using custom scripts (25). Data were processed

with a fourth-order 100 Hz low pass Butterworth filter and full-wave rectified. The mean of each

contraction was then normalized to the mean Kg-force of the highest MVC and expressed as a

percentage of MVC. The mean of each task block (five contractions) was calculated and used in

subsequent analyses.

fMRI parameters

All scans were acquired in a Siemens 3T MAGNETOM Skyra MRI (Siemens Healthcare, Erlangen,

Germany). At the start of the session a whole-brain T1-weighted structural scan was acquired using a

high-resolution magnetization prepared rapid acquisition gradient echo (MPRAGE) sequence

consisting of 192 T1-weighted echo-planar imaging slices (1 mm slice thickness with no gap), with an

in-plane resolution of 1×1 mm [field of view = 256×256; TR = 1,900 ms; echo time (TE) = 2.08 ms].

Each of the experimental fMRI runs, were acquired using T2*-weighted single-shot gradient-echo

planar imaging with an interleaved ascending sequence, consisting of 105 volumes (TR = 1,650 ms;

TE = 30 ms) of 25 axial slices of 4 mm thickness (1-mm gap) with an in-plane resolution of 2.7 mm ×

2.7 mm (field of view = 250) using a flip angle of 90°. The top 2 coil sets (16 channels) of a 20-channel

Siemens head coil (Siemens Healthcare) were used. Runs consisted of a 10-volume alternating block

design beginning with five volumes for stabilization (task, rest; 105 volumes total).

fMRI preprocessing

Functional MRI data processing was carried out using FMRI Expert Analysis Tool (FEAT) Version 6.00,

as part of FSL [FMRIB's Software Library; (26)]. Boundary based registration was used to register the

functional image to the high-resolution T1-weighted (T1w) image. Registration of the functional

images to the T1w image was carried out using FMRIB's Linear Image Registration Tool [FLIRT: (27,

28)], and then non-linearly (rigid + affine + deformable syn) registered to the MNI152_1mm brain

template using antsRegistrationSyN and antsApplyTransforms from Advanced Normalization Tools

[ANTs; (29, 30)]. The following pre-processing was applied: motion correction using Motion

Correction FMRIB's Linear Image Registration Tool [MCFLIRT; (28)]; non-brain removal using FNIRT-

based brain extraction as implemented in the fsl_anat anatomical processing script (31); spatial

smoothing using a Gaussian kernel of FWHM 6 mm; grand-mean intensity normalization of the entire

4D dataset by a single multiplicative factor (32). Next, Independent Component Analysis Automatic

Removal of Motion Artifacts (ICA-AROMA) was used remove motion-related noise from each

functional run (32). Following ICA-AROMA data clean up, data were high pass temporal filtered with

a 0.01 Hz cut off. Time-series statistical analyses were carried out using FMRIB's Improved Linear

Model (FILM) with local autocorrelation correction (33). Z (Gaussianised T/F) statistic images were

constructed non-parametrically using Gaussian Random Field theory-based maximum height

thresholding with a corrected significance threshold of p = 0.05 (34).

Region of interest signal change

FSL's featquery tool was used to extract the BOLD signal percent change from three regions of

interest (ROI) in participants' native-space. For each ROI, ANTs were used to transform ROIs from

MNI-space to native-space prior to signal change extraction. The percent signal change was extracted

from the left contralateral and right ipsilateral Pontine reticular formation nuclei (left: PnO_PnC_l;

right: PnO_PnC_r) from the Brainstem navigator toolkit (35). Additionally, the percent signal change

was extracted from left contralateral M1 using an ROI mask from the Brainnetome atlas [left M1:

A4ul_l; (36)]. The extracted values were then used in subsequent statistical analyses.

Statistical analysis

An omnibus ROI [contralateral pontine reticular formation, ipsilateral pontine reticular formation] ×

condition (25%, 50%, 75% MVC) repeated measures Analysis of Variance (RM-ANOVA) was run using

the magnitude of brain activation from each of the ROI's (i.e., BOLD signal percent change from rest)

to determine whether RST activation changed with handgrip force modulation. Data were assessed

for normality using Shapiro-Wilks in addition to $Z_{skewness}$ and $Z_{kurtosis}$. ANOVA tests are robust against

violations to normality (37, 38) and thus minor violations to normality as determined by $z_{skewness}$ and

$Z_{kurtosis}$ were considered acceptable for running the parametric model. Sphericity was assessed with

Mauchley's test of sphericity and where violated the Greenhouse-Geisser correction was used.

Hedges g effect sizes were calculated and reported to adjust for the small sample size. A Bonferroni

adjustment was used to correct for multiple comparisons where appropriate post-hoc testing was

conducted. The α-level for significance was set to α = 0.05.

Additionally, for each ROI the slope coefficient of brain activation across the three submaximal

handgrip force conditions (percent change of BOLD signal activation / percent target force) was

calculated on a subject-by-subject basis using linear regression. The slope coefficients were then

analyzed using separate one-sample t-tests to determine if each ROI scales its magnitude of

activation across the varying submaximal force outputs. A positive slope indicates activation in that

ROI increases with handgrip force. A negative slope would represent a decrease in activation with an

increase in handgrip force and a non-significant slope (i.e., not different from zero) would represent

that activation did not vary as a function of handgrip force. For this analysis, the α-level was set to α

= 0.017 (0.05 / 3 ROIs) and *p*-values ≤ 0.017 were considered significant.

Results

Magnitude of BOLD Activation Across Contraction Intensities

RM-ANOVA was performed to test the effect of force (25%, 50%, and 75% MVC) and side (ipsilateral

and contralateral) on pontine reticular nuclei activation. Mauchly's test of sphericity was violated for

Force (*p* = 0.023) and Force × Side (*p* = 0.006). There was a significant main effect for force [$F(1.337,$

$16.039) = 5.306$, *p* = 0.012, $\eta_p^2 = 0.307$], but no main effect for side [$F(1,12) = 0.413$, *p* = 0.533, $\eta_p^2 =$

0.033]. The interaction effect (Force × Side) was also not significant [$F(1.243, 14.921) = 0.957$, *p* =

0.364, $\eta_p^2 = 0.074$]. A Bonferroni post hoc analysis showed that activation of pontine nuclei at 75%

MVC was significantly greater than at 25% MVC (MD = -0.2, SE = 0.068, *p* = 0.02) as well as at 50%

MVC (MD = -0.179, SE = 0.068, *p* = 0.042). The difference in activation between 25% and 50% MVC

was not significant (MD = -0.021, SE = 0.068, *p* = 1.000; Figure 2A).

A) Reticular Formation Activation (pontine nuclei)

B) Force Scaling - Individual Regression Analysis

C) Force Scaling - Group Analysis

Figure 2. The top panel **(A)** shows the activation in the pontine nuclei of the reticular formation during contractions of varying intensities. The contralateral and ipsilateral nuclei are collapsed together because there was neither a side × force intensity interaction nor a main effect for side as a factor. Pontine activation did show a main effect for force intensity, with post-hoc pairwise comparisons shown above the box plots. The middle panel **(B)** shows the relationship between % activation and % force from a single region of interest (ROI) from a representative subject. The relationship was fit with a linear model, and the slope interpreted as the degree to which activation for that region changes with increases in force. Panel **(C)** shows the slope coefficients from each subject for each ROI. The values above the scatter plot demonstrate the percent of subjects that exhibited a positive slope (i.e., positive scaling in activation with increasing force) for each ROI. The *p*-values above the box plots are

from one-sample *t*-tests, showing that the mean of the slopes was significantly different than
zero in the pons before correcting for multiple comparisons, but not in M1. Only the
ipsilateral pontine reticular nuclei significantly scaled after statistical correction (corrected α
= 0.017). The thick gray horizontal lines in A and C extend the zero point for easy comparison.

ROI BOLD Activation Scaling with Force

Linear regression analyses were conducted on each ROI for each subject to determine the slope

coefficient, which quantifies the rate of increase in activation with increasing contraction intensity

(Figure 2B). Next, using separate one sample *t*-tests of the individual slopes, we showed at the

group-level pontine reticular activation was significantly different from zero on both the ipsilateral

[$t(12) = 2.672$, $p = 0.01$, $g = 0.712$] and contralateral [$t(12) = 1.792$, $p = 0.049$, $g = 0.478$] sides before

correction; however, after a manual Bonferroni correction for multiple comparisons, only the

ipsilateral side remained significant. The contralateral M1 was not statistically different from zero

[$t(12) = 0.245$, $p = 0.405$, $g = 0.065$], indicating it did not reliably scale with force (Figure 2C).

Discussion

The present study examined the relationship between unimanual handgrip force production and

BOLD signal activation of the contralateral and ipsilateral pontine reticular nuclei of the reticulospinal

tract, as well as the contralateral M1 for comparison. Our primary finding is that activation within the

RST consistently scaled linearly with increased force production, which was not the case for the

contralateral M1. The traditional understanding is that the RST primarily innervates axial and

proximal muscle groups in humans (39). However, over the past 15 years a small body of literature

has begun to challenge this belief (8, 9, 11, 40) with research suggesting the RST may reach distal

musculature of the arms and hands and modulate force (12–14). With this task, we have shown, for

the first time, pontine reticular nuclei BOLD signal activation significantly scales with handgrip force,

where a positive relationship between the magnitude of activation and force output are observed.

Specifically, we observe significant scaling in the ipsilateral pons (with a large effect size) after

correcting for multiple comparisons. There was also scaling with a moderate effect size in the

contralateral pons. Additionally, the repeated measures ANOVA showed that side (ipsilateral vs.

contralateral) had no effect on activation across the force conditions. These findings, paired with the

lack of scaling in contralateral M1 provide some support for the notion of the RST pathway

contributing to high force outputs, rather than the traditional belief that human distal motor control

is exclusively governed by contralateral cortical regions. Schmidt et al. (41) showed that activity in

neurons from M1 do not change with increases in force in non-human primates (Rhesus macaque

monkeys). They concluded that the CST was responsible for activating a muscle, but not modulating

its force once it was active. Our findings support their conclusion, and add to it, as our data not only

agrees that the contralateral M1 and CST dont have that force modulation role in humans either, but

also provides an alternative for which pathway does, the RST.

During isometric contractions, the recruitment of motor units tend to occur in an orderly fashion

from low-threshold to higher-threshold (42). As force demand increases, larger and higher-threshold

motor units are recruited (43–46), and the firing rates of smaller, lower-threshold motor units that

were already active increase. These recruitment properties may also affect neuromuscular plasticity,

as the repeated activation of the larger motor units with high-intensity resistance training often lead

to greater gains in strength compared to lower-intensity training (47). In addition to the impact of

high-intensity contractions on motoneuron recruitment, and strength, a recent review by Hortobágyi

et al. (48) reported high-intensity resistance training also produces greater motor-related

neuroplasticity in the neurologically intact young adult brain. However, mechanistic studies into the

specific neural pathways and adaptations to resistance training that affords this motoneuronal

recruitment remains incompletely understood. This incomplete mechanistic understanding is due in

part to research focussing on the CST, which has resulted in equivocal insight. Here, we add to the

growing body of literature that pinpoints the RST as a relevant pathway for force control with a

handgrip task, which, until recently (12–14), was thought to be exclusively controlled by the CST (49).

Specifically, we show evidence for intensity dependent activation of the RST that is correlated with

force output, which suggests this is a task relevant pathway for handgrip force. Further, based on the

principles of use-dependent plasticity, the repeated preferential activation of the RST with high-

intensity contractions over the course of a resistance training intervention may lead to the

strengthening of this pathway's integrity in a manner that facilitates strength adaptations. Recent

findings have led to proposals that the RST may have a role in modulating motor unit discharge rates,

which would in turn alter force and rate of force production (14). Škarabot et al. (14) proposed

hypothesis about the role of RST in modulating force is supported by the findings of this study.

Evaluating the RST in applied settings presents its own set of challenges, such as the need for

external stimuli (e.g., Transcranial Magnetic Stimulation or cervical medullary electrical stimulation)

and perturbations (e.g., loud acoustic startle sounds ≥ 110 dB), in addition to a high variance in

responses. Here, using fMRI, we show a non-invasive evaluation of the RST in a controlled

environment during voluntary motor behaviours. We argue this is an important contribution to the

field investigating the role of the RST, as these data provide support to previous findings without the

need for external perturbations. Further, when paired with previous research (12, 13, 15), our

findings indicate that the RST may be a functionally relevant pathway contributing to muscle force

and, by extension, possibly chronic strength training adaptations.

Limitations and future directions

Investigating patterns of brain activation during high-intensity contractions with fMRI is not a trivial

task, as head motion commonly occurs during effortful movements, and can result in the loss of data

due to motion related contamination. With that, the investigation of fMRI related BOLD signal

activation during different types of motor tasks may not be possible, and thus these experiments are

limited to simpler isolated hand movement patterns that are less likely to produce high amounts of

motion.

Though we present evidence for the role of the RST in gross distal hand movement, future research

may consider testing RST activation during finer distal movements. By focusing on dexterous

movements such as flexion, extension, abduction, adduction, and circumduction, RST activation bias

toward certain types of movement may be determined. Future research into this area should

investigate the contributions of the RST to different tasks and their adaptations to training

interventions with varying manipulations. A deeper understanding of how the RST contributes to

motor function could lead to better rehabilitative modalities for those with dysfunction of the CST,

such as individuals with stroke (8, 12). Additionally, we focused only on the pontine nuclei of the RST.

The medullary nuclei had to be excluded because the medulla was cut off from the scan for many of

the participants in favour of capturing complete cortex. Future work should examine whether the

medullary nuclei also play a potential role in force modulation. Finally, future research would benefit

from replication of this experiment but with a larger sample size to build confidence in data

interpretation and robustness of findings.

Conclusions

In a small cohort of neurologically intact young adults, we demonstrate that activation of the pontine

reticular nuclei of the reticulospinal tract scales in magnitude with handgrip force. These data

provide insight into the potential role of the RST during distal muscular force modulation, a concept

most often considered under exclusive control by the CST in humans. These data add to the growing

body of literature suggesting a potentially important role of the RST for neuromuscular strength

adaptations in humans and lays the foundation for which future research can further investigate RST

function to disentangle the neural underpinnings of force modulation and the neural adaptations

governing increases in strength.

Data availability Statement

Data are available upon reasonable request

Acknowledgements

We would like to thank Shawn Reinink, University of Saskatchewan, for his contribution to the LabView

software interface using in this experiment.

Grants

This study was funded by a Natural Sciences and Engineering Research Council of Canada (NSERC)

Discovery Grant: 2016-0529, awarded to Jonathan P. Farthing, in addition to an NSERC Discovery Grant:

183968-2013-25, awarded to Ron Borowsky.

Disclosures

The authors have none to declare

Author Contributions

J.W.A. and L.G. collected the data. J.W.A. processed the data and assisted with statistical analyses and the

writing of the manuscript. T.L.D. took the lead on writing the manuscript and performed the statistical

analyses. J.M.D. conceived the research questions for which these data were used to answer, created

figures, contributed to writing, edited and approved the final version of the manuscript. R.L.M. assisted

with writing the manuscript and approved the final version. C.E. contributed to data processing and

analysis. J.P.F. and R.B. were involved in obtaining the grant funds and ethical approvals for the larger

study of which the data from this study are from, they were also involved in the original study design,

editing and approving the final version of the manuscript.

References

1. **Kalman M**. Evolution of the brain: At the reptile-bird transition. In: *Encyclopedia of*
Neuroscience. Springer Berlin Heidelberg, p. 1305–1312.

2. **Evarts E V.** Activity of pyramidal tract neurons during postural fixation. *J Neurophysiol* 32:
375–385, 1969. doi: 10.1152/JN.1969.32.3.375.

3. **Evarts E V.** Relation of pyramidal tract activity to force exerted during voluntary movement. *J*
Neurophysiol 31: 14–27, 1968. doi: 10.1152/JN.1968.31.1.14.

4. **Lemon RN**. Descending pathways in motor control [Online]. *Annu Rev Neurosci* 31: 195–218,
2008. http://www.ncbi.nlm.nih.gov/pubmed/18558853 [5 Sep. 2019].

5. **Courtine G**, **Bunge MB**, **Fawcett JW**, **Grossman RG**, **Kaas JH**, **Lemon R**, **Maier I**, **Martin J**,
Nudo RJ, **Ramon-Cueto A**, **Rouiller EM**, **Schnell L**, **Wannier T**, **Schwab ME**, **Edgerton VR**. Can
experiments in nonhuman primates expedite the translation of treatments for spinal cord
injury in humans? *Nat Med* 13: 561–566, 2007. doi: 10.1038/NM1595.

6. **Maier MA**, **Bennett KMB**, **Hepp-Reymond MC**, **Lemon RN**. Contribution of the monkey
corticomotoneuronal system to the control of force in precision grip [Online]. *J Neurophysiol*
69: 772–785, 1993. https://pubmed.ncbi.nlm.nih.gov/8463818/ [23 May 2021].

7. **Kidgell DJ**, **Bonanno DR**, **Frazer AK**, **Howatson G**, **Pearce AJ**. Corticospinal responses
following strength training: a systematic review and meta-analysis. *Eur J Neurosci* 46: 2648–
2661, 2017. doi: 10.1111/EJN.13710.

8. **Baker SN**, **Perez MA**. Reticulospinal contributions to gross hand function after human spinal
cord injury. *J Neurosci* 37, 2017. doi: 10.1523/JNEUROSCI.3368-16.2017.

9. **Honeycutt CF**, **Kharouta M**, **Perreault EJ**. Evidence for reticulospinal contributions to
coordinated finger movements in humans. [Online]. *J Neurophysiol* 110, 2013.
http://www.ncbi.nlm.nih.gov/pubmed/23825395 [5 Sep. 2019].

10. **Maitland S**, **Baker SN**. Ipsilateral motor evoked potentials as a measure of the reticulospinal
tract in age-related strength changes. *Front Aging Neurosci* 13: 612352, 2021. doi:
10.3389/FNAGI.2021.612352/BIBTEX.

11. **Riddle CN**, **Edgley SA**, **Baker SN**. Direct and indirect connections with upper limb
motoneurons from the primate reticulospinal tract. *J Neurosci* 29: 4993–4999, 2009. doi:
10.1523/JNEUROSCI.3720-08.2009.

12. **Akalu Y**, **Frazer AK**, **Howatson G**, **Pearce AJ**, **Siddique U**, **Rostami M**, **Tallent J**, **Kidgell DJ**.
Identifying the role of the reticulospinal tract for strength and motor recovery: A scoping
review of nonhuman and human studies. *Physiol Rep* 11: e15765, 2023. doi:
10.14814/PHY2.15765.

13. **Atkinson E**, **Škarabot J**, **Ansdell P**, **Goodall S**, **Howatson G**, **Thomas K**. Does the reticulospinal
tract mediate adaptation to resistance training in humans? [Online]. *J Appl Physiol* 133: 689–
696, 2022. https://pubmed.ncbi.nlm.nih.gov/35834623/ [1 Mar. 2023].

14. **Škarabot J**, **Folland JP**, **Holobar A**, **Baker SN**, **Del Vecchio A**. Startling stimuli increase maximal
motor unit discharge rate and rate of force development in humans. *J Neurophysiol* 128: 455–
469, 2022. doi: 10.1152/JN.00115.2022/ASSET/IMAGES/LARGE/JN.00115.2022_F008.JPEG.

15. **Glover IS**, **Baker SN**. Cortical, corticospinal, and reticulospinal contributions to strength
training. *J Neurosci* 40: 5820–5832, 2020. doi: 10.1523/JNEUROSCI.1923-19.2020.

16.   **Glover IS**, **Baker SN**. Both corticospinal and reticulospinal tracts control force of contraction. *J*
391   *Neurosci* 42: 3150–3164, 2022. doi: 10.1523/JNEUROSCI.0627-21.2022.

392   17.   **Tapia JA**, **Tohyama T**, **Poll A**, **Baker SN**. The existence of the StartReact effect implies
393   reticulospinal, not corticospinal, inputs dominate drive to motoneurons during voluntary
394   movement. *J Neurosci* 42: 7634–7647, 2022. doi: 10.1523/JNEUROSCI.2473-21.2022.

395   18.   **Hammerbeck U**, **Tyson SF**, **Samraj P**, **Hollands K**, **Krakauer JW**, **Rothwell J**. The strength of
396   the corticospinal tract not the reticulospinal tract determines upper-limb impairment level
397   and capacity for skill-acquisition in the sub-acute post-stroke period. *Neurorehabil Neural*
398   *Repair* 35: 812–822, 2021. doi: 10.1177/15459683211028243.

399   19.   **Xu J**, **Ejaz N**, **Hertler B**, **Branscheidt M**, **Widmer M**, **Faria A V.**, **Harran MD**, **Cortes JC**, **Kim N**,
400   **Celnik PA**, **Kitago T**, **Luft AR**, **Krakauer JW**, **Diedrichsen J**. Separable systems for recovery of
401   finger strength and control after stroke. *J Neurophysiol* 118: 1151–1163, 2017. doi:
402   10.1152/JN.00123.2017.

403   20.   **Alagona G**, **Delvaux V**, **Gérard P**, **De Pasqua V**, **Pennisi G**, **Delwaide PJ**, **Nicoletti F**, **Maertens**
404   **De Noordhout A**. Ipsilateral motor responses to focal transcranial magnetic stimulation in
405   healthy subjects and acute-stroke patients. *Stroke* 32: 1304–1309, 2001. doi:
406   10.1161/01.STR.32.6.1304.

407   21.   **Ziemann U**, **Ishii K**, **Borgheresi A**, **Yaseen Z**, **Battaglia F**, **Hallett M**, **Cincotta M**, **Wassermann**
408   **EM**. Dissociation of the pathways mediating ipsilateral and contralateral motor-evoked
409   potentials in human hand and arm muscles. *J Physiol* 518 (Pt 3): 895–906, 1999. doi:
410   10.1111/J.1469-7793.1999.0895P.X.

411   22.   **Baker SN**. The primate reticulospinal tract, hand function and functional recovery [Online]. *J*
412   *Physiol* 589: 5603–5612, 2011. http://www.ncbi.nlm.nih.gov/pubmed/21878519 [5 Sep.
413   2019].

414   23.   **Andrushko JW**, **Gould LA**, **Renshaw DW**, **Ekstrand C**, **Hortobágyi T**, **Borowsky R**, **Farthing JP**.
415   High force unimanual handgrip contractions increase ipsilateral sensorimotor activation and
416   functional connectivity. *Neuroscience* 452: 111–125, 2021. doi:
417   10.1016/j.neuroscience.2020.10.031.

418   24.   **Ekstrand C**, **Neudorf J**, **Kress S**, **Borowsky R**. Structural connectivity predicts functional
419   activation during lexical and sublexical reading. *Neuroimage* 218, 2020. doi:
420   10.1016/J.NEUROIMAGE.2020.117008.

421   25.   **The Mathworks Inc**. MATLAB 2018a. *WwwMathworksCom/Products/Matlab* 2, 2018.

422   26.   **Jenkinson M**, **Beckmann CF**, **Behrens TEJJ**, **Woolrich MW**, **Smith SM**. FSL [Online].
423   *Neuroimage* 62: 782–790, 2012. http://www.ncbi.nlm.nih.gov/pubmed/21979382 [8 Mar.
424   2020].

425   27.   **Jenkinson M**, **Smith S**. A global optimisation method for robust affine registration of brain
426   images. *Med Image Anal* 5: 143–156, 2001. doi: 10.1016/S1361-8415(01)00036-6.

427   28.   **Jenkinson M**, **Bannister P**, **Brady M**, **Smith S**. Improved optimization for the robust and
428   accurate linear registration and motion correction of brain images. *Neuroimage* 17: 825–841,
429   2002. doi: 10.1016/S1053-8119(02)91132-8.

430   29.   **Avants B**, **Tustison N**, **Johnson H**. Advanced Normalization Tools (ANTS). .

431   30.   **Avants BB**, **Epstein CL**, **Grossman M**, **Gee JC**. Symmetric diffeomorphic image registration
432   with cross-correlation: evaluating automated labeling of elderly and neurodegenerative

brain. *Med Image Anal* 12: 26–41, 2008. doi: 10.1016/J.MEDIA.2007.06.004.

31.   **Smith SM**, **Jenkinson M**, **Woolrich MW**, **Beckmann CF**, **Behrens TEJ**, **Johansen-Berg H**,
Bannister PR, **De Luca M**, **Drobnjak I**, **Flitney DE**, **Niazy RK**, **Saunders J**, **Vickers J**, **Zhang Y**, **De**
Stefano N, **Brady JM**, **Matthews PM**. Advances in functional and structural MR image analysis
and implementation as FSL [Online]. Neuroimage.
https://pubmed.ncbi.nlm.nih.gov/15501092/ [5 Mar. 2020].

32.   **Pruim RHR**, **Mennes M**, **van Rooij D**, **Llera A**, **Buitelaar JK**, **Beckmann CF**. ICA-AROMA: A
robust ICA-based strategy for removing motion artifacts from fMRI data. *Neuroimage* 112:
267–277, 2015. doi: 10.1016/j.neuroimage.2015.02.064.

33.   **Woolrich MW**, **Ripley BD**, **Brady M**, **Smith SM**. Temporal autocorrelation in univariate linear
modeling of FMRI data. *Neuroimage* 14: 1370–1386, 2001. doi: 10.1006/nimg.2001.0931.

34.   **Worsley K**. Statistical analysis of activation images. In: *Functional MRI: An Introduction to*
Methods, edited by Jezzard P, Matthews PM, Smith SM. Oxford: Oxford University Press,
2001, p. 251–270.

35.   **Bianciardi M**. Brainstem Navigator. .

36.   **Fan L**, **Li H**, **Zhuo J**, **Zhang Y**, **Wang J**, **Chen L**, **Yang Z**, **Chu C**, **Xie S**, **Laird AR**, **Fox PT**, **Eickhoff**
SB, **Yu C**, **Jiang T**. The human brainnetome atlas: A new brain atlas based on connectional
architecture. [Online]. 26, 2016. https://dx.doi.org/10.1093/cercor/bhw157 [17 Apr. 2020].

37.   **Schmider E**, **Ziegler M**, **Danay E**, **Beyer L**, **Bühner M**. Is It really robust?: Reinvestigating the
robustness of ANOVA against violations of the normal distribution assumption [Online].
Methodology 6: 147–151, 2010. https://econtent.hogrefe.com/doi/abs/10.1027/1614-
2241/a000016 [19 Dec. 2020].

38.   **Blanca MJ**, **Alarcón R**, **Arnau J**, **Bono R**, **Bendayan R**. Non-normal data: Is ANOVA still a valid
option? [Online]. *Psicothema* 29: 552–557, 2017.
https://pubmed.ncbi.nlm.nih.gov/29048317/ [19 Dec. 2020].

39.   **Kuypers HGJM**. Anatomy of the descending pathways. .

40.   **Zaaimi B**, **Edgley SA**, **Soteropoulos DS**, **Baker SN**. Changes in descending motor pathway
connectivity after corticospinal tract lesion in macaque monkey. *Brain* 135: 2277–2289, 2012.
doi: 10.1093/BRAIN/AWS115.

41.   **Schmidt EM**, **Jost RG**, **Davis KK**. Reexamination of the force relationship of cortical cell
discharge patterns with conditioned wrist movements. *Brain Res* 83: 213–223, 1975. doi:
10.1016/0006-8993(75)90931-2.

42.   **De Luca CJ**, **Hostage EC**. Relationship between firing rate and recruitment threshold of
motoneurons in voluntary isometric contractions. *J Neurophysiol* 104: 1034–1046, 2010. doi:
10.1152/JN.01018.2009.

43.   **Milner-Brown HS**, **Stein RB**, **Yemm R**. The orderly recruitment of human motor units during
voluntary isometric contractions [Online]. *J Physiol* , 1973.
https://pubmed.ncbi.nlm.nih.gov/4350770/ [27 Oct. 2023].

44.   **Milner-Brown HS**, **Stein RB**, **Yemm R**. Changes in firing rate of human motor units during
linearly changing voluntary contractions. *J Physiol* 230: 371–390, 1973. doi:
10.1113/JPHYSIOL.1973.SP010193.

45.   **de Luca CJ**, **LeFever RS**, **McCue MP**, **Xenakis AP**. Behaviour of human motor units in different
muscles during linearly varying contractions. *J Physiol* 329: 113–128, 1982. doi:

10.1113/JPHYSIOL.1982.SP014293.

46.   **Milner-Brown HS**, **Stein RB**, **Yemm R**. The contractile properties of human motor units during
478   voluntary isometric contractions. *J Physiol* 228: 285–306, 1973. doi:
479   10.1113/JPHYSIOL.1973.SP010087.

480   47.   **Fry AC**. The role of resistance exercise intensity on muscle fibre adaptations. *Sport Med* 34:
481   663–679, 2004. doi: 10.2165/00007256-200434100-00004.

482   48.   **Hortobágyi T**, **Vetrovsky T**, **Balbim GM**, **Sorte Silva NCB**, **Manca A**, **Deriu F**, **Kolmos M**,
483   **Kruuse C**, **Liu-Ambrose T**, **Radák Z**, **Váczi M**, **Johansson H**, **dos Santos PCR**, **Franzén E**,
484   **Granacher U**. The impact of aerobic and resistance training intensity on markers of
485   neuroplasticity in health and disease. *Ageing Res Rev* 80, 2022. doi:
486   10.1016/J.ARR.2022.101698.

487   49.   **Natali AL**, **Reddy V**, **Bordoni B**. Neuroanatomy, corticospinal cord tract [Online].
488   https://www.ncbi.nlm.nih.gov/books/NBK535423/ [27 Oct. 2023].