A novel phylogenomics pipeline reveals complex pattern of reticulate evolution in

Cucurbitales

Edgardo M. Ortiz[1,2*], Alina Höwener[1,3], Gentaro Shigita[1,4], Mustafa Raza[1], Olivier Maurin[5,6],

Alexandre Zuntini[6], Félix Forest[6], William J. Baker[6,7] & Hanno Schaefer[1]

[1] Plant Biodiversity, Department Life Science Systems, Technical University of Munich

(TUM), Emil-Ramann-Str. 2, D-85354 Freising, Germany

[2] current address: Florida Museum of Natural History, University of Florida, Gainesville, FL

32611, USA

[3] current address: Systematics, Biodiversity & Evolution of Plants, Ludwig-Maximilian

University Munich (LMU), Menzingerstr. 67, D-80638 Munich, Germany.

[4] Faculty of Life and Environmental Sciences, University of Tsukuba, 305-8572 Tsukuba,

Japan

[5] Botanic Garden Meise, Nieuwedaan38, B-1860 Meise, Belgium

[6] Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3AE, United Kingdom

[7] Department of Biology, Aarhus University, Aarhus, Denmark

[*] E-mail: e.ortiz.v@gmail.com

Abstract.--- A diverse range of high-throughput sequencing data, such as target capture,

RNA-Seq, genome skimming, and high-depth whole genome sequencing, are used for

phylogenomic analyses but the integration of such mixed data types into a single

phylogenomic dataset requires a number of bioinformatic tools and significant computational

resources. Here, we present a novel pipeline, CAPTUS, to analyze mixed data in a fast and

efficient way. CAPTUS assembles these data types, allows searching of the assemblies for loci

of interest, and finally produces alignments filtered for paralogs. Compared to other software,

CAPTUS allows the recovery of a greater number of more complete loci across a larger number

of species. We apply CAPTUS to assemble a comprehensive mixed dataset, comprising the

four types of sequencing data for the angiosperm order Cucurbitales, a clade of about 3,100

species in eight mainly tropical plant families, including begonias (Begoniaceae) and gourds

(Cucurbitaceae). Our phylogenomic results support the currently accepted circumscription of

Cucurbitales except for the position of the holoparasitic Apodanthaceae, which group with

Rafflesiaceae in Malpighiales. A subset of mitochondrial gene regions supports the earlier

position of Apodanthaceae in Cucurbitales. However, the nuclear regions and majority of

mitochondrial regions place Apodanthaceae in Malpighiales. Within Cucurbitaceae, we

confirm the monophyly of all currently accepted tribes but also reveal deep reticulation

patterns both in Cucurbitales and within Cucurbitaceae. We show that contradicting results

among earlier phylogenetic studies in Cucurbitales can be reconciled when accounting for

gene tree conflict and demonstrate the efficiency of CAPTUS for complex datasets.

Keywords.--- Angiosperms353, Cucurbitaceae, Cucurbitales, gene tree discordance, paralog

filtering, genome skimming, phylotranscriptomics, nitrogen-fixing clade

Modern sequencing technology allows the rapid accumulation of large amounts of

genomic data at low cost. The analysis of these raw data, however, can still be a bottleneck

and there is strong demand for skilled bioinformaticians who can handle such complex data in

industry and in research institutions worldwide. Researchers who cannot afford to pay for a

bioinformatics expert rely on user-friendly analysis pipelines that are accessible without

specialist training. As a consequence, these pipelines are of enormous importance for the field

of phylogenomics. Among them, we find PHYLUCE (Faircloth 2016) and UCEasy (Ribeiro

et al. 2021) specialized on the analysis of Ultra Conserved Elements (UCEs), HYBPIPER

(Johnson et al. 2016) and SECAPR (Andermann et al. 2018) aimed at target capture data,

PhyloHerb (Cai et al. 2022) for genome skimming data or more generalist pipelines, such as

PhyloTol (Cerón-Romero et al. 2019), that attempt to find all the possible gene families

across a taxonomic group.

HYBPIPER and SECAPR are the most commonly used pipelines for target capture data

in plant phylogenomics, however, they are designed to process one sample at a time, only

making use of multiple computer cores at certain stages [e.g., when assembling sequences *de*

novo with SPADES (Bankevich et al. 2012)]. To process multiple samples simultaneously, the

users must rely on external tools such as GNU parallel (Tange 2021) or additional

containerization tools (Jackson et al. 2023). This lack of native parallelization capabilities can

lead to suboptimal utilization of the computing resources in high-performance clusters or

workstations and reduce accessibility for non-experts, potentially causing repeatability issues

due to command inconsistencies across samples or human error. Also, since these pipelines

were optimized for target capture data, their processing times for other data types can be

exceptionally long. In contrast, most specialized user-friendly analysis pipelines designed for

restriction site associated DNA (RAD) data, are aimed to efficiently process many samples in

parallel, e.g., IPYRAD (Eaton and Overcast 2020), DDOCENT (Puritz et al. 2014), and to a lesser

extent, Stacks (Rochette et al. 2019). For example, IPYRAD (Eaton and Overcast 2020) uses a

single or just a few consistent commands, that, combined with relatively short processing

times, allow for rapid testing of alternative settings. IPYRAD allows users to summarize results

of each processing step so that they can decide on the most appropriate settings needed for

following steps.

With these features in mind, we developed CAPTUS, a pipeline written entirely in

Python and aimed at building phylogenomic datasets from multiple types of high-throughput

sequencing data (target capture, RNA-Seq, genome skimming, and high-depth whole genome

sequencing). CAPTUS makes extensive use of Python's native parallel computing to process

many samples simultaneously in a consistent manner. This parallelization can be automated

or configured manually according to user needs. Each step has a simple basic command

syntax (although it can be customized with many options), provides extensive reports in

HTML to guide the settings of the next step, and is fully logged for repeatability. CAPTUS is

able to extract nuclear, mitochondrial, and plastid proteins considering the different genetic

codes and gene characteristics of each genomic compartment, as well as any other type of

DNA regions (such as ribosomal RNA, introns, spacers, RAD loci, etc.) in a single command.

In HYBPIPER, this would require as many runs as marker types, in SECAPR only references

composed of individual exons in nucleotides are possible. During the alignment stage,

CAPTUS can also filter paralogs by taking advantage of reference target files that contain

multiple sequences per locus such as the Angiosperms353 target file, a set of genes targeted

by a universal probe set (Johnson et al. 2019), and its more taxonomically comprehensive

version Mega353 (McLay et al. 2021). The final outputs of CAPTUS are multiple sequence

alignments (MSAs) for each reference locus found in the samples. The MSAs are organized

by genomic compartment and format (amino acid, nucleotide, etc.), and multiple versions of

each are provided (i.e., unfiltered, paralog-filtered, untrimmed, and trimmed) so the user can

select which type of alignment to analyze for phylogenetic inference. Additionally, a special

feature of CAPTUS is the search for novel conserved markers by clustering contigs that

received no hits from the reference target loci across samples, thus making use of data that

otherwise would be ignored.

We demonstrate the potential of CAPTUS with the example of the flowering plant order

Cucurbitales, according to APG IV (The Angiosperm Phylogeny Group 2016) a clade of eight

plant families with diversity centers in the Tropics: Anisophylleaceae, Apodanthaceae,

Begoniaceae, Coriariaceae, Corynocarpaceae, Cucurbitaceae, Datiscaceae, and

Tetramelaceae. Together, they include 110 genera with more than 3,100 species, about 2,000

of them in the mega-diverse genus *Begonia* in Begoniaceae (Goodall-Copestake et al. 2009)

and 1,000 in Cucurbitaceae (Stevens 2001; Schaefer 2020). The genus *Begonia* is of great

horticultural importance while Cucurbitaceae include some of the most important vegetable

and fruit crops worldwide, like cucumber and watermelon (Schaefer and Renner 2011a).

Morphologically, the taxa of Cucurbitales are rather diverse, ranging from endophytic

holoparasites in Apodanthaceae (Bellot and Renner 2014) to annual and perennial herbs in

Datiscaceae and Begoniaceae, to trees and shrubs in Anisophylleaceae, Corynocarpaceae,

Coriariaceae, and Tetramelaceae, and finally to woody or herbaceous climbers and creepers in

Cucurbitaceae (Schaefer and Renner 2011a; Schaefer 2020).

A number of studies addressed phylogenetic problems in Cucurbitales in the past three

decades. The earliest comprehensive seed plant phylogeny based on *rbc*L sequences (Chase et

al. 1993) already identified a clade uniting the genera *Coriaria* (Coriariaceae), *Datisca*

(Datiscaceae), *Luffa* (Cucurbitaceae), *Cucurbita* (Cucurbitaceae), *Begonia* (Begoniaceae), and

Tetrameles (Tetramelaceae). Zhang et al. (2006) produced the first phylogeny estimate

focused on Cucurbitales based on nine plastid, nuclear and mitochondrial loci and confirmed

the results of Chase et al. (1993) while adding Anisophylleaceae and Corynocarpaceae to the

Cucurbitales clade. The work of Barkmann et al. (2007) suggested another surprising addition

to the Cucurbitales clade: based on mitochondrial *mat*R sequence data, the holoparasitic

Apodanthaceae were placed in Cucurbitales. Filipowicz and Renner (2010) increased the

*mat*R dataset of Barkmann et al. (2007), added a nuclear 18S dataset, and confirmed that the

holoparasitic Apodanthaceae were best placed in Cucurbitales as sister lineage to all other

families. Schaefer and Renner (2011b) inferred phylogenetic relationships in the order based

on 14 DNA regions from all three genomes, confirming the position of Apodanthaceae based

on *mat*R, while their nuclear 18S-ITS-26S dataset suggested that Apodanthaceae group

outside Cucurbitales. Other studies targeted individual families: Zhang et al. (2007) provided

the first comprehensive phylogeny estimate for Anisophylleaceae, Kocyan et al. (2007) and

Schaefer et al. (2009) for Cucurbitaceae, Goodall-Copestake et al. (2009) for Begoniaceae,

and Renner et al. (2020) for Coriariaceae. In recent years, several phylogenomic studies

contributed to an even better understanding of the evolutionary relationships within the family

131 Cucurbitaceae. Bellot et al. (2020) analyzed entire plastomes plus a set of 57 single-copy

132 nuclear genes and the ITS region for 29 species from all but one of the accepted tribes and

133 detected four nodes with conflicting phylogenetic signals. With an impressive set of 136

134 transcriptomes and full genomes of Cucurbitaceae, representing c. 50% of the genera, Guo et

135 al. (2020) produced a well-supported phylogeny estimate for the family that conflicted with

136 earlier studies in several positions (Kocyan et al. 2007; Schaefer et al. 2009; Schaefer and

137 Renner 2011b). Finally, our recent angiosperm-wide genus level analysis (Zuntini et al. 2024)

138 in the framework of the Plant and Fungal Trees of Life (PAFTOL) project (Baker et al. 2022)

139 with an almost complete representation of Cucurbitales at the genus level placed the

140 holoparasitic Apodanthaceae outside Cucurbitales in Malpighiales and found a sister group

141 relationship between Cucurbitales and Rosales, challenging the results of most earlier studies.

142  In this study, we demonstrate how our new analysis pipeline CAPTUS can be used not

143 only to extract the Angiosperms353 genes (Johnson et al. 2019), but also to derive a new set

144 of thousands of nuclear genes from transcriptomic data and to extract all mitochondrial genes

145 and entire plastomes for the Cucurbitales. The resulting phylogenomic datasets were analyzed

146 with coalescent and concatenation methods to infer a complete genus-level nuclear phylogeny

147 as well as plastid and mitochondrial phylogenies and test the earlier results for the topology of

148 the order. We compare the performance and efficiency of CAPTUS against the most widely

149 used pipeline HYBPIPER, which is also able to use amino acid sequences as reference targets

150 and can handle multiple reference target sequences per locus.

152        MATERIALS & METHODS

153          *Sampling*

154  We used a total of 125 Cucurbitales samples. Of these, 113 are target capture

155 sequencing samples produced for this study, but also incorporated in broader angiosperm

156 phylogenetic analyses (Baker et al. 2022; Zuntini et al. 2024), while five target capture

samples and seven high-depth whole genome sequencing samples have not been previously

published (Table S1). Additionally, we downloaded sequence data for 327 samples

(Cucurbitales and outgroups) available in NCBI's SRA: 240 RNA-Seq samples, 48 genome

skimming samples (<35x), 31 high-depth whole genome sequencing samples (>35x), and

eight additional target capture samples (Table S2). The entire dataset comprises 342 samples

of Cucurbitales sensu APG IV, representing all 110 currently accepted genera and 249 of the

c. 3,100 species. In order to test the results of our recent angiosperm-wide analysis (Zuntini et

al. 2024) and the study of Alzate et al. (2024), who found Apodanthaceae outside

Cucurbitales as sister of Malpighiales or included in Malpighiales, we decided to use a rather

broad outgroup selection that includes 110 samples covering taxa across the three other orders

of the nitrogen-fixing clade (25 species in eight families of Rosales, 11 species in six families

of Fagales, and 28 species in three families of Fabales), as well as 31 species in 23 families of

Malpighiales, six species in six families of Malvales, three species of Huaceae (Huales), and

three species of Vitaceae (Vitales).

172   *DNA Extraction and Sample Preparation*

173   DNA samples were taken from the DNA bank of the Plant Biodiversity group at

174   Technical University of Munich and the DNA bank of Royal Botanic Gardens, Kew. Tissue

175   samples for further DNA extractions came from the herbaria of Royal Botanic Gardens, Kew

176   (K), Muséum National d'Histoire Naturelle in Paris (P), Botanische Staatssammlung

177   München (M), and Technical University of Munich (TUM).

178   Several different methods were used for DNA extraction from herbarium material

intended for target capture: (i) a CTAB-chloroform-based protocol with ethanol washes and a

cesium chloride/ethidium bromide density gradient cleaning and dialysis, (ii) a modified

CTAB protocol (Doyle and Doyle 1987) followed by an Agencourt AMPure XP bead clean-

up (Beckman Coulter, Indianapolis, Indiana, USA), (iii) an SDS-based protocol using Magen

HiPure SF Plant DNA Kit (Angen Biotech Co., Ltd, Guangzhou, China) using magnetic

beads for extraction and cleaning, and (iv) the manufacturers protocol of the CTAB-based

NucleoSpin Plant II Extraction Kit (MACHEREY-NAGEL GmbH & Co. KG, Düren,

Germany) using silica columns for binding the DNA. Depending on the available herbarium

material, we used 20-160 mg of dry material for extraction.

To measure DNA quality, we evaluated fragment size distribution using an Agilent

Technologies 4200 TapeStation System with Genomic DNA ScreenTapes (Agilent

Technologies, Santa Clara, California, USA) or by electrophoresis in 1% agarose gel. DNA

was quantified using a Quantus™ Fluorometer (Promega Corporation, Madison, WI, USA) or

with a Qubit 3.0 Fluorometer (Thermo Fisher Scientific, Invitrogen, Walthan, Massachusetts,

USA).

The seven samples intended for high-depth whole genome sequencing were extracted

with one of the following methods: (i) the standard protocol of the CTAB-based NucleoSpin

Plant II Extraction Kit, and (ii) the standard protocol of the MagBind® Plant DNA plus 96

Kit from Omega (https://www.omegabiotek.com/product/mag-bind-plant-dna-plus-96-kit/).

These seven DNA extractions were sent to GENEWIZ Germany GmbH (Leipzig, Germany),

where libraries were prepared with the NEBNext Ultra II Library Prep Kit for Illumina (New

England BioLabs, Ipswich, Massachusetts, USA) and sequenced on a NovaSeq 6000 platform

(Illumina, San Diego, California, USA) with a paired-end 2x150 configuration.

Library Preparation, Pooling, Target Enrichment, and Sequencing

Shearing was only performed for samples with fragment sizes above 350 bp using a

Covaris M220 Focused-ultrasonicator with Covaris microTUBES AFA Fiber Pre-Slit Snap-

Cap (Covaris, Woburn, Massachusetts, USA). Libraries were prepared with the DNA

NEBNext UltraTM II Library Prep Kit, including end-repair/end-prep, NEBNext Adapter

ligation, size selection of preferred DNA fragments with a length of 300 to 400 bp using SPRI

beads (Agencourt AMPure XP Bead Clean-up; Beckman Coulter, Indianapolis, IN, USA) and

amplification with NEBNext Multiplex Oligos for Illumina (Dual Index Primer Set 1 and Set

2, New England BioLabs, Ipswich, Massachusetts, USA) as indices for sample identification.

Libraries were grouped and pooled by similar DNA quality and quantity. For target

enrichment via hybridization of the pooled libraries, we used the Arbor Biosciences myBaits

Target Capture Kit, "Angiosperms353 v1" (Catalog #3081XX), following the manufacturer's

manual Version 4.01 (myBaits® Kit Manual – Arbor Biosciences 2020). Hybridized libraries

were PCR-amplified using the NEBNext Ultra II Q5 Master Mix (New England BioLabs,

Ipswich, Massachusetts, USA). Quality and quantity of the library pools were evaluated with

TapeStation and Qubit respectively as above, and then normalized and pooled for sequencing

of paired-end 2x300 reads on a MiSeq platform (Illumina, San Diego, California, USA) at the

Jodrell Laboratory, Royal Botanic Gardens, Kew or paired-end 2x150 reads at Macrogen

Europe B.V. in Amsterdam (Macrogen, Inc., Seoul, Korea) with a HiSeq system (Illumina,

San Diego, California, USA).

224   *Sequence Analysis Workflow*

CAPTUS, implemented in Python 3, is freely available and maintained through GitHub

(https://github.com/edgardomortiz/Captus) including a full documentation

(https://edgardomortiz.github.io/captus.docs/). It can be installed directly from Bioconda

(https://bioconda.github.io/recipes/captus/README.html). The workflow of CAPTUS consists

of four steps controlled by their respective modules called clean, assemble, extract,

and align, which are typically run in that order (Fig. 1).

FIGURE 1. Workflow of the analysis pipeline CAPTUS. The default thresholds shown can be

changed by the user via command options.

Alternatively, the analysis can be started at different points depending on the sample

data provided. If raw reads are provided one should start with the clean module, if the reads

have already been cleaned, they can be directly analyzed with the assemble module, and if

previously assembled data or reference genomes are provided the analysis can start using the

`extract` module. CAPTUS provides flexibility in combining samples that enter the workflow

at different stages. For example, a phylogenomic dataset could be composed of samples

represented by target capture raw sequencing reads, samples with clean RNA-Seq reads, and

genomic assemblies downloaded from GenBank in order to increase taxon sampling.

The `clean` *module.---* CAPTUS uses BBDUK from BBTOOLS (Bushnell 2022) to clean raw

reads. Cleaning is performed in two steps. First, adaptors and poly-A tails (when cleaning

RNA-Seq reads) are trimmed in two consecutive rounds. Then, adaptor-free reads are quality-

trimmed, and reads matching the phiX viral genome or known sequencing artifacts are

removed using the databases provided by BBDUK. In this step, leading and trailing bases with

PHRED scores <13 are trimmed and then trimmed reads with average PHRED score <16 are

removed, both thresholds can be altered by the user.

When cleaning is completed, FASTQC (Andrews 2019) or its faster implementation

FALCO (de Sena Brandine and Smith 2021) is run both on the raw and the cleaned reads to

evaluate and compile quality statistics. Finally, CAPTUS summarizes the cleaning statistics of

all the samples before and after cleaning in a single HTML report that allows a quick

overlook of all quality measurements.

For the Cucurbitales analysis, the default settings were used for all samples in the

`clean` module except RNA-Seq data, for which the option `--rna` was added to trim poly-A

tails.

The `assemble` *module.---* The clean reads are passed to the assembly module which uses

MEGAHIT for *de novo* assembly. This assembler is capable of handling very large datasets in

a time-efficient manner and with greatly reduced memory requirements because of the use of

a compressed representation of *de Bruijn* graphs and very efficient parallel computing

algorithms. More importantly, MEGAHIT implements the concept of *mercy-kmers*, a unique

strategy that allows removing spurious low-depth contigs while keeping authentic ones (Li et

al. 2015, 2016). Thanks to these features, MEGAHIT is not only exceptional at assembling

metagenomic datasets (its original purpose) but also single-genome datasets, and is able to

produce accurate assemblies even at low sequencing depths as shown in our previous tests

with small datasets (Raza et al. 2023).

We tested several combinations of MEGAHIT settings in order to optimize the

assembly of other types of data. These are provided in CAPTUS under three presets: (i)

`CAPSKIM` for target sequence capture, genome skimming data, or a combination of both, (ii)

`RNA` for RNA-Seq data, and (iii) `WGS` for high-depth whole genome sequencing data

(https://edgardomortiz.github.io/captus.docs/assembly/assemble/options/#--preset).

The `assemble` module also provides an option to subsample a fixed number of reads per

sample prior to assembly, which is useful when the sequencing depth is too high for a

particular sample or for speeding up assembly during trial runs.

Once assembly is completed, CAPTUS can also filter contigs based on their GC

content. For example, a maximum of 60% GC would be appropriate to remove bacterial

contamination in most eukaryotic genomes (Fierst and Murdock 2017). Finally, CAPTUS

computes assembly statistics and produces a single HTML report for all processed samples.

Target capture and genome skimming samples of the Cucurbitales were run with the

default assembly preset (`--preset CAPSKIM`), RNA-Seq samples were assembled using -

`-preset RNA`, and high-depth whole genome sequencing samples using `--preset WGS`.

To remove potential bacterial contamination, we used the option `--max_contig_gc 60`

for all assemblies.

The `extract` *module.*--- Once assemblies are completed, they can be searched for particular

loci of interest (i.e., reference target loci). CAPTUS allows the simultaneous search and

extraction of five types of markers (i) nuclear proteins (NUC), (ii) plastid proteins (PTD), (iii)

mitochondrial proteins (MIT), (iv) miscellaneous DNA regions (DNA), and (v) new

clustering-derived putative homologs (CLR).

Protein extractions (NUC, PTD, and MIT markers) use reference target loci provided as

amino acid sequences or coding sequences in nucleotides (CDS) and a different genetic code

can be specified for each separately. CAPTUS uses the program SCIPIO to perform protein

extractions. SCIPIO is able to automatically correct frameshifts that could result from

sequencing or assembly error and it can recover proteins and their putative paralogs that are

spread across several contigs by analyzing splice patterns, checking for compatible exonic

boundaries and overlaps, and for consistency in the similarity of the partial hits to the target

protein to reconstruct coherent gene models (Keller et al. 2008; Hatje et al. 2011). This is

important, given that loci assemblies resulting from target capture data can potentially be

highly fragmented, particularly when the captured target proteins contain exons that are

separated by long introns. CAPTUS comes bundled with four reference target sets, two NUC

references [curated versions of the Angiosperms353 (Johnson et al. 2019) target file and the

taxonomically expanded Mega353 (McLay et al. 2021) target file], as well as a plastid PTD

reference (SeedPlantsPTD), and a mitochondrial MIT reference (SeedPlantsMIT).

Our curation of the Angiosperms353 reference target file consisted in clustering the

sequences at 98% identity to remove highly similar sequences and duplicates while the

curation of the Mega353 reference target file consisted in removing contaminated sequences

(e.g., fungal, algal, organellar sequences) and then clustering at 78% identity to reduce

sequence redundancy. Thus, CAPTUS still takes advantage of the expanded taxonomic

representation of the Mega353 reference targets (McLay et al. 2021) while bypassing the

additional step of filtering by taxon suggested in the instructions

(https://github.com/chrisjackson-pellicle/NewTargets). The PTD and MIT target references

represent carefully curated sets of 83 proteins (represented by 2,194 sequences) and 45

proteins (represented by 167 sequences) respectively, found in most organellar genomes of

seed plants originally downloaded from GenBank. Captus also allows the utilization of

BUSCO lineage databases (e.g. https://busco-data.ezlab.org/v5/data/lineages/) as target

references and new target sets for organism groups other than seed plants are possible and

being developed.

Miscellaneous DNA extractions are aimed at recovering other types of DNA regions

(e.g., complete genes with introns, non-coding regions, ribosomal genes, individual exons,

RAD markers, etc.). In this case, CAPTUS uses the program BLAT (Kent 2002) to search the

assembly and the hits that are found across contigs are greedily assembled and concatenated

using our own code.

Additionally, CAPTUS can be used to find new putative homologs by clustering across

samples the contigs that had no hits from the reference target loci. In this case, the program

MMSEQS2 (Steinegger and Söding 2018) is used, and several of its settings are available

through CAPTUS. Once the sequence clusters have been found, they are filtered by the number

of samples in the cluster and sequence length, and only the most represented sequences per

cluster are selected as new reference targets to perform another extraction of the

miscellaneous DNA type in CAPTUS. The output from this kind of extraction is indicated by

the prefix CLR.

The user can specify the minimum sequence identity and coverage to perform the

search of target loci within the assemblies. If several matches are found for a single target

locus, they are ranked from most to least similar and complete as compared to the reference

target sequence. Therefore, the best hit is recognized in the output files by having the suffix

_00 while the rest of the copies (i.e., putative paralogs) are ranked with increasing numbers

(e.g. _01, _02, etc.).

The output files from protein extractions (NUC, PTD, MIT) are provided in four

possible formats (Fig. S1a): the protein sequence in amino acids (AA), the coding sequence in

nucleotides or CDS (`NT`), the complete gene sequence including introns (`genes`), and the

complete gene sequence flanked by a fixed number of nucleotides (`genes_flanked`).

Similarly, the output from a miscellaneous DNA extraction and clustering-derived markers

(`DNA`, `CLR`), have two possible formats (Fig. S1b): the segment of sequence that was matched

to the reference (`matches`), and the matched segments flanked by a fixed number of

nucleotides (`matches_flanked`).

Once extractions are finished, CAPTUS collects extraction statistics (e.g., recovered

marker length, similarity to the reference sequence, number of paralogs, number of contigs

used in the gene assembly, etc.) from all the processed samples and produces a single HTML

report that provides a quick overview of the recovery across regions and samples as an

interactive heatmap.

For our Cucurbitales dataset, the minimum recovery percentage was set at 20%, the

default in CAPTUS. We used the option `-n Angiosperms353` to extract nuclear proteins

using the bundled Angiosperms353 reference targets and `-n Mega353` to extract nuclear

proteins using our curated version of the taxonomically expanded Mega353. Additionally,

organellar proteins were extracted with options `-p SeedPlantsPTD` and `-m`

`SeedPlantsMIT`. Finally, we created a custom miscellaneous DNA reference by

segmenting the Cucurbitales plastomes available in GenBank in 38 pieces ranging from ~3

kbp to ~5 kbp which we stored in a FASTA file (`Plastome38.fasta`, Appendix S1) and

extracted in CAPTUS using the option `-d`.

Additionally, we created a set of 5,435 reference genes derived from publicly

available transcriptomic data. We took the 240 transcriptomic assemblies from CAPTUS and

removed the transcripts that had hits to organellar proteins, thereby retaining only putatively

nuclear transcripts. We ran CODAN (Nachtigall et al. 2021) on the nuclear transcripts in order

to find coding regions within them, searching both strands and using the `PLANTS_full`

model which only emits a CDS when a complete protein, from start to stop codon, is

identified within the transcript. Then we clustered the CDS across samples using the

`extract` module of CAPTUS, with the following options:

`-c --mmseqs2_method easy-cluster --cluster_mode 2 --`

`cl_min_identity 70 --cl_seq_id_mode 1 --cl_min_coverage 66 --`

`cl_rep_min_len 540 --cl_min_samples 144 --cl_max_copies 4`. These

CAPTUS options retained 5,435 clusters that were at least 540 bp in length, grouped at least

144 transcriptomic samples (60%), and contained at most an average of four gene copies.

After removing within-cluster redundant sequences, the resulting reference contained 13,492

sequences representing these 5,435 CDS clusters. This reference is called from this point

onwards RNA5435 (`RNA5435.fasta`, Appendix S2). The new reference targets were used

in the `extract` module of CAPTUS across all 452 samples using the option `-n`

`RNA5435.fasta --nuc_min_score 0.15 --nuc_min_identity 70` to match

the clustering identity threshold used for creating the reference .

The `align` *module.*--- The extraction output is then processed by the alignment module.

Individual markers are collected across samples and organized in separate FASTA files per

locus. By default, CAPTUS collects a maximum of five paralogs per sample and per marker to

be aligned. The reference sequences are also added to each locus file to serve as alignment

guides in case the sequences recovered from the samples are fragmentary. Then, the

alignment is performed with MAFFT (Katoh and Standley 2013) or MUSCLE5 (Edgar 2022)

using default settings or by selecting one of their specific algorithms through CAPTUS.

However, if protein sequence in amino acid (`AA`) and their corresponding coding sequence in

nucleotides (`NT`) are aligned in the same run, CAPTUS aligns the `AA` with MAFFT and then

uses the `AA` alignment as template for the `NT`, thus producing a codon-aware alignment for the

CDS. CAPTUS also allows the user to provide the sample(s) that should be considered as

outgroup, in this case the program will place those samples as the first sequence(s) in the

alignments in the order provided. This feature (`--outgroup`) takes advantage of a common

feature of many phylogenetic estimation programs (e.g., IQ-TREE, RAxML, MRBAYES)

which arbitrarily draw the first sample in the alignment at the root of their output trees.

398   Once the FASTA files are aligned, paralogs are filtered using two alternative

algorithms, `naive` and `informed`. In the `naive` filter, only the best hit (i.e., most similar

and complete as compared to reference target sequence, with suffix `_00` in output files) is

kept for each sample and no further filtering is performed. The `informed` filtering algorithm

takes advantage of reference datasets that contain multiple sequences per locus, such as the

Angiosperms353 (Johnson et al. 2019) or Mega353 (McLay et al. 2021) target files as well as

the ones developed for CAPTUS (SeedPlantsPTD and SeedPlantsMIT). For example, the

Angiosperms353 reference target set (as well as its expansion Mega353) was derived from the

1KP Project data (Matasci et al. 2014; One Thousand Plant Transcriptomes Initiative 2019),

where each gene could potentially be present in more than 1,000 species. To build a less

redundant sequence collection while still covering the entire phylogenetic diversity of the

angiosperms, each selected locus was clustered at ~70% identity, usually selecting only one

representative sequence per cluster for the final reference dataset. Thanks to this feature, one

could expect that when analyzing a single taxonomic group such as a family or genus, all

samples would have a best match to only one sequence in the reference targets collection for a

given locus (i.e., the closest relative present in the reference targets). However, a few samples

could have better matches to a different reference sequence than the rest which can be

possible when the locus in question has multiple copies (paralogs) in the studied group, and

the most common copy found in the group is absent from those few samples. In these cases,

CAPTUS compares all recovered paralogs with the reference sequence that best matches most

of the samples and keeps the copy that is most similar to that reference sequence across

samples. Finally, it could also happen that a sample presents a single copy for a specific locus

but with a sequence that is much more divergent than the average in the alignment (e.g., a

remote paralog found in a contaminated contig). By default, CAPTUS will remove any

sequence with average pairwise identity that is more than 4.0 standard deviations below the

mean pairwise identity of the entire alignment, the number of standard deviations can be

changed with the option `--tolerance`.

Recently developed coalescent methods, such as ASTRAL-PRO (Zhang et al. 2020;

Zhang and Mirarab 2022), are capable of analyzing trees of genes with paralogs (i.e., multi-

copy genes), therefore CAPTUS also provides the alignments with paralogs as well as the file

required by ASTRAL-PRO for mapping the paralog names to the names of the samples for

species tree calculation, this file is unique and can be used with any of the alignment sets

produced by CAPTUS.

The reference target sequences are then removed from the alignments. Finally, all the

produced alignments and their filtered versions are trimmed using CLIPKIT (Steenwyk et al.

2020), which can remove alignment columns based on criteria like informativeness or

proportion of missing data. By default, CAPTUS removes columns with > 90% missing data

and sequences with < 40% mean coverage. Alternatively, a minimum number of sites per

column instead of a percentage can be specified with `--min_data_per_column`. As in

previous modules, CAPTUS computes alignment statistics as well as sample occupancy

statistics from all the FASTA files along all filtering and trimming stages and produces a

comprehensive HTML report that can be used to determine outlier markers or samples that

should be removed or curated more carefully before proceeding to phylogenetic analysis.

For the Cucurbitales data, all extracted loci were aligned in CAPTUS using MAFFT's

most accurate algorithm E-INS-i (`--align_method mafft_genafpair`). For coding

regions we used the option `-f AA,NT` to take advantage of codon-aware alignments for

CDS, for Plastome38 we used `-f MA`. For trimming, we used `--`

`min_data_per_column 6`, to keep alignment columns with six or more sequences. For

the Plastome38 we also decreased the tolerance of the `informed` paralog filter to `--`

`tolerance 2.0`.

449   *Phylogenetic Analyses*

We chose the trimmed, codon-aligned coding sequences (format `NT`) from the

extractions performed with reference targets sets Angiosperms353, Mega353 and RNA5435.

For the reference targets set Plastome38 (the plastome segments) we selected the format

`matches` from the CAPTUS output. We analyzed the unfiltered alignments (i.e., including

paralogs) as well as the alignments resulting from the `naive` and `informed` paralog

filtering strategies for nuclear markers but only the alignments filtered by the `informed`

algorithm for the plastome.

Gene tree estimation.--- For each individual nuclear gene alignment we inferred a phylogeny

using IQ-TREE v. 2.2.2.6 (Minh et al. 2020b). During the run we used MODELFINDER

(Kalyaanamoorthy et al. 2017) to determine the most appropriate nucleotide substitution

model with `-m TEST`. Nodal support was inferred from 1,000 ultrafast bootstrap replicates

with option `-bb 1000` (Hoang et al. 2018). For comparison purposes, we also estimated

nuclear gene tree phylogenies using FASTTREE v. 2.1.11 (Price et al. 2010) with the manual

recommended options to increase the accuracy of the search and using the GTR substitution

model (`-pseudo -spr 6 -mlacc 3 -slownni -gtr`).

Species tree estimation.--- We estimated species trees using two methods, concatenation of

alignments followed by maximum likelihood estimation in IQ-TREE v. 2.2.2.6 (Minh et al.

2020b), and coalescent estimation by summarization of quartet frequencies in gene trees using

ASTRAL-PRO v. 1.15.1.3 (Zhang et al. 2020; Zhang and Mirarab 2022). For concatenation,

each individual locus alignment must contain a single sequence per sample (no paralogs

allowed), therefore we can only apply this method to the alignments that were filtered for

paralogs, while ASTRAL-PRO was applied to the filtered alignments as well as to the

alignments containing multiple copies. The concatenation analyses in IQ-TREE were run with

the same options as for the individual gene trees, and the loci alignments were provided as

separate files in a single directory with the option `-p`, so IQ-TREE can automatically

concatenate them into a supermatrix prior to analysis. The concatenation method was applied

only to the set 353 nuclear genes (extracted using Angiosperm353 or Mega353), to the

plastome segments (Plastome38).

For the ASTRAL-PRO analysis, the individual gene trees calculated by IQ-TREE or

FASTTREE were provided as well as the required file that maps the paralog names to their

corresponding sample name produced by CAPTUS (`captus-`

`assembly_align.astral-pro.tsv`). We also increased the number of placement and

subsampling rounds to 16 (`-R`), as well as the proportion of taxa subsampled (`--`

`proportion 0.75`) and calculated alternative quartet frequencies (`-u 3`).

Site concordance analyses.--- In order to measure concordance between alignment sites and

the species tree, we performed a concordance factor analysis for each species tree (Minh et al.

2020a). The analysis was done in IQ-TREE v. 2.2.2.6 by supplying the species tree (`-t`), the

folder containing the loci alignments (`-p`), and averaging the site concordance over 1,000

quartets (`--scfl 1000`).

Phylogenetic conflict analysis of selected nodes.--- Contentious relationships in the

Cucurbitales are centered around the relationships among tribes in the Cucurbitaceae as well

as the relationships among Cucurbitales families. In order to visualize the amount of conflict

at selected nodes in the species tree, we used the branch quartets frequencies analysis in

DISCOVISTA v. 1.0 (Sayyari et al. 2018), after annotating the species belonging to each

Cucurbitaceae tribe, to each of the families in Cucurbitales, and to the clades in the outgroup.

Phylogenetic network estimation.--- We concatenated the Angiosperms353 alignments

filtered by the `informed` method in CAPTUS keeping only the Cucurbitales families. This

concatenated alignment was used as input for SPLITSTREE v. 4.18.2 (Huson and Bryant 2006)

in order to estimate a phylogenetic network using the Neighbor Net algorithm on a matrix of

uncorrected P distances.

Mitochondrial relationships in Cucurbitales.--- The previous phylogenetic placement of the

Apodanthaceae as sister to the remaining Cucurbitales was based on the signal of a *mat*R

dataset (Filipowicz and Renner 2010, Schaefer and Renner 2011b). To compare our

contradicting nuclear phylogenomic results with a more comprehensive mitochondrial dataset,

we examined the phylogenetic relationships of all 39 mitochondrial genes found in

Cucurbitales. Since three genes are trans-spliced we segmented those in two or three parts for

a total of 43 mitochondrial gene regions.

We used CAPTUS to test minimum sequence identity thresholds between 75% and

90%, and determined that 83% was the highest threshold at which the *mat*R gene could still

be recovered from the Apodanthaceae samples. Additional mitochondrial genes (e.g., rpl2)

were not recovered for Apodanthaceae above 88% minimum sequence identity.

We extracted the mitochondrial regions from our entire dataset using CAPTUS with a

minimum sequence identity of 83%, minimum coverage of 30%, and allowing a maximum of

two additional copies per locus (`-d SeedPlantsMIT --dna_min_identity 83 --`

`dna_min_coverage 30 --max_paralogs 2`). The alignment was also performed in

CAPTUS using MAFFT's most thorough algorithm (`--align_method`

`mafft_genafpair`).

The trimmed unfiltered alignments were analyzed with IQ-TREE v2.3.6 (Minh et al.

2020b) performing standard model selection by MODELFINDER, with 1,000 ultrafast

bootstraps, and using a fixed seed (`-m TEST -bb 1000 -seed 142857`).

To distinguish between host DNA (due to contamination or perhaps horizontal gene

transfer) and parasite DNA in the Apodanthaceae samples, we analyzed the phylogenetic

affinities of the sequenced gene copies. If we recovered multiple copies per gene, those that

grouped with the known host plants [Fabaceae (Fabales) for *Pilostyles*; *Casearia* (Salicaceae,

Malpighiales) for *Apodanthes*] were considered "host DNA" sequences while extra copies

grouping elsewhere were considered "parasite DNA". If the different copies of all three

Apodanthaceae samples grouped together and each was represented by a single copy only,

they were also considered "parasite DNA". If an Apodanthaceae sample had a single gene

copy but did not cluster with the others, we decided based on sequence similarity: sequences

above 88% identity with the reference were considered as host DNA and below 88% as

parasite DNA. In a few inconclusive cases the sequences were classified as "unclear", usually

when the gene tree presented very low resolution.

Benchmarking

The locus recovery efficiency of CAPTUS on different data types was evaluated against

HYBPIPER (Johnson et al. 2016) by extracting the Angiosperms353 loci (Johnson et al. 2019)

from a selection of 80 samples (20 from each data type), representing all families of

Cucurbitales and all tribes of Cucurbitaceae (Table S3). Since the HYBPIPER workflow does

not include a read cleaning step, we used reads cleaned by CAPTUS as a common input for

both pipelines. A protein target file downloaded from

https://github.com/mossmatters/Angiosperms353 was used as a reference for both pipelines.

CAPTUS was run for the `assemble` module with the aforementioned presets optimized for

each data type, followed by the `extract` module with default settings. HYBPIPER v. 2.1.2

was run using either BLASTx (Camacho et al. 2009) or DIAMOND (Buchfink et al. 2021),

hereafter HYBPIPER-BLASTx and HYBPIPER-DIAMOND respectively, with an identity

threshold of 65% (`--thresh 65`) to equalize with that of CAPTUS. For the genome

skimming samples, HYBPIPER-BLASTx and HYBPIPER-DIAMOND were also run with a

lower coverage cutoff of 1 (`--cov_cutoff 1`) to test whether the default cutoff of 8 might

be too stringent for this data type. The locus recovery efficiency of each pipeline was

evaluated based on three metrics: running time, number of loci recovered, and total CDS

length recovered. To make the running times comparable across pipelines, all commands were

run on the same MacOS X system equipped with a 2.7 GHz Intel Xeon E5 processor with 24

threads and 64 GB RAM, allocating 6 threads per sample and processing 4 samples

concurrently. GNU parallel v. 20230122 (Tange 2021) was used to run HYBPIPER on multiple

samples concurrently and to record the running time spent on each sample in a log file (`--

joblog`). The number of loci recovered and the total CDS length recovered were calculated

from the stats file generated by each pipeline (`captus-

assembly_extract.stats.tsv` from CAPTUS and `seq_lengths.tsv` from

HYBPIPER). Additionally, HYBPIPER was run for the `paralog_retriever` command with

default settings to retrieve the detected paralog sequences. The paralog sensitivity of each

pipeline was evaluated based on the average number of paralogs per recovered locus. The

statistical significance of the differences between pipelines was assessed using the Wilcoxon

rank-sum test with Bonferroni correction.

To investigate the sequencing depth of loci recovered by CAPTUS but not by

HYBPIPER, the clean reads from each sample were mapped back to the gene sequences (exons

+ introns) recovered by CAPTUS using BBMAP v. 39.06 (Bushnell 2022) with settings

"`kfilter=22 subfilter=15 maxindel=80`". The average sequencing depth for

each locus was then calculated using the `pileup.sh` script bundled with BBMAP.

To assess the implications of differences in locus recovery efficiency among pipelines

on phylogenetic inference, we estimated coalescent species trees from the CDS recovered by

each pipeline and compared their topologies and nodal supports. For CAPTUS, three sets of

alignments with different paralog filtering strategies (`unfiltered`, `naive`, and

`informed`) were generated using the `align` module with default settings. For HYBPIPER-

BLASTX and HYBPIPER-DIAMOND, two sets of alignments were generated, one with

paralogs and one without. All alignments, regardless of the pipeline, were generated using

MAFFT v. 7.525 (Katoh and Standley 2013) with the automatic strategy selection mode (`--`

`auto`) and trimmed for columns with >90% missing data using CLIPKIT v. 2.2.4 (Steenwyk

et al. 2020) with the gappy mode (`-m gappy`). Coalescent species trees were estimated from

each set of trimmed alignments using IQ-TREE v. 2.2.2.6 (Minh et al. 2020b) and ASTRAL-

PRO v. 1.15.1.3 (Zhang et al. 2020; Zhang and Mirarab 2022) following the procedure

described above, and then visualized using TOYTREE v. 2.0.5 (Eaton 2020).

588   RESULTS

589   *Sequencing*

From our 118 library preparations for target capture, 92 (78%) worked at the first

attempt. Only eight of these successful libraries needed library concentration or increased

number of PCR cycles to improve their quality prior to sequencing. For the remaining 26

libraries (22%) that had to be repeated, we used material of the same herbarium voucher for

14, while we had to select a different specimen for 12. Among the repeated libraries, only two

needed library concentration or increased number of PCR cycles to improve their quality prior

to sequencing. Nonetheless, once the sequencing was completed, three samples had to be

discarded due to insufficient data. Two of the discarded samples corresponded to replicated

libraries of *Octomeles sumatrana* for which we had a replacement within the target capture

batch. Only one of them, *Bambekea racemosa*, needed to be replaced by additional high-depth

whole genome sequencing, which was also performed for six additional samples (for

previously missing taxa). In the end, a total of 122 samples (115 target capture and 7 high-

depth whole genome sequencing) yielded enough high-quality data to be used for the analyses

as determined by the number of loci recovered and their completeness (Table S4).

605   *Read Cleaning and Assembly*

606   The RNA-Seq data from NCBI had the largest average percentage of reads and

percentage of base pairs removed with 4.7% and 9.2% respectively. The target capture data

had the largest average percentage of raw reads with adaptors (14.6%) but adaptors were fully

removed from all data types after cleaning (Table 1: Cleaning). Average assembly size after

removing contigs with GC content > 60% was highest for high-depth whole genome

sequencing (591.1 Mbp), followed by genome skimming (289.3 Mbp), and both target capture

and RNA-Seq data with c. 44 Mbp. Despite their similar assembly sizes, target capture data

produced more fragmented assemblies than RNA-Seq as indicated by the average number of

contigs (88.3 k vs. 62.7 k respectively) and the average N50 (513.3 bp vs. 1084 bp

respectively), while high-depth whole genome sequencing data reached the largest average

N50 with 5023.7 bp. The GC content matched the expectations of the data type, being lower

for genome skimming (37.5%) and high-depth whole genome sequencing (34.8%), and higher

for the gene-centered datatypes target capture (40.8%) and RNA-Seq (42.6%) (Table 1:

Assembly).

The average GC content of the filtered contigs ranged from 59.8% to 65.4% across data types

(Table 1: Assembly). The target capture data had the highest average percentage of contigs

removed because of exceeding the threshold of 60% GC content (4%).

TABLE 1. Summary statistics by data type (CAP – target capture; GSK – genome skimming;

WGS – high-depth whole genome sequencing; RNA – RNA-seq).

	CAP N = 126		GSK N = 48		WGS N = 38		RNA N = 240	
	Mean	SD	**Mean**	SD	**Mean**	SD	**Mean**	SD
Raw reads (Millions)	**4.7**	2.6	**17.3**	12.1	**113.5**	99.4	**44.1**	33.2
Raw bases (Gbp)	**0.7**	0.4	**2.5**	1.8	**15.9**	12.2	**5.9**	4.4
Cleaning								
Removed reads (%)	**2.3**	2.7	**1.3**	2.3	**2.0**	3.7	**4.7**	10.3
Removed bases (%)	**8.4**	8.9	**4.2**	6.7	**4.3**	8.7	**9.2**	13.1
Raw reads with adaptors (%)	**14.6**	22.0	**5.6**	12.6	**1.8**	6.8	**8.5**	11.3
Clean reads with adaptors (%)	**0.0**	0.0	**0.0**	0.0	**0.0**	0.0	**0.0**	0.0
Assembly								
Contigs (Thousands)	**88.3**	61.1	**489.0**	261.4	**823.7**	654.3	**62.7**	37.9
Assembly size (Mbp)	**44.4**	31.7	**289.3**	162.2	**591.1**	256.3	**44.7**	23.8
N50 (bp)	**513.3**	116.1	**923.3**	964.8	**5023.7**	4384.0	**1084.0**	289.5
GC content (%)	**40.8**	3.6	**37.5**	2.7	**34.8**	1.5	**42.6**	2.1
Filtered contigs (Thousands)	**1.3**	3.1	**6.2**	20.1	**16.2**	49.1	**1.8**	6.1
Filtered contigs (%)	**4.0**	21.0	**1.8**	7.5	**1.5**	3.6	**1.9**	3.4
Filtered contigs GC content (%)	**59.8**	14.6	**64.1**	1.9	**65.4**	2.4	**62.9**	1.0
Extraction								
Angiosperms353								
Percentage of loci recov. (%)	**87.9**	19.3	**70.9**	30.7	**98.8**	4.1	**92.6**	14.0
Total CDS length recov. (kbp)	**183.8**	57.5	**137.2**	89.0	**255.9**	28.5	**229.2**	53.3
Mega353								
Percentage of loci recov. (%)	**88.0**	19.3	**71.3**	30.6	**98.9**	3.9	**92.7**	14.0
Total CDS length recov. (kbp)	**190.2**	60.6	**145.6**	95.1	**271.3**	30.1	**241.8**	57.1
RNA5435								
Percentage of loci recov. (%)	**20.8**	14.9	**61.2**	30.5	**97.4**	8.4	**90.5**	14.8
Total CDS length recov. (kbp)	**808.6**	713.6	**3964.6**	2757.1	**8207.7**	1136.5	**6968.8**	1832.1
SeedPlantsPTD								
Percentage of loci recov. (%)	**78.6**	24.9	**94.9**	5.3	**96.3**	0.7	**76.2**	20.6
Total CDS length recov. (kbp)	**45.9**	20.8	**64.8**	6.7	**67.0**	1.2	**39.9**	16.9
SeedPlantsMIT								
Percentage of loci recov. (%)	**73.9**	26.5	**92.4**	10.9	**93.7**	3.9	**63.8**	21.9
Total CDS length recov. (kbp)	**22.0**	10.6	**31.8**	4.1	**32.2**	0.9	**18.0**	8.2
Plastome38								
Percentage of loci recov. (%)	**88.6**	25.6	**99.0**	5.3	**100.0**	0.0	**81.3**	20.6
Total length recov. (kbp)	**145.3**	63.4	**208.3**	30.6	**211.7**	7.3	**106.6**	45.5

Since the Angiosperms353 and the Mega353 reference targets aim to recover the same

set of 353 genes, gene recovery in terms of number of loci and total CDS length was very

similar for both (Table 1: Extraction). The average percentage of genes recovered across data

types varied from c. 71% for genome skimming to c. 99% for high-depth whole genome

sequencing data while the data type with the longest total CDS lengths is high-depth whole

genome sequencing with an average of 271.3 kbp and the shortest is genome skimming with

an average of 145.6 kbp. However, the Mega353 reference targets recovered at most 0.4%

more loci than Angiosperms353 for genome skimming and at most 0.1% more for other data

types. Similarly, the Mega353 reference targets produced total CDS lengths only c. 10 kbp

longer than the Angiosperms353 reference targets on average across data types.

For the RNA5435 reference targets, target capture data had the lowest average

percentage of recovered loci (20.8%, SD 14.9%) as well as the shortest average total CDS

length (808.6 kbp, SD 713.6 kbp). For the rest of data types, the percentage of RNA5435

genes recovered ranges from 61.2% to 97.4% while the average total CDS length ranges from

3.96 Mbp to 8.2 Mbp. Recovery of organellar proteins was also high across data types,

exceeding 73% except for mitochondrial proteins from RNA-Seq data where only 63.8% of

the genes were recovered. The 38 plastome segments were successfully recovered across data

types, where the minimum was 81.3% for RNA-Seq data. The total aligned ungapped length

across samples was similar for Angiosperms353 (Fig. S2a) and Mega353 alignments (Fig.

S2b) with c. 85% of samples with lengths between c. 200 kbp and c. 290 kbp and only c.

2.5% of samples showing less than 50 kbp aligned. As for RNA5435 (Fig. S2c) around 55%

of samples had total ungapped aligned lengths between 6.0 and 8.2 Mbp, while around 20%

of the samples had less than 1 Mbp aligned, corresponding mostly to target capture samples.

Based on a preliminary extraction report from CAPTUS which indicated an unusually

high number of copies per DNA region in *Lemurosicyos variegata* (Cucurbitaceae)*,* we

concluded that this sample was contaminated with DNA from another plant family. Given the

evolutionary distance of the contaminant, we could decontaminate the *Lemurosicyos*

sequences using CAPTUS and a custom reference specific for Cucurbitaceae (Supplementary

Method). We also identified the Cucurbitaceae sample *Xerosicyos perrieri* as cross-

contaminated with *Ibervillea sonorae*, another Cucurbitaceae. In this case, decontamination

would have implied creating a *Xerosicyos*-specific reference, which was not possible, and

therefore we decided to exclude the *Xerosicyos perrieri* data from further analyses. A

preliminary plastome phylogeny revealed additional samples with chloroplast contamination

or lack of sufficient data for correct placement (Table S5).

Phylogeny estimation

Coalescent species trees estimated from the set of 353 genes (using Angiosperms353

or Mega353 reference targets sets) or the RNA5435 gene trees, regardless if inferred from

paralog-filtered or unfiltered alignments, or if estimated with FASTTREE or IQ-TREE, resulted

in topologies with monophyletic and well supported families in Cucurbitales and

monophyletic and well-supported tribes in Cucurbitaceae (Fig. 2, Fig. S3, Appendix S3).

Species trees calculated with the concatenation method for nuclear genes (Appendix S3) and

for the plastome (Fig. S5) also show monophyletic families and Cucurbitaceae tribes. The

holoparasitic Apodanthaceae are sister to Rafflesiaceae in Malpighiales in all inferred nuclear

species trees (Fig. S3, Appendix S3), due to their holoparasitic life form, we could not recover

reliable plastome data. For the mitochondrial gene regions obtained from the three

Apodanthaceae samples, the results differ remarkably between genes and gene copies (Fig. 3,

Table S6, Appendix S4). Focusing on what we classify "parasite DNA", 16 mitochondrial

regions support a phylogenetic position of Apodanthaceae in Malpighiales, while six regions

show a placement in Cucurbitales and the remaining regions lack a well-supported

phylogenetic position. For the sequences that we classify "host DNA", there is obviously a

strong affinity to the two host clades (28 *Apodanthes* sequences in 27 regions group with

Malpighiales, 25 *Pilostyles* sequences in 20 regions group with Fabales), however, in the

Pilostyles aethiopica data, we find five gene sequences grouping with Cucurbitales and seven

of unclear affinity. This indicates that the Cucurbitales link to Apodanthaceae might be

biologically meaningful and perhaps the result of an ancient host-parasite relationship that has

been lost with the extinction of an ancestral lineage in the family.

FIGURE 2. Coalescent cladogram of the Cucurbitales inferred with ASTRAL-PRO based on the

RNA5435 nuclear gene trees (Fig. S4a) with families and Cucurbitaceae tribes collapsed into

a single branch each. Gene support calculated with ASTRAL-PRO and site support calculated

using IQ-TREE is indicated at each branch, the most frequent alternative topologies are

shown with red broken lines. Branch lengths in the inset phylogram are in coalescent units.

FIGURE 3. Mitochondrial affinities in the Apodanthaceae summarized by genus. The numbers

represent the total number of copies found in the samples.

For the remaining Cucurbitales families, the relationships vary, particularly around the

nodes with ambiguous quartet support surrounding short branches (see inset in Fig. 2).

However, the quartet frequency analyses from DISCOVISTA reveal that the apparently

contradicting relationships among the major clades are all compatible with the reticulations

depicted as dotted red lines in Fig. 2 because the different species tree configurations

observed coincide with the alternative quartet configurations suggested for the nodes in

conflict (Fig. S4). The best supported alternative shows a grade, where the clade Coriariaceae

+ Corynocarpaceae is followed by Anisophylleaceae and by the remaining families, but there

is also considerable gene tree and site support for a clade in which Anisophylleaceae is sister

708 to a clade comprising Corynocarpaceae + Coriariaceae and the remainder of the order (Fig. 2).

709 The woody Tetramelaceae is resolved as sister to the two herbaceous families Begoniaceae

710 and Datiscaceae but again there is also considerable gene support for the two alternative

711 combinations within the triplet (Fig. 2). Within Cucurbitaceae, the best supported topology is

712 a clade comprising Actinostemmateae and Gomphogyneae as sister to the remaining tribes.

713 Among the genera with multiple samples, only *Kedrostis* is found to be polyphyletic (Fig. S3,

714 Appendix S3, Fig. S5). The extinct Cambodian species *Khmeriosicyos harmandii*, known

715 only from the type collection, is confidently placed in Benincaseae as sister to *Borneosicyos*

716 from Mount Kinabalu in the nuclear species trees (Fig. S3, Appendix S3) and in a clade with

717 *Borneosicyos* and the southeast Asian *Solena* in the plastome tree (Fig. S5). The network

718 analysis revealed a deep reticulation within and between Benincaseae and Cucurbiteae and

719 betweeen Schizopeponeae and Sicyoeae. There is also evidence for more recent reticulation

720 within Coniandreae and very recent within Thladiantheae (Fig. 4). The species trees resulting

721 from the concatenated analysis of the complete plastomes (Fig. S5) agree with the reticulated

722 representation of the Cucurbitales topology (Fig. 2, Fig. 4).

725   FIGURE 4. Phylogenetic network estimate for the Cucurbitales calculated with SPLITSTREE

726   using the Neighbor Net algorithm on the concatenated alignment of the Angiosperms353

727   alignments which were filtered using the informed filter of CAPTUS.

729   *Comparison with* HYBPIPER

730   The locus recovery efficiency of CAPTUS with the four different data types was

evaluated against the two methods offered by HYBPIPER based on running time, number of

loci recovered, and total CDS length recovered (Fig. 5a, Table S3). In terms of running time,

HYBPIPER-DIAMOND was the fastest, followed by CAPTUS, and then HYBPIPER-BLASTx

across all tested data types, except for genome skimming, where no significant difference was

found between CAPTUS and HYBPIPER-BLASTx. For the number and length of loci recovered,

CAPTUS consistently recovered more loci and longer sequences than both HYBPIPER-BLASTx

and HYBPIPER-DIAMOND across all tested data types. The difference in locus recovery was

most pronounced in the genome skimming samples, where HYBPIPER-BLASTx and

HYBPIPER-DIAMOND did not recover any loci from nine and eight samples, respectively,

while CAPTUS recovered at least 23.9 kb of CDS from 87 loci (Table S3). These differences

persisted even when the coverage cutoff of HYBPIPER was lowered (Fig. S6). In all tested data

types, the average sequencing depth of loci recovered only by CAPTUS was evidently lower

than that of loci recovered also by HYBPIPER-BLASTx or HYBPIPER-DIAMOND (Fig. S7),

highlighting the sensitivity of CAPTUS to samples with low sequencing depth. CAPTUS also

showed higher sensitivity in paralog detection, recovering more paralogs than HYBPIPER-

BLASTx and HYBPIPER-DIAMOND across all tested data types (Fig. S8). Collectively, these

results demonstrate that although CAPTUS is slower than HYBPIPER-DIAMOND, it can extract

more information for subsequent analyses with less bias across different data types, in a

comparable or shorter time than HYBPIPER-BLASTx.

FIGURE 5. Benchmarking of CAPTUS against HYBPIPER-BLASTX and HYBPIPER-DIAMOND

using a selection of 80 samples belonging to four different data types. **a)** Comparison of locus

recovery efficiency between pipelines for each data type, evaluating the running time required

for the assembly and extraction of the Angiosperms353 loci (top), the number of loci

recovered (middle), and the total length of coding sequences recovered (bottom). Statistical

testing was performed using the Wilcoxon rank-sum test with Bonferroni correction: * $p <$

0.05, ** $p < 0.01$. **b)** Coalescent species trees estimated from CDS alignments filtered for

paralogs, resulting from each pipeline. Vertical bar along each tree color-codes the different

tribes/families in the Cucurbitales. Red diamonds indicate conflicting or poorly supported

nodes with a local posterior probability below 0.9. Eight genome skimming samples

recovered only by CAPTUS are marked with a black dot. One tribe inferred as a polyphyletic

group in the HYBPIPER trees is indicated by its name and position. Species trees resulting

from the alternative paralog filtering strategies are shown in Fig. S9. Tree files in NEWICK

format are provided in Appendix S5.

The differences in locus recovery efficiency between the pipelines are reflected in the

results of phylogenetic inference (Fig. 5b, Fig. S9, Appendix S5). In the coalescent species

trees resulting from CAPTUS, all genera, tribes, and families were recovered as monophyletic

groups, including the genome skimming samples for which HYBPIPER did not recover any

loci. In contrast, one tribe (Schizopeponeae) was recovered as a polyphyletic group in the

trees resulting from HYBPIPER-BLASTx and HYBPIPER-DIAMOND. This polyphyly is likely

due to the misplacement of one genome skimming sample of *Herpetospermum pedunculosum*

(SB13_GSK), for which HYBPIPER-BLASTx and HYBPIPER-DIAMOND recovered only two

and one loci, respectively, whereas CAPTUS recovered 250 loci (Table S3). In terms of nodal

support, the trees resulting from CAPTUS had fewer nodes with low support (local posterior

probability < 0.9) compared to those resulting from HYBPIPER-BLASTx and HYBPIPER-

DIAMOND, suggesting a higher congruence with individual gene trees. Regardless of the

pipeline, the use of paralogs or alternative filtering strategies did not significantly affect the

species trees, except that alternative topologies were adopted for some of the conflicting

nodes (Fig. S9).

783   DISCUSSION

Our analysis of the mixed Cucurbitales sequence data shows that CAPTUS can handle

complex datasets including low-quality and contaminated sequence data and help disentangle

challenging biological phenomena like reticulation and evolution of holoparasites.

In comparison with similar existing pipelines, we find that CAPTUS is slower than

HYBPIPER-DIAMOND but with our dataset recovers a higher number of more complete genes

across a larger number of species than either HYBPIPER-BLASTx or HYBPIPER-DIAMOND

(Fig 4a). The obvious question arises if those additional and longer sequences recovered by

CAPTUS are reliable or maybe chimeric or very low-depth sequences that could result in

misleading topologies. In the case of the Cucurbitales, the tree topologies based on the

CAPTUS alignments seem more reliable than the corresponding topologies from HYBPIPER. In

a more detailed analysis using simulated reads at different sequencing depths we could show

earlier that CAPTUS is able to assemble long and accurate sequences even at low sequencing

depths [Fig. 3 in (Raza et al. 2023)]. Similar to the Cucurbitales results, a higher degree of

agreement among CAPTUS gene trees than for other pipelines in this earlier study [Fig. 6 in

(Raza et al. 2023)] also suggests that the sequence data recovered by CAPTUS is reliable.

The strong performance of CAPTUS can likely be explained by a combination of

factors. First, the *de novo* assembly by MEGAHIT allows the assembly of *all* reads in the

sample in contrast to HYBPIPER which only assembles subsets of prefiltered reads that match

the reference loci, leading to a faster but more restricted assembly potentially missing intronic

regions and small exons and decreasing assembly contiguity in general. Second, SCIPIO's

capacity to reconstruct gene models across several contigs is essential to recover longer loci

from fragmentary assemblies such as the ones resulting from target capture data and

particularly genome skimming (Keller et al. 2008; Hatje et al. 2011). Finally, thanks to its

improved reconstruction of short exons and intron splice sites as well as the ability to

automatically reconstruct genes across multiple contigs, SCIPIO (Keller et al. 2008; Hatje et al.

2011) outperforms EXONERATE (Slater and Birney 2005), the method implemented in

HYBPIPER.

The ability of CAPTUS to successfully combine different data types is confirmed by the

topological results where different kinds of sequence datasets for the same species,

consistently formed well-supported and stable clades (Supplement Trees). Also, by

assembling the entire set of reads in a sample, CAPTUS uses off-target reads efficiently instead

of excluding them a priori like HYBPIPER. Since target enrichment via DNA hybridization is

an imperfect process, it commonly carries over an important amount of off-target data that

can be useful for phylogenomics. Despite targeting only 353 genes, CAPTUS was able to find

almost three times as many nuclear genes (c. 1130 genes on average) using our RNA5435

reference targets, as well as every plastidial and mitochondrial protein and mostly complete

plastomes. Even though the number of off-target reads will greatly depend on hybridization

conditions, these are usually similar for a given set of samples and a pipeline like CAPTUS can

take better advantage of these other genomic regions that would remain unused otherwise.

Regarding the phylogeny estimate, our results show a well-supported nitrogen-fixing

clade where Cucurbitales + Rosales are sister to Fabales + Fagales (Fig. S3, Appendix S3,

Fig. S4). This topology has also been recovered by recent nuclear phylogenomic analyses

(Guo et al. 2021; Zuntini et al. 2024) but differs from phylogenetic estimates mostly based on

chloroplast data (Soltis et al. 1995; The Angiosperm Phylogeny Group 2016; Li et al. 2021)

where the topology (Fabales (Rosales (Cucurbitales, Fagales)) is recovered. Assuming the gain

of nitrogen-fixing ability evolved in the common ancestor of this clade, the net number of

independent losses inside the clade (e.g., Griesmann et al. 2018) should not be affected by this

new topology, however future studies on the subject could benefit by also interpreting gains

and losses according to this new topology. Within the Cucurbitales, all the currently accepted

families are recovered as monophyletic groups in our analyses (Fig. 2, Fig. S3, Appendix S3,

Fig. S4, Fig. S5). The relationships among families agree well with previous phylogenetic and

phylogenomic studies of Cucurbitales, except for the position of the holoparasitic

Apodanthaceae, where we find strong support for a position outside Cucurbitales, in contrast

to the earlier results of Filipowicz and Renner (2010). We show that those earlier results can

be explained by the phylogenetic signal coming from a subset of the mitochondrial genes

(including the previously analyzed *mat*R region, which coincidentally is the most divergent

gene in the mitochondria of Apodanthaceae), while most mitochondrial genes group with

Malpighiales. This affinity of Apodanthaceae with Malpighiales is strongly supported by the

nuclear data and in agreement with our recent result based on a 58% sampling of the 13,600

genera of angiosperms (Zuntini et al. 2024). However, the results of the CAPTUS analyses

indicate a highly supported sister group relationship between the Apodanthaceae and the

Rafflesiaceae (Fig. S3, Appendix S3, Fig. S4) while in our previous angiosperm-wide analysis

(Zuntini et al. 2024) this is not clear. The results of Alzate et al. (2024) with a reduced

taxonomic sampling seem to support the inclusion of Apodanthaceae in Rafflesiaceae but a

deeper phylogenomic analysis of Malpighiales that takes advantage of the increased taxon

sampling in Zuntini et al. (2024) and our increased gene sampling might be required for a

robust test of these relationships.

For the autotrophic families we find most support for a grade where the clade

Coriariaceae + Corynocarpaceae is followed by Anisophylleaceae, a clade with the triplet

Tetramelaceae plus Begoniaceae and Datiscaceae, and by Cucurbitaceae (Fig. 2, Fig. S3a,

Fig. S4). Earlier studies placed Anisophylleaceae as sister to all other except Apodanthaceae

(Zhang et al. 2006; Schaefer and Renner 2011b; Zuntini et al. 2024). Even though our

analyses show the highest support for the clade Coriariaceae + Corynocarpaceae as sister to

the remaining Cucurbitales, there is almost equal support among gene trees for

Anisophylleaceae as sister to the rest (as in previous studies) as well as for a clade

(Anisophylleaceae (Coriariaceae, Corynocarpaceae)) as sister to the rest of families in the

order (Fig. 2, Fig. S3a, Fig. S4). For Datiscaceae, Zhang et al. (2006) and Schaefer and

Renner (2011b) also found a sister group relationship to Begoniaceae, albeit with low support.

Here, we find that such a clade has the highest gene tree support but the alternatives

Datiscaceae sister to Tetramelaceae [recovered by Zuntini et al. (2024)] and Datiscaceae sister

to Begoniaceae + Tetramelaceae also receive almost equal support (Fig. 2, Fig. S4). In the

network (Fig. 4), the two dioecious families Datiscaceae + Tetramelaceae form a clade, which

is sister to the monoecious Begoniaceae.

Tribal relationships within Cucurbitaceae match well the phylotranscriptomics results

of Guo et al. (2020). The four conflicting nodes identified in Bellot et al. (2020) concerning

the position of *Luffa*, *Hodgsonia*, *Bryonia*, and *Indofevillea* are stable when coalescent and

concatenated phylogeny estimates are compared (Fig. S3, Appendix S3): *Indofevillea* is

placed as sister to the Southeast Asian Thladiantheae; the sponge gourds (*Luffa*) are sister to

all other Sicyoeae; the Asian *Hodgsonia* with the Neotropical Sicyoeae are sister to

Trichosanthes; and Bryonieae plus Schizopeponeae are sister to Sicyoeae. Looking at the

gene tree analyses, however, the considerable number of trees supporting alternative positions

of those lineages is evident (Fig. 2, indicated in red), indicating frequent hybridization events

in the evolutionary history of Cucurbitaceae.

CAPTUS allows the disentanglement of the complex pattern of deep reticulated

evolution in Cucurbitales, which seems to be prevalent across the angiosperms (Stull et al.

2023). The comparison of the gene tree frequencies obtained from the 353 captured regions

(Angiosperms353, or the taxonomically expanded Mega353) with the frequencies obtained

from the RNA5435 regions shows stable 1/3:1/3:1/3 ratio for nodes showing incomplete

lineage sorting (e.g., nodes 14, 24, 29, 32, and 41 in Fig. S4a) or 1/2:1/2:0 ratio for nodes of

hybrid origin (e.g. node 7 in Fig. S4a). This indicates that adding more genomic regions is

unlikely to change the family and tribal level relationships in Cucurbitales found in our study.

Future work should focus on the more recent evolutionary history of the clade and

phylogenetic patterns within Cucurbitaceae genera.

In conclusion, we show that our new pipeline can handle a complex phylogenomic

analysis in a very efficient way, maximizing the amount of recovered data in a reasonable

time frame while maintaining outputs organized across a large number of samples and loci.

The clustering capability of CAPTUS enables its application not only to seed plants but to any

taxonomic group, even those where a reference set of orthologous loci has not yet been

developed. Thus, CAPTUS can be used as a user-friendly universal tool for the assembly of

phylogenomic datasets, even with mixed data of different origins and degraded or

contaminated samples.

896   ACKNOWLEDGEMENTS

We are grateful to the curators of the herbarium of Royal Botanic Gardens, Kew (K),

the herbarium of Muséum National d'Histoire Naturelle in Paris (P), and the herbarium of the

Botanische Staatssammlung München (M) for permission to extract DNA from selected

specimens. We also thank A. Tellier (TUM) for access to the Population Genetics HPC and B.

Faircloth and an anonymous reviewer for comments that help to improve the manuscript. The

study was funded by the German Science Foundation DFG - SCHA 1875/4-2 within SPP

1991 Taxon-Omics (to HS), and by grants from the Calleva Foundation to the Plant and

Fungal Trees of Life Project (PAFTOL) at the Royal Botanic Gardens, Kew.

906   REFERENCES

Alzate J.F., González F.A., Pabón-Mora N. 2024. Back together: Over 1000 single-copy
nuclear loci and reproductive features support the holoendoparasitic Apodanthaceae
and Rafflesiaceae as sister lineages in the order Malpighiales. Molecular
Phylogenetics and Evolution. 201:108217.

Andermann T., Cano Á., Zizka A., Bacon C., Antonelli A. 2018. SECAPR—a bioinformatics
pipeline for the rapid and user-friendly processing of targeted enriched Illumina
sequences, from raw reads to alignments. PeerJ. 6:e5175.

Andrews S. 2019. FastQC: A quality control analysis tool for high throughput sequencing
data. Available from https://github.com/s-andrews/FastQC.

Baker W.J., Bailey P., Barber V., Barker A., Bellot S., Bishop D., Botigué L.R., Brewer G.,
Carruthers T., Clarkson J.J., Cook J., Cowan R.S., Dodsworth S., Epitawalage N.,
Françoso E., Gallego B., Johnson M.G., Kim J.T., Leempoel K., Maurin O., Mcginnie
C., Pokorny L., Roy S., Stone M., Toledo E., Wickett N.J., Zuntini A.R., Eiserhardt
W.L., Kersey P.J., Leitch I.J., Forest F. 2022. A Comprehensive Phylogenomic
Platform for Exploring the Angiosperm Tree of Life. Systematic Biology. 71:301–
319.

Bankevich A., Nurk S., Antipov D., Gurevich A.A., Dvorkin M., Kulikov A.S., Lesin V.M.,
Nikolenko S.I., Pham S., Prjibelski A.D., Pyshkin A.V., Sirotkin A.V., Vyahhi N.,
Tesler G., Alekseyev M.A., Pevzner P.A. 2012. SPAdes: A New Genome Assembly
Algorithm and Its Applications to Single-Cell Sequencing. Journal of Computational
Biology. 19:455–477.

Barkman T.J., McNeal J.R., Lim S.-H., Coat G., Croom H.B., Young N.D., dePamphilis C.W.
2007. Mitochondrial DNA suggests at least 11 origins of parasitism in angiosperms
and reveals genomic chimerism in parasitic plants. BMC Evol Biol. 7:248.

Bellot S., Mitchell T.C., Schaefer H. 2020. Phylogenetic informativeness analyses to clarify
past diversification processes in Cucurbitaceae. Sci Rep. 10:488.

Bellot S., Renner S. 2014. The systematics of the worldwide endoparasite family
Apodanthaceae (Cucurbitales), with a key, a map, and color photos of most species.
PhytoKeys. 36:41–57.

Buchfink B., Reuter K., Drost H.-G. 2021. Sensitive protein alignments at tree-of-life scale
using DIAMOND. Nat Methods. 18:366–368.

Bushnell B. 2022. BBTools: A suite of fast, multithreaded bioinformatics tools designed for
analysis of DNA and RNA sequence data. Available from https://jgi.doe.gov/data-and-
tools/software-tools/bbtools/.

Cai L., Zhang H., Davis C.C. 2022. PhyloHerb: A high‐throughput phylogenomic pipeline
for processing genome skimming data. Appl Plant Sci. 10:e11475.

Camacho C., Coulouris G., Avagyan V., Ma N., Papadopoulos J., Bealer K., Madden T.L.
2009. BLAST+: architecture and applications. BMC Bioinformatics. 10:421.

Cerón-Romero M.A., Maurer-Alcalá X.X., Grattepanche J.-D., Yan Y., Fonseca M.M., Katz
L.A. 2019. PhyloToL: A Taxon/Gene-Rich Phylogenomic Pipeline to Explore
Genome Evolution of Diverse Eukaryotes. Molecular Biology and Evolution.
36:1831–1842.

Chase M.W., Soltis D.E., Olmstead R.G., Morgan D., Les D.H., Mishler B.D., Duvall M.R.,
Price R.A., Hills H.G., Qiu Y.-L., Kron K.A., Rettig J.H., Conti E., Palmer J.D.,
Manhart J.R., Sytsma K.J., Michaels H.J., Kress W.J., Karol K.G., Clark W.D.,

Hedren M., Gaut B.S., Jansen R.K., Kim K.-J., Wimpee C.F., Smith J.F., Furnier
G.R., Strauss S.H., Xiang Q.-Y., Plunkett G.M., Soltis P.S., Swensen S.M., Williams
S.E., Gadek P.A., Quinn C.J., Eguiarte L.E., Golenberg E., Learn G.H., Graham S.W.,
Barrett S.C.H., Dayanandan S., Albert V.A. 1993. Phylogenetics of Seed Plants: An
Analysis of Nucleotide Sequences from the Plastid Gene rbcL. Annals of the Missouri
Botanical Garden. 80:528.

Doyle J.J., Doyle J.L. 1987. A rapid DNA isolation procedure for small quantities of fresh
leaf tissue. Phytochemical Bulletin. 19:11–15.

Eaton D.A.R. 2020. Toytree: A minimalist tree visualization and manipulation library for
Python. Methods Ecol Evol. 11:187–191.

Eaton D.A.R., Overcast I. 2020. ipyrad: Interactive assembly and analysis of RADseq
datasets. Bioinformatics. 36:2592–2594.

Edgar R.C. 2022. Muscle5: High-accuracy alignment ensembles enable unbiased assessments
of sequence homology and phylogeny. Nat Commun. 13:6968.

Faircloth B.C. 2016. PHYLUCE is a software package for the analysis of conserved genomic
loci. Bioinformatics. 32:786–788.

Fierst J.L., Murdock D.A. 2017. Decontaminating eukaryotic genome assemblies with
machine learning. BMC Bioinformatics. 18:533.

Filipowicz N., Renner S.S. 2010. The worldwide holoparasitic Apodanthaceae confidently
placed in the Cucurbitales by nuclear and mitochondrial gene trees. BMC Evol Biol.
10:219.

Goodall-Copestake W.P., Harris D.J., Hollingsworth P.M. 2009. The origin of a mega-diverse
genus: dating *Begonia* (Begoniaceae) using alternative datasets, calibrations and
relaxed clock methods. Botanical Journal of the Linnean Society. 159:363–380.

Griesmann M., Chang Y., Liu X., Song Y., Haberer G., Crook M.B., Billault-Penneteau B.,
Lauressergues D., Keller J., Imanishi L., Roswanjaya Y.P., Kohlen W., Pujic P.,
Battenberg K., Alloisio N., Liang Y., Hilhorst H., Salgado M.G., Hocher V., Gherbi
H., Svistoonoff S., Doyle J.J., He S., Xu Y., Xu S., Qu J., Gao Q., Fang X., Fu Y.,
Normand P., Berry A.M., Wall L.G., Ané J.-M., Pawlowski K., Xu X., Yang H.,
Spannagl M., Mayer K.F.X., Wong G.K.-S., Parniske M., Delaux P.-M., Cheng S.
2018. Phylogenomics reveals multiple losses of nitrogen-fixing root nodule symbiosis.
Science. 361:eaat1743.

Guo J., Xu W., Hu Y., Huang J., Zhao Y., Zhang L., Huang C.-H., Ma H. 2020.
Phylotranscriptomics in Cucurbitaceae Reveal Multiple Whole-Genome Duplications
and Key Morphological and Molecular Innovations. Molecular Plant. 13:1117–1133.

Guo X., Fang D., Sahu S.K., Yang S., Guang X., Folk R., Smith S.A., Chanderbali A.S., Chen
S., Liu M., Yang T., Zhang S., Liu X., Xu X., Soltis P.S., Soltis D.E., Liu H. 2021.
Chloranthus genome provides insights into the early diversification of angiosperms.
Nat Commun. 12:6930.

Hatje K., Keller O., Hammesfahr B., Pillmann H., Waack S., Kollmar M. 2011. Cross-species
protein sequence and gene structure prediction with fine-tuned Webscipio 2.0 and
Scipio. BMC Res Notes. 4:265.

Hoang D.T., Chernomor O., von Haeseler A., Minh B.Q., Vinh L.S. 2018. UFBoot2:
Improving the Ultrafast Bootstrap Approximation. Molecular Biology and Evolution.
35:518–522.

Huson D.H., Bryant D. 2006. Application of Phylogenetic Networks in Evolutionary Studies.
Molecular Biology and Evolution. 23:254–267.

Jackson C., McLay T., Schmidt‐Lebuhn A.N. 2023. hybpiper‐nf and paragone‐nf:
Containerization and additional options for target capture assembly and paralog
resolution. Appl Plant Sci. 11:e11532.

Johnson M.G., Gardner E.M., Liu Y., Medina R., Goffinet B., Shaw A.J., Zerega N.J.C.,
Wickett N.J. 2016. HybPiper: Extracting coding sequence and introns for
phylogenetics from high‐throughput sequencing reads using target enrichment.
Applications in Plant Sciences. 4:1600016.

Johnson M.G., Pokorny L., Dodsworth S., Botigué L.R., Cowan R.S., Devault A., Eiserhardt
W.L., Epitawalage N., Forest F., Kim J.T., Leebens-Mack J.H., Leitch I.J., Maurin O.,
Soltis D.E., Soltis P.S., Wong G.K., Baker W.J., Wickett N.J. 2019. A Universal
Probe Set for Targeted Sequencing of 353 Nuclear Genes from Any Flowering Plant
Designed Using k-Medoids Clustering. Systematic Biology. 68:594–606.

Kalyaanamoorthy S., Minh B.Q., Wong T.K.F., von Haeseler A., Jermiin L.S. 2017.
ModelFinder: fast model selection for accurate phylogenetic estimates. Nat Methods.
14:587–589.

Katoh K., Standley D.M. 2013. MAFFT Multiple Sequence Alignment Software Version 7:
Improvements in Performance and Usability. Molecular Biology and Evolution.
30:772–780.

Keller O., Odronitz F., Stanke M., Kollmar M., Waack S. 2008. Scipio: Using protein
sequences to determine the precise exon/intron structures of genes and their orthologs
in closely related species. BMC Bioinformatics. 9:278.

Kent W.J. 2002. BLAT—The BLAST-Like Alignment Tool. Genome Res. 12:656–664.

Kocyan A., Zhang L.-B., Schaefer H., Renner S.S. 2007. A multi-locus chloroplast phylogeny
for the Cucurbitaceae and its implications for character evolution and classification.
Molecular Phylogenetics and Evolution. 44:553–577.

Li D., Liu C.-M., Luo R., Sadakane K., Lam T.-W. 2015. MEGAHIT: an ultra-fast single-
node solution for large and complex metagenomics assembly via succinct de Bruijn
graph. Bioinformatics. 31:1674–1676.

Li D., Luo R., Liu C.-M., Leung C.-M., Ting H.-F., Sadakane K., Yamashita H., Lam T.-W.
2016. MEGAHIT v1.0: A fast and scalable metagenome assembler driven by
advanced methodologies and community practices. Methods. 102:3–11.

Li H.-T., Luo Y., Gan L., Ma P.-F., Gao L.-M., Yang J.-B., Cai J., Gitzendanner M.A.,
Fritsch P.W., Zhang T., Jin J.-J., Zeng C.-X., Wang H., Yu W.-B., Zhang R., Van Der
Bank M., Olmstead R.G., Hollingsworth P.M., Chase M.W., Soltis D.E., Soltis P.S.,
Yi T.-S., Li D.-Z. 2021. Plastid phylogenomic insights into relationships of all
flowering plant families. BMC Biol. 19:232.

Matasci N., Hung L.-H., Yan Z., Carpenter E.J., Wickett N.J., Mirarab S., Nguyen N.,
Warnow T., Ayyampalayam S., Barker M., Burleigh J.G., Gitzendanner M.A., Wafula
E., Der J.P., dePamphilis C.W., Roure B., Philippe H., Ruhfel B.R., Miles N.W.,
Graham S.W., Mathews S., Surek B., Melkonian M., Soltis D.E., Soltis P.S., Rothfels
C., Pokorny L., Shaw J.A., DeGironimo L., Stevenson D.W., Villarreal J.C., Chen T.,
Kutchan T.M., Rolf M., Baucom R.S., Deyholos M.K., Samudrala R., Tian Z., Wu X.,
Sun X., Zhang Y., Wang J., Leebens-Mack J., Wong G.K.-S. 2014. Data access for the
1,000 Plants (1KP) project. GigaSci. 3:17.

McLay T.G.B., Birch J.L., Gunn B.F., Ning W., Tate J.A., Nauheimer L., Joyce E.M.,
Simpson L., Schmidt‐Lebuhn A.N., Baker W.J., Forest F., Jackson C.J. 2021. New
targets acquired: Improving locus recovery from the Angiosperms353 probe set. Appl
Plant Sci. 9:aps3.11420.

Minh B.Q., Hahn M.W., Lanfear R. 2020a. New Methods to Calculate Concordance Factors
for Phylogenomic Datasets. Molecular Biology and Evolution. 37:2727–2733.

Minh B.Q., Schmidt H.A., Chernomor O., Schrempf D., Woodhams M.D., von Haeseler A.,
Lanfear R. 2020b. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic
Inference in the Genomic Era. Molecular Biology and Evolution. 37:1530–1534.

Nachtigall P.G., Kashiwabara A.Y., Durham A.M. 2021. CodAn: predictive models for
precise identification of coding regions in eukaryotic transcripts. Briefings in
Bioinformatics. 22:bbaa045.

One Thousand Plant Transcriptomes Initiative. 2019. One thousand plant transcriptomes and
the phylogenomics of green plants. Nature. 574:679–685.

Price M.N., Dehal P.S., Arkin A.P. 2010. FastTree 2 – Approximately Maximum-Likelihood
Trees for Large Alignments. PLoS ONE. 5:e9490.

Puritz J.B., Hollenbeck C.M., Gold J.R. 2014. *dDocent*: a RADseq, variant-calling pipeline
designed for population genomics of non-model organisms. PeerJ. 2:e431.

Raza M., Ortiz E.M., Schwung L., Shigita G., Schaefer H. 2023. Resolving the phylogeny of
Thladiantha (Cucurbitaceae) with three different target capture pipelines. BMC Ecol
Evo. 23:75.

Renner S.S., Barreda V.D., Tellería M.C., Palazzesi L., Schuster T.M. 2020. Early evolution
of Coriariaceae (Cucurbitales) in light of a new early Campanian (ca. 82 Mya) pollen
record from Antarctica. TAXON. 69:87–99.

Ribeiro C., Oliveira L., Batista R., De Sousa M. 2021. UCEasy: A software package for
automating and simplifying the analysis of ultraconserved elements (UCEs). BDJ.
9:e78132.

Rochette N.C., Rivera‑Colón A.G., Catchen J.M. 2019. Stacks 2: Analytical methods for
paired‑end sequencing improve RADseq‑based population genomics. Mol Ecol.
28:4737–4754.

Sayyari E., Whitfield J.B., Mirarab S. 2018. DiscoVista: Interpretable visualizations of gene
tree discordance. Molecular Phylogenetics and Evolution. 122:110–115.

Schaefer H. 2020. Cucurbit Website. Version 1. Available from www.cucurbit.de.

Schaefer H., Heibl C., Renner S.S. 2009. Gourds afloat: a dated phylogeny reveals an Asian
origin of the gourd family (Cucurbitaceae) and numerous oversea dispersal events.
Proc. R. Soc. B. 276:843–851.

Schaefer H., Renner S.S. 2011a. Cucurbitaceae. In: Kubitzki K., editor. The Families and
Genera of Flowering Plants. X. Flowering Plants: Eudicots. Sapindales, Cucurbitales,
Myrtaceae. Springer, Berlin. p. 112–174.

Schaefer H., Renner S.S. 2011b. Phylogenetic relationships in the order Cucurbitales and a
new classification of the gourd family (Cucurbitaceae). Taxon. 60:122–138.

de Sena Brandine G., Smith A.D. 2021. Falco: high-speed FastQC emulation for quality
control of sequencing data. F1000Res. 8:1874.

Slater G., Birney E. 2005. Automated generation of heuristics for biological sequence
comparison. BMC Bioinformatics. 6:31.

Soltis D.E., Soltis P.S., Morgan D.R., Swensen S.M., Mullin B.C., Dowd J.M., Martin P.G.
1995. Chloroplast gene sequence data suggest a single origin of the predisposition for
symbiotic nitrogen fixation in angiosperms. Proc. Natl. Acad. Sci. U.S.A. 92:2647–
2651.

Steenwyk J.L., Buida T.J., Li Y., Shen X.-X., Rokas A. 2020. ClipKIT: A multiple sequence
alignment trimming software for accurate phylogenomic inference. PLoS Biol.
18:e3001007.

Steinegger M., Söding J. 2018. Clustering huge protein sequence sets in linear time. Nat
Commun. 9:2542.

Stevens P.F. 2001. Angiosperm Phylogeny Website. Version 14, July 2017 [and more or less
continuously updated since]. Available from
http://www.mobot.org/MOBOT/research/APweb/.

Stull G.W., Pham K.K., Soltis P.S., Soltis D.E. 2023. Deep reticulation: the long legacy of
hybridization in vascular plant evolution. The Plant Journal.

Tange O. 2021. GNU Parallel 20220422 ('Буча'). Zenodo.

The Angiosperm Phylogeny Group. 2016. An update of the Angiosperm Phylogeny Group
classification for the orders and families of flowering plants: APG IV. Bot. J. Linn.
Soc. 181:1–20.

Zhang C., Mirarab S. 2022. ASTRAL-Pro 2: ultrafast species tree reconstruction from multi-
copy gene family trees. Bioinformatics. 38:4949–4950.

Zhang C., Scornavacca C., Molloy E.K., Mirarab S. 2020. ASTRAL-Pro: Quartet-Based
Species-Tree Inference despite Paralogy. Molecular Biology and Evolution. 37:3292–
3307.

Zhang L.-B., Simmons M.P., Kocyan A., Renner S.S. 2006. Phylogeny of the Cucurbitales
based on DNA sequences of nine loci from three genomes: Implications for
morphological and sexual system evolution. Molecular Phylogenetics and Evolution.
39:305–322.

Zhang L.-B., Simmons M.P., Renner S.S. 2007. A phylogeny of Anisophylleaceae based on
six nuclear and plastid loci: Ancient disjunctions and recent dispersal between South
America, Africa, and Asia. Molecular Phylogenetics and Evolution. 44:1057–1067.

Zuntini A.R., Carruthers T., Maurin O., Bailey P.C., Leempoel K., Brewer G.E., Epitawalage
N., Françoso E., Gallego-Paramo B., McGinnie C., Negrão R., Roy S.R., Simpson L.,
Toledo Romero E., Barber V.M.A., Botigué L., Clarkson J.J., Cowan R.S., Dodsworth
S., Johnson M.G., Kim J.T., Pokorny L., Wickett N.J., Antar G.M., DeBolt L.,
Gutierrez K., Hendriks K.P., Hoewener A., Hu A.-Q., Joyce E.M., Kikuchi I.A.B.S.,
Larridon I., Larson D.A., De Lírio E.J., Liu J.-X., Malakasi P., Przelomska N.A.S.,
Shah T., Viruel J., Allnutt T.R., Ameka G.K., Andrew R.L., Appelhans M.S., Arista
M., Ariza M.J., Arroyo J., Arthan W., Bachelier J.B., Bailey C.D., Barnes H.F.,
Barrett M.D., Barrett R.L., Bayer R.J., Bayly M.J., Biffin E., Biggs N., Birch J.L.,
Bogarín D., Borosova R., Bowles A.M.C., Boyce P.C., Bramley G.L.C., Briggs M.,
Broadhurst L., Brown G.K., Bruhl J.J., Bruneau A., Buerki S., Burns E., Byrne M.,
Cable S., Calladine A., Callmander M.W., Cano Á., Cantrill D.J., Cardinal-McTeague
W.M., Carlsen M.M., Carruthers A.J.A., De Castro Mateo A., Chase M.W., Chatrou
L.W., Cheek M., Chen S., Christenhusz M.J.M., Christin P.-A., Clements M.A.,
Coffey S.C., Conran J.G., Cornejo X., Couvreur T.L.P., Cowie I.D., Csiba L.,
Darbyshire I., Davidse G., Davies N.M.J., Davis A.P., Van Dijk K., Downie S.R.,
Duretto M.F., Duvall M.R., Edwards S.L., Eggli U., Erkens R.H.J., Escudero M., De
La Estrella M., Fabriani F., Fay M.F., Ferreira P.D.L., Ficinski S.Z., Fowler R.M.,
Frisby S., Fu L., Fulcher T., Galbany-Casals M., Gardner E.M., German D.A.,
Giaretta A., Gibernau M., Gillespie L.J., González C.C., Goyder D.J., Graham S.W.,
Grall A., Green L., Gunn B.F., Gutiérrez D.G., Hackel J., Haevermans T., Haigh A.,
Hall J.C., Hall T., Harrison M.J., Hatt S.A., Hidalgo O., Hodkinson T.R., Holmes
G.D., Hopkins H.C.F., Jackson C.J., James S.A., Jobson R.W., Kadereit G.,
Kahandawala I.M., Kainulainen K., Kato M., Kellogg E.A., King G.J., Klejevskaja B.,
Klitgaard B.B., Klopper R.R., Knapp S., Koch M.A., Leebens-Mack J.H., Lens F.,
Leon C.J., Léveillé-Bourret É., Lewis G.P., Li D.-Z., Li L., Liede-Schumann S.,
Livshultz T., Lorence D., Lu M., Lu-Irving P., Luber J., Lucas E.J., Luján M., Lum
M., Macfarlane T.D., Magdalena C., Mansano V.F., Masters L.E., Mayo S.J., McColl
K., McDonnell A.J., McDougall A.E., McLay T.G.B., McPherson H., Meneses R.I.,
Merckx V.S.F.T., Michelangeli F.A., Mitchell J.D., Monro A.K., Moore M.J., Mueller
T.L., Mummenhoff K., Munzinger J., Muriel P., Murphy D.J., Nargar K., Nauheimer
L., Nge F.J., Nyffeler R., Orejuela A., Ortiz E.M., Palazzesi L., Peixoto A.L., Pell
S.K., Pellicer J., Penneys D.S., Perez-Escobar O.A., Persson C., Pignal M., Pillon Y.,
Pirani J.R., Plunkett G.M., Powell R.F., Prance G.T., Puglisi C., Qin M., Rabeler
R.K., Rees P.E.J., Renner M., Roalson E.H., Rodda M., Rogers Z.S., Rokni S.,
Rutishauser R., De Salas M.F., Schaefer H., Schley R.J., Schmidt-Lebuhn A.,
Shapcott A., Al-Shehbaz I., Shepherd K.A., Simmons M.P., Simões A.O., Simões
A.R.G., Siros M., Smidt E.C., Smith J.F., Snow N., Soltis D.E., Soltis P.S., Soreng
R.J., Sothers C.A., Starr J.R., Stevens P.F., Straub S.C.K., Struwe L., Taylor J.M.,

Telford I.R.H., Thornhill A.H., Tooth I., Trias-Blasi A., Udovicic F., Utteridge
T.M.A., Del Valle J.C., Verboom G.A., Vonow H.P., Vorontsova M.S., De Vos J.M.,
Al-Wattar N., Waycott M., Welker C.A.D., White A.J., Wieringa J.J., Williamson
L.T., Wilson T.C., Wong S.Y., Woods L.A., Woods R., Worboys S., Xanthos M.,
Yang Y., Zhang Y.-X., Zhou M.-Y., Zmarzty S., Zuloaga F.O., Antonelli A., Bellot S.,
Crayn D.M., Grace O.M., Kersey P.J., Leitch I.J., Sauquet H., Smith S.A., Eiserhardt
W.L., Forest F., Baker W.J. 2024. Phylogenomics and the rise of the angiosperms.
Nature. 629:843–850.

1165

1166 **Supplementary Material**

1167 SUPPLEMENTARY METHOD. Decontamination process for target capture sample *Lemurosicyos*

1168 *variegata*.

1169

TABLE S1. List of samples sequenced for this study.

TABLE S2. List of samples with public data used in this study.

TABLE S3. Per-sample comparison of running time, number of loci recovered, and total CDS

length recovered among CAPTUS, HYBPIPER-BLASTX, and HYBPIPER-DIAMOND.

TABLE S4. Summary statistics per sample at each analysis step.

TABLE S5. Samples lacking sufficient chloroplast data for phylogenetic estimation or with

chloroplast contamination.

TABLE S6. Mitochondrial affinities in the Apodanthaceae. The number of copies found is

indicated next to the letter abbreviations for the species. A=*Apodanthes caseariae*,

Pa=*Pilostyles aethiopica*, Pt=*Pilostyles thurberi*.

FIGURE S1. The CAPTUS output formats, **a)** outputs available when a protein extraction (`NUC,`

`PTD, MIT`) is performed, **b)** outputs available when a miscellaneous DNA extraction (`DNA,`

`CLR`) is performed.

FIGURE S2. Total aligned ungapped length per sample using different nuclear reference targets

sets, **a)** Angiosperms353, **b)** Mega353, **c)** RNA5435.

FIGURE S3. Coalescent species trees estimated with ASTRAL-PRO using gene trees estimated

by IQ-TREE on alignments derived from different reference targets sets (RNA5435,

Angiosperms353, Mega353) and paralog filtering strategies (`unfiltered, informed,`

`naive`).

FIGURE S4. DISCOVISTA analyses of relative quartet frequencies imposed on the coalescent

ASTRAL-PRO topology shown in Fig. S3a using sets of gene trees derived from the different

reference target sets (RNA5435, Angiosperms353, Mega353) calculated by different

programs (IQ-TREE, FASTTREE) and filtered for paralogs (`informed`, `naive`).

FIGURE S5. Species tree estimated by IQ-TREE on the concatenated alignments derived from

the Plastome38 reference targets set.

FIGURE S6. Locus recovery efficiency at different coverage cutoffs of HYBPIPER on 20

genome skimming samples selected for benchmarking. Results from CAPTUS are shown for

comparison.

FIGURE S7. Distribution of sequencing depth for loci recovered by CAPTUS. Loci are classified

according to whether they were recovered only by CAPTUS, or also by HYBPIPER-BLASTX or

HYBPIPER-DIAMOND. Vertical dashed lines indicate the median value for each series.

FIGURE S8. Comparison of paralog sensitivity between pipelines for each data type. Statistical

testing was performed using the Wilcoxon rank-sum test with Bonferroni correction: $* p <$

0.05, $** p < 0.01$.

FIGURE S9. Coalescent species trees of 80 samples selected for the benchmarking, estimated

from CDS alignments filtered or unfiltered for paralogs. Vertical bar along each tree color-

codes the different tribes/families in the Cucurbitales. Red diamonds indicate conflicting or

poorly supported nodes with a local posterior probability below 0.9. Eight genome skimming

samples recovered only by CAPTUS are marked with a black dot. One tribe inferred as a

polyphyletic group in the HYBPIPER trees is indicated by its name and position. Tree files in

NEWICK format are provided in Appendix S5.

1211

APPENDIX S1. Plastome segments used as reference targets provided as a file in FASTA

format `Plastome38.fasta`

APPENDIX S2. Newly found nuclear putative homologs used as reference targets provided as a

file in FASTA format `RNA5435.fasta`

APPENDIX S3. Estimated species trees in NEWICK format.

APPENDIX S4. Alignments and trees of the mitochondrial regions in NEXUS format.

APPENDIX S5. Species trees in NEWICK format estimated for the benchmarking.

APPENDIX S6. Reference targets sets used in the decontamination of *Lemurosicyos variegata*

(Supplementary Method).

1221