Title: *Selection leads to false inferences of introgression using popular methods*

Authors: Megan L. Smith[1,2] and Matthew W. Hahn[2,3]

[1]Department of Biological Sciences, Mississippi State University, Starkville, MS 39762, USA

[2]Department of Biology, Indiana University, Bloomington, IN 47405, USA

[3]Department of Computer Science, Indiana University, Bloomington, IN 47405, USA

Running Title: Selection misleads inferences of migration

Key Words: geneflow, migration, background selection, selective sweeps

Correspondence: ms4438@msstate.edu (M.L. Smith)

Abstract

Detecting introgression between closely related populations or species is a fundamental objective

in evolutionary biology. Existing methods for detecting migration and inferring migration rates

from population genetic data often assume a neutral model of evolution. Growing evidence of

the pervasive impact of selection on large portions of the genome across diverse taxa suggests

that this assumption is unrealistic in most empirical systems. Further, ignoring selection has

previously been shown to negatively impact demographic inferences (e.g., of population size

histories). However, the impacts of biologically realistic selection on inferences of migration

remain poorly explored. Here, we simulate data under models of background selection, selective

sweeps, balancing selection, and adaptive introgression. We show that ignoring selection

sometimes leads to false inferences of migration in popularly used methods that rely on the site

frequency spectrum (SFS). Specifically, balancing selection and some models of background

selection result in the rejection of isolation-only models in favor of isolation-with-migration

models and lead to elevated estimates of migration rates. BPP, a method that analyzes sequence

data directly, showed false positives for all conditions at recent divergence times, but balancing

selection also led to false positives at medium divergence times. Our results suggest that such

methods may be unreliable in some empirical systems, such that new methods that are robust to

selection need to be developed.

Article Summary

Detecting migration between closely related populations is a central objective in many evolutionary biology studies. However, popular methods for detecting migration assume a simplified model of evolution. Here, we evaluate the impacts of biologically realistic natural selection, recombination, and mutation on three methods for detecting migration. We find that biological complexity leads to false inferences of migration, suggesting that results should be interpreted with caution and that new methods are needed to make robust inferences of migration across empirical systems.

Introduction

In recent years, as genomic data have become readily available for many taxa, evidence of
introgression has accumulated across the tree of life (Mallet *et al.* 2016). A growing interest in
understanding the role of introgression in diversification has led to the development of numerous
phylogenetic methods for detecting introgression (reviewed in Hibbins and Hahn 2022), but most
of these methods cannot detect introgression between sister taxa—only methods that use
population genetic data attempt to do this. While detecting introgression between sister taxa is a
difficult task, it is of central interest to many researchers. For example, understanding whether
closely related taxa exchanged genes during divergence is central to distinguishing among modes
of speciation (Payseur and Rieseberg 2016; Roux *et al.* 2016), with evidence of gene flow
between closely related taxa sometimes being interpreted as a possible signal of sympatric
speciation. Characterizing gene flow between sister species is also essential for developing null
models in scans for selection (Williamson *et al.* 2005; Nielsen *et al.* 2007; Excoffier *et al.* 2009;
Luqman *et al.* 2021). Thus, despite the difficulties of the task, many population genetic methods
have been developed (and have been widely applied) to detect gene flow between sister taxa.

Introgression should lead to increased allele-sharing between taxa and increased variance in
coalescence times compared to incomplete lineage sorting (ILS) alone, and methods to detect
introgression between sister taxa rely on these expectations. Summary-statistic methods aim to
detect particular regions of the genome that have introgressed based on the expectation that these
regions should be more similar between sister taxa than non-introgressed regions (Joly *et al.*
2009; Geneva *et al.* 2015; Rosenzweig *et al.* 2016). Other approaches focus on comparing
models with and without migration and/or estimating genome-wide migration rates. For

example, many site frequency spectrum (SFS)-based methods estimate migration rates and other

parameters by finding the parameters that maximize the composite likelihood of the SFS (e.g.,

Gutenkunst *et al.* 2009; Tellier *et al.* 2011; Excoffier *et al.* 2013), which can be computed using

diffusion approximation (e.g., ∂a∂i; Gutenkunst *et al.* 2009) or simulations (e.g., fastsimcoal2;

Excoffier *et al.* 2013). Models with and without migration can then be compared based on

estimated likelihoods. While powerful, SFS-based approaches do not take advantage of linkage

information, and other approaches exist that directly analyze sequence data rather than relying on

the SFS as a summary. For example, BPP estimates divergence times and the intensities of

introgression events from sequence data under the multispecies coalescent with introgression

(MSci) model a using Bayesian Markov chain Monte Carlo (MCMC) approach (Flouri *et al.*

2020). While these methods are powerful and can be highly accurate on simulated datasets, all

assume selection does not affect the patterns observed. Several programs attempt to relax this

assumption by allowing for variation in effective population sizes across loci, which should

mimic some effects of linked selection (e.g., Tine *et al.* 2014; Roux *et al.* 2016; Rougeux *et al.*

2017); however, the efficacy of such programs has not been tested in a wide variety of settings.

Mounting evidence suggests that selection impacts large portions of the genome (reviewed in

Cutter and Payseur 2013; Kern and Hahn 2018). Notably, selection can produce genomic signals

that mimic demographic processes. For example, linked selection can produce signals that mimic

population growth or contraction and can mislead commonly used methods for inferring

population size histories (Ewing and Jensen 2016; Schrider *et al.* 2016; Johri *et al.* 2021).

Ignoring selection may also pose a substantial problem for methods aiming to detect migration

(Cruickshank and Hahn 2014; Mathew and Jensen 2015; Roux *et al.* 2016; Fraïsse *et al.* 2021).

Selection leads to increased heterogeneity in levels of divergence among loci by either

decreasing (directional selection) or increasing (balancing selection) levels of polymorphism at

some loci. Furthermore, balancing selection may maintain polymorphisms for extended periods

of time, leading to shared polymorphisms between otherwise diverged species. Since many

methods for detecting introgression rely on these same signals, this can lead to false inferences of

migration (e.g., Cruickshank and Hahn 2014; Roux *et al.* 2016). Despite more widespread

acknowledgement of the role of selection and the shortcomings of neutral assumptions in recent

years, methods for inferring migration rates that ignore selection are still widely used.

Here, we simulate data under several evolutionary models that include selection, including

background selection, selective sweeps, balancing selection, and adaptive introgression. We

evaluate the impact of selection on inferences of migration rates in ∂a∂i, fastsimcoal2, and BPP,

and show that some types of selection lead to high rates of false inferences of migration. Our

results highlight the importance of incorporating selection into tests for migration.

Materials and Methods

Simulations

We simulated two populations that diverged at a set time in the past, T_D, and considered two

migration histories: no migration (nomig) and a pulse of migration from population 1 to

population 2 looking forward in time (p1_p2). We set all population sizes to 125,000 and

considered three divergence times: $T_D = 0.25 \times 4N, \ 1 \times 4N,$ and $4 \times 4N$ (low, medium, high).

The time since introgression, T_M, was drawn from a vector $\{0.01 \times T_D, 0.05 \times T_D, 0.10 \times$

$T_D, 0.15 \times T_D, \dots, 0.9 \times T_D\}$ and the probability of any lineage migrating, p_M, was drawn from a

vector $\{0.05, 0.1, 0.15, ..., 0.95\}$. We set the per site mutation rate, μ, to 1e-8, and the per site

recombination rate, r, to 5e-8, both per generation. To lessen the computational burden of

forward-in-time simulations, we scaled all simulations by an order of 100: population sizes were

scaled to 1250, and mutation rates, recombination rates, and selection coefficients (s; see below)

were all scaled to keep values of $N\mu$, Nr, and Ns constant. Similarly, divergence times and the

timing of migration were scaled down by an order of 100. To verify that scaling did not bias our

results, we also simulated a small number of datasets scaling only by an order of 10 (see

Results). We simulated 10,000 independent 10 kb windows for most conditions (see below for

details).

To evaluate the impact of selection on inferences of migration rates, we simulated data under six

scenarios in SLiM v4.0.1 (Haller and Messer 2019), overlaying neutral mutations with pyslim

v1.0.3 and tskit v0.5.5 (Kelleher et al., 2018; Haller et al., 2019). We considered the following

selective scenarios: 1) a neutral model; 2) background selection (BGS); 3) a selective sweep in

the ancestor of the two populations; 4) a selective sweep in population 1; 5) balancing selection

in both populations and their ancestral population; and 6) adaptive introgression. For all

scenarios except adaptive introgression, we considered both migration models described above

(nomig and p1_p2). For adaptive introgression, we considered only the p1_p2 model. For all

simulations, ancestral neutral variation was added via recapitation in pyslim. Each condition is

described in detail below:

1) Neutral model: To simulate data in the absence of selection, we overlay all mutations on

recorded tree sequences with pyslim.

2) Background selection ("BGS"): To simulate under a model of BGS, we simulated 75% of

mutations as deleterious and 25% as selectively neutral. Deleterious mutations had a

dominance value of 0.25, corresponding to partially recessive mutations. Selection

coefficients for deleterious mutations were drawn from a gamma distribution with a mean

and shape of -0.000133 and 0.35 (pre-scaling; mean of -0.0133 post-scaling), corresponding

to estimates from *Drosophila* (Huber *et al.* 2017; Schrider 2020). With the population sizes

used in our simulations, this corresponds to a mean $2Ns$ = -33.25. We included a burn-in

period in which background selection was acting for 25000 generations (post-scaling) prior

to population splitting.

3) Selective sweep in the ancestral population ("sweep ancestor"): When simulating a selective

sweep in the ancestor of the two populations, the selection coefficient was drawn from a

uniform (0.001, 0.005) distribution pre-scaling (0.1, 0.5, post-scaling) with a dominance of

1. At generation 1, a single selectively advantageous mutation was introduced into the

ancestral population at position 5000 (i.e. the middle of the locus). Then, until generation

1000 (post-scaling), we checked whether the mutation had fixed or been lost. If it had been

fixed, the two populations split at generation 1000 and the simulation proceeded. If the

mutation was lost, we restarted at generation 1 and repeated the procedure until the mutation

fixed.

4) Selective sweep in population 1 ("sweep p1"): Immediately after divergence, a selectively

advantageous mutation with a selection coefficient and dominance as in the linked ancestor

simulation was introduced into population 1 at position 5000. If the mutation was lost before

migration between populations began (or before the end of the simulation in the no

migration model), we restarted the simulation. The mutation had no fitness effect in

population 2.

5) Balancing selection ("balancing"): To simulate under a model of balancing selection, we

introduced a mutation into the common ancestor of population 1 and population 2 at the

beginning of the simulation. We used a mutation effect callback to specify the fitness of this

mutation as 1.5 minus the frequency of the mutation in the corresponding population. Thus,

when the mutation is rare in a population, it is highly beneficial, but when it is common, it

becomes deleterious. Selection should therefore favor maintaining this mutation at an

intermediate frequency. At generation 5000 (post-scaling), the two populations split. If the

mutation was lost prior to the end of the simulation, we restarted the simulation.

6) Adaptive introgression ("adaptive int"): For this model, only one migration direction was

considered (p1_p2). The selection coefficient and dominance were the same as in the model

with a selective sweep in P1, except that the mutation was also advantageous in P2. We did

not require that the advantageous mutation actually introgressed in these simulations.

We also evaluated the effects of a more biologically realistic model by including variation in

mutation rate, recombination rate, and selection coefficients across genomic segments using the

"real BGS-weak CNE" approach described in Schrider (2020). Following Schrider (2020), we

used annotation data from the University of California Santa Cruz (UCSC) Table Browser for the

D. melanogaster genome (release 5 / dm3; Adams *et al.* 2000). We also used the *D.*

melanogaster recombination map from Comeron *et al.* (2012). Briefly, each simulated replicate

was modelled after a randomly selected genomic region with selection coefficients 10-fold

smaller in conserved noncoding elements (CNEs) than in coding regions. We modelled windows

based on the *Drosophila* genome. For each 10 kb window, we selected an endpoint (constrained

to be a multiple of 10 kb). Windows with >= 75% assembly gaps were not allowed, but

otherwise windows were drawn randomly with replacement. The locations of annotated exons

and phastCons elements were recorded, and deleterious mutations occurred only at these sites.

We used recombination rates drawn from the *Drosophila* recombination map for the selected

window. We drew the mutation rate from a uniform (3.445e-9, 3.445e-8) distribution (pre-

scaling) for each simulated region. We refer to these simulations as the 'complex genomic

architecture' condition below.

We simulated 10,000 10-kb regions for the background selection and neutral conditions and

1,500 10-kb regions for each sweep condition and balancing selection. These simulated regions

were used to build datasets for downstream analyses. We sampled 20 diploid individuals (40

chromosomes) total, 10 per population. We then constructed a genotype matrix, discarding sites

that were constant in our sample. We calculated π within each population, F_{ST}, and d_{xy} from tree

sequences using functions from tskit. We also generated alignments using the

generate_nucleotides and convert_alleles functions in pyslim.

These simulated regions were used to construct test datasets (Table 1) for downstream analyses

with fastsimcoal2, ∂a∂i, and BPP. Test datasets for background selection and neutral conditions

consisted of the 10,000 regions simulated under the corresponding condition and model. For the

sweep conditions (sweep p1, sweep ancestor, and adaptive introgression) and balancing

selection, we constructed test datasets by sampling 500 (5%), 1000 (10%), or 1500 (15%)

regions simulated under the corresponding condition and the remainder of datasets (9500, 9000,

or 8500, respectively) from the corresponding set of simulations under a neutral model.

Sampling was conducted independently to generate datasets for analyses with SFS-based

methods (fastsimcoal2 and $\partial a \partial i$) and BPP. For SFS-based methods, we constructed 100 replicate

site frequency spectra (SFS) for each condition by sampling a single SNP per region in the test

dataset (see below for additional detail). For analyses with BPP, for each condition, we generated

a 500-bp alignment for each region included in the test dataset (see below for additional detail).

Comparing models and estimating migration rates in $\partial a \partial i$

To estimate migration rates in $\partial a \partial i$, we constructed SFS for all simulated datasets (54 datasets

per divergence time; Table 1). We built 100 replicate SFS for each dataset, sampling a single bi-

allelic SNP with replacement from each simulated fragment (10,000 per dataset). When

constructing the SFS for $\partial a \partial i$, we did not populate the monomorphic cell.

We estimated migration rates, population sizes, and divergence times using the split_mig model

in $\partial a \partial i$ v.2.3.0 (Gutenkunst *et al.* 2009). This model includes two populations, with sizes V_1 and

V_2 relative to the ancestral population, a divergence time, τ, and two migration rates, M_{12} and

M_{21}. We used starting parameter estimates of 0.1 for V_1 and V_2, 0.01 for M_{12} and M_{21}, and 0.5 *f*or

τ. We set the lower and upper bounds for v_1 and v_2 to 1e-3 and 5, respectively. The lower and

upper bounds for the migration rate parameters were set to 0 and 5, respectively. For τ, the

bounds depended on the divergence time of the simulated dataset being analyzed. The upper and

lower bounds were set to 0.005 and 1 for low divergence, 0.02 and 4 for medium divergence, and

0.04 and 16 for high divergence. We perturbed parameters using the perturb_params function in

∂a∂i. Then, we optimized parameters using the BOBYQA algorithm, the default algorithm in

∂a∂i. We performed a maximum of 400 evaluations.

For all datasets without migration, we also estimated parameters by maximizing the likelihood of

the same model but with migration rates set to zero. Then, we compared the likelihood of the

split_mig model to the likelihood of the model without migration using a likelihood ratio test

(LRT). We calculated the test statistic Δ as:

$\Delta = 2 * (ln\left(split_{mig}\right) - ln\left(no_{mig}\right))$. We then computed the p-value using a χ^2 distribution

with two degrees of freedom, and we rejected the null (no migration) model at a significance

level of 0.01.

A common approach for accommodating background selection is to allow for different effective

population sizes across genomic regions (Roux *et al.* 2016; Rougeux *et al.* 2017). Rougeux *et al.*

(2017) developed an approach to accommodate variation in effective population sizes and

migration rates across loci in ∂a∂i. They allowed two categories of loci with different effective

population sizes for each category, and they allowed for heterogeneous migration across the

genome by considering two categories of loci. We ran a modified version of the scripts from

Rougeux *et al.* (2017) on our high divergence datasets with a complex genomic architecture and

balancing or background selection. We considered five models: SI (strict isolation), SI2N (strict

isolation with variation in effective population sizes across loci), IM (isolation with migration),

IM2N (isolation with migration with variation in effective population sizes across loci), and

IM2m (isolation with migration with variation in migration rates across loci). As above, we

optimized parameters using the BOBYQA algorithm and performed a maximum of 400

evaluations. We also adjusted starting parameters for population sizes and divergence times to

mirror those used above, except in the IM2m model, for which we used starting values for

migration rates of 0.1 and 0.01 for the two different sets of loci. We increased the upper bound

on divergence times from the values used by Rougeux *et al.* (2017) to accommodate the deeper

divergences simulated in our study. We compared the five models using Akaike Information

Criteria (AIC).

Comparing models and estimating migration rates in fastsimcoal2

To estimate migration rates in fastsimcoal2, we used the SFS constructed for $\partial a \partial i$, except that we

included the number of monomorphic sites. To calculate the number of monomorphic sites, we

used the following equation:

$$N_{monomorphic} = \sum_{i=1}^{10000} (10000 - x_i) \times \frac{1}{x_i}$$

where x_i is the number of segregating sites in fragment *i*. This calculation accounts for the fact

that we only sampled a single segregating site per fragment.

We estimated migration rates, population sizes, and divergence times in fastimcoal2 v.2.7.0.9

(Excoffier *et al.* 2013) using a model with the same parameterization as used in $\partial a \partial i$. We set the

mutation rate to the value used in the uniform simulations (1e-8). Note that the units used in

fastsimcoal2 differ from those used in $\partial a \partial i$. We set the minimum bounds for migration rates to

zero, the minimum bounds for population sizes to 1250, and the minimum bounds for the relative

population sizes v_1 and v_2 to 1e-2. The minimum bounds on the divergence time were set to 1250

for low divergence, 5000 for medium divergence, and 10000 for high divergence. We used

100,000 simulations to estimate the expected SFS and performed 40 ECM cycles to estimate

parameters. We also compared the migration model to a model with migration rates set to zero

using an LRT with two degrees of freedom as described above for ∂a∂i.

Estimating migration rates with BPP

We estimated introgression probabilities and divergence times in BPP v4.4.0 (Flouri *et al.* 2020).

Notably, BPP uses the MSci model, which models an instantaneous introgression event, rather

than continuous migration as modelled in our simulations and by ∂a∂i and fastsimcoal2. We first

generated sequence alignments for each region equivalent to a 500-bp locus. We used the middle

500 base pairs of each of the 10,000 fragments composing the test dataset. For each simulation

condition (Table 1), we created 20 replicate datasets with 500 500-bp loci by sampling loci

without replacement from our simulated fragments. In BPP we fixed the species tree to a two-

population tree with introgression allowed in both directions and used an inverse gamma (3,

0.01) prior for the parameters θ and τ. Since the inverse gamma prior is a conjugate prior for θ,

this allowed the θ parameters to be integrated out analytically, improving run times (Hey and

Nielsen 2007). We allowed mutation rates to vary across loci using the a_mubar, b_mubar, and

a_mui priors. We set a_mubar and b_mubar to 0, so that mutation rates were relative. We set

a_mui equal to 2, and we used the iid prior. We also used the heredity scalar to allow for

variation in effective population sizes across loci. For the heredity scalar, we used a Gamma(4,4)

prior. We collected 500,000 samples from the posterior after discarding the first 20,000 samples

as burnin and sampling every 2 iterations. Some runs did not finish in 90 or 96 hours, but we

collected a minimum of 447,060 samples for all runs. These samples were used to estimate all

parameters (using the posterior mean). To assess convergence, we used effective sample size

(ESS) values. To use the results from BPP to provide a binary determination of the presence of

migration, we asked whether the highest posterior density interval (HDI) for migration

parameters included 0. An alternative approach is to use Bayes Factors to compare models with

and without introgression. The Bayes Factor is the ratio of the marginal likelihoods of two

models and requires that we approximate the marginal likelihood of each model. To do this in

BPP, we used a path-sampling approach with 8 steps, which required that we run an MCMC

algorithm over each step for each model. Given that our MCMC runs take up to 96 CPU hours to

complete, this requires ~1,536 CPU hours per dataset. Because of this, we only calculated BF for

a subset of datasets: 5 datasets each from the nomig background selection, nomig neutral, and

p1_p2 background selection datasets.

Results

Selection alters levels of diversity and divergence

Selection altered patterns of diversity and divergence in both expected and initially surprising

ways. Under the model with uniform mutation and recombination rates, the patterns observed in

summary statistics conformed to expectations (Supporting Figures S1-S3). Background selection

and (to a lesser extent) selective sweeps and adaptive introgression reduced nucleotide diversity

relative to the neutral case. Conversely, balancing selection slightly increased diversity.

Background selection also decreased divergence between populations, as expected due to

reductions in ancestral levels of diversity. Migration from population 1 into population 2

increased nucleotide diversity in population 2 and decreased divergence between populations.

Results were more complicated under the model with variation in recombination and mutation

rates. As above, selective sweeps and adaptive introgression reduced nucleotide diversity, and

balancing selection increased nucleotide diversity relative to the neutral case. However, contrary

to our initial expectations, background selection increased nucleotide diversity in many

simulations. Such a pattern might be explained by associative overdominance, which maintains

neutral diversity when strongly linked to partially recessive deleterious mutations, as in our

simulations. Associative overdominance can therefore maintain variation in a population (Ohta

1971; Pamilo and Pálsson 1998; Gilbert *et al.* 2020). To assess whether associative

overdominance could be driving our results, we simulated a smaller number of replicates

(N=1000) with a dominance coefficient of 0.5, which should eliminate the effects of associative

overdominance. As predicted, diversity was reduced relative to the neutral case (Supporting

Figure S4). When the scaling of our simulations is less extreme, the amount of nucleotide

diversity observed is very similar to the amount observed with more extreme scaling (Figure S4).

We also plotted the SFS under all models and conditions (Supporting Figures S5-S7). As

expected, migration led to an increase in the number of variants shared between populations.

Balancing selection and background selection with a complex genomic architecture also led to an

increase in the number of shared variants between populations, even in the absence of migration.

Again, in the background selection case, this could be explained by associative overdominance

in regions of low recombination. When the dominance coefficient is set to 0.5, this pattern in the

SFS disappears, again supporting associative overdominance as an explanation (Supporting

Figure S8). When the scaling of our simulations is less extreme, the SFS results in our

background selection simulations remain (Supporting Figure S8D and S8E).

Selection leads to false inferences of migration using ∂a∂i

Selection sometimes resulted in false inferences of migration in ∂a∂i using a likelihood ratio test (LRT). For the lowest divergence times, false positive rates were elevated even in the absence of selection and were not heavily impacted by selection (Figure 1a,b, Supporting Figure S9). The rates of rejection ranged from 18% to 41%, when only 1% false positives are expected at this *p*-value. When divergence times were moderate, false positive rates were not greater than 1% for any conditions (Figure 1c,d; Supporting Figure S9). Most strikingly, for the highest divergence times considered here, the isolation-only model was often erroneously rejected in favor of the isolation-with-migration model in the presence of balancing and background selection. Under the uniform recombination model, the isolation-only model was rejected in 10, 41, and 50% of replicates with 5, 10, and 15% of loci experiencing balancing selection, respectively (Figure 1e, Supporting Figure S9). Under the complex recombination model, the isolation-only model was rejected in 95, 100, and 100% of replicates with 5, 10, and 15% of loci experiencing balancing selection (Figure 1f, Supporting Figure S9). Background selection led to false positives in 100% of replicates under the complex model (Figure 1f). Under the complex model, selective sweeps also led to false positives in some conditions. Specifically, when 5% of loci experienced sweeps in population 1 or in the ancestor, the isolation-only model was rejected in 28 and 29% of replicates, respectively (Supporting Figure S9). Although the unconstrained model (including migration) should always have a higher likelihood than the constrained model (without migration), this was not always the case. Particularly in the medium divergence case, in ∂a∂i we observed instances in which the constrained model had higher likelihoods, indicating potential issues accurately approximating the likelihood for some datasets (Supporting Figure S10).

Perhaps as expected given the LRT results, selection also sometimes resulted in elevated

estimates of migration rates in $\partial a \partial i$. As with the LRT, results for the lowest divergence time lack

any clear signal related to selection: non-zero rates are observed even in the absence of selection

in very recently diverged populations (Figure 2a-b). Again, the most striking impact was seen in

the high divergence case in the presence of balancing and background selection (Figure 2e,f,

Supporting Figures S11). Migration rates were slightly overestimated under the uniform model

with balancing selection. Rates were also overestimated under the complex model with balancing

or background selection (Figure 2f), and, in the case of balancing selection, the degree of

overestimation increased with the percentage of loci experiencing balancing selection

(Supporting Figures S13-S14). When simulations included migration, migration rate estimates

tended to be higher under the complex genomic architecture (Supporting Figures S11-S14).

In the absence of migration and selection, ancestral θ was underestimated (Supporting Figure

S15). Background selection under the uniform model and selective sweeps tended to reduce

estimates of ancestral θ, while background selection with a complex genomic architecture and

balancing selection led to overestimates of ancestral θ when divergence times were high. The

presence of migration led to increased estimates of ancestral θ (Supporting Figure S15). We also

estimated the size of each population relative to the ancestral population (V_1, V_2, Supporting

Figures S16-S17). Notably, especially under the uniform genomic architecture, estimates of V_1

and V_2 tended to compensate for mistakes in the estimates of ancestral θ. In other words, when

ancestral θ was underestimated, V_1 and V_2 tended to be overestimated, and vice versa.

Divergence time estimates were fairly accurate in the absence of migration and selection,

although they were somewhat overestimated in the high divergence case (Supporting Figure

S18). When combined with a uniform genomic architecture, background selection and selective

sweeps led to overestimates of divergence times in the absence of migration, while balancing

selection led to underestimates of divergence times in the high divergence case. However, when

combined with a complex genomic architecture and moderate or high divergence times,

background selection led to underestimated divergence times. Divergence times were always

underestimated in the presence of migration, except for at the lowest divergence times with

background selection and a uniform genomic architecture.

For the two cases with the highest false positive rates (background and balancing selection with

high divergence times and a complex genomic architecture), we compared models with and

without migration, with and without variation in effective population sizes, and with or without

variation in migration rates following Rougeux *et al*. (2017). With background selection, this led

to a reduction in false positives (from 100% to 81%), but for the majority of replicates, a model

with migration was still selected as the best model, and the most commonly selected model

included two categories of migration rates (Supporting Figure S19a). For balancing selection,

this approach also reduced the false positive rate (from 100% to 83%), and the isolation-with-

migration model was selected most often (Supporting Figure S19b).

Selection leads to false inferences of migration using fastsimcoal2

Background and balancing selection often resulted in false inferences of migration in

fastsimcoal2. For the lowest divergence times, we rarely rejected the isolation-only model

(Figure 3a-b, Supporting Figure S20). When divergence times were medium, false positive rates

were elevated across all conditions, even in the absence of selection: rates of rejection ranged

from 13% to 45% (Figure 3c-d). For the highest divergence times considered here, the isolation-

only model was always erroneously rejected in favor of the isolation-with-migration model in the

presence of background selection and a complex genomic architecture (Figure 3f). Under the

complex genomic architecture, the isolation-only model was rejected in 21, 81, and 100% of

replicates with balancing selection in 5, 10, and 15% of loci, respectively. (Figure 3f, Supporting

Figure S20). There was also a slightly elevated false positive rate (4%) under a uniform genomic

architecture when 15% of loci experienced balancing selection (Figure 3e). Although the

unconstrained model (including migration) should always have a higher likelihood than the

constrained model (without migration), as with $\partial a \partial i$ this was not always the case. Particularly in

the low divergence case, in fastsimcoal2 we observed instances in which the constrained model

had higher likelihoods, indicating potential issues accurately approximating the likelihood

(Supporting Figure S21).

As with $\partial a \partial i$, selection resulted in elevated estimates of migration rates in fastsimcoal2 (Figure

4, Supporting Figures S22-S25). When divergence times were high, migration rates were slightly

overestimated with a uniform genetic architecture and balancing selection and were substantially

overestimated in the presence of a complex genomic architecture and background or balancing

selection (Figure 4f). In the case of balancing selection, the degree of overestimation increased

with the percentage of loci experiencing balancing selection (Supporting Figures S24-S25).

When simulations included migration, migration rate estimates tended to be lower under the

complex genomic architecture and were slightly elevated in the presence of background selection

with a complex genomic architecture, balancing selection, and adaptive introgression

(Supporting Figures S22-S25).

Estimates of the ancestral population sizes were fairly accurate in fastsimcoal2 (Supporting

Figure S26). The relative population sizes V_1 and V_2 were overestimated across all models and

conditions (Supporting Figures S27-S28). In the absence of selection, migration, and variation in

mutation and recombination rates divergence time estimates were accurate (Supporting Figure

S29). Divergence time estimates were higher under the complex genomic architecture compared

to the uniform genomic architecture and were reduced in the presence of background selection

(Supporting Figure S29). Divergence times tended to be underestimated in the presence of

migration (Supporting Figure S29).

Selection leads to false inferences of migration in BPP

Using BPP, non-zero migration rates were often inferred in the absence of migration (Figure 5;

Supporting Figures S30-S32). The highest posterior density interval (HDI) of the two migration

parameters, φ-*X* and φ-*Y*, rarely contained zero in the low divergence case, regardless of the

presence of selection (Figure 5a,b; Supporting Figure S30). In the medium and high divergence

cases, BPP still inferred non-zero migration often. In the medium divergence case, under the

uniform recombination and mutation model, the HDI for φ-*Y* did not include zero in 30, 30, and

40% of replicates with balancing selection in 5, 10, and 15% of loci (Figure 5c; Supporting

Figure S31). Similarly, the HDI for φ-*X* did not include zero in 10, 35, and 45% of replicates

with balancing selection in 5, 10, and 15% of loci (Figure 5c; Supporting Figure S32). Under the

complex model, the HDI for φ-*Y* did not include zero in 10, 25 and 40% of replicates with

balancing selection in 5, 10, and 15% of loci (Figure 5d; Supporting Figure S31). Similarly, the

HDI for φ-*X* did not include zero in 25, 45, and 65% of replicates (Figure 5d; Supporting Figure

S32). False positive rates were also slightly elevated across other conditions in the medium

divergence case (0-10% false positive rate). (Figure 5d; Supporting Figures S30-S32). In the

high divergence case, BPP inferred non-zero migration in 0-10% of replicates under each model

and condition, but there was no clear pattern with respect to selection and genomic architectures

(Figure 5e-f; Supporting Figures S30-S32). Although Bayesian methods do not have an

equivalent "false positive" rate to frequentist methods, we would not expect such a high

proportion of HDIs to not include the true parameter value. We also used Bayes Factors to

compare a model with introgression to a model without introgression for a subset of datasets

with high divergence (Supporting Table S1). We rejected the model without migration in 40% of

replicates without migration under both neutral and background selection conditions, and we

failed to reject the model without migration in 20% of replicates with migration and background

selection.

As expected, given that the HDIs do not overlap zero, migration rate estimates were elevated in

the presence of balancing selection (Figure 6; Supporting Figures S33). When simulations

included migration, estimates of φ-*X* and φ-*Y* were generally higher under the complex genomic

architecture (Supporting Figures S33-S34). Results were qualitatively similar whether 5, 10, or

15% of sweep datasets experienced a sweep (Supporting Figures S35-S36).

We also evaluated whether there was evidence for a lack of convergence in BPP runs by

examining ESS values. We focused on ESS values for the log likelihood, along with the φ-*X* and

φ-*Y* parameters. In the low divergence case, ESS values for the φ parameters were often low,

indicating a lack of convergence, particularly under the complex genomic architecture; however,

ESS values were generally greater than 200 in the medium and high divergence cases

(Supporting Figure S37). Under some conditions, there was evidence of a correlation between

ESS values and parameter estimates (e.g., for φ-*X* under a neutral model with a complex

genomic architecture, Supporting Figure S38). This suggests that in some (but not all) instances,

assessing convergence may allow researchers to identify problematic cases.

In the absence of selection and migration, divergence times were generally overestimated

(Supporting Figure S39). Divergence time estimates were reduced in the presence of background

selection and were elevated under the complex genomic architecture relative to the uniform

genomic architecture. The presence of migration led to underestimates of divergence times when

divergence times were high.

Discussion

Our results suggest that, while popular methods for estimating migration rates between sister

populations or species are largely robust to selective sweeps and simple models of background

selection, models with non-uniform recombination rates and models with balancing selection can

lead to high rates of false positives. The three approaches tested (fastsimcoal, ∂a∂i, and BPP) all

showed high rates of misleading results in the presence of balancing selection and background

selection with a complex genetic architecture (Figures 1, 3, 5). Given that large portions of the

genomes of many species are impacted by selection (Begun *et al.* 2007; McVicker *et al.* 2009;

Sella *et al.* 2009; Langley *et al.* 2012; Corbett-Detig *et al.* 2015; Phung *et al.* 2016; Pouyet *et al.*

2018) and that variation in mutation and recombination rates across the genome is the norm,

these results suggest that some inferences of introgression may be artifacts that do not reflect

biological reality.

Numerous studies have found that ignoring natural selection can negatively impact different

types of demographic inferences. Selection can lead to false inference of population size changes

(e.g., Ewing and Jensen 2016; Schrider *et al.* 2016; Johri *et al.* 2021) and several studies have

suggested that selection can also mislead inferences of migration (e.g., Cruickshank and Hahn

2014; Mathew and Jensen 2015; Roux *et al.* 2016). In our study, false inferences appear to be

primarily driven by the impacts of balancing selection and associative overdominance. Both

balancing selection and associative overdominance increase the number of polymorphisms

shared between two populations. This effect is clearly visible in the SFS produced under these

conditions, which resemble those produced by migration (Supporting Figures S6-S7). It remains

unclear how prevalent these patterns are likely to be in empirical systems. Notably, we did not

set out to simulate the effects of associative overdominance—our simulations were

parameterized based on the *Drosophila melanogaster* genome, and this unexpectedly led to

associative overdominance. In species with compact genomes and low recombination regions,

therefore, we may expect these impacts to potentially be widespread. In species with large

genomes, but many functional non-coding elements, the impacts may also extend throughout the

genome (e.g., Gilbert *et al.* 2020). Furthermore, while the balancing selection simulated here

may seem extreme, there are numerous examples of trans-specific polymorphisms maintained by

balancing selection (e.g., the *S*-locus in flowering plants (Wright 1939; Le Veve *et al.* 2023)).

We recommend that researchers exclude such loci and neighboring regions—if they can be

identified—when conducting demographic inference.

To account for the effects of selection, several approaches for inferring migration have been

developed that allow for heterogeneous effective population sizes among loci (Sousa *et al.* 2013;

Roux *et al.* 2016; Sethuraman *et al.* 2019; Fraïsse *et al.* 2021). Although these methods vary in

the types of inferences that can be made—from locus-specific migration rates to genome-wide

migration rates—they all model selection by allowing for variation in θ among loci. Notably,

while this may accommodate some of the simpler effects of background selection (e.g., reduced

diversity and increased variation in coalescence times), it is unlikely to accommodate the impacts

of associative overdominance or balancing selection. To evaluate this, we applied such an

approach to compare models in ∂a∂i, and, while false positive rates were reduced, they were still

high (81% and 83% for background and balancing selection, respectively). BPP also allows for

variation across loci, either by using a rate multiplier for θ and τ or for θ alone (Flouri *et al.*

2020). In our analyses, we used both rate multipliers and still found relatively high false positive

rates, particularly in the presence of balancing selection. It is not clear that the results using any

other similar methods would differ qualitatively from these (to our knowledge, none have been

tested against a no-migration scenario with selection).

Phylogenetic methods for inferring gene flow are much more robust to assumptions about

selection, largely because they often depend on asymmetries in tree topologies (Hibbins and

Hahn 2022). However, the dependence on tree asymmetry also means that they cannot be used to

detect gene flow between sister lineages. So, what is the way forward? Our results, along with

previous studies, highlight several potential possibilities. Statistical-learning approaches are

highly flexible (Schrider and Kern 2018), and one path forward involves training these

algorithms under appropriate models that incorporate selection. One such approach has

successfully used approximate Bayesian computation to jointly estimate DFEs and population

size histories (Johri *et al.* 2020). Beyond training statistical learning algorithms on more realistic

training data, techniques for domain adaptation—a subfield of machine learning that aims to

adapt an algorithm trained on the source domain (e.g., on simulations under a model of interest)

to the target domain (i.e., empirical data; reviewed in Wilson and Cook 2020) offer a promising

path forward to accommodating complex biological realities in population genomics (e.g., Mo

and Siepel 2023). Regardless of which methods are used, accurate inferences of demographic

histories will have to include the complexities introduced by selection.

Accurate inferences of introgression histories are important because they can tell us about modes

of speciation. A large number of studies supporting gene flow between closely related

populations have been interpreted as lending support to speciation-with-gene-flow models; our

results highlight that caution is warranted in these interpretations (cf. Cruickshank and Hahn,

2014), though there are other reasons to exercise caution as well (Yang *et al.* 2017). Although

many estimates were non-zero, migration rates estimated for datasets generated under models

including selection were low absolutely. In fastsimcoal2, estimated migration rates per

generation were on the order of 10^{-9}, in $\partial a \partial i$, $2 \times N_{ref} \times m_{ij}$ was on the order of 0.002 (~10^{-9}

migrants per generation), and in BPP the weight of the hybrid edge was on the order of 0.002

(Figures 2, 4, 6). On the other hand, estimates of effective population size often change multiple

orders of magnitude over extremely short periods of time in analyses of empirical data using

these same methods (e.g., Rosser *et al.* 2024), suggesting that there are other biological

complexities not captured by these methods (or by our simulations). Moving forward, we

recommend that inferences of migration between sister populations made without considering

selection be interpreted with caution, particularly when inferred rates of migration are low.

Importantly, we do not believe that the results found here with a limited set of selective scenarios

and a limited set of introgression histories can fully describe the effects of selection, mutation,

and recombination on inaccurate demographic inferences. Inferences about the presence,

direction, and timing of introgression (including whether speciation and introgression occur at

the same time—i.e. homoploid hybrid speciation) may all be affected by models that ignore

natural selection and complex genomic architectures. We hope that new methods can be

developed to overcome these obstacles.

Data Availability

Simulated data formatted for various programs are available on Figshare (DOI:

10.6084/m9.figshare.24354277). All scripts are available on GitHub

(https://meganlsmith.github.io/selectionandmigration/).

Acknowledgements

We thank Peter Ralph and two reviewers for very helpful comments and suggestions. This work

was supported by a National Science Foundation (NSF) postdoctoral fellowship to MLS (DBI-

2009989) and NSF grant to MWH (DBI-2146866).

Tables

Table 1: Simulation conditions considered in this study.

Genomic Architecture	Model	Condition
uniform	no migration	Neutral
		Background selection
		Sweep P1 (5, 10, 15%)
		Sweep Ancestor (5, 10, 15%)
		Balancing (5, 10, 15%)
	p1_p2	Neutral
		Background selection
		Sweep P1 (5, 10, 15%)
		Sweep Ancestor (5, 10, 15%)
		Balancing (5, 10, 15%)
		Adaptive Introgression (5, 10, 15%)
complex	no migration	Neutral
		Background selection
		Sweep P1 (5, 10, 15%)
		Sweep Ancestor (5, 10, 15%)
		Balancing (5, 10, 15%)
	p1_p2	Neutral
		Background selection
		Sweep P1 (5, 10, 15%)
		Sweep Ancestor (5, 10, 15%)
		Balancing (5, 10, 15%)
		Adaptive Introgression (5, 10, 15%)

Figures

Figure 1. Results of the LRT in ∂a∂i on datasets simulated without migration. Light blue

indicates cases where we failed to reject the isolation-only model, and dark blue indicates cases

where we rejected the isolation-only model in favor of the isolation-with-migration model. a)

results for T=1N with a uniform genomic architecture; b) results for T=1N with a complex

genomic architecture; c) results for T=4N with a uniform genomic architecture; d) results for

T=4N with a complex genomic architecture; e) results for T=16N with a uniform genomic

architecture; f) results for T=16N with a complex genomic architecture.

606 **Figure 2.** Estimates of M_{21} in ∂a∂i for datasets simulated without migration. Estimates are in

607 units of $2 \times N_{ref} \times m_{ij}$. a) results for T=1$N$ with a uniform genomic architecture; b) results for

608 T=1N with a complex genomic architecture; c) results for T=4N with a uniform genomic

609 architecture; d) results for T=4N with a complex genomic architecture; e) results for T=16N with

610 a uniform genomic architecture; f) results for T=16N with a complex genomic architecture.

Figure 3. Results of the LRT in fastsimcoal2 for datasets simulated without migration. Light

blue indicates cases where we failed to reject the isolation-only model, and dark blue indicates

cases where we rejected the isolation-only model in favor of the isolation-with-migration model.

a) results for T=1*N* with a uniform genomic architecture; b) results for T=1*N* with a complex

genomic architecture; c) results for T=4*N* with a uniform genomic architecture; d) results for

T=4*N* with a complex genomic architecture; e) results for T=16*N* with a uniform genomic

architecture; f) results for T=16*N* with a complex genomic architecture.

Figure 4. Estimates of m_{21} in fastsimcoal2 for datasets simulated without migration. The rate m_{ij}

is the probability of any gene moving from population i to population j backwards in time each

generation. a) results for T=1N with a uniform genomic architecture; b) results for T=1N with a

complex genomic architecture; c) results for T=4N with a uniform genomic architecture; d)

results for T=4N with a complex genomic architecture; e) results for T=16N with a uniform

genomic architecture; f) results for T=16N with a complex genomic architecture.

Figure 5. The proportions of replicates with 95% highest posterior density intervals for φ-*Y* that do not include zero in BPP for datasets simulated without migration. Light blue indicates cases where the HDI includes zero, and dark blue indicates cases where the HDI does not include zero. a) results for T=1*N* with a uniform genomic architecture; b) results for T=1*N* with a complex genomic architecture; c) results for T=4*N* with a uniform genomic architecture; d) results for T=4*N* with a complex genomic architecture; e) results for T=16*N* with a uniform genomic architecture; f) results for T=16*N* with a complex genomic architecture.

Figure 6. Mean posterior estimates of φ-Y in BPP for datasets simulated without migration. φ-Y is

the weight of the introgression edge Y in the MSci model. a) results for T=1N with a uniform

genomic architecture; b) results for T=1N with a complex genomic architecture; c) results for

T=4N with a uniform genomic architecture; d) results for T=4N with a complex genomic

architecture; e) results for T=16N with a uniform genomic architecture; f) results for T=16N with

a complex genomic architecture.

References

Adams M. D., S. E. Celniker, R. A. Holt, C. A. Evans, J. D. Gocayne, *et al.*, 2000 The genome
sequence of *Drosophila melanogaster*. Science 287: 2185–2195.

Begun D. J., A. K. Holloway, K. Stevens, L. W. Hillier, Y.-P. Poh, *et al.*, 2007 Population
genomics: whole-genome analysis of polymorphism and divergence in *Drosophila
simulans*. PLoS Biology 5: e310.

Comeron J. M., R. Ratnappan, and S. Bailin, 2012 The many landscapes of recombination in
Drosophila melanogaster. PLoS Genetics 8: e1002905.

Corbett-Detig R. B., D. L. Hartl, and T. B. Sackton, 2015 Natural selection constrains neutral
diversity across a wide range of species. PLoS Biology 13: e1002112.

Cruickshank T. E., and M. W. Hahn, 2014 Reanalysis suggests that genomic islands of
speciation are due to reduced diversity, not reduced gene flow. Molecular Ecology 23:
3133–3157.

Cutter A. D., and B. A. Payseur, 2013 Genomic signatures of selection at linked sites: unifying
the disparity among species. Nature Reviews Genetics 14: 262–274.

Ewing G. B., and J. D. Jensen, 2016 The consequences of not accounting for background
selection in demographic inference. Molecular Ecology 25: 135–141.

Excoffier L., T. Hofer, and M. Foll, 2009 Detecting loci under selection in a hierarchically
structured population. Heredity 103: 285–298.

Excoffier L., I. Dupanloup, E. Huerta-Sánchez, V. C. Sousa, and M. Foll, 2013 Robust

demographic inference from genomic and SNP data. PLoS Genetics 9: e1003905.

Flouri T., X. Jiao, B. Rannala, and Z. Yang, 2020 A Bayesian implementation of the

multispecies coalescent model with introgression for phylogenomic analysis. Molecular

Biology and Evolution 37: 1211–1223.

Fraïsse C., I. Popovic, C. Mazoyer, B. Spataro, S. Delmotte, *et al.*, 2021 DILS: Demographic

inferences with linked selection by using ABC. Molecular Ecology Resources 21: 2629–

2644.

Geneva A. J., C. A. Muirhead, S. B. Kingan, and D. Garrigan, 2015 A new method to scan

genomes for introgression in a secondary contact model. PLoS One 10: e0118621.

Gilbert K. J., F. Pouyet, L. Excoffier, and S. Peischl, 2020 Transition from background selection

to associative overdominance promotes diversity in regions of low recombination.

Current Biology 30: 101-107.e3.

Gutenkunst R. N., R. D. Hernandez, S. H. Williamson, and C. D. Bustamante, 2009 Inferring the

joint demographic history of multiple populations from multidimensional SNP frequency

data. PLoS Genetics 5: e1000695.

Haller B. C., and P. W. Messer, 2019 SLiM 3: forward genetic simulations beyond the Wright–

Fisher model. Molecular Biology and Evolution 36: 632–637.

Hey J., and R. Nielsen, 2007 Integration within the Felsenstein equation for improved Markov

chain Monte Carlo methods in population genetics. Proceedings of the National Academy

of Sciences 104: 2785–2790.

Hibbins M. S., and M. W. Hahn, 2022 Phylogenomic approaches to detecting and characterizing

introgression. Genetics 220: iyab173.

Huber C. D., B. Y. Kim, C. D. Marsden, and K. E. Lohmueller, 2017 Determining the factors

driving selective effects of new nonsynonymous mutations. Proceedings of the National

Academy of Sciences 114: 4465–4470.

Johri P., B. Charlesworth, and J. D. Jensen, 2020 Towards an evolutionarily appropriate null

model: Jointly inferring demography and purifying selection. Genetics 215: 173–192.

Johri P., K. Riall, H. Becher, L. Excoffier, B. Charlesworth, *et al.*, 2021 The impact of purifying

and background selection on the inference of population history: problems and prospects.

Molecular Biology and Evolution 38: 2986–3003.

Joly S., P. A. McLenachan, and P. J. Lockhart, 2009 A statistical approach for distinguishing

hybridization and incomplete lineage sorting. The American Naturalist 174: E54–E70.

Kern A. D., and M. W. Hahn, 2018 The neutral theory in light of natural selection. Molecular

Biology and Evolution 35: 1366–1371.

Langley C. H., K. Stevens, C. Cardeno, Y. C. G. Lee, D. R. Schrider, *et al.*, 2012 Genomic

variation in natural populations of *Drosophila melanogaster*. Genetics 192: 533–598.

Le Veve A., N. Burghgraeve, M. Genete, C. Lepers-Blassiau, M. Takou, *et al.*, 2023 Long-term

balancing selection and the genetic load linked to the self-incompatibility locus in

Arabidopsis halleri and *A. lyrata*. Molecular Biology and Evolution 40: msad120.

Luqman H., A. Widmer, S. Fior, and D. Wegmann, 2021 Identifying loci under selection via

explicit demographic models. Molecular Ecology Resources 21: 2719–2737.

Mallet J., N. Besansky, and M. W. Hahn, 2016 How reticulated are species? BioEssays 38: 140–

149.

Mathew L. A., and J. D. Jensen, 2015 Evaluating the ability of the pairwise joint site frequency

spectrum to co-estimate selection and demography. Frontiers in Genetics 6: 268.

McVicker G., D. Gordon, C. Davis, and P. Green, 2009 Widespread genomic signatures of

natural selection in hominid evolution. PLoS Genetics 5: e1000471.

Mo Z., and A. Siepel, 2023 Domain-adaptive neural networks improve supervised machine

learning based on simulated population genetic data. PLOS Genetics 19: e1011032.

Nielsen R., I. Hellmann, M. Hubisz, C. Bustamante, and A. G. Clark, 2007 Recent and ongoing

selection in the human genome. Nature Reviews Genetics 8: 857–868.

Ohta T., 1971 Associative overdominance caused by linked detrimental mutations. Genetical

Research 18: 277–286.

Pamilo P., and S. Pálsson, 1998 Associative overdominance, heterozygosity and fitness. Heredity

81: 381–389.

Payseur B. A., and L. H. Rieseberg, 2016 A genomic perspective on hybridization and

speciation. Molecular Ecology 25: 2337–2360.

Phung T. N., C. D. Huber, and K. E. Lohmueller, 2016 Determining the effect of natural

selection on linked neutral divergence across species. PLoS Genetics 12: e1006199.

Pouyet F., S. Aeschbacher, A. Thiéry, and L. Excoffier, 2018 Background selection and biased

gene conversion affect more than 95% of the human genome and bias demographic

inferences. eLife 7: e36317.

Rosenzweig B. K., J. B. Pease, N. J. Besansky, and M. W. Hahn, 2016 Powerful methods for

detecting introgressed regions from population genomic data. Molecular Ecology 25:

2387–2397.

Rosser N., F. Seixas, L. M. Queste, B. Cama, R. Mori-Pezo, *et al.*, 2024 Hybrid speciation

driven by multilocus introgression of ecological traits. Nature 628: 811–817.

Rougeux C., L. Bernatchez, and P.-A. Gagnaire, 2017 Modeling the Multiple Facets of

Speciation-with-Gene-Flow toward Inferring the Divergence History of Lake Whitefish

Species Pairs (Coregonus clupeaformis). Genome Biology and Evolution 9: 2057–2074.

Roux C., C. Fraisse, J. Romiguier, Y. Anciaux, N. Galtier, *et al.*, 2016 Shedding light on the grey

zone of speciation along a continuum of genomic divergence. PLoS Biology 14:

e2000234.

Schrider D. R., A. G. Shanku, and A. D. Kern, 2016 Effects of linked selective sweeps on

demographic inference and model selection. Genetics 204: 1207–1223.

Schrider D. R., and A. D. Kern, 2018 Supervised machine learning for population genetics: a

new paradigm. Trends in Genetics 34: 301–312.

Schrider D. R., 2020 Background selection does not mimic the patterns of genetic diversity

produced by selective sweeps. Genetics 216: 499–519.

Sella G., D. A. Petrov, M. Przeworski, and P. Andolfatto, 2009 Pervasive natural selection in the

Drosophila genome? PLoS Genetics 5: e1000495.

Sethuraman A., V. Sousa, and J. Hey, 2019 Model-based assessments of differential

introgression and linked natural selection during divergence and speciation. bioRxiv.

https://doi.org/10.1101/786038

[dataset] Smith M. L. and M. W. Hahn, 2024 Data from Selection leads to false inferences of

introgression using popular methods. Figshare. doi: 10.6084/m9.figshare.24354277

Sousa V. C., M. Carneiro, N. Ferrand, and J. Hey, 2013 Identifying loci under selection against

gene flow in isolation-with-migration models. Genetics 194: 211–233.

Tellier A., P. Pfaffelhuber, B. Haubold, L. Naduvilezhath, L. E. Rose, *et al.*, 2011 Estimating

parameters of speciation models based on refined summaries of the joint site-frequency

spectrum. PLoS One 6: e18155.

Tine M., H. Kuhl, P.-A. Gagnaire, B. Louro, E. Desmarais, *et al.*, 2014 European sea bass

genome and its variation provide insights into adaptation to euryhalinity and speciation.

Nature Communications 5: 5770.

Williamson S. H., R. Hernandez, A. Fledel-Alon, L. Zhu, R. Nielsen, *et al.*, 2005 Simultaneous

inference of selection and population growth from patterns of variation in the human

genome. Proceedings of the National Academy of Sciences 102: 7882–7887.

Wilson G., and D. J. Cook, 2020 A survey of unsupervised deep domain adaptation. ACM

Transactions on Intelligent Systems and Technology (TIST) 11: 1–46.

Wright S., 1939 The distribution of self-sterility alleles in populations. Genetics 24: 538–552.

Yang M., Z. He, S. Shi, and C.-I. Wu, 2017 Can genomic data alone tell us whether speciation

happened with gene flow? Molecular Ecology 26: 2845–2849.