Reconstitution of cytolinker-mediated crosstalk between actin and vimentin

Irene Istúriz Petitjean,[1,*] Quang D. Tran [b,2,*] Angeliki Goutou,[1] Zima Kabir,[1] Gerhard Wiche,[3,‡] Cécile Leduc,[2,§] and Gijsje H. Koenderink[1,¶]

[1]*Department of Bionanoscience, Kavli Institute of Nanoscience Delft, Delft University of Technology, 2629 HZ Delft, The Netherlands*

[2]*Université Paris Cité, CNRS, Institut Jacques Monod, F-75013 Paris, France*

[3]*Department of Biochemistry and Cell Biology, Max Perutz Laboratories, University of Vienna, Vienna, Austria.*

Abstract

Cell shape and motility are determined by the cytoskeleton, an interpenetrating network of actin filaments, microtubules, and intermediate filaments. The biophysical properties of each filament type individually have been studied extensively by cell-free reconstitution. By contrast, the interactions between the three cytoskeletal networks are relatively unexplored. They are coupled via crosslinkers of the plakin family such as plectin. These are challenging proteins for reconstitution because of their giant size and multidomain structure. Here we engineer a recombinant actin-vimentin crosslinker protein called 'ACTIF' that provides a minimal model system for plectin, recapitulating its modular design with actin-binding and intermediate filament-binding domains separated by a coiled-coil linker for dimerisation. We show by fluorescence and electron microscopy that ACTIF has a high binding affinity for vimentin and actin and creates mixed actin-vimentin bundles. Rheology measurements show that ACTIF-mediated crosslinking strongly stiffens actin-vimentin composites. Finally, we demonstrate the modularity of this approach by creating an ACTIF variant with the intermediate filament binding domain of Adenomatous Polyposis Coli. Our protein engineering approach provides a new cell-free system for the biophysical characterization of intermediate filament-binding crosslinkers and for understanding the mechanical synergy between actin and vimentin in mesenchymal cells.

Keywords: Plakins, Plectin, Cytoskeleton, Rheology, Binding kinetics

------

[b] Present address: Department of Physics, Brandeis University, Waltham, MA 02454

[*] These authors contributed equally to this work.

HIGHLIGHTS

- Engineering of a recombinant actin-vimentin crosslinker called ACTIF with the plectin intermediate filament binding domain (IFBD), calponin homology domains that mediate actin binding, and a coiled-coil linker.

- ACTIF crosslinks F-actin and vimentin and mediates their co-localization *in vitro*.

- ACTIF has a binding affinity for vimentin that is about 500 times higher than for F-actin.

- ACTIF forms composite bundles of F-actin and vimentin filaments.

- Composite F-actin/vimentin networks stiffen upon crosslinking with ACTIF.

- The design of actin-vimentin crosslinker is modular, as other IFBDs like APCn2 can also be used.

1. INTRODUCTION

The cytoskeleton serves multiple crucial functions within mammalian cells. It defines cell shape (Fletcher and Mullins 2010), provides mechanical stability (Pegoraro et al. 2017), drives cell movement and division (Nakaseko and Yanagida 2001), connects cells within tissues (Hoelzle and Svitkina 2012), and influences intracellular signaling processes (Janmey 1998). The cytoskeleton is a network of three primary types of biopolymers: actin filaments (F-actin), microtubules (MTs), and intermediate filaments (IFs), each playing distinct roles in various cellular activities. F-actin and MTs have been widely studied for their roles in regulating cell mechanics and essential processes such as cell division and migration (Dogterom and Koenderink 2019). Both biopolymers form dynamic networks that can generate forces by active (de)polymerization and by the activity of molecular motor proteins. F-actin networks drive different modes of cell migration and also drive membrane constriction in dividing cells by forming a contractile ring. At the same time, F-actin is an important determinant of cell stiffness and strength (Fletcher and Mullins 2010). MTs build the mitotic spindle that

‡ gerhard.wiche@univie.ac.at

§ cecile.leduc@cnrs.fr

¶ G.H.Koenderink@tudelft.nl

drives chromosome segregation in dividing cells, while also contributing to the overall mechanical integrity of the cytoskeleton (Dogterom and Koenderink 2019). By contrast, much less is known about the role of IFs in cell function. Traditionally, they are considered mostly as a kind of "cellular safety belt" that protects cells against large mechanical stresses (Hu et al. 2019). Indeed IFs can stretch to more than 3 times their rest length because they are built up of fibrous subunits that can unfold (Block et al. 2017, Forsting et al. 2019). However, in the past years, many additional non-mechanical functions of IFs have been discovered (Patteson et al. 2020). IF networks contribute to the functioning of several cellular organelles (Chernoivanenko et al. 2015, Styers et al. 2004) and regulate signaling pathways that control cell survival, growth, and differentiation (Ridge et al. 2022, Sjöqvist et al. 2021). Intriguingly, the IF protein family comprises 70 different proteins whose expression pattern is cell-type-specific and changes during development and in disease. This raises the interesting possibility that IFs confer cell-type-specific functions. The two most well-studied IFs are keratins, expressed in non-motile epithelial cells, and vimentin, expressed in motile mesenchymal cells. Recent biophysical studies indicate that keratin and vimentin have distinct mechanical properties tailored to their cellular functions (Lorenz et al. 2023). This mechanical difference may contribute to the invasiveness of epithelial cancer cells that have undergone a switch from keratin to vimentin expression (Berr et al. 2023).

The three cytoskeletal biopolymers have traditionally been regarded as independent systems with separate cellular tasks. However, growing evidence shows that their functions are coordinated and that this coordination is essential for many core cellular functions (Huber et al. 2015). Directional cell migration, for instance, relies on F-actin to generate the driving force, but coordination with both MTs and IFs is required to polarize the F-actin cytoskeleton and control the amplitude and directionality of force generation (Costigliola et al. 2017, Gan et al. 2016, Jiu et al. 2015, 2017, van Bodegraven and Etienne-Manneville 2020). In dividing cells, F-actin/MT interactions control the correct placement of the division plane and F-actin/IF interactions at the cell cortex are essential for normal mitotic progression (Duarte et al. 2019, Serres et al. 2020). These examples underscore the need to study the mechanisms that couple the activities of the three cytoskeletal systems.

Cytoskeletal coupling is known to involve a combination of biochemical coupling via signaling loops and transcriptional regulation (Tomar and Schlaepfer 2009) and physical coupling via crosslinking proteins (Prechova et al. 2022), motor proteins (Seetharaman and

Etienne-Manneville 2020), and joint connections to adhesion receptor complexes in the plasma membrane (Osmanagic-Myers et al. 2015) and to LINC complexes in the nuclear envelope (Tusamda Wakhloo et al. 2020). This complexity makes it difficult to understand the mechanisms of cytoskeletal coupling in cells. Cell-free reconstitution can provide a way to overcome this challenge. In this reductionist approach, components of the cytoskeleton are purified and studied in isolation or together with a limited set of regulatory proteins. In the past, cell-free reconstitution of isolated cytoskeletal biopolymers has successfully delivered quantitative models explaining how F-actin drives cell motility, how MTs organize the mitotic spindle, and how IFs provide mechanical resilience. However, coupling between different cytoskeletal subsystems has only recently begun to be investigated by reconstitution. Most of the available work has focused on F-actin/MT coordination, with studies of molecular motor motility at F-actin-MT crossings (Ali et al. 2007, Schroeder et al. 2010), interdependent F-actin/MT polymerization (Alkemade et al. 2022, Breitsprecher et al. 2012, López et al. 2014), and mechanical synergy in F-actin/MT composites (Pelletier et al. 2009, Ricketts et al. 2018, Sheung et al. 2021). There are even fewer studies of cytoskeletal composites involving IFs. A few studies have investigated the collective viscoelastic properties of two-component (IF/F-actin) (Esue et al. 2006, Golde et al. 2018, Jensen et al. 2014) and three-component composite networks (Shen et al. 2021) and interactions between IFs and MTs (Schaedel et al. 2021). In all these studies, the cytoskeletal filaments were combined in the absence of any cytolinkers.

In cells, IFs are crosslinked to F-actin and MTs via large crosslinking proteins including members of the plakin family such as plectin (Bouameur et al. 2014, Mohammed et al. 2020, Wiche and Winter 2011). Electron microscopy imaging and proximity ligation assays in cells have shown that plectin forms both IF-IF crosslinks and crosslinks of IFs to F-actin and MTs (Foisner et al. 1995, 1988, Marks et al. 2022, Svitkina et al. 1996, Wiche and Baker 1982). In motile mesenchymal cells, plectin crosslinks vimentin filaments to F-actin stress fibers (Jiu et al. 2015, Marks et al. 2022) and to F-actin found at the base of invadopodia (Schoumacher et al. 2010, Yoneyama et al. 2014). In mitotic mesenchymal cells, plectin crosslinks vimentin to the actin cortex (Serres et al. 2020). The many splice isoforms of plectin share a conserved tripartite structure with a plakin domain, central rod domain, and C-terminal domain. Depending on the isoform, plectin can have up to seven functional domains (Fig. 1A). The N-terminal motif is isoform-specific and targets plectin to specific

cellular locations such as the nucleus, hemidesmosomes, or focal adhesions. Actin-binding is mediated by calponin homology domains that form an actin-binding domain (ABD) directly following the N-terminal motif (Fontao et al. 2001, Garcia-Alvarez et al. 2003). IF-binding is mediated by four domains at the C-terminus consisting of two plakin repeat domains (PRDs) separated by a linker region and a short C-terminal extremity containing Gly-Ser-Arg (GSR) repeats (Bouameur et al. 2014). The PRDs and the linker region share a basic groove that binds the acidic coiled-coil rod domain of IFs (Choi et al. 2002, Favre et al. 2011, Fogl et al. 2016, Kang et al. 2016, Mohammed et al. 2020).

Plectin depletion has been shown to alter the mechanical behavior of cells(Bonakdar et al. 2015, Martin et al. 2016, Moch et al. 2016, Na et al. 2009, Zrelski et al. 2023) and to impair the mechanical stability of epithelial and endothelial cell sheets (Osmanagic-Myers et al. 2015, Prechova et al. 2022). Indeed plectin mutations are associated with severe skin blistering disorders and muscular dystrophy. However, plectin's specific role in these mechanical abnormalities as a cytoskeletal crosslinker has been difficult to decipher since it has several other roles besides direct crosslinking: it also anchors IFs to cell-matrix and cell-cell adhesion receptor complexes (Bhattacharya et al. 2009, Colburn and Jones 2018, De Pascalis et al. 2018, Geerts et al. 1999, Walko et al. 2015), to LINC complexes in the nuclear envelope (Wilhelmsen et al. 2005) and to mitochondria (Winter et al. 2008). There is hence a need for reconstitution studies where plectin's crosslinking role can be studied in isolation from its other functions. Unfortunately, plectin is a challenging protein for reconstitution because of its giant size (500 kDa) and multidomain structure. Full-length plectin can in principle be obtained from mammalian cells (Foisner et al. 1988), but it is difficult to obtain proteins in high yield and purity using this strategy.

Here we engineer a recombinant actin-vimentin crosslinker that we call 'ACTIF', which is designed to crosslink F-actin and IFs in a similar manner as plectin. The engineered linker has a modular design with an N-terminal Green Fluorescent Protein (GFP) tag and actin-binding domain, a C-terminal IF-binding domain, and a coiled-coil linker to induce parallel dimerisation. Its small (138 kDa) size allows for convenient recombinant expression in bacteria. We demonstrate by imaging and rheology that ACTIF with the plectin interme-diate filament binding domain (IFBD) efficiently crosslinks vimentin intermediate filaments to F-actin. Using Total Internal Reflection Fluorescence (TIRF) imaging, we quantitatively characterize the binding kinetics for the plectin-derived ACTIF crosslinker. Finally, we

demonstrate the modularity of our approach by creating an alternative ACTIF variant with an IFBD based on the armadillo repeats of the tumor suppressor protein Adenomatous Polyposis Coli (APC) (Sakamoto et al. 2013). Our work provides a powerful new reconstitution platform to study the biophysical mechanisms that contribute to the functional coupling of the F-actin and IF cytoskeleton that controls cell mechanics and migration.

FIG. 1: ACTIF crosslinker with the plectin IFBD binds F-actin and vimentin. A) Domain structures of full-length plectin (top) and ACTIF with the ABD from MACF and the IFBD from plectin (bottom). B) Interferometric scattering microscopy (iSCAT) of ACTIF (50 nM in F-buffer buffer) shows a peak at a molecular weight of 243 kDa, indicating that ACTIF forms a dimer. C) SDS-PAGE of ACTIF purification after gel filtration showing a molecular weight marker (first lane) and the final pooled sample (second lane). D) Schematic of control experiment with TIRF microscopy to probe for interactions between vimentin filaments, immobilized on the microscope coverslip using vimentin antibodies, and F-actin polymerized from 1 μM G-actin in solution. Methylcellulose (0.2 %) was added to push F-actin down toward the surface. The images (right) show no co-localization between the filaments and hence no direct interactions. E) Schematic of sequential steps in the TIRF experiments to probe ACTIF-mediated binding of F-actin to vimentin filaments (left). First ACTIF (5 nM in F-buffer) was flushed into the channel containing surface-bound vimentin filaments for 15 min. Next pre-formed F-actin (0.1 μM in F-buffer) was flushed in. Images (right) show snapshots after each step with a merged image of the final situation on the right. F) Images of F-actin attachment over time on top of ACTIF-bound vimentin filaments. G) Mean intensity over time of a representative F-actin at the cross-section marked by the white line in Fig. 1(F), showing multiple actin filaments recruited on top of ACTIF-bound vimentin. H) Distribution of pixel fluorescence intensities in the F-actin channel along vimentin filaments ($\sim$150 filaments from 2 independent experiments) at steady state. The probability distribution function (PDF) was fitted by a multiple Gaussian, with the peak positions constrained to be multiples of the position of the first peak, and common width for all the peaks except the last one. The first peak was estimated at an intensity $\sim$600, as expected for single F-actin imaged in similar conditions. Scale bar: 5 μm

.

2. RESULTS

ACTIF crosslinks F-actin and vimentin and mediates co-localization

Full-length plectin (Fig. 1A (top)) is a giant (> 500 kDa) multidomain cytolinker protein (Wiche and Winter 2011). Its isoform-specific N-terminus is followed by the actin-binding domain (ABD), an elongated plakin domain (Al-Jassar et al. 2013, Ortega et al. 2016, Sonnenberg et al. 2007), a central alpha-helical rod domain (200 nm long) that mediates plectin dimerization, six tandemly arranged plakin repeat domains (PRDs, $\sim$300 residues

each) separated by linker regions of variable length, and finally, a short tail region containing Gly-Ser-Arg (GSR) repeats. The first four PRD domains do not contribute to IF-binding (Nikolic et al. 1996, Steiner-Champliaud et al. 2010). The last two PRD domains contribute together with the PRD5-PRD6 linker region ($\sim$50 residues) and the GSR-containing C-terminal extremity to ensure efficient IF binding (Bouameur et al. 2014). The linker region is essential but not sufficient for strong IF-binding (Bouameur et al. 2014, Favre et al. 2011, Nikolic et al. 1996). In addition, a sequence of approximately 50 amino acid residues (4262-4316) within one-third of the PRD5 is indispensable for filament association (Nikolic et al. 1996).

We generated a minimal crosslinker protein designed to model the crosslinking activity of plectin (Fig. 1A (bottom)), which we call ACTIF. We based the design on an earlier engineered cytolinker called TipAct that models the actin-microtubule cytolinker protein MACF/ACF7 (Alkemade et al. 2022, López et al. 2014). The TipAct protein consisted of an N-terminal green fluorescent protein (GFP) tag, followed by the F-actin binding domain of MACF, a coiled-coil spacer domain, and a C-terminal microtubule-binding domain. We retained this domain structure but replaced the microtubule-binding domain with the C-terminal intermediate filament binding domain (IFBD) of plectin. To ensure robust IF-binding, we included the last two PRD domains (PRD5 and PRD6) and their connecting linker (Nikolic et al. 1996). The ABD from MACF consists of a tandem of two calponin-homology (CH) domains and resembles the ABD of plectin (Fontao et al. 2001, Sevcik et al. 2004, Urbániková et al. 2002). The coiled-coil spacer domain separating the ABD and IFBD was taken from cortexillin I, in order to mimic plectin's dimeric state by inducing parallel dimerization of ACTIF (Faix et al. 1996). By removing the large plakin rod and the first four PRD domains of plectin, we obtained an actin-IF cytolinker protein with a molecular weight of only 138 kDa, enabling us to express the protein in bacteria.

We first tested the oligomeric state and purity of the purified ACTIF protein. To check the oligomeric state of the ACTIF protein in solution, we performed interferometric scattering (iSCAT) microscopy (Young et al. 2018). With iSCAT the light scattering of molecules near a surface is amplified with reference light, typically the reflection of the laser on the bottom of the glass slide (Supplemental Figure 1). Background subtraction makes it possible to detect the molecules landing on or moving away from the surface. The detected signal is directly proportional to the molecular weight of the proteins (Young et al. 2018). As shown

in Fig. 1B), the mass histogram confirms that the crosslinker is dimeric, as shown by the peak of 98% of signal counts being centered around 243 ± 73 kDa, about twice the molecular weight of the monomer. To test the protein purity, we performed SDS-PAGE analysis after the final gel filtration chromatography step. As shown in Fig.1C, the SDS-PAGE gel shows a major band localized at the expected molecular weight of 138 kDa and a smaller second band at $\sim$90 kDa indicative of some protein degradation.

Before testing the crosslinking activity of ACTIF, we first wanted to test whether F-actin and vimentin have any direct interactions under our experimental conditions. This idea was motivated by earlier findings that vimentin can bind F-actin with its C-terminal tails (Esue et al. 2006). To test for interactions, we used a TIRF microscopy assay (Fig. 1D (left)). We prepared a microfluidic flow chamber made of coverglass treated with vimentin antibodies that securely trap vimentin filaments, and with passivation agents including pluronic F127 and bovine serum albumin (BSA), to prevent unwanted protein binding to the substrate. Pre-assembled vimentin filaments fluorescently labeled with Alexa Fluor 555 were next introduced into the chamber and allowed to adhere to the substrate. Next, we flowed in ATTO-647-labeled G-actin monomers in F-buffer supplemented with 0.2 % methylcellulose, to polymerize F-actin directly in the chamber. Methylcellulose was used to force the growing F-actin filaments close to the substrate with vimentin. As shown in Fig. 1D (right), we did not observe any signs of co-localization between F-actin and the fixed vimentin filaments from the beginning of the F-actin polymerization until they became long filaments. Time-lapse movies confirmed that the F-actin filaments are freely moving (Supporting Movie S1). We conclude that vimentin and F-actin do not interact directly under the conditions of our assay.

Having established that vimentin and F-actin do not interact directly, we could then test whether ACTIF induces interactions. We again used a TIRF imaging assay where we sequentially flushed in the different components (Fig. 1E (left)). We first flushed in 37 nM of pre-assembled vimentin and allowed these filaments to adhere to the vimentin antibodies on the substrate. When we next flushed in 5.3 nM of ACTIF, we observed that it rapidly bound to the vimentin filaments as seen by complete co-localization (Fig. 1E (middle)). Finally, we flushed in 0.1 μM of pre-assembled F-actin. Over time, we observed that the actin filaments settled and remained on top of the vimentin filaments (Fig. 1E (right)). At steady state, F-actin, vimentin, and ACTIF perfectly co-localized, as shown in the merged

image (Fig. 1E (far right) and Supporting Movie S2). Following the maximum intensity of the F-actin signal along a cross-section drawn perpendicularly to the vimentin filaments, we could follow the time evolution of successive F-actin binding events, which appeared as intensity steps (Fig. 1G). By quantifying pixel fluorescence intensities in the F-actin channel along vimentin filaments at steady states ($\sim$ 150 filaments from 2 independent experiments), we observed a complex distribution with multiple peaks indicating the presence of multiple actin filaments (Fig. 1H), different from the single peak distribution obtained with single vimentin (Supplemental Figure 2). As the intensity of the first peak matched the intensity expected for single F-actin, we estimated that an average of 3 actin filaments and up to 7 actin filaments could bind a vimentin filament through ACTIF crosslinking at steady state (Fig. 1H). These observations prove that ACTIF is capable of binding both F-actin and vimentin and of crosslinking the two filament types.

ACTIF robustly binds and couples vimentin and F-actin filaments

We performed a series of TIRF microscopy assays to validate the efficiency of ACTIF in crosslinking actin and vimentin filaments in real-time. In the similar TIRF microscopy and flow chamber setup as shown in Fig. 1(D and E), we flowed ACTIF at a low concentration of $c \approx 5$ nM into a channel with surface-bound vimentin filaments (Fig. 2A). We observed rapid co-localization of ACTIF with vimentin over a period of 10 min (Fig. 2B). Fluorescence quantification showed that ACTIF bound to vimentin with a reaction rate $K \approx 0.0140 \pm 0.0024$ s^{-1} (mean $\pm$ SD, averaged over 3 independent repeats) obtained by fitting the kinetics of association with a single exponential (Fig. 2D). Once the ACTIF signal reached a steady state (after 10 minutes), we performed a Fluorescence Recovery After Photobleaching (FRAP) assay (Fig. 2A) to measure the ACTIF unbinding rate, assuming that the recovery is reaction-limited (Bulinski et al. 2001). As shown in Fig. 2C and quantified in Fig. 2E, we observed about 30 % of ACTIF is mobile ($N = 3$ independent repeats). We presume that the mobile fraction corresponds to dimeric ACTIF and the immobile fraction to higher oligomerization of ACTIF induced by attachment to vimentin, which increases protein avidity. By fitting the recovery curve to a single exponential, we estimated the unbinding rate of the mobile fraction of ACTIF from vimentin filaments of $k_{\mathrm{off}} \approx 0.0032 \pm 0.0012$ s^{-1} (mean $\pm$ SD, $N = 3$ repeats). With the assumption of a first-order

FIG. 2: TIRF microscopy experiment to elucidate the crosslinking efficiency of ACTIF. A) Assay to measure the binding rate of ACTIF to surface-bound vimentin by imaging (left) and the unbinding rate by FRAP (right). B) Still images at $t = 0$, directly after flushing in 5 nM ACTIF, and at $t = 10$ min, showing ACTIF (green) co-localizing with vimentin filaments (magenta). C) Images showing the recovery of ACTIF fluorescence at $t = 20$ min after photobleaching the region boxed by white dashed lines at $t = 0$. D) Mean fluorescence intensity of ACTIF on vimentin filaments (averaged over 30 filaments) over time, obtained from the representative repeat shown in Fig. 2(B). The data is fitted by an exponential equation, $I = I_{eq1} \left(1 - e^{-Kt}\right)$. E) Recovery of fluorescence intensity of ACTIF on vimentin filaments (mean fluorescence averaged over 30 filaments) over time after photobleaching, obtained from the representative repeat shown in Fig. 2(C). The data is fitted by an exponential equation, $I = I_{eq2} \left(1 - e^{-k_{off}t}\right)$. F) Assay and images of steady-state vimentin-ACTIF-F-actin composites with surface-bound vimentin. G) Normalized fluorescence intensity of ACTIF in the actin-vimentin composites (green) over time after flushing in F-actin while flushing out solution-phase ACTIF (at $t = 0$), and on vimentin (orange) over time after flushing out solution-phase ACTIF (averaged over 30 and 50 filaments respectively). H) Assay to measure the binding rate of ACTIF to surface-bound F-actin by imaging (left) and the unbinding rate by FRAP (right). I, J) Images of ACTIF (green) co-localizing with actin filaments (cyan) at $t = 10$ min after flushing in ACTIF at a concentration of ∼5 nM (I) and ∼500 nM (J). K) Images showing the recovery of ACTIF fluorescence at 500 nM on F-actin at $t = 15$ min after photobleaching the region boxed by white dashed lines at $t = 0$. L) Fluorescence intensity of ACTIF at concentrations of 5 nM and 500 nM on F-actin (both conditions averaged over 30 filaments) over time, obtained from the representative repeat shown in Fig. 2(J). The data is fitted by an exponential equation, $I = I_{eq1} \left(1 - e^{-Kt}\right)$. M) Recovery of fluorescence intensity of ACTIF at 500 nM on F-actin (averaged over 30 filaments) over time after photobleaching, obtained from the representative repeat shown in Fig. 2(K). The data is fitted by an exponential equation, $I = I_{eq2} \left(1 - e^{-k_{off}t}\right)$. N) Assay and images of steady-state actin-ACTIF-vimentin composites with surface-bound F-actin. O) Normalized fluorescence intensity of ACTIF in the composites over time, after flushing in F-actin while flushing out solution-phase ACTIF at $t = 0$ (averaged over 30 filaments). Shaded areas in all the graphs show the standard deviations over 30 to 50 filaments. Dashed lines show exponential fits for the kinetics of association and for FRAP recovery curves. Scale bars: 5 μm.

reaction and that 30 % ACTIF is mobile, the rate K of association kinetics can be approx-

imated by: $K = k_{\text{on}}\, c + f_{\text{m}}\, k_{\text{off}}$, where k_{on} is the binding rate and f_{m} is the mobile fraction. From the concentration c, the rate K and the unbinding rate k_{off} estimated by FRAP, we estimated the binding rate of ACTIF to vimentin $k_{\text{on}} \approx (2.4 \pm 0.5) \cdot 10^6$ M^{-1}s^{-1} (mean $\pm$ SD, $N = 3$ repeats), and the affinity described by a dissociation equilibrium constant $K_d \approx 1.4 \pm 0.6$ nM ($N = 3$ repeats). We next tested the F-actin crosslinking ability of the vimentin-bound ACTIF by flowing in pre-assembled F-actin in a buffer without methylcellulose. We observed that the F-actin filaments quickly attached and localized along vimentin filaments, forming composites of F-actin-ACTIF-vimentin (Fig. 2F). Interestingly, once the composites were established, ACTIF was fixed inside the composites as its fluorescence intensity remained constant over time even though no ACTIF was supplied in solution (Fig. 2G, green curve). We verified that ACTIF partially dissociated from vimentin in the absence of F-actin when the ACTIF soluble pool was flushed out (Fig. 2G, orange curve), indicating that ACTIF is stabilized on vimentin through its binding to actin. Our findings suggest strong binding of ACTIF to vimentin filaments and stable crosslinking between vimentin and F-actin filaments.

To study the binding and unbinding rates of ACTIF on F-actin, we modified the experimental setup by immobilizing pre-assembled biotinylated actin filaments on the surface of the flow chamber, which had been coated with neutravidin, and the passivation agents F127 and BSA, respectively. Next, we flowed in ACTIF to characterize the kinetics of association to F-actin, then performed FRAP to measure the unbinding rate from F-actin and deduce the binding rate to F-actin, and finally flowed in vimentin filaments to create F-actin-ACTIF-vimentin composites (Fig. 2(H, N)). When we flowed in ACTIF at the same concentration of $\sim$5 nM, as in the earlier experiment with immobilized vimentin, we did not observe any signs of binding to the immobilized F-actin (Fig. 2(I, L) after 10 minutes. To our surprise, we needed to flow in $c \approx 500$ nM of ACTIF to achieve a final ACTIF fluorescence intensity equivalent to its intensity on vimentin achieved with a 100-fold lower ACTIF concentration (Fig. 2J). Although it is expected that the number of binding sites for ACTIF on F-actin is not the same as that on vimentin (see Discussion), the difference by a factor of 100 in concentration indicates the binding rate is much lower on actin than on vimentin. By fitting the time-dependent increase of the ACTIF fluorescence intensity to a single exponential, we obtained a reaction rate $K \approx 0.0080 \pm 0.0013$ s^{-1} (mean $\pm$ SD, $N = 3$ repeats) (Fig. 2L). FRAP measurements on the actin-bound ACTIF allowed us to

TABLE 1: Summary of (un)binding rate constants and affinities of ACTIF towards vimentin and actin filaments measured by TIRF imaging and FRAP assays. Values shown in Repeat 1-3 are means averaged over 30 filaments calculated for each independent repeat experiment. The combined "Mean $\pm$ SD" represents the averaged mean from 3 repeats with SD being the standard deviation of the means.

	Repeat #	K (s^{-1})	k_{on} (M^{-1}s^{-1})	% mobile	Mobile fraction	
				f_{m}	k_{off} (s^{-1})	K_{d} (nM)
ACTIF	1	0.0111	$2.0 \cdot 10^6$	25 %	0.0023	1.2
on	2	0.0170	$3.1 \cdot 10^6$	27 %	0.0025	0.8
vimentin	3	0.0138	$2.2 \cdot 10^6$	40 %	0.0049	2.2
$c = 5.3$ nM	Mean $\pm$ SD	0.0140 ± 0.0024	$(2.4 \pm 0.5) \cdot 10^6$	(31 ± 7) %	0.0032 ± 0.0012	1.4 ± 0.6
ACTIF	1	0.0079	$7.7 \cdot 10^3$	56 %	0.0068	883
on	2	0.0096	$13.6 \cdot 10^3$	32 %	0.0075	551
F-actin	3	0.0065	$7.2 \cdot 10^3$	55 %	0.0049	681
$c = 530$ nM	Mean $\pm$ SD	0.0080 ± 0.0013	$(9.5 \pm 2.9) \cdot 10^3$	(48 ± 11) %	0.0064 ± 0.0011	705 ± 137

estimate the mobile ACTIF unbinding rate of $k_{\mathrm{off}} \approx 0.0064 \pm 0.0011$ s^{-1} from F-actin, and a mobile fraction of about 48 % ($N = 3$) (Fig. 2K, M). From the values of K, c, k_{off} and f_{m}, we estimated the binding rate to F-actin $k_{\mathrm{on}} \approx (9.5 \pm 2.9) \cdot 10^3$ M^{-1}s^{-1} and the affinity to F-actin described by $K_d \approx 705 \pm 137$ nM. When we finally flushed in pre-assembled vimentin filaments, we again obtained complete co-localization of the vimentin filaments with the ACTIF-saturated F-actin, reaffirming the robust formation of the composite (Fig. 2N). Time-lapse imaging also again showed the ACTIF signal in the composite structure to be constant, even though no ACTIF was supplied in the solution (Fig. 2O). Overall, our results indicate the binding of ACTIF to F-actin is about 500 times lower than its affinity to vimentin (summary in Table 1).

The plectin IFBD induces vimentin bundling

To test whether the isolated plectin IFBD domain is capable of binding or bundling vimentin, we expressed and purified this domain, comprising PRD 5 and PRD 6 and the intervening linker region that provides the major IF-binding site of plectin (Nikolic et al.

1996). iSCAT imaging showed the isolated IFBD to form mostly monomers with a small (13%) fraction of dimers (Supplemental Figure 3A) and SDS-PAGE analysis showed a major band at the expected molecular weight of 71 kDa (Supplemental Figure 3B). Next, we co-polymerized the IFBD with vimentin filaments at different IFBD:vimentin molar ratios, always keeping the vimentin molar concentration fixed at 4 μM, and imaged the mixtures by electron microscopy (EM). Vimentin filaments reconstituted in the absence of the IFBD were visible in EM images as semiflexible filaments (Fig. 3A (left)). To determine the filament widths, we plotted intensity profiles perpendicular to the segment of interest at multiple locations along the filament contours. We then measured the width of the bright

FIG. 3: Electron microscopy (EM) and rheology experiments to test binding and crosslinking activities of ACTIF and its plectin IFBD. A) EM images of reconstituted vimentin filaments alone (left) or co-polymerized with 0.4 μM (middle) or 4 μM (right) of IFBD (schematic on top). B) Corresponding boxplots of vimentin bundle widths. *** $p < 0.001$ from unpaired t-test. C) EM images of co-polymerized F-actin (cyan) and vimentin filaments (pink) without ACTIF (left) or with 0.02 μM (middle) or 0.2 μM (right) of ACTIF (schematic on top). D) Corresponding boxplots of the F-actin/vimentin co-bundles. **** $p < 0.0001$ from unpaired t-test. E) (Top) Frequency dependence of the storage modulus of crosslinked (green) and non-crosslinked (grey) actin/vimentin composite networks and of one-component actin (blue) or vimentin (pink) networks. (Bottom) Corresponding loss tangent, defined as the ratio of the loss modulus G'' to the storage modulus G' (same color code). Note that the loss tangent is noisy because the applied strain and the shear moduli (especially G'') are small, so the stress is close to the sensitivity limit of the rheometer. F) Bar plots comparing the elastic modulus G' at a frequency of 1 Hz of entangled and crosslinked networks of actin, vimentin, and actin/vimentin composites. For each condition, $N = 2$, circular data points. All scale bars are 100 nm, except in the far left image of panel A, the scale bar is 1000 nm.

area between the darker borders, which corresponds to the filament diameter. We observed an average filament width of $\sim$10 μm (Supplemental Figure 4), in agreement with previous literature (Wen and Janmey 2011). Surprisingly, when vimentin was co-polymerized with the IFBD at a molar ratio of IFBD:vimentin of 1:10 (0.4 μM of IFBD), it formed thin bundles (see Fig. 3A (middle)). Bundles often contained around three to four distinguishable filaments. Using the same procedure as described above to determine the bundle widths, we found that the bundles had a variable width with an average bundle size of 33 ± 12 nm (Fig. 3B). Upon increasing the IFBD:vimentin molar ratio to 1:1 (4 μM of IFBD), vimentin filaments formed straight and well-aligned bundles (Fig. 3A (right)) of an average diameter of 58 ± 22 nm (Fig. 3B). The bundles formed at a 1:1 IFBD:vimentin ratio were significantly thicker than those formed at a 1:10 ratio ($p < 0.0001$, calculated by a statistical unpaired t-test). We conclude that the IFBD domain can by itself bundle vimentin filaments at high (> 400 nM) concentrations. Presumably, bundling requires dimerization or clustering of the IFBD domain, which we showed to be mostly monomeric (with only 13 % dimeric) in iSCAT measurements performed at a concentration of 40 nM (Supplemental Figure 3A). Potentially, at the high (> 400 nM) concentrations used in the EM assays, the IFBD domain

can oligomerize even more. As a control, we also tested co-polymerized actin and IFBD. In this case, EM images showed no signs of actin bundling, but plectin was present in the form of dense globular structures, suggesting that the IFBD can cluster at high concentrations (Supplemental Figure 5).

ACTIF forms bundles of F-actin and vimentin filaments

With the certainty based on EM data that the IFBD from plectin can bind vimentin filaments, we next imaged mixtures of F-actin and vimentin filaments with and without ACTIF, always keeping the monomer concentrations at 1 μM for actin and 1μM for vimentin. In the absence of ACTIF, actin filaments can be distinguished from vimentin filaments by their smaller diameter (8 nm vs 10 nm) (Wen and Janmey 2011)) and by their larger persistence length (10 μm vs 1 μm (Wen and Janmey 2011)). As shown in Figure 3 C(left), the two filament types can indeed be easily distinguished (superposed blue line for F-actin and superposed pink line for vimentin filaments). Upon adding ACTIF in a molar ratio of 1:100 to the sum of the vimentin and actin molar concentrations, we found that tightly packed bundles formed (Fig. 3C (middle)) with an average diameter of 66 $\pm$ 34 nm (Fig. 3D). Note that bundling of actin and vimentin by ACTIF occurs at a much lower crosslinker:monomer ratio than bundling of vimentin by the IFBD domain, consistent with ACTIF being a parallel dimer presenting two actin-binding domains on one end and two IF-binding domains on the other end. At a ten-fold larger ACTIF:monomer ratio of 1:10 (0.2 μM of ACTIF), even thicker bundles were observed that consisted of a larger number of filaments (Fig. 3C (right)). The average bundle size was significantly larger (131 $\pm$ 66 nm) than for the bundles formed at the 1:10 molar ratio (Fig. 3D), $p < 0.00001$. Many bundles were too dense to observe their constituent filaments, but in some cases we could observe bundles clearly consisting of a mixture of actin and vimentin filaments (Supplemental Figure 6A), some bundles eventually splitting into two or more sub-bundles (Supplemental Figure 6B) and some bundles apparently consisting only of actin filaments (Supplemental Figure 6C). We conclude that ACTIF drives the formation of mixed actin-vimentin bundles in a concentration-dependent manner.

Composite F-actin/vimentin networks stiffen upon crosslinking with ACTIF

To test whether ACTIF is capable of crosslinking F-actin and vimentin in dense networks, we used shear rheology to measure the viscoelastic properties of composite networks formed with or without ACTIF. The viscoelastic response of cytoskeletal networks is known to be sensitive to the presence of crosslinkers, which tend to enhance the elastic modulus and suppress viscous dissipation (Ward et al. 2008). We studied the linear viscoelastic response by applying small amplitude oscillatory shear with a strain amplitude of 1% while varying the frequency logarithmically between 0.001 and 10 Hz. We determined the network elastic (storage) shear modulus, G', and viscous (loss) modulus, G'', from the stress response. In the absence of ACTIF, the composite networks had a small elastic modulus of order ~ 10 Pa that increased very weakly as an approximate power law with frequency with an exponent of 0.07 (grey curve in Fig. 3E). In the presence of ACTIF, the networks were more than an order of magnitude stiffer (green curve in Fig. 3E), as summarized in Fig. 3F. The frequency dependence for the crosslinked network again followed a weak power law with an exponent of 0.07. Simultaneously, the network with ACTIF was more solid-like, as characterized by a smaller loss tangent, $\tan(\delta) = \frac{G''}{G'}$, compared to the corresponding network without ACTIF (Fig. 3F)). The rheology measurements hence prove that ACTIF crosslinks the composite network.

To test whether there is mechanical synergy upon combining actin and vimentin, we also performed rheology measurements for isolated actin networks (blue curve in Fig. 3E) and vimentin networks (pink curve in Fig. 3E). The stiffness of the actin/vimentin composite network formed without ACTIF was approximately equal to the sum of the elastic moduli of the two isolated networks (Fig. 3F). Interestingly, when actin was polymerized in the presence of ACTIF, it formed stiffer networks than in the absence of ACTIF (Supplemental Figure 7A), indicating that ACTIF is capable of crosslinking actin filaments. By contrast, when vimentin was polymerized in the presence of ACTIF, it formed a slightly softer network than in the absence of ACTIF (Supplemental Figure 7B). This suggests that either ACTIF does not crosslink vimentin to vimentin, or that the crosslinking effect of vimentin's C-terminal tail domains (Aufderhorst-Roberts and Koenderink 2019) dominates over the crosslinking effect of ACTIF. The stiffness of the crosslinked actin/vimentin composite network is larger than the sum of the moduli of the two isolated networks (Fig. 3F), suggesting

FIG. 4: ACTIF crosslinker with the IFBD of APC binds and crosslinks F-actin and vimentin. A) Schematic representation of the engineered crosslinker ACTIF-APCn2, which is identical to ACTIF aside from its IFBD domain consisting of the armadillo repeats of APC. B) EM image of actin 1.5 μM and vimentin 1.5 μM co-polymerized in the absence (left) and presence (right) of 0.3 μM ACTIF-APCn2. ACTIF-APCn2 forms thick straight bundles. C) Schematic and fluorescent images from the TIRF microscopy assay following the same protocol as Fig. 1(E), showing F-actin was added in solution and recruited on top of ACTIF-APCn2-bound vimentin after ACTIF-APCn2 at $\sim$5 nM was flushed into the chamber with vimentin filaments on the substrate.

the presence of some form of mechanical synergy.

ACTIF crosslinker variant with the IFBD of APC

The engineered ACTIF crosslinker has a modular design such that its functional modules can be swapped for alternative domains. To prove the benefit of this modularity, we swapped the IFBD from plectin for an alternative IFBD, taken from the Adenomatous Polyposis Coli (APC) protein. APC was reported to directly bind vimentin filaments in astrocytes, leading to crosslinking and elongation of vimentin filaments along microtubules (Sakamoto et al. 2013). IF-binding was traced to the N-terminal Armadillo repeats of APC via biochemical assays with truncated APC proteins. We hence engineered an ACTIF variant where the

IFBD from plectin was substituted by the armadillo repeats of APC, specifically, the region referred to as APCn2 (Sakamoto et al. 2013). The ACTIF-APCn2 crosslinker therefore consisted of an N-terminal GFP-tag, followed by the F-actin binding domain of MACF2, the cortexillin I coiled-coil to induce dimerization, and finally a C-terminal region containing the armadillo repeats of APC (Fig. 4A). We performed iSCAT to check the oligomeric state of the ACTIF-APCn2 crosslinker and found it to be dimeric (Supplemental Figure 3C). To test the protein purity, we performed SDS-PAGE analysis after the final gel filtration chromatography step. As shown in Supplemental Figure 3D, the SDS-PAGE gel shows a major band localized at the expected molecular weight of 116 kDa and a smaller second band at ~80 kDa indicative of some protein degradation.

We first performed EM assays to test the ability of ACTIF-APCn2 to crosslink F-actin and vimentin filaments. We compared actin and vimentin mixtures co-polymerized in the absence (Fig. 4B (left)) and presence (Fig. 4B (right)) of 0.3 μM of ACTIF-APCn2. ACTIF-APCn2 caused the formation of thick straight bundles of F-actin and vimentin filaments, showing that it is capable of crosslinking the two filament types. We next performed TIRF microscopy to validate the efficacy of ACTIF-APCn2 in crosslinking actin and vimentin filaments in real-time. After immobilizing vimentin filaments on a substrate as described earlier, we sequentially flushed in ACTIF-APCn2 (~5 nM) for 10 min, then pre-assembled F-actin. We observed the binding of ACTIF-APCn2 to vimentin, and progressive recruitment of F-actin on top of the ACTIF-APC2-decorated vimentin filaments (Fig. 4C), forming a stable actin-ACTIF-APCn2-vimentin composite. When we characterized the ACTIF-APCn2 association on vimentin at a similar concentration as for ACTIF (5 nM), we noted that ACTIF-APCn2 associated at a rate about 60 times lower than ACTIF during the initial phase of fluorescence intensity increase (Supplemental Figure 8). We conclude that the type of IFBD impacts the association rate of the ACTIF on vimentin.

3. DISCUSSION

In this work, we reconstituted specific interactions between actin and vimentin filaments by engineering a cytolinker protein that recapitulates the crosslinking activity of plectin. We found that the engineered cytolinker protein, which we named ACTIF, has a high binding affinity ($K_d \approx 1.4$ nM) for vimentin and lower binding affinity for actin ($K_d \approx 0.7$ μM). TIRF

imaging showed that ACTIF molecules very stably crosslink actin and vimentin filaments since we did not observe any dissociation when we removed the soluble ACTIF pool, while ACTIF partially dissociated when only attached to vimentin. ACTIF is thus able to form long-lived composite actin-vimentin bundles.

The affinity of ACTIF on vimentin and on actin was estimated from the mobile fraction of ACTIF in FRAP experiments, which we believe is dimeric (∼30% on vimentin and ∼50% on actin). We presume that the high proportion of immobile ACTIF results from oligomerization induced by the high density of ACTIF on the filaments, which increases the strength of the bond between the oligomers and the filaments. This could explain why we only observed one population of dimers of ACTIF with the iSCAT experiments in similar conditions (concentration and buffer), but two populations on filaments (dimers and oligomers induced by the high density on filaments). We believe that the ACTIF immobile fraction is more important on vimentin than on F-actin due to the higher density of binding sites on vimentin (at least 800 per micron length of vimentin filaments made of 10 tetramers per cross-section (Eibauer et al. 2023) and an axial repeat of 50 nm (Nunes Vicente et al. 2022), compared to 360 per micron length of F-actin (Bremer et al. 1991) if we assume one ACTIF per monomer of vimentin or actin).

The high affinity of ACTIF for vimentin is qualitatively consistent with previous biochemical measurements using different plectin fragments, showing that the PRD5 and PRD6 domains and intervening linker constitute the main vimentin binding site. In quantitative terms, ACTIF under the conditions of our assays had a higher binding affinity for vimentin as compared to values reported previously for these fragments, specifically fragments consisting of the PRD5 domain and linker ($K_\mathrm{d} = 50$ nM) (Nikolic et al. 1996), the PRD5-PRD6 (including the linker region) ($K_\mathrm{d} = 0.15 - 0.3$ μM) (Spurny et al. 2007), or the entire C-terminus starting from the PRD5 domain ($K_\mathrm{d} = 135$ nM) (Bouameur et al. 2014). Variants of ACTIF with IFBDs composed of different combinations of domains could be used in the future to sort out the contribution and synergy of the different domains of plectin implicated in controlling its affinity for vimentin. Also, ACTIF variants could be used for testing the binding affinity and crosslinking efficiency of PRD and linker domains from other plakin proteins. Sequence analysis has shown that these domains differ in the number of basic residues in their IF-binding groove, suggesting that they present a spectrum from weak to strong binding affinities (Mohammed et al. 2020). The lower binding affinity of ACTIF's ABD

for F-actin is consistent with previous measurements for TipAct (López et al. 2014), the engineered actin-microtubule crosslinker that formed the template for ACTIF. ACTIF has the same MACF-derived ABD as TipAct. Since both ACTIF and TipAct are dimers, this also further validates our assumption that the ACTIF mobile fraction is dimeric. Sequence analysis shows a 73.8% overlap between the ABD of MACF and the ABD of plectin (see sequence alignment in Supplemental Figure 9). Both ABDs consist of two calponin homology (CH) domains in tandem. The isolated CH1 supports binding to actin, albeit with a lower affinity than the tandem; in contrast, CH2 domain has a weaker affinity. The tandem binds at full capacity and has three conserved actin binding sites (Gimona and Winder 1998). The plectin ABD, with its CH domains in a closed conformation, has also been reported to have structural similarity with fimbrin (Sevcik et al. 2004), despite dissimilar sequences. Observations of plectin localization in cells suggest that the high binding affinity for vimentin and low affinity for actin of ACTIF are physiologically relevant. EM images showed that in IF-containing cells, plectin co-localized largely with the vimentin network, whereas in IF-deficient cells, it became mainly associated with the actin cytoskeleton (Foisner et al. 1995).

We showed by electron microscopy that ACTIF induces the formation of crosslinked actin/vimentin bundles and networks already at rather low molar ratios of just 1 ACTIF molecule per 100 actin/vimentin monomers. The isolated IFBD domain could induce bundling of vimentin filaments, but only at much higher (1:1) molar ratios of IFBD:monomer. This finding is consistent with previous work showing that the PRD5 domain by itself can crosslink vimentin (and cytokeratin) filaments (Steinböck et al. 2000). However, we note that contrary to that study, we did not observe any negative effect of the isolated IFBD domain on vimentin filament formation. The fact that the IFBD domain alone can bundle vimentin filaments suggests that it may oligomerize. Indeed in EM images of actin filaments mixed with the IFBD domain, we observed globular structures (non-interacting with F-actin), indicating the presence of IFBD clusters. Oligomerization of full-length plectin has been previously observed by electron microscopy images of rotary-shadowed samples, where filament-bound plectin structures were shown to be oligomeric (Foisner et al. 1988). ACTIF not only forms crosslinks between actin and vimentin filaments, but it also forms crosslinks between actin filaments. This finding is consistent with previous work showing that the ABD of plectin by itself can bundle actin filaments (Fontao et al. 2001). In this

earlier study, actin bundling relied on dimerization of the ABD. In the case of ACTIF, we enforced dimerization using a cortexillin coiled-coil spacer domain. We cannot exclude that, additionally, interactions between the ABP domains may also contribute to the observed ACTIF-mediated F-actin bundling.

The cell-free reconstitution assay we introduced here can provide a strong basis for understanding the regulation and functions of cytolinker-mediated actin/intermediate filament crosslinking in cells. First, it will be interesting to compare binding and crosslinking by ACTIF and ACTIF-APCn2 with different types of intermediate filaments (IFs), given that the expression pattern of IF proteins is cell-type specific and may convey cell-specific cellular functions. Plectin and other plakins have been shown to possess a broad binding specificity towards different types of IFs, including not only vimentin, which we studied in this work, but also cytokeratins, desmin, GFAP and neurofilaments (De Pascalis et al. 2018, Favre et al. 2011, Foisner et al. 1988, Steinböck et al. 2000), because they recognize the conserved rod domain of IF proteins. By contrast, APC was reported to interact with vimentin and GFAP, but not with keratin, although the structural basis for this is still unknown (Sakamoto et al. 2013). Our system makes it possible to perform quantitative comparisons of the kinetic (un)binding rates and the crosslinking efficiency of different IF types. Second, it is also relevant to uncover how posttranslational modifications (PTMs) of IFs such as phosphorylation (Kraxner et al. 2021), glycosyation (Tarbet et al. 2018) and electrophilic or oxidative modifications (González-Jiménez et al. 2023) may affect interactions with ACTIF and ACTIF-APCn2. Conversely, it would also be interesting to test how PTMs of plectin regulate its interactions with IFs. Cell studies have shown evidence for regulation by phosphorylation of sites like serine 4642 in the C-terminal extremity (Bouameur et al. 2013, Foisner et al. 1996, 1991) and by oxidation and nitrosylation of cysteines in plectin's IFBD (Spurny et al. 2007). Finally, a recent study showed that the association between plectin and vimentin is mechanosensitive and requires actomyosin contractility (Marks et al. 2022). Our reconstitution assay could be used to test whether this mechanosensitivity is intrinsic to plectin, for instance to its force-sensing plakin domain (Daday et al. 2017).

Our work also provides an interesting proof-of-concept for constructing cytoskeletal systems to be applied in active matter model systems (Berezney et al. 2022, Sanchez et al. 2012) or in synthetic cells (Mulla et al. 2018, Weiss et al. 2018). In these systems, designer crosslinkers like ACTIF could be used to manipulate the material's mechanical properties

and associated dynamics. An interesting expansion of our work would be to engineer a light-activatable cytolinker variant, where the crosslinking effect can be spatiotemporally controlled by illumination.

4. CONCLUSIONS

Our results reveal specific crosslinking of actin and vimentin filaments mediated by the engineered crosslinker ACTIF. Actin and vimentin filaments form composite bundles and stiff networks upon crosslinking via ACTIF. This engineered crosslinker enables biochemical, structural, and biophysical studies of crosslinker-mediated cytoskeletal interactions. The crosslinker's modular design provides an opportunity to test different IF binding domains in order to better understand intermediate filament binding, as demonstrated by our proof of concept comparing the IFBDs from plectin and APC. This approach also provides the opportunity to unravel the effect that actin-vimentin crosstalk exerts on the mechanical properties of the cytoskeleton, either in a simplified *in vitro* system as reported here, or directly in cells.

5. METHODS

5.1. Proteins

5.1.1. ACTIF Plasmid Construction

ACTIF was designed to contain an enhanced Green Fluorescent Protein (eGFP) tag followed by the ABD from MACF, the coiled-coiled linker of Cortexillin I, and the IFBD of plectin. We inserted additional residues for flexibility between the different domains, as shown in Supplemental Figure 10. As a vector, we used the TipAct construct (López et al. 2014). For the IFBD insert, we used a plasmid corresponding to residues 4030-4631 aa of the isoform Q9QXS1-1 of mouse plectin, a region that contains the plakin repeat domains 5 and 6 and intervening linker(Nikolic et al. 1996). The two fragments (vector and insert) were amplified using the primers indicated in Supplemental Table 1. PCR amplification was performed using KOD Xtreme Hot start DNA polymerase (#1975). The PCR products were run on a 1% agarose gel (Biorad #1613100EDU) and cleaned using the Wizard® SV

Gel and PCR Clean-Up System A9281. For electrophoresis, samples were prepared with TrackIt™ Cyan/Yellow Loading Buffer (Invitrogen #10482035), and TrackIt™ 1 Kb Plus DNA Ladder (Invitrogen #10488085) was used as a ladder. The electrophoresis gel was run at 100V for 20 minutes (Biorad #1613100EDU). Next, the fragments were assembled using the Gibson assembly, following the instructions of NEBuilder HiFi DNA Assembly Cloning Kit (#E5520S). Afterward, the sample was transformed into Dh5α competent *E. coli* cells (#C2987). To determine if the plasmid contained the insert, a miniprep was performed for sequencing via Macrogen. The miniprep was performed using the PureYield™ Plasmid Miniprep System A1222. After sequencing, the plasmid was transformed into *E. coli* BL21 (NEB #C2527).

5.1.2. ACTIF Protein Purification

A pre-culture of *E. coli* BL21 cells expressing ACTIF was grown from a glycerol stock in Luria-Bertani (LB) medium containing 1:1000 diluted kanamycin at 37 °C. Growth was upscaled to a total of 12 liters (separated in 4 cultures of 3 L each), adding 1:1000 diluted kanamycin, and 25 mL of ACTIF preculture. The cultures were incubated at 37 °C in flasks on a shaker platform at 200 rpm. When the Optical Density (OD) reached 0.9, we cooled the 4 cultures down in an ice bath for 30 minutes, then induced overnight expression with 1 mM Isopropyl β-D-1-thiogalactopyranoside (IPTG #11411446001) at 16 °C. The cultures were harvested by a 15 min centrifugation in the Avanti JLA8.1000 fixed angle rotor at 4000 rpm at 25 °C. The supernatant was discarded, and the cell pellets were combined in one 50 mL falcon tube. The cells were resuspended with lysis buffer (20 mM sodium phosphate pH 7.4, 10 % glycerol, 5 mM 2-mercaptoethanol, 500 mM NaCl, 45 mM imidazole), supplemented with cOmpleteTM EDTA-free protease inhibitor cocktail (SigmaAldrich, St. Louis, MO, USA #11836153001) and 1 mg/mL of lysozyme from chicken egg white (Sigma Aldrich #10837059001). The cell solution was incubated on ice for 30 minutes. The lysate was passed through the French press three times at 20 kpsi and next centrifuged at $17,000 \times g$ for 1 hour and 20 minutes at 4 °C (Avanti JXN-26). The supernatant was collected and incubated with 1 mL pre-washed nickel-IMAC resin (Thermo Fisher Scientific #88221) overnight at 4 °C. The beads were pre-washed with wash buffer (20 mM sodium phosphate pH 7.4, 10% glycerol, 5 mM 2-mercaptoethanol, 500 mM NaCl, 45 mM imidazole) in an empty

column (Econo-Pac® Chromatography Columns #7321010). The lysate was passed through the disposable column for gravity flow purification. The flow-through was collected and kept at 4 °C. The column was washed three times with wash buffer. To elute the protein from the column, the column was incubated with elution buffer (20 mM sodium phosphate pH 7.4, 10% glycerol, 5 mM 2-mercaptoethanol, 500 mM NaCl, 200 mM imidazole) for 20 minutes at 4 °C. Afterward, an analysis of protein yield and purity for the eluate was performed by SDS-PAGE analysis. To improve the protein purity, a final size exclusion chromatography step was performed on a Superdex 200 10/300 prep grade (Cytiva) column. The column was previously stored in ethanol, and it was washed with 1.5× column volume of MiliQ and then with 1.5× column volume of MRB40 buffer ($40mM$ piperazine-N,N′-bis(2-ethanesulfonic acid) (PIPES) pH 7.0, 4 mM MgCl2, and 1 mM Ethylene glycol tetraacetic acid (EGTA)). After the column was equilibrated, we injected the sample through the 5 mL loop and ran it at a flow rate of 0.5 mL/min, while collecting 0.3 mL fractions. The fractions of interest as judged from the absorbance at 280 nm were pooled and analyzed by SDS-PAGE gel electrophoresis. The sample was aliquoted at the desired molar concentration, calculated using values for the molecular weight and extinction coefficient of 138 kDa and 101,245 $M^{-1}\,cm^{-1}$, respectively, which were theoretically estimated with Expasy analysis tool (Walker 2005). The aliquots were stored at -80 °C. We typically obtained 1 mL at 7 mg from 12 L of bacterial cell culture.

5.1.3. IFBD(plectin) Construction

The IFBD(plectin) construct consisted of an N-terminal His6-tag followed by the region of the isoform Q9QXS1-1 of mouse plectin of 4030-4631 aa, corresponding to the plakin repeat domains 5 and 6 and intervening linkers, as shown in Supplemental Figure 11. The plasmid assembly was performed as reported for ACTIF, and the used primers are presented in Supplemental Table 3.

5.1.4. ACTIF-APCn2 Construction

ACTIF-APCn2 was designed as a variant of ACTIF, with the plectin-derived IFBD replaced by the armadillo repeats of APC, again using the TipAct construct (Alkemade et al.

2022, López et al. 2014) as a vector. For the insert, we used a plasmid courtesy of Sandrine Etienne-Manneville that corresponded to the APCn2 fragment of APC, specifically from residue 334 to 740 (Sakamoto et al. 2013). The assembly was performed as reported for ACTIF using the primers presented in Supplemental Table 2. The final sequence is shown in Supplemental Figure 12 .

5.1.5. IFBD(plectin) protein purification

IFBD(plectin) was purified following the same protocol as for ACTIF, but no gel filtration was required. In order to calculate the final concentration of the purified sample, the respective molecular weight and extinction coefficients are 71 kDa and 48,165 $M^{-1} cm^{-1}$, which were theoretically estimated with Expasy analysis tool (Walker 2005). We typically obtained 5 mg from 6 L of bacterial cell culture.

5.1.6. ACTIF-APCn2 protein purification

ACTIF-APCn2 was purified following the exact same protocol as ACTIF until the lysis step. Culture pellets were resuspended in 40 mL of lysis buffer (20 mM sodium phosphate pH 7.4, 500 mM NaCl, 45 mM imidazole, 5 mM bME, 10 % v/v glycerol), 1 cOmplete™, EDTA-free Protease Inhibitor Cocktail (Roche) and 1 mg/mL of lysozyme from chicken egg white (Sigma Aldrich). Lysis of the spun down cultures was performed by sonication, using a 422A tip, with a protocol following amplitude 40 % , 10 seconds on, 10 seconds off, for a duration of 10 minutes. Afterwards, the lysate was centrifuged using TI45 40,000 rpm, for 30 min at 4 °C. The supernatant was loaded on a 5mL HisTrap FF column (Cytiva) and the purification was performed using the AKTA system. The column was washed with 50 mL lysis buffer and was eluted with 50 mL lysis buffer supplemented with 200mM imidazole. The flow rate of the buffers was 1 mL/min. Fractions with the clean protein based on the SDS-gel were pooled. In order to calculate the final concentration of the purified sample, the respective molecular weight and extinction coefficients are 116 kDa and 94,770 $M^{-1} cm^{-1}$, which were theoretically estimated with Expasy analysis tool (Walker 2005). We typically obtained 1 mL at 6-8 mg from 12 L of bacterial cell culture.

5.1.7. Actin purification

For EM and rheology experiments, lyophilized rabbit alpha-skeletal muscle actin was purchased from Hypermol EK. Actin was dialyzed against G-buffer (pH 7.8) comprising 5 mM Tris-HCl, 0.1 mM $MgCl_2$, 0.2 mM ATP and 5 mM DTT. After dialysis, G-actin was aliquoted, flash-frozen in liquid nitrogen and stored at -80 °C. The final actin concentration was measured by UV-VIS absorbance measurements using an extinction coefficient at 290 nm of 26,600 $M^{-1} cm^{-1}$ and molecular weight of G-actin of 42 kDa (Houk Jr and Ue 1974).

For TIRF experiments, alpha-skeletal actin was purified from rabbit muscle acetone powder following a protocol in ref. (Wioland et al. 2017) that was based on an earlier protocol (Spudich and Watt 1971). G-actin was fluorescently labeled on accessible surface lysines using ATTO-643 succinimidyl ester (Life Technologies) as described previously (Romet-Lemonne et al. 2018). Similarly, actin was labeled with biotin using biotin succinimidyl ester (Life Technologies).

5.1.8. Vimentin purification

Wild-type human vimentin was purified from TG1 *E. coli* bacteria (Sigma-Aldrich) following an earlier protocol (Herrmann et al. 2004). In brief, we started the induction when the OD reached 1.2, and cultured the bacteria in Terrific Broth medium overnight at 37 °C. Then, we collected the bacterial cells by centrifuging the culture medium, and lysed them with lysozyme, together with DNase (Roche), RNase (Roche), and protein inhibitors (pefabloc and PMSF). We washed the released inclusion bodies 5 times by successive steps of centrifugation and resuspension using a cooled douncer. After washing, we resuspended the inclusion bodies in a denaturing buffer (8 M urea, 5 mM Tris pH 7.5, 1 mM EDTA, 1 mM DTT, 1 % PMSF) and centrifuged them at high speed ($100000 \times g$) for 1 h. Upon collecting the supernatant, we conducted vimentin purification by two sequential steps of exchange chromatography, using first an anionic (DEAE Sepharose, GE Healthcare) and then a cationic (CM Sepharose, GE Healthcare) column. We collected the vimentin protein in 2 mL Eppendorf tubes and monitored the protein concentration by Bradford assay. Only the most concentrated fractions were selected and pooled together. We stored the vimentin at -80 °C with an additional 10 mM methylamine hydrochloride solution.

For experiments, about 100 μL denatured vimentin aliquots were transferred to dialysis tubing (Servapor, molecular weight cut off at 12 kDa) and vimentin was renatured by stepwise dialysis from 8 M, 6 M, 4 M, 2 M, 1 M, 0 M urea into sodium phosphate buffer (pH 7.0, 2.5 mM sodium phosphate, 1 mM DTT) at room temperature with at least 15 minutes for each step. The final dialysis step was performed overnight at 4 °C in 2 L of the sodium phosphate buffer. The vimentin concentration was determined by UV-VIS absorbance measurements using a molecular weight of 53.65 kDa and extinction coefficient at 280 nm of 22,450 $M^{-1}\,cm^{-1}$ (Aufderhorst-Roberts and Koenderink 2019).

5.1.9. Fluorescent labeling of vimentin

We labeled vimentin proteins by coupling an Alexa Fluor 555 dye to cysteine-328, as described previously (Tran et al. 2023, Winheim et al. 2011). In short, denatured vimentin stored in 8 M urea was dialyzed for 3 h in a labeling buffer of pH 7.0 containing 50 mM sodium phosphate and 5 M urea. Next, the fluorescent dye (AF-555 C2 maleimide, ThermoFisher) dissolved in DMSO was added in a molar ratio of 5:1 of (dye:vimentin). The solution was gently mixed for 1 h at room temperature, and the reaction was then quenched by the addition of 1 mM DTT. We used a Dye Removal Column (#22858, ThermoFisher) to remove excess dye. Finally, we renatured the labeled vimentin by stepwise dialysis from 8 M, 6 M, 4 M, 2 M, 1 M, 0 M urea to sodium phosphate buffer (pH 7.0, 2.5 mM sodium phosphate, 1 mM DTT). The final product was stored at 4 °C and used within 10 days.

5.2. Total Internal Reflection Fluorescence Microscopy (TIRF-M)

5.2.1. Assembly of fluorescent vimentin and actin filaments

Prior to the experiment, we polymerized vimentin filaments from AF-555-labeled vimentin (with 10 % labeling fraction) at a concentration of 3.7 μM in V-buffer (2.5 mM sodium phosphate, pH 7.0, 100 mM KCl) for 3 h at 37 °C using a water bath. We also polymerized F-actin from 10 μM G-actin labeled with ATTO-643 (10 % labeling fraction) in F-buffer (20 mM Tris–HCl, pH 7.4, 50 mM KCl, 2 mM $MgCl_2$, 0.5 mM ATP) for 1 h at room temperature. In experiments that required F-actin immobilization, we assembled biotinylated F-actin from 10 μM G-actin labeled with both ATTO-643 (10 % labeling fraction)

and biotin (1 % labeling fraction) in F-buffer for 1 h at room temperature.

5.2.2. *TIRF-M experimental setup*

We assembled the composites of vimentin, F-actin and ACTIF in a flow chamber. The flow chamber was constructed from silanized coverslips following a previously described protocol (Leduc et al. 2012, Tran et al. 2023). Briefly, we incubated cleaned coverslips with dichlorodimethylsilane 0.05 % in trichloroethylene for 1 h at room temperature. We sonicated the coverslips 3 times in methanol, each for 15 min, to wash off the excess silane. We then heated up two silanized coverslips compressing pieces of parafilm in between to 50 °C on a hot plate. This made the parafilm melt and attach to the two coverslips, resulting in a flow chamber with a volume of about 10 microliters.

For the control experiment to test the actin-vimentin direct interaction, we first flowed vimentin antibodies (SC666260, Santa Cruz Biotechnology) at a concentration of 2 μg/mL in F-buffer into the chamber and incubated for 5 min. We passivated the flow chamber by flowing Pluronic F127 1 % in F-buffer and incubating for 15 min. We continued flowing bovine serum albumin (BSA) 5 % in F-buffer and incubated for 15 min. Finally, we thoroughly rinsed the chamber by flowing an extensive amount of F-buffer. We prepared ATTO-643-labeled G-actin (10 % labeling fraction) at 1 μM in F-buffer supplemented with an oxygen scavenger mixture (glucose 1.2 mg/mL, glucose oxidase 40 μg/mL, glucose catalase 8 μg/mL, DTT 1 mM) and 0.2 % methylcellulose 4000 cP, and gently flowed them into the chamber with vimentin filaments on substrate. We captured images of the whole process of F-actin polymerizing from G-actin and rapidly elongating to long filaments while freely moving close to the vimentin filaments.

For the TIRF-M assay in which vimentin filaments were attached to the substrate, we used a similar protocol to have the flow chamber coated with vimentin antibodies and passivated with F127 and BSA as described in the control experiment above. We then began the composite assembly by diluting pre-assembled vimentin filaments 100 times in F-buffer (final concentration: 37 nM), then flowing them into the chamber and incubating for 5 min for filament attachment onto the vimentin antibodies. Next, the chamber was rinsed, and ACTIF diluted in F-buffer supplemented with the oxygen scavenger mixture to a final concentration of 5.3 nM was flushed into the chamber while imaging had started in

advance in order to capture the early state of protein attachment to vimentin filaments. After 15 min, we performed FRAP assay by photo-bleaching a region of ACTIF binding to vimentin filaments and observing the fluorescence recovery over 20 min. Finally, pre-assembled actin filaments were diluted 100 times in F-buffer with an oxygen scavenger (final concentration: 0.1 μM) and flushed into the chamber, forming vimentin-ACTIF-actin composites.

For the assay with immobilized F-actin, we flowed neutravidin (Cat. #31000, ThermoFisher) at a concentration of 10 μg/mL in F-buffer into the chamber and incubated for 5 min. We again passivated the flow chamber with Pluronic F127 1 % in F-buffer and BSA 5 % in F-buffer, and thoroughly rinsed the chamber afterward, similar to the previous assay. Here, our first step in the composite assembly was diluting pre-assembled biotinylated F-actin 100 times in F-buffer (final concentration: 0.1 μM), then flushing them into the chamber, and incubating for 5 min. F-actin with biotin was constrained on the substrate due to the neutravidin coating. Then, we rinsed the chamber and flowed in diluted ACTIF at 5.3 nM or at 530 nM in F-buffer with oxygen scavenger while imaging the binding reaction for 15 min. We also performed a FRAP assay by photo-bleaching a region of ACTIF on F-actin and recording the recovery for 20 min. Finally, pre-assembled vimentin filaments diluted 100 times in F-buffer with oxygen scavenger (final concentration: 37 nM) were flowed into the chamber for the formation of actin/vimentin composite bundles.

TIRF microscopy was performed on a Nikon Eclipse Ti inverted microscope controlled by MicroManager, equipped with a 60× oil-immersion objective and a Kinetix sCMOS Camera (Photometrics). The TIRF illumination was performed using an Ilas2 module (GATACA Systems). The experiment temperature was maintained at 25 °C using objective and microscope stage temperature controllers.

5.3. Interferometric Scattering Microscopy

Interferometric Scattering Microscopy (iSCAT) experiments were conducted with a OneMP Mass Photometer (Refeyn). The instrument was equipped with a 525 nm laser for illumination. iSCAT measurements were carried out using CultureWell gaskets (Cat. #GBL103250, Grace Biolabs) affixed to #1.5 coverslips (Cat. #13296788, Corning). To prepare the coverslips for optimal protein adhesion, a meticulous stepwise sonication clean-

ing process was employed. The coverslips were sonicated sequentially in MilliQ water, 50% isopropanol, and again in MilliQ water, each for a duration of 5 minutes. Following the cleaning process, the coverslips were incubated with a poly-L-lysine solution (PLL, Cat. #P4832, Merck) for 30 seconds. Subsequently, the coverslips were rinsed with MilliQ water and dried under a stream of N_2 gas. The treated coverslips were stored upright in a Teflon rack within a covered beaker to protect them from dust. They were utilized within a week to maintain their cleanliness. During the iSCAT measurements, videos were recorded with a field of view measuring 10×10 μm. The recording duration ranged from 6000 to 15000 frames, with a frame rate of 300 frames per second. This ensured a minimum count of 1000 particles per video. Maintaining a dilute solution of proteins is crucial to ensure well-separated scattering patterns with no overlap. Hence protein concentrations (in the range of 10 nM to 50 nM, dilution with F-buffer from the stock concentration of the protein) were chosen to achieve adequate separation of landing events while still collecting sufficient statistics within a maximum of 15000 frames to limit data volume. The iSCAT video analysis was executed using the DiscoverMP software, a commercial software package provided by Refeyn.

5.4. Transmission Electron Microscopy

Electron microscopy analysis was conducted on samples deposited on Cu400 carbon support grids (Quantifoil, Cat. #N1-C73nCu40-01) with a JEM-1400Plus transmission electron microscope (JEOL) operating at an acceleration voltage of 120 kV. Imaging was carried out using a TemCam-F416 CMOS camera (TVIPS) with a resolution of 4k×4k pixels. The sample preparation protocol was as follows: Initially, the Cu400 grids underwent a glow discharge treatment in an oxygen plasma using the GloQube-D instrument provided by Quorum Technologies Ltd. This procedure was essential to enhance protein adsorption. Subsequently, a 4 μL droplet of the protein sample was delicately deposited onto the grid and allowed to adsorb for 1 to 2 minutes. To eliminate any surplus protein and salt, the samples underwent three washed with MilliQ water with careful blot-drying after each wash. Finally, the samples were stained for 25 seconds with a 2% aqueous solution of uranyl acetate from Electron Microscopy Sciences. Following staining, the samples were meticulously blot-dried once again to eliminate any remaining liquid.

5.5. Rheology

Rheological measurements were conducted using a stress-controlled KinexusMalvern Pro rheometer equipped with a stainless steel cone-plate geometry. The cone had a radius of 20 mm and an angle of 1°. The temperature was maintained at 37 °C by Peltier plates. F-actin-vimentin networks were polymerized between the plates of the rheometer by loading 40 μl of the sample directly after mixing the proteins into the polymerization buffer (F-buffer). To prevent solvent evaporation during the experiment, a thin layer of mineral oil Type A was carefully applied around the sample edge. Progress of network formation was monitored by applying a small oscillatory shear with a fixed strain amplitude of 0.5% and oscillation frequency of 0.5 Hz for a duration of 2 hours. Following the 2-hour polymerization period, a frequency sweep was conducted in the range of 0.01 to 10 Hz at a small strain amplitude of 1%, sampling 10 data points per decade. Data taken at frequencies exceeding 3 Hz had to be discarded because they were dominated by inertial effects from the rheometer. For each curve, we took an average over $N = 2$ independently prepared samples.

5.6. Data analysis

For TIRF microscopy data to retrieve reaction rates of ACTIF on vimentin filaments (or on F-actin), image analysis was performed by Fiji. First, the images were pre-processed with background subtraction and image registration. Then, we manually selected from 30 to 50 regions of interest (ROIs) following shapes of vimentin filaments (or F-actin) in each field of view to extract the intensity values over time for ACTIF channel. In this experiment, we performed 3 independent repeats with one field of view for each repeat. In each repeat, we calculated the mean and standard deviation of ACTIF intensity over 30 - 50 ROIs and fitted the intensity curves using GraphPad Prism. To retrieve the number of actin filaments attached to vimentin through ACTIF, we quantified the pixel fluorescence intensities in the F-actin channel along vimentin filaments ($\sim$150 filaments, $N = 2$ independent experiments) and plot the distribution of intensities. This distribution displayed multiple peaks, with the first peak at an intensity of $\sim$600 a.u., as expected for single F-actin in these imaging conditions. The distribution was fitted by a multiple Gaussian, with the peak positions constrained to be multiples of the position of the first peak, and common width for all the

peaks except the last one.

For FRAP experiment data to obtain the unbinding rate of ACTIF on vimentin filaments (or on F-actin), we also performed background subtraction and image registration in advance. Then, we also selected 30 ROIs following shapes of vimentin filaments (or F-actin) from a pre-bleached image. The selected filaments needed to be inside the photo-bleaching region. Then we extract the intensity value of ACTIF over time from the selected ROIs. In this experiment, we also performed 3 independent repeats with one field of view for each repeat. In each repeat, we obtained the mean and standard deviation of ACTIF intensity over 30 ROIs and fitted the intensity curves using GraphPad Prism.

For TEM microscopy, image analysis was performed by Fiji. In order to determine the filament widths, intensity profiles perpendicular to the segment of interest were plotted at multiple locations along the filament contours. The width of the bright area between the darker borders was measured, which corresponded to the filament diameter. For copolymerized vimentin and IFBD, $N = 28$ bundles were measured for the 1:10 IFBD:vimentin molar ratios, and $N = 17$ for the 1:1 IFBD:vimentin molar ratios. For copolymerized actin and vimentin with ACTIF, $N = 40$ bundles were measured for each ACTIF:monomer ratio.

For shear rheology, data was extracted from the rSpace Software (NETZSCH Analyzing and Testing). Custom Python scripts were used to plot the different curves of G' and $\tan(\delta)$ vs Frequency. We used an average of $N = 2$ curves of independently prepared experiments per condition.

CREDIT AUTHORSHIP CONTRIBUTION STATEMENT

IIP: Conceptualization, Methodology, Validation, Investigation, Formal analysis, Visualization, Writing – original draft preparation ; QDT: Conceptualization, Methodology, Validation, Investigation, Formal analysis, Visualization, Writing – original draft preparation ; AG: Methodology, Validation, Investigation ; ZM: Methodology, Validation, Investigation ; GW: Conceptualization, Writing – review and editing ; CL: Conceptualization, Methodology, Validation, Investigation, Formal analysis, Writing – original draft preparation; Supervision, Project administration, Funding acquisition ; GHK: Conceptualization, Methodology, Validation, Investigation, Formal analysis, Writing – original draft preparation; Supervision, Project administration, Funding acquisition.

DECLARATION OF COMPETING INTEREST

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

DATA AVAILABILITY

Data will be made available on request.

ACKNOWLEDGEMENTS

The authors would like to thank Jeffrey de Haan for protein purification and Eli van der Sluis for great teaching and discussions about cloning techniques. We would also like to thank Sandrine Etienne-Manneville for the APCn2 construct. We thank the Romet-Lemonne/Jégou lab for actin purification and labeling as well as technical help and stimulating discussions. We gratefully acknowledge funding from the NWO Talent Programme which is financed by the Dutch Research Council (project number VI.C.182.004), the Centre National de la Recherche Scientifique, and the Agence Nationale de la recherche (project number ANR-21-CE11-0004). QDT was supported by the LabEx "Who am I?" (ANR-11-LABX-0071) and the Université Paris Cité IdEx (ANR-18-IDEX-0001), funded by the French Government through its "Investments for the Future" program.

Al-Jassar, C., Bernadó, P., Chidgey, M., and Overduin, M. (2013). Hinged plakin domains provide specialized degrees of articulation in envoplakin, periplakin and desmoplakin. *PLOS One*, 8(7):e69767.

Ali, M. Y., Krementsova, E. B., Kennedy, G. G., Mahaffy, R., Pollard, T. D., Trybus, K. M., and Warshaw, D. M. (2007). Myosin va maneuvers through actin intersections and diffuses along microtubules. *Proc. Natl. Acad. Sci.*, 104(11):4332–4336.

Alkemade, C., Wierenga, H., Volkov, V. A., Preciado López, M., Akhmanova, A., Ten Wolde, P. R., Dogterom, M., and Koenderink, G. H. (2022). Cross-linkers at growing microtubule ends generate forces that drive actin transport. *Proc. Natl. Acad. Sci.*, 119(11):e2112799119.

Aufderhorst-Roberts, A. and Koenderink, G. H. (2019). Stiffening and inelastic fluidization in vimentin intermediate filament networks. *Soft Matter*, 15(36):7127–7136.

Berezney, J., Goode, B. L., Fraden, S., and Dogic, Z. (2022). Extensile to contractile transition in active microtubule–actin composites generates layered asters with programmable lifetimes. *Proc. Natl. Acad. Sci.*, 119(5):e2115895119.

Berr, A. L., Wiese, K., Dos Santos, G., Koch, C. M., Anekalla, K. R., Kidd, M., Davis, J. M., Cheng, Y., Hu, Y.-S., and Ridge, K. M. (2023). Vimentin is required for tumor progression and metastasis in a mouse model of non–small cell lung cancer. *Oncogene*, 42:2074—2087.

Bhattacharya, R., Gonzalez, A. M., DeBiase, P. J., Trejo, H. E., Goldman, R. D., Flitney, F. W., and Jones, J. C. (2009). Recruitment of vimentin to the cell surface by $\beta 3$ integrin and plectin mediates adhesion strength. *J. Cell Sci.*, 122(9):1390–1400.

Block, J., Witt, H., Candelli, A., Peterman, E. J., Wuite, G. J., Janshoff, A., and Köster, S. (2017). Nonlinear loading-rate-dependent force response of individual vimentin intermediate filaments to applied strain. *Phys. Rev. Lett.*, 118(4):048101.

Bonakdar, N., Schilling, A., Spörrer, M., Lennert, P., Mainka, A., Winter, L., Walko, G., Wiche, G., Fabry, B., and Goldmann, W. H. (2015). Determining the mechanical properties of plectin in mouse myoblasts and keratinocytes. *Exp. Cell Res.*, 331(2):331–337.

Bouameur, J.-E., Favre, B., Fontao, L., Lingasamy, P., Begré, N., and Borradori, L. (2014). Interaction of plectin with keratins 5 and 14: dependence on several plectin domains and keratin quaternary structure. *J. Invest. Dermatol.*, 134(11):2776–2783.

Bouameur, J.-E., Schneider, Y., Begré, N., Hobbs, R. P., Lingasamy, P., Fontao, L., Green, K. J., Favre, B., and Borradori, L. (2013). Phosphorylation of serine 4642 in the c-terminus of plectin by mnk2 and pka modulates its interaction with intermediate filaments. *J. Cell Sci.*, 126(18):4195–4207.

Breitsprecher, D., Jaiswal, R., Bombardier, J. P., Gould, C. J., Gelles, J., and Goode, B. L. (2012). Rocket launcher mechanism of collaborative actin assembly defined by single-molecule imaging. *Science*, 336(6085):1164–1168.

Bremer, A., Millonig, R. C., Sütterlin, R., Engel, A., Pollard, T. D., and Aebi, U. (1991). The structural basis for the intrinsic disorder of the actin filament: the" lateral slipping" model. *J. Cell Biol.*, 115(3):689–703.

Bulinski, J. C., Odde, D. J., Howell, B. J., Salmon, T. D., and Waterman-Storer, C. M. (2001). Rapid dynamics of the microtubule binding of ensconsin in vivo. *J. Cell Sci.*, 114(21):3885–3897.

Chernoivanenko, I. S., Matveeva, E. A., Gelfand, V. I., Goldman, R. D., and Minin, A. A. (2015). Mitochondrial membrane potential is regulated by vimentin intermediate filaments. *FASEB J.*, 29(3):820–827.

Choi, H.-J., Park-Snyder, S., Pascoe, L. T., Green, K. J., and Weis, W. I. (2002). Structures of two intermediate filament-binding fragments of desmoplakin reveal a unique repeat motif structure. *Nat. Struct. Biol.*, 9(8):612–620.

Colburn, Z. T. and Jones, J. C. (2018). Complexes of $\alpha6\beta4$ integrin and vimentin act as signaling hubs to regulate epithelial cell migration. *J. Cell Sci.*, 131(14):jcs214593.

Costigliola, N., Ding, L., Burckhardt, C. J., Han, S. J., Gutierrez, E., Mota, A., Groisman, A., Mitchison, T. J., and Danuser, G. (2017). Vimentin fibers orient traction stress. *Proc. Natl. Acad. Sci.*, 114(20):5195–5200.

Daday, C., Kolšek, K., and Gräter, F. (2017). The mechano-sensing role of the unique sh3 insertion in plakin domains revealed by molecular dynamics simulations. *Sci. Rep.*, 7(1):11669.

De Pascalis, C., Pérez-González, C., Seetharaman, S., Boëda, B., Vianay, B., Burute, M., Leduc, C., Borghi, N., Trepat, X., and Etienne-Manneville, S. (2018). Intermediate filaments control collective migration by restricting traction forces and sustaining cell–cell contacts. *J. Cell Biol.*, 217(9):3031–3044.

Dogterom, M. and Koenderink, G. H. (2019). Actin-microtubule crosstalk in cell biology. *Nat. Rev. Mol. Cell Biol.*, 20(1):38–54.

Duarte, S., Viedma-Poyatos, Á., Navarro-Carrasco, E., Martínez, A. E., Pajares, M. A., and Pérez-Sala, D. (2019). Vimentin filaments interact with the actin cortex in mitosis allowing normal cell division. *Nat. Commun.*, 10(1):1–19.

Eibauer, M., Weber, M. S., Kronenberg-Tenga, R., Beales, C. T., Boujemaa-Paterski, R., Turgay, Y., Sivagurunathan, S., Kraxner, J., Koester, S., Goldman, R. D., et al. (2023). Vimentin filaments integrate low complexity domains in a highly complex helical structure. *bioRxiv*, pages 2023–05.

Esue, O., Carson, A. A., Tseng, Y., and Wirtz, D. (2006). A direct interaction between actin and vimentin filaments mediated by the tail domain of vimentin. *J. Biol. Chem.*, 281(41):30393–30399.

Faix, J., Steinmetz, M., Boves, H., Kammerer, R. A., Lottspeich, F., Mintert, U., Murphy, J., Stock, A., Aebi, U., and Gerisch, G. (1996). Cortexillins, major determinants of cell shape and size, are actin-bundling proteins with a parallel coiled-coil tail. *Cell*, 86(4):631–642.

Favre, B., Schneider, Y., Lingasamy, P., Bouameur, J.-E., Begré, N., Gontier, Y., Steiner-Champliaud, M.-F., Frias, M. A., Borradori, L., and Fontao, L. (2011). Plectin interacts with the rod domain of type iii intermediate filament proteins desmin and vimentin. *Eur. J. Cell Biol.*, 90(5):390–400.

Fletcher, D. A. and Mullins, R. D. (2010). Cell mechanics and the cytoskeleton. *Nature*, 463(7280):485.

Fogl, C., Mohammed, F., Al-Jassar, C., Jeeves, M., Knowles, T. J., Rodriguez-Zamora, P., White, S. A., Odintsova, E., Overduin, M., and Chidgey, M. (2016). Mechanism of intermediate filament recognition by plakin repeat domains revealed by envoplakin targeting of vimentin. *Nat. Commun.*, 7(1):10827.

Foisner, R., Bohn, W., Mannweiler, K., and Wiche, G. (1995). Distribution and ultrastructure of plectin arrays in subclones of rat glioma c6 cells differing in intermediate filament protein (vimentin) expression. *J. Struct. Biol.*, 115(3):304–17.

Foisner, R., Leichtfried, F., Herrmann, H., Small, J., Lawson, D., and Wiche, G. (1988). Cytoskeleton-associated plectin: in situ localization, in vitro reconstitution, and binding to immobilized intermediate filament proteins. *J. Cell Biol.*, 106(3):723–733.

Foisner, R., Malecz, N., Dressel, N., Stadler, C., and Wiche, G. (1996). M-phase-specific phosphorylation and structural rearrangement of the cytoplasmic cross-linking protein plectin involve p34cdc2 kinase. *Mol. Biol. Cell*, 7(2):273–288.

Foisner, R., Traub, P., and Wiche, G. (1991). Protein kinase a-and protein kinase c-regulated interaction of plectin with lamin b and vimentin. *Proc. Natl. Acad. Sci.*, 88(9):3812–3816.

Fontao, L., Geerts, D., Kuikman, I., Koster, J., Kramer, D., and Sonnenberg, A. (2001). The interaction of plectin with actin: evidence for cross-linking of actin filaments by dimerization of the actin-binding domain of plectin. *J. Cell Sci.*, 114(11):2065–2076.

Forsting, J., Kraxner, J., Witt, H., Janshoff, A., and Köster, S. (2019). Vimentin intermediate filaments undergo irreversible conformational changes during cyclic loading. *Nano Lett.*, 19(10):7349–7356.

Gan, Z., Ding, L., Burckhardt, C., Lowery, J., Zaritsky, A., Sitterley, K., Mota, A., Costigliola, N., Starker, C., Voytas, D., Tytell, J., Goldman, R., and Danuser, G. (2016). Vimentin intermediate filaments template microtubule networks to enhance persistence in cell polarity and directed migration. *Cell Systems*, 3(3):252–263.e8.

Garcia-Alvarez, B., Bobkov, A., Sonnenberg, A., and de Pereda, J. M. (2003). Structural and functional analysis of the actin binding domain of plectin suggests alternative mechanisms for binding to f-actin and integrin $\beta 4$. *Structure*, 11(6):615–625.

Geerts, D., Fontao, L., Nievers, M. G., Schaapveld, R. Q., Purkis, P. E., Wheeler, G. N., Lane, E. B., Leigh, I. M., and Sonnenberg, A. (1999). Binding of integrin $\alpha 6\beta 4$ to plectin prevents plectin association with f-actin but does not interfere with intermediate filament binding. *J. Cell Biol.*, 147(2):417–434.

Gimona, M. and Winder, S. J. (1998). Single calponin homology domains are not actin-binding domains. *Curr. Biol.*, 8(19):R673–R675.

Golde, T., Huster, C., Glaser, M., Händler, T., Herrmann, H., Käs, J. A., and Schnauß, J. (2018). Glassy dynamics in composite biopolymer networks. *Soft Matter*, 14(39):7970–7978.

González-Jiménez, P., Duarte, S., Martínez, A. E., Navarro-Carrasco, E., Lalioti, V., Pajares, M. A., and Pérez-Sala, D. (2023). Vimentin single cysteine residue acts as a tunable sensor for network organization and as a key for actin remodeling in response to oxidants and electrophiles. *Redox Biol.*, page 102756.

Herrmann, H., Kreplak, L., and Aebi, U. (2004). Isolation, characterization, and in vitro assembly of intermediate filaments. In *Methods Cell Biol.*, volume 78, pages 3–24. Elsevier.

Hoelzle, M. K. and Svitkina, T. (2012). The cytoskeletal mechanisms of cell-cell junction formation in endothelial cells. *Mol. Biol. Cell.*, 23(2):310–323.

Houk Jr, T. W. and Ue, K. (1974). The measurement of actin concentration in solution: a comparison of methods. *Anal. Biochem.*, 62(1):66–74.

Hu, J., Li, Y., Hao, Y., Zheng, T., Gupta, S. K., Parada, G. A., Wu, H., Lin, S., Wang, S., Zhao, X., et al. (2019). High stretchability, strength, and toughness of living cells enabled by hyperelastic vimentin intermediate filaments. *Proc. Natl. Acad. Sci.*, 116(35):17175–17180.

Huber, F., Boire, A., López, M. P., and Koenderink, G. H. (2015). Cytoskeletal crosstalk: when three different personalities team up. *Curr. Opin. Cell Biol.*, 32:39–47. Cell architecture.

Janmey, P. A. (1998). The cytoskeleton and cell signaling: Component localization and mechanical coupling. *Physiol. Rev.*, 78(3):763–781.

Jensen, M. H., Morris, E. J., Goldman, R. D., and Weitz, D. A. (2014). Emergent properties of composite semiflexible biopolymer networks. *Bioarchitecture*, 4(4-5):138–143.

Jiu, Y., Lehtimäki, J., Tojkander, S., Cheng, F., Jäälinoja, H., Liu, X., Varjosalo, M., Eriksson, J., and Lappalainen, P. (2015). Bidirectional interplay between vimentin intermediate filaments and contractile actin stress fibers. *Cell Rep.*, 11(10):1511–1518.

Jiu, Y., Peränen, J., Schaible, N., Cheng, F., Eriksson, J. E., Krishnan, R., and Lappalainen, P. (2017). Vimentin intermediate filaments control actin stress fiber assembly through GEF-H1 and RhoA. *J. Cell Sci.*, 130(5):892–902.

Kang, H., Weiss, T. M., Bang, I., Weis, W. I., and Choi, H.-J. (2016). Structure of the intermediate filament-binding region of desmoplakin. *PLOS One*, 11(1):e0147641.

Kraxner, J., Lorenz, C., Menzel, J., Parfentev, I., Silbern, I., Denz, M., Urlaub, H., Schwappach, B., and Köster, S. (2021). Post-translational modifications soften vimentin intermediate filaments. *Nanoscale*, 13(1):380–387.

Leduc, C., Padberg-Gehle, K., Varga, V., Helbing, D., Diez, S., and Howard, J. (2012). Molecular crowding creates traffic jams of kinesin motors on microtubules. *Proc. Natl. Acad. Sci.*, 109(16):6100–6105.

López, M. P., Huber, F., Grigoriev, I., Steinmetz, M. O., Akhmanova, A., Koenderink, G. H., and Dogterom, M. (2014). Actin–microtubule coordination at growing microtubule ends. *Nat. Commun.*, 5(1):1–9.

Lorenz, C., Forsting, J., Style, R. W., Klumpp, S., and Köster, S. (2023). Keratin filament mechanics and energy dissipation are determined by metal-like plasticity. *Matter*, 6(6):2019–2033.

Marks, P. C., Hewitt, B. R., Baird, M. A., Wiche, G., and Petrie, R. J. (2022). Plectin linkages are mechanosensitive and required for the nuclear piston mechanism of three-dimensional cell migration. *Mol. Biol. Cell*, 33(12):ar104.

Martin, I., Moch, M., Neckernuss, T., Paschke, S., Herrmann, H., and Marti, O. (2016). Both monovalent cations and plectin are potent modulators of mechanical properties of keratin k8/k18 networks. *Soft Matter*, 12(33):6964–6974.

Moch, M., Windoffer, R., Schwarz, N., Pohl, R., Omenzetter, A., Schnakenberg, U., Herb, F., Chaisaowong, K., Merhof, D., Ramms, L., et al. (2016). Effects of plectin depletion on keratin network dynamics and organization. *PLOS One*, 11(3):e0149106.

Mohammed, F., Trieber, C., Overduin, M., and Chidgey, M. (2020). Molecular mechanism of intermediate filament recognition by plakin proteins. *Biochim. Biophys. Acta Mol. Cell Res.*, 1867(11):118801.

Mulla, Y., Aufderhorst-Roberts, A., and Koenderink, G. H. (2018). Shaping up synthetic cells. *Phys. Biol.*, 15(4):041001.

Na, S., Chowdhury, F., Tay, B., Ouyang, M., Gregor, M., Wang, Y., Wiche, G., and Wang, N. (2009). Plectin contributes to mechanical properties of living cells. *Am. J. Physiol. Cell Physiol.*, 296(4):C868–C877.

Nakaseko, Y. and Yanagida, M. (2001). Cytoskeleton in the cell cycle. *Nature*, 412(6844):291–292.

Nikolic, B., Mac Nulty, E., Mir, B., and Wiche, G. (1996). Basic amino acid residue cluster within nuclear targeting sequence motif is essential for cytoplasmic plectin-vimentin network junctions. *J. Cell Biol.*, 134(6):1455.

Nunes Vicente, F., Lelek, M., Tinevez, J.-Y., Tran, Q. D., Pehau-Arnaudet, G., Zimmer, C., Etienne-Manneville, S., Giannone, G., and Leduc, C. (2022). Molecular organization and mechanics of single vimentin filaments revealed by super-resolution imaging. *Sci. Adv.*, 8(8):eabm2696.

Ortega, E., Manso, J. A., Buey, R. M., Carballido, A. M., Carabias, A., Sonnenberg, A., and de Pereda, J. M. (2016). The structure of the plakin domain of plectin reveals an extended rod-like shape. *J. Biol. Chem.*, 291(36):18643–18662.

Osmanagic-Myers, S., Rus, S., Wolfram, M., Brunner, D., Goldmann, W. H., Bonakdar, N., Fischer, I., Reipert, S., Zuzuarregui, A., Walko, G., and Wiche, G. (2015). Plectin reinforces vascular integrity by mediating crosstalk between the vimentin and the actin networks. *J. Cell Sci.*, 128(22):4138–4150.

Patteson, A. E., Vahabikashi, A., Goldman, R. D., and Janmey, P. A. (2020). Mechanical and non-mechanical functions of filamentous and non-filamentous vimentin. *BioEssays*, 42(11):2000078.

Pegoraro, A. F., Janmey, P., and Weitz, D. A. (2017). Mechanical properties of the cytoskeleton and cells. *Cold Spring Harb. Perspect. Biol.*, 9(11).

Pelletier, V., Gal, N., Fournier, P., and Kilfoil, M. L. (2009). Microrheology of microtubule solutions and actin-microtubule composite networks. *Phys. Rev. Lett.*, 102(18):188303.

Prechova, M., Adamova, Z., Schweizer, A.-L., Maninova, M., Bauer, A., Kah, D., Meier-Menches, S. M., Wiche, G., Fabry, B., and Gregor, M. (2022). Plectin-mediated cytoskeletal crosstalk controls cell tension and cohesion in epithelial sheets. *J. Cell Biol.*, 221(3):e202105146.

Ricketts, S. N., Ross, J. L., and Robertson-Anderson, R. M. (2018). Co-entangled actin-microtubule composites exhibit tunable stiffness and power-law stress relaxation. *Biophys. J.*, 115(6):1055–1067.

Ridge, K. M., Eriksson, J. E., Pekny, M., and Goldman, R. D. (2022). Roles of vimentin in health and disease. *Genes Dev.*, 36(7-8):391–407.

Romet-Lemonne, G., Guichard, B., and Jégou, A. (2018). *Using Microfluidics Single Filament Assay to Study Formin Control of Actin Assembly*, pages 75–92. Springer New York, New York, NY.

Sakamoto, Y., Boëda, B., and Etienne-Manneville, S. (2013). APC binds intermediate filaments and is required for their reorganization during cell migration. *J. Cell Biol.*, 200(3):249–258.

Sanchez, T., Chen, D. T., DeCamp, S. J., Heymann, M., and Dogic, Z. (2012). Spontaneous motion in hierarchically assembled active matter. *Nature*, 491(7424):431–434.

Schaedel, L., Lorenz, C., Schepers, A. V., Klumpp, S., and Köster, S. (2021). Vimentin intermediate filaments stabilize dynamic microtubules by direct interactions. *Nat. Commun.*, 12(1):3799.

Schoumacher, M., Goldman, R. D., Louvard, D., and Vignjevic, D. M. (2010). Actin, microtubules, and vimentin intermediate filaments cooperate for elongation of invadopodia. *J. Cell Biol.*, 189(3):541–556.

Schroeder, H. W., Mitchell, C., Shuman, H., Holzbaur, E. L., and Goldman, Y. E. (2010). Motor number controls cargo switching at actin-microtubule intersections in vitro. *Curr. Biol.*, 20(8):687–696.

Seetharaman, S. and Etienne-Manneville, S. (2020). Cytoskeletal crosstalk in cell migration. *Trends Cell Biol.*, 30(9):720–735.

Serres, M. P., Samwer, M., Quang, B. A. T., Lavoie, G., Perera, U., Görlich, D., Charras, G., Petronczki, M., Roux, P. P., and Paluch, E. K. (2020). F-actin interactome reveals vimentin as a key regulator of actin organization and cell mechanics in mitosis. *Dev. Cell*, 52(2):210–222.

Sevcik, J., Urbanikova, L., Kost'an, J., Janda, L., and Wiche, G. (2004). Actin-binding domain of mouse plectin. crystal structure and binding to vimentin. *Eur. J. Biochem.*, 271(10):1873–1884.

Shen, Y., Wu, H., Lu, P. J., Wang, D., Shayegan, M., Li, H., Shi, W., Wang, Z., Cai, L. H., Xia, J., Zhang, M., Ding, R., Herrmann, H., Goldman, R., Mackintosh, F. C., Moncho-Jordá, A., and Weitz, D. A. (2021). Effects of vimentin intermediate filaments on the structure and dynamics of in vitro multicomponent interpenetrating cytoskeletal networks. *Phys. Rev. Lett.*, 127(10):108101.

Sheung, J. Y., Achiriloaie, D. H., Currie, C., Peddireddy, K., Xie, A., Simon-Parker, J., Lee, G., Rust, M. J., Das, M., Ross, J. L., et al. (2021). Motor-driven restructuring of cytoskeleton composites leads to tunable time-varying elasticity. *ACS Macro Lett.*, 10(9):1151–1158.

Sjöqvist, M., Antfolk, D., Suarez-Rodriguez, F., and Sahlgren, C. (2021). From structural resilience to cell specification — intermediate filaments as regulators of cell fate. *FASEB J.*, 35(1):e21182.

Sonnenberg, A., Rojas, A. M., and de Pereda, J. M. (2007). The structure of a tandem pair of spectrin repeats of plectin reveals a modular organization of the plakin domain. *J. Mol. Biol.*, 368(5):1379–1391.

Spudich, J. A. and Watt, S. (1971). The regulation of rabbit skeletal muscle contraction: I. biochemical studies of the interaction of the tropomyosin-troponin complex with actin and the proteolytic fragments of myosin. *J. Biol. Chem.*, 246(15):4866–4871.

Spurny, R., Abdoulrahman, K., Janda, L., Ruönzler, D., Koöhler, G., Castanon, M. J., and Wiche, G. (2007). Oxidation and nitrosylation of cysteines proximal to the intermediate filament (if)-binding site of plectin: effects on structure and vimentin binding and involvement in if collapse. *J. Biol. Chem.*, 282(11):8175–8187.

Steinböck, F. A., Nikolic, B., Coulombe, P. A., Fuchs, E., Traub, P., and Wiche, G. (2000). Dose-dependent linkage, assembly inhibition and disassembly of vimentin and cytokeratin 5/14 filaments through plectin's intermediate filament-binding domain. *J. Cell Sci.*, 113(3):483–491.

Steiner-Champliaud, M.-F., Schneider, Y., Favre, B., Paulhe, F., Praetzel-Wunder, S., Faulkner, G., Konieczny, P., Raith, M., Wiche, G., Adebola, A., et al. (2010). Bpag1 isoform-b: Complex distribution pattern in striated and heart muscle and association with plectin and α-actinin. *Exp. Cell Res.*, 316(3):297–313.

Styers, M. L., Salazar, G., Love, R., Peden, A. A., Kowalczyk, A. P., and Faundez, V. (2004). The endo-lysosomal sorting machinery interacts with the intermediate filament cytoskeleton. *Mol. Biol. Cell*, 15(12):5369.

Svitkina, T. M., Verkhovsky, A. B., and Borisy, G. G. (1996). Plectin sidearms mediate interaction of intermediate filaments with microtubules and other components of the cytoskeleton. *J. Cell Biol.*, 135(4):991–1007.

Tarbet, H. J., Dolat, L., Smith, T. J., Condon, B. M., O'Brien III, E. T., Valdivia, R. H., and Boyce, M. (2018). Site-specific glycosylation regulates the form and function of the intermediate filament cytoskeleton. *eLife*, 7:e31807.

Tomar, A. and Schlaepfer, D. D. (2009). Focal adhesion kinase: switching between gaps and gefs in the regulation of cell motility. *Curr. Opin. Cell Biol.*, 21(5):676.

Tran, Q. D., Sorichetti, V., Pehau-Arnaudet, G., Lenz, M., and Leduc, C. (2023). Fragmentation and entanglement limit vimentin intermediate filament assembly. *Phys. Rev. X*, 13:011014.

Tusamda Wakhloo, N., Anders, S., Badique, F., Eichhorn, M., Brigaud, I., Petithory, T., Vassaux, M., Milan, J.-L., Freund, J.-N., Rühe, J., Davidson, P. M., Pieuchot, L., and Anselme, K. (2020). Actomyosin, vimentin and linc complex pull on osteosarcoma nuclei to deform on micropillar topography. *Biomaterials*, 234:119746.

Urbániková, L., Janda, L., Popov, A., Wiche, G., and Ševčík, J. (2002). Purification, crystallization and preliminary x-ray analysis of the plectin actin-binding domain. *Acta Crystallogr. D*, 58(8):1368–1370.

van Bodegraven, E. J. and Etienne-Manneville, S. (2020). Intermediate filaments against actomyosin: the david and goliath of cell migration. *Curr. Opin. Cell Biol.*, 66:79–88.

Walker, J. M. (2005). *The proteomics protocols handbook*. Springer.

Walko, G., Castañón, M. J., and Wiche, G. (2015). Molecular architecture and function of the hemidesmosome. *Cell Tissue Res.*, 360:363–378.

Ward, S. M. V., Weins, A., Pollak, M. R., and Weitz, D. A. (2008). Dynamic viscoelasticity of actin cross-linked with wild-type and disease-causing mutant α-actinin-4. *Biophys. J.*, 95(10):4915–4923.

Weiss, M., Frohnmayer, J. P., Benk, L. T., Haller, B., Janiesch, J.-W., Heitkamp, T., Börsch, M., Lira, R. B., Dimova, R., Lipowsky, R., et al. (2018). Sequential bottom-up assembly of mechanically stabilized synthetic cells by microfluidics. *Nat. Mater.*, 17(1):89–96.

Wen, Q. and Janmey, P. A. (2011). Polymer physics of the cytoskeleton. *Curr. Opin. Solid State Mater. Sci.*, 15(5):177–182.

Wiche, G. and Baker, M. A. (1982). Cytoplasmic network arrays demonstrated by immunolocalization using antibodies to a high molecular weight protein present in cytoskeletal preparations from cultured cells. *Exp. Cell Res.*, 138(1):15–29.

Wiche, G. and Winter, L. (2011). Plectin isoforms as organizers of intermediate filament cytoarchitecture. *Bioarchitecture*, 1(1):14–20.

Wilhelmsen, K., Litjens, S. H., Kuikman, I., Tshimbalanga, N., Janssen, H., van den Bout, I., Raymond, K., and Sonnenberg, A. (2005). Nesprin-3, a novel outer nuclear membrane protein, associates with the cytoskeletal linker protein plectin. *J. Cell Biol.*, 171(5):799–810.

Winheim, S., Hieb, A. R., Silbermann, M., Surmann, E.-M., Wedig, T., Herrmann, H., Langowski, J., and Mücke, N. (2011). Deconstructing the late phase of vimentin assembly by total internal reflection fluorescence microscopy (tirfm). *PLOS One*, 6(4).

Winter, L., Abrahamsberg, C., and Wiche, G. (2008). Plectin isoform 1b mediates mitochondrion–intermediate filament network linkage and controls organelle shape. *J. Cell Biol.*, 181(6):903–911.

Wioland, H., Guichard, B., Senju, Y., Myram, S., Lappalainen, P., Jégou, A., and Romet-Lemonne, G. (2017). Adf/cofilin accelerates actin dynamics by severing filaments and promoting their depolymerization at both ends. *Curr. Biol.*, 27(13):1956–1967.

Yoneyama, M. S., Hatakeyama, S., Habuchi, T., Inoue, T., Nakamura, T., Funyu, T., Wiche, G., Ohyama, C., and Tsuboi, S. (2014). Vimentin intermediate filament and plectin provide a scaffold for invadopodia, facilitating cancer cell invasion and extravasation for metastasis. *Eur. J. Cell Biol.*, 93(4):157–169.

Young, G., Hundt, N., Cole, D., Fineberg, A., Andrecka, J., Tyler, A., Olerinyova, A., Ansari, A., Marklund, E. G., Collier, M. P., Chandler, S. A., Tkachenko, O., Allen, J., Crispin, M., Billington, N., Takagi, Y., Sellers, J. R., Eichmann, C., Selenko, P., Frey, L., Riek, R., Galpin, M. R., Struwe, W. B., Benesch, J. L. P., and Kukura, P. (2018). Quantitative mass imaging of single biological macromolecules. *Science*, 360(6387):423–427.

Zrelski, M. M., Hösele, S., Kustermann, M., Fichtinger, P., Kah, D., Athanasiou, I., Esser, P. R., Wagner, A., Herzog, R., Kratochwill, K., et al. (2023). Plectin deficiency in fibroblasts deranges intermediate filament and organelle morphology, migration, and adhesion. *J. Invest. Dermatol.*

6. APPENDIX A: SUPPLEMENTARY MATERIAL

6.1. Supplementary tables and figures

Supplemental Table 1: Primers used for Gibson Assembly of ACTIF construct

Construct	Fragment	Forward primer	Reverse primer
ACTIF	Vector	5'- TGAGATCCGGCTGCTAACAAAGCC -3'	5'- CCCTCTGATTCTGGTGGCTTCGAT -3'
	Insert	5'- AAAGTTCCTGAGGGTGGAGAAGGGAAAGCTCGT TTAGAGAGTAGCAAA -3' 5'- TTGTTGTTCGAGTTTTGATTTTTCATCGATGGCTTCAGCTA ATTTCAA -3'	

Supplemental Table 2: Primers used for Gibson Assembly of ACTIF-APCn2 construct

Construct	Fragment	Forward primer	Reverse primer
ACTIF-APCn2	Vector	5'- GAAAAATCAAAACTCGAACAACAA -3'	5'- CCCTTCTCCACCCTCAGGAACTTT -3'
	Insert	5'- ATCGAAGCCACCAGAATCAGAGGGCTAGCTATGTCTAGCTCCCAAGAC-3' 5'- GGCTTTGTTAGCAGCCGGATCTCAGGCATCCTTGTACTTCGCAGGCCT -3'	

Supplemental Table 3: Primers used for Gibson Assembly of IFBD(plectin) construct

Construct	Fragment	Forward primer	Reverse primer
IFBD	Vector	5'- GATCCGGCTGCTAACAAAGCCCGA-3'	5'- GCGACCCATTTGCTGTCCACCAGT-3'
	Insert	5'- ACTGGTGGACAGCAAATGGGTCGCATCGAAGCCACCAGAATCAGAGGG-3' 5'-TCGGGCTTTGTTAGCAGCCGGATCTCAGGCGGTCGAGCCAGAGCCACT -3'	

Supplemental Figure 1: Frames from a time-lapse video recorded for interferometric scattering microscopy of a sample of 40nM ACTIF. Different time frames are shown in seconds. The area of detection shown is of 18 μM^2

Supplemental Figure 2: Distribution of pixel fluorescence intensities along vimentin filaments. The distribution appears as a single peak with the average intensity expected for single filaments under these imaging conditions, indicating that vimentin exists mainly as single filaments when attached to the substrate. Sample size, ~100 filaments. PDF, Probability density function.

Supplemental Figure 3: iSCAT and SDS-PAGE of ACTIF-APCn2 and IFBD. (A) iSCAT histogram count of IFBD (plectin), portraying a mostly monomeric population with a 13% dimeric population count. (B) SDS-PAGE of the pooled fractions of IFBD(plectin) after purification, with an arrow marking the band of interest. (C) iSCAT histogram count of APCn2, portraying a 95% dimeric population. (D) SDS-PAGE of the pooled fractions of APCn2, with an arrow marking the band of interest.

Supplemental Figure 4: Histogram of diameters of randomly sampled vimentin filaments (N=50).

Supplemental Figure 5: Transmission electron microscopy of (A) actin filaments at 2 μM concentration. (B) Actin filaments at 2 μM concentration and 20 muM IFBD (plectin). No apparent interaction between actin and the IFBD, which tends to oligomerize (see globular structures not present on (A)). Scale bars indicate 100 nm.

Supplemental Figure 6: Details from transmission electron microscopy of copolymerized actin, vimentin and ACTIF. (A) Mixed actin and vimentin within a bundle (blue arrows for actin, magenta for vimentin). (B) Detail on different bundle packing. (C) Bundle of only distinguishable actin filaments, based on their diameter. Scale bars indicate 100 nm.

Supplemental Figure 7: A) Frequency sweep for actin and crosslinked actin networks (top), and loss factor for both actin and crosslinked actin (bottom). B) Frequency sweep for vimentin and crosslinked vimentin networks (top), and loss factor for both actin and crosslinked actin (bottom).

Supplemental Figure 8: Binding affinity of ACTIF-APCn2 on vimentin filaments. (A) Schematic of the binding assay and images of ACTIF-APCn2 (green) at 5 nM co-localized with vimentin filaments (magenta) attached to the substrate, at $t = 10$ min after flushing ACTIF-APCn2 into the chamber. (B) Fluorescence intensity of ACTIF-APCn2 at 5 nM (dark green), in comparison with ACTIF at the same concentration of 5 nM (light green), on vimentin filaments over time (averaged over 30 filaments). Black lines represent linear regression of the curves. The shaded area represents standard deviations. Scale bar: 5 μm.

```
PLEC    1    RVQKKTFTKWVNKHLIKAQRHISDLYEDLRDGHNLISLLEVLSGDSLPREKGRMRFHKLQ    60
ACTIF   1    RVQKKTFTKWVNKHLMKVRKHINDLYEDLRDGHNLISLLEVLSGIKLPREKGRMRFHRLQ    60

PLEC    61   NVQIALDYLRHRQVKLVNIRNDDIADGNPKLTLGLIWTIILHFQ------------MTAK    108
ACTIF   61   NVQIALDFLKQRQVKLVNIRNDDITDGNPKLTLGLIWTIILHFQISDIYISGESGDMSAK    120

PLEC    109  EKLLLWSQRMVEGYQGLRCDNFTSSWRDGRLFNAIIHRHKPLLIDMNKVYRQTNLENLDQ    168
ACTIF   121  EKLLLWTQKVTAGYTGIKCTNFSSCWSDGKMFNALIHRYRPDLVDMERVQIQSNRENLEQ    180

PLEC    169  AFSVAERDLGVTRLLDPEDVDVPQPDEKSIITYVSSLYDAMP    210
ACTIF   181  AFEVAER-LGVTRLLDAEDVDVPSPDEKSVITYVSSIYDAFP    221
```

Supplemental Figure 9: Sequence alignment of the calponin homology domains of plectin (top) and ACTIF (bottom), where the CH tandem of ACTIF has been taken from MACF. In red, sequence differences.

A

ACTIF

eGFP   CH   CH   Cortexillin coiled-coil   IFBD

ABD

B

```
MGSSHHHHHHSSGLVPRGSHMASMTGGQQMGRMVSKGEELFTGVVPILVELDGDVNGHKF   60
SVSGEGEGDATYGKLTLKFICTTGKLPVPWPTLVTTLTYGVQCFARYPDHMKQHDFFKSA  120
MPEGYVQERTIFFKDDGNYKTRAEVKFEGDTLVNRIELKGIDFKEDGNILGHKLEYNYNS  180
HKVYITADKQKNGIKVNFKTRHNIEDGSVQLADHYQQNTPIGDGPVLLPDNHYLSTQSAL  240
SKDPNEKRDHMVLLEFVTAAGITLGMDELYKSGLMDERDRVQKKTFTKWVNKHLMKVRKH  300
INDLYEDLRDGHNLISLLEVLSGIKLPREKGRMRFHRLQNVQIALDFLKQRQVKLVNIRN  360
DDITDGNPKLTLGLIWTIILHFQISDIYISGESGDMSAKEKLLLWTQKVTAGYTGIKCTN  420
FSSCWSDGKMFNALIHRYRPDLVDMERVQIQSNRENLEQAFEVAERLGVTRLLDAEDVDV  480
PSPDEKSVITYVSSIYDAFPKVPEGGEGKARLESSKNEMANRLAGLENSLESEKVSREQL  540
IKQKDQLNSLLASLESEGAEREKRLRELEAKLDETLKNLELEKLARMELEARLAKTEKDR  600
AILELKLAEAIDEKSKLEQQIEATRIRGRVPGAQQFFRGLRKQITVEELVRSQVMDEATA  660
LQLQEGLTSIEEVTKNLQKFLEGTSCIAGVFVDATKERLSVYQAMKKGIIRPGTAFELLE  720
AQAATGYVIDPIKGLKLTVEEAVRMGIVGPEFKDKLLSAERAVTGYKDPYSGKLISLFQA  780
MKKGLILKDHGIRLLEAQIATGGIIDPEESHRLPVEVAYKRGLFDEEMNEILTDPSDDTK  840
GFFDPNTEENLTYLQLMERCITDPQTGLCLLPLKEKKRERKTSSKSSVRKRRVVIVDPET  900
GKEMSVYEAYRKGLIDHQTYLELSEQECEWEEITISSSDGVVKSMIIDRRSGRQYDIDDA  960
ITKNLIDRSALDQYRAGTLSITEFADMLSGNAGGFRSRSSSVGSSSSYPISSAGPRTQLA 1020
SWSDPTEETGPVAGILDTETLEKVSITEAMHRNLVDNITGQRLLEAQACTGGIIDPSTGE 1080
RFPVTEAVNKGLVDKIMVDRINLAQKAFCGFEDPRTKTKMSAAQALKKGWLYYEAGQRFL 1140
EVQYLTGGLIEPDTPGRVSLDEALQRGTVDARTAQKLRDVSAYSKYLTCPKTKLKISYKD 1200
ALDRSMVEEGTGLRLLEAAAQSSKGYYSPYSVSGSGSTA*                     1239
```

Supplemental Figure 10: ACTIF's amino-acid sequence and domain organization. (A) ACTIF domain organization with the amino-acid numbers indicated from the sequence in (B). eGFP, enhanced green fluorescent protein; ABD, actin-binding domain; CC, coiled-coil; IFBD, intermediate filament binding domain (B) Amino-acid sequence of ACTIF. The color code corresponds to the domains indicated on the diagram in (A) and the spacers between domains are indicated in light grey letters.

A

IFBD (plectin)

B

MGSSHHHHHHSSGLVPRGSHMASMTGGQQMGRIEATRIRG**RVPGAQQFFRGLRKQITVEE**   **60**

LVRSQVMDEATALQLQEGLTSIEEVTKNLQKFLEGTSCIAGVFVDATKERLSVYQAMKKG   **120**

IIRPGTAFELLEAQAATGYVIDPIKGLKLTVEEAVRMGIVGPEFKDKLLSAERAVTGYKD   **180**

PYSGKLISLFQAMKKGLILKDHGIRLLEAQIATGGIIDPEESHRLPVEVAYKRGLFDEEM   **240**

NEILTDPSDDTKGFFDPNTEENLTYLQLMERCITDPQTGLCLLPLKEKKRERKTSSKSSV   **300**

RKRRVVIVDPETGKEMSVYEAYRKGLIDHQTYLELSEQECEWEEITISSSDGVVKSMIID   **360**

RRSGRQYDIDDAITKNLIDRSALDQYRAGTLSITEFADMLSGNAGGFRSRSSSVGSSSSY   **420**

PISSAGPRTQLASWSDPTEETGPVAGILDTETLEKVSITEAMHRNLVDNITGQRLLEAQA   **480**

CTGGIIDPSTGERFPVTEAVNKGLVDKIMVDRINLAQKAFCGFEDPRTKTKMSAAQALKK   **540**

GWLYYEAGQRFLEVQYLTGGLIEPDTPGRVSLDEALQRGTVDARTAQKLRDVSAYSKYLT   **600**

CPKTKLKISYKDALDRSMVEEGTGLRLLEAAAQSSKGYYSPYSVSGSGSTA*   **651**

Supplemental Figure 11: IFBD's domain organization and amino-acid sequence. (A) Domain organization of IFBD (plectin). (B) Amino-acid sequence of IFBD (plectin) domain. The color code corresponds to the domains indicated on the diagram in (A).

A

ACTIF-APCn2

eGFP   CH   CH   Cortexillin   Armadillo
                    coiled-coil   repeats

              ABD

B

MGSSHHHHHHSSGLVPRGSHMASMTGGQQMGRMVSKGEELFTGVVPILVELDGDVNGHKF   60

SVSGEGEGDATYGKLTLKFICTTGKLPVPWPTLVTTLTYGVQCFARYPDHMKQHDFFKSA   120

MPEGYVQERTIFFKDDGNYKTRAEVKFEGDTLVNRIELKGIDFKEDGNILGHKLEYNYNS   180

HKVYITADKQKNGIKVNFKTRHNIEDGSVQLADHYQQNTPIGDGPVLLPDNHYLSTQSAL   240

SKDPNEKRDHMVLLEFVTAAGITLGMDELYKSGLMDERDRVQKKTFTKWVNKHLMKVRKH   300

INDLYEDLRDGHNLISLLEVLSGIKLPREKGRMRFHRLQNVQIALDFLKQRQVKLVNIRN   360

DDITDGNPKLTLGLIWTIILHFQISDIYISGESGDMSAKEKLLLWTQKVTAGYTGIKCTN   420

FSSCWSDGKMFNALIHRYRPDLVDMERVQIQSNRENLEQAFEVAERLGVTRLLDAEDVDV   480

PSPDEKSVITYVSSIYDAFPKVPEGGEGKARLESSKNEMANRLAGLENSLESEKVSREQL   540

IKQKDQLNSLLASLESEGAEREKRLRELEAKLDETLKNLELEKLARMELEARLAKTEKDR   600

AILELKLAEAIDEKSKLEQQIEATRIRGLAMSSSQDSCISMRQSGCLPLLIQLLHGNDKD   660

SVLLGNSRGSKEARARASAALHNIIHSQPDDKRGRREIRVLHLLEQIRAYCETCWEWQEA   720

HEPGMDQDKNPMPAPVEHQICPAVCVLMKLSFDEEHRHAMNELGGLQAIAELLQVDCEMY   780

GLTNDHYSITLRRYAGMALTNLTFGDVANKATLCSMKGCMRALVAQLKSESEDLQQVIAS   840

VLRNLSWRADVNSKKTLREVGSVKALMECALEVKKESTLKSVLSALWNLSAHCTENKADI   900

CAVDGALAFLVGTLTYRSQTNTLAIIESGGGILRNVSSLIATNEDHRQILRENNCLQTLL   960

QHLKSHSLTIVSNACGTLWNLSARNPKDQEALWDMGAVSMLKNLIHSKHKMIAMGSAAAL   1020

RNLMANRPAKYKDA*

Supplemental Figure 12: ACTIF-APCn2's amino-acid sequence and domain organization. (A) Amino-acid sequence of ACTIF-APCn2. The color code corresponds to the domains indicated on the diagram in (B) and the spacers between domains are indicated in light grey letters. (B) ACTIF-APCn2 domain organization with the amino-acid numbers indicated from the sequence in (A). eGFP, enhanced green fluorescent protein; ABD, actin-binding domain; CC, coiled-coil; armadillo repeats.

6.2. Supplementary movie captions

Supporting movie 1

Control experiment by TIRF microscopy showing no direct interaction between vimentin and actin filaments in our experimental conditions. Pre-assembled vimentin filaments (magenta) were stabilized on the coverslip substrate coated with vimentin antibodies, while F-actin (cyan) was growing from G-actin at 1 μM in F-buffer supplemented with 0.2 % methylcellulose, and moving freely on top of vimentin filaments with no signs of co-localization for 20 min.

Supporting movie 2

TIRF microscopy experiment to probe ACTIF-mediated binding of F-actin to vimentin filaments. First, ACTIF (green) at 5 nM in F-buffer was flushed into the chamber containing settled pre-assembled vimentin filaments (magenta) on the substrate for 10 min. Then, pre-assembled F-actin at 0.1 μM in F-buffer, without the addition of ACTIF, was flushed into the chamber. F-actin rapidly attached and co-localized with ACTIF-bound vimentin filaments.