1

2 **Title**: Myogenic artifacts masquerade as neuroplasticity in the auditory frequency-following

3 response (FFR)

Author(s): Gavin M. **Bidelman**[1,2,3,*], Alexandria **Sisson**[1], Rose **Rizzi**[1,2,] Jessica **MacLean**[1,2],

and Kaitlin **Baer**[4,5]

Affiliations:

[1]Department of Speech, Language and Hearing Sciences, Indiana University, Bloomington,

IN, USA

[2]Program in Neuroscience, Indiana University, Bloomington, IN, USA

[3]Cognitive Science Program, Indiana University, Bloomington, IN, USA

[4]School of Communication Sciences & Disorders, University of Memphis, Memphis, TN, USA

[5]VA Medical Center, Memphis, TN, USA

Author contributions: GMB designed the experiment, AS, RR, JM, and KB collected the

data, GMB analyzed the data, and GMB, AS, RR, JM, and KB wrote the paper.

25
26 **Competing interests**: The authors declare no competing interests.
27
28
29

[*]**Address for editorial correspondence:**
Gavin M. Bidelman, PhD
Department of Speech, Language and Hearing Sciences
Indiana University
2631 East Discovery Parkway
Bloomington IN, 47408
TEL: (812) 855-9339
Email: gbidel@indiana.edu

Abstract

The frequency-following response (FFR) is an evoked potential that provides a neural index of complex sound encoding in the brain. FFRs have been widely used to characterize speech and music processing, experience-dependent neuroplasticity (e.g., learning, musicianship), and biomarkers for hearing and language-based disorders that distort receptive communication abilities. It is widely assumed FFRs stem from a mixture of phase-locked neurogenic activity from brainstem and cortical structures along the hearing neuraxis. Here, we challenge this prevailing view by demonstrating upwards of ~50% of the FFR can originate from a non-neural source: contamination from the postauricular muscle (PAM) vestigial startle reflex. We measured PAM, transient auditory brainstem responses (ABRs), and sustained frequency-following response (FFR) potentials reflecting myogenic (PAM) and neurogenic (ABR/FFR) responses in young, normal-hearing listeners with varying degrees of musical training. We first establish PAM artifact is present in all ears, varies with electrode proximity to the muscle, and can be experimentally manipulated by directing listeners' eye gaze toward the ear of sound stimulation. We then show this muscular noise easily confounds auditory FFRs, spuriously amplifying responses by 3-4x fold with tandem PAM contraction and even explaining putative FFR enhancements observed in highly skilled musicians. Our findings expose a new and unrecognized myogenic source to the FFR that drives its large inter-subject variability and cast doubt on whether changes in the response typically attributed to neuroplasticity/pathology are solely of brain origin.

Keywords: Auditory brainstem response (ABR); auditory evoked potentials (AEPs); frequency-following response (FFR); post-auricular muscle reflex (PAMR); experience-dependent plasticity

Introduction

Neuroelectric brain recordings have been indispensable in demonstrating neuroplasticity at all

levels of the auditory system. In particular, the frequency-following response (FFR), a scalp-

recorded neurophonic reflecting phase-locked activity along the auditory neuroaxis to periodic

signals, has served as a neural index of sound coding in the EEG. The precision of the FFR is

evidenced by the fact these brain potentials can support intelligible speech when they are

replayed (i.e., sonified) as audio stimuli (Bidelman, 2018a). The strength to which FFRs

capture voice pitch (i.e., fundamental frequency; F0) and harmonic timbre cues of complex

signals is also related to listeners' perception of speech material (Weiss and Bidelman, 2015).

FFRs are also enhanced by various experiential factors including native language experience

(Krishnan et al., 2010; Zhao and Kuhl, 2018), musical abilities (Wong et al., 2007; Mankel and

Bidelman, 2018), and perceptual learning (Reetzke et al., 2018). Conversely, FFRs reveal

deficiencies in neural processing in a variety of auditory, literacy, and neurodevelopmental

disorders (e.g., dyslexia, autism, hearing loss, aging; Russo et al., 2008; Chandrasekaran et

al., 2009; Anderson et al., 2013; White-Schwoch et al., 2015; Bidelman et al., 2017a).

Collectively, a wealth of studies implies FFRs might be a valuable auditory biomarker for

tracking both positive and maladaptive auditory plasticity in the brain that either bolsters or

compromises the perceptual organization of speech and musical sounds.

92   Despite an abundance of FFR studies and potential applications to understanding brain

plasticity and normal and disordered central auditory processing, the anatomical origins of the

FFR remain highly contentious (Coffey et al., 2016; Holmes and Herrmann, 2017; Bidelman,

2018b; Coffey et al., 2019; White-Schwoch et al., 2019). Historically described as a brainstem

potential (Smith et al., 1975; Sohmer and Pratt, 1977; Chandrasekaran and Kraus, 2010), it is

now recognized FFRs reflect a mixture of phase-locked activity from brainstem and cortical

structures throughout the auditory pathway (Bidelman, 2018b; Coffey et al., 2019). While

different neuroimaging techniques emphasize brainstem- and cortico-centric contributions to

the response, the FFR has always been described, unequivocally, as a *brain response* of

auditory-neurogenic origin (Hoormann et al., 1992; Chandrasekaran and Kraus, 2010;

Bidelman and Powers, 2018). However, anecdotal observations as earlier as the late 1970s

(Sohmer et al., 1977)[1], and our own experience over the past decade reveal occasional

---

[1]Sohmer et al. (1977) describe two cases studies from "normal subjects with an unusually large
response from the post-auricular muscle (p.656)." They observed, "the amplitude of the post-auricular
responses to click stimuli and this frequency following of the post-auricular muscle were both
susceptible to head position, that is to the degree of neck tension (p.660)" and concluded the PAM

listeners that produce unusually large FFRs that far exceed the amplitudes expected for an

auditory-neurogenic potential in humans. Such enigmatic responses are easily obscured in

grand average data but can differ from the FFRs' normal operating range (Chandrasekaran

and Kraus, 2010) by an order of magnitude (i.e., 1-2 µV vs. 100-200 nV range), and thus

warrant further explanation. Here, we expose a heretofore unrecognized *myogenic* source of

the FFR that explains substantial individual differences in the response including waveform

enhancements usually attributed to plasticity in central auditory nervous system function

(Krishnan et al., 2005; Musacchia et al., 2007; Wong et al., 2007; Kraus et al., 2009; Parbery-

Clark et al., 2009; Krizman et al., 2012; Kraus et al., 2014; Coffey et al., 2017; Mankel and

Bidelman, 2018; Zhao and Kuhl, 2018).

Situated behind the ear, the postauricular muscle (PAM) is part of a vestigial startle

reflex that once acted to retract the pinna to protect hearing (Bérzin and Fortinguerra, 1993).

The muscle produces large (>100 µV), bilateral contraction with peak latency of 9-15 ms

following brief sounds (Thornton, 1975; O'Beirne and Patuzzi, 1999) and is highly variable

across listeners (Picton et al., 1974). Anesthetic blockage of the muscle and facial nerve

abolishes PAM confirming its non-neural origin (Bickford et al., 1964). While PAM activation

declines with increasing stimulus rates (Geisler et al., 1958), the reflex can be elicited by

surprisingly fast periodic stimuli without fatigue or habituation (Jacobson et al., 1964) (but see

Yoshie and Okudaira, 1969). Indeed, motor units underlying the PAM reflex can be driven at

rates up to 100-200 Hz (Kiang et al., 1963; Jacobson et al., 1964), resulting in steady-state

muscle potentials that, like the FFR, phase-lock to periodic signals (Kiang et al., 1963).

Problematically, these rates closely coincide with the low-F0 (~100 Hz) pitched stimuli used in

most auditory FFR studies (Coffey et al., 2016). This raises the possibility that FFRs to low-

frequency sounds, including the voice pitch and low harmonics of speech, might be partially

driven by muscular rather than auditory neurophonic structures. This is particularly relevant in

the context of typical FFR recording approaches, which place electrodes behind the ear where

PAM reflex is optimally recorded (O'Beirne and Patuzzi, 1999). Indeed, in our informal survey

of FFR studies published in the past 50 years (N=314 studies), nearly half (42%=131 papers;

see *SI Materials*) used a mastoid reference electrode montage that promotes extraneous

pickup of PAM muscle artifact (**Fig. 1a**). The problem thus appears widespread in the

literature. More critically, identifying undocumented muscular confound in the FFR would be

particularly germane to interpreting plasticity studies and the unchallenged assumption that

amplified FFRs (e.g., as in musicians, bilinguals) necessarily reflect increased fidelity of

---

contributes to the FFR. Unfortunately, no quantitative analysis was undertaken so these observations
remain qualitative.

auditory neural coding due to listening experience (cf. Musacchia et al., 2007; Wong et al.,

2007; Kraus et al., 2009; Parbery-Clark et al., 2009; Bidelman et al., 2011; Krizman et al.,

2012; Coffey et al., 2017; Mankel and Bidelman, 2018; Zhao and Kuhl, 2018).

Among the more widely reported neuroplastic effects captured in the FFR are the

putative enhancements reported in musicians (Schneider et al., 2002; Musacchia et al., 2007;

Wong et al., 2007; Kraus and Chandrasekaran, 2010). Speech-FFRs are stronger and shorter

in latency in musicians relative to their nonmusicians peers (Kraus et al., 2009; Parbery-Clark

et al., 2009), providing a neural account of their enhanced speech perception observed

behaviorally. Such enhancements are typically interpreted as reflecting stronger neural

representation for speech shaped by experience-dependent plasticity from the enriched sonic

environment afforded by musical engagement (Wong et al., 2007; Kraus and

Chandrasekaran, 2010; Herholz and Zatorre, 2012). However, recent studies have challenged

this notion by demonstrating even nonmusicians with superior music-listening abilities (i.e.,

"musical sleepers") have "enhanced" FFRs that mirror those of highly trained musicians

(Mankel and Bidelman, 2018). Such findings reveal innate differences in auditory brain

function (and possibly other unmeasured factors) can easily masquerade as plasticity in

studies on the brain benefits of music (Musacchia et al., 2007; Wong et al., 2007; Tierney et

al., 2015). The present study was not intended to refute the possible connections between

musicianship and FFR enhancements observed in both cross-sectional and longitudinal

studies (Musacchia et al., 2007; Wong et al., 2007; Tierney et al., 2015; Nan et al., 2018).

Rather, we aimed to identify whether PAM muscle activity naturally induced by sound

stimulation might not only be larger in musicians but partially mediate the auditory processing

enhancements reported in studies on music-related plasticity.

To this end, we evaluated whether unmeasured muscle artifact might explain

substantial inter-subject variability in the FFR and account for at least some of the neural

enhancements in speech processing frequently reported in highly skilled listeners (e.g.,

musicians). A major source of variation in the strength of PAM elicitation stems from

uncontrolled eye movements (Patuzzi and O'Beirne, 1999). Gaze directed toward the ear of

sound presentation potentiates PAM contraction and thus pickup of the muscle artifact in EEG

(O'Beirne and Patuzzi, 1999; Patuzzi and O'Beirne, 1999). By experimentally manipulating

PAM with a gaze paradigm, we show here that repeated stimulation of the muscle produces

"following-like" potentials in the EEG via simple linear superposition of overlapping PAM

wavelets (cf. Bidelman, 2015). Moreover, we find this muscle noise easily masquerades as

the auditory FFR and spuriously amplifies the response by 3-4x fold, partially accounting for

FFR enhancements observed in highly skilled musicians. As a solution for recording artifact-

free FFRs, we further show PAM-FFR contamination is easily circumvented by using high-
frequency stimuli and relocating electrodes to a non-cephalic (neck) site.

Materials and Methods

Participants. The sample included N = 20 young adults (age [$\mu \pm \sigma$]: 24.9 ± 2.8 years; 4 male,
16 female). This sample size was determined *a priori* to match comparable studies on auditory
plasticity and the FFR (Parbery-Clark et al., 2009; Bidelman et al., 2014; Coffey et al., 2016).
All had normal hearing (i.e., pure-tone air-conduction thresholds ≤ 25 dB HL; 250-8000 Hz),
similar levels of education (19.1 ± 2.3 years), and reported no previous history of
neuropsychiatric illness. All but one individual were right-handed (77 ± 45% laterality)
(Oldfield, 1971). The sample included a range of formal musical training ($\mu \pm \sigma$: 6.5 ± 7.3
years; range 0-23 years) to assess whether putative enhancements in the FFR reported in
experienced listeners (cf. Musacchia et al., 2007; Wong et al., 2007; Skoe and Kraus, 2012;
Mankel and Bidelman, 2018) might instead result from undocumented PAM muscle artifact
rather than auditory neuroplasticity, *per se.* Music training was treated as a continuous rather
than binary variable since there is disagreement on what constitutes the definition of a
"musician" (Zhang et al., 2020) and we used self-report (as in previous FFR studies) rather
than a formal tests of music listening skills (cf. Mankel and Bidelman, 2018). Each gave
written informed consent in compliance with a protocol approved by Indiana University IRB.

Stimuli. Auditory brainstem responses (ABRs) were recorded to 100 μs clicks. Frequency-
following responses (FFRs) were elicited by complex pitch stimuli (periodic click trains) with
fundamental frequencies (F0s) of either 100 or 200 Hz (Bidelman, 2015). Pulse trains were
constructed using a periodic series of impulses $h(t) = \sum_{n=1}^{\frac{d}{T}} \delta(t - nT),$ where *d* is the duration
of the desired pulse train (here, 100 ms), and *T* is the period between successive impulses
(i.e., 1/F0)**.** Each pulse of the train was constructed using identical clicks (bandwidth, pulse
with, amplitude) to those described for eliciting the ABR. This allowed us to directly test the
assumption that a large amplitude periodic FFR response is a linear superposition of repeated
transient PAM waves (cf. Bidelman, 2015).

EEG recordings. Participants sat in an electro-acoustically shielded booth in a recliner chair
positioned 1 m from the wall. Numbered signs (labeled 1-5) were positioned in the front
hemifield demarcating angles of -70, -35, 0, +35, +70 degrees (left to right) relative to head

center (see **Fig. 1b**)[2]. For each condition, we directed participants' eye gaze to one of the five

angles by cueing the respective number. Compliance was monitored by the experimenter via

a window into the chamber. FFRs were recorded with eye gaze directed at each of the 5

azimuths.

FFRs were recorded using a two-channel vertical montage with Ag/AgCl disc

electrodes placed on the mid-hairline (i.e., midway between Fpz and Fz) referenced to (i) right

(ipsi) mastoid (M2) and (ii) 7[th] cervical vertebra (mid-forehead= ground). Due to technical

error, data from the second channel were not recorded in N=2 listeners and were treated as

missing values. Impedances were ≤ 5 kΩ. EEGs were digitized at 20 kHz using a SmartEP

EEG system (Intelligent Hearing Systems; Miami, FL) with an online passband of 50 - 3000

Hz (+ 60 Hz notch filter) and 100K amplifier gain. Evoked responses were elicited from each

participant in response to right ear presentation at 80 dB SPL through electromagnetically

shielded insert earphones (Ultra-shielded ER3A inserts, 300Ω; IHS, Miami, FL) that eradicated

electromagnetic stimulus artifact from contaminating biological responses (Price and

Bidelman, 2021). Stimuli were presented with fixed, rarefaction polarity at a repetition rate of

9.09/s. Presentation order was randomized both within and across participants. Continuous

EEGs were epoched (*ABR*: 0-24.7 ms; *FFR:* 0-127 ms) and ensemble averaged across trials

to derive evoked responses per condition. Sweeps > ±50 µV were automatically rejected

during online averaging. A total of 2000 artifact-free sweeps were collected per stimulus and

eye gaze position.

Model simulating FFRs from PAM artifact. To evaluate the degree to which FFRs are

explained by PAM artifact, we compared the empirical FFR recordings with derived FFRs,

simulated via simple convolution (e.g., Goldstein and Kiang, 1958; Janssen et al., 1991;

Bidelman, 2015; Carter and Bidelman, 2023). The model presumes the sustained following

response is generated by a series of overlapping onset responses such that the FFR is an

iterated ABR/PAM (see Fig. 1 of ref. Bidelman, 2015). We simulated FFRs by convolving each

listener's PAM-contaminated ABR (see Fig. 1) with a periodic click train of 100 Hz spacing

(i.e., the FFR stimulus F0). This process generated a new ABR/PAM signature at each

stimulus pulse which when strung across time, yielded a complex waveform closely mirroring

the actual FFR. We then assessed correspondence between model-predicted and true FFR

recordings via cross-correlation (20 ms lag search window) (Galbraith et al., 2000; Bidelman,

2015).

---

[2]In practice, 70[0] rotation is difficult to achieve as human eye movements rarely exceed 50-60[0] (Lee et al., 2019). In these extreme conditions, listeners simply panned their eye gaze left/right as far as possible.

Statistical analysis. We analyzed the data using a 2x2 mixed-model ANOVA in *R* (R Core
Team, 2020) and the *lme4* package (Bates et al., 2015). Fixed effects were eye gaze position
(5 levels) and channel montage (2 levels). Subjects served as a random effect. We used
Satterthwaite's method to compute degrees of freedom. The data were SQRT-transformed to
improve normality and homogeneity of variance assumptions necessary for parametric
analyses. Effect sizes are reported as η_p^2.

We conducted regression analyses to assess whether a linear combination of
listeners' neuro-behavioral measures (i.e., years of music training, PAM amplitudes) predicted
the strength of their FFR [e.g., FFR$_{rms}$~ music*PAM]. We then used leverage plots (Sall,
1990), partial correlations to assess the relative strength of music vs. artifact regressors in
driving putative musicianship-FFR relations (e.g., Musacchia et al., 2007; Wong et al., 2007;
Skoe and Kraus, 2012; Mankel and Bidelman, 2018). Mediation analysis was used to evaluate
relationships between neural and behavioral measures. We used an efficient, bootstrapping
implementation of the Sobel statistic (Sobel, 1982; Preacher and Hayes, 2004) (N = 1000
resamples) to determine whether PAM artifact fully or partially mediates the strength of
listeners' FFR. The Sobel test contrasts the strength of regressions between a pairwise vs.
triplet (mediation) model (i.e., *X→Y* vs. *X→M→Y).* Mediator *M* is said to mediate the relation
between the *X→Y* if (i) *X* first predicts *Y* on its own, (ii) *X* predicts *M*, and (iii) the functional
relation between *X→Y* is rendered insignificant after controlling for *M* (Preacher and Hayes,
2004).
255

Results

Postauricular muscle (PAM) artifact is present but variable across ears. We first
confirmed PAM could be successfully elicited in individual ears and experimentally
manipulated by directing listeners' eye gaze towards their stimulated ear (**Fig. 1b**). **Figure 1c**
shows grand average click-ABRs recorded using montages with ipsilateral mastoid vs. midline
(C7) electrode reference. In both traces, eye gaze was directed toward the right (ipsilateral)
ear, which was expected to elicit maximal PAM contraction (O'Beirne and Patuzzi, 1999;
Patuzzi and O'Beirne, 1999). Positive deflections in the first 6 ms form the canonical waves of
the auditory brainstem response (I = 1.5 ms, III=3.5 ms, V=5.5 ms) reflecting serial activation
of the auditory nerve and major nuclei along the ascending auditory pathway. Following the
ABR, the myogenic PAM artifact was evident between 12.5-15 ms and was especially
prominent in mastoid-referenced recordings (i.e., Fpz – M2). PAM artifact was not evident at
C7 where the reference electrode is distanced from the muscle. Consequently, PAM artifact
was easily isolated from the neurogenic ABR by simple waveform subtraction of mastoid vs.

neck recordings (**Fig. 1d**). Although PAM artifact was present in all ears to some degree, the

response was highly variable, ranging from 0.18-9.7 µV (peak-to-peak amplitude; µ±σ: 2.38 ±

2.36 µV) across individuals.

Figure 1: Eye gaze toward the ear introduces PAM artifact in ABRs. (**a**) In our literature review of N=314 FFR studies, nearly half (42%=131 papers) use a mastoid reference electrode montage that can promote inadvertent pickup of muscle artifact (see *SI Materials*). (**b**) Schematic of the visual gaze paradigm to direct listeners' eyes during ABR/FFR recording. Listeners were positioned 1 m from the booth wall and were cued to direct their eye gaze to one of 5 positions spanning ±70 degrees from head center relative to the ipsilateral ear of stimulus presentation (right ear). (**c**) Grand average ABR waveforms recorded with a vertical ipsilateral montage between a non-inverting electrode at the high forehead (~Fpz) referenced to an inverting electrode placed on the (i) ipsilateral mastoid (M2) or (ii) 7th cervical vertebra (C7). Eye gaze was directed to position 5 (70 deg) in both cases. Strong PAM is recorded for the channel with electrode over the ipsilateral mastoid. Moving the reference to C7 (i.e., distal to the PAM muscle) eradicates pickup of the artifact. (**d**) Individual data, with PAM responses isolated via subtraction of the traces in panel C. PAM strength ranges from 0.5-10 µV (peak-to-peak) across individuals. Anatomy adapted from Feneis (1994). Shading = ± 1 s.e.m.

PAM contraction confounds the auditory FFR but declines with stimulus frequency and

electrode montage. Having established PAM artifact was easily recordable and could be

manipulated experimentally via eye gaze, we then tested whether FFR strength might also

systematically vary with eye gaze direction and thus expose a myogenic source of the neural

response. Confirming our intuition, FFRs to complex tones varied parametrically in amplitude

when listeners were cued to direct their gaze across a ±70^0 range (**Fig. 2a**). Hard rotation of

the eyes to either the left or right ear enhanced FFRs by 3-4x, particularly between 100-200

Hz where the PAM resides within the EEG spectrum (O'Beirne and Patuzzi, 1999). FFR

amplitude depended strongly on both eye gaze and recording location [position x channel:

$F_{4,152.16} = 3.72$, $P = 0.0064$; $\eta_p^2 = 0.09$]. Comparisons between the C7 and mastoid channel

showed a ~180^0 phase shift, which is common for FFRs recorded from brainstem (highlighted

in the C7 reference electrode) vs. auditory nerve (highlighted in the M2 reference electrode)

(see Fig 10 in Bidelman, 2018b). For the mastoid channel, FFR strength decreased for gaze

directed toward the midline (0^0) and increased with gaze toward either ear (quadratic contrast:

$t_{152}=6.82$, $P<0.0001$). A nearly identical pattern was observed for the 200 Hz stimulus (data

not shown; position x channel: $F_{1, 50.4} = 8.52$, $P = 0.0052$; $\eta_p^2 = 0.14$). Moreover, a direct

comparison of frequency effects revealed stronger FFRs in the 100 Hz vs. 200 Hz condition

regardless of channel [main effect of frequency: $F_{1, 117.99} = 4.66$, $P = 0.033$; $\eta_p^2 = 0.04$],

indicating PAM-related changes in FFR declined with increasing stimulus frequency.

Myogenic-induced enhancements were not apparent in electrodes positioned far from the

mastoid and FFRs were invariant to eye gaze when recorded from the Fpz-C7 channel (linear

and quadratic contrasts: $P > 0.17$) (**Fig. 2b**).

We also found PAM peak-to-peak amplitude was highly correlated with FFR root-

mean-squared (rms) amplitude [Pearson's-$r = 0.72$, $P = 0.0005$] (**Fig. 2c**). This suggests

listeners' FFR directly scaled with the size of their muscle artifact. Collectively, these findings

reveal FFR strength depends critically on eye gaze position and moreover, that PAM activity

overlays true neural responses, potentially masquerading as the neurophonic FFR commonly

described as having brainstem origin (Chandrasekaran and Kraus, 2010; Bidelman, 2018b;

Coffey et al., 2019).

Sustained FFRs are explained by serial PAM contractions: Model simulations. Single-unit motor neurons underlying the PAM reflex can be driven at rates up to 100-200 Hz (Kiang et al., 1963; Jacobson et al., 1964). This raises the possibility that the FFR, and what appears to be a sustained brain potential, might at least partially reflect a series of overlapping transient PAM artifacts evoked by the individual pitch pulses of low-frequency, periodic sounds. To test this possibility, we simulated FFRs via simple convolution of listeners' isolated PAM response (i.e., Fig. 1) with a 100 Hz impulse train. This train had identical periodicity as the complex tones used in our FFR experiment. This model regenerates the PAM at each pitch period of the F0, such that responses temporally overlap and produce a quasi-steady state auditory response (Galambos et al., 1981). Repeating the PAM wavelet across time (**Fig. 3a**) yielded a sustained waveform that was strikingly similar to the empirical FFR recordings (**Fig. 3b**). Actual and PAM-derived FFRs were highly correlated (r_{xcorr} = 0.71 ± 0.21; *inset*), suggesting upwards of 50.4% of the variance (R^2) in the FFR could be explained by a non-neuronal source.

PAM artifact mediates music-related plasticity in FFRs. Having established that FFRs are

partly confounded by muscular influences, we next asked whether putative enhancements

reported in the literature (e.g., due to musical training, bilingualism (Krishnan et al., 2005;

Musacchia et al., 2007; Wong et al., 2007; Krizman et al., 2012; Skoe and Kraus, 2012;

Mankel and Bidelman, 2018)) might be explained not in terms of auditory plasticity but rather,

differences in tonic PAM contraction among listeners. To this end, we conducted multiple

regression analyses to assess whether a linear combination of listeners' neuro-behavioral

measures (i.e., years of music training, PAM amplitudes) predicted the strength of their FFR

[e.g., FFR_{rms}~ music*PAM]. **Figure 4** shows leverage plots (Sall, 1990) illustrating the relative

effects of music training and PAM artifact amplitudes on FFR strength. Variance inflation

factors (VIFs) were < 2 for both predictor variables, indicating negligible multicollinearity in the

data (Lüdecke, 2020). We found that music training strongly predicted FFR strength [r = 0.34,

P = 0.00076] (**Fig. 4a**), consistent with prior studies suggesting musicianship enhances the

brain's early neural encoding of complex sounds (cf. Musacchia et al., 2007; Wong et al.,

2007; Skoe and Kraus, 2012; Mankel and Bidelman, 2018). However, even after partialing out

musical training, the strength of listeners' PAM remained strongly correlated with FFR

amplitude [r = 0.49, P < 0.0001] (**Fig. 4b**). Similarly, musicianship was independently

correlated with enhanced FFR amplitudes after controlling PAM contributions [r = 0.21, P =

0.037]. Indeed, a regression model that included the interaction between music training and

PAM amplitudes was a significant improvement in describing FFR variance over one

containing music alone [$F_{1,92}$ = 29.01, P < 0.001]. Taken together, these data confirm that the

strength of the FFR is driven by an interaction between neuroplasticity (i.e., related to listening

experience) and artifactual origins.

Figure 4: PAM artifact can masquerade as experience-dependent plasticity in the FFR. (**a**) (*left*) FFR is
stronger in listeners with more musical training, suggestive of the experience-dependent plasticity reported in
trained musicians (cf. Musacchia et al., 2007; Wong et al., 2007; Skoe and Kraus, 2012; Mankel and Bidelman,
2018). Both PAM (*middle*) and musical training (*right*) independently correlate with FFR strength after partialing out
the other variable. Correlation data are aggregated from the 5 eye gaze positions. (**b**) Mediation analysis. The
putative FFR-music relation is severely reduced after accounting for PAM [Sobel test: z=2.72, P=0.006] suggesting

To further evaluate the confounding link between PAM-related muscle noise and

putative music-induced FFR plasticity (e.g., Musacchia et al., 2007; Wong et al., 2007; Skoe

and Kraus, 2012; Mankel and Bidelman, 2018), we conducted mediation analyses (Sobel,

1982; Preacher and Hayes, 2004) to determine the degree to which PAM artifact mediated

FFR strength (**Fig. 4b**). Mediation contrasts the strength of regressions between a pairwise

vs. triplet (mediation) model (i.e., $X{\rightarrow}Y$ vs. $X{\rightarrow}M{\rightarrow}Y$). Mediator M is said to mediate the

relation between the $X{\rightarrow}Y$ if (i) X first predicts Y on its own, (ii) X predicts M, and (iii) the

functional relation between $X{\rightarrow}Y$ is rendered insignificant after controlling for M (Preacher and

Hayes, 2004). We found PAM satisfied the first two criteria for a mediating variable (**Fig. 4b**).

However, accounting for the strength of listeners' PAM artifact severely reduced—but did not

render insignificant—relations between music training and FFR strength (Sobel test: z=2.72,

P=0.006) (Sobel, 1982). These results suggest PAM artifact is a partial (but not fully)

mediating variable describing music-related enhancements observed in the FFR.

FFR timing is independent of PAM. The analyses thus far revealed links been amplitude

properties of the FFR and PAM reflex. However, plasticity in the FFR has also been described

in the latency of the response to speech and musical sounds, with faster neural timing

following long-term musicianship (Parbery-Clark et al., 2012; Bidelman et al., 2014; Mankel

and Bidelman, 2018) and short-term auditory training (Anderson et al., 2013). To explore the

possibility that variation in FFR timing might also be due to undocumented PAM artifact, we

examined correlations between the onset latency of the FFR [measure in the 6-12 ms search

window (Mankel and Bidelman, 2018)] and PAM artifact [8-18 ms; see Fig. 1d]. However,

neither latency [r = -0.10, P = 0.69] nor amplitude [r = 0.14, P = 0.57] of the PAM predicted

FFR timing to the 100 Hz stimuli. Similar null correspondence was observed at 200 Hz [all Ps

> 0.09]. Thus, in stark contrast to FFR amplitude measures, fine timing precision of the FFR

appears less susceptible to artifactual influence.

Discussion

Our findings expose a strong influence of myogenic activity on the auditory FFR. We

demonstrate uncontrolled eye gaze systematically alters the apparent strength of FFRs with

tandem PAM contraction. More critically, we show this muscular noise can easily masquerade

as neural enhancements, accounting for large variability in the response previously ascribed

to auditory-sensory plasticity. Importantly, these data do not negate the possibility that certain

human experiences (e.g., music training, language expertise, learning) can confer experience-

dependent plasticity that manifests in spectrotemporal changes in the FFR (Krishnan et al.,

2005; Kraus et al., 2014; Mankel and Bidelman, 2018). Rather, we argue that undue muscle

confounds external to the brain may play a larger role in generating the FFR and its use as a

biomarker of plasticity than conventionally thought.

While typically considered an artifactual nuisance to auditory EEG and evoked

potential recordings, the PAM and related myogenic potentials do find several important

neuro-otological applications including hearing threshold (Yoshie and Okudaira, 1969) and

vestibular assessment (Rosengren et al., 2019). Yet, in the context of FFRs whose purpose is

to evaluate central auditory neural processing and the brain's sensory representation of

complex sounds, we show concomitant PAM muscle activation easily confounds these

neurogenic responses, spuriously amplifying the FFR by 3-4x fold. Our findings are

reminiscent of other microflexes shown to confound oscillatory EEG responses (e.g.,

microsaddes Yuval-Greenberg et al., 2008) and suggest uncontrolled eye movements can

systematically alter sustained auditory FFR potentials via PAM muscle engagement.

We found PAM-related artifact was largest for hard rotation of the eyes toward or way

from the ipsilateral ear of sound presentation but parametrically scaled in direct portion with

listeners' lateral eye gaze. While it is important to note that PAM was evoked up to a rather

extreme eye gaze rotation (representing a "worst case scenario"), we also find the artifact

scales in direct proportion away from midline and persists at gaze angles typical of normal eye

movements. This point is particularly salient in light of speech-FFR studies that use passive

listening paradigms where listeners are allowed to watch muted subtitled movies (Wong et al.,

2007; Coffey et al., 2016; Mankel and Bidelman, 2018; Price and Bidelman, 2021). Videos are

thought to provide added control during electrophysiological procedures, inducing a calm yet

wakeful state that yields cleaner electrophysiological recordings (Wong et al., 2007; Coffey et

al., 2016). However, subtended gaze angles when viewing a monitor at typical seated

distance can exceed ~30-35 degrees (Feng and Spence, 2008). Our data clearly show the

FFR is nearly double its nominal size at these typical viewing angles (Fig. 2b). Consequently,

while seemingly an innocuous control task, undue eye movements during video watching

could easily confound the FFR by artificially inflating response amplitude. Along these lines,

there is evidence that musicians show reduced fixation dwell times and more saccades during

viewing tasks (Perra et al., 2022). Such increased eye movement could explain the larger

PAM activation (and artifactual FFR) we find in musically trained individuals (Fig. 4). Similar

augments could extend to bilinguals, who also show increased visual search (Ratiu et al.,

2017) and more robust speech FFRs (Krishnan et al., 2005; Krizman et al., 2012).

While we acknowledge our gaze-induced manipulation of PAM could represent an
overestimate of its activation, we note that PAM can be elicited down to hearing threshold
(Purdy et al., 2005). Moreover, the strong relation between PAM and FFR amplitudes we find
would remain unaffected since changes in scale affect neither the magnitude nor sign of a
correlation. Thus, the fact that 50%, let alone *any* portion of variance in the FFR is explained
by a myogenic artifact is a cause for concern for FFR recordings.

While there is overwhelming evidence in humans and animals to suggest PAM artifact
is generated from the bipolar activation of muscle fibers behind the auricle of the ear (Kiang et
al., 1963; O'Beirne and Patuzzi, 1999; Patuzzi and O'Beirne, 1999), it is conceivable that other
non-auditory mechanisms contribute to FFR contamination. For example, it is well known that
FFRs require delicate recording strategies to prevent inadvertent pickup of stimulus
headphone artifact that can easily swamp the biological response (Price and Bidelman, 2021).
Electromagnetic shielding (used here) prevents this possibility (Price and Bidelman, 2021) and
can be further ruled out in the present data by the fact that FFRs showed gaze- and electrode-
dependent changes that would not occur if waveforms were mere stimulus bleed (Fig. 2b).
However, bogus FFR amplification could also result from other biological sources. Indeed,
bone-conducted FFRs have been recorded in profoundly deaf listeners suggesting non-
auditory pathways of response generation (Ribarić et al., 1984). These might include direct or
indirect contributions from the vestibular system [e.g., evoked myogenic potentials (Ribarić et
al., 1984; Prevec and Ribarić-Jankes, 1996; Lawlor et al., 2022)] and/or somatosensory
contributions near the electrode-skin interface which can persist up to rates of 287 (Lawlor et
al., 2022) and 130 Hz (Prevec and Ribarić-Jankes, 1996), respectively. Regardless of the
underlying mechanism, our data clearly demonstrate physiological noise outside the auditory
system easily conflates neurogenic FFR signals.

Several studies have shown enhanced voice pitch (F0) and timbre (harmonic)
encoding in musically trained listeners (Musacchia et al., 2007; Wong et al., 2007; Kraus et
al., 2009; Parbery-Clark et al., 2009; Kraus and Chandrasekaran, 2010; Mankel and
Bidelman, 2018), though not always consistently (e.g., see Strait et al., 2012; Bidelman and
Alain, 2015). Our data confirm a robust and independent relation between musical training
and the strength of listeners' auditory FFR (Musacchia et al., 2007; Wong et al., 2007; Skoe
and Kraus, 2012; Mankel and Bidelman, 2018) (Fig. 4a). However, in a departure from prior
studies, we show such effects are easily confounded. PAM contraction was both larger in
listeners with more extended music training and partially mediated enhancements in their FFR
(Fig. 4b). At first glance, these data therefore cast doubt on whether speech-FFR
enhancements associated with musical training (and perhaps other experiential factors) are

solely experience-driven (cf. Munte et al., 2002; Wong et al., 2007; Herholz and Zatorre, 2012)

or even entirely auditory-neurogenic in nature. Undue PAM influence might explain why some

studies fail to observe FFR advantages in musicians (Strait et al., 2012; Bidelman and Alain,

2015; MacLean et al., in press). Nevertheless, we show that even after controlling for PAM,

music training remains a robust predictor of FFR enhancements. This indicates musicianship

independently bolsters the strength of the FFR and does so above and beyond any artifactual

sources.

Surprisingly, we found musicians have stronger PAM vestigial muscle reflexes than

their nonmusician peers. This finding broadly converges with notions that intense auditory

experiences might fortify tonic engagement of reflexive pathways that provide feedback

control to peripheral hearing processes (Brashears et al., 2003; Bidelman et al., 2017b). Still,

it remains to be seen whether other long-term listening experiences that also enhance FFR

similarly elevate PAM and account for neuroplastic effects observed in bilingualism and tone-

language expertise (Krishnan et al., 2005; Krizman et al., 2012; Zhao and Kuhl, 2018).

Regardless, the intrusive nature of PAM mandates caution for interpretating even short-term

changes in the FFR, as observed in rapid perceptual learning studies (Anderson et al., 2013;

Reetzke et al., 2018). More successful learners could show increased arousal while acquiring

novel auditory information, leading to increased eye movements that exaggerate PAM artifact,

and in turn false amplification of FFRs. FFRs are modulated by listeners' arousal state at the

single-trial level (Lai et al., 2022; Carter and Bidelman, 2023) and eye movements change

during the learning process (Laamerad et al., 2020). Thus, while it awaits empirical

confirmation, our data raise the possibility that rapid plasticity observed in FFRs during short-

term auditory/speech learning tasks might be due to not improvements in the brain's sensory

representation of sound, *per se*, but an unmeasured artifactual source (Anderson et al., 2013;

Reetzke et al., 2018).

Fortunately, our data offer a direct solution to easily thwart PAM-related confounds in

the FFR and allow for untainted assessment of neuroplasticity. FFRs are frequently recorded

using a differential electrode montage and non-inverting reference electrode placed on the

bony mastoid process of the ipsilateral ear of stimulation (e.g., Fpz-M1/M2; see Fig. 1c, inset).

We show this common practice is highly problematic, leading to extraneous pickup of the

adjacent PAM muscle and overinflation of FFR that, by our estimates, has affected nearly half

the published literature (see Fig. 1a and *SI Materials*). Repositioning electrodes to the earlobe

(another common technique) (Chandrasekaran and Kraus, 2010; Skoe and Kraus, 2010)

attenuates but does not fully eliminate PAM pickup (O'Beirne and Patuzzi, 1999). Instead, we

find clean FFRs are easily recorded with electrodes placed on the upper neck (C7 vertebra),

distal to the pinna and PAM muscle fibers. A midline Fpz-C7 montage has the further

advantage that it (i) reduces pickup of more peripheral (cochlear) sources of the FFR and (ii)

is optimally oriented with the vertical dipolar sources in the brainstem that dominate

generation of the neurogenic FFR (Galbraith et al., 2000; Bidelman, 2015; 2018b). At present,

it appears only 16% of studies use this optimal C7/neck montage (Fig. 1a). Secondly, we

showed that FFR latency measures were largely independent of artifactual influences. Thus,

timing characteristics of the FFR might provide a more veridical index of neuroplasticity

(Parbery-Clark et al., 2012; Anderson et al., 2013; Bidelman et al., 2014; Mankel and

Bidelman, 2018) than amplitude measures alone when tracking brain changes due to

experience-dependent factors and novel sound learning (but see Mankel and Bidelman,

2018). Thirdly, controlling participants' eye gaze straight forward (e.g., with a fixation cross) or

instructing them to close their eyes could help minimize PAM contributions to the FFR. Still,

we note that changes in eye gaze within only a small viewing angle could affect FFR and eye

closer does not prevent occular movement altogether as the eyes are still free to move under

the eyelid. Lastly, we found that PAM-related changes in FFR declined with increasing

stimulus frequency. This suggests that in addition to isolating brainstem-centric sources of the

response without cortical contributions (Bidelman, 2018b), the use of auditory stimuli with

higher F0s (e.g., > 200 Hz) might be further advantageous in protecting against PAM

confounds in the FFR.

In conclusion, our data provide new evidence that myogenic sources can artificially

inflate the sound-evoked FFR and might account for certain *de novo* enhancements in the

response. Provocatively, they also show that formally trained musicians have additional gains

in sensory brain processing above and beyond PAM-related confounds. Our results

emphasize the critical need for future studies to safeguard against ocular and myogenic

confounders before claiming changes in FFR track with improvements in auditory brain

function due to experience-dependent factors or neurobehavioral interventions. Of practical

implication, we recommend that to measure true neurogenic FFRs without undue

electromyographic contamination, investigators should (i) adopt a midline, necked-referenced

recording approach, (ii) use high-frequency F0 stimuli (> 200 Hz) where PAM contributions are

minimal, and (iii) focus primarily on response latency quantification to index disordered

auditory processing and/or neuroplastic changes due to intervention.

Acknowledgments

Work supported by the National Institutes of Health (NIH/NIDCD R01DC016267).

Data availability

Data associated with this manuscript are provided in the *Supplemental Material*.

Competing interests

The authors have no relevant financial or non-financial interests to disclose.

References

Anderson, S., White-Schwoch, T., Parbery-Clark, A., and Kraus, N. (2013). Reversal of age-
related neural timing delays with training. *Proceedings of the National Academy of*
Sciences of the United States of America 110**,** 4357-4362.

Bates, D., Mächler, M., Bolker, B., and Walker, S. (2015). Fitting linear mixed-effects models
using lme4. *Journal of Statistical Software* 67**,** 1-48.

Bérzin, F., and Fortinguerra, C.R. (1993). EMG study of the anterior, superior and posterior
auricular muscles in man. *Ann Anat* 175**,** 195-197.

Bickford, R.G., Jacobson, J.L., and Cody, D.T. (1964). Nature of average evoked potentials to
sound and other stimuli in man. *Annals of the New York Academy of Sciences* 112**,**
204-223.

Bidelman, G., and Powers, L. (2018). Response properties of the human frequency-following
response (FFR) to speech and non-speech sounds: level dependence, adaptation and
phase-locking limits. *International Journal of Audiology* 57**,** 665–672.

Bidelman, G.M. (2015). Multichannel recordings of the human brainstem frequency-following
response: Scalp topography, source generators, and distinctions from the transient
ABR. *Hearing Research* 323**,** 68-80.

Bidelman, G.M. (2018a). Sonification of scalp-recorded frequency-following responses (FFRs)
offers improved response detection over conventional statistical metrics. *Journal of*
Neuroscience Methods 293**,** 59-66.

Bidelman, G.M. (2018b). Subcortical sources dominate the neuroelectric auditory frequency-
following response to speech. *NeuroImage* 175**,** 56–69.

Bidelman, G.M., and Alain, C. (2015). Musical training orchestrates coordinated
neuroplasticity in auditory brainstem and cortex to counteract age-related declines in
categorical vowel perception. *Journal of Neuroscience* 35**,** 1240 –1249.

Bidelman, G.M., Gandour, J.T., and Krishnan, A. (2011). Cross-domain effects of music and
language experience on the representation of pitch in the human auditory brainstem.
Journal of Cognitive Neuroscience 23**,** 425-434.

Bidelman, G.M., Lowther, J.E., Tak, S.H., and Alain, C. (2017a). Mild cognitive impairment is
characterized by deficient hierarchical speech coding between auditory brainstem and
cortex. *Journal of Neuroscience* 37**,** 3610-3620.

Bidelman, G.M., Schneider, A.D., Heitzmann, V.R., and Bhagat, S.P. (2017b). Musicianship
enhances ipsilateral and contralateral efferent gain control to the cochlea. *Hearing*
Research 344**,** 275-283.

Bidelman, G.M., Weiss, M.W., Moreno, S., and Alain, C. (2014). Coordinated plasticity in
brainstem and auditory cortex contributes to enhanced categorical speech perception
in musicians. *European Journal of Neuroscience* 40**,** 2662-2673.

Brashears, S.M., Morlet, T.G., Berlin, C.I., and Hood, L.J. (2003). Olivocochlear efferent
suppression in classical musicians. *Journal of the American Academy of Audiology* 14**,**
314-324.

Carter, J.A., and Bidelman, G.M. (2023). Perceptual warping exposes categorical
representations for speech in human brainstem responses. *NeuroImage* 269**,** 119899.

Chandrasekaran, B., Hornickel, J., Skoe, E., Nicol, T., and Kraus, N. (2009). Context-
dependent encoding in the human auditory brainstem relates to hearing speech in
noise: Implications for developmental dyslexia. *Neuron* 64**,** 311-319.

Chandrasekaran, B., and Kraus, N. (2010). The scalp-recorded brainstem response to
speech: Neural origins and plasticity. *Psychophysiology* 47**,** 236-246.

Coffey, E.B., Herholz, S.C., Chepesiuk, A.M., Baillet, S., and Zatorre, R.J. (2016). Cortical
contributions to the auditory frequency-following response revealed by MEG. *Nature*
Communications 7**,** 11070.

Coffey, E.B.J., Chepesiuk, A.M.P., Herholz, S.C., Baillet, S., and Zatorre, R.J. (2017). Neural
correlates of early sound encoding and their relationship to speech-in-noise
perception. *Frontiers in Neuroscience* 11.

Coffey, E.B.J., Nicol, T., White-Schwoch, T., Chandrasekaran, B., Krizman, J., Skoe, E.,
Zatorre, R.J., and Kraus, N. (2019). Evolving perspectives on the sources of the
frequency-following response. *Nature Communications* 10**,** 5036.

Feneis, H. (1994). *Pocket Atlas of Human Anatomy.* Thieme Publishing Group.

Feng, J., and Spence, I. (2008). "Attending to large dynamic displays", in: *CHI '08 Extended*
Abstracts on Human Factors in Computing Systems. (Florence, Italy: Association for
Computing Machinery).

Galambos, R., Makeig, S., and Talmachoff, P.J. (1981). A 40-Hz auditory potential recorded
from the human scalp. *Proceedings of the National Academy of Sciences of the United*
States of America 78**,** 2643-2647.

Galbraith, G., Threadgill, M., Hemsley, J., Salour, K., Songdej, N., Ton, J., and Cheung, L.
(2000). Putative measure of peripheral and brainstem frequency-following in humans.
Neuroscience Letters 292**,** 123-127.

Geisler, C.D., Frishkopf, L.S., and Rosenblith, W.A. (1958). Extracranial responses to acoustic
clicks in man. *Science* 128**,** 1210-1211.

Goldstein, M.H., and Kiang, N.Y.S. (1958). Synchrony of neural activity in electric responses
evoked by transient acoustic stimuli. *Journal of the Acoustical Society of America* 30**,**
107–114.

Herholz, S.C., and Zatorre, R.J. (2012). Musical training as a framework for brain plasticity:
Behavior, function, and structure. *Neuron* 76**,** 486-502.

Holmes, E., and Herrmann, B. (2017). Revisiting the contribution of auditory cortex to
frequency-following responses. *Journal of Neuroscience* 37**,** 5218-5220.

Hoormann, J., Falkenstein, M., Hohnsbein, J., and Blanke, L. (1992). The human frequency-
652   following response (FFR): Normal variability and relation to the click-evoked brainstem
653   response. *Hearing Research* 59**,** 179-188.

654   Jacobson, J.L., Cody, D.T., Lambert, E.H., and Bickford, R.G. (1964). Physiological Properties
655   of the Post-Auricular Responses (Sonomotor) in Man. *The Physiologist* 7**,** 167-167.

656   Janssen, T., Steinhoff, H.J., and Böhnke, F. (1991). Zum entstehungsmechanismus der
657   frequenzfolgepotentiale. *Otorhinolaryngol. Nova* 1**,** 16-24.

658   Kiang, N.Y., Crist, A.H., French, M.A., and Edwards, A.G. (1963). Postauricular electrical
659   response to acoustic stimuli in humans. *Massachusetts Institute of Technology*
660   *Quarterly Progress Report* No. 68**,** 218-225.

661   Kraus, N., and Chandrasekaran, B. (2010). Music training for the development of auditory
662   skills. *Nature Reviews Neuroscience* 11**,** 599-605.

663   Kraus, N., Skoe, E., Parbery-Clark, A., and Ashley, R. (2009). Experience-induced malleability
664   in neural encoding of pitch, timbre, and timing. *Annals of the New York Academy of*
665   *Sciences* 1169**,** 543-557.

666   Kraus, N., Slater, J., Thompson, E.C., Hornickel, J., Strait, D.L., Nicol, T., and White-
667   Schwoch, T. (2014). Music enrichment programs improve the neural encoding of
668   speech in at-risk children. *Journal of Neuroscience* 34**,** 11913-11918.

669   Krishnan, A., Gandour, J.T., and Bidelman, G.M. (2010). The effects of tone language
670   experience on pitch processing in the brainstem. *Journal of Neurolinguistics* 23**,** 81-95.

671   Krishnan, A., Xu, Y., Gandour, J.T., and Cariani, P. (2005). Encoding of pitch in the human
672   brainstem is sensitive to language experience. *Brain Research Cognitive Brain*
673   *Research* 25**,** 161-168.

674   Krizman, J., Marian, V., Shook, A., Skoe, E., and Kraus, N. (2012). Subcortical encoding of
675   sound is enhanced in bilinguals and relates to executive function advantages.
676   *Proceedings of the National Academy of Sciences of the United States of America*
677   109**,** 7877-7881.

678   Laamerad, P., Guitton, D., and Pack, C.C. (2020). Eye movements shape visual learning.
679   *Proc Natl Acad Sci U S A* 117**,** 8203-8211.

680   Lai, J., Price, C.N., and Bidelman, G.M. (2022). Brainstem speech encoding is dynamically
681   shaped online by fluctuations in cortical α state. *NeuroImage* 263**,** 119627.

682   Lawlor, K.J., Clinard, C.G., and Piker, E.G. (2022). Temporal Modulation Transfer Functions
683   of Amplitude-Modulated Cervical Vestibular-Evoked Myogenic Potentials in Young
684   Adults. *Ear and Hearing* 43.

685   Lee, W.J., Kim, J.H., Shin, Y.U., Hwang, S., and Lim, H.W. (2019). Differences in eye
686   movement range based on age and gaze direction. *Eye (Lond)* 33**,** 1145-1151.

687   Lüdecke, D. (2020). "sjPlot: Data visualization for statistics in social science. R package
688   version 2.8.6. https://CRAN.R-project.org/package=sjPlot".).

Maclean, J.A., Stirn, J.R., Sisson, A.E., and Bidelman, G.M. (in press). Short- and long-term
neuroplasticity interact during the perceptual learning of concurrent speech. *Cerebral*
Cortex 10.1093/cercor/bhad543.

Mankel, K., and Bidelman, G.M. (2018). Inherent auditory skills rather than formal music
training shape the neural encoding of speech. *Proceedings of the National Academy of*
Sciences of the United States of America 115**,** 13129-13134.

Munte, T.F., Altenmuller, E., and Jancke, L. (2002). The musician's brain as a model of
neuroplasticity. *Nature Reviews Neuroscience* 3**,** 473-478.

Musacchia, G., Sams, M., Skoe, E., and Kraus, N. (2007). Musicians have enhanced
subcortical auditory and audiovisual processing of speech and music. *Proceedings of*
the National Academy of Sciences of the United States of America 104**,** 15894-15898.

Nan, Y., Liu, L., Geiser, E., Shu, H., Gong, C.C., Dong, Q., Gabrieli, J.D.E., and Desimone, R.
(2018). Piano training enhances the neural processing of pitch and improves speech
perception in Mandarin-speaking children. *Proceedings of the National Academy of*
Sciences of the United States of America 115**,** E6630-E6639.

O'beirne, G.A., and Patuzzi, R.B. (1999). Basic properties of the sound-evoked post-auricular
muscle response (PAMR). *Hearing Research* 138**,** 115-132.

Oldfield, R.C. (1971). The assessment and analysis of handedness: The Edinburgh inventory.
Neuropsychologia 9**,** 97-113.

Parbery-Clark, A., Anderson, S., Hittner, E., and Kraus, N. (2012). Musical experience offsets
age-related delays in neural timing. *Neurobiology of aging* 33**,** 1483 e1481-1484.

Parbery-Clark, A., Skoe, E., and Kraus, N. (2009). Musical experience limits the degradative
effects of background noise on the neural processing of sound. *Journal of*
Neuroscience 29**,** 14100-14107.

Patuzzi, R.B., and O'beirne, G.A. (1999). Effects of eye rotation on the sound-evoked post-
auricular muscle response (PAMR). *Hearing Research* 138**,** 133-146.

Perra, J., Latimier, A., Poulin-Charronnat, B., Baccino, T., and Drai-Zerbib, V. (2022). A Meta-
analysis on the Effect of Expertise on Eye Movements during Music Reading. *J Eye*
Mov Res 15.

Picton, T.W., Hillyard, S.A., Krausz, H.I., and Galambos, R. (1974). Human auditory evoked
potentials. I. Evaluation of components. *Electroencephalography and Clinical*
Neurophysiology 36**,** 179-190.

Preacher, K.J., and Hayes, A.F. (2004). SPSS and SAS procedures for estimating indirect
effects in simple mediation models. *Behavior Research Methods, Instruments, &*
Computers 36**,** 717-731.

Prevec, T.S., and Ribarić-Jankes, K. (1996). Can somatosensory system generate frequency
following response? *Pflugers Arch* 431**,** R301-302.

Price, C.N., and Bidelman, G.M. (2021). Attention reinforces human corticofugal system to aid
speech perception in noise. *NeuroImage* 235**,** 118014.

Purdy, S.C., Agung, K.B., Hartley, D., Patuzzi, R.B., and O'beirne, G.A. (2005). The post-
auricular muscle response: an objective electrophysiological method for evaluating
hearing sensitivity. *Int J Audiol* 44**,** 625-630.

R-Core-Team (2020). "R: A language and environment for statistical computing". R
Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-
project.org/).

Ratiu, I., Hout, M.C., Walenchok, S.C., Azuma, T., and Goldinger, S.D. (2017). Comparing
visual search and eye movements in bilinguals and monolinguals. *Attention,*
Perception, & Psychophysics 79**,** 1695-1725.

Reetzke, R., Xie, Z., Llanos, F., and Chandrasekaran, B. (2018). Tracing the trajectory of
sensory plasticity across different stages of speech learning in adulthood. *Current*
Biology 28**,** 1419-1427.e1414.

Ribarić, K., Prevec, T.S., and Kozina, V. (1984). Frequency-following response evoked by
acoustic stimuli in normal and profoundly deaf subjects. *Audiology* 23**,** 388-400.

Rosengren, S.M., Colebatch, J.G., Young, A.S., Govender, S., and Welgampola, M.S. (2019).
Vestibular evoked myogenic potentials in practice: Methods, pitfalls and clinical
applications. *Clin Neurophysiol Pract* 4**,** 47-68.

Russo, N.M., Skoe, E., Trommer, B., Nicol, T., Zecker, S., Bradlow, A., and Kraus, N. (2008).
Deficient brainstem encoding of pitch in children with autism spectrum disorders.
Clinical Neurophysiology 119**,** 1720-1731.

Sall, J. (1990). Leverage Plots for General Linear Hypotheses. *The American Statistician* 44**,**
308-315.

Schneider, P., Scherg, M., Dosch, H.G., Specht, H.J., Gutschalk, A., and Rupp, A. (2002).
Morphology of Heschl's gyrus reflects enhanced activation in the auditory cortex of
musicians. *Nature Neuroscience* 5**,** 688-694.

Skoe, E., and Kraus, N. (2010). Auditory brain stem response to complex sounds: A tutorial.
Ear and Hearing 31**,** 302-324.

Skoe, E., and Kraus, N. (2012). A little goes a long way: How the adult brain is shaped by
musical training in childhood. *Journal of Neuroscience* 32**,** 11507-11510.

Smith, J.C., Marsh, J.T., and Brown, W.S. (1975). Far-field recorded frequency-following
responses: Evidence for the locus of brainstem sources. *Electroencephalography and*
Clinical Neurophysiology 39**,** 465-472.

Sobel, M.E. (1982). Asymptotic confidence intervals for indirect effects in structural equation
models. *Sociological Methodology* 13**,** 290-312.

Sohmer, H., and Pratt, H. (1977). Identification and separation of acoustic frequency following
responses (FFRs) in man. *Electroencephalography and Clinical Neurophysiology* 42**,**
493-500.

Sohmer, H., Pratt, H., and Kinarti, R. (1977). Sources of frequency-following responses (FFR)
in man. *Electroencephalography and Clinical Neurophysiology* 42**,** 656-664.

Strait, D.L., Parbery-Clark, A., Hittner, E., and Kraus, N. (2012). Musical training during early
childhood enhances the neural encoding of speech in noise. *Brain and Language* 123**,**
191-201.

Thornton, A.R.D. (1975). The use of post-auricular muscle responses. *The Journal of*
Laryngology & Otology 89**,** 997-1010.

Tierney, A.T., Krizman, J., and Kraus, N. (2015). Music training alters the course of
adolescent auditory development. *Proceedings of the National Academy of Sciences*
of the United States of America 112**,** 10062-10067.

Weiss, M.W., and Bidelman, G.M. (2015). Listening to the brainstem: Musicianship enhances
intelligibility of subcortical representations for speech. *Journal of Neuroscience* 35**,**
1687-1691.

White-Schwoch, T., Anderson, S., Krizman, J., Nicol, T., and Kraus, N. (2019). Case studies
in neuroscience: Subcortical origins of the frequency-following response. *Journal of*
Neurophysiology 122**,** 844-848.

White-Schwoch, T., Woodruff Carr, K., Thompson, E.C., Anderson, S., Nicol, T., Bradlow,
A.R., Zecker, S.G., and Kraus, N. (2015). Auditory processing in noise: A preschool
biomarker for literacy. *PLOS Biology* 13**,** e1002196.

Wong, P.C., Skoe, E., Russo, N.M., Dees, T., and Kraus, N. (2007). Musical experience
shapes human brainstem encoding of linguistic pitch patterns. *Nature Neuroscience*
10**,** 420-422.

Yoshie, N., and Okudaira, T. (1969). Myogenic evoked potential responses to clicks in man.
Acta Otolaryngol Suppl 252**,** 89-103.

Yuval-Greenberg, S., Tomer, O., Keren, A.S., Nelken, I., and Deouell1, L.Y. (2008). Transient
induced gamma-band response in eeg as a manifestation of miniature saccades.
Neuron 58**,** 429-441.

Zhang, J.D., Susino, M., Mcpherson, G.E., and Schubert, E. (2020). The definition of a
musician in music psychology: A literature review and the six-year rule. *Psychology of*
Music 48**,** 389-409.

Zhao, T.C., and Kuhl, P.K. (2018). Linguistic effect on speech perception observed at the
brainstem. *Proceedings of the National Academy of Sciences of the United States of*
America 115**,** 8716-8721.
798