Molecular evaluation of *Conogethes punctiferalis* (Guenée, 1854): Species status and intraspecific divergence

Jin An[1,2], Ya-Lin Yao[1], Ping Gao[3], Minghua Xiu[1,2], Cheng-Min Shi[1,2,*]

[1]College of Plant Protection, Hebei Agricultural University, Baoding 071000

[2]State Key Laboratory of North China Crop Improvement and Regulation, Baoding 071000

[3]College of Plant Protection, Shenyang Agricultural University, Shenyang 110866

[*]Correspondence; shichengmin@hebau.edu.cn

Abstract: Species are not only the fundamental units of taxonomy but also the basic units of pest management. Insects of *Conogethes* are important agricultural and forestry pests. However, species boundaries within *Conogethes* often appear obscure. In the present study, we re-evaluated the species status of *Conogethes* by applying three species delimitation approaches based on the mitochondrial DNA sequences, with particular emphasis on the yellow peach moth *C. punctiferalis* (Guenée, 1854). We first optimized species delimitation and inter-species genetic divergence threshold using a DNA barcoding dataset. Our results revealed that several nominal species of *Conogethes* species harbored deeply diverged mitochondrial lineages which were recognized as independent species by the species delimitation methods. The p-distance between the delimited putative species ranged from 0.0159 to 0.1321 with a mean of 0.0841. Then we refined the species status of *C. punctiferalis* using the smallest interspecific distance threshold based on a geographically comprehensive population-scale dataset. This procedure narrowed the species concept of *C. punctiferalis* to a genetically coherent unit. Further investigation of its intraspecific divergence in the geographic context revealed that the refined *C. punctiferalis* was still widely distributed with the same or highly similar mitochondrial haplotypes occurring across South and East Asia.

Key words: species delimitation, DNA barcoding, genetic distances, mitochondrial DNA, *Conogethes punctiferalis*

1 Introduction

Species are not only the fundamental unit of taxonomy and biodiversity inventory but also the basic unit of management for those that pose threats to agriculture, forestry, and public health. For pest insects, how species are defined and delimited has profound implications for pest management practices. A basic, but often unstated, assumption of modern pest management is that members of the same target species share similar life cycles, host range, habitat preferences, and other ecological traits that confer similar responses to management tactics. As it has been widely testified that distinct species react differently to control methods, the presence of cryptic pest species (two or more distinct species that are erroneously classified and hidden under one species name) can lead to failure of control efforts for no apparent reason (Walter, 2005). There are many examples where the misidentification of a cryptic species complex as one species has thwarted control efforts. However, most, if not all, agricultural pest insects were initially recognized only from morphological perspectives. Now it is well acknowledged that morphological species do not always present a natural unit that is reproductively isolated from other such units or has a coherent evolutionary history. Cryptic species are common in generalist pest insects, for instance, in the neotropical skipper butterfly *Astraptes fulgerator* (Hebert et al., 2004), and the whitefly *Bemisia tabaci* (Vyskočilová et al., 2018). The consequence of cryptic species on pest management is particularly profound for species that have developed insecticide resistance which appears common in agroecosystems (Horowitz et al., 2020).

The yellow peach moth *Conogethes punctiferalis* (Guenée, 1854), belonging to the specious family Crambidae (Insecta: Lepidoptera), is an important agricultural pest. It was first described as *Astura punctiferalis*, then revised as *Dichocrocis punctiferalis* before getting its current name. It is widespread in subtropical and tropical Asia and Oceania, and of particular concern in agriculture in China, India, Indonesia, Korea, Japan, Malaysia, New Guinea, Vietnam and Australia (EPPO, 2023). This species is highly polyphagous feeding on more than 100 host plants from >30 different families, many of which are economically important crops and fruits (Lu et al., 2010; Rojas-Sandoval, 2022). Larvae of *C. punctiferalis* have long been considered a serious pest of peach, apple, pear, chestnut and other fruits across its range, and recently emerged as a devastating pest of maize and sunflower in north China (Wang et al., 2006). The wide geographic distribution combined with extreme polyphagy implies cryptic divergence within the species.

Fueled by the increasing availability of DNA sequences, the discoveries of cryptic species have increased exponentially over the past three decades (Bickford et al., 2007). Indeed, two ecotypes of *C. punctiferalis* with different host plant preference observed in Japan was recognized as two distinct species (Honda & Mitsuhashi, 1989). An oligophagous Pinaceae-feeding type (PFT) mainly feeding on pines was recognized as *C. pinicolalis*; a still polyphagous fruit-feeding type (FFT) was re-described as *C. punctiferalis* (Inoue & Yamanaka, 2006). The validity of the recognized species was recently supported by divergence in morphology, ecology and genetics in terms of both

mitochondrial DNA and nuclear DNA (Jeong et al., 2021). Substantial genetic divergences were detected between the two species with an average of 5.46% for the mitochondrial COI genes and 2.10% for the nuclear EF1α sequences (Jeong et al., 2021). Both of these two species, referred to here as *C. punctiferalis s. str.* and *C. pinicolalis*, were identified in China based on mitochondrial DNA sequences (Wang et al., 2014). Similarly, *C. punctiferalis* in India feeding on castor and cardamom represented two different lineages (Chakravarthy et al., 1991). The cardamom population was lately recognized as a distinct species *C. sahyadriensis* which differed from *C. punctiferalis* by genetic distances of no less than 6.00% (Shashank et al., 2018). Thus, it is clear that *C. punctiferalis* represents a species complex including multiple species of very similar morphology, variable ecotypes, and overlapping host ranges (Armstrong, 2010).

Although several cryptic species with restricted host plants and narrow geographic range have been recognized in the traditional morphological species *C. punctiferalis*, its remaining members (*C. punctiferalis s. str.*) still appear extremely host-rich and widespread. To better inform the management of this pest insect in the future, a global re-evaluation of its species status and intraspecific genetic variation is warranted. Here we performed such a molecular re-evaluation based on mitochondrial DNA sequence data. We first delimitated species by applying three statistical species delimitation methods and optimized species recognition threshold. Using the threshold obtained, we re-evaluated the species status and intraspecific genetic variation of *C. punctiferalis s. str.* based on a global dataset.

2 Materials and methods

2.1 Data curation

Two DNA datasets of the mitochondrial COI gene (mtCOI) were compiled in the present study. The first dataset, referred to as ConCOI, was used to examine the performance of species delimitation methods and thereby optimize the threshold of genetic distance for species recognition. The dataset included mtCOI sequences for all species of *Conogethes* currently recognized. We derived the ConCOI dataset by querying the GenBank through BLASTn with mtCOI sequences of the putative species recognized by the taxonomic treatment of Shashank et al. (2018). We filtered the hits from each of blast search with identity of 95%~100% and coverage of 98%~100%. Sequences were aligned using MUSCLE (Edgar, 2004). The aligned ConCOI dataset included 297 mtCOI sequences of *Conogethes* species and one sequence (HQ952111) for *Oxycanus niphadias* as an outgroup. After trimming, the 297 mtCOI sequences of *Conogethes* species harbored 110 unique sequences (haplotypes) of 566-bp in length.

The second dataset, referred to as CpunCOI, was used to re-evaluation species status and intraspecific genetic variation. This dataset was compiled by downloading all available mtCOI with taxon label of *Conogethes punctiferalis* from GenBank and BOLD databases. After filtering using the within-species genetic distance of *C. punctiferalis* inferred from the ConCOI dataset (see below),

the CpunCOI dataset included 420 mtCOI sequences of 233 bp long after trimming to the same length. We preferred to include more individuals over selecting longer sequences as it has been demonstrated that there were no significant differences in performance for species-level identification between full-length and mini-barcodes as long as they are > 200 bp in length (Yeo et al., 2020).

2.2 Tests of species limits and optimization of genetic threshold

We tested species limits in *Conogethes* using the ConCOI haplotype dataset by employing both the genetic clustering method and the phylogenetic tree-based approaches. Our contention is that robust species delimitations will be consistently supported by signals from multiple species delimitation approaches. For the genetic clustering method, we used the assembling species by automatic partitioning (ASAP) method which employs pairwise genetic distances for hierarchical clustering (Puillandre et al., 2021). The genetic distances were computed using K80 (a.k.a. K2P) substitution model with the transition vs. transversion ratio (ts/tv = 3.74) being estimated from the data using the maximum likelihood method. The best species partition was identified according to the ASAP score. Following the recommendation of Puillandre et al. (2021), we considered the partitions with the top two best ASAP scores.

Two tree-bases methods were applied. First, potential species were delimited using the generalized mixed Yule Coalescent (GMYC) method (Fujisawa & Barraclough, 2013) based on the maximum clade credibility (MCC) tree inferred using BEAST v.1,10.4 (Suchard et al., 2018). We only used the single threshold model because we expected the result to be transferable among datasets. BEAST was run with TN93+ G4 substitution model and strict molecular model with a Yule process as speciation prior. Markov chain Monte Carlo (MCMC) searches were run for 2×10^7 generations with trees and parameters being sampled every 2000 generations. We used Tracer 1.7 (Rambaut et al. 2018) to check the stationarity and the convergence of the MCMC chains. MCC tree was summarized from the sampled trees using TreeAnnotator 1.10 (Suchard et al. 2018), with the first 10% of samples being discarded as burn-in. Second, species were delimited using the Poisson tree process (PTP) model (Zhang et al., 2013) based on the maximum likelihood tree inferred using IQ-TREE v. 2.3 (Nguyen et al., 2015) and applying a single threshold. We used both the maximum likelihood and Bayesian implementation of the PTP model (mPTP and bPTP, respectively). The analysis was run for 100000 generations with a thinning parameter of 100 and burn-in of 0.1.

Pairwise genetic distances (*p*-distance and K2P-distance) among afore-delimited species were calculated using MEGA 11(Tamura et al., 2021). The smallest interspecific distance was used as a putative genetic threshold for further demarcation of species boundary of *Conogethes* species. It has been reported that the use of the mean distance exaggerates the size of interspecific divergences and leads to misidentification (Meier et al., 2008)

2.3 Analyses of intraspecific genetic variations for *C. punctiferalis s. str.*

For the clade/cluster or delimited species corresponding to *C. punctiferalis* (sensu Shashank et al. 2018), the pairwise genetic distances (*p*-distance) among mtCOI haplotypes were calculated using MEGA 11(Tamura et al., 2021). To further evaluate intraspecific genetic variation, we used these haplotypes as queries and recursively blasted the GenBank with the smallest interspecific distance found among the species delimited in the present study as a filtering threshold. After removing repeated sequences, aligning with above haplotypes and trimming to the same length, these sequences made up of the CpunCOI dataset. We determined the unique haplotypes and calculated population genetic summary statistics using DnaSP version 6.12.03 (Rozas et al., 2017). Median-joining networks (Bandelt et al., 1999) were constructed using NETWORK 10.2.0.0 (https://www.fluxus-engineering.com/) to illustrate intraspecies relationships, and geographic distribution of unique haplotypes.

3 Results

3.1 Delimitation of *Conogethes* species

The ConCOI dataset included 297 mtCOI sequences (566 bp) which were collapsed into 110 unique haplotypes. From this dataset, the number of species delimitation by three different methods ranged from 15 to 23 (Figure 1). The best partition found by ASAP clustered the 110 haplotypes of *Conogethes* into 19 subsets (species) with an ASAP score of 4.00 and a distance threshold of 0.0143 (ASAP-1 in Figure 1). The second-best partition (ASAP score = 4.50) clustered the *Conogethes* haplotypes into 14 subsets with a distance threshold of 0.0343 (ASAP-2 in Figure 1). The maximum likelihood partition found by the maximum likelihood search of the PTP model (mPTP) delimited 20 species. The most supported partition found by Bayesian PTP search (bPTP) delimited 21 species. One species (ANIC631-06, HQ953112) delimited by mPTP was split into two independent species. We note that these two sequences were poorly differentiated in the Bayesian MCC tree (Figure 1). Applying the single threshold, GMYC delimited 23 species (confidence interval: 18~27). The species delimitation model was significantly better than the null model (likelihood ratio: 26.60, result of LR test: 1.67e-06). The consensus species delimitation among the three methods recognized 19 species (Figure 1). All of these species formed strongly supported monophyletic clades. Except for *C. pinicolalis*, all other 18 species included identical or highly similar mtCOI sequences to those used by recent taxonomic treatment by Shashank et al. (2018), so we named each species following Shashank et al. (2018) to make it easy for cross-references.

The genetic distances calculated from mtCOI sequences of the 19 recognized consensus species are summarized in Table 1. The interspecific p-distances ranged from 0.0159 to 0.1321, with a mean of 0.0841. The corresponding K2P distances were 0.0161 to 0.1455, and 0.900, respectively. The maximum within species genetic distances were 0.0066 and 0.0067 for p-distances and K2P

distances, respectively.

3.2 Intraspecific genetic divergence of *Conogethes punctiferalis*

The p-distances between 33 mtCOI haplotypes (566 bp) of *C. punctiferalis* from ConCOI

dataset ranged from 0.0018 to 0.0159 with an average distance of 0.0066 (Table 2). To further

evaluate within species genetic divergence of *C. punctiferalis*, we filtered the CpunCOI dataset with

the minimum inter-species genetic distance found among the delimited species (0.0159, *C. evaxalis*

1 vs. *C. evaxalis* 2; Table 1). The final CpunCOI dataset included 420 sequences of 233 bp long.

These 420 sequences harbored 39 haplotypes defined by 30 variable sites. The haplotype diversity

and nucleotide diversity were 0.6100 and 0.0053, respectively. The p-distances between the 39 short

haplotypes (233 bp) of *C. punctiferalis* ranged from 0.0043 to 0.0343 with an average of 0.0126

(Table 2).

The relationships among haplotypes are shown by the median-joining network in Figure 2. The

haplotype frequencies range from 1 to 257 (H02). Twenty-two haplotypes are singletons that

appeared only once in the dataset. The majority of haplotypes were connected to the dominant

haplotype H02 by one mutation step. The second most frequent haplotype (H14; frequency, 44) was

connected to H02 by three mutation steps and two intermediate haplotypes. The 39 haplotypes were

found in six countries (China, Japan, South Korea, India, Pakistan and Malaysia). Ten haplotypes

were shared at least by two counties. Haplotype H02 was the most widely spread one, occurring in

all six countries. Haplotype H14 was found in South Korea, China, India, Pakistan and Japan.

Haplotypes H04, H08, H11and H15 were shared by East Asian countries (China, Japan and South

Korea). Haplotype H03 and H18 were shared between East Asia (China and/or Japan and South

Korea) and South Asia (India). In summary, we found extremely wide geographic distributions of

the same or highly similar mtCOI haplotypes of *C. punctiferalis* across South and East Asia.

4 Discussion

4.1 Multiple divergent mitochondrial lineages challenge species recognition of *Conogethes*

Species are often treated as the basic unit in the management of pest insects and cryptic species

present a challenge to pest management practice (Walter, 2005). It has been long suggested that

some members of *Conogethes* represent species complexes including multiple morphologically

similar species (Armstrong, 2010). However, species boundaries for most members of *Conogethes*

remain to be systematically evaluated using a consistent standard. Here, we bridged this knowledge

gap by performing molecular species delimitation using a comprehensive dataset that included all

available mtCOI sequences of *Conogethes* species. Species delimitation using three different

approaches (5 methods) consistently revealed that several nominal species of *Conogethes* included

multiple highly diverged mitochondrial lineages which warrant independent species (Figure 1). For

instance, nominal species *C. punctiferalis* was divided into two distantly related monophyletic

clades (*C. punctiferalis* vs. *C.cf. punctiferalis*) which differed by 5.91% in mtCOI sequences (Table 1). Similarly, the nominal species *C. semifascialis* was divided into three deeply diverged lineages (*C. semifascialis* 1, *C. semifascialis* 2 and *C. cf. semifascialis*). Besides, several deeply diverged mtCOI lineages (*C.* sp. 1~4), potentially representing distinct species, await to be formally described. Unexpectedly, we found *C. punctiferalis* was more closely related to *C. semifascialis* 1 than to any species that were recently recognized in the *C. punctiferalis* species complex, i.e. *C. pinicolalis* and *C. sahyadriensis* (Figure 1). These findings together signified that DNA-based identification of *Conogethes* species is essential for precision pest management practice for this group.

Considering the consensus species delimitation reported here and species recognition by Shashank et al. (2018) as valid, the minimum inter-species genetic difference was as low as 1.59%, although the overall average of pairwise between-species genetic difference was more than 8% (Table 1). This observation reinforced the earlier finding that use of mean instead of the smallest interspecific distance exaggerates the threshold of the barcoding gap and leads to the over-lumping of species (Meier et al., 2008). We thus suggest the use of the smallest inter-species genetic distance (1.59%) as a putative threshold for molecular recognition of *Conogethes* pest species in management practice.

4.2 Redefine species concept of *Conogethes punctiferalis*

The two datasets (ConCOI and CpunCOI) compiled in the present study allowed us to refine the species boundary of *C. punctiferalis* from genetic and geographic perspectives. From a phylogenetic point of view, we redefined *C. punctiferalis* as a strongly supported monophyletic mitochondrial lineage which was most closely related to *C. semifascialis* 1 (Figure 1). The genetic distance between *C. punctiferalis* and its most close relative was 2.30% (p-distance, and 2.35% for K2P distance). Only the GMYC method delimited two sub-lineages within *C. punctiferalis* as independent species (Figure 1). However, these sub-lineages were not well supported in the phylogenetic analyses. Thus, we treated them as intraspecific variations. The species concept of *C. punctiferalis* redefined here was consistent with Shashank et al. (2018).

From a geographic perspective, the refined *C. punctiferalis* is still widely spread. It occurred in China, Japan, South Korea, India, Pakistan, and Malaysia, straddling the temperate, subtropical and tropical zones. The cross-regional sharing of the same mitochondrial haplotypes suggests that this species may have high dispersal capacity and that long-distance dispersal might have been rampant. Such possibility was also supported by the detection of *C. punctiferalis* in the autumn migrating insect troops across northern China by radar observatories (Feng et al., 2003). Therefore, we think *C. punctiferalis* we redefined here represents a coherent genetic unit which may withstand tests from other aspects.

4.3 Limitations and perspectives

We should note that we have delimitated species *Conogethes* species, and *C. punctiferalis* in
particular, only from a maternal (mitochondrial) perspective. Now, it is increasingly recognized that
mitochondrial DNA records the incomplete history of species (Ballard & Whitlock, 2004), and in
some cases, mitochondrial DNA can mislead species recognition (Rubinoff & Holland, 2005).
Despite the aforementioned and other obvious drawbacks of using mitochondrial DNA alone for
species delimitation, the success of using mitochondrial DNA in distinguishing species from a range
of taxa and revealing cryptic species is remarkable (Dasmahapatra & Mallet, 2006). Our results
therefore can serve as a working hypothesis for further tests. Future efforts should be made to
delimitate *Conogethes* species using nuclear DNA data and more robust methods that consider both
the speciation process between species and the coalescence process within species (Rannala & Yang
2020). As the advantages and limitations of DNA-based species identification become apparent in
pest management practice, it is clear that taxonomic approaches integrating DNA data,
morphological characteristics, ecological features, and damage symptoms will achieve optimal
efficiency in species identification.
264

265 **Funding**

266 This study was supported by a starting fund from Hebei Agricultural University and the State Key

267 Laboratory of North China Crop Improvement and Regulation (YJ2020028).

269 Table 1. Estimates of genetic divergence between the delimited species of *Conogethes*. The p-distances were shown above the diagonal and the K2P distances were shown

270 blow the diagonal. The numbers on the diagonal represented the within species K2P distances.

Species	Cpun	Csem1	Ccfsem	Cpin	Csah	Ccfpun	Csp1	Csem2	Cplu	Ccfhae	Cers	Ceva2	Csp2	Cdim	Chae	Csp4	Ceva1	Ctha	Csp3
Cpun	**0.0066**	0.0230	0.0365	0.0500	0.0518	0.0519	0.0549	0.0561	0.0591	0.0791	0.0815	0.0850	0.0859	0.0856	0.0894	0.0887	0.0931	0.0974	0.1078
Csem1	0.0235	**0.0024**	0.0318	0.0503	0.0496	0.0409	0.0526	0.0480	0.0570	0.0773	0.0699	0.0813	0.0808	0.0775	0.0855	0.0797	0.0866	0.0965	0.1005
Ccfsem	0.0375	0.0326	**n.c.**	0.0547	0.0624	0.0481	0.0601	0.0590	0.0548	0.0901	0.0789	0.0936	0.0972	0.0883	0.0954	0.0954	0.0919	0.1056	0.1113
Cpin	0.0522	0.0526	0.0572	**0.0060**	0.0615	0.0636	0.0592	0.0603	0.0561	0.0810	0.0664	0.0801	0.0849	0.0751	0.0863	0.0951	0.0872	0.1100	0.1025
Csah	0.0544	0.0521	0.0661	0.0651	**0.0012**	0.0691	0.0607	0.0710	0.0527	0.0942	0.0760	0.0989	0.0954	0.0819	0.0901	0.1013	0.1042	0.0991	0.1119
Ccfpun	0.0545	0.0424	0.0500	0.0671	0.0738	**0.0034**	0.0641	0.0408	0.0593	0.0852	0.0690	0.1019	0.0966	0.0834	0.0908	0.0905	0.0970	0.1042	0.1108
Csp1	0.0575	0.0550	0.0631	0.0621	0.0640	0.0677	**n.c.**	0.0643	0.0526	0.0954	0.0736	0.1025	0.0972	0.0813	0.0872	0.0954	0.1042	0.1038	0.1095
Csem2	0.0589	0.0500	0.0620	0.0633	0.0757	0.0424	0.0677	**0.0044**	0.0654	0.0905	0.0862	0.1011	0.1003	0.0939	0.0864	0.0952	0.1029	0.1080	0.1100
Cplu	0.0625	0.0603	0.0575	0.0591	0.0557	0.0626	0.0550	0.0693	**0.0025**	0.1014	0.0674	0.1009	0.1009	0.0800	0.0917	0.0972	0.0961	0.1127	0.1133
Ccfhae	0.0837	0.0816	0.0961	0.0858	0.1009	0.0905	0.1022	0.0966	0.1093	**n.c.**	0.0907	0.0813	0.0777	0.1007	0.0595	0.0883	0.0830	0.1038	0.1007
Cers	0.0871	0.0739	0.0842	0.0701	0.0809	0.0730	0.0780	0.0927	0.0712	0.0969	**0.0035**	0.0995	0.1013	0.0230	0.0801	0.0978	0.0942	0.1104	0.1037
Ceva2	0.0902	0.0861	0.1002	0.0847	0.1064	0.1099	0.1105	0.1089	0.1088	0.0861	0.1074	**0.0000**	0.0247	0.1042	0.0878	0.0813	0.0159	0.1140	0.0989
Csp2	0.0913	0.0856	0.1043	0.0903	0.1023	0.1037	0.1043	0.1078	0.1088	0.0821	0.1095	0.0253	**n.c.**	0.1060	0.0895	0.0901	0.0318	0.1184	0.1007
Cdim	0.0919	0.0827	0.0952	0.0799	0.0877	0.0896	0.0868	0.1018	0.0855	0.1086	0.0235	0.1131	0.1152	**n.c.**	0.0836	0.1007	0.1095	0.1184	0.1095
Chae	0.0952	0.0909	0.1022	0.0917	0.0961	0.0970	0.0927	0.0919	0.0979	0.0621	0.0849	0.0934	0.0955	0.0889	**0.0024**	0.0819	0.0919	0.1092	0.0948
Csp4	0.0945	0.0843	0.1023	0.1020	0.1093	0.0966	0.1023	0.1020	0.1044	0.0942	0.1053	0.0861	0.0962	0.1087	0.0869	**n.c.**	0.0866	0.1113	0.0989
Ceva1	0.0996	0.0921	0.0981	0.0929	0.1127	0.1041	0.1126	0.1109	0.1031	0.0881	0.1012	0.0161	0.0327	0.1195	0.0982	0.0922	**0.0000**	0.1170	0.1042
Ctha	0.1045	0.1034	0.1139	0.1191	0.1065	0.1124	0.1120	0.1168	0.1226	0.1118	0.1199	0.1238	0.1290	0.1296	0.1182	0.1207	0.1274	**0.0041**	0.1321
Csp3	0.1166	0.1080	0.1207	0.1104	0.1217	0.1202	0.1186	0.1192	0.1234	0.1083	0.1118	0.1063	0.1084	0.1188	0.1017	0.1065	0.1126	0.1455	**n.c.**

271 Species codes refer to Figure 1--- Cpun: *C. punctiferalis*; Ccfpun: *C. cf. punctiferalis*; Csem1: *C. semifascialis* 1; Csem2: *C. semifascialis* 2; Ccfsem: *C. cf. semifascialis*;

272 Cpin: *C. pinicolali*s; Csah: *C. sahyadriensi*s; Cplu: *C. pluto*; Chae: *C. haemactalis*; Ccfhae: *C. cf. haemactalis*; Cers: *C. ersealis*; Ceva1: *C. evaxalis* 1; Ceva2: *C. evaxalis* 2;

273 Cdim: *C. diminutive*; Ctha: *C. tharsalea*; Csp1~4: *C.* sp. 1~4.

275 Table 2. Estimates of genetic divergence between mtCOI haplotypes of *C. punctiferalis*. n: sample
276 size; H: number of haplotypes.

Datasets	Length	n	H	p-distance (mean [min., max.])	K2P-distance (mean [min., max.])
ConCOI	566 bp	147	33	0.0066 [0.0018, 0.0159]	0.0067 [0.0018, 0.0161]
CpuCOI	233 bp	420	39	0.0126 [0.0043, 0.0343]	0.0128 [0.0043, 0.0352]

References

References

Armstrong, K. 2010. DNA barcoding: a new module in New Zealand's plant biosecurity diagnostic
toolbox. *EPPO Bulletin*, 40(1): 91-100.

Ballard, J.W.O., Whitlock, M.C. 2004. The incomplete natural history of mitochondria. *Molecular*
Ecology, 13(4): 729-744.

Bandelt, H.J., Forster, P., Röhl, A. 1999. Median-joining networks for inferring intraspecific
phylogenies. *Molecular Biology and Evolution*, 16(1): 37-48.

Bickford, D., Lohman, D.J., Sodhi, N.S., Ng, P.K.L., Meier, R., Winker, K., Ingram, K.K., Das, I.
2007. Cryptic species as a window on diversity and conservation. *Trends in Ecology &*
Evolution, 22(3): 148-155.

Chakravarthy, A.K., Honda, H., Thyagaraj, N.E. 1991. Comparison of containers for larval rearing
in stalk and fruit feeding type of *Conogethes punctiferalis* (Guen.) (Lepidoptera:
Pyralidae). *International Symposium on Plantation Crops*: 9, 127-131.

Dasmahapatra, K.K., Mallet, J. 2006. Taxonomy: DNA barcodes: recent successes and future
prospects. *Heredity*, 97(4): 254-255.

Edgar, R.C. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput.
Nucleic Acids Research, 32(5): 1792-1797.

EPPO. 2023. *Dichocrocis punctiferalis* (DICHPU). https://gd.eppo.int/taxon/DICHPU

Feng, H.Q., Wu, K.M., Cheng, D.F., Guo, Y.Y. 2003. Radar observations of the autumn migration
of the beet armyworm *Spodoptera exigua* (Lepidoptera: Noctuidae) and other moths in
northern China. *Bulletin of Entomological Research*, 93(2): 115-124.

Fujisawa, T., Barraclough, T.G. 2013. Delimiting species using single-locus data and the generalized
mixed Yule coalescent approach: A revised method and evaluation on simulated data sets.
Systematic Biology, 62(5): 707-724.

Hebert, P.D.N., Penton, E.H., Burns, J.M., Janzen, D.H., Hallwachs, W. 2004. Ten species in one:
DNA barcoding reveals cryptic species in the neotropical skipper butterfly *Astraptes*
fulgerator. *Proceedings of the National Academy of Sciences*, 101(41): 14812-14817.

Honda, H., Mitsuhashi, W. 1989. Morphological and morphometrical differences between the fruit-
and Pinaceae-feeding type of yellow peach moth, *Conogethes punctiferalis* (Guenee)
(Lepidoptera : Pyralidae). *Applied Entomology and Zoology*, 24: 1-10.

Horowitz, A. R., Ghanim, M., Roditakis, E., Nauen, R., Ishaaya, I. 2020. Insecticide resistance and
its management in *Bemisia tabaci* species. *Journal of Pest Science*, 93(3): 893-910.

Inoue, H., Yamanaka, H. 2006. Redescription of *Conogethes punctiferalis* (Guenée, 1854) and
descriptions of two new closely allied species from Eastern Palaearctic and Oriental
Regions (Pyralidae: Pyraustinae). *Tinea*, 19(2): 80-91.

Jeong, N.-R., Kim, M.-J., Kim, S.-S., Choi, S.-W., Kim, I.-S. 2021. Morphological, ecological, and
molecular divergence of *Conogethes pinicolalis* from *C. punctiferalis* (Lepidoptera:
Crambidae). *Insects*, 12(5): 455.

Lu, J., Wang, Z., He, K., Liu, Y. 2010. Research history, progresses and prospects in the yellow
peach moth, *Conogethes punctiferalis*. *Plant Protection*, 36(2): 31-38.

Meier, R., Zhang, G., Ali, F. 2008. The use of mean instead of smallest interspecific distances
exaggerates the size of the "barcoding gap" and leads to misidentification. *Systematic*
Biology, 57(5): 809-813.

Nguyen, L.-T., Schmidt, H.A., von Haeseler, A., Minh, B.Q. 2015. IQ-TREE: A fast and effective

stochastic algorithm for estimating maximum likelihood phylogenies. *Molecular Biology*
and Evolutions, 32: 268-274.

Puillandre, N., Brouillet, S., Achaz, G. 2021. ASAP: assemble species by automatic partitioning.
Molecular Ecology Resources, 21(2): 609-620.

Rambaut, A., Drummond, A.J., Xie, D., Baele, G., Suchard, M.A. 2018. Posterior summarization in
Bayesian phylogenetics using Tracer 1.7. *Systematic Biology*, 67:901–904.

Rannala, B., Yang, Z. 2020. Species Delimitation. In Scornavacca, C., Delsuc, F., Galtier, N., editors,
Phylogenetics in the Genomic Era, Chapter 5.5, pp. 5.5:1-18.

Rojas-Sandoval, J. 2022. *Conogethes punctiferalis* (yellow peach moth) (Vol. CABI Compendium).
CABI.

Rozas, J., Ferrer-Mata, A., Sánchez-DelBarrio, J.C., Guirao-Rico. S., Librado, P., Ramos-Onsins,
S.E., Sánchez-Gracia, A. 2017. DnaSP 6: DNA sequence polymorphism analysis of large
data sets. *Molecular Biology and Evolution* 34:3299-3302.

Rubinoff, D., Holland, B.S. 2005. Between two extremes: Mitochondrial DNA is neither the panacea
nor the nemesis of phylogenetic and taxonomic inference. *Systematic Biology*, 54(6), 952-
961.

Shashank, P.R., Kammar, V., Mally, R., Chakravarthy, A.K. 2018. A new Indian species of shoot
and capsule borer of the genus *Conogethes* (Lepidoptera: Crambidae), feeding on
cardamom. *Zootaxa*, 4374(2): 215-234.

Suchard, M.A., Lemey, P., Baele, G., Ayres, D.L., Drummond, A.J., Rambaut, A. 2018. Bayesian
phylogenetic and phylodynamic data integration using BEAST 1.10. *Virus Evolutio*n, 4(1):
vey016.

Tamura, K., Stecher, G., Kumar, S. 2021. MEGA11: Molecular evolutionary genetics analysis
version 11. *Molecular Biology and Evolution*, 38(7): 3022-3027.

Vyskočilová, S., Tay, W.T., van Brunschot, S., Seal, S., Colvin, J. 2018. An integrative approach to
discovering cryptic species within the *Bemisia tabaci* whitefly species complex. *Scientific*
Reports, 8(1): 10886.

Walter, G.H. 2005. Insect Pest Management and Ecological Research. Cambridge University Press.

Wang, J., Zhang, T.T., Wang , Z.Y., He, K.L., Liu, Y., Li, J. 2014. Molecular taxonomy of
Conogethes punctiferalis and *Conogethes pinicolalis* (Lepidoptera: Crambidae) based on
mitochondrial DNA sequences. *Journal of Integrative Agriculture*, 13(9): 1982-1989.

Wang, Z., He, K.S., Ma, S. 2006. Causes deterioration *Conogethes punctiferalis* damage on maize
and strategies for its control. *Plant Protection*, 32(2): 67-69.

Yeo, D., Srivathsan, A., Meier, R. 2020. Longer is not always better: Optimizing barcode length for
large-scale species discovery and identification. *Systematic Biology*, 69(5): 999-1015.

Zhang, J., Kapli, P., Pavlidis, P., Stamatakis, A. 2013. A general species delimitation method with
applications to phylogenetic placements. *Bioinformatics*, 29(22): 2869-2876.

Figure 1. Species delimitation and phylogenetic relationship of *Conogethes* mitochondrial haplotypes. The ultrametric phylogenetic tree was inferred using BEAST with a strict molecular clock model and a speciation prior of Yule process. Strongly supported nodes with posteriors ≥ 0.95 and maximum likelihood bootstrapping support values ≥ 70 were indicated by black dots. The bars on the right depict the putative species delimited by different methods. Species names identified following Shashank et al. (2018).

367
368 Figure 2. Median-joining network of mitochondrial haplotypes of refined *Conogethes punctiferalis*.
369 Each node represents a unique haplotype identified from 420 mtCOI sequences. Node size is
370 proportional to the frequency of relevant haplotypes, and branch length is proportional to the
371 mutations that occurred between haplotypes. Color depicts the geographic origins of the harboring
372 mtCOI sequences.