TITLE

Phenotypic complexities of rare heterozygous neurexin-1 deletions

AUTHORS

Michael B. Fernando[1,2,4,8], Yu Fan[3]*, Yanchun Zhang[3]*, Alex Tokolyi[5], Aleta N. Murphy[1,2,4], Sarah

Kammourh[3], P.J. Michael Deans[8], Sadaf Ghorbani[7,8], Ryan Onatzevitch[2], Adriana Pero[3],

Christopher Padilla[3], Sarah Williams[1,2,4], Erin K. Flaherty[1,2,4], Iya A. Prytkova[1,2,4], Lei Cao[3] , David

A. Knowles[5,6,], Gang Fang[3#], Paul A. Slesinger,[1,2#] & Kristen J. Brennand[2-4,8#]

AFFILIATIONS

[1]Graduate School of Biomedical Science, Icahn School of Medicine at Mount Sinai, New York,
NY 10029

[2]Nash Family Department of Neuroscience, Icahn School of Medicine at Mount Sinai, New York,
NY 10029

[3]Department of Genetics and Genomics, Icahn School of Medicine at Mount Sinai, New York, NY
10029

[4]Friedman Brain Institute, Black Family Stem Cell Institute, Pamela Sklar Division of Psychiatric
Genomics, Icahn School of Medicine at Mount Sinai, New York, NY 10029

[5]New York Genome Center, New York, NY, 10013

[6]Departments of Computer Science, Systems Biology, and Data Science Institute, Columbia
University, New York, NY, USA, 10027

[7]Haukeland University Hospital, Bergen, Norway

[8]Department of Psychiatry, Yale School of Medicine, New Haven, CT, 06520

*These authors contributed equally

#Correspondence: kristen.brennand@yale.edu, paul.slesinger@mssm.edu and
gang.fang@mssm.edu

KEYWORDS

Human induced pluripotent stem cells; *NRXN1*; alternative splicing; glutamatergic neurons;

GABAergic neurons; genomics; neuropsychiatric disorder; disease modeling; precision medicine

ABSTRACT

Given the large number of genes significantly associated with risk for neuropsychiatric disorders,

a critical unanswered question is the extent to which diverse mutations --sometimes impacting

the same gene-- will require tailored therapeutic strategies. Here we consider this in the context

of rare neuropsychiatric disorder-associated copy number variants (2p16.3) resulting in

heterozygous deletions in *NRXN1*, a pre-synaptic cell adhesion protein that serves as a critical

synaptic organizer in the brain. Complex patterns of *NRXN1* alternative splicing are fundamental

to establishing diverse neurocircuitry, vary between the cell types of the brain, and are

differentially impacted by unique (non-recurrent) deletions. We contrast the cell-type-specific

impact of patient-specific mutations in *NRXN1* using human induced pluripotent stem cells, finding

that perturbations in *NRXN1* splicing result in divergent cell-type-specific synaptic outcomes. Via

distinct loss-of-function (LOF) and gain-of-function (GOF) mechanisms, *NRXN1*$^{+/-}$ deletions cause

decreased synaptic activity in glutamatergic neurons, yet increased synaptic activity in GABAergic

neurons. Reciprocal isogenic manipulations causally demonstrate that aberrant splicing drives

these changes in synaptic activity. For *NRXN1* deletions, and perhaps more broadly, precision

medicine will require stratifying patients based on whether their gene mutations act through LOF

or GOF mechanisms, in order to achieve individualized restoration of *NRXN1* isoform repertoires

by increasing wildtype, or ablating mutant isoforms. Given the increasing number of mutations

predicted to engender both LOF and GOF mechanisms in brain disorders, our findings add

nuance to future considerations of precision medicine.

MAIN

Neurexins are pre-synaptic cell adhesion proteins that act as synaptic organizers[1]. There are

three neurexin genes (*NRXN1*, *NRXN2*, and *NRXN3*) in mammals and each is highly alternatively

spliced to produce hundreds of isoforms, primarily categorized into long alpha and shorter beta

isoforms[2]. The complex alternative splicing of neurexins expands protein-protein interaction

capabilities[3], allowing neurexins to interact with diverse post-synaptic ligands to establish and

maintain neurotransmission. *NRXN1α* splice variants are specific to brain regions[4] and between

cell types[5], but the cell-type-specific functional impact of individual isoforms remains unclear.

Although rare in the population, large copy number variations (deletions or duplications) at the

NRXN1 locus 2p16.3, particularly those deleting exonic regions, are highly penetrant, pleiotropic,

and are strongly associated with several neuropsychiatric diseases, including schizophrenia

(odds ratio 14.4[6]), autism spectrum disorder (odds ratio 14.9[7]), epilepsy (odds ratio 9.91[8]),

intellectual disability (odds ratio 7.47[9]) and Tourette's syndrome (odds ratio 20.3[10]). Deletions in

NRXN1 are non-recurrent (that is, they vary in size and location), making it difficult to determine

the molecular mechanisms underlying their diverse clinical symptoms (e.g., diagnosis, severity,

prognosis, and age-of-onset). In rodent studies, gene knockouts (KO) of *NRXNs* are sufficient to

produce an array of excitatory and inhibitory synaptic phenotypes[1]. However, heterozygous

deletions yield only modest behavioral and physiological changes *in vivo*[11]. By contrast, *in vitro*

studies of engineered heterozygous *NRXN1*$^{+/-}$ human neurons reveal robust changes in excitatory

neurotransmission that are not recapitulated in matched *NRXN1*$^{+/-}$ mouse neurons[12,13], and

studies that utilize patient derived *NRXN1*$^{+/-}$ neurons have yet to deconvolute complex phenotypes

in a cell-type and genotype specific manner[12,14]. Altogether, neurexins possess unique human

neurobiology, and therefore, the impact of distinct patient-specific *NRXN1*$^{+/-}$ must be specifically

evaluated in human models with an important consideration of how unique *NRXN1*$^{+/-}$ deletions

impact splicing patterns and neuronal function across distinct cell-types.

Human induced pluripotent stem cell (hiPSC)-derived neurons provide an ideal platform to study

NRXN1α alternative splicing. Previously we established that hiPSC-derived forebrain cultures,

comprised of a mixture of glutamatergic and GABAergic neurons with astroglia, recapitulate the

diversity of *NRXN1α* alternative splicing observed in the human brain, cataloguing 123 high-

confidence *NRXN1α* isoforms[15]. Furthermore, using patient-derived *NRXN1*$^{+/-}$ hiPSCs with unique

5'- or 3'-deletions in the gene, we uncovered wide-scale reduction in wildtype *NRXN1α* isoform

levels and, robust expression of dozens of novel isoforms from the 3'-deletion allele only[15].

Overexpression of individual wildtype isoforms (WT) ameliorated reduced neuronal activity in

patient-derived *NRXN1*[+/-] hiPSC-neurons in a genotype-dependent manner, whereas mutant

isoform (MT) expression decreased neuronal activity levels in control hiPSC-neurons[15]. We

therefore hypothesized that 5'-deletions of the promoter region represent classical loss-of-

function (LOF), while robust expression of novel 3'-specific MT isoforms confer a gain-of-function

(GOF) effect that cannot be rescued by overexpression of WT isoforms. Although *NRXN1* splicing

varies between the cell types of the brain, the impact of non-recurrent *NRXN1*[+/-] deletions on cell-

type-specific splicing patterns and synaptic function remains untested.

Neurexin signaling impacts both glutamatergic and GABAergic synapse properties[3,16], suggesting

that neurexins may regulate excitatory and inhibitory balance, which is strongly implicated across

neuropsychiatric disorders[17]. Given its multifaceted roles, the lack of mechanistic understanding

of how aberrant *NRXN1* splicing impacts neuronal physiology in a cell-type *and* genotype-

dependent manner presents a significant challenge for therapeutic targeting. To discover the

disease mechanisms that underpin *NRXN1*[+/-] deletions, we compared excitatory and inhibitory

neurons, across LOF and GOF deletions, to specifically evaluate cell-autonomous phenotypes

arising from distinct *NRXN1*[+/-] deletions. We identified points of phenotypic divergence across

glutamatergic and GABAergic neurons, which were independently validated in isogenic

experiments, thereby establishing causal relationships between aberrant splicing and synaptic

dysfunction. Finally, we evaluated novel therapeutic agents for *NRXN1*[+/-] deletions based on

LOF/GOF stratified patient mechanisms.

RESULTS

Patient-specific *NRXN1*[+/-] mutations produces differential *NRXN1* splicing patterns across

glutamatergic and GABAergic neurons

To examine the impact of *NRXN1*[+/-] deletions in two different neuronal cell types, we used

transcription-factor based lineage conversion of hiPSCs to generate excitatory glutamatergic

neurons or inhibitory GABAergic neurons from our previous cohort[14] (all available clinical and

experimental information reported in **Supplementary Table 1**). Two cases sharing a ~115-kb

deletion in the 5'-region of *NRXN1* (5'-Del), affecting the alpha-transcript promoter, represent the

LOF condition. Two additional cases sharing a ~136-kb deletion in the 3'-region of *NRXN1* (3'-

Del), impacting two alternative splice sites (SS4, SS5) across three exons (21-23), robustly

express unique MT isoforms from the affected allele and therefore represent the GOF condition[15].

For controls, we used passage-matched hiPSCs from four healthy sex-balanced subjects (**Fig.**

1a,b). Transient overexpression of *NGN2* in hiPSCs produced iGLUT neurons that were >95%
pure glutamatergic neurons, robustly expressed glutamatergic genes, released glutamate, and
produced spontaneous synaptic activity by day 21 in vitro[18,19]. On the other hand, transient
overexpression of *ASCL1* and *DLX2* yielded iGABA neurons that were >80% positive for
expressing GABA and GAD1/2 by day 35 in vitro, and possessed mature physiologic properties
of inhibitory neurons by day 42[20,21]. Immunostaining confirmed expression of neurotransmitter
transporters, vGLUT1 and vGAT for iGLUT and iGABA neurons, respectively (**Fig. 1c,h**). RNA-
sequencing (RNAseq) further validated iGLUT (DIV21) and iGABA (DIV35) neuronal induction in
all donors (**Fig. 1d,i**).

Despite cell-type-specific *NRXN1* isoform profiles reported in healthy brains[5] and neurons[15],
aberrant *NRXN1* alpha/beta exon expression patterns appeared similar between *NRXN1*[+/-] iGLUT
and iGABA neurons (**Fig. 1e,j**), whereby alpha exon expression (exons 2-18) decreased for 5'-
Del and increased for 3'-Del in both types of neurons (iGLUT 5'-Del p = 0.0031; 3'-Del, p = 0.0086,
1-way ANOVA, Dunnett's test; iGABA 5'-Del p = 0.0004; 3'-Del, p < 0.0001, 1-way ANOVA,
Dunnett's test). To investigate changes in alternative splicing, we applied LeafCutter[22], an
annotation free analysis to estimate differential splicing events visualized by splicegraphs
denoting change of percent-spliced-in (ΔPSI) ratios across the *NRXN1* gene at annotated splice
sites (SS1-6) and novel unannotated junctions (**Fig 1b**). Differential splicing analysis between
iGLUT and iGABA neurons produced a single significant cluster (out of nine identified), centered
around SS3 (**Extended Data Fig. 1a**, p = 0.01960, Bonferroni corrected), suggesting that the
majority of isoforms were conserved between the two cell-types, as previously reported[15].

Specifically, we detected both canonical and noncanonical splicing patterns in 5'-Del and 3'-Del
neurons, relative to controls. Splicing of *NRXN1* transcript was perturbed at two locations in 5'-
Del neurons. The first change in splicing occurred at the alternative transcription start site,
reducing splicing of beta-exon transcripts into alpha-exon transcripts (henceforth referred to as
β→α), as expected given the affected alpha promoter. We observed a significant reduction in 5'-
Del iGABA neurons (ΔPSI -0.187, p = 0.0146, Bonferroni corrected), and a more modest decrease
in iGLUT neurons (ΔPSI -0.086, p = 0.1821, Bonferroni corrected), consistent with a LOF. In 5'-
Del iGLUT neurons, we also detected differential splicing at SS4 inclusion (ΔPSI -0.046, p =
0.006, Bonferroni corrected) (**Fig. 1f,k**). For 3'-Del neurons, we observed a robust increase in a
mutant splice junction between exon 20-24 in iGLUT (ΔPSI 0.227, p = 1.09E-19) and iGABA
(ΔPSI 0.23, p = 6.93E-23, Bonferroni corrected) neurons, concurrent with reduced wildtype
splicing around SS4 and SS5 (**Fig. 1g,l**), consistent with a GOF. Interestingly, STAR-family RNA-

binding proteins that regulate *NRXN1* splicing at SS4[23–26] were also dysregulated across cell-

types and genotypes **Extended Data Fig**. **1b,f**). Taken together, these differential splicing

patterns support our hypothesis that 5'-Del and 3'-Del mutations confer LOF/GOF phenotypes,

either by reducing $\beta \rightarrow \alpha$ splicing or producing novel MT isoforms, respectively.

***NRXN1*[+/-] mutations broadly impact synaptic and neurodevelopmental pathways in induced**

and organoid-derived glutamatergic and GABAergic neurons

To unbiasedly evaluate the transcriptomic impact of *NRXN1*[+/-] deletions, and their unique splicing

patterns, we utilized differential expression analysis[27] in iGLUT and iGABA neurons, which

revealed overlap of FDR-corrected differentially expressed genes (DEGs) across cell

types/genotypes (e.g., *FAM66D, TTC34, GALNT9*), and represented gene sets related to synaptic

function (via SynapseGO[28]) (**Fig. 2a-d, j-m**). To test for pathway enrichment among top DEGs

(filtered by ±1.5 Log$_2$FC), we performed gene-ontology analyses via ClusterProfiler[29], against a

background of all expressed genes to avoid exclusion of biologically relevant low abundance

genes. 5'-Del neurons revealed a robust enrichment of terms related to neurotransmission and

synaptic function in iGLUT neurons, and ligand-gated anion channel activity in iGABA neurons

(**Supplementary Table 2**). 3'-Del neurons were enriched for terms related to neurodevelopment,

significant in iGABA neurons, but only nominally enriched in iGLUT neurons. DEGs were distinct

between genotypes (**Extended Data Fig**. **1c,d,g,h**), with different DEGs in hierarchical clustered

clades enriched for DNA-binding related GO terms (**Extended Data Fig. 1e,i**). Risk enrichments

for schizophrenia, bipolar disorder, and autism spectrum disorder[30,31] were most enriched in 5'-

Del iGLUT neurons and 3'-Del iGABA neurons (**Extended Data Fig. 2**).

To explore aberrant *NRXN1* splicing in a more complex neurodevelopmental system, we studied

the effect of 5'-Del and 3'-Del in organoids. We applied dorsal forebrain patterning to yield human

cortical organoids (hCOs) that resembled the pallium, or applied ventral forebrain patterning to

generate human subpallial organoids (hSOs) that resembled the subpallium (**Fig. 2e,n and**

Extended Data Fig. 3a-c,h-j)[32–34]. To stratify LOF and GOF genotypes, we confirmed exclusive

MT *NRXN1* isoforms in 3'-Del organoids (**Extended Data Fig. 3d-e, k-l**). We performed single

cell RNA-seq (n = 47,460 cells from hCOs, n= 35,563 cells from hSOs using 10x genomics) at 6

months, a timepoint with well characterized neural activity[33,35], and subsequently identified

clusters of cell-types within hCOs and hSOs (**Fig.2f,o**), without significant differences in cell

frequencies across pooled genotypes (quasibinomial regression model **Fig. 2g,p**). Differentially

expressed gene sets in hCO-glutamatergic and hSO-GABAergic clusters identified robust

enrichment of GO terms related to RNA splicing (from upregulated genes), and

neurodevelopment and synaptic function (from downregulated genes) (**Fig. 2h,i,q,r**). Taken

together, these results reinforce the hypothesis that perturbations in *NRXN1* splicing converge on

synaptic function[14,36].

Patient-specific alterations in spontaneous neural activity occur with minimal changes in

passive and excitable membrane properties.

To evaluate the functional consequence of *NRXN1*[+/-] deletions in iGLUT and iGABA neurons, we

next conducted a population-level analysis of spontaneous neuronal activity using a multi-

electrode array (MEA) and an examination of passive membrane properties using whole-cell

patch-clamp electrophysiology. Spontaneous network activity (weighted mean firing rate, wMFR)

in both 5'- and 3'-Del *NRXN1*[+/-] iGLUT neurons, which increased over time in a linear fashion (**Fig**

3a), was reduced by over 40% across two independent time points, at weeks post induction

(WPI)4 (**Fig. 3b** 5'-Del p = 0.0005; 3'-Del, p < 0.0001, 1-way ANOVA, Dunnett's test) and WPI6

(**Fig 3c** 5'-Del p = 0.0129, 3'-Del p = 0.0069 1-way ANOVA, Dunnett's test). The cell capacitance,

membrane resistance and resting membrane potentials did not significantly differ between 5'- and

3'-Del cases and control iGLUT neurons (**Fig. 3d**). To examine the intrinsic excitability, we

compared the input-output curve for induced firing and found slightly lower firing in 5'-Del iGLUT

neurons, but no difference in 3'-Del iGLUT neurons (**Fig. 3e**). The voltage-dependent sodium and

potassium current densities were similar across 5'-Del, 3'-Del and control iGLUT neurons

(**Extended Data Fig**. **4a**). Taken together, these results suggest that changes in passive or

intrinsic excitability membrane properties cannot fully explain the reduced firing observed on

MEAs for 5'-Del and 3'-Del in iGLUT neurons.

In parallel, we generated iGABA neurons with the same 5'-Del, 3'-Del, and control hiPSCs, as

above. We observed that immature *NRXN1*[+/-] iGABA neurons exhibited a robust ~2-fold increase

in population-wide wMFR activity (~WPI2) from both 5'-Del and 3'-Del cases (5'-Del p = 0.0029,

3'-Del p = 0.0015 1-way ANOVA, Dunnett's test) (**Fig. 3f,g**). Though unexpected for GABA

neurons, the finding is consistent with activation of ionotropic GABA receptors leading to

depolarization due to low KCC2 expression and high chloride levels[37]. Indeed, immature iGABA

neurons expressed higher (10-fold increase compared to WPI2 neurons) levels of *SLC12A5* (the

gene encoding *KCC2*) (**Extended Data Fig. 5a** p < 0.0001, 2-way ANOVA, Dunnett's test).

Furthermore, this transient hyperexcitability was pharmacologically inhibited by 10μM gabazine,

a selective GABA$_A$ antagonist (**Extended Data Fig. 5b,c**). In mature iGABA neurons (WPI6),

however, the average wMFR decreased in 5'-Del and 3'-Del neurons (5'-Del p = 0.0323; 1-way
ANOVA, Dunnett's test) (**Fig. 3h**). Like iGLUT neurons, the passive and intrinsic excitability
membrane properties of mature iGABA neurons were not different between *NRXN1*[+/-] 5'-Del or
3'-Del and controls (**Fig. 3i,j and Extended Data Fig. 4d**). Overall, these data suggest that
patient-specific changes in spontaneous neural activity are not fully explained by differences in
passive and excitable membrane properties in iGLUT or iGABA neurons from *NRXN1*[+/-] 5'-Del
and 3'-Del patients.

***NRXN1*[+/-] 5' and 3' deletions result in divergent synaptic transmission deficits.**

To further dissect the factors mediating phenotypes in altered spontaneous firing, we investigated
the efficacy of synaptic transmission by patch-clamp electrophysiology. Voltage-clamp recordings
of spontaneous excitatory post-synaptic currents (sEPSCs, no TTX) in iGLUT neurons revealed
decreased frequency of events for both for 5'-Del and 3'-Del iGLUT neurons (**Fig. 4a**). The
cumulative probabilities of inter-event-intervals (IEI) for both 3'-Del (p = 4.88E-5, Levene's test
with Bonferroni correction) and 5'-Del iGLUTs (p = 2.54E-11, Levene's test with Bonferroni
correction) (**Fig. 4a,b**) was significantly increased, compared to controls. The sEPSC amplitude
increased for 5'-Del neurons (p = 3.88E-4, Levene's test with Bonferroni correction) (**Fig. 4a,c**).
Miniature excitatory post-synaptic currents (mEPSCs, +TTX) showed similar trends in IEI but no
changes in amplitude sizes across genotypes (IEI 5'-Del p = 0.0411, 1-way ANOVA, Dunnett's
test) (**Extended Data Fig. 4b,c**). These reductions in synaptic transmission are consistent with
the transcriptomic signatures; pre-synaptic (SynGO) genes showed a larger change (than post-
synaptic) in synaptic gene expression signatures (5'-Del Pre-SynGO, Log$_2$FC = 0.1252; 3'-Del
Pre-SynGO, Log$_2$FC= -0.0156) (**Fig. 4d,e**). We further probed transcriptional signatures of known
NRXN1 trans-synaptic interaction partners that mediate synapse formation, function, plasticity
and are frequently linked with neuropsychiatric disease[16], and found they were represented in
DEGs, including *CBLN2*, *LRRTM4* and *NXPH1*, suggesting these synaptic effects are largely
driven by change in *NRXN1* expression (**Fig. 4f**).

By contrast, synaptic transmission in iGABA neurons appeared to be enhanced. The frequency
of spontaneous inhibitory post-synaptic currents (sIPSCs) increased, marked by a significant
decrease in IEI in 3'-Del iGABA neurons (p = 1.00E-4 by Levene's test with Bonferroni correction)
(**Fig. 4i,j**). There was no change in sIPSC amplitude (**Fig. 4i,k**). Miniature inhibitory post-synaptic
currents (mIPSCs) recorded in the presence of TTX and CNQX (AMPA/Kinate receptor
antagonist) revealed similar trends in IEI, with no change in mIPSC amplitudes (**Extended Data**

Fig. 4e,f). Similarly, SynGO analysis revealed concordant changes in synaptic transmission and
transcriptomic signatures at 3'-Del (Pre-SynGO Log$_2$FC = 0.0324) and represented DEGs among
NRXN1 trans-synaptic interaction partners *CBLN2*, *NLGN1*, *NXPH1, CASK,* and *LRRTM2-4* (**Fig.**
4l-n).

Quantification of synaptic puncta (via immunostaining against synapsin-1, SYN1), normalized to
dendritic length (via immunostaining somatodendritic marker MAP2) uncovered a bidirectional
decrease in iGLUT neurons (**Fig. 4g,h**), and an increase in iGABA neurons (**Fig. 4o,p**). Thus,
divergent neurotransmission phenotypes appear to correlate with synapse number. Overall,
patient-specific alterations in spontaneous neural activity were driven by synaptic deficits, with the
cell type-specific impact of aberrant *NRXN1* splicing resulting in divergent neurotransmission
phenotypes and changes in synapse number. Furthermore, non-recurrent LOF and GOF
presented unequal effect sizes between cell-types, with 5'-Del neurons being most impacted by
excitatory transmission, but 3'-Del neurons being affected by both excitatory and inhibitory
transmission. Given that deletions affected iGLUT and iGABA neurons in opposing directions,
these findings implicate *NRXN1* as a key mediator of excitatory/inhibitory balance, a prevalent
theme among neuropsychiatric disorders[38].

Isogenic validation of bidirectional excitatory-inhibitory (E-I) synaptic deficits.

To demonstrate a direct link between aberrant *NRXN1* splicing and synaptic dysfunction, we
designed an experiment to specifically target splicing patterns. We utilized short hairpin RNAs
(shRNAs) to knockdown wildtype splice isoforms in control lines, mimicking a LOF phenotype.
Targeted knockdown of constitutively expressed exon 9 (expressed in alpha isoforms but not beta
isoforms) achieved mRNA knockdown of ~55% in iGLUT neurons, and ~75% in iGABA neurons
across one or more isogenic pairs (compared to a non-targeting (NT) control shRNA) (**Extended**
Data Fig. 6a,e). These knockdowns would roughly mimic a 5'-Del heterozygote. Differential
splicing analysis confirmed the reduction of β→α splicing in both iGLUT and iGABA neurons by
ΔPSI -0.191 (*p* = 0.0083*)* and -0.208 (*p* = 0.0202), respectively (**Fig. 5a, i**), without significantly
altering other *NRXN1* splice sites. Functionally, we observed decreased synaptic transmission in
iGLUT neurons (i.e., increased sEPSC IEI, *p* = 2.2E-16 by Levene's Test) (**Fig. 5b,c**), and
increased synaptic transmission in iGABA neurons (decreased IEI, *p* = 3.65E-8 by Levene's Test
and *p* = 0.0092 by Student's t-test) (**Fig. 5j,k**), similar to the changes with 5'-Del neurons (**Fig.**
4b,j). Transcriptomic profiles of isogenic lines further validated *NRXN1* knockdown (**Extended**
Data Fig 6c,g) and demonstrated DEGs related to multiple aspects of synaptic function in both

iGLUT (29/258 DEGs; 1.274161-fold, p = 0.1049) and iGABA neurons (384/3525 genes;

1.234861-fold, p = 1.307E-6) (**Fig. 5d,i and Supplementary Table S2**). Altogether, knockdown

of wildtype splicing recapitulated cell-type-specific differences in $NRXN1^{+/-}$ 5'-Del neurons,

causally implicating decreased wildtype $NRXN1\alpha$ expression (LOF) as a driver of cell-type-

specific phenotypes.

For GOF $NRXN1$ 3'-Del neurons, we took a different approach and applied shRNAs to knockdown

mutant splice isoforms, in an attempt to reverse the GOF phenotype. We designed a shRNA

against the mutant splice junction overlapping exons 20 and 24, expressed in all 3'-Del unique

$NRXN1$ alpha and beta isoforms, and achieved targeted knockdown of mutant splice isoforms by

95% in iGLUT, and 25% in iGABA neurons in both donors (**Extended Data Fig 6b,f**). Differential

splicing analysis again confirmed the selective reduction of GOF splicing in both iGLUT and

iGABA neurons, by ΔPSI -0.108 (p = 0.0623) and -0.074 (p = 0.0258), respectively (**Fig**. **5e,m**),

without significantly altering other $NRXN1$ splice sites. In the donor with the most robust

knockdown in both iGLUT (90%) and iGABA (37%) neurons, electrophysiological recordings

revealed a reversal of synaptic transmission phenotypes, achieving IEI levels similar to control

iGLUT and iGABA neurons (**Fig. 4b,j**). In iGLUT neurons, shRNA MT decreased sEPSC IEI (p <

2.2E-16 by Levene's Test and p = 0.0194) (**Fig. 5f,g**) and in iGABA neurons increased sIPSC IEI

(p = 0.042 by Levene's Test) (**Fig 5h,o**), as compared to shRNA-NT. Parallel to the LOF isogenic

transcriptomic profiles, the GOF isogenic line had modest changes in iGLUT neurons (3/28 DEGs;

1.214533-fold, p = 0.4541 by Levene's test), but more pronounced synaptic DEGs in iGABA

neurons (128/1077 DEGs; 1.347225-fold, p = 2.694E-4) (**Fig. 5h,p and Extended Data Fig.**

6d,h).

In summary, knockdown of WT $NRXN1$ in iGLUT and iGABA neurons recapitulated the

consequences of reduced β→α splicing on neurotransmitter phenotypes of 5' LOF $NRXN1$

deletions, while knockdown of MT splicing appeared to mitigate the negative effect of the 3' GOF

$NRXN1$ deletions. Furthermore, the bidirectional effect of $NRXN1$ deletions on iGLUTs

(decrease) and iGABAs (increase), i.e., E-I balance, was reversed by shRNA treatments.

Transcriptomic profiling of shRNA treated isogenic lines revealed ~10% of DEGs related to

synaptic function across all four experimental conditions. Altogether, shRNA mediated

perturbations among WT and MT isoforms causally implicate $NRXN1$ splicing to synaptic

dysfunction.

Framework for precision medicine against stratified LOF/GOF phenotypes

As a proof-of-principle therapeutic intervention, we tested methods to directly and indirectly
manipulate *NRXN1* expression, separately targeting LOF and GOF mechanisms, and focusing
on reversing decreased excitation phenotypes in iGLUT neurons.

For LOF patients, we hypothesized that increasing transcription of the WT *NRXN1α* allele would
restore *NRXN1* levels and reverse the signature of reduced β→α splicing (**Fig. 6a**). β-estradiol
reversed *NRXN1* LOF neurogenesis deficits in *xenopus* and human NPC models[39]. Although the
mechanism is unknown, chromatin immunoprecipitation with sequencing (ChIP-Seq) in mouse
brain tissue identified estrogen receptor alphaα (ERα) binding sites at the *NRXN1* alpha locus
(**Extended Data Fig. 7a,b**)[40]. We predicted *NRXN1* to be a target of ERα, and report that acute
treatment with β-estradiol (10nM or 30nM, 3-5 days) significantly increased *NRXN1α* expression
in iGLUT neurons (p = 0.0297 by Student's t-test) derived from 5'-Del patients (**Fig. 6b**), but not
in controls (**Extended Data Fig. 7c**). The functional effect of chronic treatment of post-mitotic 5'-
Del neurons with 30nM β-estradiol (relative to DMSO vehicle) was evaluated across spontaneous
neural activity (MEA), synaptic transmission (patch-clamp) and gene expression (RNA-
sequencing). MEA recordings at WPI3 revealed a significant increase in wMFR activity (p =
0.0057 via student's t-test) in the β-estradiol treated condition (**Fig. 6c**). Transcriptional profiling
of the β-estradiol treatment detected an inversion of the β→α splicing signature (ΔPSI 0.045), and
more subtle changes in global gene expression (7 upregulated and 34 downregulated DEGs)
(**Fig. 6d,e**). Likewise, patch-clamp recordings revealed that β-estradiol treatment ameliorated the
sEPSC IEI phenotype in 5'-Del neurons (p = 4.62E-4 by Levene's Test, Bonferroni corrected),
but not in vehicle treated 5'-Del neurons (p = 0.9055 by Levene's Test), compared to healthy
controls (**Fig. 6f,g**).

Towards a therapeutic strategy for treating GOF patients, we investigated the utility of anti-sense
oligonucleotides (ASOs), recently used to treat several neurological diseases[41], to target a
specific RNA splice site. We designed an "alternative splice matrix" by juxtaposing splice donor
and acceptor RNA sequences from each *NRXN1α* exon, generating all possible combinations of
canonical and non-canonical splice junctions (**Fig. 6h,i**). Then, for the sequence covering the
20/24 mutant splice junction, we designed an ASO to facilitate targeted RNAseH1-dependent
degradation of 3-Del mutant isoforms. ASO treatment (1μM) for 72hrs decreased total mutant
isoforms by ~55% in iGLUT neurons (p < 0.001 by student's t-test), as compared to a non-
targeting control ASO (**Fig. 6j**). Differential splicing analysis confirmed the reduction of GOF
splicing (ΔPSI -0.103) and increased ratios of SS4+. We also observed a significant reduction in

$\beta \rightarrow \alpha$ splicing (ΔPSI -0.253, *p* = 0.007, Bonferroni corrected), perhaps resulting from reduced MT

isoform expression, contrasting the elevated baseline of *NRXN1α* in 3'-Del (**Fig. 6k**).

Transcriptomic profiling revealed robust DEGs, enriched for synaptic properties, neurotransmitter

signaling, and neurodevelopmental pathways **(Supplementary Table S3)** (231/1844 Bonferroni

corrected DEGs annotated in SynGO; 1.420028-fold, *p* = 1.016E-08), (**Fig. 6c-d**). To demonstrate

the broader applicability of an ASO strategy, we report mutant splice isoforms in post-mortem

brain tissue from an unrelated *NRXN1[+/-]* case diagnosed with autism spectrum disorder (ASD)

(**Supplemental Fig. 8**). By integrating long-read and short-read sequencing[15], we identified nine

high-confidence isoforms that are predicted to be translated: two of which contained a novel splice

junction overlapping the deletion encompassing exon 14/19, targetable by an ASO tailored to the

mutant splice junction.

Altogether, increased *NRXN1* expression in LOF neurons and knockdown of mutant splicing in

GOF neurons can each rescue the splicing defects that produce opposing cell-type-specific

case/control differences in *NRXN1[+/-]* neurons. We propose the aberrant *NRXN1α* splicing is a key

a driver of complex *NRXN1[+/-]* phenotypes and that modulators of *NRXN1* expression and/or

splicing represent novel targeted therapies.

DISCUSSION

NRXN1[+/-] deletions are non-recurrent between patients, linked to diverse clinical outcomes that

cannot be explained by the size or boundaries of the deletion itself[6]. Here, we show that *NRXN1*

splicing alters excitatory-inhibitory (E-I) imbalance[1,32], a common theme among neuropsychiatric

disorders, by bidirectionally regulating synaptic transmission, with a decrease in frequency of

sEPSCs in 5'-Del and 3'-Del iGLUT neurons but an increase in frequency of sIPSCs in 3'-Del

iGABA neurons. Using a case/control *NRXN1[+/-]* cohort as well as isogenic manipulations of the

NRXN1 isoform repertoire, we report distinct phenotypic effects in human iGLUT and iGABA

neurons that predominately manifest in changes in the frequency of synaptic function. These

results suggest causal relationships between aberrant *NRXN1* splicing and synaptic dysfunction,

dictated by unique patient-specific mechanisms.

The regulation of *NRXN1* splicing, including the formation of aberrant splice sites by specific

RBPs, is poorly understood; >100 RBPs are predicted to interact with *NRXN1* mRNA in a cell-

type[42,43] and neuronal activity[23] dependent-manner. For example, KH-domain STAR-family

RBPs regulate SS4+ in a neuronal activity dependent manner[23–25], mediating trans-synaptic

signaling by varying interactions among a host of post-synaptic ligands. Although patient

transcriptomic profiles nominate STAR proteins as potential drivers of aberrant GOF splicing,

further investigation is required to identify the biochemical mechanisms involved. Nevertheless,

we posit that direct manipulation of splicing may achieve therapeutic benefit in some cases.

Neurexins are expressed across all synapses and among certain non-neuronal cell-types, such

as astrocytes, perhaps with distinct cell-type-specific functional roles[37]. Mechanistically, our study

did not resolve whether mutant GOF isoforms shifted the stoichiometry of alternative splicing

against wildtype RNA isoforms or if they altered trans-synaptic protein-protein interactions. In

either case, *NRXN1* encodes a pre-synaptic molecule but traditional neuropharmacological

agents typically target specific receptors. Given that patient-specific patterns of aberrant *NRXN1*

splicing across non-recurrent mutations are unlikely to be reversed by a pharmacological

approach, precision therapies that target aberrant splicing are needed to effectively reverse

defects in the multiple cell types affected by patient-specific deletions.

The opposing effects of *NRXN1*[+/-] in glutamate and GABA neurons provides the foundation for

evaluating proof-of-concept therapeutics that target distinct LOF or GOF mechanisms. For LOF

mutations, mechanisms for upregulation of wild-type allelic expression and/or restoration of

proper *NRXN1* splicing are required (**Fig. 6a**). While direct approaches to increase *NRXN1*

expression may be ideal, such as small activating RNAs[44], further work is required to carefully

evaluate safety and efficacy in clinical trials. On the other hand, indirect approaches have the

advantage of prioritizing drugs from a list of already clinically approved molecules, selected based

on their predicted ability to alter the expression of a target gene (e.g., LINCs[45]). With relevance to

the ability of estradiol to increase *NRXN1* and rescue synaptic deficits, steroid-based

pharmaceuticals are already used to treat neuroinflammation-related conditions[46], with recent

studies revealing estrogen-mediated roles in neuroprotection[39], demonstrating the safety and

feasibility of this approach. For GOF, ASOs designed to target a mutant splice junction and

facilitate degradation of MT isoforms may represent a viable therapeutic approach (**Fig. 6h**). Not

all in-frame coding mutations will be GOF; for example, unique mutations at non-canonical splice

donor/acceptor sites will require further characterization prior to assigning LOF/GOF status. In all

cases, maintaining a stoichiometric balance between MT and WT isoforms may require

combination treatments (e.g., ASO and transcriptional activator) for therapeutic benefit. Cases of

muscular dystrophies treated via splicing modulating ASOs have proven effective[47], but overall,

gene therapy in brain disease has shown mixed successes[48]. Critically, although our proof-of-

principle *in vitro* findings suggest a novel therapeutic avenue, translational studies will be required

to confirm the efficacy of this framework *in vivo*, and ultimately, in the clinic.

Several technical limitations warrant acknowledgement. The iGLUT and iGABA neuron
populations studied herein are not just immature, but also comprise diverse neuronal subtypes[49];
thus, the more discrete impacts of *NRXN1[+/-]* on synaptic physiology across different subtypes of
neurons (e.g., SST versus PV expressing GABA neurons) remain unresolved. Moreover, future
studies in more physiologically relevant and circuit-like models may uncover novel non-
autonomous and activity-dependent phenotypes. Mechanistically, it will be important to probe the
biochemical and proteomic interactions of *NRXN1*; for example, unbiased proximity-labelling
methods (e.g., BioID[50]) could define perturbations in protein-protein interaction profiles between
wildtype and mutant isoforms.

Precision medicine seeks to tailor treatments to individual patients,[51] regardless of differential
penetrance of genetic mutations. Therefore, we focused on stratifying mutations in the same gene
based on LOF and GOF mechanisms, in contrast to clinical diagnosis (bipolar disorder or
schizophrenia), with the goal of one day providing the right therapy at the right time to the right
patient. The most successful examples of precision medicine to date have been in cancer[52] and
monogenic disease[53], whereby genetic analyses revealed molecular subtypes that benefited from
specific treatments. In the context of brain disorders, patient stratification is particularly
challenging, reflecting in part the myriad rare and common variants linked to disease[54]. Disease-
agnostic analyses reveal that transcription factors and nucleic acid binding proteins are
overwhelmingly driven by LOF mutations, whereas signaling molecules, enzymes, receptors and
transporters more frequently incur dominant GOF mutations[55]. When both are possible, distinct
mutations in the same genes can result in pleiotropic phenotypic effects. A binary therapeutic
approach, similar to what we proposed here for *NRXN1*, may prove suitable for dual LOF an GOF
mechanisms linked to mutations in other neuropsychiatric disorder-related synaptic genes (e.g.,
NLGN3[56], *CACNA1D*[57], *CACNA1C*[58], *SCN1A*[59], *SCN2A*[60]) as well as broadly in
neurodegenerative disease (e.g., *SOD1*[61], *TDP43*[61], *FUS/TLS*[61], *C9ORF72*[62], *AR*[63]), and many
short tandem repeat disorders[64]. Taken together, our work advances precision medicine,
demonstrating the necessity of functionally dissecting the phenotypic impact of diverse patient-
specific genetic variants across cellular contexts, in order to resolve candidate therapies across
stratified disease mechanisms.

METHODS

Plasmid designs and molecular cloning

i. TetO-Ascl1-Neo

The _ASCL1_ insert from TetO-Ascl1-Puro (Addgene #97329) was synthesized as a gBLOCK
flanked by EcoR1 cut sites, and cloned into TetO-hNgn2-Neo using EcoR1 to remove _NGN2._ The
recipient vector was dephosphorylated with shrimp alkaline phosphatase (rSAP NEB #M0371S)
during the digest, column purified and ligated at a 1:1 vector to insert ratio using the QuickLig Kit
(NEB #M2200S). The _ASCL1_ stop codon was subsequently mutated using the QuickChange II-
XL site-directed mutagenesis kit (Agilent #200523) and verified via whole plasmid sequencing
from plasmidsaurus.

ii. shRNA RNA interference constructs

All shRNAs were designed and produced by Sigma-Aldrich via custom submitted sequences
against wildtype _NRXN1α_ (constitutively expressed exon 9) and mutant _NRXN1α_ (mutant 20/24
splice junction) cloned into TCR2 pLKO.5-puro.

Cell Culture

i. hiPSC culture

Passage matched (±3) human induced pluripotent stem cells (hiPSCs) were cultured in StemFlex
media (Life technologies #A3349401) on Matrigel (Corning, #354230). At ~70-80% hiPSCs were
clump passaged using 0.5mM EDTA in PBS without Mg/Ca (Life technologies #10010-031). Cells
were washed once and incubated for 4-7 min with 0.5μM EDTA at RT. The EDTA was aspirated,
and the remaining colonies were lifted off with 1mL of StemFlex and re-distributed at varying
densities in a 6–well maintenance plate. hiPSC lines were split every 4-6 days. For
neuronal/organoid differentiation, wells of similar confluence across all hiPSC donors were
resuspended and seeded onto a Matrigel coated 10cm dish and expanded until ~70-80%
confluency.

ii. HEK293T culture and lenti-viral production

HEK293T cells were maintained in 15cm dishes and cultured in DMEM supplemented with 10%
standard FBS (Life technologies #16000069). 3rd Gen lenti-viral particles were produced using

previously described methods and REV, MDL and VSV-G plasmid ratios[18], each transfected with
PEImax (Polysciences #24765-1). Each PEImax batch was volumetrically titrated at total µgDNA:
µLPEI for optimal transfection efficiency.

iii. Primary mouse glia production

All mouse work was performed under approved IACUC protocols at the Icahn School of Medicine
at Mount Sinai. C57BL/6 mice were used as breeding pairs. For glial preps, dissected cortices
from 3 pups (at p0-3) were dissociated using papain (R&D #LS003126) and plated on 10cm
dishes in MEF medium (DMEM, 10% Cosmic Calf Serum (Fisher #SH3008703HI), 1x Antibiotic-
antimycotic (Life technologies #15240), 1x Sodium Pyruvate (Life technologies #11360070), 1x
MEM Non-Essential Amino Acids Solution (Life technologies #11140050), 4uL 2-Mercaptoethanol
(Sigma #60-24-2), supplemented with 100µg/mL Normocin (InvivoGen #ant-nr-2). Glial cells were
recovered and propagated for 7 days, and expanded into three 10cm dishes. To promote neuronal
health and synapse maturation, we utilized mouse glia from well-established protocols that
significantly outperform human astrocytes for co-culture experiments[19,21,65]. All glial preps were
tested twice for mycoplasma (Normocin withdrawn) (Lonza, #LT07-318) prior to freezing or
neuronal co-culture. At day 14, one 10cm dish with mouse glia were distributed to two MEA, 12-
or 24- well plates, and subsequently inactivated with 4µM Ara-C (Sigma #C1768) prior to or during
re-seeding of induced neurons.

iv. iGLUT induction and astrocyte co-culture

At day -1 hiPSCs expanded in 10cm dishes were dissociated with accutase (StemCell
Technologies, #07920), washed and pelleted with a 1:4 ratio of accutase to DMEM for 5 min at
1000 rcf, and re-suspended with StemFlex media containing ROCK inhibitor, THX (10µM/mL;
Sigma Aldrich, SML1045). The hiPSCs are then co-transduced with TetO-Ngn2-Puro (Addgene
#79049) or TetO-Ngn2-Neo (Addgene# 99378) and UCb-rtTA (legacy vector from the lab of Fred
Gage) and seeded at $1.0\text{-}1.5\times10^6$ cells in 1.5mL per well on 6-well plates were coated with 2x
Matrigel for at least one hour at 37°C. The hiPSC-viral mixture was then incubated overnight. The
following morning (day 0), a full media change with iGLUT induction media was performed with
the following recipe: Neurobasal Media: 1x N2 (Life technologies #17502-048), 1x B-27 minus
Vitamin A (Life technologies #12587-010), 1x Antibiotic-Antimycotic, 1x Sodium Pyruvate, 1x
GlutaMax (Life technologies #35050), 500µg/mL cyclic-AMP (Sigma #D0627), 200µM L-ascorbic
acid (Sigma #A0278), 20ng/ml BDNF (Peprotech #450-02), 20ng/ml GDNF (Peprotech #450-10),

1µg/ml natural mouse laminin (Life technologies #23017015). On days 1-2, iGLUT cells were
treated with respective antibiotic concentrations at 1µg/mL puromycin (Sigma# P7255) or
0.5µg/mL neomycin (Life technologies #11811-031). On day 3, antibiotic medium was withdrawn
and iGLUT cells were treated with 4µM Ara-C. On Day 4, iGLUT cells were dissociated with
accutase for 15 min, washed and pelleted with a 1:4 ratio of accutase to DMEM for 5 min at 800
rcf, and re-suspended with iGLUT media containing ROCK inhibitor, Ara-C and 2% low-
hemoglobin FBS (R&D systems #S11510). iGLUT neurons were distributed among wells (500-
750k cells per 24wp or 0.75-1.5E6 cell per 12wp) pre-seeded with confluent mouse glia. The
following day, iGLUT neurons received a full media change with Brainphys maturation media
(Neurobasal Media, 1x N2, 1x B-27 minus Vitamin A, 1x Antibiotic-Antimycotic, 500µg/mL cyclic-
AMP, 200µM Ascorbic Acid, 20ng/ml BDNF, 20ng/ml GDNF, 2% low-hemoglobin FBS, 1µg/ml
Mouse Laminin) supplemented with Ara-C and were subsequently monitored for growth of non-
neuronal/glial cells. Ara-C treatment was tittered down with half-media changes (without Ara-C)
every 3-4 days until used for experiments.

v. iGABA induction and astrocyte co-culture

iGABA production paralleled the methods aforementioned. hiPSCs were instead co-transduced
with TetO-Ascl1-puro (Addgene #97330) or TetO-Ascl1-neo (Addgene #TBD), TetO-Dlx2-hygro
(Addgene #97329) and UCb-rtTA and seeded at 0.8-1.2x10^6 cells in 1.5mL per well on 6-well
plates similarly prepared. The following morning (day 0), a full media change with iGABA induction
media (DMEM/F-12 + Glutamax, 1x N2, 1x B-27 minus Vitamin A, 1x Antibiotic-Antimycotic) was
performed. On days 1-2 iGABA cells were selected with respective antibiotic concentrations at
1µg/mL puromycin or 0.5µg/mL neomycin and 0.25µg/mL hygromycin (Life Technologies
#10687010), followed by antibiotic withdrawal and Ara-C treatment on day 3. iGABA neurons
were re-seeded identically to iGLUT cells, at 150-250k cells per 24wp well. iGABA cultures were
morphologically QC'ed prior to all experiments, with uncharacteristic batches being discarded.

vi. Cortical and subpallial organoid differentiation

Cortical organoids were generated according to the protocol described by Sloan et. al.[66], with
several modifications. hiPSCs were first aggregated into embryoid bodies (EBs) using an
AggreWell™800 Microwell Culture Plate system (Stemcell Tech #34850). Expanded hiPSCs
were rinsed twice with DPBS without Ca/Mg, and then dissociated using accutase. 3x10E6
hiPSCs were added to a single well in the prepared AggreWell and allowed to aggregate in a

37°C incubator for 24 hours. The following day (day 0), EBs were dislodged from the AggreWell

plate using a cut p1000 pipette tip and passed over a 40µm strainer, and washed with excess

DMEM. The strainer was inverted over an Ultra-Low Attachment 10 cm culture dish and the EBs

were collected in spheroid induction media, which contained Stemflex supplemented with two

SMAD inhibitors, SB-431542 (SB) and LDN193189 (LDN), and THX. The following day (day 1),

the media THX was withdrawn. From d2-d6, induction media was replaced daily, and no longer

contained the Stemflex supplement (only base Stemflex media with SB and LDN). On day 6, the

media was replaced with organoid patterning media, formulated with Neurobasal-A medium, 1x

B-27 minus Vitamin A, 1x GlutaMAX, and 1x Antibiotic-Antimycotic. From d6-d24, organoid

maturation media was supplemented with 20ng/ml of EGF (R&D Systems, #236-EG) and 20ng/ml

FGF2 (R&D Systems, #233-FB-01M). Media was changed every day from d6 – d15, after which

media was changed every other day. From d25-d43, the organoid maturation media was

supplemented with 20ng/ml of NT-3 (PeproTech, #450-03) and 20ng/ml BDNF. From d43

onwards, organoids received organoid maturation media with no supplements, and media was

changed every 4 days or as needed. Subpallial organoids were generated in the same way as

cortical organoids, but with additions in media formulations. From d4-d23, hSOs received

spheroid induction media or organoid maturation media supplemented with 5µM of the Wnt

inhibitor, IWP-2 (Selleckchem, #S7085). From d12-d23, hSO organoids received neuronal

differentiation media supplemented with 100nM of the SHH agonist, SAG (Selleckchem, #S7779).

Electrophysiology

i. Multi-electrode array (MEA)

The Axion Maestro (Middleman) system was used to perform all MEA recordings. Following

iGLUT and iGABA inductions, 80-100k cells were re-plated on each MEA well and measurements

began as early as DIV9 for both iGLUT and iGABA co-cultures. For time course experiments,

MEA plates were recorded every 2-3 days per week with a full media change prior to each

recording. Comparisons for iGLUT neurons were made at WPI4 and WPI6, as well characterized

timepoints for synaptic activity. For iGABA neurons, comparisons were made at WPI2 and WPI6

to include both timepoints of elevated neuronal activity. For acute drug treatments with gabazine

(Tocris, #1262), a baseline recording (pre-treatment) was first obtained, followed by an immediate

addition of a small volume of concentrated stock. A second, post-treatment recording was then

obtained to evaluate the difference in activity before and after drug treatment. A full media change

was performed one day prior to the day of recording. MEA wells were visually QC'ed for similar

densities prior to recording, with high/low density wells being excluded from recordings.

ii. Whole-cell patch-clamp electrophysiology

For whole-cell patch-clamp recordings, iGLUT (300k/well) or iGABA (250k/well) human-mouse
glia co-cultures were recorded at 4-6 weeks following dox-induction (time points specified in figure
legends), with a full media change one day prior to recording. Only coverslips with similar densities
were selected for recording. Cells were visualized on a Nikon inverted microscope equipped with
fluorescence and Hoffman optics. Neurons were recorded with an Axopatch 200B amplifier
(Molecular Devices), digitized at 10 kHz using a Digidata 1320a (Molecular Devices) and filtered
between 1-10 kHz, using Clampex 10 software (Molecular Devices). Patch pipettes were pulled
from borosilicate glass electrodes (Warner Instruments) to a final tip resistance of 3-5 MΩ using
a vertical gravity puller (Narishige). To sustain the baseline activity of neurons from extended
cultures, coverslips were recorded in base Brainphys medium (external solution, Osm 305, pH
7.49). Each coverslip was equilibrated to room temperature for 10min, with 3-5 neurons were
recorded for no more than a total of 75min per coverslip. For measurements of passive and
excitable properties, an internal patch solution was used containing (in mM): K-d-gluconate, 140;
NaCl, 4; MgCl2, 2; EGTA, 1.1; HEPES, 5; Na2ATP, 2; sodium creatine phosphate, 5; Na3GTP,
0.6, at a pH of 7.4. Osmolarity was 290-295 mOsm. Neurons were chosen at random using DIC
and all recordings were made at room temperature (~22°C). Current-clamp measurement
occurred across -10pA to +50pA steps, with a maximum stimulus of +60pA, whereas voltage-
clamp measurements occurred across -50mV to +50mV steps, normalized to cell capacitance (to
control for variable neuronal size). Current clamp measurements were corrected for the junction
potential (~-15.5mV). For sEPSC/sIPSC recordings, the internal solution was replaced with (in
mM): Cesium-Chloride, 135; HEPES-CoOH, 10; QX-314, 5; EGTA, 5. Osmolarity was 290-295
mOsm. All mEPSC measurements were recorded under the presence of 100nM TTX-citrate
(Tocris Cat# 1069). mIPSC measurements were made using 100nM TTX-citrate and CNQX-
disodium salt (Tocris Cat#1045/1) to pharmacologically inhibit ionotropic glutamate receptors. All
chemicals were purchased from Sigma-Aldrich Co. (St. Louis, MO). All toxic compounds were
handled in accordance with ISMMS EHS standards.

iii. Patch-clamp data analysis

All patch-clamp data were analyzed on Clampfit (v11) and Easy-Electrophysiology (v2.4.1 or beta-
versions). Briefly, for voltage-clamp data, files were opened in Easy-Electrophysiology and two

bins were assigned for Na+/K+ measures for minimum and maximum current values, respectively.

For current-clamp data, files were opened in Easy-Electrophysiology and action potential (AP)

analysis was used to automatically determine spike number and properties. For gap-free

recordings, all data was post-hoc adjusted to a baseline of zero on ClampFit, and subsequently

analyzed in Easy-Electrophysiology by template and linear threshold detection. For case v. control

experiments, a minimum cutoff of 10 events for the duration of the recording (3min) was used as

QC. For typical EPSC events, a single template from a randomly chosen recording was used to

analyze all traces (with a 30ms decay cutoff). For IPSC events, three templates were used to

detect variable GABA receptor kinetics, for all traces (with a 60ms decay cutoff). An amplitude cut

off of 7pA was used to call positive events. For cumulative probabilities, each cell was binned by

experiment and averaged across all cells for a representative curve (GraphPad Prism).

RNA-Sequencing and bioinformatic analyses

i. Bulk RNA sequencing and DEG analysis of iGLUT and iGABA neurons

iGLUT (DIV21) and iGABA (DIV14 or DIV35) co-cultured with primary mouse glia (to match

functional experiments) were harvested in Trizol and submitted to the New York Genome Center

for high-throughput RNA extraction and quality control, library prep using the Kapa Hyper library

prep with ribo-erase (Roche #KK8541) and subsequently sequenced on Illumina NovaSeq.

Similarly, shRNA samples were harvested at DIV21-24, and DIV35-49 for iGLUT and iGABA

neurons, respectively. Returned raw data files were first processed to remove mouse reads in the

RNA-seq data, a combined genome reference (hg38 and mm10) was created using the "mkref"

command in cellranger (v6.1.2). The raw sequencing reads were then aligned to the combined

genome reference using STAR (v2.7.2a)[67]. Reads that mapped specifically to the human

reference genome were extracted from the resulting BAM files for subsequent gene expression

analysis. Gene-level read counts were obtained using the Subread (v2.0.1) package

featureCount[68], and RPKM values were calculated using the R package edgeR[69]. To confirm

sample identity, variants were called from RNA-seq bam files by HaplotypeCaller and

GenotypeGVCFs in GATK (v4.2.0). Then bcftools (v1.15) was used

to examine variants concordance with variants from whole-exome sequencing data from the

same donor. Following donor identity confirmation, the differential gene expression analysis

followed the methods as described previously[70]. First, CibersortX[71] was utilized to predict

differences in cell type composition across all samples. The R package variancePartition

(v1.30.2)[72] was then employed to investigate the contribution of specific variables to the variance

in gene expression. The limma/voom package[27] was used for differential expression analysis,
with the regression of fibroblast and hiPSC cell type compositions. The analysis began with a
comparison between the case and control groups. Subsequently, within each case vs control
group, subgroup comparisons were conducted for all four pairs (two donors each for 3'-Del and
5-Del patients and two healthy controls) of samples, accounting for heterogeneity between
different donors. Genes with an FDR less than 0.1, and a fold change above 1.5 and below -1.5
in all four pairs of subgroup comparisons were defined as the final set of differentially expressed
genes, unless otherwise specified. Kallisto (v0.46.1)[73] was used to calculate the *NRXN1* exon
usage ratios.

ii. Analysis of alternative splicing estimates via LeafCutter

Reads were aligned to a combined GRCh38 human and GRCm38 mouse reference genome
using STAR (v2.7.1.a), with an index built against GRCh38 Gencode GTF (v92) using the option
-sjdbOverhang 100. To allow the discovery of novel splice junctions and increase mapping
accuracy, STAR was run in two-pass mode with standard options. Reads mapping exclusively to
GRCh38 were subsequently extracted and replicates merged with samtools (v1.6). Splice
junctions were extracted from the resulting bam files with regtools v0.5.2 `junctions extract` with
parameters "-a 8", "-m 50", and "-M 1200000" due to potential for the genomic deletions to cause
larger intervals between junctions. Junctions were clustered using the leafcutter pipeline
(v2c9907e) script "leafcutter_cluster_regtools.py" with option "-l 1200000". For differential splicing
analysis, leafcutter was run with "-i 2 -g 0" to reflect the samples used. Splicegraph visualizations
were constructed using a modified version of the leafviz pipeline. All statistical tests utilized a
Dirichlet-multinomial generalized linear model, and were corrected for multiple comparisons when
necessary via Bonferroni adjustment.

iii. Pathway and Network Analysis of DEGs

For hierarchical clustering of DEGs based on gene expression fold change, the Pheatmap
function in R was used cluster gene expression fold change of DEGs combined from both 5'-Del
and 3'-Del conditions. R package clusterProfiler[29] was used to performed the gene set enrichment
analysis (GSEA). For each cell type, ranked gene expression of each genotype (5'-Del or 3'-Del)
against control were used as background and DEGs from both genotypes (5'-Del and 3'-Del) were
used as query gene sets. For Gene Ontology (GO) enrichment analysis, the enrichGO function
within R package clusterProfiler was used. For GO with a default background, query gene list was

converted from 'ENSEMBL' format to 'ENTREZID' format by bitr function in clusterProfiler and the
OrgDb parameter was set as 'org.Hs.eg.db'. When customized genes were used as background,
both background and query gene lists were kept as 'ENSEMBL' format. For SynapseGO (SynGO
v1.2), DEG or prioritized DEG lists (Log$_2$FC filtered) gene lists were tested for enrichment with
"brain expressed" background set selected, containing 18035 unique genes in total of which 1591
overlap with SynGO annotated genes. Sunburst plots were exported via the web-based
application. The ASD, BP, and SCZ risk gene lists were extracted from previously curated gene
lists[30], Genes with the top 200 smallest FDR values and a fold change larger than 1.5 in the case
vs control comparison were selected for the protein interaction network analysis. Then,
overlapping genes between the selected gene list, the disease risk gene sets, and the proteins in
the SIGNOR database were utilized to query the SIGNOR database and build the interaction
network, using the "connect + add bridge proteins" search mode in The SIGNOR Cytoscape App
(v1.2)[74].

v. Dissociation and 10x Single-Cell RNA sequencing of organoids

Whole organoids were dissociated to the single cell level in preparation for single-cell RNA
sequencing using the Papain Worthington Kit (Worthington, LK003150). All solutions included in
the kit were prepared according to the manufacturer's instructions. 4-6 organoids were transferred
to one well of a low attachment 6 well plate and were washed with PBS without Ca^{2+} and Mg^{2+}.
Organoids were cut with a small scalpel into smaller pieces for easier dissociation. 800μl of papain
solution (supplied in the kit) was added per well. Samples were incubated at 37°C for about two
hours or until single cell suspension was achieved. Every 15 minutes, the mixture was pipetted
up and down with a cut P1000 pipette tip. Once single cell suspension was reached, 500 μl of
inhibitor solution (supplied in the kit) was added to the well. The solution was gently mixed, filtered
through a 70μm-pore sieve, and transferred to a 15 ml centrifuge tube. The cells were pelleted by
centrifugation at 300 x g for 5 minutes at room temperature. Cell pellets were resuspended in
500μl of ice-cold 0.04% BSA diluted in PBS without Ca^{2+} and Mg^{2+}. scRNA-seq was performed
on 4-6 pooled 6-month-old organoids per donor line, per condition (hCS or hSS) for a total of 48
organoids. A minimum of 10,000 dissociated cells were submitted for sequencing. The library was
prepared using the Chromium platform (10x Genomics) with the 3' gene expression (3' GEX)
V3/V3.1 kit. Libraries were sequenced on an Illumina NovaSeq sequencer with an S4 flow cell,
targeting a minimum depth of 20,000 reads per cell. The lengths for read parameters Read 1, i7
index, i5 index, and Read 2 were 100, 8, 0, 100, respectively.

vi. Bioinformatic analysis of scRNASeq data

The raw sequencing data, represented as base call (BCL) files produced by the Illumina
sequencing platform, were subjected to demultiplexing and subsequent conversion into FASTQ
format using CellRanger software (version 6.0.0, 10x Genomics) with default parameters. The
software then mapped the FASTQ reads to the human reference genome (GRCh38) with default
parameters. Following this, the 'count' command in the Cell Ranger (v6.0.0) software was utilized
for the quantification of gene expression. For alignment and counting, the reference genome
refdata-gex-GRCh38-2020-A was used, which was procured from the official 10x Genomics
website. We performed QC and normalization using the Seurat (v3) R package[75]. For QC, we
filtered out low-quality cells using the following criteria: (i) cells with unique gene counts outside
the range of 200 to 6000; (ii) cells with more than 30% mitochondrial gene content; and cells with
less than 2000 unique molecular identifiers[76,77]. Post QC, we carried out normalization, scaling
the gene expression measurements for each cell by the total expression, multiplied by a scale
factor (10,000 by default), and log-transformed the results. We extracted the expression profiles
of the 338 genes identified by Birey et al[33], to reduce the dimensionality of the dataset through
principal component analysis (PCA), and identify statistically significant PCs using a JackStraw
permutation test. This was followed by non-linear dimensional reduction using the UMAP (Uniform
Manifold Approximation and Projection) technique for visualization. Cells were clustered based
on their PCA scores using the Shared Nearest Neighbor (SNN) modularity optimization-based
clustering algorithm in Seurat. After dimensionality reduction, we used known marker genes to
guide the clustering of cells. Each cluster was then annotated using cell type markers identified
by Birey et al[33]. To account for variability among individual donors we pooled genotypes, similar
to our strategies in all other experiments, and performed a quasibinomial regression model to test
if cell frequencies significantly differed between genotypes. Finally, we conducted differential
expression analysis across the defined cell clusters using the FindAllMarkers function in Seurat,
which employs a Wilcoxon Rank-Sum test to control for the variability within and between groups.
The significantly differentially expressed genes were then used to interpret the biological
significance of cell clusters. For each identified cell type, we conducted an enrichment analysis
using the WebGestalt (WEB-based GEne SeT AnaLysis Toolkit) online tool, with the Human
genome (GRCh38) as the reference set, employing a hypergeometric statistical method and the
Benjamini & Hochberg method for multiple test adjustment.

vii. Analysis of ChIP-sequencing data from β-estradiol treated rodent brains.

Tracks in bigWig format were downloaded from the GEO dataset GSE144718[40]. Two peaks within

the *NRXN1* gene region in the mm10 genome were visualized using Spark (v2.6.2)[78]

viii. Generation of long-read sequencing data from post-mortem brain tissue

All aspects of sample processing (tissue handing, RNA extraction, library prep, QC, sequencing

and bioinformatic analysis) was performed as previously described[15].

Therapeutic Treatments of iNeurons

i. β-estradiol treatment

β-estradiol was reconstituted in DMSO and subsequently diluted in Brainphys maturation media

for a final concentration of 10 or 30nM. Neurons were treated for 3-5 consecutive days, with fresh

drug or vehicle control replenished daily (in half media changes). iGLUT and iGABA neurons were

treated starting from ~DIV14-18. On the final day of treatment, cells were harvested ~4 hours post

dosage. Chronic treatments for functional experiments extended daily β-estradiol replenishments

for ~14 days and withdrawn 2-3 days prior to patch-clamp recordings. For RNASeq, the same

chronic paradigm was performed and harvested on DIV21 ~4 hours post final dosage.

ii. Antisense oligonucleotide treatment

A single HPLC-grade ASO was designed from Qiagen (LNA GapmeR) against the mutant (Ex

20/24) splice junction containing a phosphorothioate modified backbone with or without a 5'-FAM

label for fluorescent visualization. All experiments were performed matched with a non-targeting

ASO as the control group. ASOs were delivered using Lipofectamine RNAiMAX Transfection

Reagent (Thermo, #13778075). The Lipofectamine RNAiMAX was diluted in Opti-MEM Medium

(Thermo, #31985070). ASO was then diluted in Opti-MEM Medium at 1µM. The diluted ASO was

added in a 1:1 ratio to the diluted Lipofectamine RNAiMAX and incubated at room temperature

for 5 minutes. The ASO-lipid complex was added to cells, and incubated for 72 hours post-

transfection, until RNA harvest. ASO samples for RNA-Seq were harvested on DIV17.

Molecular Biology and Imaging

i. RNA extraction and RT-qPCR

For the isolation of RNA, 2D cells were lysed using TRIzol (Life Technologies #15596026) and
purified using the Qiagen miRNeasy Kit (Qiagen Cat# 74106) according to the manufacturer's
instructions. For 3D organoids, pooled (early timepoints) or single organoids were washed and
lysed using TRIzol, by manual homogenization with a pestle in a 1.5mL centrifuge tube. Following
purification, samples were normalized within each experiment (15-50ng) and subsequently used
for RT-qPCR assays using the *Power* SYBR Green RNA-to-Ct 1-Step Kit (Thermo REF 4389986).
Relative transcript abundance was determined using the ΔΔCT method and normalized to the
ACTB housekeeping gene. All primer sequences are listed below.

shRNA, primers and oligonucleotide probe sequences

Target	Sequences (5'-3')	Supplier
RT-qPCR Primer Probes		
NRXN1 (wildtype) Forward	AGAAAGATGCCAAGCACCCA	ThermoFisher
NRXN1 (wildtype) Reverse	CCCATGTCCAGGAGGAGGTA	ThermoFisher
NRXN1-20/24 (mutant) Forward	GCTACCCTGCAGCCAACC	ThermoFisher
NRXN1-20/24 (mutant) Reverse	GACCATACCCGTGGTGCTG	ThermoFisher
ACTB Forward	TGTCCCCCAACTTGAGATGT	ThermoFisher
ACTB Reverse	TGTGCACTTTTATTCAACTGGTC	ThermoFisher
MAP2 Forward	AAACTGCTCTTCCGCTCAGACACC	ThermoFisher
MAP2 Reverse	GTTCACTTGGGCAGGTCTCCACAA	ThermoFisher
NEUROD1 Forward	GGTGCCTTGCTATTCTAAGACGC	ThermoFisher
NEUROD1 Reverse	GCAAAGCGTCTGAACGAAGGAG	ThermoFisher
SLC17A7 Forward	CGCATCATGTCCACCACCAACGT	ThermoFisher
SLC17A7 Reverse	GAGTAGCCGACCACCAACAGCAG	ThermoFisher

SLC17A6 Forward	TCAACAACAGCACCATCCACCGC	ThermoFisher
SLC17A6 Reverse	GTTTCCGGGTCCCAGTTGAATTTGG	ThermoFisher
GAD65 Forward	CTATGACACTGGAGACAAGGC	ThermoFisher
GAD65 Reverse	CAAACATTTATCAACATGCGCTTC	ThermoFisher
DLX5 Forward	ACAGAGACTTCACGACTCCCAG	ThermoFisher
DLX5 Reverse	TGTGGGGCTGCTCTGGTCTA	ThermoFisher

shRNA Sequences

NRXN1 wildtype mRNA (Exon 9)	ATGGAGTGGTGGCATTTAAAT	Sigma
NRXN1 mutant mRNA (Overlapping Exon 20/24)	CGCTACCCTGCAGCCAACCCA	Sigma

Antisense Oligonucleotides (LNA Gapmers)

Non-targeting ASO	AACACGTCTATACGC/36-FAM/	Qiagen
NRXN1-Splice ASO	GGTTGGCTGCAGGGTA/36-FAM/	Qiagen

ii. Bright Field Imaging

For organoid perimeter analyses, brightfield microscope images of organoids were taken with a
2x objective. Image analysis was performed in ImageJ, with best fitting ovals or ellipses were
selected around an organoid, and perimeter was measured.

iii. Immunostaining and imaging 2D cultures

For immunostaining of 2D monocultures, iGLUT and iGABA neurons seeded on acid-etched
coverslips coated with PEI buffered with boric acid and 4x Matrigel. Samples were washed with
DPBS $Ca^{2+}/Mg2^+$ and fixed using cold, fresh 16% paraformaldehyde (PFA, (Life Technologies,
#28908)), diluted to 4% for 12 minutes at RT. Coverslips were then blocked and permeabilized
with 2% donkey serum in DPBS $Ca^{2+}/Mg2^+$ supplemented with 0.1% Triton-X (Sigma, #93443-

100ML) (blocking buffer), for one hour at RT. Primary antibody solutions were prepared in
blocking buffer and incubated overnight at 4°C. The following day, samples were washed three
times with PBS, and treated with secondary antibodies diluted in blocking buffer, for 1 hour in a
dark chamber. Finally, samples were washed three times, and stained with DAPI for 15min at RT
during the final wash. Coverslips were mounted with antifade (Vectashield #H-1000-10) onto
glass slides and stored at 4°C until imaging using an upright Zeiss LSM 780 confocal microscope.

iv. High content imaging of 2D cultures

For high-content imaging, 100k cells/well were plated in an optically clear olefin bottom 96-well
plate (Revvity Health Sciences, #6055302). At WPI4 (iGLUT) and WPI5 (iGABA), cultures were
double fixed in PFA and 100% ice-cold methanol for 10min each with at least 2 washes of DPBS
$Ca^{2+}/Mg2^{+}$ between and after fixation steps. Fixed cultures were washed twice in PBS and
permeabilized and blocked two hours, followed by incubation with primary antibody solution
overnight at 4°C. Cultures were then washed 3 times with PBS and incubated with secondary
antibody solution for 1 hour at RT. Cultures were washed a further 3 timess with PBS with the
second wash containing 1µg/ml DAPI. Fixed cultures were then imaged on a CellInsight CX7 HCS
Platform with a 20x objective (0.4 NA) and neurite tracing and synaptic puncta detection
performed using the synaptogenesis module in the Thermo Scientific HCS Studio 4.0 Cell
Analysis Software to determine Syn1+ puncta density per µm of Map2+ve neurite length. 10 wells
were imaged per donor with 9 images acquired per well for neurite tracing analysis. Wells with
<10 annotated synapses, were excluded from the analysis.

Target	Antibody Dilution	Supplier/ CAT#
Primary Antibodies		
(Rabbit) vGAT	1:500	Synaptic Systems, #135-303
(Rabbit) vGLUT1	1:500	Synaptic Systems, #131-002
(Chicken) MAP2	1:1000	Abcam, #ab5392
(Mouse) Synapsin-1	1:500	Synaptic Systems, #106 011
DAPI	1:1000	Sigma, #D9542

Secondary Antibodies		
(Donkey) 488-Mouse	1:200	Jackson ImmunoResearch, # 711-545-152
(Donkey) 568-Rabbit	1:200	Abcam, #ab175700
(Donkey) 647-Chicken	1:500	Life technologies, #A10042
(Donkey) 568-Mouse	1:500	Life technologies, #A10037
(Donkey) 488-Rabbit	1:1000	Life Technologies, #A-21206

Statistics

No statistical power estimation analyses were used to predetermine sample sizes, where were

chosen to match previous publications[14,21] and field standards. All experimental statistics were

performed in Prism *v10.1.1* and R *v4.2.0.* Bioinformatic analyses were performed in R *v3.5.3* (Bulk

RNASeq DEG), *v4.1.2* (Bulk RNASeq GO) and *v4.1.2* (scRNAseq).

Data and Code Availability

All source donor hiPSCs have been deposited at the Rutgers University Cell and DNA Repository

(study 160; http://www.nimhstemcells.org/) and all bulk and single-cell transcriptome sequencing

data are being prepared for deposits to GEO. To facilitate improved reproducibility of our data

analyses, custom scripts has been deposited to github (https://github.com/mbfernando/NRXN1).

Source data will be provided with this manuscript.

Acknowledgements

MBF was supported by a Gilliam Fellowship from the Howard Hughes Medical Institute. This work

was supported by the National Institute of Mental Health grants RO1 MH121074-02 (KJB, GF and

PAS) and RO1 MH125579-02 (GF and KJB). DAK was supported by the National Science

Foundation under Grant No. DBI2146398. Any opinions, findings, and conclusions or

recommendations expressed in this material are those of the authors and do not necessarily

reflect the views of the National Science Foundation. We thank the Stem Cell Engineering Core

at the Icahn School of Medicine at Mount Sinai. We are grateful to the labs of Nan Yang (Ruiqi

Hu and Xiaoting Zhou) and Samuele Marro (Madel Durens) for assistance in primary glial
preparations. We are especially thankful to Kayla G. Townsley and Mark G. Baxter for advice on
statistical testing, and Liang Yang for critical advice on scRNASeq analyses. We acknowledge
Daniel Weinberger and the Lieber Institute for Brain Development at Johns Hopkins School of
Medicine, for sharing post-mortem materials. Finally, the authors thank all members of the
Brennand, Slesinger and Fang labs for critical feedback and discussions throughout the course
of this work.

Author Information

MBF, SK, ANM, RO, CP, AP, SG performed and analyzed experiments supervised by PAS and
KJB. YZ and YF performed bioinformatic analyses supervised by GF. AT performed bioinformatic
analysis of alternative splicing estimates, supervised by DAK. SW produced virus for the
generation of iGABA neurons, and MD performed high-content imaging experiments. EKF
processed post-mortem tissue and generated long-read data. IAP assisted in statistical analyses.
MBF, GF, PAS and KJB wrote the paper with input from all authors.

Ethics declarations / Competing interest statement.

All authors have no competing interests to declare.

Figures

Figure 1. *Aberrant NRXN1 splicing across human iPSC derived glutamatergic (iGLUT) and*
GABAergic (iGABA) neurons. (**a**) Brief description of clinical information of all hiPSC lines used
in this study, and (**b**) schematic of *NRXN1* gene structure as a splicegraph, denoting splice sites
(SS1-6), with red and blue shades corresponding on 3'-Del and 5'-Del genotypes, respectively.

Arrows indicate relative internal promoter positions. (**c, h**) Induction timeline and factors to

generate iGLUT and iGABA neurons, with immunostaining validation of neuronal identity (MAP2),

glutamate identity (vGLUT1), and GABA identity (vGAT). (**d, i**). Gene expression panel confirming

abundance of neuronal markers and neurotransmitter identity in iGLUT (n: Control = 6/2; 5'-Del =

6/2; 3'-Del = 6/2 | 1 batch) and in iGABA neurons (n: Control = 5/2; 5'-Del = 6/2; 3'-Del = 6/2 | 1

batch). (**e, j**) Mapped percent of alpha *NRXN1* exon reads in iGLUT, compared via a 1-way

ANOVA, with Dunnett's test ($F_{2, 15}$ = 25.70; 5'-Del p = 0.0031, 3'-Del p = 0.0086), and iGABA

neurons ($F_{2, 14}$ = 92.92; 5'-Del p < 0.001, 3'-Del p = 0.0004), with beta *NRXN1* exon reads,

calculated by subtracting alpha-specific reads against total reads. (**f, k**) Splicegraphs displaying

significant gene wide splicing clusters, compared via Dirichlet-multinomial generalized linear

model with Bonferroni corrections, for 5'-Del iGLUT (SS4 and SS5 cluster *p* = 0.006) and iGABA

neurons ($\beta \rightarrow \alpha$ cluster *p* = 0.0146), and for (**g, l**) 3'-Del iGLUT (SS4 and SS5 cluster *p* = 1.09E-

19) and iGABA neurons (SS4 and SS5 cluster *p* = 6.39E-23). n reported as samples/donors |

independent   batches.

Figure 2: *Transcriptomic impact of NRXN1$^{+/-}$ in induced (iGLUT/iGABA) and organoid-derived*
hCO-glutamatergic and hSO-GABAergic neurons. (**a, c**) Volcano plots of differential gene
expression (DE) analysis across both genotypes in iGLUT and (**j, l**) iGABA neurons. Vertical
dashed lines represent DE genes ±1.5 Log2FC. Horizontal dashed lines represent FDR = 0.1
cutoff (lower) and Bonferroni corrected cutoff (upper). (**b, d**) Sunburst plots of all FDR corrected
DEGs with SynGO annotated synapse function for iGLUT and (**k, m**) iGABA neurons. (**e, n**)
Timeline of neural organogenesis for hCO and hSOs (**f, o**), UMAPs of hCO and hSO organoid
samples sequenced at 6 months, annotated by cell clusters, and (**g, p**) relative proportions of cell
clusters across genotypes, (hCO = 47,460 cells) and (hSO = 35,563 cells). (**h, q**) validation of
regionalization across forebrain (*FOXG1*), dorsal (*EMX1*) and ventral (*DLX2*) regions, with
NRXN1 expression across all cells. (**i, r**) Gene ontological analysis results using DEGs from
scRNASeq. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, Wilcoxon's rank sum test, FDR = 0.05. Data
represented as mean ± sem. n reported as samples/donors | independent batches.

Figure 3. *Spontaneous, passive, and excitable properties are minimally changed from NRXN1$^{+/-}$* *induced neurons.* (**a, f**) Timelapse of multi-electrode array recordings every 2-3 days apart starting near ~DIV12 for a single representative induction. Tiles represent averaged wMFR values across genotypes during a single recording session. (**b**) MEA quantification of iGLUT neuronal activity, compared via a 1-way ANOVA, with Dunnett's test (n: Control = 52/2; 5'-Del = 42/2; 3'-Del = 38/2 | 3 batches) at WPI4 ($F_{2, 129}$ = 12.77; 5'-Del p = 0.0005, 3'-Del p < 0.0001), and (**c**) WPI6 ($F_{2, 129}$ = 5.737; 5'-Del p = 0.0129, 3'-Del p = 0.0069). (**d**) intrinsic properties of iGLUT neurons (n: Control = 51/2; 5'-Del = 51/2; 3'-Del = 48/2 | 8 batches): compared via 1-way ANOVA, with Dunnett's test, Capacitance ($F_{2, 147}$ = 1.505; 5'-Del p = 0.1565, 3'-Del p = 0.7633), Input resistance ($F_{2, 147}$ = 1.311; 5'-Del p = 0.1949, 3'-Del p = 0.7906), and RMP (n: Control = 16/2; 5'-Del = 14/2; 3'-Del = 16/2 | 2 batches) compared via 1-way ANOVA, with Dunnett's test ($F_{2, 43}$ = 0.8234; 5'-Del p = 0.6378, 3'-Del p = 0.8468). (**e**) Input-output curves of excitable properties (n: Control = 16/2; 5'-Del = 14/2; 3'-Del = 16/2 | 2 batches), with representative traces (right), compared via Step x Genotype 2-way ANOVA; Dunnett's Test ($F_{26, 559}$ = 1.690, 5'-Del p < 0.01 at Step 6, 7 and 9, 3'-Del p = n.s. on all steps). (**g**) MEA quantification of iGABA neuronal activity, compared via a 1-way ANOVA, with Dunnett's test (n: Control = 71/2; 5'-Del = 48/2; 3'-Del = 55/2 | 5 batches) at WPI2 ($F_{2, 171}$ = 7.805; 5'-Del p = 0.0029, 3'-Del p = 0.0015), and (**h**) WPI5 (n: Control = 27/4; 5'-Del = 23/2; 3'-Del = 23/2 | 2 batches) compared via 1-way ANOVA, with Dunnett's test ($F_{2, 70}$ = 3.158; 5'-Del p = 0.0323, 3'-Del p = 0.1824). (**i**) intrinsic properties of iGABA neurons (n: Control = 39/2; 5'-Del = 34/2; 3'-Del = 37/2 | 6 batches): compared via 1-way ANOVA, with Dunnett's test, Capacitance ($F_{2, 107}$ = 0.04256; 5'-Del p = 0.9425, 3'-Del p = 0.9941), Input resistance (n: Control = 39/2; 5'-Del = 34/2; 3'-Del = 36/2 | 6 batches, $F_{2, 106}$ = 1.451; 5'-Del p = 0.8162, 3'-Del p = 0.1703) and RMP (n: Control = 18/2; 5'-Del = 16/2; 3'-Del = 16/2 | 2 batches) compared via 1-way ANOVA, with Dunnett's test ($F_{2, 47}$ = 0.02842; 5'-Del p = 0.9893, 3'-Del p = 0.9601). (**j**) Input-output curves of excitable properties (n: Control = 16/2; 5'-Del = 14/2; 3'-Del = 16/2 | 2 batches), with representative traces (right), compared via Step x Genotype 2-way ANOVA; Dunnett's Test ($F_{26, 572}$ = 0.5137, 5'-Del and 3'-Del p = n.s. on all steps). Data represented as mean ± sem. n reported as samples/donors | independent batches.

Figure 4: *Divergent impact on neurotransmission from NRXN1$^{+/-}$ induced neurons.* (**a**) Representative traces of iGLUT sEPSCs. (**b**) Cumulative probabilities and log-scaled cell averages of inter-event-internals (IEIs) across genotypes (n: Control = 39/4; 5'-Del = 33/3; 3'-Del = 29/2 | 6), compared by Levene's Test with Bonferroni correction for averaged distributions (5'-Del F = 46.635, df = 1, *p* = 2.54E-11; 3'-Del F = 17.928, df = 1, *p* = 4.88E-5), and 1-way ANOVA, with Dunnett's test for inset ($F_{2, 98}$ = 3.117; 5'-Del *p* = 0.0285, 3'-Del *p* = 0.2929). (**c**) Cumulative probabilities and log-scaled cell averages of amplitude size across genotypes, compared by Levene's Test with Bonferroni correction for averaged distributions (5'-Del F = 14.15, df = 1, *p* = 3.88E-4; 3'-Del F = 1.0851, df = 1, *p* = 0.5964), and 1-way ANOVA, with Dunnett's test for inset ($F_{2, 98}$ = 2.839; 5'-Del *p* = 0.0634, 3'-Del *p* = 0.9994). (**d**) Transformed p-values of Levene's Test and (**e**) SynGO gene-set averaged log$_2$FC values across pre- or post- synaptic genes. (**f**) Gene expression panel (z-scores), of canonical *NRXN1* binding partners in iGLUT neurons, boxes indicate reaching genome wide significance. (**g**) Representative images of iGLUT synaptic puncta traced (SYN1) onto dendrites (MAP2), and (**h**) normalized fold change of SYN1 puncta to length of MAP2 ratio (Syn:Neu), (n: Control = 40/2; 5'-Del: 39/2; 3'-Del: 40/2), compared via 1-way ANOVA, with Dunnett's test ($F_{2, 116}$ = 7.538, 5'-Del *p* = 0.0205, 3'-Del *p* = 0.0005). (**i**) Representative traces of iGABA sIPSCs. (**j**) Cumulative probabilities and log-scaled cell averages of inter-event-internals (IEIs) across genotypes (n: Control = 26/2; 5'-Del = 25/3; 3'-Del = 22/2 | 4), compared by Levene's Test with Bonferroni correction for averaged distributions (5'-Del F = 3.4002, df = 1, *p* = 0.1306; 3'-Del F = 16.501, df = 1, *p* = 1.00E-04), and 1-way ANOVA, with Dunnett's test for inset ($F_{2, 70}$ = 0.8296; 5'-Del *p* = 0.7986, 3'-Del *p* = 0.339). (**k**) Cumulative probabilities and log-scaled cell averages of amplitude size across genotypes, compared by Levene's Test for averaged distributions (5'-Del F = 0.1, df = 1, *p* = 0.9204; 3'-Del F = 0.0006, df = 1, *p* = 0.9801), and 1-way ANOVA, with Dunnett's test for inset ($F_{2, 70}$ = 0.08143; 5'-Del *p* = 0.9431, 3'-Del *p* = 0.9904). (**l**) Transformed p-values of Levene's Test and (**m**) SynGO gene-set averaged log$_2$FC values across pre- or post- synaptic genes. (**n**) Gene expression panel (z-scores), of canonical *NRXN1* binding partners in iGABA neurons, boxes indicate reaching genome wide significance. (**o**) Representative images of iGABA synaptic puncta traced (SYN1) onto dendrites (MAP2), and (**p**) normalized fold change of SYN1 puncta to length of MAP2 ratio (Syn:Neu), (n: Control = 33/2; 5'-Del: 36/2; 3'-Del: 40/2), compared via 1-way ANOVA, with Dunnett's test ($F_{2, 101}$ = 12.59, 5'-Del *p* < 0.0001, 3'-Del *p* = 0.0271). **p* < 0.05, ***p* < 0.01, ****p* <0.001, Wilcoxon's rank sum test, FDR = 0.05. Data represented as mean ± sem. n reported as samples/donors | independent batches.

Figure 5: *Isogenic recapitulation and rescue of neurotransmission phenotypes.* (**a**) Differential splicing of β→α cluster in shWT, compared via Dirichlet-multinomial generalized linear model (*p* = 0.0083) in iGLUT neurons. (**b**) Representative traces of iGLUT WT knockdown effects, with (**c**) cumulative probabilities of sEPSC IEI distributions, with insets of cell-averaged IEI and amplitude measures (n: shNT = 12/1; shWT = 7/1 | 2 batches). Curves were compared via a Levene's test (F = 70.78, df =1, *p* < 2.2E-16), and insets were compared via a Student's t-test (IEI t=1.166, df=19, *p* = 0.2579; AMP t=0.5661, df=19, *p* = 0.578). (**d**) SynGO (biological process) sunburst plots showing enrichment of DEGs associated with synaptic function for iGLUT WT-KD compared to a brain expressed background via Fisher's exact test (1.274161-fold, *p* = 0.1049). (**e**) Splicing of MT exon 20→24 cluster in shMT, compared via Dirichlet-multinomial generalized linear model (*p* = 0.0623). (**f**) Representative traces of iGLUT MT knockdown effects, with (**g**) cumulative probabilities of sEPSC IEI distributions, with insets of cell-averaged IEI and amplitude measures (n:shNT = 14/1; shMT = 18/1 | 2 batches). Curves were compared via a Levene's test (F = 230.6; df = 1, *p* < 2.2E-16), and insets were compared via a Student's t-test (IEI t=2.471, df=30, *p* = 0.0194; AMP t=1.812, df=30, *p* = 0.08). (**h**) SynGO (biological process) sunburst plots showing enrichment of DEGs associated with synaptic function for iGLUT MT-KD compared to a brain expressed background via Fisher's exact test (1.214533-fold, *p* = 0.4541). (**i**) Differential splicing of β→α cluster in shWT, compared via Dirichlet-multinomial generalized linear model (*p* = 0.0203) in iGABA neurons. (**j**) Representative traces of iGABA WT knockdown effects, with (**k**) cumulative probabilities of sIPSC IEI distributions, with insets of cell-averaged, IEI and amplitude measures (shNT = 16/3; shWT = 19/3 | 2 batches). Curves were compared via a Levene's test (F = 30.879, df =1, *p* < 3.66E-08), and insets were compared via a Student's t-test (IEI t=2.768, df=33, *p* = 0.0092; AMP t=0.7856, df=33, *p* = 0.4377). (**l**) SynGO (biological process) sunburst plots for iGABA WT-KD compared to a brain expressed background via Fisher's exact test (1.234861-fold, *p* = 1.307E-6). (**m**) Splicing of MT exon 20→24 cluster in shMT, compared via Dirichlet-multinomial generalized linear model (*p* = 0.0258) in iGABA neurons. (**n**) Representative traces of iGABA MT knockdown effects, with (**o**) cumulative probabilities of sIPSC IEI distributions, with insets of cell-averaged, IEI and amplitude measures (shNT = 16/1; shMT = 19/1 | 2 batches). Curves were compared via a Levene's test (F = 4.1324; df = 1, *p* < 0.04226), and insets were compared via a Student's t-test (IEI t=1.305, df=25, *p* = 0.2038; AMP t=0.9292, df=25, *p* = 0.3617). (**p**) SynGO (biological process) sunburst plots for iGABA WT-KD compared to a brain expressed background via Fisher's exact test (1.347225-fold, *p* = 2.694E-4). Data represented as mean ± sem. n reported as samples/donors | independent batches.

Figure 6: *Precise therapeutic targeting of stratified GOF- and LOF-NRXN1$^{+/-}$ in iGLUT neurons.*
(**a**) Model and proposed mechanism of rescue for LOF patients ameliorating loss of wildtype isoforms. (**b**) RT-qPCR of WT *NRXN1* gene expression in 5-Del patients, post-acute treatment (3-5 days) (n: DMSO = 19/2; β-estradiol = 18/2 | 3 batches) compared via Student's t-test t=2.293, df=35, *p* = 0.028. (**c**) Quantification of iGLUT neuronal activity at WPI3 across vehicle and treatment (n: DMSO = 72/3; β-estradiol = 74/3 | 3 batches) compared via Student's t-test (t=2.804, df=144, *p* = 0.0057). (**d**) Splicing of β→α cluster and (**e**) volcano plot of DEGs in β-estradiol treated 5'-Del iGLUT neurons, compared to vehicle treated 5'-Del iGLUT neurons. (**f**) Representative patch-clamp traces and (**h**) cumulative probabilities of sEPSC IEI distributions, with insets of cell-averaged, IEI and amplitude measures (n: Control + DMSO = 18/1; 5'-Del + DMSO = 33/2 5'-Del + β-estradiol = 35/2). Curves were compared via Levene's test with Bonferroni corrections (Control + DMSO v. 5'-Del + DMSO F = 13.151; df = 1, *p* = 8.763E-4; 5'-Del + DMSO v 5'-Del + β-estradiol F = 14.359; df = 1, *p* = 4.62E-04; Control + DMSO v 5'-Del + β-estradiol, F = 0.0141; df = 1, *p* = 0.955). Insets were compared via Student's t-test (IEI t=1.161, df=66, *p* = 0.2499). (**h**) Model and proposed mechanism of rescue for GOF patients expressing mutant isoforms. (**i**) Schematic of a *NRXN1* ASO matrix, with the selective sequence targeting 20/24 splice junction. (**j**) RT-qPCR of MT *NRXN1* gene expression in 3'-Del patients' post-acute treatment for 72hrs (ASO-NT = 8/2; ASO-MT = 10/2 | batches),compared via Student's t-test t=12.91, df=16 *p* < 0.0001. (**k**) Splicing of MT Exon 20→24 cluster and (**l**) volcano plot of DEGs in ASO-MT treated 3'-Del iGLUT neurons, compared to ASO-NT treated 3'-Del iGLUT neurons.(**m**) Biological process and (**n**) cellular compartment GO terms demonstrating enrichment (-log$_{10}$[adj *p* value]) of selected synapse related pathways. Data represented as mean ± sem. n reported as samples/donors | independent batches.

 Extended Data Figures

Extended Data Figure 1: (**a**) Splicegraph displaying significant gene wide splicing clusters at

NRXN1 SS3 (p = 0.0196), compared via Dirichlet-multinomial generalized linear model with

Bonferroni corrections. (**b, f**) Gene expression fold-change of select *NRXN1* predicted RNA-

binding proteins (RBP) across patients and control iGLUT and iGABA neurons, with statistical

comparisons for STAR-Family RBPs. (**c, g**) Overlap of DEGs and (**d,h**) gene set enrichment

analysis (GSEA) between genotypes. (**e,i**) Distinct gene expression patterns by hierarchal

clustering of all patient specific DEGs. Sample information correspond to Fig. 1

	iGLUT 5'-Del		iGLUT 3'-Del		iGABA 5'-Del		iGABA 3'-Del	
	Overlapped DEGs with risk genes	Overlapped DEGs with risk genes & in SIGNOR database	Overlapped DEGs with risk genes	Overlapped DEGs with risk genes & in SIGNOR database	Overlapped DEGs with risk genes	Overlapped DEGs with risk genes & in SIGNOR database	Overlapped DEGs with risk genes	Overlapped DEGs with risk genes & in SIGNOR database
ASD	11	**7**	4	**0**	3	**1**	6	**2**
BP	16	**9**	3	**1**	2	**1**	5	**1**
SCZ	26	**14**	10	**1**	9	**3**	14	**3**
Total	37	**22**	12	**2**	9	**3**	18	**4**

iGLUT

iGABA

Extended Data Figure 2: *Extended transcriptomics analysis on disease risk associated genes.* (a) Summary table of overlapping DEGs with risk enrichments across publicly curated datasets for autism (ASD), bipolar disorder (BD) and schizophrenia. (**b, e**) Enrichment of genes across neuropsychiatric disorders for iGLUT and iGABA neurons. (**c**) Interaction maps of risk genes for 5'-Del iGLUT, (**d**) 3'-Del iGLUT, (**f**) 5'-Del iGABA and (**g**) 5'-iGABA. Sample information correspond to Fig. 1

human Cortical Organoid (hCO)

c 4 months RT-qPCR validation panel

g 6 months scRNASeq gene expression panel

human Subpallial Organoid (hSO)

j 4 months RT-qPCR validation panel

n 6 months scRNASeq gene expression panel

Extended Data Figure 3: *Extended data on human organoid generation and characterization.*
(**a,h**) Representative images of hiPSC aggregation and immature spheroids post dislodging. (**b,i**)
Normalized organoid perimeters over time (compared to averaged control), hCO (n = 6 donors |
2 batches | 72-161 organoids) and hSO (n = 6 donors | 2 batches | 46-134 organoids). (**c, j**) RT-
qPCR results from 4-month organoids of genes for pluripotency, neuronal, and cell-type specific
markers. (d-e, k-l) RT-qPCR results of *NRXN1* WT and MT expression hCO (n = 6 donors | 1
representative batch | 12 samples) and hSO (n = 6 donors | 1 representative batch | 12 samples).
Statistical tests used were 1-way ANOVAs with Dunnett's test. (f, m) relative proportions of cell
clusters across individual donors, (hCO = 47,460 cells) and (hSO = 35,563 cells). (**g, n**)
Comprehensive gene expression panel across sub-clusters of hCO and hSO samples across
neuronal, cortical, subpallial and astroglia markers. Data corresponds to Fig 2.

1003
Extended Data Figure 4: *Extended data on electrophysiological properties of 5'-Del and 3'-Del*
neurons. (**a**) Voltage-gated potassium and channel kinetics across genotypes for iGLUT neurons
(n = 6 donors | 2 inductions | 45 neurons). (**b**) Comparative mEPSC kinetics of IEI and (**c**)

amplitude size from iGLUT neurons (n= 6 donors | 4 inductions | 47 neurons), compared via a 1-

way ANOVA with Dunnett's test. (**d**) Voltage-gated potassium and channel kinetics across

genotypes for iGABA neurons (n = 6 donors | 2 inductions | 34 neurons). (**e**) Comparative mIPSC

kinetics of (a) IEI and (**f**) amplitude size from iGABA neurons (n= 6 donors | 3 inductions | 27

neurons), compared via a 1-way ANOVA with Dunnett's test. See Supplementary Table 5 for all

summary statistics.

1013

1014

1015   **Extended Data Figure 5:** *Extended KCC2 related data from immature GABA neurons.* (a)

1016   Transcriptomic comparison of *SLC12A5* expression across DIV14 and DIV35 RNASeq

1017   timepoints. (b,c) MEA tests from pre- and post- treatment of 10uM GABAzine. (n = 2 donors | 1

1018   representative induction | 28 MEA wells) Statistical tests are paired student's t-test for time-linked

1019   comparison and unpaired student's t-test for pre/post activity foldchange.

1020

1021

1022

1023   **Extended Data Figure 6:** *shRNA knockdown validation.* (a) Extent of shRNA knockdown on WT

1024   and (b) MT *NRXN1* expression in iGLUT neurons (n = 2 donors | 1-3 inductions). (c) Extent of

1025   shRNA knockdown on WT and (d) MT *NRXN1* expression in iGABA neurons, (n = 2-3 donors | 1-

1026   3 inductions) Statistical tests used were Student's t-test.

1027

1028   **Extended Data Figure 7:** *ChIP-seq enrichment of ER1 binding at NRXN1 loci in rodent brain.* (**a**)

1029   Female and (**b**) male mus musculus ChIP tracts of *NRXN1* locus, with red dashed areas

1030 highlighting binding enrichment across vehicle and estradiol treated mice. (**c**) Effect of beta-
1031 estradiol on control donors (n = 16/4 | Representative).

1032
1033
Extended Data Figure 8: *In-vivo validation of MT isoform expression from an unrelated autism*
NRXN1$^{+/-}$ patient. (a) Schematic of novel NRXN1 autism patient, and GOF therapeutic targeting
pipeline, with (b) schematic of the *NRXN1α* isoform structures, with each row representing a
unique *NRXN1α* isoform and each column representing a *NRXN1* exon. The colored isoforms
(navy, wildtype; peach, patient-specific) are spliced into the transcript while the blank exons are
spliced out. (c) The abundance of each *NRXN1α* isoform by sample.
1040

1041

1042
1043
1044

1045

1046

References

References

1. Sudhof, T. C. Synaptic Neurexin Complexes: A Molecular Code for the Logic of Neural

Circuits. *Cell* **171**, 745–769 (2017).

2. De Wit, J. & Ghosh, A. Specification of synaptic connectivity by cell surface interactions.

Nat. Rev. Neurosci. **17**, 4–4 (2016).

3. Gomez, A. M., Traunmüller, L. & Scheiffele, P. Neurexins: molecular codes for shaping

neuronal synapses. *Nat. Rev. Neurosci.* **22**, 137–151 (2021).

4. Schreiner, D. *et al.* Targeted Combinatorial Alternative Splicing Generates Brain Region-

Specific Repertoires of Neurexins. *Neuron* **84**, 386–398 (2014).

5. Fuccillo, M. V. *et al.* Single-Cell mRNA Profiling Reveals Cell-Type-Specific Expression of

Neurexin Isoforms. *Neuron* **87**, 326–340 (2015).

6. Marshall, C. R. *et al.* Contribution of copy number variants to schizophrenia from a genome-

wide study of 41,321 subjects. *Nat Genet* **49**, 27–35 (2017).

7. Matsunami, N. *et al.* Identification of Rare Recurrent Copy Number Variants in High-Risk

Autism Families and Their Prevalence in a Large ASD Population. *PLoS ONE* **8**, e52239

(2013).

8. Moller, R. S. *et al.* Exon-disrupting deletions of NRXN1 in idiopathic generalized epilepsy.

Epilepsia **54**, 256–64 (2013).

9. Ching, M. S. L. *et al.* Deletions of *NRXN1* (neurexin-1) predispose to a wide spectrum of

developmental disorders. *Am. J. Med. Genet. B Neuropsychiatr. Genet.* **153B**, 937–947

(2010).

10. Huang, A. Y. *et al.* Rare Copy Number Variants in NRXN1 and CNTN6 Increase Risk for

Tourette Syndrome. *Neuron* **94**, 1101-1111 e7 (2017).

11. Grayton, H. M., Missler, M., Collier, D. A. & Fernandes, C. Altered Social Behaviours in

Neurexin 1α Knockout Mice Resemble Core Symptoms in Neurodevelopmental Disorders.

PLoS ONE **8**, e67114 (2013).

12. Pak, C. *et al.* Cross-platform validation of neurotransmitter release impairments in

schizophrenia patient-derived *NRXN1* -mutant neurons. *Proc. Natl. Acad. Sci.* **118**,

e2025598118 (2021).

13. Pak, C. *et al.* Human Neuropsychiatric Disease Modeling using Conditional Deletion

Reveals Synaptic Transmission Defects Caused by Heterozygous Mutations in NRXN1. *Cell*

Stem Cell **17**, 316–328 (2015).

14. Sebastian, R. *et al.* Schizophrenia-associated NRXN1 deletions induce developmental-

timing- and cell-type-specific vulnerabilities in human brain organoids. *Nat. Commun.* **14**,

3770 (2023).

15. Flaherty, E. *et al.* Neuronal impact of patient-specific aberrant NRXN1alpha splicing. *Nat*

Genet **51**, 1679–1690 (2019).

16. Boxer, E. E. & Aoto, J. Neurexins and their ligands at inhibitory synapses. *Front. Synaptic*

Neurosci. **14**, 1087238 (2022).

17. Taoufik, E., Kouroupi, G., Zygogianni, O. & Matsas, R. Synaptic dysfunction in

neurodegenerative and neurodevelopmental diseases: an overview of induced pluripotent

stem-cell-based disease models. *Open Biol.* **8**, 180138 (2018).

18. Ho, S.-M. *et al.* Rapid Ngn2-induction of excitatory neurons from hiPSC-derived neural

progenitor cells. *Methods* **101**, 113–124 (2016).

19. Zhang, Y. *et al.* Rapid single-step induction of functional neurons from human pluripotent

stem cells. *Neuron* **78**, 785–98 (2013).

20. Barretto, N. *et al.* ASCL1- and DLX2-induced GABAergic neurons from hiPSC-derived

NPCs. *J Neurosci Methods* **334**, 108548 (2020).

21. Yang, N. *et al.* Generation of pure GABAergic neurons by transcription factor programming.

Nat Methods (2017) doi:10.1038/nmeth.4291.

22. Li, Y. I. *et al.* Annotation-free quantification of RNA splicing using LeafCutter. *Nat Genet* **50**,

151–158 (2018).

23. Iijima, T. *et al.* SAM68 regulates neuronal activity-dependent alternative splicing of neurexin-

1. *Cell* **147**, 1601–14 (2011).

24. Iijima, T., Iijima, Y., Witte, H. & Scheiffele, P. Neuronal cell type-specific alternative splicing

is regulated by the KH domain protein SLM1. *J Cell Biol* **204**, 331–42 (2014).

25. Traunmuller, L., Gomez, A. M., Nguyen, T. M. & Scheiffele, P. Control of neuronal synapse

specification by a highly dedicated alternative splicing program. *Science* **352**, 982–6 (2016).

26. Traunmüller, L. *et al.* A cell-type-specific alternative splicing regulator shapes synapse

properties in a trans-synaptic manner. *Cell Rep.* **42**, 112173 (2023).

27. Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing and

microarray studies. *Nucleic Acids Res.* **43**, e47–e47 (2015).

28. Koopmans, F. *et al.* SynGO: An Evidence-Based, Expert-Curated Knowledge Base for the

Synapse. *Neuron* **103**, 217-234.e4 (2019).

29. Wu, T. *et al.* clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. *The*

Innovation **2**, 100141 (2021).

30. Seah, C. *et al.* Modeling gene × environment interactions in PTSD using human neurons

reveals diagnosis-specific glucocorticoid-induced gene expression. *Nat. Neurosci.* **25**,

1434–1445 (2022).

31. Lo Surdo, P. *et al.* SIGNOR 3.0, the SIGnaling network open resource 3.0: 2022 update.

Nucleic Acids Res. **51**, D631–D637 (2023).

32. Schafer, S. T. *et al.* Pathological priming causes developmental gene network

heterochronicity in autistic subject-derived neurons. *Nat. Neurosci.* **22**, 243–255 (2019).

33. Birey, F. *et al.* Assembly of functionally integrated human forebrain spheroids. *Nature* **545**,

54–59 (2017).

34. Yoon, S.-J. *et al.* Reliability of human cortical organoid generation. *Nat. Methods* **16**, 75–78

(2019).

35. Khan, T. A. *et al.* Neuronal defects in a human cellular model of 22q11.2 deletion syndrome.

Nat. Med. **26**, 1888–1898 (2020).

36. Zeng, L. *et al.* Functional Impacts of NRXN1 Knockdown on Neurodevelopment in Stem Cell

Models. *PLoS ONE* **8**, e59685 (2013).

37. Virtanen, M. A., Uvarov, P., Mavrovic, M., Poncer, J. C. & Kaila, K. The Multifaceted Roles

of KCC2 in Cortical Development. *Trends Neurosci.* **44**, 378–392 (2021).

38. Sohal, V. S. & Rubenstein, J. L. R. Excitation-inhibition balance as a framework for

investigating mechanisms in neuropsychiatric disorders. *Mol. Psychiatry* **24**, 1248–1257

(2019).

39. Willsey, H. R. *et al.* Parallel in vivo analysis of large-effect autism genes implicates cortical

neurogenesis and estrogen in risk and resilience. *Neuron* **109**, 1409 (2021).

40. Gegenhuber, B., Wu, M. V., Bronstein, R. & Tollkuhn, J. Gene regulation by gonadal

hormone receptors underlies brain sex differences. *Nature* **606**, 153–159 (2022).

41. Roberts, T. C., Langer, R. & Wood, M. J. A. Advances in oligonucleotide drug delivery. *Nat.*

Rev. Drug Discov. **19**, 673–694 (2020).

42. Saito, Y. *et al.* Differential NOVA2-Mediated Splicing in Excitatory and Inhibitory Neurons

Regulates Cortical Development and Cerebellar Function. *Neuron* **101**, 707-720.e5 (2019).

43. Paz, I., Kosti, I., Ares, M., Cline, M. & Mandel-Gutfreund, Y. RBPmap: a web server for

mapping binding sites of RNA-binding proteins. *Nucleic Acids Res.* **42**, W361–W367 (2014).

44. Ghanbarian, H., Aghamiri, S., Eftekhary, M., Wagner, N. & Wagner, K.-D. Small Activating

RNAs: Towards the Development of New Therapeutic Agents and Clinical Treatments. *Cells*

10, 591 (2021).

45. Subramanian, A. *et al.* A Next Generation Connectivity Map: L1000 Platform and the First

1,000,000 Profiles. *Cell* **171**, 1437-1452.e17 (2017).

46. Yilmaz, C. *et al.* Neurosteroids as regulators of neuroinflammation. *Front. Neuroendocrinol.*

55, 100788 (2019).

47. Brunet De Courssou, J.-B., Durr, A., Adams, D., Corvol, J.-C. & Mariani, L.-L. Antisense

therapies in neurological diseases. *Brain* **145**, 816–831 (2022).

48. Ingusci, S., Verlengia, G., Soukupova, M., Zucchini, S. & Simonato, M. Gene Therapy Tools

for Brain Diseases. *Front. Pharmacol.* **10**, 724 (2019).

49. Lin, H.-C. *et al.* NGN2 induces diverse neuron types from human pluripotency. *Stem Cell*

Rep. **16**, 2118–2127 (2021).

50. Zafra, F. & Piniella, D. Proximity labeling methods for proteomic analysis of membrane

proteins. *J. Proteomics* **264**, 104620 (2022).

51. Bell, J. Stratified medicines: towards better treatment for disease. *The Lancet* **383**, S3–S5

(2014).

52. Tsimberidou, A. M. *et al.* Molecular tumour boards — current and future considerations for

precision oncology. *Nat. Rev. Clin. Oncol.* **20**, 843–863 (2023).

53. Zhang, H., Colclough, K., Gloyn, A. L. & Pollin, T. I. Monogenic diabetes: a gateway to

precision medicine in diabetes. *J. Clin. Invest.* **131**, e142244 (2021).

54. Sullivan, P. F. & Geschwind, D. H. Defining the Genetic, Genomic, Cellular, and Diagnostic

Architectures of Psychiatric Disorders. *Cell* **177**, 162–183 (2019).

55. Gerasimavicius, L., Livesey, B. J. & Marsh, J. A. Loss-of-function, gain-of-function and

dominant-negative mutations have profoundly different effects on protein structure. *Nat.*

Commun. **13**, 3895 (2022).

56. Wang, L. *et al.* Analyses of the autism-associated neuroligin-3 R451C mutation in human

neurons reveal a gain-of-function synaptic mechanism. *Mol. Psychiatry* **29**, 1620–1635

(2024).

57. Pinggera, A. *et al.* New gain-of-function mutation shows CACNA1D as recurrently mutated

gene in autism spectrum disorders and epilepsy. *Hum. Mol. Genet.* **26**, 2923–2932 (2017).

58. Clark, M. B. *et al.* Long-read sequencing reveals the complex splicing profile of the

psychiatric risk gene CACNA1C in human brain. *Mol. Psychiatry* **25**, 37–47 (2020).

59. Brunklaus, A. *et al.* Gene variant effects across sodium channelopathies predict function

and guide precision therapy. *Brain* **145**, 4275–4286 (2022).

60. Sanders, S. J. *et al.* Progress in Understanding and Treating SCN2A-Mediated Disorders.

Trends Neurosci. **41**, 442–456 (2018).

61. Kim, G., Gautier, O., Tassoni-Tsuchida, E., Ma, X. R. & Gitler, A. D. ALS Genetics: Gains,

Losses, and Implications for Future Therapies. *Neuron* **108**, 822–842 (2020).

62. Balendra, R. & Isaacs, A. M. C9orf72-mediated ALS and FTD: multiple pathways to disease.

Nat. Rev. Neurol. **14**, 544–558 (2018).

63. Prakasam, R. *et al.* LSD1/PRMT6-targeting gene therapy to attenuate androgen receptor

toxic gain-of-function ameliorates spinobulbar muscular atrophy phenotypes in flies and

mice. *Nat. Commun.* **14**, 603 (2023).

64. Chintalaphani, S. R., Pineda, S. S., Deveson, I. W. & Kumar, K. R. An update on the

neurological short tandem repeat expansion disorders and the emergence of long-read

sequencing diagnostics. *Acta Neuropathol. Commun.* **9**, 98 (2021).

65. Rhee, H. J. *et al.* An Autaptic Culture System for Standardized Analyses of iPSC-Derived

Human Neurons. *Cell Rep.* **27**, 2212-2228.e7 (2019).

66. Sloan, S. A., Andersen, J., Pașca, A. M., Birey, F. & Pașca, S. P. Generation and assembly

of human brain region–specific three-dimensional cultures. *Nat. Protoc.* **13**, 2062–2085

(2018).

67. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).

68. Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for

assigning sequence reads to genomic features. *Bioinformatics* **30**, 923–930 (2014).

69. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR : a Bioconductor package for

differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140

(2010).

70. Hoffman, G. E. *et al.* Transcriptional signatures of schizophrenia in hiPSC-derived NPCs

and neurons are concordant with post-mortem adult brains. *Nat. Commun.* **8**, 2225 (2017).

71. Newman, A. M. *et al.* Determining cell type abundance and expression from bulk tissues

with digital cytometry. *Nat. Biotechnol.* **37**, 773–782 (2019).

72. Hoffman, G. E. & Schadt, E. E. variancePartition: interpreting drivers of variation in complex

gene expression studies. *BMC Bioinformatics* **17**, 483 (2016).

73. Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Near-optimal probabilistic RNA-seq

quantification. *Nat. Biotechnol.* **34**, 525–527 (2016).

74. De Marinis, I., Lo Surdo, P., Cesareni, G. & Perfetto, L. SIGNORApp: a Cytoscape 3

application to access SIGNOR data. *Bioinformatics* **38**, 1764–1766 (2022).

75. Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902.e21

(2019).

76. Urresti, J. *et al.* Cortical organoids model early brain development disrupted by 16p11.2

copy number variants in autism. *Mol. Psychiatry* **26**, 7560–7580 (2021).

77. Yang, X. *et al.* Identification and validation of genes affecting aortic lesions in mice. *J. Clin.*

Invest. **120**, 2414–2422 (2010).

78. Kurtenbach, S. & William Harbour, J. SparK: A Publication-quality NGS Visualization Tool.

Preprint at https://doi.org/10.1101/845529 (2019).

1219