TITLE

Precise Therapeutic Targeting of Distinct *NRXN1*[+/-] Mutations

AUTHORS

Michael B. Fernando[1,2,4,5], Yu Fan[3]*, Yanchun Zhang[3]*, Sarah Kammourh[3], Aleta N. Murphy[1,2,4],

Sadaf Ghorbani[5,6], Ryan Onatzevitch[2], Adriana Pero[3], Christopher Padilla[3], Erin K. Flaherty[1,2], Iya

K. Prytkova[2], Lei Cao[3], Sarah Williams[1-4], Gang Fang[3#], Paul A. Slesinger[2#] & Kristen J.

Brennand[2-5#]

AFFILIATIONS

[1] Graduate School of Biomedical Science, Icahn School of Medicine at Mount Sinai, New York,
NY 10029

[2] Nash Family Department of Neuroscience, Icahn School of Medicine at Mount Sinai, New York,
NY 10029

[3] Department of Genetics and Genomics, Icahn School of Medicine at Mount Sinai, New York, NY
10029

[4] Friedman Brain Institute, Black Family Stem Cell Institute, Pamela Sklar Division of Psychiatric
Genomics, Icahn School of Medicine at Mount Sinai, New York, NY 10029

[5] Department of Psychiatry, Yale School of Medicine, New Haven, CT, 06520

[6] Haukeland University Hospital, Bergen, Norway

*These authors contributed equally

[#]Correspondence: kristen.brennand@yale.edu, paul.slesinger@mssm.edu and
gang.fang@mssm.edu

KEYWORDS

Human induced pluripotent stem cells; *NRXN1*; alternative splicing; glutamatergic neurons;
GABAergic neurons; genomics; neuropsychiatric disorder; disease modeling; precision medicine

ABSTRACT

As genetic studies continue to identify risk loci that are significantly associated with risk for

neuropsychiatric disease, a critical unanswered question is the extent to which diverse mutations

--sometimes impacting the same gene-- will require tailored therapeutic strategies. Here we

consider this in the context of rare neuropsychiatric disorder-associated copy number variants

(2p16.3) resulting in heterozygous deletions in *NRXN1*, a pre-synaptic cell adhesion protein that

serves as a critical synaptic organizer in the brain. Complex patterns of *NRXN1* alternative

splicing are fundamental to establishing diverse neurocircuitry, vary between the cell types of the

brain, and are differentially impacted by unique (non-recurrent) deletions. We contrast the cell-

type-specific impact of patient-specific mutations in *NRXN1* using human induced pluripotent

stem cells, finding that perturbations in *NRXN1* splicing result in divergent cell-type-specific

synaptic outcomes. Via distinct loss-of-function (LOF) and gain-of-function (GOF) mechanisms,

NRXN1$^{+/-}$ deletions cause decreased synaptic activity in glutamatergic neurons, yet increased

synaptic activity in GABAergic neurons. Stratification of patients by LOF and GOF mechanisms

will facilitate individualized restoration of *NRXN1* isoform repertoires; towards this, antisense

oligonucleotides knockdown mutant isoform expression and alters synaptic transcriptional

signatures, while treatment with β-estradiol rescues synaptic function in glutamatergic neurons.

Given the increasing number of mutations predicted to engender both LOF and GOF mechanisms

in brain disease, our findings add nuance to future considerations of precision medicine.

MAIN

Neurexins are pre-synaptic cell adhesion proteins that act as a synaptic organizer[1]. There are

three neurexin genes (*NRXN1*, *NRXN2*, and *NRXN3*) in mammals and each is highly alternatively

spliced to produce hundreds of isoforms, primarily categorized into long alpha and shorter beta

isoforms[2]. The complex alternative splicing of neurexins allows for widespread expansion of

protein-protein interaction capabilities[3], providing the basis for neurexins to interact with diverse

post-synaptic ligands to establish and maintain neurotransmission. *NRXN1α* splice variants are

specific to brain regions[4] and between cell types[5], but the cell-type-specific functional impact of

individual isoforms remains unclear. Although rare in the population, large copy number variations

(deletions or duplications) at the *NRXN1* locus 2p16.3, particularly those deleting exonic regions

are highly penetrant, pleiotropic, and are strongly associated with several neuropsychiatric

diseases, including schizophrenia (odds ratio 14.4(ref. [6]), autism spectrum disorder (odds ratio

14.9ref. [7]), epilepsy (odds ratio 9.91ref. [8]), intellectual disability (odds ratio 7.47ref. [9]) and

Tourette's syndrome (odds ratio 20.3ref. [10]). These exonic deletions in *NRXN1* are non-recurrent

(that is, they vary in size and location), making it difficult to determine the molecular mechanisms

underlying their diverse clinical symptoms (e.g., diagnosis, severity, prognosis, and age-of-onset).

In rodent studies, gene knock outs (KO) of *NRXNs* is sufficient to produce an array of excitatory

and inhibitory synaptic phenotypes[1]. However, heterozygous KO produce modest behavioral and

physiological changes *in-vivo*[11]. By contrast, however, *in-vitro* studies of engineered heterozygous

NRXN1[+/-] human neurons reveal robust changes in excitatory neurotransmission that are not

observed in matched *NRXN1*[+/-] mouse neurons[12,13]. While these studies do not evaluate how

NRXN1[+/-] impacts the complex alternative splicing of the gene or the non-recurrent nature of the

deletions, they collectively suggest that neurexins possess unique human neurobiology, and

therefore, the impact of the distinct patient-specific *NRXN1*[+/-] must be specifically evaluated in

human models.

Human induced pluripotent stem cell (hiPSC)-derived neurons provide an ideal platform to study

NRXN1α alternative splicing. Previously we established that hiPSC-derived forebrain cultures,

comprised of a mixture of glutamatergic and GABAergic neurons with astroglia, recapitulate the

diversity of *NRXN1α* alternative splicing observed in the human brain, cataloguing 123 high-

confidence *NRXN1α* isoforms[14]. Furthermore, using patient-derived *NRXN1*[+/-] hiPSCs with unique

5' or 3' deletions in the gene, we uncovered wide-scale reduction in wildtype *NRXN1α* isoform

levels and, robust expression of dozens of novel isoforms from the 3' mutant allele[14].

Overexpression of individual wildtype isoforms ameliorated reduced neuronal activity in patient-

derived *NRXN1*[+/-] hiPSC-neurons in a genotype-dependent manner, whereas mutant isoform

expression decreased neuronal activity levels in control hiPSC-neurons[14]. We therefore

hypothesized that 5' deletions of the promoter region represent classical loss-of-function (LOF),

while robust expression of 3' specific novel mutant isoforms confer a gain-of-function (GOF) effect

that cannot be rescued by overexpression of wildtype isoforms. Although *NRXN1* splicing varies

between the cell types of the brain, the impact of non-recurrent *NRXN1*[+/-] deletions on cell-type-

specific splicing patterns and synaptic function remains untested.

Neurexin signaling impacts both glutamatergic and GABAergic synapse properties[15,16],

suggesting that neurexins may regulate excitatory and inhibitory balance, which is strongly

implicated across neuropsychiatric disorders [17]. Given its multifaceted roles, the lack of

mechanistic understanding of how aberrant *NRXN1* splicing impacts neuronal physiology in a

cell-type *and* genotype-dependent manner presents a significant challenge for therapeutic

targeting. To discover the disease mechanisms that underpin *NRXN1*[+/-] deletions, we leverage

advanced 2D lineage conversion, and 3D organoid models to simultaneously compare excitatory

and inhibitory neurons, across 5' LOF and 3' GOF deletions. Our experiments were conducted

with separate populations of glutamate and GABA neurons, to specifically evaluate cell-

autonomous phenotypes arising from distinct $NRXN1^{+/-}$ deletions. We identified points of

phenotypic divergence across glutamatergic and GABAergic neurons, which were independently

validated in isogenic experiments to establish causal relationships between aberrant splicing and

synaptic dysfunction. Finally, we inform and test the design of novel therapeutic agents for

$NRXN1^{+/-}$ deletions on LOF/GOF stratified patient mechanisms.

RESULTS

Shared aberrant splicing in patient-specific glutamatergic and GABAergic neurons.

To examine the impact of $NRXN1^{+/-}$ deletions in two different neuronal cell types, we used lineage

conversion of patient hiPSCs for generating either excitatory glutamatergic neurons or inhibitory

GABAergic neurons. Transient overexpression of $NGN2$ in hiPSCs produces iGLUT neurons that

are >95% pure glutamatergic neurons, robustly express glutamatergic genes, release glutamate,

and produce spontaneous synaptic activity by day 21 in vitro[18-21]. On the other hand, transient

overexpression of $ASCL1$ and $DLX2$ yields iGABA neurons that are >80% positive for expressing

GABA and GAD1/2 by day 35 in vitro, and possess mature physiologic properties of inhibitory

neurons by day 42[19,22]. We applied these highly efficient iGLUT and iGABA induction protocols

to hiPSCs from patients with rare heterozygous intragenic deletions in $NRXN1$[14]. Two cases

containing the same ~136-kb deletion in the 3'-region of $NRXN1$ (3'-Del) and two cases with ~115-

kb deletions in the 5'-region of $NRXN1$ (5'-Del)*. For controls, we used passage-matched hiPSCs

from four healthy sex-balanced subjects (**Fig. 1a,b**). Immunostaining confirmed expression of

neurotransmitter transporters, vGLUT1 and vGAT for iGLUT and iGABA neurons, respectively

(**Fig. 1c,h**). RNA-sequencing (RNAseq) further validated iGLUT (DIV21) and iGABA (DIV35)

neuronal induction in all donors (**Fig. 1d,i**).

We first compared the transcriptomics of $NRXN1^{+/-}$ deletions with controls for both iGLUT and

iGABA neurons. RNAseq analyses revealed overlap of the top differentially expressed genes

(DEGs) across cell types/genotypes (e.g., $FAM66D, TTC34, GALNT9$), whereas the majority of

case/control DEGs were genotype-specific (**Fig. 1e,j**). Gene-ontology analyses revealed a robust

enrichment of developmental and synaptic terms for the 5'-Del iGLUT condition over all other

conditions (**Supplementary Table S1**). Using publicly curated gene lists[23,24], risk enrichments for

schizophrenia, bipolar disorder, and autism spectrum disorder were most enriched in iGLUT
neurons from 5'-Del cases and iGABA neurons from 3'-Del cases (**Fig. 1f,k and Extended Data**
Fig. 2). Baseline transcriptional comparisons therefore suggested the potential of strong
phenotypes for 5'-Del iGLUT neurons. Despite cell-type-specific *NRXN1* isoform profiles reported
in healthy brains[5] and neurons[14], aberrant *NRXN1* alpha/beta exon usages were shared between
NRXN1+/- iGLUT and iGABA neurons (**Fig. 1g,l**), whereby alpha exon usage decreased for 5'-Del
and increased for 3'-Del in both types of neurons. These observations with specific iGLUT and
iGABA neurons parallel those from our previous study using hiPSC-derived forebrain neurons,
which was a mixture of glutamatergic and GABAergic neurons with astroglia[14].

Patient-specific alterations in spontaneous neural activity occur without changes in
passive and excitable membrane properties.

To evaluate the functional consequence of *NRXN1+/-* deletions on neuronal activity, we conducted
a population-level analysis of spontaneous neuronal activity using a multi-electrode array (MEA)
with 5'-Del, 3'-Del or control neurons co-cultured with primary mouse glial cells for six weeks to
promote maturation. Interestingly, spontaneous network activity (weighted mean firing rate,
wMFR) in *NRXN1+/-* iGLUT neurons was reduced by over 40% in two independent time points, at
weeks post induction (WPI)4 (**Fig. 2a,b** 5'-Del $p = 0.0011$; 3'-Del, $p < 0.0001$, 1-way ANOVA,
Dunnett's test) and WPI6 (**Fig 2c** 5'-Del $p = 0.0238$, 3'-Del $p = 0.0211$ 1-way ANOVA, Dunnett's
test). We next examined the passive membrane properties using whole-cell patch-clamp
electrophysiology. The cell capacitance, membrane resistance and resting membrane potentials
did not significantly differ between 5'- and 3'-Del cases and control iGLUT neurons (**Fig. 2d**). To
examine the intrinsic excitability, we compared the input-output curve for induced firing and found
no statistical difference in the maximal number of evoked action potentials (**Fig. 2e**). Furthermore,
the voltage-dependent sodium and potassium current densities were similar between 5'-Del, 3'-
Del and control iGLUT neurons (**Extended Data Fig**. **3a**). Taken together, these results suggest
that changes in passive or intrinsic excitability membrane properties cannot explain the reduced
firing observed on MEAs for 5'-Del and 3'-Del in iGLUT neurons.

In parallel, we generated iGABA neurons with the same 5'-Del, 3'-Del, and control hiPSCs, as
above. We observed that immature *NRXN1+/-* iGABA neurons exhibited a robust ~2-fold increase
in population-wide wMFR activity (WPI2-3) from both 5'-Del and 3'-Del cases (5'-Del $p = 0.0381$,
3'-Del $p = 0.0047$ 1-way ANOVA, Dunnett's test) (**Fig. 2f,g**). Though unexpected for GABA
neurons, the finding is consistent with activation of ionotropic GABA receptors leading to

depolarization due to little KCC2 expression and reduction of internal chloride levels[25]. Indeed,

immature iGABA neurons expressed higher (10-fold increase compared to WPI2 neurons) levels

of *SLC12A5* (the gene encoding *KCC2*) (**Extended Data Fig. 4a** *p* < 0.0001, 2-way ANOVA,

Dunnett's test). Furthermore, this transient hyperexcitability was pharmacologically inhibited by

10μM gabazine, a selective GABA$_A$ antagonist (**Extended Data Fig. 4b,c**). In mature iGABA

neurons (>5 weeks), post-*KCC2* switch, the average wMFR decreased in 5'-Del and 3'-Del

neurons (5'-Del *p* = 0.0809; 1-way ANOVA, Dunnett's test) (**Fig. 2h**). Like iGLUT neurons, the

passive and intrinsic excitability membrane properties of iGABA neurons were not different

between *NRXN1$^{+/-}$* 5'-Del or 3'-Del and controls (**Fig. 2i,j and Extended Data Fig. 3b**).

Overall, these data suggest that patient-specific changes in spontaneous neural activity are not

driven by differences in passive and excitable membrane properties in iGLUT or iGABA neurons

from *NRXN1$^{+/-}$* 5'-Del and 3'-Del patients. Furthermore, these deletions did not appear to alter

maturation of iGLUT or iGABA neurons, as evidenced by similar induced firing rates and resting

membrane potentials.

***NRXN1$^{+/-}$* 5' and 3' deletions result in divergent synaptic transmission deficits.**

To further dissect the factors mediating phenotypes in altered spontaneous firing, we next

investigated the efficacy of synaptic transmission in iGLUT and iGABA neurons. Voltage-clamp

recordings of spontaneous excitatory post-synaptic currents (sEPSCs, no TTX) in iGLUT neurons

revealed decreased frequency of events for both for 5'-Del and 3'-Del cases (**Fig. 3a**). The

cumulative probabilities of inter-event-intervals (IEI) for both 3'-Del (*p* = 1.27E-11, Levene's test)

and 5'-Del iGLUTs (5'-del *p* = 2.44E-5, Levene's test) (**Fig. 3a,b**) was significantly increased,

compared to controls. The sEPSC amplitude increased for 5'-Del neurons (**Fig. 3a,c**). Miniature

excitatory post-synaptic currents (mEPSCs, +TTX) showed similar trends in IEI but no changes

in amplitude sizes across genotypes (IEI 5'-del *p* = 0.0411, 1-way ANOVA, Dunnett's test)

(**Extended Data Fig. 5a,b**). These reductions in synaptic transmission are consistent with the

transcriptomic signatures; pre-synaptic (SynGO) genes showed a larger change (than post-

synaptic) in synaptic gene expression signatures (5'-del Pre-SynGO, Log$_2$FC = 0.1252; 3'-del Pre-

SynGO, Log$_2$FC= -0.0156) (**Fig. 3d,e**). We further probed transcriptional signatures of known

NRXN1 trans-synaptic interaction partners that mediate synapse formation, function, plasticity

and are frequently linked with neuropsychiatric disease[16], and found they were represented in

DEGs, including *CBLN2*, *LRRTM4* and *NXPH1*, suggesting these synaptic effects are largely

driven by change in *NRXN1* expression (**Fig. 3f**).

By contrast, synaptic transmission in iGABA neurons appeared to be enhanced. The frequency

of spontaneous inhibitory post-synaptic currents (sIPSCs) increased, marked by a significant

decrease in IEI in 3'-Del iGABA neurons (3'-del p = 5.05E-5 by Levene's test) (**Fig. 3g,h**). There

was no change in sIPSC amplitude (**Fig. 3g,i**). Miniature inhibitory post-synaptic currents

(mIPSCs) recorded in the presence of TTX and CNQX (AMPA/Kinate receptor antagonist)

revealed similar trends in IEI, with no change in mIPSC amplitudes (**Extended Data Fig. 5c,d**).

Similarly, SynGO analysis revealed concordant changes in synaptic transmission and

transcriptomic signatures at 3'-del Pre-SynGO (Log$_2$FC = 0.0324), and represented DEGs among

NRXN1 trans-synaptic interaction partners *CBLN2, NLGN1, NXPH1, CASK,* and *LRRTM2-4* (**Fig.**

3j-l).

Overall, patient-specific alterations in spontaneous neural activity were driven by synaptic deficits,

with the cell type-specific impact of aberrant *NRXN1* splicing resulting in divergent

neurotransmission phenotypes. Furthermore, non-recurrent LOF and GOF presented unequal

effect sizes between cell-types, with 5'-Del neurons being most impacted by excitatory

transmission, but 3'-Del neurons being affected by both excitatory and inhibitory transmission.

Given that deletions affected iGLUT and iGABA neurons in opposing directions, these findings

may implicate *NRXN1* as a key mediator of excitatory/inhibitory balance, a prevalent theme

among neuropsychiatric disorders[26].

Region-specific organoids resolve non-autonomous defects of neurogenesis and

neuronal development.

To explore the impact of aberrant *NRXN1* splicing on neural patterning[27] within a self-organized

microenvironment, we applied dorsal forebrain patterning to yield 5'-Del, 3'-Del and control human

cortical organoids (hCOs) that resemble the pallium, or applied ventral forebrain patterning to

generate human subpallial organoids (hSOs) that resemble the subpallium (**Fig. 4a,h**)[28,29].

Organoid sizes varied at early time points, but maintained consistency across genotypes by 4

months (**Extended Data Fig. 6a,b,g,h**). RT-qPCR validated dorsal and ventral patterning, and

further confirmed robust expression of mutant isoforms exclusively from the 3'-Del patients, with

no expression from 5'-Del and control organoids (**Extended Data Fig. 6c-e, i-k**). To unbiasedly

characterize transcriptomics of cortical and subpallial organoids, we performed single cell RNA-

seq (n = 47,460 cells from hCOs, n= 35,563 cells from hSOs using 10x genomics) at 6 months, a

timepoint with well characterized neural activity[28,30]. We identified subclusters of cell-types within

hCOs and hSOs (**Fig.4b,i**), and confirmed consistent distributions of cell clusters across

genotypes (**Fig. 4c,j**). Strong expression of *NRXN1* and *FOXG1* confirmed forebrain identity (and

excluded one of six cell lines), while *EMX1* expression confirmed dorsalization of remaining hCOs.

Likewise, *DLX2* expression confirmed ventralization of hSOs, with an enrichment of GABAergic-

fated neurons (**Fig. 4d,k and Extended Data Fig. 6f,l**).

A recent study demonstrated that KO of *NRXN1* increased replication of NPCs in human cells

and in the *xenopus* telencephalon[31]. To test if *NRXN1*[+/-] deletions could impact neurogenesis, we

quantified neural rosette proliferation detected with immunofluorescence by calculating the ratio

of *KI67* protein to *SOX2* protein expression, as markers of proliferative cells and multipotent

neural stem cells. In the hCO subtype (DIV55-62), 3'-Del patient rosettes exhibited increased

KI67:SOX2 compared to control (p = 0.0063, by 1-Way ANOVA) (**Fig. 4e,f**), while in the hSO

subtype, 5'- and 3'-Del rosettes had reduced *KI67:SOX2* ratios (5'-del p = 0.0004; 3'-del p =

0.0004, by 1-Way ANOVA) (**Fig. 4l,m**). These results may explain the variation in NPC:neuron

proportions across genotypes (**Fig. c,j**). Differentially expressed gene sets identified cell-type-

specific effects in hCO-glutamatergic, hSO-GABAergic, and astroglia clusters, with robust

enrichment of GO terms related to RNA splicing (from upregulated genes), and

neurodevelopment and synaptic function (from downregulated genes) (**Fig. 4g,n**). Much of the

enriched GO terms overlapped among hCO-glutamatergic and hSO-GABAergic neurons, and

unexpectedly certain GO terms also reached significance in hCO-astroglia, but not hSO-astroglia

clusters (**Extended Data Fig. 7**). Overall, these data suggest that patterns of divergent cell-type

specific effects, particularly at the level of neural precursor states, persisted in our 3D models of

forebrain development, resulting in significant changes to glutamatergic and GABAergic neurons.

Isogenic validation of bidirectional excitatory-inhibitory (E-I) synaptic deficits.

To further support our findings of opposing, cell-type-specific changes in glutamatergic and

GABAergic synaptic transmission, with unequal effect size between non-recurrent *NRXN1*

deletions, we utilized a more direct approach using isogenic lines to causally link aberrant splicing

to synaptic dysfunction and develop candidate therapies. We utilized short hairpin RNAs

(shRNAs) to knockdown wildtype or mutant splice isoforms to mimic LOF or rescue GOF

phenotypes, respectively. Targeted knockdown of constitutively expressed exon 9 (expressed in

all alpha isoforms but not beta isoforms) achieved mRNA knockdown of ~55% in iGLUT neurons,

and ~75% in iGABA neurons across two or more isogenic pairs (compared to a non-targeting

(NT) control shRNA) (**Extended Data Fig. 8a,e**). In iGLUT neurons, we observed decreased

synaptic transmission (i.e., increased sEPSC IEI, p = 2.2E-16 by Levene's Test) (**Fig. 5a,b**),

whereas in iGABA neurons we observed increased synaptic transmission (decreased IEI, p =
3.65E-8 by Levene's Test and p = 0.0092 by Student's t-test) (**Fig. 5g,h**), similar to the changes
with 5'-Del neurons (**Fig. 3b,h**). Transcriptomic profiles of isogenic recreation lines further
validated total *NRXN1* knockdown (**Extended Data Fig 7c,g)** and demonstrated DEGs related to
multiple aspects of synaptic function in both iGLUT (29/258 DEGs; 1.274161-fold, p = 0.1049)
and iGABA neurons (384/3525 genes; 1.234861-fold, p = 1.307E-6) (**Fig. 5c,l and**
Supplementary Table S2). Altogether, knockdown of wildtype splicing recapitulated cell-type-
specific differences in *NRXN1*$^{+/-}$ 5'-Del neurons, causally implicating decreased wildtype *NRXN1α*
expression (LOF) as a key driver of opposing cell-type-specific phenotypes, and displayed
evidence of reshaped synaptic properties.

For *NRXN1 3*'-Del neurons, we reasoned that reducing the expression of the GOF transcript
should restore the changes in synaptic transmission for both iGLUT and iGABA neurons. We
designed a shRNA against the mutant splice junction overlapping exons 20 and 24, expressed in
all 3'-Del unique *NRXN1* alpha and beta isoforms, and achieved targeted knockdown of mutant
splice isoforms by 95% in iGLUT, and 25% in iGABA neurons in both donors (**Extended Data Fig**
6b,f). In the donor with most robust knockdown in both iGLUT (90%) and iGABA (37%) neurons,
electrophysiological recordings revealed reversal of changes in synaptic transmission. In iGLUT
neurons, shRNA MT decreased sEPSC IEI (p < 2.2E-16 by Levene's Test and p = 0.0194) (**Fig.**
5d,e) and in iGABA neurons increased sIPSC IEI (p = 0.042 by Levene's Test) (**Fig 5j,k**), as
compared to shRNA-NT. Notably, the IEIs for rescued neurons appeared similar to those for
control iGLUT and iGABA neurons (**Fig. 3b,h**). Parallel to the LOF isogenic transcriptomic
profiles, the GOF isogenic line had modest changes in iGLUT neurons (3/28 DEGs; 1.214533-
fold, p = 0.4541 by Levene's test), but more pronounced synaptic DEGs in iGABA neurons
(128/1077 DEGs; 1.347225-fold, p = 2.694E-4) (**Fig. f,l and Extended Data Fig. d,h**).

In summary, knockdown of WT *NRXN1* in iGLUT and iGABA neurons recapitulated
neurotransmitter phenotypes of 5' LOF *NRXN1* deletions, while knockdown of MT isoforms
rescued them, appearing to mitigate the negative effect of the 3' GOF NRXN1 deletions.
Furthermore, the bidirectional effect of *NRXN1* deletions on iGLUTs (decrease) and iGABAs
(increase), i.e., E-I balance, was reversed by shRNA treatments. Transcriptomic profiling of
shRNA treated isogenic lines revealed ~10% of DEGs related to synaptic function across all four
experimental conditions. Altogether, shRNA mediated perturbations among WT and MT isoforms
causally implicate *NRXN1* splicing to synaptic dysfunction.

Precise therapeutic targeting of stratified LOF/GOF binary phenotypes

Based on our causal linkage of unique LOF/GOF aberrant $NRXN1\alpha$ splicing patterns to E-I synaptic dysfunction, we stratified our patients to systematically explore novel therapeutic strategies. Our findings across case/control and isogenic experiments highlight impaired E-I signaling, notably a bias against excitation, via decreased glutamatergic and increased GABAergic synaptic activity. As proof of principle, we tested methods to manipulate $NRXN1$ expression, focusing on reversing phenotypes in iGLUT neurons, to mitigate decreased excitation.

For GOF patients, we investigated the utility of using anti-sense oligonucleotides (ASOs), which have been recently used to treat different neurological diseases[32], to target a specific RNA splice site. We first designed an "alternative splice matrix," by juxtaposing splice donor and acceptor RNA sequences from each $NRXN1\alpha$ exon, which generates all possible combinations of canonical and non-canonical splice junctions (**Fig. 6a**). We noted the sequence that covers the 20/24 mutant splice junction and designed an ASO to facilitate targeted RNAseH1-dependent degradation of 3-Del mutant isoforms. ASO treatment ($1\mu M$) for 72hrs decreased total mutant isoforms by ~55% in iGLUT neurons ($p < 0.001$ by student's t-test), as compared to a non-targeting control ASO (**Fig. 6b**). Transcriptomic profiling revealed robust DEGs, enriched for synaptic properties and neurotransmitter signaling and neurodevelopmental pathways **(Supplementary Table S3)** (231/1844 Bonferroni corrected DEGs annotated in SynGO; 0.420028-fold, $p = 1.016E-08$), (**Fig. 6c-d**). Supporting the broader applicability of an ASO strategy, we investigated the presence of mutant splice isoforms in post-mortem brain tissue from an unrelated $NRXN1^{+/-}$ case diagnosed with autism spectrum disorder (ASD) (**Supplemental Fig. 9**). By integrating long-read and short-read sequencing[14], we identified nine high-confidence isoforms that are predicted to be translated: two of which contained a novel splice junction overlapping the deletion encompassing exon 14/19, targetable by an ASO tailored to mutant splice junction.

To develop a therapeutic strategy for treating LOF patients, we took a different approach. We hypothesized that enhancing transcription of the WT $NRXN1\alpha$ allele might be sufficient to restore $NRXN1$ levels. β-estradiol reversed $NRXN1$ LOF neurogenesis deficits in *xenopus* and human NPC models[31]. The mechanism is unknown, but chromatin immunoprecipitation with sequencing (ChIP-Seq) in mouse brain tissue identified estrogen receptor alphaα (ERα) binding sites at the $NRXN1$ alpha locus (**Extended Data Fig. 10**)[33]. We predicted $NRXN1$ to be a target of ERα, and

found that acute treatment with β-estradiol (10nM or 30nM, 3-5 days) significantly increased

NRXN1α expression in iGLUT neurons (*p* = 0.0297 by Student's t-test) derived from 5'-Del

patients (**Fig. 6f**). To test the hypothesis that chronic exposure to β-estradiol would ameliorate

phenotypes of iGLUT synaptic transmission, we treated post-mitotic 5'-Del neurons with either

30nM β-estradiol or vehicle (DMSO), and measured spontaneous neural activity with MEA and

synaptic transmission with patch-clamping. MEA recordings at DIV21-13 revealed a significant

increase in wMFR activity (*p* = 0.0057 via student's t-test) in the β-estradiol treated condition (**Fig.**

6g). In addition, β-estradiol treatment ameliorated the change in sEPSC IEIs in 5'-Del neurons (*p*

= 1.54E-4 by Levene's Test), but not in vehicle treated 5'-Del neurons (*p* = 0.9055 by Levene's

Test) **(Fig. 6h,i)**.

Altogether, increased *NRXN1* expression in LOF neurons and knockdown of mutant splicing in

GOF neurons can each rescue the opposing cell-type-specific case/control differences in

NRXN1[+/-] neurons, implicating *NRXN1α* expression as a key a driver of *NRXN1* mutant

phenotypes, and that modulators of *NRXN1* expression and splicing may be utilized for targeted

therapies.

DISCUSSION

NRXN1[+/-] deletions are linked to diverse clinical outcomes, all of which have been associated with

excitatory-inhibitory (E-I) imbalance[1,27,30]. While several studies have shed insights into the

regulation of synaptic properties, the molecular regulators of E-I balance has yet to be

characterized. Here, we show that *NRXN1* splicing bidirectionally regulates E-I synaptic

transmission, with the 5'-Del and 3'-Del patients reducing sEPSC but increasing sIPSC IEIs in

iGLUT and iGABA neurons, respectively. These changes likely play a critical role in altering the

E-I balance, a common theme among neuropsychiatric disorders. Using a case/control *NRXN1*[+/-]

cohort as well as isogenic manipulations of the *NRXN1* isoform repertoire, we report distinct

phenotypic effects in human iGLUT and iGABA neurons, which predominately manifest in

changes in the frequency of synaptic transmission. These results suggest causal relationships

between aberrant splicing and synaptic dysfunction, dictated by unique patient-specific

mechanisms. A recent report evaluated the impact of *NRXN1*[+/-] on forebrain organoids by single

cell transcriptomic characterization over time, and highlighted specific perturbations in the

developmental trajectories of early neural progenitors[34]. Likewise, we report divergent

proliferation dynamics, with an increase in glutamatergic enriched hCOs, but a decrease in

GABAergic-enriched hSOs. While it is mechanistically unclear how mutations in *NRXN1* impact

neurogenesis, prior evidence also exists for this phenomenon[31,35], altogether suggesting roles for

NRXN1 beyond mediating synaptic organization and transmission, perhaps similar to

SYNGAP1[36], a gene canonically thought to have exclusive synaptic function. Nevertheless, our

causal demonstration of manipulating *NRXN1* splicing, which resulted in the recreation/rescue of

synaptic phenotypes in post-mitotic neurons suggests that direct manipulation of splicing may

achieve therapeutic benefit in cases.

Neurexins are expressed across all synapses and among certain non-neuronal cell-types, such

as astrocytes, which may confer distinct functional roles[37]. With respect to GOF mechanisms, it

remains unclear whether mutant isoforms shift the stoichiometry of alternative splicing against

wildtype RNA isoforms or alter trans-synaptic protein-protein interactions. Unique patterns of

aberrant splicing amongst non-recurrent mutations would be challenging to target with a common

pharmacological agent for all *NRXN1*[+/-] carriers. Given that *NRXN1* fundamentally encodes for

pre-synaptic cell adhesion molecules, traditional neuropharmacological agents that target specific

receptors within single neurotransmission systems would be unlikely to broadly ameliorate the

many levels of neural circuitry impacted across neurodevelopment, especially within dysfunctional

synapses. Instead, precision therapies that target aberrant splicing are needed to effectively

reverse defects in the multiple cell types affected by patient-specific deletions.

Our results show that the opposing effects of *NRXN1*[+/-] in glutamate and GABA neurons provides

the foundation for evaluating proof-of-concept therapeutics that target aberrant splicing, once the

LOF or GOF phenotype has been determined. ASOs designed to target a mutant splice junction

to facilitate degradation of mutant isoforms represent a therapeutic approach with which to

ameliorate the toxic effects of GOF mutations (**Fig. 6e**). While gene therapy in brain disease has

resulted in mixed successes[38], cases of muscular dystrophies which have been treated via

splicing modulating ASOs have proved effective[39].

For LOF mutations, mechanisms for upregulation of wild-type allelic expression and restoration

of proper *NRXN1* splicing are required (**Fig. 6j**). Steroid-based pharmaceuticals for treating LOF

mutations is feasible, as they have been used to treat neuroinflammation-related conditions[40],

with recent studies revealing estrogen-mediated roles in neuroprotection[31], demonstrating the

safety and feasibility of this approach. Altogether, precision medicine provides therapeutic

promise for treating neuropsychiatric conditions.

Our findings with estrogen-dependent regulation of *NRXN1* expression also hint at possible

mechanistic explanations for the dramatic sex differences in the prevalence and severity of autism
spectrum disorders, for which the biological basis for the prominent male bias in diagnosis is
unknown[41]. At the genetic level, evidence exists for a female protective effect – the observation
that females have a higher threshold for reaching affectation status and require greater mutational
burden to express ASD phenotypes due to female-specific protective factors that lead to
resilience[42,43]. Estradiol ameliorates neurodevelopmental phenotypes in *xenopus* and zebrafish
genetic models of ASD[31,44], suggesting that estradiol might be modulating relevant circuits in the
setting of ASD risk genes to mitigate a range of ASD gene mutations. Estradiol modulates
excitatory and inhibitory circuits in the developing brain, regulating the switch from excitatory to
inhibitory GABAergic signaling in the hippocampus and playing a neuroprotective role against
glutamate-induced excitotoxicity[45,46]. Estrogens therefore may modulate neurodevelopment of
cells and circuits in a manner that confers resilience to ASD.

While our work carefully considers the cell-type specific impact of *NRXN1* splicing on non-
recurrent deletions by utilizing autonomous culture paradigms, it is important to acknowledge
certain limitations inherent in our study. Notably, the neuronal subtypes that are yielded from
lineage conversion can be diverse and immature, as recently characterized[47]. Furthermore, the
impact of $NRXN1^{+/-}$ on synaptic physiology across different subtypes of neurons (e.g., SST versus
PV expressing GABA neurons), remains to be determined. Future studies will need to
comprehensively assess the impact of $NRXN1^{+/-}$ across more discrete cell types and complex co-
culture systems, which may elucidate non-autonomous and activity dependent phenotypes of
$NRXN1^{+/-}$. Furthermore, it will be important to also probe the biochemical and proteomic
interactions of *NRXN1*. Unbiased proximity-labelling methods such as BioID[48], which may provide
mechanistic insights into *NRXN1* mediated synapse function, by defining alternations in binding
profiles from wildtype and mutant isoforms.

Precision medicine seeks to tailor treatments to individual patients[49]. Here we focus on the
possibility of stratifying cases based on inherited DNA mutations, with the goal of one day
providing the right therapy at the right time to the right patient. The most successful examples of
precision medicine to date have been in cancer[50], and monogenic disease[51], where genetic
analysis reveals molecular subtypes that benefit from specific treatments. In the context of brain
disorders, patient stratification is particularly challenging, reflecting in part the myriad rare and
common variants linked to disease[52]. An additional challenge, as discussed herein, is that
different mutations in the same genes can result in distinct phenotypic effects. Notably, genome-
wide disease-agnostic analyses reveal that transcription factors and nucleic acid binding proteins

are overwhelmingly driven by LOF mutations; whereas signaling molecules, enzymes, receptors

and transporters by dominant GOF mutations[53]. Therefore, a binary therapeutic approach, similar

to what we propose for *NRXN1*, may prove suitable for dual loss- and gain-of-function

mechanisms linked to mutations in other neuropsychiatric disorder-related synaptic genes (e.g.,

NLGN3[54], *CACNA1D*[55], *CACNA1C*[56], *SCN1A*[57], *SCN2A*[58]) as well as broadly in

neurodegenerative disease (e.g., *SOD1*[59] *TDP43*[59] *FUS/TLS*[59] *C9ORF72*[60], *AR*[61]), and many

short tandem repeat disorders[62]. Taken together, our work highlights important implications for

precision medicine, demonstrating the necessity of functionally dissecting the phenotypic impact

of diverse patient-specific genetic variants across cellular contexts, in order to resolve candidate

therapies across stratified disease mechanisms.

METHODS

Plasmid designs and molecular cloning

i. TetO-Ascl1-Neo

The *ASCL1* insert from TetO-Ascl1-Puro (Addgene #97329) was synthesized as a gBLOCK

flanked by EcoR1 cut sites, and cloned into TetO-hNgn2-Neo using EcoR1 to remove *NGN2.* The

recipient vector was dephosphorylated with shrimp alkaline phosphatase (rSAP NEB #M0371S)

during the digest, column purified and ligated at a 1:1 vector to insert ratio using the QuickLig Kit

(NEB #M2200S). The *ASCL1* stop codon was subsequently mutated using the QuickChange II-

XL site-directed mutagenesis kit (Agilent #200523) and verified via whole plasmid sequencing

from plasmidsaurus.

ii. shRNA RNA interference constructs

All shRNAs were designed and produced by Sigma-Aldrich via custom submitted sequences

against wildtype *NRXN1α* (constitutively expressed exon 9) and mutant *NRXN1α* (mutant 20/24

splice junction) cloned into TCR2 pLKO.5-puro.

Cell Culture

i. hiPSC culture

Passage matched (±3) human induced pluripotent stem cells (hiPSCs) were cultured in StemFlex

media (Life technologies #A3349401) on Matrigel (Corning, #354230). At ~70-80% hiPSCs were

clump passaged using 0.5mM EDTA in PBS without Mg/Ca (Life technologies #10010-031). Cells

were washed once and incubated for 4-7 min with EDTA at RT. The EDTA was aspirated, and

the remaining colonies were lifted off with 1mL of StemFlex and re-distributed at varying densities

in a 6–well maintenance plate. hiPSC lines were split every 4-6 days. For neuronal/organoid

differentiation, wells of similar confluence across all hiPSC donors were resuspended and seeded

onto a Matrigel coated 10cm dish and expanded until ~70-80% confluency.

ii. HEK293T culture and lenti-viral production

HEK293T cells were maintained in 15cm dishes and cultured in DMEM supplemented with 10%

standard FBS (Life technologies #16000069). 3rd Gen lenti-viral particles were produced using

previously described methods and REV, MDL and VSV-G plasmid ratios[18], each transfected with

PEImax (Polysciences #24765-1). Each PEIMax batch was volumetrically titrated at total

ugDNA:uLPEI for optimal transfection efficiency.

iii. Primary mouse glia production

All mouse work was performed under approved IACUC protocols at the Icahn School of Medicine

at Mount Sinai. C57BL/6 mice were used as breeding pairs. For glial preps, dissected cortices

from 3 pups (at p0-3) were dissociated using papain (R&D #LS003126) and plated on 10cm

dishes in MEF medium (DMEM, 10% Cosmic Calf Serum (Fisher #SH3008703HI), 1x Antibiotic-

antimycotic (Life technologies #15240), 1x Sodium Pyruvate (Life technologies #11360070), 1x

MEM Non-Essential Amino Acids Solution (Life technologies #11140050), 4uL 2-Mercaptoethanol

(Sigma #60-24-2), supplemented with 100µg/mL Normocin (InvivoGen #ant-nr-2). Glial cells were

recovered and propagated for 7 days, expanded into three 10cm dishes and were used for co-

culture experiments. All glial preps were tested twice for mycoplasma (Normocin withdrawn)

(Lonza, #LT07-318) prior to freezing or neuronal co-culture. At day 14, one 10cm dish with mouse

glia were distributed to two MEA, 12- or 24- well plates, and subsequently inactivated with 4µM

Ara-C (Sigma #C1768) prior to or during re-seeding of induced neurons.

iv. iGLUT induction and astrocyte co-culture

At day -1 hiPSCs expanded in 10cm dishes were dissociated with accutase (StemCell

Technologies, #07920), washed and pelleted with a 1:4 ratio of accutase to DMEM for 5 min at

1000 rcf, and re-suspended with StemFlex media containing ROCK inhibitor, THX (10µM/mL;

Sigma Aldrich, SML1045). The hiPSCs are then co-transduced with TetO-Ngn2-Puro (Addgene

#79049) or TetO-Ngn2-Neo (Addgene# 99378) and UCb-rtTA (legacy vector from the lab of Fred
Gage) and seeded at 1.0-1.5x10^6 cells in 1.5mL per well on 6-well plates were coated with 2x
Matrigel for at least one hour at 37C. The hiPSC-viral mixture was then incubated overnight. The
following morning (day 0), a full media change with iGLUT induction media was performed with
the following recipe: Neurobasal Media: 1x N2 (Life technologies #17502-048), 1x B-27 minus
Vitamin A (Life technologies #12587-010), 1x Antibiotic-Antimycotic, 1x Sodium Pyruvate, 1x
GlutaMax (Life technologies #35050), 500µg/mL cyclic-AMP (Sigma #D0627), 200nM L-ascorbic
acid (Sigma #A0278), 20ng/ml BDNF (Peprotech #450-02), 20ng/ml GDNF (Peprotech #450-10),
1µg/ml natural mouse laminin (Life technologies #23017015). On days 1-2, iGLUT cells were
treated with respective antibiotic concentrations at 1ug/mL puromycin (Sigma# P7255) or
0.5ug/mL neomycin (Life technologies #11811-031). On day 3, antibiotic medium was withdrawn
and iGLUT cells were treated with 4µM Ara-C. On Day 4, iGLUT cells were dissociated with
accutase for 15 min, washed and pelleted with a 1:4 ratio of accutase to DMEM for 5 min at 800
rcf, and re-suspended with iGLUT media containing ROCK inhibitor, Ara-C and 2% low-
hemoglobin FBS (R&D systems #S11510). iGLUT neurons were distributed among wells (500-
750k cells per 24wp or 0.75-1.5E6 cell per 12wp) pre-seeded with confluent mouse glia. The
following day, iGLUT neurons received a full media change with Brainphys maturation media
(Neurobasal Media, 1x N2, 1x B-27 minus Vitamin A, 1x Antibiotic-Antimycotic, 500µg/mL cyclic-
AMP, 200nM Ascorbic Acid, 20ng/ml BDNF, 20ng/ml GDNF, 2% low-hemoglobin FBS, 1µg/ml
Mouse Laminin) supplemented with Ara-C and were subsequently monitored for growth of non-
neuronal/glial cells. Ara-C treatment was tittered down with half-media changes (without Ara-C)
every 3-4 days until used for experiments.

v. iGABA induction and astrocyte co-culture

iGABA production paralleled the methods aforementioned. hiPSCs were instead co-transduced
with TetO-Ascl1-puro (Addgene #97330) or TetO-Ascl1-neo (Addgene #TBD), TetO-Dlx2-hygro
(Addgene #97329) and UCb-rtTA and seeded at 0.8-1.2x10^6 cells in 1.5mL per well on 6-well
plates similarly prepared. The following morning (day 0), a full media change with iGABA induction
media (DMEM/F-12 + Glutamax, 1x N2, 1x B-27 minus Vitamin A, 1x Antibiotic-Antimycotic) was
performed. On days 1-2 iGABA cells were selected with respective antibiotic concentrations at
1µg/mL puromycin or 0.5ug/mL neomycin and 0.25ug/mL hygromycin (Life Technologies
#10687010), followed by antibiotic withdrawal and Ara-C treatment on day 3. iGABA neurons
were re-seeded identically to iGLUT cells, at 150-250k cells per 24wp well. iGABA cultures were

morphologically QC'ed prior to all experiments, with uncharacteristic batches being discarded.

vi. Cortical and subpallial organoid differentiation

Cortical organoids were generated according to the protocol described by Sloan et. al.[63], with
several modifications. hiPSCs were first aggregated into embryoid bodies (EBs) using an
AggreWell™800 Microwell Culture Plate system (Stemcell Tech #34850). Expanded hiPSCs
were rinsed twice with DPBS without Ca/Mg, and then dissociated using accutase. 3x10E6
hiPSCs were added to a single well in the prepared AggreWell and allowed to aggregate in a 37
°C incubator for 24 hours. The following day (day 0), EBs were dislodged from the AggreWell
plate using a cut p1000 pipette tip and passed over a 40µm strainer, and washed with excess
DMEM. The strainer was inverted over an Ultra-Low Attachment 10 cm culture dish and the EBs
were collected in spheroid induction media, which contained Stemflex supplemented with two
SMAD inhibitors, SB-431542 (SB) and LDN193189 (LDN), and THX. The following day (day 1),
the media THX was withdrawn. From d2-d6, induction media was replaced daily, and no longer
contained the Stemflex supplement (only base Stemflex media with SB and LDN). On day 6, the
media was replaced with organoid patterning media, formulated with Neurobasal-A medium, 1x
B-27 minus Vitamin A, 1x GlutaMAX, and 1x Antibiotic-Antimycotic. From d6-d24, organoid
maturation media was supplemented with 20 ng/ml of EGF (R&D Systems, #236-EG) and 20
ng/ml FGF2 (R&D Systems, #233-FB-01M). Media was changed every day from d6 – d15, after
which media was changed every other day. From d25-d43, the organoid maturation media was
supplemented with 20 ng/ml of NT-3 (PeproTech, #450-03) and 20 ng/ml BDNF. From d43
onwards, organoids received organoid maturation media with no supplements, and media was
changed every 4 days or as needed. Subpallial organoids were generated in the same way as
cortical organoids, but with additions in media formulations. From d4-d23, hSOs received
spheroid induction media or organoid maturation media supplemented with 5 µM of the Wnt
inhibitor, IWP-2 (Selleckchem, #S7085). From d12-d23, hSO organoids received neuronal
differentiation media supplemented with 100 nM of the SHH agonist, SAG (Selleckchem,
#S7779).

Electrophysiology

i. Multi-electrode array (MEA)

The Axion Maestro (Middleman) system was used to perform all MEA recordings. Measurements
began as early as DIV9 for both iGLUT and iGABA co-cultures. For time course experiments,

MEA plates were recorded every 2-3 days per week with a full media change prior to each
recording.

ii. Whole-cell patch-clamp electrophysiology

For whole-cell patch-clamp recordings, iGLUT or iGABA human-mouse glia co-cultures were
recorded at 4-6 weeks following dox-induction (time points specified in figure legends), with a full
media change one day prior to recording. Cells were visualized on a Nikon inverted microscope
equipped with fluorescence and Hoffman optics. Neurons were recorded with an Axopatch 200B
amplifier (Molecular Devices), digitized at 10 kHz using a Digidata 1320a (Molecular Devices) and
filtered between 1-10 kHz, using Clampex 10 software (Molecular Devices). Patch pipettes were
pulled from borosilicate glass electrodes (Warner Instruments) to a final tip resistance of 3-5 MΩ
using a vertical gravity puller (Narishige). To sustain the baseline activity of neurons from
extended cultures, coverslips were recorded in base Brainphys medium (external solution, Osm
305, pH 7.49). Each coverslip was equilibrated to room temperature for 10min, with 3-5 neurons
were recorded for no more than a total of 75min per coverslip. For measurements of passive and
excitable properties, an internal patch solution was used containing (in mM): K-d-gluconate, 140;
NaCl, 4; MgCl2, 2; EGTA, 1.1; HEPES, 5; Na2ATP, 2; sodium creatine phosphate, 5; Na3GTP,
0.6, at a pH of 7.4. Osmolarity was 290-295 mOsm. Neurons were chosen at random using DIC
and all recordings were made at room temperature (~22°C). Current-clamp measurement
occurred across -10pA to +50pA steps, with a maximum stimulus of +60pA, whereas voltage-
clamp measurements occurred across -50mV to +50mV steps, normalized to cell capacitance (to
control for variable neuronal size). Current clamp measurements were corrected for the junction
potential (~-15.5mV). For sEPSC/sIPSC recordings, the internal solution was replaced with (in
mM): Cesium-Chloride, 135; HEPES-CoOH, 10; QX-314, 5; EGTA, 5. Osmolarity was 290-295
mOsm. All mEPSC measurements were recorded under the presence of 100nM TTX-citrate
(Tocris Cat# 1069). mIPSC measurements were made using 100nM TTX-citrate and CNQX-
disodium salt (Tocris Cat#1045/1) to pharmacologically inhibit ionotropic glutamate receptors. All
chemicals were purchased from Sigma-Aldrich Co. (St. Louis, MO). All toxic compounds were
handled in accordance with ISMMS EHS standards.

iii. Patch-clamp data analysis

All patch-clamp data were analyzed on Clampfit (v) and Easy-Electrophysiology (v2.4.1 or beta-
versions). Briefly, for voltage-clamp data, files were opened in Easy-Electrophysiology and two

bins were assigned for Na+/K+ measures for minimum and maximum current values, respectively.

For current-clamp data, files were opened in Easy-Electrophysiology and action potential (AP)

analysis was used to automatically determine spike number and properties. For gap-free

recordings, all data was post-hoc adjusted to a baseline of zero on ClampFit, and subsequently

analyzed in Easy-Electrophysiology by template and linear threshold detection. For case control

experiments, a minimum cutoff of 10 events for the duration of the recording (3min) was used as

QC. For typical EPSC events, a single template from a randomly chosen recording was used to

analyze all traces (with a 30ms decay cutoff). For IPSC events, three templates were used to

detect variable GABA receptor kinetics, for all traces (with a 60ms decay cutoff). An amplitude cut

off of 7pA was used to call positive events. For cumulative probabilities, each cell was binned by

experiment and averaged across all cells for a representative curve (GraphPad Prism).

RNA-Sequencing and bioinformatic analyses

i. Bulk RNA sequencing and DEG analysis of iGLUT and iGABA neurons

iGLUT (DIV21) and iGABA (DIV14 or DIV35) co-cultured with primary mouse glia (to match

functional experiments) were harvested in Trizol and submitted to the New York Genome Center

[64] for high-throughput RNA extraction and quality control, library prep using the Kapa Hyper library

prep with ribo-erase (Roche #KK8541) and subsequently sequenced on Illumina NovaSeq.

Returned raw data files were first processed to remove mouse reads in the RNA-seq data, a

combined genome reference (hg38 and mm10) was created using the "mkref" command in

cellranger v6.1.2. The raw sequencing reads were then aligned to the combined genome

reference using STAR v2.7.2a[65]. Reads that mapped specifically to the human reference genome

were extracted from the resulting BAM files for subsequent gene expression analysis. Gene-level

read counts were obtained using the Subread v2.0.1 package featureCount[66], and RPKM values

were calculated using the R package edgeR[67]. To confirm sample identity, variants were called

from RNA-seq bam files by HaplotypeCaller and GenotypeGVCFs in GATK v4.2.0.0. Then

bcftools v1.15 was used to examine variants concordance with variants from whole-exome

sequencing data from the same donor. Following donor identity confirmation, the differential gene

expression analysis followed the methods as described previously[68]. First, CibersortX[69] was

utilized to predict differences in cell type composition across all samples. The R package

variancePartition v1.30.2[70] was then employed to investigate the contribution of specific variables

to the variance in gene expression. The limma/voom package[71] was used for differential

expression analysis, with the regression of fibroblast and hiPSC cell type compositions. The

analysis began with a comparison between the case and control groups. Subsequently, within
each case vs control group, subgroup comparisons were conducted for all four pairs (two donors
each for 3'-Del and 5-Del patients and two healthy controls) of samples, accounting for
heterogeneity between different donors. Genes with a fold change larger than 1.5 between the
patient and control group, an FDR less than 0.1, and a fold change larger than 1.5 in all four pairs
of subgroup comparisons were defined as the final set of differentially expressed genes. Kallisto
v0.46.1[72] was used to calculate the *NRXN1* exon usage ratios.

ii. Network Analysis of top DEGs overlapping with disease risk genes

The ASD, BP, and SCZ risk gene lists were extracted from previously curated gene lists [23], Genes
with the top 200 smallest FDR values and a fold change larger than 1.5 in the case vs control
comparison were selected for the protein interaction network analysis. Then, overlapping genes
between the selected gene list, the disease risk gene sets, and the proteins in the SIGNOR
database[24] were utilized to query the SIGNOR database and build the interaction network, using
the "connect + add bridge proteins" search mode in The SIGNOR Cytoscape App v1.2.[73]

iii. Gene ontology analysis

iv. Dissociation and 10x Single-Cell RNA sequencing of organoids

Whole organoids were dissociated to the single cell level in preparation for single-cell RNA
sequencing using the Papain Worthington Kit (Worthington, LK003150). All solutions included in
the kit were prepared according to the manufacturer's instructions. 4-6 organoids were transferred
to one well of a low attachment 6 well plate and were washed with PBS without Ca^{2+} and Mg^{2+}.
Organoids were cut with a small scalpel into smaller pieces for easier dissociation. 800 µl of
papain solution (supplied in the kit) was added per well. Samples were incubated at 37°C for
about two hours or until single cell suspension was achieved. Every 15 minutes, the mixture was
pipetted up and down with a cut P1000 pipette tip. Once single cell suspension was reached, 500
µl of inhibitor solution (supplied in the kit) was added to the well. The solution was gently mixed,
filtered through a 70 µm-pore sieve, and transferred to a 15 ml centrifuge tube. The cells were
pelleted by centrifugation at 300 x g for 5 minutes at room temperature. Cell pellets were
resuspended in 500 µl of ice-cold 0.04% BSA diluted in PBS without Ca^{2+} and Mg^{2+}. scRNA-seq
was performed on 4-6 pooled 6-month-old organoids per donor line, per condition (hCS or hSS)
for a total of 48 organoids. A minimum of 10,000 dissociated cells were submitted for sequencing.
The library was prepared using the Chromium platform (10x Genomics) with the 3' gene

expression (3' GEX) V3/V3.1 kit. Libraries were sequenced on an Illumina NovaSeq sequencer

with an S4 flow cell, targeting a minimum depth of 20,000 reads per cell. The lengths for read

parameters Read 1, i7 index, i5 index, and Read 2 were 100, 8, 0, 100, respectively.

iv. Bioinformatic analysis of scRNASeq data

The raw sequencing data, represented as base call (BCL) files produced by the Illumina

sequencing platform, were subjected to demultiplexing and subsequent conversion into FASTQ

format using CellRanger software (version 6.0.0, 10x Genomics) with default parameters. The

software then mapped the FASTQ reads to the human reference genome (GRCh38) with default

parameters. Following this, the 'count' command in the Cell Ranger v6.0.0 software was utilized

for the quantification of gene expression. For alignment and counting, the reference genome

refdata-gex-GRCh38-2020-A was used, which was procured from the official 10x Genomics

website. We performed QC and normalization using the Seurat v3 R package[74]. For QC, we

filtered out low-quality cells using the following criteria: (i) cells with unique gene counts outside

the range of 200 to 6000; (ii) cells with more than 30% mitochondrial gene content; and [75] cells

with less than 2000 unique molecular identifiers[76]. Post QC, we carried out normalization, scaling

the gene expression measurements for each cell by the total expression, multiplied by a scale

factor (10,000 by default), and log-transformed the results. We extracted the expression profiles

of the 338 genes identified by Birrey et al[28], to reduce the dimensionality of the dataset through

principal component analysis (PCA), and identify statistically significant PCs using a JackStraw

permutation test. This was followed by non-linear dimensional reduction using the UMAP (Uniform

Manifold Approximation and Projection) technique for visualization. Cells were clustered based

on their PCA scores using the Shared Nearest Neighbor (SNN) modularity optimization-based

clustering algorithm in Seurat. After dimensionality reduction, we used known marker genes to

guide the clustering of cells. Each cluster was then annotated using cell type markers identified

by Birey et al[28]. Finally, we conducted differential expression analysis across the defined cell

clusters using the FindAllMarkers function in Seurat, which employs a Wilcoxon Rank-Sum test.

The significantly differentially expressed genes were then used to interpret the biological

significance of cell clusters. For each identified cell type, we conducted an enrichment analysis

using the WebGestalt (WEB-based GEne SeT AnaLysis Toolkit) online tool, with the Human

genome (GRCh38) as the reference set, employing a hypergeometric statistical method and the

Benjamini & Hochberg method for multiple test adjustment.

v. Analysis of ChIP-sequencing data from β-estradiol treated rodent brains.

Tracks in bigWig format were downloaded from the GEO dataset GSE144718[33]. Two peaks within

the *NRXN1* gene region in the mm10 genome were visualized using Spark v2.6.2[77]

vi. Generation of long-read sequencing data from post-mortem brain tissue

All aspects of sample processing (tissue handing, RNA extraction, library prep, QC, sequencing

and bioinformatic analysis) was performed as previously described (Flaherty et al).

<u>*Therapeutic Treatments of iNeurons*</u>

i. β-estradiol treatment

β-estradiol was reconstituted in DMSO and subsequently diluted in Brainphys maturation media

for a final concentration of 10 or 30nM. Neurons were treated for 3-5 consecutive days, with fresh

drug or vehicle control replenished daily (in half media changes). iGLUT and iGABA neurons were

treated starting from ~DIV14-18. On the final day of treatment, cells were harvested ~4 hours post

dosage. Chronic treatments for functional experiments extended daily β-estradiol replenishments

for ~14 days, and withdrawn 2-3 days prior to patch-clamp recordings.

ii. Antisense Oligonucleotide treatment

A single HPLC-grade ASO was designed from Qiagen (LNA GapmeR) against the mutant (Ex

20/24) splice junction containing a phosphorothioate modified backbone with or without a 5'-FAM

label for fluorescent visualization. All experiments were performed matched with a non-targeting

ASO as the control group. ASOs were delivered using Lipofectamine RNAiMAX Transfection

Reagent (Thermo, #13778075). The Lipofectamine RNAiMAX was diluted in Opti-MEM Medium

(Thermo, #31985070). ASO was then diluted in Opti-MEM Medium. The diluted ASO was added

in a 1:1 ratio to the diluted Lipofectamine RNAiMAX and incubated at room temperature for 5

minutes. The ASO-lipid complex was added to cells, and incubated for 72 hours post-transfection,

until RNA harvest.

<u>*Molecular Biology and Imaging*</u>

i. RNA extraction and RT-qPCR

For the isolation of RNA, 2D cells were lysed using TRIzol (Life Technologies #15596026) and

purified using the Qiagen miRNeasy Kit (Qiagen Cat# 74106) according to the manufacturer's

instructions. For 3D organoids, pooled (early timepoints) or single organoids were washed and

lysed using TRIzol, by manual homogenization with a pestle in a 1.5mL centrifuge tube. Following

purification, samples were normalized within each experiment (15-50ng) and subsequently used

for RT-qPCR assays using the *Power* SYBR Green RNA-to-Ct 1-Step Kit (Thermo REF 4389986).

Relative transcript abundance was determined using the ΔΔCT method and normalized to the

ACTB housekeeping gene. All primer sequences are listed below.

shRNA, primers and oligonucleotide probe sequences

Target	Sequences (5'-3')	Supplier
RT-qPCR Primer Probes		
NRXN1 (wildtype) Forward	AGAAAGATGCCAAGCACCCA	ThermoFisher
NRXN1 (wildtype) Reverse	CCCATGTCCAGGAGGAGGTA	ThermoFisher
NRXN1-20/24 (mutant) Forward	GCTACCCTGCAGCCAACC	ThermoFisher
NRXN1-20/24 (mutant) Reverse	GACCATACCCGTGGTGCTG	ThermoFisher
ACTB Forward	TGTCCCCCAACTTGAGATGT	ThermoFisher
ACTB Reverse	TGTGCACTTTTATTCAACTGGTC	ThermoFisher
MAP2 Forward	AAACTGCTCTTCCGCTCAGACACC	ThermoFisher
MAP2 Reverse	GTTCACTTGGGCAGGTCTCCACAA	ThermoFisher
NEUROD1 Forward	GGTGCCTTGCTATTCTAAGACGC	ThermoFisher
NEUROD1 Reverse	GCAAAGCGTCTGAACGAAGGAG	ThermoFisher
SLC17A7 Forward	CGCATCATGTCCACCACCAACGT	ThermoFisher
SLC17A7 Reverse	GAGTAGCCGACCACCAACAGCAG	ThermoFisher

SLC17A6 Forward	TCAACAACAGCACCATCCACCGC	ThermoFisher
SLC17A6 Reverse	GTTTCCGGGTCCCAGTTGAATTTGG	ThermoFisher
GAD65 Forward	CTATGACACTGGAGACAAGGC	ThermoFisher
GAD65 Reverse	CAAACATTTATCAACATGCGCTTC	ThermoFisher
DLX5 Forward	ACAGAGACTTCACGACTCCCAG	ThermoFisher
DLX5 Reverse	TGTGGGGCTGCTCTGGTCTA	ThermoFisher

shRNA Sequences

NRXN1 wildtype mRNA (Exon 9)	ATGGAGTGGTGGCATTTAAAT	Sigma
NRXN1 mutant mRNA (Overlapping Exon 20/24)	CGCTACCCTGCAGCCAACCCA	Sigma

Antisense Oligonucleotides (LNA Gapmers)

Non-targeting ASO	AACACGTCTATACGC/36-FAM/	Qiagen
NRXN1-Splice ASO	GGTTGGCTGCAGGGTA/36-FAM/	Qiagen

ii. Bright Field Imaging

For organoid perimeter analyses, brightfield microscope images of organoids were taken with a
2x objective. Image analysis was performed in ImageJ, with best fitting ovals or ellipses were
selected around an organoid, and perimeter was measured.

iii. Immunostaining of 2D cultures

For immunostaining of 2D monocultures, iGLUT and iGABA neurons seeded on acid-etched
coverslips coated with PEI buffered with boric acid and 4x Matrigel. Samples were washed with
DPBS $Ca^{2+}/Mg2^{+}$ and fixed using cold, fresh 16% paraformaldehyde (Life Technologies, #28908),
diluted to 4% for 12 minutes at RT. Coverslips were then blocked and permeabilized with 2%
donkey serum in DPBS $Ca^{2+}/Mg2^{+}$ supplemented with 0.1% Triton-X (Sigma, #93443-100ML)

(blocking buffer), for one hour at RT. Primary antibody solutions were prepared in blocking buffer

and incubated overnight at 4°C. The following day, samples were washed three times with PBS,

and treated with secondary antibodies diluted in blocking buffer, for 1 hour in a dark chamber.

Finally, samples were washed three times, and stained with DAPI for 15min at RT during the final

wash. Coverslips were mounted with antifade (Vectashield #H-1000-10) onto glass slides and

stored at 4°C until imaging using an upright Zeiss LSM 780 confocal microscope.

iv. Immunostaining 3D cultures

For immunostaining of 3D organoids, samples were first washed with DPBS $Ca^{2+}/Mg2^+$ and fixed

using cold 4% PFA for 2 hours at 4°C. Organoids were then washed with DPBS $Ca^{2+}/Mg2^+$

supplemented with 10% sucrose (Sigma, #S0389-500G) for one hour. Sucrose percent was

increased to 20% for an additional hour, and finally 30% for an overnight incubation. The following

day, upon observing the organoids saturated sunk to the bottom of the 1.5mL tube, organoids

were transferred to OCT (Sakura #4583) twice in order to remove residual sucrose solution, and

snap frozen in a cryopreservation mold with dry ice and stored at –80°C until cryosectioned.

Organoids were sectioned at 20 µm thickness onto a Superfrost *Plus* slide (Fisher Scientific, #22-

037-246) and stored at -20°C until immunohistochemistry staining. Boundaries were drawn

around tissue sections with a lipid pen (Fisher Scientific, #NC9545623) to contain solution and

slides were washed once with DPBS with Ca^{2+}/Mg^{2+} to remove excess OCT compound. Slides

were then washed twice with 0.01% Triton-X in DPBS with Ca^{2+}/Mg^{2+} (DPBS-TX) and were

blocked in 2% Donkey Serum in DPBS-TX for one hour at RT. Primary antibody incubation

mixtures were diluted in 2% Donkey Serum in DPBS-TX (see below for a list of primary and

secondary antibodies and dilutions used). 100 µl of primary antibody mixtures was added per

slide, and allowed to incubate overnight in a humidified chamber at 4°C. Pieces of parafilm were

added on top of primary antibody mixtures to further prevent dehydration of slides. The following

day, slides were washed three times with DPBS-TX. Secondary antibodies were diluted in DPBS-

TX and were allowed to incubate for an hour. Slides were again washed three times with DPBS-

TX. A 1:2000 DAPI solution diluted in DPBS-TX was applied to slides for 10 minutes. Secondary

and DAPI stainings took place in a covered chamber in order to minimize exposure to light. Slides

were washed twice with DPBS-TX, and then once with DPBS with Ca^{2+} and Mg^{2+}. Slides were

mounted with coverslips using Aqua Polymount solution (Polysciences #18606-20). Slides were

stored at -20 °C until they were imaged.

Target	Antibody Dilution	Supplier/ CAT#
Primary Antibodies		
MAP2AB	1:500	Sigma, #M1406
(Rabbit) vGAT	1:500	Synaptic Systems, #135-303
(Rabbit) vGLUT1	1:500	Synaptic Systems, #131-002
SOX2	1:200	Santa Cruz Biotechnology, #sc-365964
(Rabbit) KI67	1:300	Abcam, #ab15580
(Chicken) MAP2	1:250	Aves Labs, #MAP
DAPI	1:1000	Sigma, #D9542
Secondary Antibodies		
(Donkey) 488-Mouse	1:200	Jackson ImmunoResearch, # 711-545-152
(Donkey) 568-Rabbit	1:200	Abcam, #ab175700
(Donkey) 647-Chicken	1:200	Jackson ImmunoResearch, # 703-605-155
(Donkey) 568-Mouse	1:500	Invitrogen (A10037)
(Donkey) 488-Rabbit	1:1000	Life Technologies (A-21206)

Statistics

No statistical power estimation analyses were used to predetermine sample sizes, where were

chosen to match previous publications[14,21] and field standards. All experimental statistics were

performed in Prism *v10.1.1* and R *v4.2.0.* Bioinformatic analyses were performed in R *v3.5.3* (Bulk

RNASeq DEG), *v4.1.2* (Bulk RNASeq GO) and *v4.1.2* (scRNAseq).

Data and Code Availability

All source donor hiPSCs have been deposited at the Rutgers University Cell and DNA Repository

(study 160; http://www.nimhstemcells.org/) and all bulk and single-cell transcriptome sequencing

data are being prepared for deposits to GEO. To facilitate improved reproducibility of our data
analyses, custom scripts will be deposited to github post peer-review.

Acknowledgements

MBF is supported by a Gilliam Fellowship from the Howard Hughes Medical Institute. This work
is supported by the National Institute of Mental Health grants RO1 MH121074-02 (KJB, GF and
PAS) and RO1 MH125579-02 (GF and KJB). We thank the Stem Cell Engineering Core at the
Icahn School of Medicine at Mount Sinai. The authors are grateful to the labs of Nan Yang (Ruiqi
Hu and Xiaoting Zhou) and Samuele Marro (Madel Durens) for assistance in primary glial
preparations. The authors thank Kayla G. Townsley and Mark G. Baxter for advice on statistical
testing. We acknowledge Daniel Weinberger and the Lieber Institute for Brain Development at
Johns Hopkins School of Medicine, for sharing post-mortem materials. Finally, the authors thank
all members of the Brennand, Slesinger and Fang labs for critical feedback and discussions
throughout the course of this work.

Author Information

MBF, SK, ANM, RO, CP, AP performed and analyzed experiments supervised by PAS and KJB.
YZ and YF performed bioinformatic analyses supervised by GF. SG contributed to RNASeq
analyses. SW produced virus for the generation of iGABA neurons. EKF processed post-mortem
tissue and generated long-read data. IKP assisted in statistical analyses. MBF, GF, PAS and KJB
wrote the paper with input from all authors.

Ethics declarations / Competing interest statement.

All authors have no competing interests to declare.

Figures

Figure 1. *Changes in NRXN1 splicing are conserved across human iPSC derived glutamatergic (iGLUT) and GABAergic (iGABA) neurons.* (**a**) Brief description of clinical information of all hiPSC

lines used in this study, and (**b**) schematic of *NRXN1* gene structure with red and blue shades
corresponding on 3'-Del and 5'-Del genotypes, respectively. Arrows indicate relative internal
promoter positions with purple bars corresponding to alpha exons, and green bars correlating to
beta exons. (**c, h**) Induction timeline and factors to generate iGLUT and iGABA neurons, with
immunostaining validation of neuronal identity (MAP2), glutamate identity (vGLUT1), and GABA
identity (vGAT). (**d, i**). Gene expression panel confirming abundance of neuronal markers and
neurotransmitter identity in iGLUT (n: Control = 6/2; 5'-Del = 6/2; 3'-Del = 6/2 | 1 batch) and in
iGABA neurons (n/d: Control = 5/2; 5'-Del = 6/2; 3'-Del = 6/2 | 1 batch). (**e, j**) Volcano plots of
differential gene expression (DE) analysis across both genotypes in iGLUT and iGABA neurons.
Vertical dashed lines represent DE genes above/below 1.5 Log_2FC. Horizontal dashed lines
represent FDR = 0.1 cutoff (lower) and Bonferroni corrected cutoff (upper). (**f, k**) Overlap of DEGs
between genotypes and enrichment of genes across neuropsychiatric disorders. (**g, l**) Percent
usage of alpha *NRXN1* exon usage in iGLUT, compared via a 1-way ANOVA, with Dunnett's test
($F_{2, 15}$ = 25.70; 5'-Del p = 0.0031, 3'-Del p = 0.0086), and iGABA neurons ($F_{2, 14}$ = 92.92; 5'-Del p
< 0.001, 3'-Del p = 0.0004), with beta *NRXN1* exon usage, calculated by subtracting alpha-
specific reads against total reads. Data represented as mean ± sem. n reported as n (samples)/d
(donors) | independent batches.

Figure 2. *Spontaneous, passive, and excitable properties are minimally changed from NRXN1$^{+/-}$ induced neurons.* (**a, f**) Timelapse of multi-electrode array recordings every 2-3 days apart starting near ~DIV12 for a single representative induction. Tiles represent averaged wMFR values across genotypes during a single recording session. (**b**) MEA quantification of iGLUT neuronal activity

(n/d: Control = 52/2; 5'-Del = 42/2; 3'-Del = 38/2 | 3 batches) at WPI4 ($F_{2,\,129}$ = 11.64; 5'-Del p =

0.0001, 3'-Del p = 0.0011), and (**c**) WPI6 ($F_{2,\,129}$ = 4.560; 5'-Del p = 0.0238, 3'-Del p = 0.0211).

(**d**) intrinsic properties of iGLUT neurons (n/d: Control = 51/2; 5'-Del = 51/2; 3'-Del = 48/2 | 8

batches): compared via 1-way ANOVA, with Dunnett's test, Capacitance ($F_{2,\,147}$ = 1.505; 5'-Del

p = 0.1565, 3'-Del p = 0.7633), Input resistance ($F_{2,\,147}$ = 1.311; 5'-Del p = 0.1949, 3'-Del p =

0.7906), and RMP (n/d: Control = 16/2; 5'-Del = 14/2; 3'-Del = 16/2 | 2 batches) compared via 1-

way ANOVA, with Dunnett's test ($F_{2,\,43}$ = 0.8234; 5'-Del p = 0.6378, 3'-Del p = 0.8468). (**e**) Input-

output curves of excitable properties (n/d: Control = 16/2; 5'-Del = 14/2; 3'-Del = 16/2 | 2 batches),

with representative traces (right), compared via Step x Genotype 2-way ANOVA; Dunnett's Test

($F_{26,\,559}$ = 1.690, 5'-Del p < 0.01 at Step 6, 7 and 9, 3'-Del p = n.s. on all steps). (**g**) MEA

quantification of iGABA neuronal activity at WPI2 (n/d: Control = 71/2; 5'-Del = 48/2; 3'-Del = 55/2

| 5 batches) compared via 1-way ANOVA, with Dunnett's test ($F_{2,\,171}$ = 5.422; 5'-Del p = 0.0381,

3'-Del p = 0.0047), and (**h**) WPI5 (n/d: Control = 27/4; 5'-Del = 23/2; 3'-Del = 23/2 | 2 batches)

compared via 1-way ANOVA, with Dunnett's test ($F_{2,\,70}$ = 2.132; 5'-Del p = 0.0809, 3'-Del p =

0.4134). (**i**) intrinsic properties of iGABA neurons (n/d: Control = 39/2; 5'-Del = 34/2; 3'-Del = 37/2

| 6 batches): compared via 1-way ANOVA, with Dunnett's test, Capacitance ($F_{2,\,107}$ = 0.04256; 5'-

Del p = 0.9425, 3'-Del p = 0.9941), Input resistance (n/d: Control = 39/2; 5'-Del = 34/2; 3'-Del =

36/2 | 6 batches, $F_{2,\,106}$ = 1.451; 5'-Del p = 0.8162, 3'-Del p = 0.1703) and RMP (n/d: Control =

18/2; 5'-Del = 16/2; 3'-Del = 16/2 | 2 batches) compared via 1-way ANOVA, with Dunnett's test

($F_{2,\,47}$ = 0.02842; 5'-Del p = 0.9893, 3'-Del p = 0.9601). (**j**) Input-output curves of excitable

properties (n/d: Control = 16/2; 5'-Del = 14/2; 3'-Del = 16/2 | 2 batches), with representative traces

(right), compared via Step x Genotype 2-way ANOVA; Dunnett's Test ($F_{26,\,572}$ = 0.5137, 5'-Del

and 3'-Del p = n.s. on all steps). Data represented as mean ± sem. n reported as n (samples)/d

(donors) | independent batches.

iGLUT

a Representative sEPSCs

b sEPSC Frequency

c sEPSC Amplititude

d sEPSC Score

e SynGO Score

f Trans-Synaptic Neurexin-1 Binding Partners

iGABA

g Representative sIPSCs

h sIPSC Frequency

i sIPSC Amplititude

j sIPSC Score

k SynGO Score

l Trans-Synaptic Neurexin-1 Binding Partners

Figure 3: *Divergent impact on neurotransmission from NRXN1$^{+/-}$ induced neurons.* (**a**) Representative traces of iGLUT sEPSCs. (**b**) Cumulative probabilities and log-scaled cell averages of inter-event-internals (IEIs) across genotypes (n/d: Control = 39/4; 5'-Del = 33/3; 3'-Del = 29/2 | 6), compared by Levene's Test for averaged distributions (5'-Del F = 46.635, df = 1, *p* = 1.27E-11; 3'-Del F = 17.928, df = 1, *p* = 2.44E-5), and 1-way ANOVA, with Dunnett's test for inset (F$_{2,\ 98}$ = 3.117; 5'-Del *p* = 0.0285, 3'-Del *p* = 0.2929). (**c**) Cumulative probabilities and log-scaled cell averages of amplitude size across genotypes, compared by Levene's Test for averaged distributions (5'-Del F = 14.15, df = 1, *p* = 1.94E-4; 3'-Del F = 1.0851, df = 1, *p* = 0.2982),

and 1-way ANOVA, with Dunnett's test for inset ($F_{2, 98}$ = 2.839; 5'-Del p = 0.0634, 3'-Del p =

0.9994). (**d**) Transformed p-values of Levene's Test and (**e**) SynGO gene-set averaged $\log_2$FC

values across pre- or post- synaptic genes. (**f**) Gene expression panel (z-scores), of canonical

NRXN1 binding partners in iGLUT neurons, boxes indicate reaching genome wide significance.

(**g**) Representative traces of iGABA sIPSCs. (**h**) Cumulative probabilities and log-scaled cell

averages of inter-event-internals (IEIs) across genotypes (n/d: Control = 26/2; 5'-Del = 25/3; 3'-

Del = 22/2 | 4), compared by Levene's Test for averaged distributions (5'-Del F = 3.4002, df = 1,

p = 0.0653; 3'-Del F = 16.501, df = 1, p = 5.00E-05), and 1-way ANOVA, with Dunnett's test for

inset ($F_{2, 70}$ = 0.7986; 5'-Del p = 0.7986, 3'-Del p = 0.339). (**i**) Cumulative probabilities and log-

scaled cell averages of amplitude size across genotypes, compared by Levene's Test for

averaged distributions (5'-Del F = 0.1, df = 1, p = 0.9204; 3'-Del F = 0.0006, df = 1, p = 0.9801),

and 1-way ANOVA, with Dunnett's test for inset ($F_{2, 70}$ = 0.08143; 5'-Del p = 0.9431, 3'-Del p =

0.9904). (**j**) Transformed p-values of Levene's Test and (**k**) SynGO gene-set averaged $\log_2$FC

values across pre- or post- synaptic genes. (**l**) Gene expression panel (z-scores), of canonical

NRXN1 binding partners in iGABA neurons, boxes indicate reaching genome wide significance.

*p < 0.05, **p < 0.01, ***p <0.001, Wilcoxon's rank sum test, FDR = 0.05. Data represented as

mean ± sem. n (samples)/d (donors) | independent batches.

Figure 4: *NRXN1$^{+/-}$ impacts patterning of region-specific human brain organogenesis.* (**a, h**)

Timeline of organogenesis protocols (**b, i**), UMAPs of hCO and hSO organoid samples sequenced

at 6 months, annotated by cell clusters, and (**c, j**) relative proportions of cell clusters across

genotypes, (hCO = 47,460 cells) and (hSO = 35,563 cells). (**d, k**) validation of regionalization

across forebrain (*FOXG1*), dorsal (*EMX1*) and ventral (*DLX2*) regions, with *NRXN1* expression

across all cells. (**e, l**) Representative images of neural rosettes, immunostained with SOX2, KI67

and DAPI, across genotypes with (**f, m**) quantified intensity ratios for hCO (n/d: Control = 92/2; 5'-

Del = 84/3; 3'-Del = 108/2 | across 3-4 organoids per group from 2 batches), compared via 1-way

ANOVA, with Dunnett's test ($F_{2, 281}$ = 4.490; 5'-Del p = 0.4094, 3'-Del p = 0.0063), and (n/d: Control

= 73/2; 5'-Del = 79/3; 3'-Del = 79/2| across 3-4 organoids per from 2 batches), compared via 1-

way ANOVA, with Dunnett's test ($F_{2, 228}$ = 9.366; 5'-Del p = 0.0004, 3'-Del p = 0.0004). (g, n) Gene

ontological analysis results using DEGs from scRNASeq. *P < 0.05, **P < 0.01, ***P <0.001,

Wilcoxon's rank sum test, FDR = 0.05. Data represented as mean ± sem. n reported as n

(samples)/d (donors) | independent batches.

Figure 5: *Isogenic recapitulation and rescue of neurotransmission phenotypes.* (**a**) Representative traces of iGLUT WT knockdown effects, with (**b**) cumulative probabilities of sEPSC IEI distributions, with insets of cell-averaged IEI and amplitude measures (n/d: shNT = 12/1; shWT = 7/1 | 2 batches). Curves were compared via a Levene's test (F = 70.78, df =1, *p* <

2.2E-16), and insets were compared via a Student's t-test (IEI t=1.166, df=19, p = 0.2579; AMP

t=0.5661, df=19, p = 0.578). (**c**) SynGO (biological process) sunburst plots showing enrichment

of DEGs associated with synaptic function for iGLUT WT-KD compared to a brain expressed

background via Fisher's exact test (Fold =1.274161, p = 0.1049). (**d**) Representative traces of

iGLUT MT knockdown effects, with (**e**) cumulative probabilities of sEPSC IEI distributions, with

insets of cell-averaged IEI and amplitude measures (n/d: shNT = 14/1; shMT = 18/1 | 2 batches).

Curves were compared via a Levene's test (F = 230.6; df = 1, p < 2.2E-16), and insets were

compared via a Student's t-test (IEI t=2.471, df=30, p = 0.0194; AMP t=1.812, df=30, p = 0.08).

(**f**) SynGO (biological process) sunburst plots showing enrichment of DEGs associated with

synaptic function for iGLUT MT-KD compared to a brain expressed background via Fisher's exact

test (Fold = 1.214533, p = 0.4541). (**g**) Representative traces of iGABA WT knockdown effects,

with (**h**) cumulative probabilities of sIPSC IEI distributions, with insets of cell-averaged, IEI and

amplitude measures (n/d: shNT = 16/3; shWT = 19/3 | 2 batches). Curves were compared via a

Levene's test (F = 30.879, df =1, p < 3.66E-08), and insets were compared via a Student's t-test

(IEI t=2.768, df=33, p = 0.0092; AMP t=0.7856, df=33, p = 0.4377). (**i**) SynGO (biological process)

sunburst plots for iGABA WT-KD compared to a brain expressed background via Fisher's exact

test (Fold = 1.234861, p = 1.347E-06). (**j**) Representative traces of iGABA MT knockdown effects,

with (**k**) cumulative probabilities of sIPSC IEI distributions, with insets of cell-averaged, IEI and

amplitude measures (n/d: shNT = 16/1; shMT = 19/1 | 2 batches). Curves were compared via a

Levene's test (F = 4.1324; df = 1, p < 0.04226), and insets were compared via a Student's t-test

(IEI t=1.305, df=25, p = 0.2038; AMP t=0.9292, df=25, p = 0.3617). (**l**) SynGO (biological process)

sunburst plots for iGABA WT-KD compared to a brain expressed background via Fisher's exact

test (Fold = 1.347225, p = 2.694E-04). Data represented as mean ± sem. n reported as n

(samples)/d (donors) | independent batches.

Figure 6: *Precise therapeutic targeting of GOF- and LOF-NRXN1*[+/-] *in iGLUT neurons.* (**a**)

Schematic of a *NRXN1* ASO matrix, with the selective sequence targeting 20/24 splice junction.

(**b**) RT-qPCR of MT *NRXN1* gene expression in 3'-Del patients' post-acute treatment for 72hrs

(n/d: ASO-NT = 8/2; ASO-MT = 10/2 | batches),compared via Student's t-test t=12.91, df=16 $p <$

0.0001. (**c**) RNASeq analysis on ASO treated patients (n/d: ASO-NT = 4/2; ASO-MT = 4/2 |

representative batch), demonstrating biological process and (**d**) cellular compartment GO terms

demonstrating enrichment ($\log_{10}$[adj p value]) of selected synapse related pathways. (**e**) Model

and proposed mechanism of rescue for GOF patients expressing mutant isoforms. (**f**) RT-qPCR

of WT *NRXN1* gene expression in 5-Del patients, post-acute treatment (3-5 days) (n/d: DMSO =

19/2; β-estradiol = 18/2 | 3 batches) compared via Student's t-test t=2.293, df=35p = 0.028. (**g**)

Quantification of iGLUT neuronal activity at WPI3 across vehicle and treatment (n/d: DMSO =

72/3; β-estradiol = 74/3 | 3 batches)) compared via Student's t-test (t=2.804, df=144, p = 0.0057).

(**h**) Representative patch-clamp traces and (**i**) cumulative probabilities of sEPSC IEI distributions,

with insets of cell-averaged, IEI and amplitude measures (n/d: Control + DMSO = 18/1; 5'-Del +

DMSO = 33/2 5'-Del + β-estradiol = 35/2). Curves were compared via Levene's test (Control +

DMSO v 5'-Del + DMSO F = 13.151; df = 1, p = 2.921E-4; 5'-Del + DMSO v 5'-Del + β-estradiol

F = 14.359; df = 1, p = 1.54E-04; Control + DMSO v 5'-Del + β-estradiol, F = 0.0141; df = 1, p =

0.955). Insets were compared via Student's t-test (IEI t=1.161, df=66, p = 0. 0.2499). (**j**) Model

and proposed mechanism of rescue for LOF patients ameliorating loss of wildtype isoforms. Data

represented as mean ± sem. n reported as n (samples)/d (donors) | independent batches.

		CONTROL				5'-DEL		3'-DEL		Genotype
										Data Point Color
Reference	Experiment	553-3	2607-4	3182-3	690-2	972-5	973-5	581-5	641-6	Donor Cell Line-Clone
Fig. 1 & ED Fig. S2	iGLUT/ iGABA generation	✓	✓	✓	✓	✓	✓	✓	✓	
	Immunostaining	✓	✓							
	Case/Control RNASeq	✓	✓			✓	✓	✓	✓	
Fig. 2 & ED Fig. S3	Case/Control MEA iGLUT	✓	✓			✓	✓	✓	✓	
	Case/Control MEA iGABA	✓	✓	✓	✓	✓	✓	✓	✓	
	Patch-Clamp iGLUT (Excitability)	✓	✓			✓	✓	✓	✓	
	Patch-Clamp iGABA (Excitability)	✓	✓			✓	✓	✓	✓	
ED Fig. S4	KCC2 MEA (GABAZine)			✓	✓					
Fig. 3	Patch-Clamp iGLUT (Neurotransmitter release)	✓	✓	✓	✓	✓	✓	✓	✓	
Fig. 3	Patch-Clamp iGABA (Neurotransmitter release)	✓	✓			✓	✓	✓	✓	
ED Fig S5	TTX Experiments iGLUT	✓	✓			✓	✓	✓	✓	
ED Fig S5	TTX Experiments iGABA	✓	✓		✓	✓	✓	✓	✓	
Fig. 4, ED Fig. S6 & S7	Organoid generation	✓	✓			✓	✓	✓	✓	
	Organoid characterization	✓	✓			✓	✓	✓	✓	
	Organoid rosette imaging	✓	✓			✓	✓	✓	✓	
	Organoid single cell sequencing	✓*	✓			✓	✓	✓	✓	
Fig. 5 & ED FigS8.	shRNA testing iGLUT-WT	✓	✓	✓	✓					
	shRNA patch-clamp iGLUT-WT				✓					
	shRNA testing iGABA-WT	✓	✓	✓	✓					
	shRNA patch-clamp iGABA-WT		✓	✓	✓					
	shRNA testing iGLUT-MT							✓	✓	
	shRNA patch-clamp iGLUT-MT							✓		
	shRNA testing iGABA-MT							✓	✓	
	shRNA patch-clamp iGABA-MT							✓		
	shRNA RNASeq			✓	✓			✓		
Fig. 6	Beta-Estradiol				✓	✓	✓			
Fig. 6	Anti-sense Oligos							✓	✓	

Extended Data Figure 1: *Specific cell lines used in each experiment throughout this study.* Reference table of all cell lines-clones used per experiment with corresponding data point colors. Check marks indicate inclusion for experiments, *represents dropped donor due to failed QC.

	iGLUT 5'-Del		iGLUT 3'-Del		iGABA 5'-Del		iGABA 3'-Del	
	Overlapped DEGs with risk genes	Overlapped DEGs with risk genes & in SIGNOR database	Overlapped DEGs with risk genes	Overlapped DEGs with risk genes & in SIGNOR database	Overlapped DEGs with risk genes	Overlapped DEGs with risk genes & in SIGNOR database	Overlapped DEGs with risk genes	Overlapped DEGs with risk genes & in SIGNOR database
ASD	11	**7**	4	**0**	3	**1**	6	**2**
BP	16	**9**	3	**1**	2	**1**	5	**1**
SCZ	26	**14**	10	**1**	9	**3**	14	**3**
Total	37	**22**	12	**2**	9	**3**	18	**4**

Extended Data Figure 2: *Extended transcriptomics analysis on disease risk associated genes.*
(**a**) Summary table of overlapping DEGs with risk enrichments across publicly curated datasets for autism (ASD), bipolar disorder (BD) and schizophrenia. Interaction maps of risk genes for (**b**) 3'-Del iGLUT, (**c**) 3'-Del iGABA, (**d**) 5'-Del iGABA and (**e**) 5'-iGLUT. Sample information correspond to Fig. 1

Extended Data Figure 3: *Extended data excitable properties of 5'-Del and 3'-Del neurons.* Voltage-gated potassium and channel kinetics across genotypes for (**a**) iGLUT and (**b**) iGABA neurons, sample information correspond Fig. 2. Data represented as mean ± sem.

Extended Data Figure 4: *Extended KCC2 related data from immature GABA neurons.* (**a**)
Transcriptomic comparison of *SLC12A5* expression across DIV14 and DIV35 RNASeq timepoints
sample information correspond Fig. 1 (iGABA), compared via mixed effects model-time ($F_{1, 28}$ =
499.1 *p* <0.0001). (**b,c**) MEA tests from pre- and post- treatment of 10μM GABAzine. (n/d = 28/2
| 1 representative batch) compared via paired student's t-test for time-linked comparison (t=9.739,
df=27; *p* < 0.0001) and unpaired student's t-test for pre/post activity foldchange (t=2.380, df=54;
0.0209). Data represented as mean ± sem. n reported as n (samples)/d (donors) | independent
batches.

Extended Data Figure 5: *Miniature neurotransmitter release (+TTX) from iGLUT and iGABA neurons.* Comparative GLUT mEPSC kinetics of (**a**) IEI (n/d: Control = 22/2; 5'-Del = 18/2; 3'-Del = 7/2 | 4 batches) compared via 1-way ANOVA, with Dunnett's test ($F_{2, 44}$ = 2.880; 5'-Del *p* = 0.0411, 3'-Del *p* = 0.8229, and (**b**) amplitude size (n/d: Control = 24/2; 5'-Del = 22/2; 3'-Del = 10/2 | 4 batches) compared via 1-way ANOVA, with Dunnett's test ($F_{2, 53}$ = 0.1996; 5'-Del *p* = 0.7806, 3'-Del *p* = 0.9972). (**c**) Comparative iGABA mIPSC kinetics of IEI (n/d: Control = 10/2; 5'-Del = 9/2; 3'-Del = 8/2 | 3 batches) compared via 1-way ANOVA, with Dunnett's test ($F_{2, 24}$ = 0.9632; 5'-Del *p* = 0.6452, 3'-Del *p* = 0.7565, and (**d**) amplitude size (n/d: Control = 13/2; 5'-Del = 10/2; 3'-Del = 9/2 | 3 batches) compared via 1-way ANOVA, with Dunnett's test ($F_{2, 29}$ = 0.5097; 5'-Del *p* = 0.5141, 3'-Del *p* = 0.8618). Data represented as mean ± sem. n reported as n (samples)/d (donors) | independent batches.

Extended Data Figure 6: *Extended data on human organoid generation and characterization.*

(**a,g**) Representative images of hiPSC aggregation and immature spheroids post dislodging. (**b,h**)

Normalized organoid perimeters over time (relative to averaged control) for hCO (n/d: Control =

136/2; 5'-Del = 128/2; 3'-Del = 150/2 | 2 batches) and hSO (n/d: Control = 118/2; 5'-Del = 92/2;

3'-Del = 120/2 | 2 batches). (**c**) hCO RT-qPCR results from 4-month organoids of genes for

pluripotency, neuronal, and cell-type specific markers (n/d: Control = 4/2; 5'-Del = 4/2; 3'-Del =

4/2 | 2 batches), compared via 1-way ANOVA, with Dunnett's test (*SOX2* $F_{2, 9}$ = 3.921, 5'-Del p =

0.065, 3'-Del p = 0.9981; *MAP2AB* $F_{2, 9}$ = 0.7201, 5'-Del p = 0.5479, 3'-Del p = 0.4688; *vGLUT1*

$F_{2, 9}$ = 1.594, 5'-Del p = 0.6353, 3'-Del p = 0.6331; *vGLUT2* $F_{2, 9}$ = 1.710, 5'-Del p = 0.3992, 3'-Del

p = 0.7901). (**d,e**) hCO RT-qPCR results of *NRXN1* WT and MT expression (n/d: Control = 4/2;

5'-Del = 4/2; 3'-Del = 4/2 | Representative batch) compared via 1-way ANOVA, with Dunnett's test

(*NRXN1-WT* $F_{2, 9}$ = 0.5873, 5'-Del p = 0.9256, 3'-Del p = 0.6961) (**f**) Comprehensive gene

expression panel across sub-clusters of hCO samples across neuronal, cortical, subpallial and

astroglia markers. (**i**) hSO RT-qPCR results from 4-month organoids of genes for pluripotency,

neuronal, and cell-type specific markers (n/d: Control = 4/2; 5'-Del = 4/2; 3'-Del = 4/2 | 2 batches),

compared via 1-way ANOVA, with Dunnett's test (*SOX2* $F_{2, 9}$ = 5.683, 5'-Del p = 0.1563, 3'-Del p

= 0.0152; *MAP2AB* $F_{2, 9}$ = 4.800, 5'-Del p = 0.0875, 3'-Del p = 0.0285; *GAD65* $F_{2, 9}$ = 1.641, 5'-

Del p 0.3573, 3'-Del p = 0.8806; *DLX5* $F_{2, 9}$ = 0.01755, 5'-Del p = 0.981, 3'-Del p = 0.9809). (**j,k**)

hSO RT-qPCR results of *NRXN1* WT and MT expression (n/d: Control = 4/2; 5'-Del = 4/2; 3'-Del

= 4/2 | Representative batch) compared via 1-way ANOVA, with Dunnett's test (*NRXN1-WT* $F_{2, 9}$

= 4.818, 5'-Del p = 0.7447, 3'-Del p = 0.0824) (**l**) Comprehensive gene expression panel across

sub-clusters of hSO samples across neuronal, cortical, subpallial and astroglia markers. Data

corresponds to Fig 4. Data represented as mean ± sem. n (samples)/d (donors) | independent

batches.

Extended Data Figure 7: *Extended analysis on astroglia subclusters from hCO and hSOs.* (a)

hCO subcluster and (b) hCO subcluster GO analysis of overlapped terms with hCO-glutamatergic

and hSO-GABAergic enrichments. *P < 0.05, **P < 0.01, ***P <0.001, Wilcoxon's rank sum test,

FDR = 0.05.

tailed t-test (t=4.186, df=9, *p* = 0.0024), and (b) MT *NRXN1* (n/d: shNT = 8/2; shMT = 7/2)

compared via two-tailed t-test (t=9.342, df=13, *p* < 0.0001). (c,d) iGLUT RNASeq DEG plots of

WT (n/d: shNT = 3/1; shWT = 3/1) and MT (n/d: shNT = 2/1; shMT = 2/1) knockdowns performed

in isogenic analyses. (a) iGABA RT-qPCR of shRNA knockdown on WT *NRXN1* (n/d: shNT =

8/3; shWT = 8/3) compared via two-tailed t-test (t=5.294, df=15, *p* < 0.0001), and (b) MT *NRXN1*

(n/d: shNT = 9/2; shMT = 10/2) compared via two-tailed t-test (t=2.011, df=17, *p* = 0.0604). (c,d)

iGABA RNASeq DEG plots of WT (n/d: shNT = 3/1; shWT = 3/1) and MT (n/d: shNT = 2/1; shMT

= 2/1 | 2-4 wells pooled across 2 batches) knockdowns performed in isogenic analyses. Data

represented as mean ± sem. n (samples)/d (donors).

Extended Data Figure 9: *In-vivo validation of MT isoform expression from an unrelated autism NRXN1$^{+/-}$ patient.* (a) Schematic of novel NRXN1 autism patient, and GOF therapeutic targeting pipeline, with (b) schematic of the *NRXN1α* isoform structures, with each row representing a unique *NRXN1α* isoform and each column representing a *NRXN1* exon. The colored isoforms (navy, wildtype; peach, patient-specific) are spliced into the transcript while the blank exons are spliced out. (c) The abundance of each *NRXN1α* isoform by sample.

Data from Gegenhuber et al., *Nature*, 2022
Fold = 3.61, *p* = 9.39E-07, FDR = 9.33E-05

999 **Extended Data Figure 10:** *ChIP-seq enrichment of ER1 binding at NRXN1 loci in rodent brain.*

1000 (a) Female and (b) male mus musculus ChIP tracts of *NRXN1* locus, with red dashed areas

1001 highlighting binding enrichment across vehicle and estradiol treated mice.

1002

References

1   Sudhof, T. C. Synaptic Neurexin Complexes: A Molecular Code for the Logic of Neural Circuits. *Cell* **171**, 745-769 (2017). https://doi.org:10.1016/j.cell.2017.10.024

2   de Wit, J. & Ghosh, A. Specification of synaptic connectivity by cell surface interactions. *Nat Rev Neurosci* **17**, 22-35 (2016). https://doi.org:10.1038/nrn.2015.3

3   Yang, X. *et al.* Widespread Expansion of Protein Interaction Capabilities by Alternative Splicing. *Cell* **164**, 805-817 (2016). https://doi.org:10.1016/j.cell.2016.01.029

4   Schreiner, D. *et al.* Targeted combinatorial alternative splicing generates brain region-specific repertoires of neurexins. *Neuron* **84**, 386-398 (2014). https://doi.org:10.1016/j.neuron.2014.09.011

5   Fuccillo, M. V. *et al.* Single-Cell mRNA Profiling Reveals Cell-Type-Specific Expression of Neurexin Isoforms. *Neuron* **87**, 326-340 (2015). https://doi.org:10.1016/j.neuron.2015.06.028

6   Marshall, C. R. *et al.* Contribution of copy number variants to schizophrenia from a genome-wide study of 41,321 subjects. *Nat Genet* **49**, 27-35 (2017). https://doi.org:10.1038/ng.3725

7   Matsunami, N. *et al.* Identification of rare recurrent copy number variants in high-risk autism families and their prevalence in a large ASD population. *PLoS One* **8**, e52239 (2013). https://doi.org:10.1371/journal.pone.0052239

8   Moller, R. S. *et al.* Exon-disrupting deletions of NRXN1 in idiopathic generalized epilepsy. *Epilepsia* **54**, 256-264 (2013). https://doi.org:10.1111/epi.12078

9   Ching, M. S. *et al.* Deletions of NRXN1 (neurexin-1) predispose to a wide spectrum of developmental disorders. *Am J Med Genet B Neuropsychiatr Genet* **153B**, 937-947 (2010). https://doi.org:10.1002/ajmg.b.31063

10   Huang, A. Y. *et al.* Rare Copy Number Variants in NRXN1 and CNTN6 Increase Risk for Tourette Syndrome. *Neuron* **94**, 1101-1111 e1107 (2017). https://doi.org:10.1016/j.neuron.2017.06.010

11   Grayton, H. M., Missler, M., Collier, D. A. & Fernandes, C. Altered social behaviours in neurexin 1alpha knockout mice resemble core symptoms in neurodevelopmental disorders. *PLoS One* **8**, e67114 (2013). https://doi.org:10.1371/journal.pone.0067114

12   Pak, C. *et al.* Human Neuropsychiatric Disease Modeling using Conditional Deletion Reveals Synaptic Transmission Defects Caused by Heterozygous Mutations in NRXN1. *Cell stem cell* **17**, 316-328 (2015). https://doi.org:10.1016/j.stem.2015.07.017

13   Pak, C. *et al.* Cross-platform validation of neurotransmitter release impairments in
schizophrenia patient-derived NRXN1-mutant neurons. *Proc Natl Acad Sci U S A* **118**
(2021). https://doi.org:10.1073/pnas.2025598118

14   Flaherty, E. *et al.* Neuronal impact of patient-specific aberrant NRXN1alpha splicing. *Nat*
Genet **51**, 1679-1690 (2019). https://doi.org:10.1038/s41588-019-0539-z

15   Boxer, E. E. & Aoto, J. Neurexins and their ligands at inhibitory synapses. *Front Synaptic*
Neurosci **14**, 1087238 (2022). https://doi.org:10.3389/fnsyn.2022.1087238

16   Gomez, A. M., Traunmüller, L. & Scheiffele, P. Neurexins: molecular codes for shaping
neuronal synapses. *Nature Reviews Neuroscience* **22**, 137-151 (2021).
https://doi.org:10.1038/s41583-020-00415-7

17   Taoufik, E., Kouroupi, G., Zygogianni, O. & Matsas, R. Synaptic dysfunction in
neurodegenerative and neurodevelopmental diseases: an overview of induced pluripotent
stem-cell-based disease models. *Open Biology* **8**, 180138 (2018).
https://doi.org:10.1098/rsob.180138

18   Ho, S. M. *et al.* Rapid Ngn2-induction of excitatory neurons from hiPSC-derived neural
progenitor cells. *Methods* **101**, 113-124 (2016).
https://doi.org:10.1016/j.ymeth.2015.11.019

19   Barretto, N. *et al.* ASCL1- and DLX2-induced GABAergic neurons from hiPSC-derived
NPCs. *J Neurosci Methods* **334**, 108548 (2020).
https://doi.org:10.1016/j.jneumeth.2019.108548

20   Zhang, Y. *et al.* Rapid single-step induction of functional neurons from human pluripotent
stem cells. *Neuron* **78**, 785-798 (2013). https://doi.org:10.1016/j.neuron.2013.05.029

21   Schrode, N. *et al.* Synergistic effects of common schizophrenia risk variants. *Nat Genet*
51, 1475-1485 (2019). https://doi.org:10.1038/s41588-019-0497-5

22   Yang, N. *et al.* Generation of pure GABAergic neurons by transcription factor
programming. *Nat Methods* (2017). https://doi.org:10.1038/nmeth.4291

23   Seah, C. *et al.* Modeling gene x environment interactions in PTSD using human neurons
reveals diagnosis-specific glucocorticoid-induced gene expression. *Nat Neurosci* **25**,
1434-1445 (2022). https://doi.org:10.1038/s41593-022-01161-y

24   Lo Surdo, P. *et al.* SIGNOR 3.0, the SIGnaling network open resource 3.0: 2022 update.
Nucleic Acids Res **51**, D631-d637 (2023). https://doi.org:10.1093/nar/gkac883

25   Virtanen, M. A., Uvarov, P., Mavrovic, M., Poncer, J. C. & Kaila, K. The Multifaceted Roles
of KCC2 in Cortical Development. *Trends in neurosciences* **44**, 378-392 (2021).
https://doi.org:https://doi.org/10.1016/j.tins.2021.01.004

Sohal, V. S. & Rubenstein, J. L. R. Excitation-inhibition balance as a framework for investigating mechanisms in neuropsychiatric disorders. *Mol Psychiatry* **24**, 1248-1257 (2019). https://doi.org:10.1038/s41380-019-0426-0

Schafer, S. T. *et al.* Pathological priming causes developmental gene network heterochronicity in autistic subject-derived neurons. *Nat Neurosci* **22**, 243-255 (2019). https://doi.org:10.1038/s41593-018-0295-x

Birey, F. *et al.* Assembly of functionally integrated human forebrain spheroids. *Nature* **545**, 54-59 (2017). https://doi.org:10.1038/nature22330

Yoon, S. J. *et al.* Reliability of human cortical organoid generation. *Nat Methods* **16**, 75-78 (2019). https://doi.org:10.1038/s41592-018-0255-0

Khan, T. A. *et al.* Neuronal defects in a human cellular model of 22q11.2 deletion syndrome. *Nat Med* **26**, 1888-1898 (2020). https://doi.org:10.1038/s41591-020-1043-9

Willsey, H. R. *et al.* Parallel in vivo analysis of large-effect autism genes implicates cortical neurogenesis and estrogen in risk and resilience. *Neuron* **109**, 1409 (2021). https://doi.org:10.1016/j.neuron.2021.03.030

Roberts, T. C., Langer, R. & Wood, M. J. A. Advances in oligonucleotide drug delivery. *Nature Reviews Drug Discovery* **19**, 673-694 (2020). https://doi.org:10.1038/s41573-020-0075-7

Gegenhuber, B., Wu, M. V., Bronstein, R. & Tollkuhn, J. Gene regulation by gonadal hormone receptors underlies brain sex differences. *Nature* **606**, 153-159 (2022). https://doi.org:10.1038/s41586-022-04686-1

Sebastian, R. *et al.* Schizophrenia-associated NRXN1 deletions induce developmental-timing- and cell-type-specific vulnerabilities in human brain organoids. *Nature Communications* **14**, 3770 (2023). https://doi.org:10.1038/s41467-023-39420-6

Zeng, L. *et al.* Functional impacts of NRXN1 knockdown on neurodevelopment in stem cell models. *PLoS One* **8**, e59685 (2013). https://doi.org:10.1371/journal.pone.0059685

Birtele, M. *et al.* The autism-associated gene SYNGAP1 regulates human cortical neurogenesis. *bioRxiv*, 2022.2005.2010.491244 (2022). https://doi.org:10.1101/2022.05.10.491244

Trotter, J. H. *et al.* Astrocytic Neurexin-1 Orchestrates Functional Synapse Assembly. *BioRxiv* (2020). https://doi.org:10.1101/2020.08.21.262097

Ingusci, S., Verlengia, G., Soukupova, M., Zucchini, S. & Simonato, M. Gene Therapy Tools for Brain Diseases. *Frontiers in pharmacology* **10** (2019). https://doi.org:10.3389/fphar.2019.00724

39 Brunet de Courssou, J.-B., Durr, A., Adams, D., Corvol, J.-C. & Mariani, L.-L. Antisense
therapies in neurological diseases. *Brain* **145**, 816-831 (2022).
https://doi.org:10.1093/brain/awab423

40 Yilmaz, C. *et al.* Neurosteroids as regulators of neuroinflammation. *Frontiers in*
Neuroendocrinology **55**, 100788 (2019).
https://doi.org:https://doi.org/10.1016/j.yfrne.2019.100788

41 National Academies of Sciences, E. & Medicine. *Sex Differences in Brain Disorders:*
Emerging Transcriptomic Evidence: Proceedings of a Workshop. (The National
Academies Press, 2021).

42 Jack, A. *et al.* A neurogenetic analysis of female autism. *Brain* **144**, 1911-1926 (2021).
https://doi.org:10.1093/brain/awab064

43 Werling, D. M. The role of sex-differential biology in risk for autism spectrum disorder. *Biol*
Sex Differ **7**, 58 (2016). https://doi.org:10.1186/s13293-016-0112-8

44 Hoffman, E. J. *et al.* Estrogens Suppress a Behavioral Phenotype in Zebrafish Mutants of
the Autism Risk Gene, CNTNAP2. *Neuron* **89**, 725-733 (2016).
https://doi.org:10.1016/j.neuron.2015.12.039

45 McCarthy, M. M. Estradiol and the developing brain. *Physiol Rev* **88**, 91-124 (2008).
https://doi.org:10.1152/physrev.00010.2007

46 Woolley, C. S. Acute Effects of Estrogen on Neuronal Physiology. *Annual Review of*
Pharmacology and Toxicology **47**, 657-680 (2007).
https://doi.org:10.1146/annurev.pharmtox.47.120505.105219

47 Lin, H. C. *et al.* NGN2 induces diverse neuron types from human pluripotency. *Stem Cell*
Reports **16**, 2118-2127 (2021). https://doi.org:10.1016/j.stemcr.2021.07.006

48 Zafra, F. & Piniella, D. Proximity labeling methods for proteomic analysis of membrane
proteins. *Journal of Proteomics* **264**, 104620 (2022).
https://doi.org:https://doi.org/10.1016/j.jprot.2022.104620

49 Bell, J. Stratified medicines: towards better treatment for disease. *Lancet* **383 Suppl 1**,
S3-5 (2014). https://doi.org:10.1016/s0140-6736(14)60115-x

50 Tsimberidou, A. M. *et al.* Molecular tumour boards — current and future considerations
for precision oncology. *Nature Reviews Clinical Oncology* **20**, 843-863 (2023).
https://doi.org:10.1038/s41571-023-00824-4

51 Zhang, H., Colclough, K., Gloyn, A. L. & Pollin, T. I. Monogenic diabetes: a gateway to
precision medicine in diabetes. *The Journal of Clinical Investigation* **131** (2021).
https://doi.org:10.1172/JCI142244

52   Sullivan, P. F. & Geschwind, D. H. Defining the Genetic, Genomic, Cellular, and
Diagnostic Architectures of Psychiatric Disorders. *Cell* **177**, 162-183 (2019).
https://doi.org:10.1016/j.cell.2019.01.015

53   Gerasimavicius, L., Livesey, B. J. & Marsh, J. A. Loss-of-function, gain-of-function and
dominant-negative mutations have profoundly different effects on protein structure. *Nature*
Communications **13**, 3895 (2022). https://doi.org:10.1038/s41467-022-31686-6

54   Wang, L. *et al.* Analyses of the autism-associated neuroligin-3 R451C mutation in human
neurons reveal a gain-of-function synaptic mechanism. *Molecular psychiatry* (2022).
https://doi.org:10.1038/s41380-022-01834-x

55   Pinggera, A. *et al.* New gain-of-function mutation shows CACNA1D as recurrently mutated
gene in autism spectrum disorders and epilepsy. *Hum Mol Genet* **26**, 2923-2932 (2017).
https://doi.org:10.1093/hmg/ddx175

56   Clark, M. B. *et al.* Long-read sequencing reveals the complex splicing profile of the
psychiatric risk gene CACNA1C in human brain. *Molecular psychiatry* **25**, 37-47 (2020).

57   Brunklaus, A. *et al.* Gene variant effects across sodium channelopathies predict function
and guide precision therapy. *Brain* **145**, 4275-4286 (2022).
https://doi.org:10.1093/brain/awac006

58   Sanders, S. J. *et al.* Progress in Understanding and Treating SCN2A-Mediated Disorders.
Trends in neuroscience **41**, 442-456 (2018).
https://doi.org:https://doi.org/10.1016/j.tins.2018.03.011

59   Kim, G., Gautier, O., Tassoni-Tsuchida, E., Ma, X. R. & Gitler, A. D. ALS Genetics: Gains,
Losses, and Implications for Future Therapies. *Neuron* **108**, 822-842 (2020).
https://doi.org:10.1016/j.neuron.2020.08.022

60   Balendra, R. & Isaacs, A. M. C9orf72-mediated ALS and FTD: multiple pathways to
disease. *Nat Rev Neurol* **14**, 544-558 (2018). https://doi.org:10.1038/s41582-018-0047-2

61   Prakasam, R. *et al.* LSD1/PRMT6-targeting gene therapy to attenuate androgen receptor
toxic gain-of-function ameliorates spinobulbar muscular atrophy phenotypes in flies and
mice. *Nature Communications* **14**, 603 (2023). https://doi.org:10.1038/s41467-023-
36186-9

62   Chintalaphani, S. R., Pineda, S. S., Deveson, I. W. & Kumar, K. R. An update on the
neurological short tandem repeat expansion disorders and the emergence of long-read
sequencing diagnostics. *Acta Neuropathologica Communications* **9**, 98 (2021).
https://doi.org:10.1186/s40478-021-01201-x

63 Sloan, S. A., Andersen, J., Pasca, A. M., Birey, F. & Pasca, S. P. Generation and assembly
of human brain region-specific three-dimensional cultures. *Nat Protoc* (2018).
https://doi.org:10.1038/s41596-018-0032-7

64 Nicolas, A. *et al.* Genome-wide Analyses Identify KIF5A as a Novel ALS Gene. *Neuron*
97, 1268-1283 e1266 (2018). https://doi.org:10.1016/j.neuron.2018.02.027

65 Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21
(2013). https://doi.org:10.1093/bioinformatics/bts635

66 Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics* **30**, 923-930 (2014).
https://doi.org:10.1093/bioinformatics/btt656

67 Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139-
140 (2010). https://doi.org:10.1093/bioinformatics/btp616

68 Hoffman, G. E. *et al.* Transcriptional signatures of schizophrenia in hiPSC-derived NPCs
and neurons are concordant with post-mortem adult brains. *Nat Commun* **8**, 2225 (2017).
https://doi.org:10.1038/s41467-017-02330-5

69 Newman, A. M. *et al.* Determining cell type abundance and expression from bulk tissues
with digital cytometry. *Nature biotechnology* **37**, 773-782 (2019).
https://doi.org:10.1038/s41587-019-0114-2

70 Hoffman, G. E. & Schadt, E. E. variancePartition: interpreting drivers of variation in
complex gene expression studies. *BMC bioinformatics* **17**, 483 (2016).
https://doi.org:10.1186/s12859-016-1323-z

71 Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing
and microarray studies. *Nucleic Acids Res* **43**, e47 (2015).
https://doi.org:10.1093/nar/gkv007

72 Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Near-optimal probabilistic RNA-seq
quantification. *Nature biotechnology* **34**, 525-527 (2016). https://doi.org:10.1038/nbt.3519

73 De Marinis, I., Lo Surdo, P., Cesareni, G. & Perfetto, L. SIGNORApp: a Cytoscape 3
application to access SIGNOR data. *Bioinformatics* **38**, 1764-1766 (2022).
https://doi.org:10.1093/bioinformatics/btab865

74 Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902 e1821
(2019). https://doi.org:10.1016/j.cell.2019.05.031

75   Urresti, J. *et al.* Correction: Cortical organoids model early brain development disrupted
1204   by 16p11.2 copy number variants in autism. *Mol Psychiatry* **26**, 7581 (2021).
1205   https://doi.org:10.1038/s41380-021-01289-6
1206   76   Yang, X. *et al.* Identification and validation of genes affecting aortic lesions in mice. *J Clin*
1207   *Invest* **120**, 2414-2422 (2010). https://doi.org:10.1172/JCI42742
1208   77   Stefan, K. & Harbour, J. W. SparK: A Publication-quality NGS Visualization Tool. *bioRxiv*,
1209   845529 (2019). https://doi.org:10.1101/845529
1210