Title

Retinoic acid-dependent loss of synaptic output from bipolar cells impairs visual information processing in inherited

retinal degeneration.

Author List

Logan Ganzen[1†], Shubhash Chandra Yadav[1†], Mingxiao Wei[1], Hong Ma[1], Scott Nawy[1], Richard H Kramer[1]

1. University of California Berkeley, Department of Molecular and Cell Biology. Berkeley, CA, USA

† Authors contributed equally to this work.

Abstract

In retinitis pigmentosa (RP), rod and cone photoreceptors degenerate, depriving downstream neurons of light-sensitive

input, leading to vision impairment or blindness. Although downstream neurons survive, some undergo morphological

and physiological remodeling. Bipolar cells (BCs) link photoreceptors, which sense light, to retinal ganglion cells (RGCs),

which send information to the brain. While photoreceptor loss disrupts input synapses to BCs, whether BC output

synapses remodel is unknown. Here we report that synaptic output from BCs plummets in mouse models of RP owing to

loss of voltage-gated Ca^{2+} channels. Remodeling reduces the dynamic range and the reliability of the BC output synapse

to repeated optogenetic stimuli, eliminating RGC firing at high stimulus frequencies. However, the consequences of

functional BC remodeling can be reversed by inhibiting the retinoic acid receptor (RAR). RAR inhibitors targeted to BCs

present a new therapeutic opportunity for mitigating detrimental effects of remodeling on signals initiated by surviving

photoreceptors or vision-restoring tools.

Introduction

Rod and cone photoreceptors degenerate in retinitis pigmentosa (RP) and age-related macular degeneration (AMD),

depriving the visual system of light-evoked signals necessary for sight. The remaining retinal neurons can survive long

after the rods and cones are gone. Critically, retinal ganglion cells (RGCs), the output neurons of the retina, remain

connected to the brain [1,2]. This enables responses initiated by surviving photoreceptors to remain detectable late into

disease progression [3]. The maintained connectivity of RGCs also provides a lifeline for restoring vision in profound

blindness, either through regeneration of lost photoreceptors from stem cells, or introduction of light sensitivity in

surviving retinal neurons with optoelectronic [4], optogenetic [5,6], or opto-pharmacological tools [7,8]. Whether direct or

indirect, if a light stimulus can alter RGC firing appropriately, the response should be transmitted to the brain and result

in visual perception.

Unfortunately, downstream neurons undergo remodeling that can corrupt the proper encoding of information and

exacerbate vision loss [9]. Morphological remodeling can require months to develop in mice [10], while in contrast,

physiological remodeling begins almost immediately when photoreceptors die [11,12]. Many RGCs become hyperactive,

spontaneously firing bursts of action potentials that obscure responses initiated by surviving photoreceptors. Patch

clamp recordings show that the intrinsic electrical properties of RGCs are altered by photoreceptor degeneration [8] and

transcriptional analysis shows altered expression of voltage-gated ion channels [13]. In addition, the plasma membrane of

RGCs exhibits increased permeability to large dye molecules and azobenzene photoswitches, events attributed to

increased expression of ATP-gated P2X receptors, which have a large pore [13].

How does photoreceptor death in the outer retina lead to physiological remodeling of RGCs in the inner retina? Retinoic

acid (RA) is the key signal that triggers RGC hyperactivity and hyperpermeability [12]. Inhibitors of retinaldehyde

dehydrogenase (RALDH), the enzyme that synthesizes RA, or RAR, the nuclear retinoic acid receptor, reduce

hyperactivity and hyperpermeability. These agents also unmask responses of RGCs to dim light, sharpen responses of

visual cortical neurons to patterns of illumination, and improve behavioral visual perception in vision-impaired mice with

partial photoreceptor degeneration [9].

Bipolar cells (BCs) are located between photoreceptors and RGCs. During photoreceptor degeneration, input synapses

exhibit morphological remodeling. BC dendrites, which normally invaginate the synaptic terminals of rods and cones,

begin to retract [14], disconnecting the synapse. Synaptic transmission is also impaired by the loss or redistribution of

dendritic mGluR6 receptors [15] and TRPM1 channels [16], which normally underlie the ON-BC response to the

photoreceptor neurotransmitter glutamate. Despite these negative consequences, not all remodeling is counter-

productive. The retina shows resilience by employing mechanisms that partially compensate for photoreceptor loss.

Over time, new dendrites sprout from BCs and can re-establish synaptic connections in retinas with degenerated

photoreceptors [17]. Experimental ablation of half of the rods in adults rebalances excitatory and inhibitory synaptic

strength to BCs to restore proper RGC responses [18]. In young mice, partial ablation of cones leads to dendritic rewiring in

certain BCs to restore the original number of input synapses [19].

In contrast, the question of whether the efficacy of the BC output synapse is changed by photoreceptor degeneration

has remained unanswered. This is critical, as changes in BC output will alter the response properties of downstream

RGCs, the only cell type that provides visual input to the brain. Here we evaluate the effects of photoreceptor

degeneration on output synapses of two types of ON-BCs; rod bipolar cells (RBC), critical for scotopic vision and the

most numerous BC type in mouse, and cone bipolar cell type-6 (CBC6), which plays a crucial role in all vision, from

scotopic, to mesopic, to photopic. We find a dramatic reduction of voltage-gated Ca^{2+} current and neurotransmitter

release from both RBCs and CBC6s, weakening transmission and decreasing the frequency response of the BC to RGC

synapse. These effects can be reduced by inhibiting RA signaling, providing a potential therapeutic approach for

abrogating the detrimental effects of BC remodeling in RP and AMD.

Results

Voltage-gated calcium current in ON-BCs is reduced by photoreceptor degeneration.

Voltage-gated calcium channels are critical for evoked neurotransmitter release at the BC output synapse [20]. To assess whether photoreceptor degeneration alters activity of voltage-gated calcium channels, we measured membrane current in whole-cell voltage-clamped BCs in retinal slices. We compared voltage-gated Ca^{2+} current in wildtype (WT) mice to those in rd1 and rd10 mice, widely used animal models of RP. We focused on the Type-6 ON cone bipolar cell (CBC6), which synapses directly onto RGCs, and rod bipolar cells (RBCs), which drive RGCs indirectly through a serial synapse on the CBC output synaptic terminal (Figures 1a,d). Candidate BCs of each type were identified with cell-type selective gene promoters that drive expression of a fluorescent protein reporter fused with channelrhodopsin-2 (ChR2-eYFP), namely the CCK promoter for CBC6 cells and the PCP2 promoter for RBCs. The identity of individual BCs was confirmed by dye-filling the cell to localize their terminals to specific sublamina in the inner plexiform layer (IPL) (Figure 1g).

We found that the voltage-gated Ca^{2+} current in CBC6 cells was 3-fold larger in WT mice than in rd1 or rd10 mice at maximal activation (Figures 1b,c). Similarly, voltage-gated Ca^{2+} current in RBCs was 2-fold larger than in rd1 or rd10 mice at maximal activation (Figures 1e,f). In both BC types, steady-state current vs. voltage curves showed reduced current amplitude without any change in voltage-dependent activation (Figures 1c,f), consistent with a change in Ca^{2+} channel number rather than a change in gating properties or channel type. We found that the L-type Ca^{2+} channel inhibitor nimodipine eliminated the current in CBC6 cells (Figure S1) consistent with previous findings that L-type Ca^{2+} channels predominate in BCs [21]. Photoreceptor degeneration occurs early in postnatal development in rd1 mice and later in adulthood in rd10 mice, making certain aspects of morphological remodeling more severe in the rd1 strain [14]. Despite this, the loss of Ca^{2+} current in CBC6 cells and RBCs was nearly identical in rd1 and rd10 mice.

Figure 1. Voltage-gated calcium current of ON-BCs is reduced in photoreceptor degenerated retinas. A, B) Retinal circuit diagrams showing whole-cell patch-clamped CBC6 cells or RBCs with accompanying fluorescent dye fills (Alexa-Fluor 594) of a CBC6 cell or an RBC in a retinal slice. Scale bar = 5 µm. C, D) Inward calcium currents in BCs from WT, rd1, and rd10 retinas. Current was activated with a depolarizing voltage step (50 ms) from -60 mV to -20 mV. E, F) Steady-state current vs. voltage (I-V) curves, elicited with a series of 10 mV incrementing depolarizing steps from -60 to 0 mV (for CBC6 cells WT: N = 9; rd1: N = 5; rd10: N = 6; for RBCs, WT:N = 9; rd1:N = 6; rd10:N = 10). I-V curves from rd1 and rd10 BCs shows that the voltage-gated calcium current is lower than in WT BCs (Pairwise Wilcoxon Rank Sum Exact Test (PWT) with false discovery rate (FDR) correction (for CBC6: WT – rd1, p = 0.017; WT – rd10, p = 0.024. For RBC: WT – rd1, p= 0.006; WT – rd10, p-value = 0.001). Error bars indicate ± SEM.

Transmission at the BC output synapse is reduced by photoreceptor degeneration.

Neurotransmitter release is triggered by intracellular Ca^{2+}, hence the reduced Ca^{2+} current in BCs suggests that synaptic

transmission will also be reduced. ON-BCs are ordinarily depolarized in the light, but in the absence of photoreceptors

we can use optogenetics as an alternative to depolarize the cells and evoke synaptic release. We leveraged the cell-type

selective expression of ChR2-eYFP to optogenetically depolarize either RBCs or CBC6s, in the WT, rd1, or rd10

background. To interrogate the BC output synapse in isolation, we blocked photoreceptor inputs onto BCs with the

mGluR6 agonist L-AP4 and the kainate receptor antagonist ACET, which together eliminate light responses in all BCs [22].

We also blocked inhibitory synaptic inputs onto BCs with antagonists of ionotropic GABA and glycine receptors and

electrical synapses with the gap junction uncoupler MFA.

We stimulated CBC6 cells with light flashes while we recorded evoked excitatory postsynaptic currents (EPSCs) from ON

α-RGCs from whole-mount retinas (Figure 2a). Amongst the various BCs, this promoter enables transgene expression

specifically in CBC6 cells, although sparse expression has been noted in non-BC cell types, including some ON type RGCs

[23]. ON α-RGCs were identified by soma size and dendritic layer termination in the IPL, mapped after dye-filling. If a

ChR2-eYFP expressing RGC was inadvertently targeted, it was removed from further consideration based on the

exceedingly short optogenetic response delay and lack of susceptibility to the AMPAR antagonist DNQX (Figure S2).

Optogenetically evoked EPSCs in ON α-RGCs were smaller and slower in rd1 and rd10 than in WT retinas (Figure 2b).

EPSCs were reduced across all stimulation strengths, including stimuli that saturated the synaptic response, consistent

with diminished synaptic efficacy. EPSCs were ~40% smaller in rd1 and rd10 retinas with a saturating stimulus (Figures

2c,d). In addition, EPSC kinetics were slower in both rd1 and rd10, with a longer time-to-peak (Figure 2e). Patch clamp

recordings of CBC6 cells expressing eYFP-ChR2 showed that the light-elicited current was nearly identical in WT, rd1, and

rd10 retinas (Figure S3b), consistent with equal ChR2 expression. Furthermore, test flashes depolarized the cells equally

(Figure S3). Hence the reduced EPSCs cannot be explained by reduced presynaptic depolarization in the degenerated

retina, but rather by reduced synaptic efficacy. The output synapse also exhibited a striking loss of sensitivity in

degenerated retina, most easily observed as a rightward shift in the plot of normalized responses vs. stimulus strength

(Figure S4a). The stimulus required to evoke a half-maximal EPSC was nearly three times stronger in rd1 compared to

WT retina (Figure S4c). Thus, the CBC6 output synapse exhibited a decrease in both peak response and sensitivity.

For examining the output synapse of RBCs, we expressed ChR2-EYFP under control of the Pcp2 promoter [24]. We

optogenetically depolarized RBCs in retinal slices, recording EPSCs from their postsynaptic partner the AII amacrine cell.

AII amacrine cells are electrically coupled to CBC6, indirectly driving RGC firing (Figure 2f). We recorded EPSCs in AII cells,

which were identified after dye-filling by their distinctive cell body shape and dendritic terminations in the IPL. Synaptic

output of RBCs onto AII cells was also reduced in rd1 and rd10 retinas as compared to WT (Figure 2g). Responses to

saturating light at the RBC synapse were reduced by ~75% in rd1 retinas and ~65% in rd10 retinas as compared to WT

with a saturating stimulus (Figures 2h,i). There was also a loss of synaptic sensitivity, although less striking than the loss

at the CBC6 output synapse (Figures S4b,d). RBCs ordinarily exhibit highly synchronous release, resulting in rapid and

transient postsynaptic responses [25], but the responses were slower to rise and more prolonged in degenerated retina,

implying that degeneration promotes more asynchronous transmitter release from RBCs. The time-to-peak of evoked

EPSCs was 2- or 3-fold longer, for rd10 and rd1 retinas as compared to WT (Figure 2e,j). In summary, the output

synapses of both CBC6 cells and RBCs show dramatically reduced synaptic efficacy, consistent with reduced Ca^{2+}-

dependent neurotransmitter release.

Figure 2. Synaptic output from ON-BCs is reduced in photoreceptor degenerated retinas. A,F) Retinal circuit diagrams
showing optogenetically stimulated CBC6 cells or RBCs. EPSCs were recorded from a postsynaptic ON α-RGC or AII
amacrine cell, respectively. B,G) EPSCs in CBC6 cells (B) and RBCs (G). Note the reduced EPSCs in rd1 and rd10 retinas
compared to WT. The green dot represents the light flash. C,H) At all effective optogenetic stimulus strengths, EPSCs
from CBC6 cells (C) and RBCs (H) were smaller in rd1 and rd10 than in WT (For CBC6 cells: rd1 vs. WT: p<0.001; rd10 vs.
WT: p = 0.009; rd1:N=15, rd10:N=11, WT:N=13. For RBCs: rd1 vs. WT: p<0.001; rd10 vs. WT: p<0.001; WT:N=17,
rd1:N=10, rd10:N=14). D,I) EPSCs elicited by a saturating stimulus (dashed box in panels C and H) were smaller in rd1
and rd10 than in WT (For CBC6 cells: rd1= 1.27 nA, rd10 = 1.48 nA, WT = 2.12 nA; rd1 vs. WT: p <0.001; rd10 vs. WT: p =
0.008; for RBCs: rd1 = 152 pA, rd10 = 198 pA, WT = 546 pA; rd1 vs. WT: p <0.001; rd10 vs. WT: p = 0.007).   E,J)Rise time
of EPSCs is slower in rd1 and rd10 than in WT retinas for both the CBC6 output synapse (rd1 = 11.92 ms, rd10 = 10.99
ms, WT = 7.72 ms; rd1 vs. WT, p <0.001; rd10 vs. WT, p = 0.011) and the RBC output synapse (rd1 = 21.17 ms, rd10 =
19.19 ms, WT = 9.24 ms; rd1 vs. WT, p <0.001; rd10 vs. WT, p <0.001). For panels D, E, I and J, each data point represents
peak EPSC from an individual recording; horizontal lines represent median EPSC values; error bars indicate ± SEM. All
statistical comparisons were made with PWT with FDR correction.

Morphological analysis shows that the number of output synapses from CBC6 cells is unaltered by photoreceptor degeneration.

Reduced synaptic transmission could either be caused by a decrease in the number of synaptic contacts between pre- and postsynaptic neurons or caused by a decrease in the synaptic efficacy with no change in synapse number. To discriminate between these possibilities, we used immunohistochemistry and fluorescence microscopy to quantify the density of CBC6 synaptic contacts onto ON α-RGC dendrites in the IPL. We used antibodies against secretagogin to visualize the axon terminals of CBC6 cells. Secretagogin labels CBC types 2,3,4, 5, 6 in the mouse retina. CBC2-4 types are OFF-BCs that terminate in distant off sublamina, CBC5 stratify in IPL 3, so CBC6 terminals can be identified unambiguously in the IPL 4 [26]. CBC6 cells constitute more than 70% of bipolar synaptic inputs to ON α-RGC [27] while other ON-CBC types sharing stratification with ON α-RGCs lack secretagogin expression [28]. BC terminals possess synaptic ribbons, which we labeled with a fluorescent antibody against ribeye, the main structural protein in the ribbon. Finally, we visualized the dendritic tree of an individual ON α-RGC by dye-filling the cell (Figure 3a).

Confocal microscope images from the IPL show CBC6 terminals overlapping with ON α-RGC dendrites (Figure 3b). To restrict our analysis to functional synapses, we examined the overlap between ON α-RGC dendrites and ribeye-labeled puncta (CtBP2) within CBC6 terminals (Figures 3c-e). We found that the number of putative synapses onto the ON α-RGC was the same in WT and rd1 retinas (Figure 3f). The number of synapses per unit area of the ON α-RGC was also the same (Figure 3g). By focusing through the depth of the IPL, we reconstructed the terminal tree of an individual CBC6 cell, which is displayed as a flattened 2D projection (Figure 3h). Imaging ribeye puncta overlapping with the terminal tree revealed the total number of presumptive ribbon synapses formed by that CBC6 cell (Figure 3i,j). Using this approach, we found no difference in the number of CBC6 cell output synapses between WT and rd1 retinas (Figure 3k). Additionally, there was no difference in the density of ribbons per axon terminal (Figure 3l). We conclude that there is no difference in the number of presumptive CBC6 synapses between WT and rd1 retina.

Figure 3. CBC6 cell output synapse morphology is unaltered by photoreceptor degeneration. A) Soma and dendrites of an ON α-RGC, labeled by intracellular injection with Alexa Fluor 568. Image is a 3-D projection from a stack of confocal optical sections. B-E) Magnified view of one ON α-RGC dendrite (magenta), overlapping with CBC6 axon terminals (B), labeled with anti-secretagogin (cyan), C) overlapping with synaptic ribbons, labeled with anti-CtBP2 (yellow), D) overlapping with synaptic ribbons exclusively in CBC6 terminals, identified by co-labeling with anti-secretagogin and anti-CtBP2. E) Arrows point to sites of co-localization between the dendrite, CBC6 axon terminals, and CtBP2, i.e., presumptive ribbon synapses between CBC6 cells and the ON α-RGC. F, G) Quantification of the total number of CBC6 ribbon synapses across all optical sections of the dendritic tree of an ON α-RGC (F) and the density of CBC6 ribbon synapses per unit area of the ON α-RGC. (G). There is no significant difference in the number (WT vs. rd1: p = 1) or density of ribbon synapses (WT vs. rd1: p = 0.33) in rd1 vs. WT retinas. WT: N=5, rd1: N=6. H)2D projection of the 3D structure of axon terminals of CBC6 cells in WT and rd1 retinas. I) CBC6 axon terminal overlapping with CtBP2. J) CBC6 axon terminals with yellow dots denoting thresholded points of overlap with CtBP2. The red outline indicates the axon terminal tree of one individual CBC6 cell. K, L) Quantification of the number of CtBP2 sites per CBC6 axon terminal tree (K) and the density of CtBP2 sites per CBC6 axon terminal tree (L). There is no significant difference in the number (WT vs. rd1: p = 0.33) or density (WT vs. rd1: p = 1) of ribbon synapses made by an individual CBC cell onto an ON α-RGCs in rd1 vs. WT retinas. WT: N=8, rd1: N=8. Scale bars: A (50 μm); B-E, H-J (5 μm). All statistical comparisons were made with PWT with FDR correction.

Electrically evoked glutamate release from BC terminals is reduced by photoreceptor degeneration.

The described results suggest that decreased synaptic efficacy from BC to postsynaptic cells in the photoreceptor-degenerated retina is caused by decreased release of the neurotransmitter glutamate. To directly confirm reduced glutamate release, we used the genetically encoded glutamate indicator iGluSnFR, delivered to RGCs with an AAV vector that transduces diverse types of RGCs indiscriminately. After allowing 2-3 weeks for viral transduction and gene expression, we isolated and mounted the retina in a chamber that enables simultaneous trans-retinal electrical stimulation and 2-photon imaging (Figure 4a). By focusing on different depths through the inner retina, we obtained virtual cross-sections of the IPL, which showed similar baseline levels of iGluSnFR expression in WT and rd1 retinas (Figure 4b). We placed electrodes across the thickness of the retina to enable electrical stimulation of all cells. To examine neurotransmitter release from BCs output synapse in isolation, we pharmacologically blocked all input synapses to BCs. Under these conditions, we found that a brief electrical stimulus (10 ms) evoked a transient increase in iGluSnFR fluorescence in the ON-layer of the IPL that was smaller in rd1 than in WT (Figure 4c). The peak change in fluorescence over background ($\Delta F/F_0$) was 75% smaller in rd1 than in WT (Figures 4d,e). We were concerned that retinas lacking photoreceptors might have altered tissue resistance, which could impact the effectiveness of electrical stimulation of BCs. To address this point, we varied electrical stimulus strength to generate a range of responses, from no detectable iGluSnFR signal to a saturating signal (Figure 5f). We found that while the magnitude of the response to all stimuli (including the saturating stimulus) was lower in rd1, the sensitivity was the same, indicating that the reduced response could not be attributed to less effective electrical stimulation (Figures 4f). Furthermore, peak iGluSnFR $\Delta F/F$ in response to application of 10 mM glutamate was no different in WT and rd1 retinas, ruling out the possibility of impaired iGluSnFR expression in rd1 retina (Figure 4g). Taken together, these results suggest that reduced synaptic efficacy in degenerated retina is caused by reduced evoked glutamate release from ON-BCs.

Figure 4. Synaptic release of glutamate from ON-BCs is reduced in retinas with photoreceptor degeneration. A)
Apparatus for simultaneous trans-retinal electrical stimulation and 2-photon imaging of glutamate, the BC
neurotransmitter. B) Virtual cross-sections (z-stacks) of retinas imaged with 2-photon microscopy. The genetically
encoded fluorescent glutamate indicator iGluSnFR (cyan) was virally expressed throughout the IPL of WT) and rd1
retinas. The dashed line represents the ON Layer imaging plane, 10 μm into the IPL from the GCL border. Cell body
layers were counterstained with a DNA-binding dye (NUCLEAR-ID, red). C) An optical section within the ON layer of the
IPL before and 15 ms after stimulation, consisting of a 50 V, 2.5 ms shock in WT and rd1. D) The change in iGluSnFr
fluorescence over background (ΔF/F), proportional to the change in glutamate concentration. Note that the
fluorescence change upon stimulation is smaller in rd1 than in WT. E) Time course of the fluorescence change elicited by
a single 2.5 ms, 50 V shock in a WT and rd1 retina. F) Peak iGluSnFR ΔF/F in response to varying shock strength. G) Peak
iGluSnFR ΔF/F in response to a saturating 50 V, 2.5 ms shock is lower in rd1 than in WT ($p = 0.02$, N = 7 retinas). Peak
iGluSnFR ΔF/F in response to application of 10 mM glutamate is no different between WT and rd1 retinas ($p = 0.71$, N = 7
retinas). Barplots indicate median value; error bars indicate ± SEM. Scale bar: 5 μm. All statistical comparisons were
made with PWT with FDR correction.

Retinoic acid mediates physiological remodeling of BCs

RA triggers physiological remodeling of RGCs in the degenerated retina. To explore whether RA also underlies physiological remodeling of BCs we treated mice with BMS 493, an inverse agonist of the RA receptor, RAR [29]. Because the effects of RA are mediated by changes in gene expression, which may require days to take effect, we injected BMS 493 into the vitreous 5-7 days before retinal recording. First, we measured voltage-gated Ca^{2+} current from BCs in retinal slices. BMS 493 increased the current by ~3-fold in both CBC6 cells from rd1 mice and in RBCs from rd10 mice at maximal activation while causing no change in voltage-dependent activation, as shown in steady-state I-V curves (Figure 5a,b). In contrast, BMS 493 caused no change in Ca^{2+} current in CBC6 cells from WT mice (Figures S5). Next, we examined the effect of inhibiting RA signaling on the synaptic output of BCs in degenerated retinas. BMS 493 increased optogenetically-elicited synaptic transmission from CBC6 cells by ~35% in rd1 retina and ~45% in rd10 retina with a saturating stimulus (Figure 5c), nearly completely reversing the deficit caused by photoreceptor degeneration (see Figure 2c). BMS 493 also increased synaptic transmission from RBCs, by 30% in rd1 retina and 50% in rd10 with a saturating stimulus (Figure 5d), reducing the photoreceptor degeneration deficit (see Figure 2c).

Inhibiting RA signaling also had dramatic effects on short-term synaptic plasticity, which is altered in photoreceptor degenerated retina. In WT retinas, the BC output synapse exhibits profound paired-pulse depression, consistent with the large Ca^{2+} current eliciting so much neurotransmitter release that the readily-releasable pool of synaptic vesicles is depleted by the first stimulus, leaving little to be released by the second stimulus (Figure 5e,f) [30,31]. In rd1 or rd10 retinas, there is little paired-pulse depression, consistent with smaller Ca^{2+} current evoking less release, preserving the readily releasable vesicle pool. BMS 493 caused more synaptic depression in and vesicle depletion in rd1 retinas similar to WT. Injecting all-trans RA (ATRA), a relatively stable isomer of RA, into WT eyes reduced the amplitude of optogenetically-evoked EPSCs and reduced paired-pulse depression, suggesting that RA signaling is sufficient to trigger changes in the BC output synapse. Taken together, these findings implicate RA as the key signal that induces physiological modeling of BCs.

Figure 5. RA signaling is necessary and sufficient for physiological remodeling of BCs. A) Inward calcium currents in BCs treated with BMS 493 from rd1 CBC6 or rd10 RBCs. Current was activated with a depolarizing voltage step (50 ms) from -60 mV to -20 mV. B) Steady-state current vs. voltage (I-V) curves, elicited with a series of 10 mV incrementing depolarizing steps from -60 mV. I-V curves show that the BMS 493 significantly rescues voltage-gated calcium current in both CBC6 cells in rd1 retina (p <0.001) and RBCs in the rd10 retina (p <0.001). CBC6 cells: rd1, N = 5; rd1 + BMS 493, N = 5. RBCs: rd10, N = 10, rd10 + BMS 493, N = 7. C) EPSC magnitude vs. optogenetic stimulus strength. BMS 493 increased EPSC magnitude in ON α-RGCs in rd1 (p=0.006) and rd10 retinas (p<0.001). RGCs: rd1, N = 15; rd1 + BMS 493, N = 11; rd10 N = 11; rd10 + BMS 493, N = 8. D) BMS 493 also increased EPSC magnitude in AII amacrine cells in rd1 (p=0.001) and rd10 retinas (p<0.001). AII cells, rd1: N=10, rd1 + BMS 493: N=9, rd10: N=14, rd10 + BMS 493: N=12. E) EPSCs from ON α-RGCs in response to paired pulse flashes given with a 50 ms interval. Green dots indicate light flash. WT shows complete paired-pulse depression and rd1 shows no paired-pulse depression. Note that BMS 493 induces paired-pulse depression in rd1 and ATRA eliminates paired-pulse depression in WT. F) Quantification of the paired pulse ratio (EPSC 2 / EPSC 1). Compared to WT, paired EPSCs exhibited less synaptic depression in rd1 (p<0.001), rd10 (p=0.004), and ATRA-treated WT retinas (p<0.001) denoted with ***. BMS 493 treatment increased synaptic depression in rd1 retinas (p<0.001) denoted with ###. WT: N=16, rd1: N=13, rd10: N=16, rd1 + BMS 493: N=11, WT + ATRA: N=5. Horizontal lines indicate median values: WT=0.02, rd1=1.38, rd10=0.43, rd1 + BMS 493=0.10, WT + ATRA=0.30 . Error bars indicate ± SEM. All statistical comparisons were made with PWT with FDR correction.

Physiological remodeling of BCs alters spatiotemporal information processing.

In the WT retina, sensory information from an array of photoreceptors signals through a series of retinal neurons, ultimately converging onto an RGC to produce a classical receptive field. In a photoreceptor-degenerated retina with the responses of a particular type of BC under optogenetic control, each RGC receives a projective field, reflecting the convergence of information from that specific BC cell type. Here we compare rd1 and WT retinas to ask whether photoreceptor degeneration alters the CBC6 to ON α-RGC projective field. We optogenetically stimulated small clusters of CBC6 cells with 50 μm spots of light to yield a spatial map of BCs projecting onto an individual ON α-RGC (Figures 6a-c). We found that the spatial map in rd1 retinas is nearly identical to the map in WT retina, but scaled down linearly (Figures 6d,e), presumably owing to decreased synaptic transmission. This finding is consistent with previous work showing little change in cone-mediated RGC receptive field size following rod degeneration, in a different mouse model of RP [32].

We next compared the temporal properties of the CBC6 to ON α-RGC network in WT and photoreceptor degenerated retinas. We optogenetically stimulated populations of CBC6 cells at different rates and recorded either EPSCs in whole-cell patch recordings or spiking in cell-attached patch recordings from individual RGCs (Figure 7a). In the WT retina, a low frequency train of stimuli (0.25 Hz) evoked a non-decrementing train of EPSCs (Figures 7b,c). Even with higher frequency stimulation (up to 7.5 Hz), EPSCs followed reliably, although their amplitudes declined during the train (Figures 7b-e). In rd1 and rd10 retinas, EPSCs were non-decrementing at low frequency (0.25 Hz), but their amplitudes declined rapidly at higher frequencies, with EPSCs failing completely by the end of the 7.5 Hz stimulation train. Comparing across all stimulation rates, the frequency at which EPSCs declined by 90% (-10 dB) at steady state was 5.25 Hz for WT , 0.68 Hz for rd1, and 2.47 Hz for rd10. (Figure 7f).

Decrementing synaptic efficacy would be expected to adversely impact action potential reliability in RGCs. To test this, we made cell-attached patch-clamp recordings to non-invasively monitor spike activity from ON α-RGCs. In the WT retina, RGC firing followed optogenetic BC stimulation reliably up to 7.5 Hz (Figure 7g), consistent with temporal tuning of normal light responses in ON α-RGCs in WT retinas [22]. In the rd1 and rd10 retina, firing failed to follow even at 2.5 Hz stimulation frequency. We compiled firing reliability data over a wide range of stimulus frequencies and train durations. RGCs failed to follow BC stimulation at a low frequency in rd10 and an even lower frequency in rd1 (Figure 7h). The decline of reliability in the CBC6 to ON α-RGC network can be attributed entirely to changes in the BCs, as there is no difference in the response of ON α-RGC to firing induced by direct current injection in WT and degenerated retinas (Figure S6). These findings illustrate the severe deficit that BC physiological remodeling imposes on information transfer to RGCs in the degenerated retina.

Equivalent experiments on RBC output synapse showed an even more extreme frequency-dependent decline in photoreceptor degenerated retina (Figures 8a,b). Even at 0.25 Hz, EPSCs from rd1 and rd10 retinas began decrementing while WT did not (Figure 8b,c). Higher stimulation frequency led to the near complete loss of EPSCS in rd1 and rd10 at

steady state (Figure 8b,d,e). Comparing all stimulation rates, the frequency at which EPSCs declined by 90% (-10 dB) at steady state was 5.14 Hz for WT, 0.69 Hz for rd1, and 1.34 Hz for rd10. (Figure 8f). Since the vast majority of ON BC types receive direct or indirect input from RBCs, the decrementing responses in photoreceptor degenerated retinas is expected to adversely impact a wide variety of ON-RGCs beyond just the ON α-RGC subtype.

Figure 6. The spatial map of the BC to RGC network is preserved in photoreceptor degenerated retina but scaled down linearly. A) Schematic diagram of the experiment. To map the projective field corresponding to many CBC6 cells converging onto a single ON α-RGC, we used 50 μm non-overlapping spots of light to optogenetically stimulate clusters of BCs, covering the entire dendritic tree of the ON α-RGCs. B-C) Example spatial projective field map from a WT (B) and rd1 (C) retina. Detectable EPSCs were elicited over ~200 μm^2 of the 650 μm^2 mapped area. D) Group data from ON α-RGCs (N=9 each), with the projective field represented as mean EPSC vs. distance from the cell soma. Data from all directions from the cell center were averaged and are shown duplicated for clarity. Note that EPSCs were smaller in rd1 than WT (p=0.012), but there was no difference in spatial weighting of the linearly scaled projection field (p=0.47). Variability ribbons indicate 95% confidence intervals. All statistical comparisons were made with PWT with FDR correction.

Figure 7. Photoreceptor degeneration reduces the frequency response in the cone pathway. A) CBC6 cells were optogenetically stimulated at different rates while EPSCs were recorded in an ON α-RGC. B) EPSCs elicited by stimuli repeated at 0.25 Hz, 2.5 Hz, and 7.5 Hz. Green dots indicate the time of light flash. C − E) EPSCs in rd1 and rd10 retinas diminish faster with increasing stimulation frequency. F) EPSC decline expressed in dB ($10Log\frac{EPSC_{Final}}{EPSC_{Initial}}$) for each stimulation frequency. The continuous line is the fit with a linear function. (WT: N=7, $y = -0.143 - 1.93x$, R^2= 1.00; rd1: N = 9, $y = 1.42 - 12.5x$, R^2 = 0.99,; rd10: N=10 $y = 0.272 - 3.93x$, R^2= 0.95). The dotted line is $y = -10\ dB$. EPSCs for rd1 at 7.5 and 10 Hz become immeasurably small at steady state. G) Raster of cell-attached spike recordings from stimulation at 0.25 Hz, 2.5 Hz, and 7.5 Hz. Green dots indicate light flash. H) The ratio of stimuli that produced an action potential to the total number of stimuli per frequency. Compared to WT, rd1 and rd10 ON α-RGCs successfully encoded fewer stimuli as stimulation frequency increased. WT: N=6, rd1, N=6, rd10, N = 5 cells.(rd1: p<0.001, rd10: p=0.034. PWT with FDR correction).

Figure 8. Photoreceptor degeneration reduces the frequency response in the rod pathway. A) RBCs cells were

optogenetically stimulated at different rates while EPSCs were recorded in an AII amacrine cell. B) EPSCs elicited by

stimuli repeated at 0.25 Hz, 2.5 Hz, and 5 Hz. Green dots indicate the time of light flash. C – E) EPSCs in rd1 and rd10

retinas diminish faster with increasing stimulation frequency. F) EPSC decline expressed in dB ($10Log \frac{EPSC_{Final}}{EPSC_{Initial}}$) for each

stimulation frequency. The continuous line is the fit with a linear function. (WT: N=6, $y = -0.28 - 1.89x$, R^2= 0.90;

rd1: N=5, $y = 2.63 - 18.4x$, R^2= 0.99,; rd10: N=11, $y = 0.579 - 7.91x$, R^2= 0.94). The dotted line is $y = -10 \, dB$.

EPSCs for rd1 at 5 Hz become immeasurably small at steady state.

Discussion

This paper reports the surprising finding that photoreceptor degeneration leads to dramatic functional weakening of the output synapses of ON-BCs. RBCs, the most numerous type of BC in the rod-dominant mouse retina, and CBC6 cells, which receive information from both rod and cone pathways, are both affected. The net effect of these changes in BCs is to reduce the amplitude and narrow the frequency response of light onset signals sent to RGCs and onward to the brain. Our imaging results with iGluSnFr suggests that glutamate release from OFF-BCs might also be reduced in the degenerated retina, which might also reduce signals at light offset. New cell-type selective promoters for particular OFF-BCs could help address this point.

Previous experiments on a rat model of RP showed that ON-RGCs exhibit reduced spontaneous firing, paralleling the loss of the RGC light response [33]. The most obvious explanation for the decline in spontaneous and evoked ON-RGC activity is that the diminishing number of photoreceptors drives a declining amount of excitatory synaptic input. However, the possibility that changes intrinsic to ON-BCs also contributed to changes in ON-RGC activity remained unexplored. Our discovery that the output synapse of ON-BCs is dramatically weakened adds an important new mechanism that may contribute to changes in ON-RGC activity across species, not only corrupting retinal signal processing, but also contributing to the pathophysiology of vision loss.

Another consequence of photoreceptor loss is the emergence of membrane potential oscillations in the inner retina, ultimately resulting in spontaneous burst firing in RGCs [34]. The oscillation is initiated in the electrically coupled network of AII amacrine cells [35], but it remains to be determined if RA-mediated events are involved.

Mechanism of physiological remodeling

Our findings suggest that RA is necessary and sufficient for physiological remodeling of the ON-BC output synapse. Adding exogenous ATRA mimics changes in synaptic release that are characteristic of degenerated retinas (Fig. 4e). Inhibiting RAR with BMS 493 reverses changes in synaptic release and restores voltage-gated Ca^{2+} current to near WT levels (Fig. 4). Our previous studies utilizing an RA-induced reporter demonstrated elevated RA-induced gene expression in degenerated retina [12].

It remains unclear where the excessive RA is produced. RA has been implicated in physiological remodeling not only in the output synapses of BCs but also in RGCs [12]. Moreover, RA is critical for morphological remodeling of BC dendrites, at their input synapses [17]. The fact that multiple cell types are affected in a degenerated retina suggests that RA might act in a paracrine fashion, produced by one cell type (perhaps RPE cells), and diffusing throughout the retina to trigger remodeling in many cell types. Immunocytochemistry and enzyme assays show that the retina, choroid, and RPE all express isoforms of RALDH, the enzyme that synthesizes RA [36]. In the RPE, RALDH normally converts all-trans retinaldehyde released from photobleached rhodopsin into RA [37]. The loss of photoreceptors might increase the concentration of retinaldehyde, leading to increased production of RA by the RPE. In this scenario, RA would be an

intercellular signal, diffusing from the outer to the inner retina to exert its actions. Alternatively, it is possible that RA is

synthesized within particular retinal neurons and serves as an intracellular messenger. In neurons in the hippocampus,

local production of RA in dendrites regulates local gene translation, a key event underlying homeostatic synaptic

plasticity [38] .

How does elevated RA decrease voltage-gated Ca^{2+} current and synaptic transmission? The size of the Ca^{2+} current in

ON-BCs is reduced by photoreceptor degeneration, but the activation and inactivation properties are unchanged,

suggesting a decreased level of channel protein rather than a change in channel type. RAR is an enhancer of gene

transcription, which often results in increased expression of specific proteins. However, there are many examples of RA-

induced down-regulation of proteins, mediated by a sign-inverting gene regulatory cascade [39].

L-type Ca^{2+} channels are complexed with accessory proteins, including alpha-2/delta subunits that determine Ca^{2+}

current density and activation/inactivation kinetics [40], in addition to controlling vesicle release probability at synapses [41].

It is possible that functional down-regulation of Ca^{2+} channels is caused by misregulation of alpha-2/delta, as occurs in

sensory neurons in dorsal root ganglia after nerve injury [42]. Many other RA-induced biochemical events could also result

in decreased Ca^{2+} current, including altered post-translational modification or altered membrane trafficking of Ca^{2+}

channel proteins [42].

Is the loss of Ca^{2+} channels sufficient to account for the decline in synaptic release, without invoking additional changes

in the release machinery? We find no decline in the number of synaptic contacts between BCs and RGCs and no loss of

synaptic ribbons, a key component of the BC synaptic machinery. Hence, at least at this age (p60) in rd1 mice,

photoreceptor degeneration causes no obvious structural changes in the BC output synapse. We did observe one

interesting phenomenon concerning the frequency-dependence of the ON-BC output synapse in the degenerated retina,

but it also may be explained by decreased Ca^{2+} influx during stimulation. We found that paired-pulse depression was

dramatically decreased in rd1 and rd10 retinas (Fig. 4f), whereas there was a great increase in failure of the synapse with

more sustained, high-frequency stimulation (Fig. 7h). Intracellular Ca^{2+} regulates synaptic vesicle fusion to the plasma

membrane, but it is also thought to regulate other steps in the synaptic vesicle cycle, such as vesicle docking and priming

[43]. In fact, there is evidence for multiple Ca^{2+}-dependent steps in the vesicle cycle at BC ribbon synapses [31]. If Ca^{2+} influx

were sufficient to cause vesicle fusion but insufficient to accelerate vesicle resupply, repetitive stimulation might deplete

the pool of readily releasable vesicles before they could be replenished. Different isoforms of synaptotagmin mediate

fusion and resupply [44], providing opportunities for understanding how reduced Ca^{2+} influx in remodeled BCs affects

synaptic transmission at the molecular level.

Functional impact

Our results imply that physiological remodeling of ON-BCs will constrain information processing further along in the

visual system, limiting visual perception. The deficit in temporal coding caused by decreased ON-BC output was

particularly severe, which is expected to have a particularly large impact on perception of object motion. Humans with

RP often fail to detect moving objects or report motion in the opposite direction of actual movement [45], even if they

retain significant visual acuity for detecting static objects [46]. RP is a progressive disease that causes gradual vision loss,

but the distinct contributions of decreased photoreceptor density vs. physiological remodeling to vision decline is

unclear. Our previous results [9] showed that treatments that reverse physiological remodeling could substantially rescue

visual contrast sensitivity in rd10 mice at 2-3 months of age, when photoreceptor degeneration is already severe. We

now know that physiological remodeling applies not only to RGCs, but also to BCs, but whether remodeling of one cell

type or the other predominates in altering visual perception is also unknown.

In certain respects, the physiological remodeling that we characterized in ON-BCs might be considered a homeostatic

mechanism, analogous to other compensatory events caused by partial photoreceptor loss (Lee et al., 2022). In the

healthy retina, photoreceptors tonically release glutamate, which keeps ON-BCs hyperpolarized. Hypothetically, by

eliminating glutamate release, photoreceptor death should depolarize the ON-BCs, chronically activating their voltage-

gated Ca^{2+} channels, and increasing their tonic release of glutamate onto RGCs. If this chronic synaptic excitation of RGCs

were sufficient to trigger firing, it could impair encoding of synaptic information from other BCs that had yet to undergo

remodeling. One way that ON-BCs could compensate for excessive depolarization is by down-regulating their Ca^{2+}

channels. While the loss of Ca^{2+} channels might be adaptive, by preventing excessive output from ON-BCs, it comes at a

cost. We have demonstrated that physiological remodeling limits the dynamic range of the synaptic output of CBC6 cells

(Fig. 2c) and RBCs (Fig. 3c) and severely limits temporal reliability (Fig. 7), maladaptive for encoding visual information.

The degree of photoreceptor loss may determine whether physiological remodeling is adaptive or maladaptive.

Our findings that physiological remodeling in ON-BCs, as well as RGCs, can be reversed by inhibiting RA has important

implications for reviving sight in low-vision patients with RP and perhaps other photoreceptor degenerative disorders.

They also have important implications for vision restoration in advanced RP patients with nearly complete

photoreceptor loss and little remaining light perception. One treatment approach for these patients is to install light

responses in surviving neurons of the outer retina, including ON-BCs, either through optogenetic stimulation [6,47] or

electrical stimulation with sub-retinal implants [48]. Physiological remodeling of the ON-BC output synapse is predicted to

seriously limit the effectiveness of direct or indirect stimulation of BCs on downstream information processing and visual

perception. Our results indicate that reduced frequency-response of the ON-BC output synapse, seen as a failure of

EPSCs, severely impairs RGC spike-rate coding. In the clinical context, this may severely limit the effectiveness of

optogenetic therapies targeted to ON-BCs or upstream remnant photoreceptors in degenerated retina. The same

problems may also apply to signals initiated by stem-cell derived regenerated photoreceptors inserted into the outer

retina. Inhibiting RA synthesis or signaling may remove these problems and augment the effectiveness of any vision

restoration technology that delivers light-elicited signals to the blind retina. RA inhibition can be accomplished with

pharmacological inhibitors of RALDH or RAR [9], or with targeted AAV vectors that deliver genetic-based inhibitors of

RALDH and RAR to specific retinal cell types, for example ON-BCs or RGCs [5,47].

Methods

Animals

Mice were handled in accordance with protocols approved by the UC Berkeley Institutional Animal Care and Use Committee (AUP-2016-04-8700-1) and conformed to the NIH Guide for the Care and Use of Laboratory Animals. C3H/HeJ mice (Jackson 000659) were used as the rd1 model (*Pde6b^{rd1}*). B6.CXB1-Pde6b^{rd10}/J mice were used as the rd10 model (Jackson 004297). CCK-ires-Cre knock-in mice (Jackson 012706, abbreviated CCK-Cre) were used to drive Cre recombinase under the cholecystokinin promoter to label CBC6 cells. BAC-Pcp2-IRES-Cre mice (Jackson 010536, abbreviated Pcp2-Cre) were used to drive Cre recombinase under the Pcp2 promoter to label RBCs. Ai32 mice (Jackson 024109, B6.Cg-*Gt(ROSA)26Sor^{tm32(CAG-COP4*H134R/EYFP)Hze}*/J) were used to drive channelrhodopsin-2/eYFP fusion protein in Cre-expressing CBC6 or RBC cells. Both rd1 and rd10 mice strains were homozygous for their respective mutations in the *Pde6b* locus. The CCK-Cre, Pcp2-Cre, and Ai32 alleles were always used in mice in a hemizygous state. Mice of both sexes were used interchangeably. For experiments with WT and rd1 mice, animals were used within 3 days of postnatal day 60 (p60). For experiments with rd10 mice, animals were used within 3 days of p90.

Retinal Dissection

Mice were euthanized via isoflurane exposure and internal decapitation. Retinal dissection was performed with enucleated eyes in oxygenated ACSF (in mM: 119 NaCl, 26.2 $NaHCO_3$, 11 Dextrose, 2.5 KCl mM, 1 K_2HPO_4, 1.3 $MgCl_2$* $6H_2O$, 2.5 $CaCl_2$, 293 mOsm/Kg), (5% CO_2/95% O_2) in room light conditions with a dissection microscope. For retinal whole-mount experiments, retinas were flattened with 4 relieving cuts. For retinal slice experiments, flattened retinas were mounted on 13 mm diameter 0.45 µm filter paper discs (MF-Millipore) and sliced to a thickness of 250 µm with a Stoelting Tissue Slicer. Retinal slices were rotated 90 degrees and mounted within a vacuum grease well.

Electrophysiology

For RGC recordings, retinas were isolated and incubated in ACSF containing hyaluronidase (9,800 U/mL) and collagenase (2,500 U/mL) for 10 minutes to facilitate penetrating of the inner limiting membrane allowing access to RGC somas for patch clamp recording [49]. Retinas were mounted ganglion cell side up in a recording chamber and held in place with a harp (Warner Instruments), superfused with ACSF oxygenated with 5% CO_2/95% O_2 at a rate of 5 ml/minute at 34 degrees C and viewed under DIC optics with an upright microscope (Olympus). Reagents for ACSF were purchased from Fisher Scientific.

For retinal slice experiments, slices were made as described and perfused in the same way as whole-mount retinas. Electrodes were made with boroscilicate capillary glass tubing with OD = 1.5 mm, ID = 1.17 mm (Warner G150TF-4). For RGC recording, glass was pulled to a resistance of 5-6 MΩs using a Narishige pipette puller. For BCs and amacrine cell recording, electrodes were made with a resistance of 7-8 MΩs with a Sutter Instruments P1000. For voltage clamp

experiments, the pipette solution contained (in mM): 123 Cs gluconate, 8 NaCl, 1 CaCl$_2$, 10 EGTA, 10 HEPES, 10 glucose, 5 Mg^{2+} ATP, 5 QX 314 (Tocris Bioscience), 0.01 Alexa 594, pH 7.4 with CsOH, 293 mOsm/Kg. For current clamp experiments in BCs to measure optogenetic responses, the pipette solution contained (in mM): 125 K$^+$ Gluconate, 10 EGTA, 10 KCl, 10 HEPES, 4 Mg^{2+} ATP, 0.01 Alexa 594, 293 mOsm/Kg. Alexa Fluor 594 Hydrazide (ThermoFisher Scientific) at 10 µM was used to dye fill all cells during recording for confirmation of cell identity. Whole cell recordings were sampled at 20 kHz and filtered at 2 kHz with a Multiclamp 700B amplifier, digitized with a 1440a or 1550a Digidata A-D converter and analyzed offline with Axograph X or Clampfit. Recordings with series resistance above 10% of the cell membrane resistance or reached above 30 mΩ were discarded.

To block photoreceptor responses to light, the kainate receptor antagonist ACET (1 µM) and the mGluR6 agonist L-AP4 (10 µM) were added to ACSF. Gabazine (10 µM) and strychnine (10 µM) were also added to block ionotropic GABA and glycine receptors. MFA (100 µM) was used to inhibit gap junctions. Pharmacological agents were purchased from Tocris Biosciences.

Optogenetic stimulation of CBC6s and RBCs was provided by a 470 nm LED delivering 2.1X10^8 photons/s, (Lumencore or CoolLED pE-4000) measured at the plane of the retina with a spectrophotometer (Thor Labs). For figures 2 and 3, stimulation intensity was varied by changing the duration of the stimulus from 0.1 to 10 ms. Thus, the number of photons delivered during the optogenetic stimulus ranged from 2.1X10^7 (100%) to 2.1X10^5 (1%).

Mapping ON α-RGC Virtual Spatial Receptive Rield

Spots of light with a 50 µm diameter were generated with a Texas Instruments Digital Light Projector (CEL1015 Light Engine, Digital Light innovations) controlled with CELconductor Control Software. Briefly, an ON α-RGC was identified and patch clamped as described above. The cell body was focused on with an Olympus LUMPlanFl 20x/0.5 W ∞/0 objective and centered within a 650 µm x 650 µm field of view. Each spot projection and simultaneous EPSC recording was controlled via Digidata.

Trans-Retinal Stimulation

To achieve electrical trans-retinal stimulation, a custom chamber was created. A 65 mm petri dish was shaved down in height and embedded within a 100 mm petri dish with RTV sealant. The 65 mm dish was elevated ~ 3 mm from the bottom of the 100 mm dish. An approximately 1 mm hole was melted in the center of the 65 mm dish with a 21 gauge needle. To mount a retina, a ~ 7.5 mm x 7.5 mm piece of ion-permeant dialysis membrane was placed over the 1 mm hole to supply an even surface to place a flat-mount retina, RGC side up. A ~ 7.5 mm x 7.5 mm piece of flexible cellophane with an ~ 1mm hole punched into the center was placed over the retina as a coverslip. The retina is then sealed between the membrane and coverslip with vacuum grease so that the only electrical path between the two dishes is through the retina. Two copper wires were placed in each dish below and above the retina which were

505 connected to an ISO-Flex stimulus isolation unit. Electrical stimulation was achieved by setting the desired voltage on
506 the ISO-Flex and triggering the stimulus via Digidata.

2-Photon iGluSnFR Imaging

508 2-Photon fluorescence imaging was performed on a Sutter Instruments moveable objective microscope with an Olympus
509 XLUMPlanFl 20x/0.95 W ∞/0 objective. iGluSnFR excitation was achieved with a Coherent Chameleon Ultra II tuned to
510 910 nm. Imaging was performed with ScanImage software. Scanning field of view was 25 μm x 25 μm. Images collected
511 in Fig 4c-d were scanned at 256 pixels per line x 256 lines (frame rate = 1.48 Hz, 2 ms per line). Higher time resolution of
512 iGluSnFR flouresence for Fig 4e-g was performed at 256 pixels per line x 6 lines (frame rate = 63.13 Hz, 2 ms per line).

Immunohistochemistry

ON α-RGCs were injected with Alexa Fluor 568 hydrazide (A10441, Thermo Fisher) as described previously [50]. In short,
somas of RGCs in the whole-mount retina were visualized by acridine orange labeling [51]. Potential ON α-RGCs were
identified based on their large soma size (>20 μm) and targeted with sharp microelectrodes containing Alexa Fluor 568.
The dye injected retinas were fixed with 4% PFA and the following immunolabeling was carried out as described before
[52] . In brief, the whole-mount retinas were incubated into primary and secondary antibodies for 3 and 1 days,
respectively. CtBP2 (mouse, AB_399431, BD Biosciences) monoclonal and secretagogin (sheep, AB_2034062, BioVendor)
polyclonal antibodies were used at a dilution of 1:500 and 1:750, respectively. All immunostainings were performed at
room temperature. Images were scanned with 40x/1.4 oil objective using Zeiss LSM780 Laser Scanning Confocal
Microscope. The z-stacks were acquired at a constant pixel size of 60nm*60nm and a z-step of 0.3μm unless stated
otherwise. All raw images thus obtained were deconvolved with Huygens Essential software (Scientific Volume Imaging,
Netherlands). The deconvolution was achieved by using a theoretical point spread function. The resulting images were
further processed and analyzed in Fiji (https://fiji.sc/) [53]. The z-stacks were normalized using stack histograms prior to
analysis. The colocalization analysis was carried out within each optical section unless stated otherwise. Brightness and
contrast of the images were adjusted post-analysis in Fiji.

Colocalization between channels was performed with Fiji utilizing the colocalization highlighter plugin as previously
described [54]. Briefly, deconvoluted z-stacks were segmented using stack histogram. Secretagogin and CtBP2 signals were
thresholded with Otsu's method, and dye injected RGCs were manually thresholded. With colocalization highlighter, an
intensity ratio of 50% was applied to estimate colocalization between Secretagogin and CtBP2. Colocalization of
Secretagogin and CtBP2 with RGCs was performed sequentially with an intensity ratio of 10%. Resultant 8-bit images of
puncta were generated where at least 5 or more connected voxels were considered as colocalization. These images
were overlayed with either RGCs or CBC6 axon terminals, and the 3D Objects Counter plugin was used to quantify
putative synaptic contacts. To isolate individual CBC6 axon terminals, axonal endings were traced back from a common
point in the z-stack 8 μm into the IPL.

AAV Delivery of iGluSnFR

Plasmid containing iGluSnFR(A184S) under the control of the human synapsin promoter was purchased from Addgene

(#106174) and packaged in the AAV8-Y733F serotype [55]. Viral constructs were generated in the Gene Delivery module of

the Vision Science Core at UC Berkeley. One month prior to imaging, 1.5 µl of virus (1×10^{15} vg/mL) was injected into

intravitreally into the eye of rd1 or WT mice.

Data Visualization and Statistics

All reported statistics and diagrams were generated with R 4.2.2. Retinal and experimental diagrams were created with

Biorender.com.

References

1.   Mazzoni, F., Novelli, E., and Strettoi, E. (2008). Retinal ganglion cells survive and maintain normal dendritic

morphology in a mouse model of inherited photoreceptor degeneration. The Journal of neuroscience : the

official journal of the Society for Neuroscience *28*, 14282-14292. 10.1523/JNEUROSCI.4968-08.2008.

2.   Medeiros, N.E., and Curcio, C.A. (2001). Preservation of ganglion cell layer neurons in age-related macular

degeneration. Investigative ophthalmology & visual science *42*, 795-803.

3.   Ellis, E.M., Paniagua, A.E., Scalabrino, M.L., Thapa, M., Rathinavelu, J., Jiao, Y., Williams, D.S., Field, G.D., Fain,

G.L., and Sampath, A.P. (2023). Cones and cone pathways remain functional in advanced retinal degeneration.

Current biology : CB *33*, 1513-1522.e1514. 10.1016/j.cub.2023.03.007.

4.   Humayun, M.S., Dorn, J.D., da Cruz, L., Dagnelie, G., Sahel, J.A., Stanga, P.E., Cideciyan, A.V., Duncan, J.L., Eliott,

D., Filley, E., et al. (2012). Interim results from the international trial of Second Sight's visual prosthesis.

Ophthalmology *119*, 779-788. 10.1016/j.ophtha.2011.09.028.

5.   Bi, A., Cui, J., Ma, Y.P., Olshevskaya, E., Pu, M., Dizhoor, A.M., and Pan, Z.H. (2006). Ectopic expression of a

microbial-type rhodopsin restores visual responses in mice with photoreceptor degeneration. Neuron *50*, 23-33.

10.1016/j.neuron.2006.02.026.

6.   Sahel, J.A., Boulanger-Scemama, E., Pagot, C., Arleo, A., Galluppi, F., Martel, J.N., Esposti, S.D., Delaux, A., de

Saint Aubert, J.B., de Montleau, C., et al. (2021). Partial recovery of visual function in a blind patient after

optogenetic therapy. Nature medicine *27*, 1223-1229. 10.1038/s41591-021-01351-4.

7.   Polosukhina, A., Litt, J., Tochitsky, I., Nemargut, J., Sychev, Y., De Kouchkovsky, I., Huang, T., Borges, K., Trauner,

D., Van Gelder, R.N., and Kramer, R.H. (2012). Photochemical restoration of visual responses in blind mice.

Neuron *75*, 271-282. 10.1016/j.neuron.2012.05.022.

8.   Tochitsky, I., Polosukhina, A., Degtyar, V.E., Gallerani, N., Smith, C.M., Friedman, A., Van Gelder, R.N., Trauner,

D., Kaufer, D., and Kramer, R.H. (2014). Restoring visual function to blind mice with a photoswitch that exploits

electrophysiological remodeling of retinal ganglion cells. Neuron *81*, 800-813. 10.1016/j.neuron.2014.01.003.

9.   Telias, M., Sit, K.K., Frozenfar, D., Smith, B., Misra, A., Goard, M.J., and Kramer, R.H. (2022). Retinoic acid

inhibitors mitigate vision loss in a mouse model of retinal degeneration. Science advances *8*, eabm4643.

10.1126/sciadv.abm4643.

10.   Pfeiffer, R.L., Marc, R.E., and Jones, B.W. (2020). Persistent remodeling and neurodegeneration in late-stage

retinal degeneration. Progress in retinal and eye research *74*, 100771. 10.1016/j.preteyeres.2019.07.004.

11.   Stasheff, S.F., Shankar, M., and Andrews, M.P. (2011). Developmental time course distinguishes changes in

spontaneous and light-evoked retinal ganglion cell activity in rd1 and rd10 mice. Journal of neurophysiology *105*,

3002-3009. 10.1152/jn.00704.2010.

12.   Telias, M., Denlinger, B., Helft, Z., Thornton, C., Beckwith-Cohen, B., and Kramer, R.H. (2019). Retinoic Acid
Induces Hyperactivity, and Blocking Its Receptor Unmasks Light Responses and Augments Vision in Retinal
Degeneration. Neuron *102*, 574-586 e575. 10.1016/j.neuron.2019.02.015.

13.   Tochitsky, I., Helft, Z., Meseguer, V., Fletcher, R.B., Vessey, K.A., Telias, M., Denlinger, B., Malis, J., Fletcher, E.L.,
and Kramer, R.H. (2016). How Azobenzene Photoswitches Restore Visual Responses to the Blind Retina. Neuron
92, 100-113. 10.1016/j.neuron.2016.08.038.

14.   Marc, R.E., Jones, B.W., Watt, C.B., and Strettoi, E. (2003). Neural remodeling in retinal degeneration. Progress in
retinal and eye research *22*, 607-655. 10.1016/s1350-9462(03)00039-9.

15.   Dunn, F.A. (2015). Photoreceptor ablation initiates the immediate loss of glutamate receptors in postsynaptic
bipolar cells in retina. The Journal of neuroscience : the official journal of the Society for Neuroscience *35*, 2423-
2431. 10.1523/JNEUROSCI.4284-14.2015.

16.   Gayet-Primo, J., and Puthussery, T. (2015). Alterations in Kainate Receptor and TRPM1 Localization in Bipolar
Cells after Retinal Photoreceptor Degeneration. Front Cell Neurosci *9*, 486. 10.3389/fncel.2015.00486.

17.   Lin, Y., Jones, B.W., Liu, A., Tucker, J.F., Rapp, K., Luo, L., Baehr, W., Bernstein, P.S., Watt, C.B., Yang, J.H., et al.
(2012). Retinoid receptors trigger neuritogenesis in retinal degenerations. FASEB journal : official publication of
the Federation of American Societies for Experimental Biology *26*, 81-92. 10.1096/fj.11-192914.

18.   Care, R.A., Kastner, D.B., De la Huerta, I., Pan, S., Khoche, A., Della Santina, L., Gamlin, C., Santo Tomas, C., Ngo,
J., Chen, A., et al. (2019). Partial Cone Loss Triggers Synapse-Specific Remodeling and Spatial Receptive Field
Rearrangements in a Mature Retinal Circuit. Cell reports *27*, 2171-2183.e2175. 10.1016/j.celrep.2019.04.065.

19.   Shen, N., Wang, B., Soto, F., and Kerschensteiner, D. (2020). Homeostatic Plasticity Shapes the Retinal Response
to Photoreceptor Degeneration. Current biology : CB *30*, 1916-1926.e1913. 10.1016/j.cub.2020.03.033.

20.   Tachibana, M., Okada, T., Arimura, T., Kobayashi, K., and Piccolino, M. (1993). Dihydropyridine-sensitive calcium
current mediates neurotransmitter release from bipolar cells of the goldfish retina. The Journal of Neuroscience
13, 2898-2909. 10.1523/jneurosci.13-07-02898.1993.

21.   Heidelberger, R., and Matthews, G. (1992). Calcium influx and calcium current in single synaptic terminals of
goldfish retinal bipolar neurons. J Physiol *447*, 235-256. 10.1113/jphysiol.1992.sp019000.

22.   Tien, N.-W., Soto, F., and Kerschensteiner, D. (2017). Homeostatic Plasticity Shapes Cell-Type-Specific Wiring in
the Retina. Neuron *94*, 656-665.e654. 10.1016/j.neuron.2017.04.016.

23.   Zhu, Y., Xu, J., Hauswirth, W.W., and DeVries, S.H. (2014). Genetically targeted binary labeling of retinal neurons.
The Journal of neuroscience : the official journal of the Society for Neuroscience *34*, 7845-7861.
10.1523/JNEUROSCI.2960-13.2014.

24.   Liang, C.-Q., Zhang, G., Zhang, L., Chen, S.-Y., Wang, J.-N., Zhang, T.-T., Singer, J.H., and Ke, J.-B. (2021).
Calmodulin Bidirectionally Regulates Evoked and Spontaneous Neurotransmitter Release at Retinal Ribbon
Synapses. eneuro *8*, ENEURO.0257-0220.2020. 10.1523/eneuro.0257-20.2020.

25.   Singer, J.H., Lassová, L., Vardi, N., and Diamond, J.S. (2004). Coordinated multivesicular release at a mammalian
ribbon synapse. Nature Neuroscience *7*, 826-833. 10.1038/nn1280.
26.   Puthussery, T., Gayet-Primo, J., and Taylor, W.R. (2010). Localization of the calcium-binding protein secretagogin
in cone bipolar cells of the mammalian retina. The Journal of comparative neurology *518*, 513-525.
10.1002/cne.22234.
27.   Schwartz, G.W., Okawa, H., Dunn, F.A., Morgan, J.L., Kerschensteiner, D., Wong, R.O., and Rieke, F. (2012). The
spatial structure of a nonlinear receptive field. Nat Neurosci *15*, 1572-1580. 10.1038/nn.3225.
28.   Helmstaedter, M., Briggman, K.L., Turaga, S.C., Jain, V., Seung, H.S., and Denk, W. (2013). Connectomic
reconstruction of the inner plexiform layer in the mouse retina. Nature *500*, 168-174. 10.1038/nature12346.
29.   Germain, P., Gaudon, C., Pogenberg, V., Sanglier, S., Van Dorsselaer, A., Royer, C.A., Lazar, M.A., Bourguet, W.,
and Gronemeyer, H. (2009). Differential Action on Coregulator Interaction Defines Inverse Retinoid Agonists and
Neutral Antagonists. Chemistry & Biology *16*, 479-489. https://doi.org/10.1016/j.chembiol.2009.03.008.
30.   Singer, J.H., and Diamond, J.S. (2006). Vesicle depletion and synaptic depression at a mammalian ribbon
synapse. Journal of neurophysiology *95*, 3191-3198. 10.1152/jn.01309.2005.
31.   Gomis, A., Burrone, J., and Lagnado, L. (1999). Two actions of calcium regulate the supply of releasable vesicles
at the ribbon synapse of retinal bipolar cells. The Journal of neuroscience : the official journal of the Society for
Neuroscience *19*, 6309-6317. 10.1523/jneurosci.19-15-06309.1999.
32.   Scalabrino, M.L., Thapa, M., Chew, L.A., Zhang, E., Xu, J., Sampath, A.P., Chen, J., and Field, G.D. (2022). Robust
cone-mediated signaling persists late into rod photoreceptor degeneration. Elife *11*. 10.7554/eLife.80271.
33.   Sekirnjak, C., Jepson, L.H., Hottowy, P., Sher, A., Dabrowski, W., Litke, A.M., and Chichilnisky, E.J. (2011). Changes
in physiological properties of rat ganglion cells during retinal degeneration. Journal of neurophysiology *105*,
2560-2571. 10.1152/jn.01061.2010.
34.   Trenholm, S., and Awatramani, G.B. (2015). Origins of spontaneous activity in the degenerating retina. Frontiers
in Cellular Neuroscience *9*. 10.3389/fncel.2015.00277.
35.   Choi, H., Zhang, L., Cembrowski, M.S., Sabottke, C.F., Markowitz, A.L., Butts, D.A., Kath, W.L., Singer, J.H., and
Riecke, H. (2014). Intrinsic bursting of AII amacrine cells underlies oscillations in the rd1 mouse retina. Journal of
neurophysiology *112*, 1491-1504. 10.1152/jn.00437.2014.
36.   Harper, A.R., Wiechmann, A.F., Moiseyev, G., Ma, J.X., and Summers, J.A. (2015). Identification of active
retinaldehyde dehydrogenase isoforms in the postnatal human eye. PloS one *10*, e0122008.
10.1371/journal.pone.0122008.
37.   McCaffery, P., Mey, J., and Dräger, U.C. (1996). Light-mediated retinoic acid production. Proc Natl Acad Sci U S A
93, 12570-12574. 10.1073/pnas.93.22.12570.
38.   Thapliyal, S., Arendt, K.L., Lau, A.G., and Chen, L. (2022). Retinoic acid-gated BDNF synthesis in neuronal
dendrites drives presynaptic homeostatic plasticity. Elife *11*. 10.7554/eLife.79863.

39. Balmer, J.E., and Blomhoff, R. (2002). Gene expression regulation by retinoic acid. Journal of Lipid Research *43*,
1773-1808. 10.1194/jlr.R100015-JLR200.
40. Dolphin, A.C. (2009). Calcium channel diversity: multiple roles of calcium channel subunits. Current opinion in
neurobiology *19*, 237-244. 10.1016/j.conb.2009.06.006.
41. Hoppa, M.B., Lana, B., Margas, W., Dolphin, A.C., and Ryan, T.A. (2012). α2δ expression sets presynaptic calcium
channel abundance and release probability. Nature *486*, 122-125. 10.1038/nature11033.
42. D'Arco, M., Margas, W., Cassidy, J.S., and Dolphin, A.C. (2015). The upregulation of α2δ-1 subunit modulates
activity-dependent Ca2+ signals in sensory neurons. The Journal of neuroscience : the official journal of the
Society for Neuroscience *35*, 5891-5903. 10.1523/jneurosci.3997-14.2015.
43. Wan, Q.F., and Heidelberger, R. (2011). Synaptic release at mammalian bipolar cell terminals. Visual
neuroscience *28*, 109-119. 10.1017/S0952523810000453.
44. Weingarten, D.J., Shrestha, A., Juda-Nelson, K., Kissiwaa, S.A., Spruston, E., and Jackman, S.L. (2022). Fast
resupply of synaptic vesicles requires synaptotagmin-3. Nature *611*, 320-325. 10.1038/s41586-022-05337-1.
45. Turano, K., and Wang, X. (1992). Motion thresholds in retinitis pigmentosa. Investigative ophthalmology & visual
science *33*, 2411-2422.
46. Alexander, K.R., Derlacki, D.J., Xie, W., Fishman, G.A., and Szlyk, J.P. (1998). Discrimination of spatial
displacements by patients with retinitis pigmentosa. Vision research *38*, 1171-1181.
https://doi.org/10.1016/S0042-6989(97)00235-6.
47. Lagali, P.S., Balya, D., Awatramani, G.B., Münch, T.A., Kim, D.S., Busskamp, V., Cepko, C.L., and Roska, B. (2008).
Light-activated channels targeted to ON bipolar cells restore visual function in retinal degeneration. Nat
Neurosci *11*, 667-675. 10.1038/nn.2117.
48. Roux, S., Matonti, F., Dupont, F., Hoffart, L., Takerkart, S., Picaud, S., Pham, P., and Chavane, F. (2016). Probing
the functional impact of sub-retinal prosthesis. Elife *5*. 10.7554/eLife.12687.
49. Schmidt, T.M., and Kofuji, P. (2011). An Isolated Retinal Preparation to Record Light Response from Genetically
Labeled Retinal Ganglion Cells. JoVE, e2367. doi:10.3791/2367.
50. Tetenborg, S., Yadav, S.C., Hormuzdi, S.G., Monyer, H., Janssen-Bienhold, U., and Dedek, K. (2017). Differential
Distribution of Retinal Ca(2+)/Calmodulin-Dependent Kinase II (CaMKII) Isoforms Indicates CaMKII-β and -δ as
Specific Elements of Electrical Synapses Made of Connexin36 (Cx36). Frontiers in molecular neuroscience *10*,
425. 10.3389/fnmol.2017.00425.
51. Meyer, A., Yadav, S. C., and Dedek, K. (2018). Phenotyping of gap-junctional coupling in the mouse retina.
Methods Mol. Biol. 1753, 249–259. doi: 10.1007/978-1-4939-7720-8_17.
52. Yadav, S.C., Tetenborg, S., and Dedek, K. (2019). Gap Junctions in A8 Amacrine Cells Are Made of Connexin36 but
Are Differently Regulated Than Gap Junctions in AII Amacrine Cells. Frontiers in molecular neuroscience *12*, 99.
10.3389/fnmol.2019.00099.

53.   Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C.,

Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source platform for biological-image analysis. Nat Methods *9*,

676-682. 10.1038/nmeth.2019.

54.   Behrens, C., Yadav, S.C., Korympidou, M.M., Zhang, Y., Haverkamp, S., Irsen, S., Schaedler, A., Lu, X., Liu, Z.,

Lause, J., et al. (2022). Retinal horizontal cells use different synaptic sites for global feedforward and local

feedback signaling. Current biology : CB *32*, 545-558.e545. 10.1016/j.cub.2021.11.055.

55.   Dalkara, D., Byrne, L.C., Klimczak, R.R., Visel, M., Yin, L., Merigan, W.H., Flannery, J.G., and Schaffer, D.V. (2013).

In vivo-directed evolution of a new adeno-associated virus for therapeutic outer retinal gene delivery from the

vitreous. Science translational medicine *5*, 189ra176. 10.1126/scitranslmed.3005708.

Supplemental Figures

Supplemental Figure 1. Voltage-gated Ca^{2+} current is eliminated by the L-type Ca^{2+} channel blocker nimodipine. A)
Retinal circuit diagrams showing whole-cell patch-clamped CBC6 cells. B) Application of 10 μM nimodipine eliminated
voltage-gated Ca^{2+} current in a WT cell, indicating the observed voltage-gated Ca^{2+} current is generated by L-type channels
in CBC6 cells.

Supplemental Figure 2. ON α-RGCs do not express ChR2-eYFP. A) An ON α-RGC patch clamped using bright field microscopy does not express ChR2-eYFP visualized with fluorescence microscopy. An other RGC, that is not an ON α-RGC, patch clamped using bright field microscopy is expressing ChR2-eYFP visualized with fluorescence microscopy. B) Optogenetic stimulation of ON a-RGCs evoked EPSCs at latencies consistent with monosynaptic transmission. Blocking synaptic transmission with DNQX application fully eliminates the EPSC. Optogenetic stimulation of other ChR2-eYFP expressing RGCS generate currents with extremely short latency that are mostly insensitive to synaptic blockade with DNQX.

Supplemental Figure 3. CBC6 cells from WT and rd1 retinas have the same ChR2 current density and depolarize the
same amount in response to optogenetic light. A) Retinal circuit diagram showing whole-cell patch-clamp and
simultaneous optogenetic stimulation of a CBC6 cell. B) ChR2-induced inward current elicited by optogenetic stimulation
of CBC6 cells in WT and rd1. C) There is no difference in ChR2 current density in CBC6s in WT and rd1 retinas (p = 0.15).
D) ChR2-induced membrane depolarization elicited by optogenetic stimulation of CBC6 cells in WT and rd1. E) There is
no difference in the magnitude of ChR2-induced depolarization in CBC6s in WT and rd1 retinas (p = 0.11). All statistical
comparisons were made with PWT with FDR correction.

Supplemental Figure 4. Reduced sensitivity of ON bipolar cell synaptic transmission in degenerated retinas. A,B)
Stimulus vs. response plots from Fig. 2 scaled to the maximal EPSC response. The shift along the x-axis indicates reduced
sensitivity of cone (A) and rod (B) BC synapses in degenerated retinas. C,D) Summary of the stimulus strength required
to elicit half-maximal EPSCs in CBC6 (C) and AII ACs (D). In all cases, 100% stimulus strength was a 10 ms 470 nm flash
delivering $2.1X10^7$ photons. *** indicates p<0.0001, * indicates p=0.02 rd1 vs. WT, p=0.05 rd10 vs. WT. ### indicates
p<0.0001 vs. rd1. All statistical comparisons were made with PWT with FDR correction.

Supplementary Figure 5. BMS 493 does not increase voltage-gated calcium current in WT retina. A) Retinal circuit
diagrams showing whole-cell patch-clamped CBC6 cells. B) Inward calcium currents from CBC6 cells from WT and BMS
493-treated retinas. Current was activated with a depolarizing voltage step (50 ms) from -60 mV to -20 mV. C) Steady-
state current vs. voltage (I-V) curves, elicited with a series of 10 mV incrementing depolarizing steps from -60 mV. There
was no significant difference between WT and BMS 493-treated retinas. All statistical comparisons were made with PWT
with FDR correction.

Supplemental Figure 6. **Degeneration does not alter the ability of ON a-RGCs to fire in response to high frequency**

stimulation. A) Retinal circuit diagrams showing whole-cell patch-clamped ON a-RGCs. B,C) Action potentials from a WT

and rd1 ON a-RGC in response to 2.5 ms, 500 pA direct current injection at a frequency of 1 Hz (B) and at 7.5 Hz (C). D,E)

There is no difference in ON a-RGCs from WT and rd1 responding to super-threshold current injection at a frequency of

1 Hz (p =1) and at 7.5 Hz (p=1). N = 6. All statistical comparisons were made with PWT with FDR correction.