1   # Soil health is linked to primary productivity across Europe

2
3   Ferran Romero*[1], Maëva Labouyrie[1,2,3], Alberto Orgiazzi[3,4], Cristiano Ballabio[3], Panos Panagos[3], Arwyn
4   Jones[3], Leho Tedersoo[5], Mohammad Bahram[6,7], Linnea C. Smith[8,9], Carlos A. Guerra[8,9], Emanuele
5   Lugato[3], Nico Eisenhauer[8,10], Manuel Delgado-Baquerizo[11], Marcel G.A. van der Heijden*[1,2]
6
[1]Plant-Soil Interactions group, Agroscope, Reckenholzstrasse 191, 8046 Zurich, Switzerland

[2]Department of Plant and Microbial Biology, University of Zurich, Zollikerstrasse 107, 8008 Zurich,
Switzerland

[3]European Commission, Joint Research Centre Ispra (JRC Ispra), Via Enrico Fermi 2749, 21027 Ispra, Italy

[4]European Dynamics, Brussels, Belgium

[5]Mycology and Microbiology Center, University of Tartu, Liivi 2, 50409 Tartu, Estonia

[6]Department of Botany, Institute of Ecology and Earth Sciences, University of Tartu, Liivi 2, 50409 Tartu,
Estonia

[7]Department of Ecology, Swedish University of Agricultural Sciences, Uppsala, Sweden

[8]German Center for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Leipzig, Germany

[9]Institute of Biology, martin Luther University Halle Wittenberg, Halle (Saale), Germany

[10]Institute of Biology, Leipzig University, Leipzig, Germany

[11]Laboratorio de Biodiversidad y Funcionamiento Ecosistémico. Instituto de Recursos Naturales y
Agrobiología de Sevilla (IRNAS), CSIC, Av. Reina Mercedes 10, E-41012, Sevilla, Spain.
21

*Authors for correspondence: Ferran Romero (ferran.romeroblanch@agroscope.admin.ch) and Marcel
van der Heijden (marcel.vanderheijden@agroscope.admin.ch)

Abstract

The role of soil biodiversity and soil health in regulating primary productivity across different
land use types is still poorly understood, hindering our ability to predict the impact of soil
degradation on essential ecosystem services such as food provision. To address this gap, we
conducted a pan-European observational field study using data from 588 sites and 27 countries
to investigate the linkages between soil microbial diversity, soil health, and primary productivity
across three major land use types: woodlands, grasslands, and croplands. We demonstrate that
a combination of climatic and edaphic factors, together with soil biodiversity parameters,
explained between 20.76% and 31.20% of the variability in primary productivity across
contrasting land use types. Our study further revealed a general positive effect of soil health
(i.e., microbial biomass, soil nitrogen and carbon content, and microbial diversity) on primary
productivity, particularly for croplands ($R^2 = 0.16$) and grasslands ($R^2 = 0.18$). The effects of soil
microbial diversity depended on the microbial group and the land use type considered. In
general, higher diversity of beneficial groups (e.g., Acidobacteria, Mortierellomycota) was linked
to increased productivity, while the opposite was found for plant pathogens (particularly in
croplands). Together, our results provide insights into the importance of soil biodiversity and
soil health for maintaining essential ecosystem functions across contrasting land use types and
highlight the need for land-use specific management strategies for preserving belowground
diversity and primary productivity.

Introduction

Plant productivity is a crucial ecosystem process that supports essential services necessary for human welfare on Earth (Barrios, 2007). A range of studies have shown that plant diversity in ecosystems promotes plant productivity and yield (Dietrich et al., 2023; Fraser et al., 2015; Isbell et al., 2011; Le Provost et al., 2023; Tilman et al., 1997). On the contrary, it is still poorly understood whether the diversity of soil biota (i.e., belowground organisms) influences plant productivity. In addition to this, our understanding of how soil health relates to primary productivity at large scales remains poorly understood.

Soil health is a composite measure and refers to the capacity of soils to support plant productivity and other functions (Lehmann et al., 2020; Qiao et al., 2022). Soil health assessments have traditionally focused on soil properties including microbial biomass, available nutrients, and organic carbon (Bünemann et al., 2018). However, advanced molecular methods such as next-generation sequencing techniques offer the possibility to include a microbial component (e.g., soil biodiversity) into soil health indices, providing a more comprehensive view of the wide range of properties that characterize soils and influence ecosystem functioning. Although there is much interest globally on soil health, current scientific evidence that demonstrates the importance of soil health at large scale is missing (Lehmann et al., 2020; Rinot et al., 2019). Despite of this, a recent study assessing croplands in China demonstrated that soil health is positively linked to plant yield (Li et al., 2021). However, systematic comparisons of the relative importance that soil health has across contrasting ecosystems (i.e., woodlands, grasslands) have received less attention. Moreover, the studies performed so far have largely focused on chemical and physical properties, often ignoring biological measure (Van Bruggen & Semenov, 2000). The few studies including biological measures in soil quality have mostly focused on the activity of soil microorganisms (e.g., enzymatic activities), but biodiversity *per se* has received less attention (Alkorta et al., 2003).

Experiments performed in experimental microcosms where soil biodiversity can be manipulated (i.e., reduced) have demonstrated that a reduction of microbial diversity (e.g., number of bacterial and fungal taxa) leads to altered microbial mediated functions, less nutrient uptake by plants, and reduced productivity (Chen et al., 2020; Luo et al., 2023; Romero, Hilfiker, et al., 2023; van der Heijden et al., 1998; Wagg et al., 2019). However, only recently medium to large-scale monitoring studies have started to uncover the role that belowground diversity plays in ecosystem productivity (Anthony et al., 2022; Delgado-Baquerizo et al., 2016, 2020; Fan et al., 2021; Hu et al., 2021; Liu et al., 2022). Most of these studies have so far focused on natural or

semi-natural land use types (e.g., grasslands), or contained a limited number of study sites

(Delgado-Baquerizo et al., 2020; Fan et al., 2023; Hu et al., 2021). Moreover, the specific drivers

of plant productivity, especially in intensively managed ecosystems (i.e., croplands) are still

poorly understood, and it is widely debated whether soil microbial diversity and microbiome

complexity are important for plant productivity in these ecosystems (van der Heijden & Wagg,

2013; Xiong & Lu, 2022).

Studies to date have primarily focused on microbial-mediated functions such as carbon storage

and nitrogen cycling, leaving a significant knowledge gap in understanding the broader effects

of belowground diversity on other functions (i.e., plant productivity) across different ecosystems

(Bissett et al., 2011; Huang et al., 2023; Wu et al., 2023). Malik and collaborators (2018) used 21

paired sites, comprising low- and high-intensity management practices (permanent grasslands

vs. fertilized croplands), and observed higher carbon storage capacity attributed to enhanced

microbial growth efficiency in undisturbed sites (grasslands) compared to heavily fertilized

croplands. Similarly, Edlinger and collaborators (2023) observed lower soil aggregate stability

and soil organic carbon content in croplands compared with neighboring grassland sites across

a 3000 Km European gradient. Also recently, it was observed that soil microbial diversity at the

European scale is highest in croplands compared to woodlands and grasslands (Karimi et al.,

2019; Köninger et al., 2023; Labouyrie et al., 2023; Szoboszlay et al., 2017), but the question of

whether productivity is similarly driven by belowground diversity across contrasting land use

types (e.g., woodland and grasslands vs. cropland) remains unexplored, as a systematic large-

scale comparison among different land uses is lacking and large scale soil monitoring efforts are

scarce.

In this article, we tested the hypothesis of whether the relationship between soil health and

primary productivity differs across contrasting land-use types. To that purpose, we used a pan-

European observational field study to test the linkages between soil microbial diversity, soil

properties, and primary productivity. We used fresh soil samples collected from 588 sites in 2018

and covering 27 countries and the three major land-use types in the European continent:

woodlands (n = 185), grasslands (n = 126), and croplands (n = 277). Using next-generation

sequencing methods targeting ribosomal marker genes, we obtained site-level information on

the diversity of bacteria and fungi, as well as their major phyla, and functional groups (e.g.,

fungal pathogens, nitrogen-fixing bacteria). The analysis demonstrated that soil microbial

diversity differed across land use types in Europe (Labouyrie et al., 2023). Here we used data

from that earlier study and linked this with 11 variables representing information on climatic

and edaphic factor. Moreover, we used remote sensing to obtain primary productivity data
across the 588 sites. Subsequently we explored how soil health, soil microbial diversity and a
range of biotic and abiotic factors relate to primary productivity across contrasting land use
types.

Methods

Field survey and soil sampling

This study was carried out on the EU Statistical Office's Land Use and Coverage Area frame
Survey (LUCAS), the largest pan-European scheme for assessing soil characteristics in relation to
land cover and land use (Orgiazzi et al., 2018). As part of the 2018 LUCAS survey, 881 fresh soil
samples were collected across Europe and stored at -20°C before further processing (Orgiazzi et
al., 2022). The sampling followed a composite strategy: the final sample collected at each
location comprised five topsoil (0-20 cm) subsamples that are mixed to form a single composite
sample. The first subsample is taken at the precise GIS (geographical information system)
coordinates of the pre-established LUCAS point, and the remaining four subsamples are taken 2
m from the central one following the cardinal directions (North, East, South, and West). To avoid
seasonality and time effects, all samples were collected over the shortest possible timeframe
(spring 2018). For this study, we focused on 277 croplands, 126 grasslands, and 185 woodlands
from 881 LUCAS sites, across 27 countries (Figure 1). The selection was based on those samples
from which all variables (soil diversity, edaphic factors, and primary productivity) were available.
Sampling was based on a regular grid defined by the intersection points of a 2 km x 2 km grid
covering the European territory, and each sampling location was assigned to one of three land
use types (woodland, grassland, cropland) using orthophotos or satellite images (Orgiazzi et al.,
2018).

Soil microbial diversity

Soil DNA was extracted from each of the 588 composite soil samples using the Qiagen DNeasy
PowerSoil HTP 96 Kit Q12955-4. Three 0.2 g aliquots per sample were extracted and pooled
together. Negative and positive controls were included to determine the presence of any
external contamination. DNA extracts were quantified and checked for quality using a Qubit™
1X dsDNA HS Assay Kit and a Qubit™ 3 Fluorometer (Invitrogen). Primer sets for barcode
amplification were 515F (GTGYCAGCMGCCGCGGTAA) and 926R (GGCCGYCAATTYMTTTRAGTTT)
for the bacterial 16S region (Caporaso et al., 2011; Parada et al., 2016), and ITS9mun

(GTACACACCGCCCGTCG) and ITS4ngsUni (CGCCTSCSCTTANTDATATGC) for the fungal ITS region

(Tedersoo & Anslan, 2019). Sequencing was performed by Illumina MiSeq platform with 2 x 300

paired-end mode for bacterial data and PacBio Sequel II platform for fungal data. Additional

information including DNA amplification and library preparation is available in Labouyrie et al.,

2023.

The Illumina and PacBio amplicon data (for bacteria and fungi, respectively) were demultiplexed

using LotuS2 (Özkurt et al., 2022), and paired-end reads were assembled setting a minimum

overlap of 10 bp using FLASH 1.2.10 (Magoč & Salzberg, 2011). Reads failing to merge were

removed from the final dataset. For bacteria, operational taxonomic units (OTUs) were

generated with the usearch option (version 10.0.024) available in UPARSE (Edgar, 2013). Merged

reads were truncated up to the 16S primer sequences (i.e., 515F and 926R) and filtered for the

presence of primer sequences (allowing a maximum of 2 mismatches). Merged reads were

further quality-filtered with usearch by removing sequences with more than one expected error,

and duplicated sequences were collapsed with fqtrim version 0.9.7 (Pertea, 2015). Obtained

OTUs were annotated with the taxonomy data available from the Ribosomal Database Project

(Cole et al., 2014). To avoid sequencing artifacts, only OTUs with at least 50 read counts and

present in at least 10 samples were kept. For fungi, 98%-OTUs were obtained as described in

Tedersoo et al., 2021. Fungal taxonomic annotation was performed using BLAST+ version 2.11.0

(Camacho et al., 2009) against the UNITE 9.1 database (Nilsson et al., 2019). Bacterial and fungal

datasets were further normalized to comparable sequencing depths (bacteria: 40,109 reads;

fungi: 502 reads) using the Scaling with Ranked Subsampling (SRS) method (Beule & Karlovsky,

2020). Normalized datasets were used to obtain values of total richness (i.e., OTU number),

community composition (extracted values from the first dimension of NMDS), and the richness

of major microbial groups (i.e., phyla). Normalized datasets were also used to obtain the richness

of major functional groups as retrieved from the FAPROTAX (bacteria) and FungalTraits (fungi)

databases (Louca et al., 2016; Põlme et al., 2020). Further details on bacterial/fungal sequence

processing, filtering, and taxonomic/functional annotation are available in Labouyrie et al.,

2023.

Edaphic and climatic factors

The following measurements were undertaken for each sampling site to ensure a

comprehensive understanding of climatic and edaphic factors. Total nitrogen and phosphorus

content were assessed to determine nitrogen and phosphorus availability in the soil. Soil organic

carbon was quantified to estimate the amount of carbon stored in the soil organic matter. Soil

pH and microbial biomass were used as measures of acidity and activity of the living component

of the soil, respectively. The bulk density was used as a surrogate of soil compactness. Soil

texture was characterized by measuring silt, sand, and clay content. Additionally, to capture the

broader climatic context of each site, mean annual precipitation and mean annual temperature

were retrieved from the WorldClim database (www.worldclim.org) . These parameters

collectively provided a detailed profile of the environmental and soil conditions at each sampling

site. Specific details on the methods employed for these measurements are available in

Labouyrie et al., 2023 and Smith et al., 2021.

Primary productivity

The normalized difference vegetation index (NDVI) was calculated from images taken with the

NASA satellites Terra and Aqua using moderate resolution imaging spectroradiometer (MODIS)

technology. The NVDI is a simple indicator for the health and abundance of vegetation in a

particular area and is calculated using the difference between the amount of light reflected by

vegetation in the near-infrared range and the amount of light reflected in the red range

(Pettorelli et al., 2005). The NDVI is widely accepted as a proxy for plant biomass and

photosynthetic efficiency (Camps-Valls et al., 2021; Jobbágy et al., 2002; Joiner et al., 2018;

Paruelo et al., 1997) across different ecosystems, and has previously been used to investigate

the relationship between soil biodiversity and plant productivity (Delgado-Baquerizo et al.,

2016, 2020; Fan et al., 2023; Liu et al., 2022). Here, we used averaged NVDI values obtained for

the months before, during, and after sampling at each location. This index was calculated in the

same way for croplands, grasslands, and woodlands.

Statistics

Statistical analyses were performed using R software, version 4.2.1. Only soil samples for which

all explanatory variables (i.e., soil biodiversity and climatic, edaphic, and spatial factors) were

available were used for this study, so none of the values in any explanatory variables was

inferred or calculated from the median value of the respective dataset. A total of 588 sites were

used for this study, including 277 croplands, 126 grasslands, and 185 woodlands. To assess the

most important drivers of primary productivity within the three land-use systems, we performed

a three-step model-selection approach (Edlinger et al., 2022).

First, we selected a wide range of explanatory variables including biodiversity indicators (i.e.,

OTU richness, diversity, community composition), together with climatic and edaphic factors.

Total number of explanatory variables was 34 (Table S1). We then investigated the correlation
among all variables to ensure there was no strong co-correlation among them, and reduced
variables that were strongly correlated (i.e., Spearman rank correlation coefficient higher than
+0.90, or lower than -0.90). Correlations were run using the corr function within the corrplot
package (Wei & Simko, 2017). This approach reduced the 34 explanatory variables to 29 (Table
S1). The five variables removed included Shannon's diversity index for bacteria and fungi (strong
correlation with OTU richness), total nitrogen (strong correlation with soil organic carbon),
richness of chemoheterotrophic bacteria (strong correlation with Actinobacteria and total
bacteria), and Proteobacteria (strong correlation with total bacterial richness and
Bacteroidetes). Some of the resulting variables (selection based on Lehmann et al., 2020, see
Table S2) were used to build a soil health index: this was calculated using z-score transformation:
each variable was standardized to have a mean of 0 and a standard deviation of 1. The soil health
index for each site was then computed by summing its respective z-scores. This approach
allowed us to integrate multiple soil health indicators into a single composite index, facilitating
a comprehensive assessment.

The second step involved a multi-model inference approach using the R package glmulti
(Calcagno & de Mazancourt, 2010). This package automates the process of exploring different
combinations of explanatory variables and selects the most appropriate model to explain
variability in each outcome (here, primary productivity). Multi-model inference approach
resulted in a high number of possible linear models, ranked by Akaike information criterion (AIC).
In the last step, the best models that fell within an AIC range of two were averaged to obtain a
conservative estimate and relative importance of the most important predictors using the
MuMin package (Barton, 2012). Briefly, this package combines the results of multiple models to
get more robust and reliable estimates. We used ΔAIC values to compute the Akaike weight of
each of the selected models and computed a weighted-average model. The relative importance
of each explanatory variable was estimated by taking all the models into consideration as the
sum of the weights/probabilities for the models in which the variable appears.

To further clarify the relative contribution of environmental factors and soil biodiversity to
explain patterns in primary productivity, we conducted a classification Random Forest (RF)
analysis. The RF analyses were performed independently for soil biodiversity, environmental
parameters, and both combined. The RF were also performed for each land use type
independently (croplands, grasslands, and woodlands). Therefore, a total of nine RF analyses
were performed using the rfPermute package (Archer, 2016). For each RF, we obtained the

variation (as %) explained by the different predictors (soil biodiversity, environmental
parameters, or both combined).

Results and discussion

This study represents the first continental-scale comparison of patterns on soil biodiversity-
primary productivity across contrasting land uses. We surveyed 588 sites across the European
continent including 277 croplands, 126 grasslands, and 185 woodlands (Figure 1). For each site,
we gathered information on environmental parameters (i.e., climatic, and edaphic factors),
together with information on soil microbial (i.e., bacterial, and fungal) diversity, and primary
productivity (Table 1). Primary productivity was significantly lower in croplands, compared to
grasslands and woodlands, in line with previous research (Krause et al., 2022) (Figure S1).
Overall, our results demonstrate that a combination of climatic and edaphic factors, together
with soil biodiversity, explain patterns in primary productivity across contrasting land use types
at the European scale: our predictors explained 20.76% of the variability in primary productivity
in grasslands, 23.80% in woodlands, and 31.20% in croplands (Figure 2). Moreover, we show
that a soil health index built from soil chemical properties but also biodiversity (Table S2)
positively correlates to primary productivity, especially in croplands (R^2 = 0.16) and grasslands
(R^2 = 0.18).

We show that soil biodiversity accounts for a unique proportion of the variation in primary
productivity, especially in grasslands. Among soil biodiversity components, bacterial richness
(particularly the richness of Acidobacteria and Firmicutes) was positively associated to primary
productivity in croplands and grasslands, while woodlands relied more on fungi (saprotrophic
fungi and the phylum Mortierellomycota). Bacterial diversity (i.e., OTU number, Shannon's
diversity index) was higher in croplands, followed by grasslands, and woodlands (Table 1, Figure
S2). Fungal richness was highest in grasslands and lowest in woodlands. Fungal diversity
(Shannon's index) was not significantly different between grasslands and croplands (Table 1,
Figure S3). Regarding community composition, both bacterial and fungal communities markedly
differed across land use types (Table 1, Figure S2, Figure S3). By-phylum richness analysis
revealed that Acidobacteria, Basidiomycota, and Leotiomycetes (Ascomycota) were more
numerous in woodlands, whereas Verrucomicrobia were more numerous in grasslands.
Croplands showed the highest richness of Sordariomycetes (Ascomycota), Actinobacteria,
Proteobacteria, Nitrospirae, Bacteroidetes, and Firmicutes (Table 1, Figure S4, Figure S5). At the
functional level, nitrogen-fixing bacteria where more abundant in woodlands and grasslands

than in croplands. Croplands, however, displayed the highest abundance of bacterial
chemoheterotrophs and fungal plant pathogens (Table 1, Figure S6).

We used a multi-model inference approach followed by random forest regression to evaluate
the performance of predictor combinations in explaining patterns in primary productivity across
land use types. Moreover, we used 8 variables known to be often associated to soil health to
build a soil health index and observed positive correlations between this index and primary
productivity, especially in grasslands (R^2 = 0.18), and croplands (R^2 = 0.16) (Figure 3). Among the
different variables included in the index (see Table S2), the strongest correlations with primary
productivity were found for total nitrogen content, soil organic content, and microbial biomass
(Figure S7) Multi-model inference resulted in a high number of possible linear models, ranked
by their Akaike Information Criterion (AIC). We weighted the best models into an average model
(Table 2) and obtained a conservative estimate and relative importance of the most important
predictors (Figure 4) (Cade, 2015; Edlinger et al., 2022). In croplands, the model best explaining
patterns in primary productivity (AIC = 4780.66, p-value < 0.001) included the richness (OTU
number) of several microbial groups (i.e., Acidobacteria, Actinobacteria, Firmicutes,
Chytridiomycota) as well as three edaphic properties (sand content, pH, and microbial biomass).
Our results confirm previous evidence on the role of pH in driving crop productivity; accordingly,
we observed a clear decrease in cropland productivity in highly alkaline (pH > 8) soils (Figure S7)
(Couto, 2018).

Soil microbial biomass was also selected as an important variable explaining plant productivity.
Interestingly, a recent study indicated that high microbial biomass in croplands enhances
nutrient use efficiency and nutrient cycling of plants, so we argue that further work is necessary
to assess the importance of microbial biomass for agroecosystem performance (Toda et al.,
2023). Predictor ranking based on relative importance following multi-model averaging
indicated that the best predictor of primary productivity in croplands was the richness of
Acidobacteria (Figure 4). Acidobacteria are abundant soil-borne microorganisms known to be
involved in nutrient mineralization and organic matter degradation (Banerjee et al., 2016; Kielak
et al., 2016; Sikorski et al., 2022). Here we demonstrate for the first time that the positive effect
of Acidobacteria on primary productivity is comparatively higher in croplands than in woodlands
or grasslands (Figure S8). Our results align with a recent study where Acidobacteria positively
correlated with cropland productivity in a 4-decade fertilization experiment (Fan et al., 2021).
Finally, the richness of Chytridiomycota negatively correlated to productivity in croplands
(Figure S9). While this fungal group is best known for parasitizing vertebrates, some species are

responsible for plant disease in Europe (van de Vossenberg et al., 2022). In line with this, we
observed a negative correlation between fungal plant pathogens and cropland productivity
(Figure S10).

Our multi-model inference approach points towards soil organic carbon as a major predictor of
grassland productivity (Figure 4). Grasslands store one third of the terrestrial carbon stock, and
most of it is stored belowground as soil organic carbon (Conant, 2012). Recent studies show that
plant diversity increases soil organic carbon storage in grasslands by elevating carbon inputs to
belowground biomass and promoting microbial necromass contribution to soil organic carbon
storage (Lange et al., 2015). Our results show that not only plant diversity is interlinked with soil
organic carbon, but also plant productivity at the continental scale. We did not differentiate
between necromass and living biomass in our study, but we also found positive correlations
between bacterial richness/ microbial biomass and grassland productivity. As for croplands, pH
and grassland productivity described a hump-shaped response, with productivity peaking at pH
≈ 6.

Soil organic carbon was also retained as an important predictor of primary productivity in
woodlands (Figure 4). Other relevant environmental factors in woodland productivity were
temperature, phosphorus, and clay content. However, the main predictors of primary
productivity in woodlands according to our multi-model inference approach were bacterial and
fungal community composition (NMDS1). Our results align with a recent study suggesting that
fungal community composition outcompetes richness (number of fungal OTUs) as a driver of soil
functioning in woodlands (Anthony et al., 2022; Osburn et al., 2023). We argue, however, that
future studies at a large scale should include the bacterial compartment in their analyses, as
both previous research and the findings of this study show that the distribution of bacterial
groups plays a role in woodland productivity.

Previous studies have suggested that functional groups (e.g., mycorrhizal fungi, nitrogen-fixing
bacteria, plant pathogens) might be better predictors of ecosystem functioning than the
richness or diversity of taxonomy-based microbial groups (i.e., OTUs) (Delgado-Baquerizo et al.,
2020; Liu et al., 2022). For example, Delgado-Baquerizo and collaborators (2020) observed
positive relationships between the richness of mycorrhizal fungi and soil functions such as
phosphorus mineralization and soil respiration. The same study observed no relationship
between mycorrhizal or saprophytic fungi and primary productivity. Another large-scale study
observed a positive relationship between saprophytic fungi and the stability of primary
productivity, and no relationship between productivity and mycorrhizal fungi (Liu et al., 2022).

Our findings indicate a positive trend of increased productivity associated with higher
mycorrhizal richness, especially within grassland ecosystems (Figure S11). We argue that the
variability in outcomes across various large-scale studies underscores the urgency for additional
research conducted under controlled conditions, aiming to elucidate the precise mechanisms
through which mycorrhizal diversity fosters plant productivity (Romero et al., 2023) . Finally, our
study aligns with previous research showing a decrease in cropland productivity with increasing
pathogen richness (Liu et al., 2022).

In summary, our results provide compelling evidence that a combination of environmental
factors and soil biodiversity relate to primary productivity at the continental scale. This study
shows that between 20.76% and 31.2% of the variability in primary productivity can be explained
from information on microbial (fungal and bacterial) community composition, and edaphic/
climatic parameters. We further demonstrate that bacterial community composition exerts a
significant influence on primary productivity across diverse land use types, whereas in
woodlands, fungal community composition also governs primary productivity. Overall, our study
emphasizes the importance of addressing relationships between soil biodiversity and primary
productivity across contrasting land use types, and further highlights the need to include
microorganisms in soil monitoring schemes to gain a comprehensive understanding of
ecosystem functioning in terrestrial ecosystems.

Figure 1. Study site locations. Geographic distribution across the European continent of the 589 locations used in this study. Three major land use types were covered in the survey: cropland (n = 277, yellow), grassland (n = 126, green), and woodland (n = 185, purple).

Figure 2. Importance of soil biodiversity and environmental parameters in explaining patterns in primary productivity. Variance in primary productivity (%) as obtained from random forest regression explained by soil biodiversity (bacteria and fungi), environmental parameters (climate and edaphic factors), or the combination of both, across land use types (croplands, grasslands, and woodlands).

Figure 3. Relationship between soil health and primary productivity. Linear regressions were fitted to explore the relationship between the soil health index and primary productivity across land use types (woodlands, grasslands, and croplands). Explained variation is expressed as R^2, and significance is indicated by asterisks (***; p-value < 0.001, **; p-value < 0.010, *; p-value < 0.050). Non-significant (p-value > 0.05) regressions are denoted with "n.s.".

Figure 4. Relative importance of the main predictors on primary productivity. Relative

importance of main predictors for primary productivity based on multi-model inference

analysis in woodlands (n = 185), grasslands (n = 126), and croplands (n = 277). Dark colors

indicate positive predictor effects on primary productivity, and light colors indicate negative

effects. Asterisks indicate a significant effect based on the averaged model coefficients (***;

p-value < 0.001, **; p-value < 0.010, *; p-value < 0.050).

		Units	Croplands (n = 277)		Grasslands (n = 126)		Woodlands (n = 185)		Kruskal-Wallis test (Chi-squared)	Kruskal-Wallis test (p-value)
Climatic factors	Precipitation	mm	48,54 ± 23,64	a	68,89 ± 28,57	b	64,07 ± 27,97	b	56,82	< 0,001
	Temperature	°C	17,3 ± 4,91	a	15,13 ± 3,5	b	14,44 ± 4,85	b	44,61	< 0,001
Edaphic factors	Clay content	(%)	23,31 ± 13,31	a	22,23 ± 11,35	a	14,73 ± 12,33	b	64,64	< 0,001
	Sand content	(%)	35,41 ± 26	a	36,61 ± 20,97	a	47,72 ± 24,54	b	30,57	< 0,001
	Silt content	(%)	41,13 ± 19,43	a	40,86 ± 16,76	a	37,53 ± 17,27	a	3,97	> 0,05
	pH	unitless	7,07 ± 1,02	a	6,29 ± 1,02	b	5,16 ± 1,12	c	222,42	< 0,001
	Microbial biomass	µg C/g soil	278,05 ± 156,11	a	455,45 ± 272,69	b	394,4 ± 335,85	c	44,53	< 0,001
	Organic carbon	g/Kg	18,61 ± 14,93	a	36,5 ± 33,8	b	59,23 ± 64,51	c	170,23	< 0,001
	Phosphorus content	mg/Kg	36,65 ± 30,45	a	22,81 ± 23,82	b	19,59 ± 21,41	b	60,48	< 0,001
	Nitrogen content	mg/Kg	1,82 ± 1,15	a	3,36 ± 2,54	b	3,25 ± 2,71	b	81,00	< 0,001
	Bulk density	g/cm3	1,23 ± 0,2	a	1,06 ± 0,23	b	0,77 ± 0,36	c	202,79	< 0,001
Biodiversity (fungi)	Total richness	OTU number	120,46 ± 26,67	a	129,48 ± 32,4	b	104,69 ± 30,2	c	57,91	< 0,001
	Shannon's index	unitless	3,86 ± 0,53	a	3,93 ± 0,64	a	3,48 ± 0,66	b	55,91	< 0,001
	Community composition (NMDS1)	unitless	0,10 ± 0,07	a	0,03 ± 0,09	b	-0,16 ± 0,16	c	277,43	< 0,001
	Basidiomycota	OTU number	17,34 ± 7,33	a	23,17 ± 10,94	b	33,23 ± 11,8	c	210,27	< 0,001
	Chytridiomycota	OTU number	8,61 ± 4,61	a	7,8 ± 4,87	a	2,39 ± 3,27	b	222,66	< 0,001
	Mucoromycota	OTU number	2,08 ± 1,97	a	3,72 ± 3,55	b	3,22 ± 3,06	b	30,47	< 0,001
	Ascomycota	OTU number	71,69 ± 17,66	a	68,52 ± 23,64	a	51,94 ± 20,51	b	100,01	< 0,001
	Mortierellomycota	OTU number	9,19 ± 4,02	a	9,4 ± 4,59	a	6,54 ± 4,84	b	58,42	< 0,001
Biodiversity (bacteria)	Total richness	OTU number	7417,12 ± 1122,83	a	6959,97 ± 1079,26	b	5797,98 ± 1648,39	c	111,54	< 0,001
	Shannon's index	unitless	7,92 ± 0,3	a	7,75 ± 0,32	b	7,34 ± 0,52	c	164,98	< 0,001
	Community composition (NMDS1)	unitless	-0,13 ± 0,14	a	-0,02 ± 0,15	b	0,21 ± 0,21	c	232,37	< 0,001
	Acidobacteria	OTU number	965,81 ± 251,36	a	1035,03 ± 230,47	b	1089,4 ± 210,19	c	30,44	< 0,001
	Actinobacteria	OTU number	1694,05 ± 328,01	a	1522,75 ± 428,15	b	1012,31 ± 561,86	c	154,85	< 0,001
	Proteobacteria	OTU number	2385,02 ± 426,75	a	2149,83 ± 402,81	b	1862,84 ± 587,83	c	102,79	< 0,001
	Bacteroidetes	OTU number	643,71 ± 214,04	a	529,41 ± 182,11	b	369,04 ± 228,53	c	138,8	< 0,001
	Nitrospirae	OTU number	19,6 ± 9,33	a	14,44 ± 9,73	b	6,27 ± 8,35	c	197,23	< 0,001
	Verrucomicrobia	OTU number	403,29 ± 134,63	a	441,89 ± 144,25	b	388,28 ± 126,25	a	11,95	< 0,001
	Firmicutes	OTU number	152,34 ± 81,8	a	118,16 ± 54,12	b	55,24 ± 47,4	c	228,84	< 0,001
Biodiversity (functional groups)	Bacterial chemoheterotrophs	OTU number	1676,78 ± 288,57	a	1339,74 ± 335,02	b	929,82 ± 472,29	c	246,99	< 0,001
	Nitrogen-fixing bacteria	OTU number	9,04 ± 4,68	a	9,87 ± 5,23	ab	9,94 ± 3,7	b	10,28	< 0,001
	Fungal saprotrophs	OTU number	71,44 ± 15,75	a	67,11 ± 19,95	a	50,41 ± 19,02	b	129,11	< 0,001
	Fungal plant pathogens	OTU number	11,01 ± 4,98	a	7,12 ± 4,81	b	1,55 ± 2,28	c	324,32	< 0,001
	Mycorrhizal fungi	OTU number	3,26 ± 4,52	a	6,61 ± 6,03	b	20,63 ± 12,05	c	286,89	< 0,001

Table 1. Mean values of spatial, climatic, and edaphic factors (± standard deviation) in soils from cropland (n = 277), grassland (n = 126), and woodland (n = 185) sites used in this study. Mean values of biodiversity measures are also shown for bacteria, fungi, and predicted functional groups. Kruskal-Wallis test was used to assess the overall effect of land use type on explanatory variables (chi-squared and p-value are indicated for each test). Grouping information following pairwise Wilcoxon test is indicated above each boxplot, different letters indicate significant differences (p < 0.05).

Land Use	Sites	Best model	AIC	p-value
Croplands	277	NPP ~ 1 + Acidobacteria + Actinobacteria + Firmicutes + Sand content + pH + Microbial biomass + Chytridiomycota	4780.66	< 0.001
Grasslands	126	NPP ~ 1 + Firmicutes + Bacterial richness + Clay content + Sand content + pH + Soil organic C + Phosphorus content + Soil density + Basidiomycota + Chytridiomycota + Mycorrhizal richness	2168.70	< 0.001
Woodlands	185	NPP ~ 1 + Bacteroidetes + Bacteria (NMDS1) + Clay content + Soil organic C + Phosphorus content + Temperature + Fungi (NMDS1) + Chytridiomycota + Mortierellomycota + Saprotrophic fungi	3208.55	< 0.001

Table 2. Output of model selection. Best models explaining variation in primary productivity across land use types following multi-model averaging inference.

References

Alkorta, I., Aizpurua, A., Riga, P., Albizu, I., Amézaga, I., & Garbisu, C. (2003). Soil
enzyme activities as biological indicators of soil health. *Reviews on Environmental*
Health, *18*(1), 65–73.
Anthony, M. A., Crowther, T. W., van der Linde, S., Suz, L. M., Bidartondo, M. I., Cox, F.,
Schaub, M., Rautio, P., Ferretti, M., Vesterdal, L., De Vos, B., Dettwiler, M.,
Eickenscheidt, N., Schmitz, A., Meesenburg, H., Andreae, H., Jacob, F., Dietrich, H.-
P., Waldner, P., … Averill, C. (2022). Forest tree growth is linked to mycorrhizal
fungal composition and function across Europe. *The ISME Journal*, *16*(5), 1327–
1336. https://doi.org/10.1038/s41396-021-01159-7
Archer, E. (2016). rfPermute: estimate permutation p-values for random forest
importance metrics. *R Package Version*, *1*(2).
Banerjee, S., Kirkby, C. A., Schmutter, D., Bissett, A., Kirkegaard, J. A., & Richardson, A.
E. (2016). Network analysis reveals functional redundancy and keystone taxa
amongst bacterial and fungal communities during organic matter decomposition
in an arable soil. *Soil Biology and Biochemistry*, *97*, 188–198.
https://doi.org/https://doi.org/10.1016/j.soilbio.2016.03.017
Barrios, E. (2007). Soil biota, ecosystem services and land productivity. *Ecological*
Economics, *64*(2), 269–285.
https://doi.org/https://doi.org/10.1016/j.ecolecon.2007.03.004
Barton, K. (2012). MuMIn: Multi-model inference. R package version 1.7. 2.
Http://CRAN. R-Project. Org/Package= MuMIn.
Beule, L., & Karlovsky, P. (2020). Improved normalization of species count data in
ecology by scaling with ranked subsampling (SRS): application to microbial
communities. *PeerJ*, *8*, e9593.
Bissett, A., Richardson, A. E., Baker, G., & Thrall, P. H. (2011). Long-term land use
effects on soil microbial community structure and function. *Applied Soil Ecology*,
51, 66–78.
Bünemann, E. K., Bongiorno, G., Bai, Z., Creamer, R. E., De Deyn, G., De Goede, R.,
Fleskens, L., Geissen, V., Kuyper, T. W., & Mäder, P. (2018). Soil quality–A critical
review. *Soil Biology and Biochemistry*, *120*, 105–125.
Cade, B. S. (2015). Model averaging and muddled multimodel inferences. *Ecology*,
96(9), 2370–2382. https://doi.org/https://doi.org/10.1890/14-1639.1
Calcagno, V., & de Mazancourt, C. (2010). glmulti: an R package for easy automated
model selection with (generalized) linear models. *Journal of Statistical Software*,
34, 1–29.
Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., &
Madden, T. L. (2009). BLAST+: architecture and applications. *BMC Bioinformatics*,
10, 1–9.
Camps-Valls, G., Campos-Taberner, M., Moreno-Martínez, Á., Walther, S., Duveiller, G.,
Cescatti, A., Mahecha, M. D., Muñoz-Marí, J., García-Haro, F. J., & Guanter, L.
(2021). A unified vegetation index for quantifying the terrestrial biosphere.
Science Advances, *7*(9), eabc7447.
Caporaso, J. G., Lauber, C. L., Walters, W. A., Berg-Lyons, D., Lozupone, C. A.,
Turnbaugh, P. J., Fierer, N., & Knight, R. (2011). Global patterns of 16S rRNA

diversity at a depth of millions of sequences per sample. *Proceedings of the*
National Academy of Sciences, *108*(supplement_1), 4516–4522.

Chen, Q.-L., Ding, J., Zhu, Y.-G., He, J.-Z., & Hu, H.-W. (2020). Soil bacterial taxonomic
diversity is critical to maintaining the plant productivity. *Environment*
International, *140*, 105766.
https://doi.org/https://doi.org/10.1016/j.envint.2020.105766

Cole, J. R., Wang, Q., Fish, J. A., Chai, B., McGarrell, D. M., Sun, Y., Brown, C. T., Porras-
Alfaro, A., Kuske, C. R., & Tiedje, J. M. (2014). Ribosomal Database Project: data
and tools for high throughput rRNA analysis. *Nucleic Acids Research*, *42*(D1),
D633–D642.

Conant, R. T. (2012). Grassland soil organic carbon stocks: status, opportunities,
vulnerability. *Recarbonization of the Biosphere: Ecosystems and the Global Carbon*
Cycle, 275–302.

Couto, W. (2018). Soil pH and plant productivity. In *Handbook of agricultural*
productivity (pp. 71–84). CRC Press.

Delgado-Baquerizo, M., Maestre, F. T., Reich, P. B., Jeffries, T. C., Gaitan, J. J., Encinar,
D., Berdugo, M., Campbell, C. D., & Singh, B. K. (2016). Microbial diversity drives
multifunctionality in terrestrial ecosystems. *Nature Communications*, *7*(1), 10541.
https://doi.org/10.1038/ncomms10541

Delgado-Baquerizo, M., Reich, P. B., Trivedi, C., Eldridge, D. J., Abades, S., Alfaro, F. D.,
Bastida, F., Berhe, A. A., Cutler, N. A., Gallardo, A., García-Velázquez, L., Hart, S. C.,
Hayes, P. E., He, J.-Z., Hseu, Z.-Y., Hu, H.-W., Kirchmair, M., Neuhauser, S., Pérez,
C. A., … Singh, B. K. (2020). Multiple elements of soil biodiversity drive ecosystem
functions across biomes. *Nature Ecology & Evolution*, *4*(2), 210–220.
https://doi.org/10.1038/s41559-019-1084-y

Dietrich, P., Eisenhauer, N., & Roscher, C. (2023). Linking plant diversity–productivity
relationships to plant functional traits of dominant species and changes in soil
properties in 15-year-old experimental grasslands. *Ecology and Evolution*, *13*(3),
e9883. https://doi.org/https://doi.org/10.1002/ece3.9883

Edgar, R. C. (2013). UPARSE: highly accurate OTU sequences from microbial amplicon
reads. *Nature Methods*, *10*(10), 996–998.

Edlinger, A., Garland, G., Banerjee, S., Degrune, F., García-Palacios, P., Herzog, C.,
Pescador, D. S., Romdhane, S., Ryo, M., Saghaï, A., Hallin, S., Maestre, F. T.,
Philippot, L., Rillig, M. C., & van der Heijden, M. G. A. (2023). The impact of
agricultural management on soil aggregation and carbon storage is regulated by
climatic thresholds across a 3000 km European gradient. *Global Change Biology*,
29(11), 3177–3192. https://doi.org/https://doi.org/10.1111/gcb.16677

Edlinger, A., Garland, G., Hartman, K., Banerjee, S., Degrune, F., García-Palacios, P.,
Hallin, S., Valzano-Held, A., Herzog, C., Jansa, J., Kost, E., Maestre, F. T., Pescador,
D. S., Philippot, L., Rillig, M. C., Romdhane, S., Saghaï, A., Spor, A., Frossard, E., &
van der Heijden, M. G. A. (2022). Agricultural management and pesticide use
reduce the functioning of beneficial plant symbionts. *Nature Ecology & Evolution*,
6(8), 1145–1154. https://doi.org/10.1038/s41559-022-01799-8

Fan, K., Chu, H., Eldridge, D. J., Gaitan, J. J., Liu, Y.-R., Sokoya, B., Wang, J.-T., Hu, H.-W.,
He, J.-Z., Sun, W., Cui, H., Alfaro, F. D., Abades, S., Bastida, F., Díaz-López, M.,
Bamigboye, A. R., Berdugo, M., Blanco-Pastor, J. L., Grebenc, T., … Delgado-
Baquerizo, M. (2023). Soil biodiversity supports the delivery of multiple

ecosystem functions in urban greenspaces. *Nature Ecology & Evolution*, *7*(1), 113–
126. https://doi.org/10.1038/s41559-022-01935-4

Fan, K., Delgado-Baquerizo, M., Guo, X., Wang, D., Zhu, Y., & Chu, H. (2021).
Biodiversity of key-stone phylotypes determines crop production in a 4-decade
fertilization experiment. *The ISME Journal*, *15*(2), 550–561.
https://doi.org/10.1038/s41396-020-00796-8

Fraser, L. H., Pither, J., Jentsch, A., Sternberg, M., Zobel, M., Askarizadeh, D., Bartha, S.,
Beierkuhnlein, C., Bennett, J. A., Bittel, A., Boldgiv, B., Boldrini, I. I., Bork, E.,
Brown, L., Cabido, M., Cahill, J., Carlyle, C. N., Campetella, G., Chelli, S., … Zupo, T.
(2015). Worldwide evidence of a unimodal relationship between productivity and
plant species richness. *Science*, *349*(6245), 302–305.
https://doi.org/10.1126/science.aab3916

Hu, W., Ran, J., Dong, L., Du, Q., Ji, M., Yao, S., Sun, Y., Gong, C., Hou, Q., Gong, H.,
Chen, R., Lu, J., Xie, S., Wang, Z., Huang, H., Li, X., Xiong, J., Xia, R., Wei, M., …
Deng, J. (2021). Aridity-driven shift in biodiversity–soil multifunctionality
relationships. *Nature Communications*, *12*(1), 5350.
https://doi.org/10.1038/s41467-021-25641-0

Huang, J., Liu, X., Liu, J., Zhang, Z., Zhang, W., Qi, Y., Li, W., & Chen, Y. (2023). Changes
of soil bacterial community, network structure, and carbon, nitrogen and sulfur
functional genes under different land use types. *CATENA*, *231*, 107385.

Isbell, F., Calcagno, V., Hector, A., Connolly, J., Harpole, W. S., Reich, P. B., Scherer-
Lorenzen, M., Schmid, B., Tilman, D., van Ruijven, J., Weigelt, A., Wilsey, B. J.,
Zavaleta, E. S., & Loreau, M. (2011). High plant diversity is needed to maintain
ecosystem services. *Nature*, *477*(7363), 199–202.
https://doi.org/10.1038/nature10282

Jobbágy, E. G., Sala, O. E., & Paruelo, J. M. (2002). Patterns and controls of primary
production in the Patagonian steppe: a remote sensing approach. *Ecology*, *83*(2),
307–319.

Joiner, J., Yoshida, Y., Zhang, Y., Duveiller, G., Jung, M., Lyapustin, A., Wang, Y., &
Tucker, C. J. (2018). Estimation of terrestrial global gross primary production
(GPP) with satellite data-driven models and eddy covariance flux data. *Remote
Sensing*, *10*(9), 1346.

Karimi, B., Dequiedt, S., Terrat, S., Jolivet, C., Arrouays, D., Wincker, P., Cruaud, C.,
Bispo, A., Chemidlin Prévost-Bouré, N., & Ranjard, L. (2019). Biogeography of soil
bacterial networks along a gradient of cropping intensity. *Scientific Reports*, *9*(1),
3812.

Kielak, A. M., Cipriano, M. A. P., & Kuramae, E. E. (2016). Acidobacteria strains from
subdivision 1 act as plant growth-promoting bacteria. *Archives of Microbiology*,
198(10), 987–993. https://doi.org/10.1007/s00203-016-1260-2

Köninger, J., Ballabio, C., Panagos, P., Jones, A., Schmid, M. W., Orgiazzi, A., & Briones,
M. J. I. (2023). Ecosystem type drives soil eukaryotic diversity and composition in
Europe. *Global Change Biology*.

Krause, A., Papastefanou, P., Gregor, K., Layritz, L. S., Zang, C. S., Buras, A., Li, X., Xiao,
J., & Rammig, A. (2022). Quantifying the impacts of land cover change on gross
primary productivity globally. *Scientific Reports*, *12*(1), 18398.
https://doi.org/10.1038/s41598-022-23120-0

Labouyrie, M., Ballabio, C., Romero, F., Panagos, P., Jones, A., Schmid, M. W.,
Mikryukov, V., Dulya, O., Tedersoo, L., & Bahram, M. (2023). Patterns in soil
microbial diversity across Europe. *Nature Communications*, *14*(1), 3311.
Lange, M., Eisenhauer, N., Sierra, C. A., Bessler, H., Engels, C., Griffiths, R. I., Mellado-
Vázquez, P. G., Malik, A. A., Roy, J., Scheu, S., Steinbeiss, S., Thomson, B. C.,
Trumbore, S. E., & Gleixner, G. (2015). Plant diversity increases soil microbial
activity and soil carbon storage. *Nature Communications*, *6*(1), 6707.
https://doi.org/10.1038/ncomms7707
Le Provost, G., Schenk, N. V, Penone, C., Thiele, J., Westphal, C., Allan, E., Ayasse, M.,
Blüthgen, N., Boeddinghaus, R. S., Boesing, A. L., Bolliger, R., Busch, V., Fischer,
M., Gossner, M. M., Hölzel, N., Jung, K., Kandeler, E., Klaus, V. H., Kleinebecker, T.,
… Manning, P. (2023). The supply of multiple ecosystem services requires
biodiversity across spatial scales. *Nature Ecology & Evolution*, *7*(2), 236–249.
https://doi.org/10.1038/s41559-022-01918-5
Lehmann, J., Bossio, D. A., Kögel-Knabner, I., & Rillig, M. C. (2020). The concept and
future prospects of soil health. *Nature Reviews Earth & Environment*, *1*(10), 544–
553.
Li, P., Li, Y., Xu, L., Zhang, H., Shen, X., Xu, H., Jiao, J., Li, H., & Hu, F. (2021). Crop yield-
soil quality balance in double cropping in China's upland by organic amendments:
A meta-analysis. *Geoderma*, *403*, 115197.
https://doi.org/https://doi.org/10.1016/j.geoderma.2021.115197
Liu, S., García-Palacios, P., Tedersoo, L., Guirado, E., van der Heijden, M. G. A., Wagg,
C., Chen, D., Wang, Q., Wang, J., Singh, B. K., & Delgado-Baquerizo, M. (2022).
Phylotype diversity within soil fungal functional groups drives ecosystem stability.
Nature Ecology & Evolution, *6*(7), 900–909. https://doi.org/10.1038/s41559-022-
01756-5
Louca, S., Parfrey, L. W., & Doebeli, M. (2016). Decoupling function and taxonomy in
the global ocean microbiome. *Science*, *353*(6305), 1272–1277.
Luo, J., Banerjee, S., Ma, Q., Liao, G., Hu, B., Zhao, H., & Li, T. (2023). Organic
fertilization drives shifts in microbiome complexity and keystone taxa increase the
resistance of microbial mediated functions to biodiversity loss. *Biology and*
Fertility of Soils, 1–18.
Magoč, T., & Salzberg, S. L. (2011). FLASH: fast length adjustment of short reads to
improve genome assemblies. *Bioinformatics*, *27*(21), 2957–2963.
Malik, A. A., Puissant, J., Buckeridge, K. M., Goodall, T., Jehmlich, N., Chowdhury, S.,
Gweon, H. S., Peyton, J. M., Mason, K. E., van Agtmaal, M., Blaud, A., Clark, I. M.,
Whitaker, J., Pywell, R. F., Ostle, N., Gleixner, G., & Griffiths, R. I. (2018). Land use
driven change in soil pH affects microbial carbon cycling processes. *Nature*
Communications, *9*(1), 3591. https://doi.org/10.1038/s41467-018-05980-1
Nilsson, R. H., Larsson, K.-H., Taylor, A. F. S., Bengtsson-Palme, J., Jeppesen, T. S.,
Schigel, D., Kennedy, P., Picard, K., Glöckner, F. O., & Tedersoo, L. (2019). The
UNITE database for molecular identification of fungi: handling dark taxa and
parallel taxonomic classifications. *Nucleic Acids Research*, *47*(D1), D259–D264.
Orgiazzi, A., Ballabio, C., Panagos, P., Jones, A., & Fernández-Ugalde, O. (2018). LUCAS
Soil, the largest expandable soil dataset for Europe: a review. *European Journal of*
Soil Science, *69*(1), 140–153. https://doi.org/https://doi.org/10.1111/ejss.12499

Orgiazzi, A., Panagos, P., Fernández-Ugalde, O., Wojda, P., Labouyrie, M., Ballabio, C., Franco, A., Pistocchi, A., Montanarella, L., & Jones, A. (2022). LUCAS Soil Biodiversity and LUCAS Soil Pesticides, new tools for research and policy development. *European Journal of Soil Science*, *73*(5), e13299. https://doi.org/https://doi.org/10.1111/ejss.13299

Osburn, E. D., Yang, G., Rillig, M. C., & Strickland, M. S. (2023). Evaluating the role of bacterial diversity in supporting soil ecosystem functions under anthropogenic stress. *ISME Communications*, *3*(1), 66.

Özkurt, E., Fritscher, J., Soranzo, N., Ng, D. Y. K., Davey, R. P., Bahram, M., & Hildebrand, F. (2022). LotuS2: an ultrafast and highly accurate tool for amplicon sequencing analysis. *Microbiome*, *10*(1), 1–14.

Parada, A. E., Needham, D. M., & Fuhrman, J. A. (2016). Every base matters: assessing small subunit rRNA primers for marine microbiomes with mock communities, time series and global field samples. *Environmental Microbiology*, *18*(5), 1403–1414.

Paruelo, J. M., Epstein, H. E., Lauenroth, W. K., & Burke, I. C. (1997). ANPP estimates from NDVI for the central grassland region of the United States. *Ecology*, *78*(3), 953–958.

Pertea, G. (2015). fqtrim: v0. 9.4 release. *Zenodo: Geneva, Switzerland*.

Pettorelli, N., Vik, J. O., Mysterud, A., Gaillard, J.-M., Tucker, C. J., & Stenseth, N. Chr. (2005). Using the satellite-derived NDVI to assess ecological responses to environmental change. *Trends in Ecology & Evolution*, *20*(9), 503–510. https://doi.org/https://doi.org/10.1016/j.tree.2005.05.011

Põlme, S., Abarenkov, K., Henrik Nilsson, R., Lindahl, B. D., Clemmensen, K. E., Kauserud, H., Nguyen, N., Kjøller, R., Bates, S. T., & Baldrian, P. (2020). FungalTraits: a user-friendly traits database of fungi and fungus-like stramenopiles. *Fungal Diversity*, *105*, 1–16.

Qiao, L., Wang, X., Smith, P., Fan, J., Lu, Y., Emmett, B., Li, R., Dorling, S., Chen, H., & Liu, S. (2022). Soil quality both increases crop production and improves resilience to climate change. *Nature Climate Change*, *12*(6), 574–580.

Rinot, O., Levy, G. J., Steinberger, Y., Svoray, T., & Eshel, G. (2019). Soil health assessment: A critical review of current methodologies and a proposed new approach. *Science of the Total Environment*, *648*, 1484–1491.

Romero, F., Argüello, A., de Bruin, S., & van der Heijden, M. G. A. (2023). The plant–mycorrhizal fungi collaboration gradient depends on plant functional group. *Functional Ecology*, *37*(9), 2386–2398.

Romero, F., Hilfiker, S., Edlinger, A., Held, A., Hartman, K., Labouyrie, M., & van der Heijden, M. G. A. (2023). Soil microbial biodiversity promotes crop productivity and agro-ecosystem functioning in experimental microcosms. *Science of The Total Environment*, 163683.

Sikorski, J., Baumgartner, V., Birkhofer, K., Boeddinghaus, R. S., Bunk, B., Fischer, M., Fösel, B. U., Friedrich, M. W., Göker, M., & Hölzel, N. (2022). The evolution of ecological diversity in Acidobacteria. *Frontiers in Microbiology*, *13*, 78.

Smith, L. C., Orgiazzi, A., Eisenhauer, N., Cesarz, S., Lochner, A., Jones, A., Bastida, F., Patoine, G., Reitz, T., & Buscot, F. (2021). Large-scale drivers of relationships between soil microbial properties and organic carbon across Europe. *Global Ecology and Biogeography*, *30*(10), 2070–2083.

Szoboszlay, M., Dohrmann, A. B., Poeplau, C., Don, A., & Tebbe, C. C. (2017). Impact of
land-use change and soil organic carbon quality on microbial diversity in soils
across Europe. *FEMS Microbiology Ecology*, *93*(12), fix146.

Tedersoo, L., & Anslan, S. (2019). Towards PacBio-based pan-eukaryote metabarcoding
using full-length ITS sequences. *Environmental Microbiology Reports*, *11*(5), 659–
668.

Tedersoo, L., Mikryukov, V., Anslan, S., Bahram, M., Khalid, A. N., Corrales, A., Agan, A.,
Vasco-Palacios, A.-M., Saitta, A., & Antonelli, A. (2021). The Global Soil
Mycobiome consortium dataset for boosting fungal diversity research. *Fungal*
Diversity, *111*, 573–588.

Tilman, D., Lehman, C. L., & Thomson, K. T. (1997). Plant diversity and ecosystem
productivity: theoretical considerations. *Proceedings of the National Academy of*
Sciences, *94*(5), 1857–1861.

Toda, M., Walder, F., & van der Heijden, M. G. A. (2023). Organic management and soil
health promote nutrient use efficiency. *Journal of Sustainable Agriculture and*
Environment.

Van Bruggen, A. H. C., & Semenov, A. M. (2000). In search of biological indicators for
soil health and disease suppression. *Applied Soil Ecology*, *15*(1), 13–24.

van de Vossenberg, Bart. T. L. H., Prodhomme, C., Vossen, J. H., & van der Lee, T. A. J.
(2022). Synchytrium endobioticum, the potato wart disease pathogen. *Molecular*
Plant Pathology, *23*(4), 461–474.
https://doi.org/https://doi.org/10.1111/mpp.13183

van der Heijden, M. G. A., Klironomos, J. N., Ursic, M., Moutoglis, P., Streitwolf-Engel,
R., Boller, T., Wiemken, A., & Sanders, I. R. (1998). Mycorrhizal fungal diversity
determines plant biodiversity, ecosystem variability and productivity. *Nature*,
396(6706), 69–72. https://doi.org/10.1038/23932

van der Heijden, M. G. A., & Wagg, C. (2013). Soil microbial diversity and agro-
ecosystem functioning. *Plant and Soil*, *363*, 1–5.

Wagg, C., Schlaeppi, K., Banerjee, S., Kuramae, E. E., & van der Heijden, M. G. A.
(2019). Fungal-bacterial diversity and microbiome complexity predict ecosystem
functioning. *Nature Communications*, *10*(1), 4841.
https://doi.org/10.1038/s41467-019-12798-y

Wei, T., & Simko, V. (2017). *R package "corrplot": Visualization of a Correlation Matrix*
(Version 0.84). Vienna.

Wu, T., Sabula, M., Milner, H., Strickland, G., & Liu, G. (2023). Agricultural practice
negatively affects soil bacterial diversity and nitrogen functional genes comparing
to adjacent native forest soils. *Applied Soil Ecology*, *186*, 104856.

Xiong, C., & Lu, Y. (2022). Microbiomes in agroecosystem: Diversity, function and
assembly mechanisms. *Environmental Microbiology Reports*, *14*(6), 833–849.

658