Proteomic evidence of depression-associated astrocytic dysfunction in the human male olfactory bulb

Reza Rahimian[1,2†], Kelly Perlman[1,3†], Gohar Fakhfouri[4†], Refilwe Mpai[1,3], Vincent R. Richard[5], Christa Hercher[1,3], Lucy Penney[4], Maria Antonietta Davoli[1,2], Corina Nagy[1,2,3], René P. Zahedi[6,7,8,9], Christoph H. Borchers[5,10,11,12], Bruno Giros[4], Gustavo Turecki[1,2,3], Naguib Mechawar[1,2,3*]

[1]McGill Group for Suicide Studies, Douglas Mental Health University Institute, Verdun, QC, Canada.
[2]Department of Psychiatry, McGill University, Montreal, QC, Canada.
[3]Integrated Program in Neuroscience, McGill University, Montreal, QC, Canada.
[4]Department of Psychiatry, Douglas Hospital, McGill University; Montreal, QC, Canada
[5]Segal Cancer Proteomics Centre, Lady Davis Institute for Medical Research, Jewish General Hospital, McGill University, Montréal, QC, Canada.
[6]Department of Internal Medicine, University of Manitoba, Winnipeg, MB, Canada.
[7]Manitoba Centre for Proteomics and Systems Biology, University of Manitoba, Winnipeg, MB, Canada.
[8]Department of Biochemistry and Medical Genetics, University of Manitoba, Winnipeg, MB, Canada.
[9]CancerCare Manitoba Research Institute, Winnipeg, MB, Canada.
[10]Gerald Bronfman Department of Oncology, McGill University, Montréal, Québec, Canada
[11]Department of Pathology, McGill University, Montréal, QC, Canada
[12] Department of Experimental Medicine, McGill University, Montreal, QC, Canada

† These authors contributed equally

*Corresponding author:
Naguib Mechawar, PhD
Douglas Mental Health University Institute, 6875 Boulevard LaSalle, Montréal, Quebec Canada, H4H 1R3
e-mail: naguib.mechawar@mcgill.ca

Abstract

The olfactory bulb (OB), a major structure of the limbic system, has been understudied in human investigations of psychopathologies such as depression. To explore more directly the molecular features of the OB in depression, a global comparative proteome analysis was carried out with human post-mortem OB samples from 11 males having suffered from depression and 12 healthy controls. We identified 188 differentially abundant proteins (with adjusted $p<0.05$) between depressed cases and controls. Gene ontology and gene enrichment analyses suggested that these proteins are involved in biological processes including the complement and coagulation cascades. Cell type enrichment analysis displayed a significant reduction in several canonical astrocytic proteins in OBs from depressed patients. Furthermore, using RNA-fluorescence *in-situ* hybridization, we observed a decrease in the percentage of ALDH1L1^{+} cells expressing canonical astrocytic markers including *ALDOC*, *NFIA*, *GJA1 (connexin 43)* and *SLC1A3 (EAAT1)*. These results are consistent with previous reports of downregulated astrocytic marker expression in other brain regions in depressed patients. We also conducted a comparative phosphoproteomic analysis of OB samples and found a dysregulation of proteins involved in neuronal and astrocytic functions. To determine whether OB astrocytic abnormalities is specific to humans, we also performed proteomics on the OB of socially defeated male mice, a commonly used model of depression. Cell-type specific analysis revealed that in socially defeated animals, the most striking OB protein alterations were associated with oligodendrocyte-lineage cells rather than with astrocytes, highlighting an important species difference. Overall, this study further highlights cerebral astrocytic abnormalities as a consistent feature of depression in humans.

Keywords: Olfactory bulb (OB); Depression; Limbic system; Proteomic; Phosphoproteomic; Astrocyte, Oligodendrocyte

66
67 **INTRODUCTION**
68
The olfactory bulb (OB) is a major component of the limbic system, a group of brain structures

involved in motivation, emotion, learning, and memory (Milardi et al., 2017). The OB transmits

olfactory information to other limbic areas, including the amygdala, hippocampus, and the

anterior cingulate cortex (Milardi et al., 2017). Different brain regions that participate in the

processing of olfactory information have been implicated in depression (Croy and Hummel,

2017; Kohli et al., 2016; Rottstaedt et al., 2018). Interestingly, humans with OB aplasia

(congenital anosmia) have higher rates of depressive symptoms (Croy et al., 2012; Rottstaedt et

al., 2018), and patients with depression have been reported to display decreased olfactory

sensitivity and discrimination (Lombion-Pouthier et al., 2006; Negoias et al., 2010). Moreover,

structural MRI has revealed a smaller OB volume in depressed patients compared to healthy

controls (Negoias et al., 2010). In rodents, bilateral olfactory bulbectomy results in a depression-

like phenotype and chronic antidepressant administration can rescue the associated behavioral

and physiological changes (Kelly et al., 1997; Song and Leonard, 2005). Given the highly

evolutionarily conserved OB-limbic circuitry, and the resulting functional connections between

the olfactory, salience, and emotional brain networks (Croy and Hummel, 2017; Kohli et al.,

2016; Rottstaedt et al., 2018; Takahashi et al., 2016), these observations strengthen the

hypothesis that depression is associated with impaired OB structure and function. The underlying

cellular and molecular changes, however, remain largely unexplored in humans. To examine the

molecular alterations occurring in the OB in depression, the present study aimed at performing

the first unbiased global proteome and phosphoproteome analysis of post-mortem OB samples

from depressed individuals and matched healthy controls. For comparative purposes, we the

same proteomics analysis experiments in the OB of socially defeated mice. Our main hypothesis

was that the protein profile of canonical glial markers would be altered in the OBs of humans

having suffered from depression and in the OBs of socially defeated mice.

MATERIAL AND METHODS

Human OB samples

Post-mortem OB samples from depressed patients (D) and healthy controls (CTRL) were

obtained from the Douglas-Bell Canada Brain Bank (DBCBB; Montreal, Canada). The DBCBB

collaborates closely with the Quebec Coroner's Office, and all phenotypic information was

obtained via standardized psychological autopsies, with informed consent provided by the next

of kin (Dumais et al., 2005). Either presence or suspected presence of any neurological or

neurodegenerative disorders based on clinical files was an exclusion criterion for this study.

Information from the Coroner's office, medical records, social services records, and

toxicological analysis were complemented with proxy-based interviews with one or more

informants best acquainted with the donor. Clinical vignettes realized by combining all this

information were evaluated by a panel of clinicians, using Diagnostic and Statistical Manual of

Mental Disorders (DSM-IV) criteria, to establish a diagnosis (Dumais et al., 2005; Tanti et al.,

2022).

Three distinct sets of fresh-frozen OB samples from depressed patients and healthy controls were

used in this study: the first for proteomic (12 CTRL, 11 D) and phosphoproteomic (analyses of 9

samples/group), the second for RNA-fluorescent *in situ* hybridization (5 CTRL, 3 D), and the

third for immunofluorescence (4 CTRL, 5 D). Unfortunately, although the same OB lysate was

used for proteomic and phosphoproteomic analyses, phosphoprotein enrichments were only

successful in 9 samples per group. Group characteristics, including age, post-mortem interval

(PMI), tissue pH, substance dependence, and antidepressant use, are presented in Supplementary

Tables 1, 2, 3, and 4.

Chronic social defeat stress and mouse OB samples

Animal experiments were conducted with the approval of the Douglas Animal Care Committee.

Male C57BL/6 mice were subjected to a social defeat paradigm, consisting of a daily 5-minute

defeat with a different CD-1 aggressor mouse on each of 10 consecutive days (Golden et al.,

2011) (all details are provided in the Supplementary Methods). Twenty-four hours following the

last behavioral test, socially defeated and control mice were euthanized, the OB pairs were

extracted on an ice-cold metal surface, snap frozen and kept at -80˚C for proteomics analysis.

For the social defeated group, only susceptible mice consistently displaying susceptibility

throughout the behavioral tests (Open Field, Social Interaction and Elevated Plus Maze; all

details are provided in the Supplementary Methods) following the chronic stress paradigm were

chosen. Every two pairs of OBs were pooled to serve as a single biological replicate.

Proteomic and phosphoproteomic experiments and analyses

For both human and mouse samples, proteins were extracted from whole frozen OBs in tissue

lysis buffer containing 5% SDS (Fisher Chemical), 100 mM TRIS (Fisher BioReagents™) pH

8.5 supplemented with PhosStop phosphatase inhibitor as well as cOmplete™ mini protease

inhibitor cocktail (Roche). Samples were subjected to probe-based sonication using a Thermo

Sonic Dismembrator and subsequently heated at 95° C for 10 minutes. A portion of the sample

(less than 5% by volume) was reserved for total protein quantitation by BCA assay

(Thermo/Pierce). Disulfide bonds were reduced and alkylated with 20 mM TCEP and 25 mM

iodoacetamide. 50 μg of protein per sample was digested with trypsin (Promega) overnight at

37°C using S-TRAP micro cartridges (Protifi LLC,). Peptide-containing samples were desalted

using Waters Oasis HLB SPE cartridges. 10% of the sample was reserved for measurement of

the total proteome, and the remainder was used for Fe-(III)-NTA based immobilized metal

affinity chromatography (IMAC)-based automated phosphopeptide enrichment using an Agilent

Bravo liquid handling robot and Agilent AssayMap IMAC cartridges (Beck et al., 2017;

Burkhart et al., 2012; Gonczarowska-Jorge et al., 2017). Samples were analyzed by nano Liquid

Chromatography with tandem mass spectrometry (LC-MS/MS) by data dependent acquisition

(DDA) using an Easy nLC 1200 coupled with a Q Exactive Plus mass spectrometer (Thermo

Fisher Scientific).

147
Samples were first loaded onto a pre-column (Acclaim PepMap 100 C18, 3 μm particle size, 75

μm inner diameter x 2 cm length) in 0.1% formic acid (buffer A). Peptides were then separated

using either a 100-min binary gradient for total ranging from 3-40% B (84% acetonitrile, 0.1%

formic acid) on the analytical column (Acclaim PepMap 100 C18, 2 μm particle size, 75 μm

inner diameter x 25 cm length) at 300 nL/min. MS spectra were acquired from m/z 350-1,500 at

a resolution of 70,000, with an automatic gain control (AGC) target of 1×10^6 ions and a

maximum injection time of 50 ms. The 15 most intense ions (charge states +2 to +4) were

isolated with a window of m/z 1.2, an AGC target of 2×10^4 and a maximum injection time of

64 ms and fragmented using a normalized higher-energy collisional dissociation (HCD) energy

of 28. MS/MS spectra were acquired at a resolution of 17,500 and the dynamic exclusion was set

to 30 s. DDA MS raw data was processed with Proteome Discoverer 2.5 (Thermo Scientific) and

searched using Sequest HT against a FASTA file containing all reviewed protein sequences of

the canonical human proteome without isoforms downloaded from Uniprot. The enzyme

specificity was set to trypsin with a maximum of 2 missed cleavages. Carbamidomethylation of

cysteine was set as static modification and methionine oxidation as variable modification. The

precursor ion mass tolerance was set to 10 ppm, and the product ion mass tolerance was set to

0.02 Da. The percolator node was used, and the data was filtered using a false discovery rate

(FDR) cut-off of 1% at both the peptide and protein level. The Minora feature detector node of

Proteome Discoverer was used for precursor-based label free quantitation (Beck et al., 2017;

Burkhart et al., 2012; Gonczarowska-Jorge et al., 2017). Only peptides that are unique to a

protein were used for quantitation. In cases where a protein was not found in a particular sample,

we imputed the missing areas using Low Abundance Resampling, an algorithm that randomly

uses an area value from the lowest 5% of those detected (Gonczarowska-Jorge et al., 2017).

For human samples, proteins or phosphoprotein modifications that were not detected in at least

50% of subjects in both cases and controls were removed. This conservative approach was

chosen to increase the robustness of our results given the small sample size, as it ensured that

each protein had absolute quantities that were as representative of the group as possible (i.e., no

imputation required) without excluding too many proteins. For mouse samples, we elected to

remove any protein that was not detected in at least 1/3 of the samples from both groups, since

each sample was derived from the combination of 2 OBs and inter-subject variability was lower

(6 OBs in control and 6 OBs in social defeated mice). The underlying data structure was

visualized using PCA and Grubb's test was used to identify outliers for removal. Differentially

abundant proteins (DAP), referring to proteins whose abundances are significantly different

between cases and controls, were computed with t-tests that have "background" adjusted p-

values which means that a weighting factor is applied to the p-values that takes into account the

distribution of ratios for all other proteins as well as the protein abundances themselves (Navarro

et al., 2014). P-values were corrected for multiple comparisons with the Benjamini-Hochberg

method. This background t-test is designed to fulfill the assumption of data normality as it

handles data such that the variance is constant for the reference points of the background. All

downstream analyses and data visualizations were conducted in R. Gene ontology and gene set

enrichment analyses were conducted with the clusterProfiler and DOSE packages (Korotkevich

et al., 2021; Yu et al., 2012; Yu et al., 2015). Ridgeplots are used visualize top terms as

determined by the gseGO function, which was ran with the parameter values used in the online

vignette   (https://learn.gencore.bio.nyu.edu/rna-seq-analysis/gene-set-enrichment-analysis/).

DAPs were compared with the literature using the DisGeNet database (Piñero et al., 2017) and

assessed for overlap between our DAPs and genes that were established to be differentially

expressed in the following conditions: Mental Depression (CUI: C0011570), Unipolar

Depression (CUI: C0041696), Endogenous depression (CUI: C0011573), Severe depression

(CUI: C0588008), Depression and Suicide (CUI: C1524032), clinical depression (CUI:

C2362914), Major Depressive Disorder (CUI: C1269683), Major depression (CUI: C1269683),

provided that the gda score was greater than 0.01. Over enrichment of cell types based on DAPs

was calculated by comparing published sets of gene lists and using fisher's exact test to look for

statistically significant overabundance. The list of genes for each major cell type was formed by

taking the union of the markers from the McKenzie et al. (2018) and Qiu et al. (2021), with

corresponding MarkerPen tool, plus genes selected to complete the cell type categories which

were previously deemed insufficient, primary the oligodendrocytes (OLs).

RNA-fluorescence in-situ hybridization (FISH) in human OB samples

Frozen OBs were cut in serial 10 µm-thick dorsoventral sections with a cryostat, and sections

were collected on Superfrost charged slides. *In situ* hybridization was performed using Advanced

Cell Diagnostics RNAscope® probes and reagents following the manufacturer's instructions

(Reemst et al., 2022). Sections were first fixed in cold 10% neutral buffered formalin for

15 min, dehydrated by increasing gradient of ethanol baths, and air-dried for 5 min.

Endogenous peroxidase activity was quenched with hydrogen peroxide for 10 min followed by

protease digestion for 30 min at room temperature (Tanti et al., 2022). The following probes

were then hybridized for 2 h at 40 °C in a humidity-controlled oven: Hs-ALDH1L1 (ACDbio,

REF;438881), Hs-ALDOC (ACDbio, REF;407031), Hs-GJA1(ACDbio, REF;444281), Hs-NFIA

(ACDbio, REF;442781) and Hs-SLC1A3 (ACDbio, REF;461081). Amplifiers were added using

the AMP reagents and the signal visualized through probe-specific HRP-based detection by

tyramide signal amplification (TSA, ACDbio) with Opal dyes (Opal 520 and Opal 570; Perkin

Elmer) diluted 1:900. In order to eliminate endogenous autofluorescence from lipofuscin and

cellular debris, sections were incubated with TrueBlack (Biotium, Cat; 23007) for 30 seconds.

Slides were then cover slipped with Vectashield mounting medium with 4′,6-diamidino-2-

phenylindole (DAPI) for nuclear staining (Vector Laboratories, H-1800) and kept at 4 °C until

imaging. Both positive and negative controls provided by the supplier (ACDbio) were used on

separate sections to confirm signal specificity. The slides were imaged at 20X magnification

using the Olympus VS120 virtual slide scanner and the scans were transferred to QuPath

(v.0.3.0) for further analysis. OB area (whole section) was demarcated manually and QuPath was

employed for automated cell detection based on DAPI staining and RNAscope signal

quantification. For each probe, only cells bearing three or more fluorescent puncta were counted

as positively labeled.

Immunofluorescence in human OB samples

OBs were placed in 10% neutral buffered formalin for 2 hours. Samples were then stored in a 30% sucrose solution until they were processed for cryosectioning and flash-frozen in isopentane. Samples were then sectioned at 30 μm on a cryostat, collected on charged Super Frost Plus slides and stored at -80□. Slides were rinsed in phosphate-buffered saline (PBS) and incubated in a blocking solution of PBS, 0.2% Triton-X and 10% normal donkey serum (NDS) for an hour at 37□. Following this, sections were briefly rinsed in PBS before being incubated overnight at 4□ with an anti-ALDH1L1 antibody (Millipore, 1:250, MABN495 | Proteintech, 1:500, 17390-1-AP) in a blocking solution of PBS/0.2% Triton-X/5% NDS. Slides were then rinsed and incubated for 1 hour at room temperature with the secondary Alexa-647 anti-mouse antibody (Jackson ImmunoResearch, 1:500, 715-605-151) diluted in the same blocking solution. Next, sections were rinsed and endogenous autofluorescence from lipofuscin and cellular debris were quenched with TrueBlack (Biotium) and coverslipped with Vectashield mounting medium (Vector Laboratories, H-1800). For ALDH1L1 quantification, sections were scanned using a Slide Scanner Olympus VS120 and ALDH1L1$^+$ astrocytes then quantified using Qupath (Bankhead, 2017). For each subject, two OB sections per subject were quantified. To minimize variability the dorsal most and ventral most sections from each OB were used. Within each section, two ROIs were selected such that both the right and the left part of each tissue section was sampled. Each ROI comprised of 28 250x250 μm grid squares. Within each ROI, all ALDH1L1$^+$ cells were counted in the same z-plane. The presence of a DAPI-labelled nucleus

overlapping with the ALDH1L1$^+$ immunolabelling was required for positive identification of an

astrocyte. Group characteristics, including age, PMI, tissue pH, substance dependence, and

medication, are presented in Supplementary Tables 4.

Statistical analysis

R and GraphPad Prism (v8) were employed to perform statistical analyses. For animal studies

and FISH experiments, unpaired Student t tests were used to compare different behavioral

parameters and quantifications, respectively. For FISH quantifications, double- labeled

astrocytes were assessed using QuPath (v0.3.2) and reported as a percentage of total astrocytes

(ALDH1L1$^+$). $p < 0.05$ was set as the threshold for statistical significance in all analyses.

RESULTS
262
Altered expression of canonical astrocyte proteins in OB of depressed individuals

We identified 188 DAPs (with corrected p<0.05 for group differences) between cases and

controls, which was narrowed down to 115 proteins when using a 2-sigma fold change cut-off

(Figure 1A).

Supplementary Table 5 shows the number of proteins detected before and after the selected

conservative cut-off described in the methods. Gene set enrichment analyses suggested that the

DAPs are involved in general biological processes such as cellular detoxification (e.g., GSTM1,

HBG1, HBD) and nucleotide metabolic process (e.g., CMPK2, CDAKD, ALDOB, SULT1A1)

but the KEGG enrichment tool suggested specific enrichment for complement and coagulation

cascade (e.g., C3, F2, PLG, C4BPA, CFH, CFB). Given the scarcity of molecular OB research in

depression, we sought to assess the extent to which the list of DAPs overlapped with

differentially expressed genes reported in the literature for other brain regions in humans. To this

end, we used the DisGeNeT database and considered the differentially expressed genes that were

common to studies of depressive disorders. We then interrogated the overlap (i.e., intersection)

of those genes with the DAPs identified in the current study and found 38 overlapping candidates

from our initial total list of 188 proteins (Figure 1D). These 38 candidates included genes

involved in the complement cascade, lipoprotein regulation, and response to oxidative stress.

Since the proteomics analyses were performed on bulk OB tissue, we examined whether the

DAP list would suggest any cell type specificity to the alterations. We assessed the overlap

between broad cell type specific marker genes (astrocyte, endothelial, microglial, OL-lineage,

and neuronal) provided in Supplementary Table 6, constructed as described in the Methods

section. We applied hypergeometric tests on this list to determine cell type enrichment and found

that the only significantly enriched cell types were the astrocyte and endothelial cells (astrocyte,

$p = 0.0128$; endothelial, $p = 0.0356$; microglia, $p = 0.220$; OL-lineage, $p = 0.893$; neuron, $p =$

0.993; Figure 1E). However, since the PMI was not matched between groups (Supplementary

Table 2), we elected to validate our findings with an alternative analysis. We ran linear models

with the following structure: Protein ~ Group + Age + PMI. We selected proteins whose models

had p<0.05 for the group factor as the DAPs. We then conducted the cell type marker enrichment

analysis with these new DAPs and found that only astrocyte markers reached significance

(astrocyte, $p = 0.0249$; endothelial, $p = 0.896$; microglia, $p = 0.118$; OL-lineage, $p = 0.986$;

neuron, $p = 0.495$) (Supplementary Figure 1). Given that astrocyte markers were found to be

significantly enriched with both analysis methods, we considered this as sufficient evidence to

focus primarily on astrocytes going forward.

Canonical markers GFAP ($p = 0.00508$) and SLC1A3 ($p = 0.0254$) - a glial high affinity

glutamate transporter - were among the markers significantly downregulated in depressed OBs in

the proteomic analysis while astrocyte-specific markers ALDH1L1, ALDOC, NFIA and GJA1

showed a downregulation trend (Figure 2A). To confirm these observations in situ, at the

transcriptional level, with a highly sensitive method, we carried out multiplexed fluorescence in

situ hybridization (FISH; RNAScope®) to assess canonical astrocytic marker transcripts. We

used the pan-astrocytic marker ALDH1L1 and counted positive cells as cells that expressed both

the target transcript and ALDH1L1. These experiments confirmed that ALDOC, SLC1A3

(EAAT1), GJA1 (Connexin 43) and NFIA are primarily expressed in human OB astrocytes

(Supplementary Figure 2). Furthermore, the number of ALDH1L1-positive cells was not

significantly different between groups ($p = 0.1406$). However, the number of astrocytes

expressing ALDOC ($p = 0.0272$), GJA1 ($p = 0.0021$), NFIA ($p = 0.0003$) and SLC1A3 ($p =$

0.0030) were significantly reduced in the OB of depressed individuals compared to controls

(Figure2B-F). As ALDOC is a glycolytic enzyme, its downregulation in depression can be

related to the reduction of astrocytic glutamate transporters such as SLC1A3, which was also

observed in the OB of cases. According to the astrocyte-neuron lactate-shuttle hypothesis,

astrocytes upregulate glycolysis to produce lactate for neurons when glutamate is taken up from

the synapse (O'Leary and Mechawar, 2021; Pellerin and Magistretti, 1994; Zhao et al., 2016).

Given the results obtained with FISH showing no significant group difference in cells expressing

ALDH1L1, we examined this further using IF with an antibody against ALDH1L1. As illustrated

in Supplementary figure 3, results were consistent, revealing no significant group difference in

the density of ALDH1L1-immunoreactive cells in the dorsal ($p = 0.556$) nor ventral ($p = 0.905$)

OB sections. Similarly, pooling data for both OB regions led to similar densities between cases

and controls ($p = 0.905$).

We then accessed a single nucleus RNA sequencing (snRNA-seq) dataset recently generated by

our group with human dorsolateral PFC (dlPFC) from depressed suicides and control adult males

(Maitra et al., 2023) to compare the differentially expressed genes in its astrocyte clusters to our

proteome results. Despite differences in brain regions, we found overlapping genes between the

astrocyte 1 and 2 dlPFC cluster from the Maitra et al. study (in a re-analysis of the male data

from Nagy et al., 2020) and the proteomics data presented here, with all 8 overlapping genes

displaying a downregulation compared to controls, according to log fold change (Supplemental

Table 7) which indicates a strong concordance between RNA and protein changes (Maitra et al.,

2023; Nagy et al., 2020). Some of these proteins, such as AQP1, MT1X and S1PR1, are highly

expressed by astrocytes, and elicit crucial roles in homeostatic and immune-related functions.

Namely, S1PR1 is implicated in proliferation and trafficking of immune cells (Rothhammer et

al., 2017). By generating high levels of antioxidants including metallothionein-I (MT1X),

astrocytes play a key role in offsetting oxidative stress associated with inflammation (Waller et

al., 2018).

We then verified that these depression-associated changes in OB astrocytic proteins were not

driven by confounders such as child abuse history and antidepressant treatment. As illustrated in

Supplementary Figure 4, no significant relationship was found between protein quantities and

either of these potential confounders using the Mann-Whitney U-Test (child abuse (CA):

ALDOC, $p = 0.456$; ALDH1L1, $p = 0.611$; GFAP, $p = 0.2505$; GJA1, $p = 0.845$; NFIA, $p =$

0.785; SLC1A3, $p = 0.667$. Antidepressant treatment: ALDOC, $p = 0.413$; ALDH1L1, $p = 0.928$;

GFAP, $p = 0.133$; GJA1, $p = 0.413$; NFIA, $p = 0.880$; SLC1A3, $p = 0.316$).

Dysregulation of phosphoproteins in the OB of depressed individuals

The phosphoproteome is more dynamic than the proteome, thus we expected some discordance

with the proteome results. We found 57 differentially abundant phosphopeptides, corresponding

to 50 individual proteins, all of which met the 2-sigma fold change cut-off (Figure 3A). Gene

ontology indicated that the corresponding proteins of the differentially abundant

phosphopeptides are involved in biological processes such as synaptic vesicle cycle (e.g.,

AMPH, SYN1, SYN2, PPFIA2), phosphorus metabolic processes (e.g., PNPO, ATP2B4,

SLC4A1), and RNA splicing (e.g., TRA2A, SRSF1, SRRM1, THRAP3) (Figure 3B). We

assessed overlap between this list of 57 phosphopeptides and DisGeNET differentially expressed

genes for depression and found 10 overlapping candidates (Figure 3C). We performed the same

cell-type specific hypergeometric tests as in the proteomic analysis and that both neurons (p =

0.0125), and astrocytes (p = 0.0281) were significantly enriched, with neurons having the highest

enrichment score (endothelial, p = 0.665; microglia, p = 1.000; OL-lineage, p = 0.733; Figure

3D).

We also examined the relationship between the proteome and phosphoproteome differential

abundance. As listed in Supplemental Table 8, there are eight proteins overlapping between the

two analyses, including GFAP and AQP1, which are fairly specific to astrocytes in the adult

human brain. SLC1A4 had two differentially abundant phosphopeptides, both upregulated, as

was observed for the protein. In fact, the regulation direction was found to be consistent for all

proteins. For instance, both GFAP protein and a GFAP phosphopeptide were downregulated in

cases compared to controls and was identified in the DisGeNET dataset as well.

Dysregulation of oligodendrocyte proteins in OB of socially defeated mice

To examine whether human OB findings could be observed in a mouse model of depression, we

performed the same proteomics analyses with socially defeated mice and controls. Proteins were

extracted from 3 control and 3 socially-defeated OB pairs (each sample was derived from 2 OBs,

yielding 6 subjects per group in total). As presented in the Methods section, only susceptible

mice consistently showing susceptibility throughout the behavioral tests (Open Field, Social

Interaction and Elevated Plus Maze) following chronic stress were chosen for proteomic analyses

(Figure 4A-C). The analysis revealed 261 significantly DAPs, 193 of which also met the fold

change cut-off (Figure 4D). Gene ontology analyses suggested that these proteins are involved in

functions such as fatty acid biosynthetic process (e.g., ptgds, anxa1, acsm3) and neurogenesis

(e.g. snapin, ndel1, cit) (Figure 4E). We found 35 DAPs overlapping with the DisGeNET

database, indicating overlap with genes previously implicated in MDD (Figure 4F). Strikingly,

we found a different pattern of cell type enrichment in the DAPs, as the only cell type reaching

our significance cut-off was OL-lineage cells ($p = 0.0453$), with endothelial cells showing a trend

(endothelial $p = 0.0679$, astrocyte $p = 0.733$, microglia $p = 0.627$, neuronal $p = 0.375$) (Figure

4G). For instance, plp1 ($p = 0.0106$), anln ($p = 0.0299$), apc ($p = 0.000765$), ptgds ($p =$

0.00000309) and tf ($p = 0.000000189$) were among the strongly dysregulated OL markers in

socially defeated mice. These genes have all been implicated in the process of myelination, with

plp1 being the most abundant protein in myelin of the central nervous system (Jahn et al., 2009).

We then conducted a species comparison and found 13 overlapping DAPs including cat

(involved in oxidative stress) and apoa1 (involved in lipid metabolism). According to log fold

change, 11 of these 13 proteins were regulated in the same direction in both species

(Supplemental Table 9).

DISCUSSION

This is the first study to compare adult OB proteomic and phosphoproteomic profiles between depressed individuals and healthy controls. We observed that depressed OBs presented DAPs that are enriched for astrocyte markers, which led us to explore whether these findings were reflective of changes in the astrocyte population. While the percentage of ALDH1L1$^+$ astrocytes remained similar between cases and controls, the percentages of astrocytes expressing the canonical markers ALDOC, GJA1, NFIA, and SLC1A3 were significantly reduced in the OB of cases. In addition to these cell type-enriched changes, we also found that there was a substantial overlap between depression-associated dysregulated OB proteins and genes previously implicated in depression across other brain regions. Moreover, parallel analyses conducted with a mouse model of depression revealed that, in contrast to humans, differentially abundant proteins in the OB of socially stressed animals were enriched for oligodendrocyte-lineage markers.

Astrocytic abnormalities have been consistently implicated in depression and suicide (Torres-Platas et al., 2016), with reported downregulations of astrocyte-specific genes, decreased immunoreactivity to astrocyte-canonical markers, and reduced blood vessel coverage by astrocytic endfeet in postmortem brain samples from depressed patients (Rajkowska et al., 2013). Surprisingly, and despite neuroimaging and psychometric studies implicating the OB in depression, this major limbic region had never been examined at the cellular/molecular level in depressed patients. Our results provide the first evidence that in depression, astrocytic abnormalities are present in the OB, in agreement with our previous reports of astrocytic abnormalities in other cortico-limbic brain regions of depressed suicides (O'Leary et al., 2021; O'Leary and Mechawar, 2021; Torres-Platas et al., 2016; Xiong et al., 2019), as well as with

independent animal studies having implicated astrocytes in depression-like behavior (Abbink et

al., 2020; Codeluppi et al., 2021; Du Preez et al., 2021; Rurak et al., 2021; Simard et al., 2018).

Depression-associated changes in astrocytic numbers and functions could result from a variety

factors, such as chronic glucocorticoid exposure or altered neurotrophic or neurotransmitter

signaling (Rajkowska and Miguel-Hidalgo, 2007). The current study indirectly implicates the

role of astrocytic transcription factor NFIA, which has been reported to regulate astrocyte

morphology and function in the rodent hippocampus (Huang et al., 2020). In addition, these

authors found that NFIA-deficient astrocytes or simply hypo-functional NFIA can impair

neuron-astrocyte communication *in vitro* (Huang et al., 2020). Interestingly, within 5 days,

transient NFIA expression *in vitro* activates glial potential in human pluripotent stem cells,

transforming them into astrocytes upon exposure to glial-promoting factors. These NFIA-

induced astrocytes not only foster synaptogenesis but also exhibit neuroprotective properties

(Tchieu et al., 2019). We can speculate that the reduction in NFIA measured here in the OB of

depressed individuals impairs the morpho-functional properties of astrocytes, having in turn an

impact on OB volume and olfactory sensitivity, which have both been documented in depressed

patients (Negoias et al., 2010; Pause et al., 2001). A smaller OB volume has been associated with

higher rates of depressive episodes and with decreased signaling with other limbic areas

(Athanassi et al., 2021; Croy and Hummel, 2017). Altered olfaction in depression has been also

hypothesized to be linked to a decreased receptor turnover rate in the olfactory epithelium

(Athanassi et al., 2021; Croy and Hummel, 2017). Our finding of astrocytic disruptions in the

OB of depressed individuals might be associated with this phenomenon and contribute directly to

changes in perception, since mature OB astrocytes, at least in mice, are required for normal

physiological sensory processing (Ung et al., 2021).

To better understand the effects of protein regulation in depression, our study also compared

differentially phosphorylated proteins in OB samples from depressed patients compared to

healthy controls. Changes in phosphorylation almost always reflects a change in protein activity,

although many such modifications remain uncharacterized (Ardito et al., 2017). In agreement

with our DAP analysis, which highlighted an enrichment of astrocyte proteins in the OB of

depressed individuals, the phosphoproteomic analysis revealed a corresponding enrichment in

differentially phosphorylated astrocytic proteins. Interestingly, it also revealed enrichment in

differentially phosphorylated neuronal proteins. It is well established that astrocytes are an

essential component of synapses, regulating both synaptic and circuit plasticity and function

(Perez-Catalan et al., 2021). In consequence, the astrocytic dysregulation suggested by our study

may impact neuronal functions at various levels, including through post-translational

modifications of synaptic proteins. Furthermore, given the converging evidence implicating

alterations in synaptic density, synapse microenvironment, and synaptic vesicle release (Duman

et al., 2016; Holmes et al., 2019; Ren and Guo, 2021) in depression, it is not surprising that some

of our most prominent differentially abundant phosphoproteins are localized at the synapse. This

can be illustrated by synapsin (SYN) 1 and 2, which were found to be significantly different

between cases and controls. Synapsins are ubiquitous phosphoproteins on synaptic vesicles,

where they regulate the trafficking of vesicles in nerve terminals, a critical parameter in short-

term synaptic plasticity (Sun et al., 2006). A previous study of depression in postmortem dlPFC

from males and females pointed to synaptic function as the main biological process associated

with differentially abundant phosphoproteins (Martins-de-Souza et al., 2012). Interestingly, 10

out of the 50 phosphoproteins we found to be differentially abundant were similarly

differentially abundant in the latter study, despite the regional difference. In our results, SYN 1

was one of the top differentially abundant phosphoproteins with three significantly

downregulated phosphopeptides, indicating hypophosphorylation of SYN 1 in depressed

individuals compared to controls. This finding was mirrored by significantly downregulated

SYN 1 phosphopetides in Martins-de-Souza et al. (2012) (Martins-de-Souza et al., 2012). The

sizable overlap between these two studies further strengthens the evidence for post-translational

modifications occurring at the synapse in depression.

We then conducted the same OB proteomic analysis in a mouse model of depression. These

experiments in the social defeat mouse model indicated that the enriched astrocyte signature

observed in humans with depression is not shared by chronically stressed mice, possibly

highlighting a limitation of this model which recapitulates many of the molecular signatures of

depression in regions such as the medial prefrontal cortex (mPFC) and nucleus accumbens

(Scarpa et al., 2020).

The social defeat model of human depression comes with some limitations. This model involves

daily defeat and thus fights, which involve stress, wounds, and local inflammatory events. The

model also differs in many ways from the post-mortem human brains from depressed patients.

Namely, depression-like states in mice appear on vastly different timescales than those of disease

processes implicated in depressive symptoms in humans. Moreover, there exist differences in

pathological and adaptive mechanisms when comparing this animal model and depressed

humans. Finally, social defeat is performed almost exclusively in adult animals, which cannot

inform on the contribution of early-life events to the depression in adulthood.

Our observation adds to the extensive breadth of evidence supporting OL lineage abnormalities

that have been reported in post-mortem and animal model studies of depression (Liu et al., 2012;

Makinodan et al., 2012; Nagy et al., 2020; Rahimian et al., 2022). In a study of chronic social

defeat stress, decreases in myelin proteins were observed in the nucleus accumbens and mPFC,

with the mPFC also displaying thinner myelin and decreased internode length (Bonnefil et al.,

2019). In a subsequent study, this model led long lasting reductions of OL precursor cells (OPC),

abnormal differentiation rates of OPC to OLs, and extensive hypomyelination (Kokkosis et al.,

2022). In fact, these findings are not limited to social defeat stress, as myelin-related molecular

and ultrastructural changes have been reported in several models of stress including social

isolation and chronic variable stress (Bonnefil et al., 2019; Liu et al., 2012; Rahimian et al.,

2022). While our OL enrichment findings may point to differences in myelin content or

regulation in socially defeated mice, further experiments are needed to better understand the

biological significance of these results. It is also a possibility that discordance in results might

reflect differences in OB cellular composition between species. In summary, while there was

overlap between human and mouse DAPs, as well as an overlap with previous findings from

DisGeNET, the cell type enrichment and gene ontology analyses indicated considerable

divergence between results in humans and mice, indicating possible mechanistic differences

between the murine social defeat model and human depression, at least in what concerns the OB.

It is plausible to assume that this divergence is at least partially related to the disproportionately

greater OB/brain volume in mice compared to humans.

The current study contributes to a better understanding of the cellular and molecular factors

associated with depression, which is required for the design of impactful translational studies,

including research on diagnostic or prognostic biomarkers as well as on emerging therapeutics. A

growing body of evidence suggests that astrocyte-derived extracellular vesicles can be obtained

non-invasively from the serum of clinically depressed patients and inform on the functional state

of astrocytes *in vivo* (Xie et al., 2023). It remains to be determined if peripheral levels of these

proteins are correlated with astrocytic deficits in one or multiple regions of the CNS and with

one or more astrocytic subtypes.

The main limitations of our study were its sample size and the fact that all the OBs were from

male donors. It is challenging to collect postmortem OB samples, and particularly samples that

are of sufficient quality for proteomics, and this becomes even more difficult by the very low

number of OBs collected from female middle-aged depressed individuals over the years. As a

result, we were unable to include female samples in a meaningful way. Given the distinct male

vs female molecular signatures of depression that have been reported in other brain regions,

especially in astrocytes and microglia (Maitra et al., 2023) and the sexual dimorphism in OB

cellular composition in humans (Oliveira-Pinto et al., 2014), future investigations will need to

circumvent these challenges to explore whether the OB is similarly affected in females affected

by depression. Lastly, since all but one of the cases included in this study died by suicide, we

cannot exclude that the proteomic signature reported in this study is more representative of male

depressed suicides than of all depressed individuals (Peng et al., 2023).

Overall, the findings of our study further implicate astrocytes in depression. The consistency of

cortico-limbic astrocyte findings in depression research lends support to the notion that this cell

type might play a key role in the pathophysiology of this heterogeneous illness. Furthermore, our

study raises the possibility that the chronic social defeat stress model of depressive-like behavior

might not sufficiently recapitulate the molecular neurobiology of human depression and suicide,

at least in the OB. Understanding the strengths and weaknesses of animal models of depression

will help to better position pre-clinical research questions such that their translatability to human

depression may be enhanced.

528   Data availability

529   The mass spectrometry proteomics data have been deposited to the ProteomeXchange

530   Consortium via the PRIDE partner repository with the dataset identifier PXD045011

532   **Project Name:** Proteomic evidence of depression-associated astrocytic dysfunction in the human

533   male olfactory bulb

534   **Project accession:** PXD045011

535
536

537   The processed spreadsheets for human proteomics and phosphoproteomics, as well as the mouse

538   proteomics have been provided as Supplementary Tables 10 (human proteomics). Supplementary

539   Table 11 (human phosphoproteomics), and Supplementary Table 12 (mouse proteomics). The

540   cell type marker list for the enrichment analysis has been as Supplementary Table 6 as well.

542   REFERENCES

Abbink, M.R., Kotah, J.M., Hoeijmakers, L., Mak, A., Yvon-Durocher, G., van der Gaag, B.,
Lucassen, P.J., Korosi, A., 2020. Characterization of astrocytes throughout life in
wildtype and APP/PS1 mice after early-life stress exposure. J. Neuroinflammation 17, 91.
Ardito, F., Giuliani, M., Perrone, D., Troiano, G., Lo Muzio, L., 2017. The crucial role of protein
phosphorylation in cell signaling and its use as targeted therapy (Review). Int. J. Mol.
Med. 40, 271-280.
Athanassi, A., Dorado Doncel, R., Bath, K.G., Mandairon, N., 2021. Relationship between
depression and olfactory sensory function: a review. Chem. Senses 46.
Beck, F., Geiger, J., Gambaryan, S., Solari, F.A., Dell'Aica, M., Loroch, S., Mattheij, N.J.,
Mindukshev, I., Potz, O., Jurk, K., Burkhart, J.M., Fufezan, C., Heemskerk, J.W., Walter,
U., Zahedi, R.P., Sickmann, A., 2017. Temporal quantitative phosphoproteomics of ADP
stimulation reveals novel central nodes in platelet activation and inhibition. Blood 129,
e1-e12.
Bonnefil, V., Dietz, K., Amatruda, M., Wentling, M., Aubry, A.V., Dupree, J.L., Temple, G.,
Park, H.J., Burghardt, N.S., Casaccia, P., Liu, J., 2019. Region-specific myelin

differences define behavioral consequences of chronic social defeat stress in mice. Elife
8.
Burkhart, J.M., Schumbrutzki, C., Wortelkamp, S., Sickmann, A., Zahedi, R.P., 2012.
Systematic and quantitative comparison of digest efficiency and specificity reveals the
impact of trypsin quality on MS-based proteomics. J. Proteomics 75, 1454-1462.
Codeluppi, S.A., Chatterjee, D., Prevot, T.D., Bansal, Y., Misquitta, K.A., Sibille, E., Banasr,
M., 2021. Chronic Stress Alters Astrocyte Morphology in Mouse Prefrontal Cortex. Int.
J. Neuropsychopharmacol. 24, 842-853.
Croy, I., Hummel, T., 2017. Olfaction as a marker for depression. J. Neurol. 264, 631-638.
Croy, I., Negoias, S., Novakova, L., Landis, B.N., Hummel, T., 2012. Learning about the
functions of the olfactory system from people without a sense of smell. PLoS One 7,
e33365.
Du Preez, A., Onorato, D., Eiben, I., Musaelyan, K., Egeland, M., Zunszain, P.A., Fernandes, C.,
Thuret, S., Pariante, C.M., 2021. Chronic stress followed by social isolation promotes
depressive-like behaviour, alters microglial and astrocyte biology and reduces
hippocampal neurogenesis in male mice. Brain. Behav. Immun. 91, 24-47.
Dumais, A., Lesage, A.D., Lalovic, A., Seguin, M., Tousignant, M., Chawky, N., Turecki, G.,
2005. Is violent method of suicide a behavioral marker of lifetime aggression? Am. J.
Psychiatry 162, 1375-1378.
Duman, R.S., Aghajanian, G.K., Sanacora, G., Krystal, J.H., 2016. Synaptic plasticity and
depression: new insights from stress and rapid-acting antidepressants. Nat. Med. 22, 238-
249.
Golden, S.A., Covington, H.E., 3rd, Berton, O., Russo, S.J., 2011. A standardized protocol for
repeated social defeat stress in mice. Nat. Protoc. 6, 1183-1191.
Gonczarowska-Jorge, H., Loroch, S., Dell'Aica, M., Sickmann, A., Roos, A., Zahedi, R.P., 2017.
Quantifying missing (phospho)proteome regions with the broad-specificity protease
subtilisin. Anal. Chem. 89, 13137-13145.
Holmes, S.E., Scheinost, D., Finnema, S.J., Naganawa, M., Davis, M.T., DellaGioia, N., Nabulsi,
N., Matuskey, D., Angarita, G.A., Pietrzak, R.H., Duman, R.S., Sanacora, G., Krystal,
J.H., Carson, R.E., Esterlis, I., 2019. Lower synaptic density is associated with depression
severity and network alterations. Nat Commun 10, 1529.
Huang, A.Y., Woo, J., Sardar, D., Lozzi, B., Bosquez Huerta, N.A., Lin, C.J., Felice, D., Jain,
A., Paulucci-Holthauzen, A., Deneen, B., 2020. Region-Specific Transcriptional Control
of Astrocyte Function Oversees Local Circuit Activities. Neuron 106, 992-1008 e1009.
Jahn, O., Tenzer, S., Werner, H.B., 2009. Myelin proteomics: molecular anatomy of an
insulating sheath. Mol. Neurobiol. 40, 55-72.
Kelly, J.P., Wrynn, A.S., Leonard, B.E., 1997. The olfactory bulbectomized rat as a model of
depression: an update. Pharmacol. Ther. 74, 299-316.
Kohli, P., Soler, Z.M., Nguyen, S.A., Muus, J.S., Schlosser, R.J., 2016. The Association
Between Olfaction and Depression: A Systematic Review. Chem. Senses 41, 479-486.
Kokkosis, A.G., Madeira, M.M., Mullahy, M.R., Tsirka, S.E., 2022. Chronic stress disrupts the
homeostasis and progeny progression of oligodendroglial lineage cells, associating
immune oligodendrocytes with prefrontal cortex hypomyelination. Mol. Psychiatry 27,
2833-2848.
Korotkevich, G., Sukhov, V., Budin, N., Shpak, B., Artyomov, M.N., Sergushichev, A., 2021.
Fast gene set enrichment analysis. bioRxiv, 060012.

Liu, J., Dietz, K., DeLoyht, J.M., Pedre, X., Kelkar, D., Kaur, J., Vialou, V., Lobo, M.K., Dietz,
D.M., Nestler, E.J., Dupree, J., Casaccia, P., 2012. Impaired adult myelination in the
prefrontal cortex of socially isolated mice. Nat. Neurosci. 15, 1621-1623.
Lombion-Pouthier, S., Vandel, P., Nezelof, S., Haffen, E., Millot, J.L., 2006. Odor perception in
patients with mood disorders. J. Affect. Disord. 90, 187-191.
Maitra, M., Mitsuhashi, H., Rahimian, R., Chawla, A., Yang, J., Fiori, L.M., Davoli, M.A.,
Perlman, K., Aouabed, Z., Mash, D.C., Suderman, M., Mechawar, N., Turecki, G., Nagy,
C., 2023. Cell type specific transcriptomic differences in depression show similar patterns
between males and females but implicate distinct cell types and genes. Nat Commun 14,
2912.
Makinodan, M., Rosen, K.M., Ito, S., Corfas, G., 2012. A critical period for social experience-
dependent oligodendrocyte maturation and myelination. Science 337, 1357-1360.
Martins-de-Souza, D., Guest, P.C., Vanattou-Saifoudine, N., Rahmoune, H., Bahn, S., 2012.
Phosphoproteomic differences in major depressive disorder postmortem brains indicate
effects on synaptic function. Eur. Arch. Psychiatry Clin. Neurosci. 262, 657-666.
McKenzie, A.T., Wang, M., Hauberg, M.E., Fullard, J.F., Kozlenkov, A., Keenan, A., Hurd,
Y.L., Dracheva, S., Casaccia, P., Roussos, P., Zhang, B., 2018. Brain cell type specific
gene expression and co-expression network architectures. Sci. Rep. 8, 8868.
Milardi, D., Cacciola, A., Calamuneri, A., Ghilardi, M.F., Caminiti, F., Cascio, F., Andronaco,
V., Anastasi, G., Mormina, E., Arrigo, A., Bruschetta, D., Quartarone, A., 2017. The
Olfactory System Revealed: Non-Invasive Mapping by using Constrained Spherical
Deconvolution Tractography in Healthy Humans. Front. Neuroanat. 11, 32.
Nagy, C., Maitra, M., Tanti, A., Suderman, M., Theroux, J.F., Davoli, M.A., Perlman, K., Yerko,
V., Wang, Y.C., Tripathy, S.J., Pavlidis, P., Mechawar, N., Ragoussis, J., Turecki, G.,
2020. Single-nucleus transcriptomics of the prefrontal cortex in major depressive disorder
implicates oligodendrocyte precursor cells and excitatory neurons. Nat. Neurosci. 23,
771-781.
Navarro, P., Trevisan-Herraz, M., Bonzon-Kulichenko, E., Nunez, E., Martinez-Acedo, P.,
Perez-Hernandez, D., Jorge, I., Mesa, R., Calvo, E., Carrascal, M., Hernaez, M.L.,
Garcia, F., Barcena, J.A., Ashman, K., Abian, J., Gil, C., Redondo, J.M., Vazquez, J.,
2014. General statistical framework for quantitative proteomics by stable isotope
labeling. J. Proteome Res. 13, 1234-1247.
Negoias, S., Croy, I., Gerber, J., Puschmann, S., Petrowski, K., Joraschky, P., Hummel, T., 2010.
Reduced olfactory bulb volume and olfactory sensitivity in patients with acute major
depression. Neuroscience 169, 415-421.
O'Leary, L.A., Belliveau, C., Davoli, M.A., Ma, J.C., Tanti, A., Turecki, G., Mechawar, N.,
2021. Widespread Decrease of Cerebral Vimentin-Immunoreactive Astrocytes in
Depressed Suicides. Front Psychiatry 12, 640963.
O'Leary, L.A., Mechawar, N., 2021. Implication of cerebral astrocytes in major depression: A
review of fine neuroanatomical evidence in humans. Glia 69, 2077-2099.
Oliveira-Pinto, A.V., Santos, R.M., Coutinho, R.A., Oliveira, L.M., Santos, G.B., Alho, A.T.,
Leite, R.E., Farfel, J.M., Suemoto, C.K., Grinberg, L.T., Pasqualucci, C.A., Jacob-Filho,
W., Lent, R., 2014. Sexual dimorphism in the human olfactory bulb: females have more
neurons and glial cells than males. PLoS One 9, e111733.
Pause, B.M., Miranda, A., Goder, R., Aldenhoff, J.B., Ferstl, R., 2001. Reduced olfactory
performance in patients with major depression. J. Psychiatr. Res. 35, 271-277.

Pellerin, L., Magistretti, P.J., 1994. Glutamate uptake into astrocytes stimulates aerobic
glycolysis: a mechanism coupling neuronal activity to glucose utilization. Proc. Natl.
Acad. Sci. U. S. A. 91, 10625-10629.
Peng, S., Zhou, Y., Xiong, L., Wang, Q., 2023. Identification of novel targets and pathways to
distinguish suicide dependent or independent on depression diagnosis. Sci. Rep. 13, 2488.
Perez-Catalan, N.A., Doe, C.Q., Ackerman, S.D., 2021. The role of astrocyte-mediated plasticity
in neural circuit development and function. Neural Dev 16, 1.
Piñero, J., Bravo, À., Queralt-Rosinach, N., Gutiérrez-Sacristán, A., Deu-Pons, J., Centeno, E.,
García-García, J., Sanz, F., Furlong, L.I., 2017. DisGeNET: a comprehensive platform
integrating information on human disease-associated genes and variants. Nucleic Acids
Res. 45, D833-d839.
Qiu, Y., Wang, J., Lei, J., Roeder, K., 2021. Identification of cell-type-specific marker genes
from co-expression patterns in tissue samples. Bioinformatics 37, 3228-3234.
Rahimian, R., Perlman, K., Canonne, C., Mechawar, N., 2022. Targeting microglia-
oligodendrocyte crosstalk in neurodegenerative and psychiatric disorders. Drug Discov
Today 27, 2562-2573.
Rajkowska, G., Hughes, J., Stockmeier, C.A., Javier Miguel-Hidalgo, J., Maciag, D., 2013.
Coverage of blood vessels by astrocytic endfeet is reduced in major depressive disorder.
Biol. Psychiatry 73, 613-621.
Rajkowska, G., Miguel-Hidalgo, J.J., 2007. Gliogenesis and glial pathology in depression. CNS
Neurol. Disord. Drug Targets 6, 219-233.
Reemst, K., Kracht, L., Kotah, J.M., Rahimian, R., van Irsen, A.A.S., Congrains Sotomayor, G.,
Verboon, L.N., Brouwer, N., Simard, S., Turecki, G., Mechawar, N., Kooistra, S.M.,
Eggen, B.J.L., Korosi, A., 2022. Early-life stress lastingly impacts microglial
transcriptome and function under basal and immune-challenged conditions. Transl
Psychiatry 12, 507.
Ren, F., Guo, R., 2021. Synaptic Microenvironment in Depressive Disorder: Insights from
Synaptic Plasticity. Neuropsychiatr. Dis. Treat. 17, 157-165.
Rothhammer, V., Kenison, J.E., Tjon, E., Takenaka, M.C., de Lima, K.A., Borucki, D.M., Chao,
C.C., Wilz, A., Blain, M., Healy, L., Antel, J., Quintana, F.J., 2017. Sphingosine 1-
phosphate receptor modulation suppresses pathogenic astrocyte activation and chronic
progressive CNS inflammation. Proc. Natl. Acad. Sci. U. S. A. 114, 2012-2017.
Rottstaedt, F., Weidner, K., Strauss, T., Schellong, J., Kitzler, H., Wolff-Stephan, S., Hummel,
T., Croy, I., 2018. Size matters - The olfactory bulb as a marker for depression. J. Affect.
Disord. 229, 193-198.
Rurak, G.M., Woodside, B., Aguilar-Valles, A., Salmaso, N., 2021. Astroglial cells as
neuroendocrine targets in forebrain development: Implications for sex differences in
psychiatric disease. Front. Neuroendocrinol. 60, 100897.
Scarpa, J.R., Fatma, M., Loh, Y.E., Traore, S.R., Stefan, T., Chen, T.H., Nestler, E.J., Labonte,
B., 2020. Shared Transcriptional Signatures in Major Depressive Disorder and Mouse
Chronic Stress Models. Biol. Psychiatry 88, 159-168.
Simard, S., Coppola, G., Rudyk, C.A., Hayley, S., McQuaid, R.J., Salmaso, N., 2018. Profiling
changes in cortical astroglial cells following chronic stress. Neuropsychopharmacology
43, 1961-1971.
Song, C., Leonard, B.E., 2005. The olfactory bulbectomised rat as a model of depression.
Neurosci. Biobehav. Rev. 29, 627-647.

Sun, J., Bronk, P., Liu, X., Han, W., Sudhof, T.C., 2006. Synapsins regulate use-dependent
synaptic plasticity in the calyx of Held by a Ca2+/calmodulin-dependent pathway. Proc.
Natl. Acad. Sci. U. S. A. 103, 2880-2885.
Takahashi, T., Nishikawa, Y., Yucel, M., Whittle, S., Lorenzetti, V., Walterfang, M.,
Sasabayashi, D., Suzuki, M., Pantelis, C., Allen, N.B., 2016. Olfactory sulcus
morphology in patients with current and past major depression. Psychiatry Res
Neuroimaging 255, 60-65.
Tanti, A., Belliveau, C., Nagy, C., Maitra, M., Denux, F., Perlman, K., Chen, F., Mpai, R.,
Canonne, C., Théberge, S., McFarquhar, A., Davoli, M.A., Belzung, C., Turecki, G.,
Mechawar, N., 2022. Child abuse associates with increased recruitment of perineuronal
nets in the ventromedial prefrontal cortex: a possible implication of oligodendrocyte
progenitor cells. Mol. Psychiatry 27, 1552-1561.
Tchieu, J., Calder, E.L., Guttikonda, S.R., Gutzwiller, E.M., Aromolaran, K.A., Steinbeck, J.A.,
Goldstein, P.A., Studer, L., 2019. NFIA is a gliogenic switch enabling rapid derivation of
functional human astrocytes from pluripotent stem cells. Nat. Biotechnol. 37, 267-275.
Torres-Platas, S.G., Nagy, C., Wakid, M., Turecki, G., Mechawar, N., 2016. Glial fibrillary
acidic protein is differentially expressed across cortical and subcortical regions in healthy
brains and downregulated in the thalamus and caudate nucleus of depressed suicides.
Mol. Psychiatry 21, 509-515.
Ung, K., Huang, T.W., Lozzi, B., Woo, J., Hanson, E., Pekarek, B., Tepe, B., Sardar, D., Cheng,
Y.T., Liu, G., Deneen, B., Arenkiel, B.R., 2021. Olfactory bulb astrocytes mediate
sensory circuit processing through Sox9 in the mouse brain. Nat Commun 12, 5230.
Waller, R., Murphy, M., Garwood, C.J., Jennings, L., Heath, P.R., Chambers, A., Matthews,
F.E., Brayne, C., Ince, P.G., Wharton, S.B., Simpson, J.E., Cognitive, F., Ageing
Neuropathology Study, G., 2018. Metallothionein-I/II expression associates with the
astrocyte DNA damage response and not Alzheimer-type pathology in the aging brain.
Glia 66, 2316-2323.
Xie, X.H., Lai, W.T., Xu, S.X., Di Forti, M., Zhang, J.Y., Chen, M.M., Yao, L.H., Wang, P.,
Hao, K.K., Rong, H., 2023. Hyper-inflammation of astrocytes in patients of major
depressive disorder: Evidence from serum astrocyte-derived extracellular vesicles. Brain.
Behav. Immun. 109, 51-62.
Xiong, W., Cao, X., Zeng, Y., Qin, X., Zhu, M., Ren, J., Wu, Z., Huang, Q., Zhang, Y., Wang,
M., Chen, L., Turecki, G., Mechawar, N., Chen, W., Yi, G., Zhu, X., 2019. Astrocytic
epoxyeicosatrienoic acid signaling in the medial prefrontal cortex modulates depressive-
like behaviors. J. Neurosci. 39, 4606-4623.
Yu, G., Wang, L.G., Han, Y., He, Q.Y., 2012. clusterProfiler: an R package for comparing
biological themes among gene clusters. OMICS 16, 284-287.
Yu, G., Wang, L.G., Yan, G.R., He, Q.Y., 2015. DOSE: an R/Bioconductor package for disease
ontology semantic and enrichment analysis. Bioinformatics 31, 608-609.
Zhao, J., Verwer, R.W., van Wamelen, D.J., Qi, X.R., Gao, S.F., Lucassen, P.J., Swaab, D.F.,
2016. Prefrontal changes in the glutamate-glutamine cycle and neuronal/glial glutamate
transporters in depression with and without suicide. J. Psychiatr. Res. 82, 8-15.

739
740

741 **AUTHOR CONTRIBUTIONS**

ACKNOWLEDGMENTS

This study was funded by an ERA NET Neuron grant and a CIHR Project grant to N.M. R.R.

and G.F. received fellowships and K.P. a scholarship from the FRQ-S. K.P. currently holds a

CIHR doctoral scholarship and R.M. a scholarship from the Botswana government. The

Douglas-Bell Canada Brain Bank is supported in part by platform support grants from the

Réseau Québécois sur le Suicide, les Troubles de l'Humeur et les Troubles Associés (FRQ-S),

Healthy Brains for Healthy Lives (CFREF), and Brain Canada. This study used the services of

the Molecular and Cellular Microscopy Platform (MCMP) at the Douglas Research Centre.

This work was also supported by funding to CHB from Genome Canada's Genomics

Technology Platform (264PRO). CHB is also grateful for support from the Segal McGill Chair

in Molecular Oncology at McGill University (Montreal, Quebec, Canada) for support from

the Terry Fox Research Institute, the Warren Y. Soper Charitable Trust, and the Alvin Segal

Family Foundation to the Jewish General Hospital (Montreal, Quebec, Canada).

This work was done under the auspices of a Memorandum of Understanding between McGill

and the U.S. National Cancer Institute's International Cancer Proteogenome Consortium (ICPC).

ICPC encourages international cooperation among institutions and nations in proteogenomic

cancer research in which proteogenomic datasets are made available to the public. This work was

also done in collaboration with the U.S. National Cancer Institute's Clinical Proteomic Tumor

Analysis Consortium (CPTAC).

Lastly, the authors wish to thank the donors and their families for their invaluable gift.

COMPETING INTERESTS

CHB is the CSO of MRM Proteomics Inc., and the VP of Proteomics at Molecular You. RPZ is

CEO of MRM Proteomics. The remaining authors declare no conflict of interest.

776
777 **ADDITIONAL INFORMATION**
778
Supplementary information

Figure legends

Figure 1. Human OB proteomics reveal enrichment in astrocyte-specific differentially

abundant proteins in depressed individuals. (**A**) Volcano plot with log base 2 abundance ratio

between cases and controls on the x-axis and the $-\log_{10}$(p-value) on the y-axis. Each dot

represents a protein, and they are colored by direction of regulation with downregulation in pink

and upregulation in turquoise. The log base 2 abundance ratio cutoff is |2.5| for visualization

purposes and the p-value is cutoff is $-\log_{10}$(0.05). D = Depression, CTRL = Control. (**B**)

Graphical representation of the DAPs contributing to the enrichment of the complement and

coagulation cascade biological process identified by the enrichKEGG function. The node color

of the DAP represents the log2 fold change. (**C**) Ridge plot with the enrichment of the gene

ontology (GO) terms on the x-axis, and each GO biological process listed on the y-axis. This plot

shows the over enriched GO terms in the list of differentially abundant proteins between cases

and controls. The ridges are colored by adjusted p-value of the enrichment tests as demonstrated

by the color bar. The positive enrichment distribution values indicate enrichment of the GO term

in cases compared to controls, and the negative enrichment distribution values indicate

enrichment in controls compared to cases. (**D**) Table of overlap of proteins in list of differentially

abundant proteins and genes found in the DisGeNET database. Proteins are listed in descending

order by log base 2-fold change between cases and controls. Proteins with positive fold change

values are colored in pink and negative fold change values are colored in blue. (**E**) Bar plot

showing cell type enrichment scores of differentially abundant proteins. The x-axis shows the

broad cell types of the brain, and the y-axis shows the $-\log_{10}(\text{p-value})$, where the p-value refers to

the enrichment p-value from hypergeometric tests. The dotted line refers to the significance

cutoff which is $-\log_{10}(0.05)$. Only the astrocytes surpass this significant threshold.

Figure 2. Translational deregulation of canonical astrocytic markers in OBs of depressed

individuals (**A**) Heat map illustrating differentially abundant canonical astrocytic markers in

OBs from CTRL individuals as opposed to those from depressed subjects. GFAP and SLC1A3

proteins were significantly downregulated in MDD OBs (respectively, $p = 0.0013$ and $p =$

0.017), while ALDOC, NFIA, ALDH1L1, and GJA1 showed a downregulation trend. (**B**) Left:

Representative microphotographs of FISH experiments on CTRL and MDD OBs using

ALDH1L1 probe. Right: percentage of ALDH1L1$^+$ cells, $p = 0.1406$. **C-F**) Left: Representative

microphotographs of FISH experiments on control (CTRL) and depressed patients (D) OBs

using ALDH1L1 probe along with (**C**) ALDOC, (**D**) GJA1, (**E**) NFIA, or (**F**) SLC1A3 probes.

Right: percentage of double positive astrocytes over the total number of astrocytes (ALDH1L1$^+$

cells). (**C**) $p = 0.0272$; (**D**) $p = 0.0021$; (**E**) $p = 0.0003$; (**F**) $p = 0.0030$. (**B-F**) CTRL: n = 3;

MDD: n = 5; scale bars: 20µm.

Figure 3. Human OB differential phosphoproteomics in depressed individuals. (**A**) Volcano

plot with log base 2 abundance ratio between cases and controls on the x-axis and the $-\log_{10}$(p-

value) on the y-axis. Each dot represents a protein that showed significant phosphopeptide

differential regulation between cases and controls, and the points are colored by direction of the

regulation of the abundance of the phosphopeptide modification with downregulation in pink and

upregulation in turquoise. The log base 2 abundance ratio cutoff is |3| for visualization purposes

and the p-value is cutoff is $-\log_{10}$(0.05). D = Depression, CTRL = Control. (**B**) Ridge plot with

the enrichment of the gene ontology (GO) terms on the x-axis, and each GO biological process

listed on the y-axis. This plot shows the over enriched GO terms in the list of differentially

abundant proteins between cases and controls. The ridges are colored by adjusted p-value of the

enrichment tests as demonstrated by the color bar. The positive enrichment distribution values

indicate enrichment of the GO term in cases compared to controls, and the negative enrichment

distribution values indicate enrichment in controls compared to cases (**C**) Table of overlap of

proteins in list of proteins with differentially abundant phosphopeptide and genes found in the

DisGeNET database. Proteins are listed in descending order by log base 2-fold change between

cases and controls. Abundance of the differentially abundant phosphopeptide modification for

the protein between cases and controls is represented by log base 2-fold change, phosphopeptide

with positive fold change values are colored in pink and negative fold change values are colored

in blue. In cases where there were multiple phosphopeptide regulations for a given protein, the

average fold change across all modifications was used. (**D**) Bar plot showing cell type

enrichment scores of differentially abundant proteins. The x-axis shows the broad cell types of

the brain, and the y-axis shows the $-\log_{10}$(p-value), where the p-value refers to the enrichment p-

value from hypergeometric tests. The dotted line refers to the significance cutoff which is -

$\log_{10}(0.05)$. Only the neurons and astrocytes surpass this significance threshold.

Figure 4. OB proteomics of social defeat stress mice do not recapitulate human OB

proteomics of depressed individuals. Plots **A**-**C** demonstrate the behavioral tests conducted on

control and socially defeated mice, CTRL, n = 6; SD, n = 8. (**A**) $p = 0.002$; (**B**) $p = 0.012$; (**C**) p

$= 0.012$. (**D**) Volcano plot with log base 2 abundance ratio between cases and controls on the x-

axis and the $-\log_{10}$(p-value) on the y-axis. Each dot represents a protein, and they are colored by

direction of regulation with downregulation in pink and upregulation in turquoise. The log base 2

abundance ratio cutoff is |2.5| for visualization purposes and the p-value is cutoff is $-\log_{10}(0.05)$.

(**E**) Ridge plot with the enrichment of the gene ontology (GO) terms on the x-axis, and each GO

biological process listed on the y-axis. This plot shows the over enriched GO terms in the list of

differentially abundant proteins between cases and controls. The ridges are colored by adjusted

p-value of the enrichment tests as demonstrated by the color bar. The positive enrichment

distribution values indicate enrichment of the GO term in SD compared to controls, and the

negative enrichment distribution values indicate enrichment in controls compared to SD. (**F**)

Table of overlap of proteins in list of differentially abundant proteins and genes found in the

DisGeNET database. Proteins are listed in descending order by log base 2-fold change between

cases and controls. Proteins with positive fold change values are colored in pink and negative

fold change values are colored in blue. (**G**) Bar plot showing cell type enrichment scores of

differentially abundant proteins. The x-axis shows the broad cell types of the brain, and the y-

axis shows the $-\log_{10}$(p-value), where the p-value refers to the enrichment p-value from

hypergeometric tests. The dotted line refers to the significance cutoff which is $-\log_{10}(0.05)$. Only

the OL-lineage cells surpass this significance threshold.

883 **Figures**

884 **Figure 1**

893   **Figure 2**

Figure 3

A

B

C

D

Name	log₂FC D/CTRL
SYN2	4.94
HSP90AB1	2.66
GFAP	-2.12
NEFM	-2.34
MBP	-2.51
SGIP1	-2.88
DPYSL2	-3.57
SYN1	-3.8
NDRG2	-5.07
STK32C	-6.26

905   **Figure 4**

D

Name	log$_2$FC D/CTRL
SYN2	4.94
HSP90AB1	2.66
GFAP	-2.12
NEFM	-2.34
MBP	-2.51
SGIP1	-2.88
DPYSL2	-3.57
SYN1	-3.8
NDRG2	-5.07
STK32C	-6.26

Name	logFC SD/CTRL
Commd3	2.63
Aass1	1.48
Pion	0.68
Hmgb1	0.68
Hdac6	0.66
NccL	0.65
Arb	0.57
Ntrk2	0.55
Nrp1	0.52
Nans	0.52
Atg7	0.50
Hspa2	0.50
Ampa2	0.49
Crt	0.42
Stfts1	0.39
Rida	0.27
Th	0.19
Plp1	0.18
Nqo1	0.17
Krfm	0.14
Actb	-0.21
Cbln1	-0.26
Actg1	-0.32
Gja1	-0.37
Nign	-0.39
Mora	-0.39
Pbgds	-0.59
Sort1	-0.76
Crt	-0.76
NccB	-0.87
Pam	-0.93
Ndn1	-0.95
Vgf	-1.42
Pcmsd1	-1.42
AcxB	-3.06